The Spatial Atlas of Human Anatomy (SAHA): A Multimodal Subcellular-Resolution Reference Across Human Organs

Jiwoon Park[1,2], Roberto De Gregorio[1,3], Erika Hissong[1,4], Elif Ozcelik[1], Nicholas Bartelo[1], Felipe Segato Dezem[5,13], Luke Zhang[5,6], Maycon Marção[5,6], Hannah Chasteen[5,6], Yimin Zheng[7], Ernesto Abila[7,8], Junbum Kim[1], Jacqueline Proszynski[1], Akua A. Agyemang[1], Mohith Reddy Arikatla[1], Arjumand Wani[5,6], Yutian Liu[5,6], Evelyn Metzger[9], Stefan Rogers[9], Prajan Divakar[9], Parambir S. Dulai[10,11], Jason Reeves[9], Yan Liang[9], Liuliu Pan[9], Sayani Bhattacharjee[9], Michael Patrick[9], Kimberly Young[9], Ashley Heck[9], Mithra Korukonda[9], Dan McGuire[9], Lidan Wu[9], Aster Wardhani[9], Joseph Beechem[9], George Church[2], Steven M Lipkin[1], Sanjay Patel[1,4], Fabio Socciarelli[1,4], Rohit Chandwani[1], Sebastien Monette[12], Brian Robinson[1,4], Massimo Loda[1,4], Olivier Elemento[1,3], Luciano Martelotto[13,*], Jasmine Plummer[5,6,14,15,*], André F. Rendeiro[7,8,*], Alicia Alonso[1,3,*], Robert E. Schwartz[1,*], Shauna Lee Houlihan[1,3,*], Christopher E. Mason[1,*]

Affiliations:
[1] Weill Cornell Medicine, New York, NY, USA
[2] Harvard Medical School, Boston, MA, USA
[3] Caryl and Israel Englander Institute for Precision Medicine, Weill Medical College of Cornell University, New York, USA
[4] Department of Pathology and Laboratory Medicine, Weill Medical College of Cornell University, New York, NY, USA
[5] Center for Spatial Omics, St. Jude Children's Research Hospital, Memphis, TN, USA
[6] Department of Developmental Neurobiology, St. Jude Children's Research Hospital, Memphis, TN, USA
[7] CeMM Research Center for Molecular Medicine of the Austrian Academy of Sciences, Vienna, Austria
[8] Ludwig Boltzmann Institute for Network Medicine at the University of Vienna, Austria
[9] Bruker Spatial Biology, Seattle, WA, USA
[10] Department of Medicine, Division of Gastroenterology and Hepatology, Northwestern University Feinberg School of Medicine
[11] Center for Human Immunobiology, Northwestern University Feinberg School of Medicine
[12] Memorial Sloan Kettering Cancer Center, New York, NY, USA
[13] University of Adelaide, Adelaide, Australia
[14] Comprehensive Cancer Center, St. Jude Children's Research Hospital, Memphis, TN, USA
[15] Department of Cellular & Molecular Biology, St. Jude Children's Research Hospital, Memphis, TN, USA

*Corresponding author(s); Main contact: Christopher E. Mason (chm2042@med.cornell.edu)

ABSTRACT

The Spatial Atlas of Human Anatomy (SAHA) represents the first multimodal, subcellular-resolution reference of healthy adult human tissues across multiple organ systems. Integrating spatial transcriptomics, proteomics, and histological features across over 15 million cells from more than 100 donors, SAHA maps conserved and organ-specific cellular niches in gastrointestinal and immune tissues. High-resolution profiling using CosMx SMI, 10x Xenium, RNAscope, GeoMx DSP, and single-nucleus RNA-seq reveals spatially organized cell states, rare adaptive immune populations, and tissue-specific cell-cell interactions. Comparative analyses with colorectal cancer and inflammatory bowel disease demonstrate the power of SAHA to detect disease-associated spatial disruptions, including crypt dedifferentiation, perineural invasion, and therapy-resistant immune remodeling. All data are openly accessible through a FAIR-compliant interactive portal to support exploration, benchmarking, and machine learning model training. Through SAHA, we provide a foundational framework for spatial diagnostics and next-generation precision medicine grounded in a comprehensive human tissue atlas.

INTRODUCTION

Human tissues normally maintain structural integrity and function across diverse physiological states, but how these architectures break down in disease, aging, and stress remains a fundamental yet largely unresolved question in biology and medicine. The advent of spatial omics technologies has transformed our ability to map gene and protein expression within intact tissues, offering subcellular-resolution insights into cellular interactions and tissue architecture [1], [2]. Yet a critical gap remains: there is no comprehensive spatial reference of healthy human tissues across multiple organ systems. Existing atlases, including the Human Cell Atlas [3], Human Protein Atlas, and Human Tumor Atlas Network [4], have advanced our understanding of single-cell biology, but are largely derived from dissociated cells or disease-specific samples, limiting their utility as healthy spatial baselines [5].

To address this critical need [6], we established the Spatial Atlas of Human Anatomy (SAHA)—the first multimodal, subcellular-resolution reference atlas of healthy adult human tissues spanning gastrointestinal and immune systems. SAHA integrates spatial transcriptomic, proteomic, and histological data from over 100 diverse donors, capturing cellular organization at ~50 nm resolution. Unlike prior efforts, SAHA implements a rigorously standardized pipeline for tissue procurement, processing, quality control, and cross-platform integration. Our initiative enables consistent, reproducible analysis across organs, individuals, and spatial profiling technologies, while preserving the architectural context often lost in dissociated single-cell studies.

SAHA not only provides a high-resolution baseline of normal tissue organization but also facilitates comparative analyses with diseased tissues, enabling identification of spatially defined pathologies such as immune-epithelial remodeling, crypt dedifferentiation, and rare invasive niches. Through an open-access, FAIR-compliant [7] data portal (link to be updated

83 upon acceptance), SAHA aims to serve as a foundational benchmark for translational research,
84 computational modeling, and spatially informed diagnostics [8].
85
Here, we report the first phase of SAHA encompassing 16 gastrointestinal and immune tissue
types and demonstrate its utility for spatially resolved analysis of diseases, such as colorectal
cancer, pancreatic cancer, and inflammatory bowel disease. SAHA represents the largest
subcellular resolution, multimodal transcriptomic and proteomic dataset of healthy human
tissues to date, enabling the characterization of both conserved and organ-specific spatial
niches. Through comparative analyses with diseased cohorts, we illustrate SAHA's translational
potential as a reference framework for identifying pathological alterations. By bridging spatial
molecular profiling with clinical precision medicine, SAHA establishes a new paradigm for
functional tissue atlasing and a blueprint for the next generation of spatial omics research.
95
96
RESULTS
1. Overview of the first gastrointestinal and immune spatial organ atlas
The first phase of the Spatial Atlas of Human Anatomy (SAHA) systematically mapped major
digestive and immune organs at subcellular resolution, capturing over 15 million cells from over
100 healthy adult donors (ages 27-81 years, **Supplementary Table 1**). Using multiple high-plex
spatial platforms including CosMx™ Spatial Molecular Imager, Xenium, RNAscope, and
GeoMx® Digital Spatial Profiler, we captured high-resolution profiles of stomach, liver,
pancreas, small intestine (ileum), large intestine (descending colon and appendix), bone
marrow, and lymph nodes. To provide a molecular bridge between spatially resolved tissue
architecture and a readily available systemic readout, we also included peripheral blood
mononuclear cells (PBMCs), which represent a non-solid tissue component (**Fig. 1,**
Supplementary Table 2). Across these efforts, we profiled 15,915,616 cells, including 9.5
million tissue-resident cells and 6 million circulating blood cells, at a spatial resolution of 50
nanometers. For the normal reference tissues, the CosMx RNA (1000-plex) dataset included 94
tissue samples, profiling 2.9 million cells and capturing over 524 million individual transcript
measurements. This 1000-gene spatial atlas represents a substantial advancement in
transcriptomic resolution—enabling detailed cell type phenotyping and cell–cell interaction
analysis while preserving subcellular spatial fidelity. The corresponding CosMx protein dataset
(67-plex) profiled 3.5 million cells and yielded over 17.8 billion quantified protein features. In
contrast, disease and PBMC samples were analyzed using expanded RNA panels (6000-plex or
19,000-plex), providing deeper molecular coverage for characterizing rare populations and
disease-specific states.
119
Standardized tissue collection and rigorous quality control were implemented to minimize batch
effects and cross-platform variability, ensuring robust data integration and downstream analysis
[9] (See **Methods**, **Extended Data Fig. 1, Supplementary Table 2**). This enabled joint analysis
of transcriptomic profiles across organs, revealing both conserved and organ-specific cellular
identities. Uniform Manifold Approximation and Projection (UMAP) of combined CosMx RNA
data demonstrated distinct clustering of major epithelial lineages—such as hepatocytes (liver),
acinar and ductal cells (pancreas), and gastric parietal and chief cells (stomach)—alongside

broad immune populations including T cells, B cells, macrophages, and dendritic cells (**Fig. 2a,**
Extended Data Fig. 2a). In total, over 50 distinct cell types were annotated using canonical
marker genes, encompassing broad epithelial, immune, stromal, and neuronal lineages,
highlighting both conserved and organ-specialized programs of cellular diversity and relative
abundance (**Fig. 2b-d**).
132
Subcellular resolution enabled precise localization of transcripts and proteins within individual
cellular compartments, allowing us to map the anatomical distribution of billions of biomolecules
across tissue sections. While higher resolution can increase technical variability and noise, our
quality control strategy (i.e., histopathological review, quantification of control probes,
RNAscope-based RNA integrity assessment, and platform cross-validation) ensured robust
reproducibility across tissues and donors (**Extended Data Fig. 3**). For example, RNAscope
control probes (i.e., *PPIB*) benchmarked RNA quality, and Spearman correlation analyses
demonstrated a strong monotonic relationship ($\rho = 0.86$, $p = 0.0007$) between RNAscope and
CosMx transcript measurements, without significant systematic bias, while CosMx maintained
greater transcriptome coverage due to its higher plex capacity (**Extended Data Fig. 3a-c**).
Together, these analyses establish the technical reproducibility and biological validity of SAHA
datasets and cell annotations, enabling robust cross-organ and cross-platform comparisons and
providing a reliable reference for future spatial studies. Quality metrics across sample cohorts
were highly consistent (**Extended Data Fig. 3d**), underscoring the robustness of the integrated
data resource. This inaugural phase of SAHA establishes the largest high-quality, multimodal
spatial reference of the human digestive and immune systems at subcellular resolution—
providing a foundational resource for investigating tissue architecture, cellular diversity, and
microenvironmental organization in health and disease.
151
152
2. Cellular neighborhoods and interactions across intestinal and immune structures
Leveraging an integrated embedding of over 15 million spatially profiled cells, SAHA enables
both cross-organ comparisons and high-resolution analysis of spatial architecture within and
across organ systems, tissue types, and cell populations (**Fig. 2**). This flexible framework
supports multiscale investigation of tissue organization and microenvironmental interactions. In
the gastrointestinal system, high-plex spatial profiling revealed nuanced, layer-specific
architectures and discrete cellular niches spanning the epithelium and underlying submucosa
(**Fig. 3a**). In the colon, for instance, epithelial crypts were distinctly segmented, with *PIGR* and
MHC-I expression localized to mucosal and submucosal regions, and smooth muscle markers
(e.g., *ACTA2*) confined to the muscularis layer. Cell-level adjacency mapping further uncovered
localized immune infiltration at crypt boundaries, suggesting active epithelial-immune crosstalk
that is not readily detectable in dissociated single-cell datasets.
165
SAHA provides a topographical link between structure and single-cell activity. Analysis of
regionally enriched gene programs demonstrated spatial specialization along tissue depth axis
(**Fig. 3b**). Surface regions were enriched for keratinization-associated signatures, whereas
deeper crypt zones expressed genes involved in antigen-processing and presentation. Spatial
mapping of cell cycle regulators (e.g., *CDKN1A, CCND1*) and stress-response genes (e.g.,

TFEB) showed selective enrichment at immune-rich epithelial interfaces, suggesting localized
zones of proliferative and immunomodulatory adaptation (**Fig. 3c**). Unbiased clustering of cell-
cell adjacency networks further delineated distinct cellular neighborhoods across crypt and
submucosal compartments (**Fig. 3d-f**). Discrete microenvironments dominated by epithelial cell
to T cells, B cells, or myofibroblast interactions emerged along the crypt-luminal axis,
emphasizing substantial spatial heterogeneity even within morphologically similar layers. These
findings highlight the unique power of subcellular-resolution spatial profiling to uncover
microanatomical, structural features that are inaccessible through dissociated single-cell
approaches.
180
Beyond crypt-associated immune niches, gastrointestinal organs also revealed highly organized
lymphoid structures. In the appendix (**Fig. 3g, Extended Data Fig. 4**), we identified large
lymphoid aggregates composed of interspersed T and B cells, surrounded by mesenchymal and
myeloid populations. Force-directed graph visualizations of cell-cell adjacency networks
demonstrated dense immune-immune interactions within these aggregates, consistent with their
role as specialized hubs for adaptive immune activation (**Fig. 3h**). UMAP embeddings
annotated by both broad and fine-grained cell identities highlighted tight spatial co-localization of
functionally distinct immune subsets, including T follicular helper cells and germinal follicle
center B cells (**Fig. 3i**). Ligand-receptor interaction analyses further revealed frequent paracrine
signaling among these populations, suggesting coordinated immunological activity within these
spatially confined niches (**Fig. 3j**).
192
These lymphoid aggregates occupy anatomically defined regions within gastrointestinal tissues
and serve as sites of antigen presentation, lymphocyte activation, and localized immune
coordination. To assess the generalizability of such spatially organized immune niches, we
compared them with canonical germinal centers (GCs) in lymph nodes (LNs). Unsupervised
clustering and spatial mapping of LN datasets revealed distinct compartmentalization between
GC-resident B cells and surrounding immune and stromal populations (**Fig. 4a, Extended Data
Fig. 5a-b**). Differential expression analysis further highlighted this organization, with B cell and
antigen presentation genes (e.g., *CD74, CD79A, HLA-DRA*) enriched in GC cores (q-values =
5.47×10^{-34}, 1.89×10^{-10}, 1.20×10^{-20}), while stromal and chemotactic transcripts (*CCL21, CXCL12,
VIM*) localized to adjacent zones (q-values = 3.18×10^{-99}, 7.21×10^{-24}, 4.84×10^{-20}) (**Extended Data
Fig. 5c-f**).
204
To compare gut-associated lymphoid structures with canonical germinal centers, we analyzed
anatomically paired epithelial (APE) lymphoid aggregates in the gastrointestinal tract. Similar to
lymph node GCs, intra-GC B cells in APE structures expressed *IGHG1* and *IGKC*, while
surrounding stromal regions were enriched for *APOE* and extracellular matrix genes (**Fig. 4c,
Extended Data Fig. 5g**). However, unlike the sharply zonated architecture of lymph nodes,
APE lymphoid structures were embedded within a more intermixed environment of epithelial
and mesenchymal cells (**Fig. 4d-e**), suggesting greater potential for immune-epithelial crosstalk.
conserved and tissue-specific transcriptomic gradients across GC and peri-GC regions in LN
versus APE, highlighting organ-specific spatial organization (**Extended Data Fig. 5h**).
214

Expanding our analysis across all SAHA gastrointestinal tissues—including appendix (APE),
ileum (ILE), colon (COL), and stomach (STO)—we profiled 43 patient samples comprising 948
fields of view (FOVs) and over 1.15 million spatially resolved cells (**Fig. 4g**). Integrative
clustering of transcriptomic profiles combined with spatial adjacency information identified three
major niche types: (1) organ-shared epithelial-immune interaction zones, (2) organ-specific
immune microenvironments within lymphoid structures, and (3) rare cell populations with distinct
spatial distributions (**Fig. 4h-k, Extended Data Fig. 6**). Shared epithelial-immune
neighborhoods, characterized by dense T- and B-cell infiltration adjacent to crypt structures,
were recurrently observed in both colon and ileum samples (**Fig. 4h-i, Extended Data Fig. 6a-**
d). In contrast, the stomach exhibited parietal-chief cell clusters unique to gastric glands, and
the appendix contained specialized adaptive immune hubs, reflecting tissue-specific adaptations
of immune-epithelial architecture (**Fig. 4j-k**).
227
To investigate the functional roles of these spatial niches, we leveraged the unique ability of
spatial data to profile the ligand-receptor interaction networks between shared and organ-
specific microenvironments. Conserved immune-epithelial niches were dominated by paracrine
signaling between compartments, whereas tissue-specific structures exhibited distinct adaptive
immune signaling profiles (**Extended Data Fig. 6e**). Complementary spatial variance analysis
revealed that genes involved in immune activation (e.g., *CD74*, *IGHM*), epithelial barrier integrity
(e.g., *PIGR*), and chemokine signaling (e.g., *CCL21*) were highly localized within discrete spatial
clusters (**Fig. 4l**). These findings suggest that spatial neighborhoods are defined not only by
their cellular composition but also by localized transcriptional programs. Collectively, these
analyses demonstrate that spatial information is essential for uncovering hierarchical tissue
organization, functionally specialized microenvironments, and organ-specific immune
architecture in the human gastrointestinal tract.
240
241
242   **3. Validation and integration of multimodal spatial data**
243
The detailed spatial mapping of gastrointestinal tissues highlights the power of subcellular-
resolution atlases to uncover structured microenvironments and immune-epithelial interactions.
To ensure the robustness and biological fidelity of these insights across platforms and tissues,
we systematically validated SAHA datasets through multimodal integration of spatial
transcriptomic and proteomic data; we performed cross-platform validation using orthogonal
spatial assays, including CosMx RNA and protein profiling, 10x Genomics Xenium, RNAscope,
GeoMx DSP, and histopathology-based annotations (**Supplementary Table 2**).
251
In particular, integration of CosMx RNA and protein data enabled single-cell-level validation of
biological consistency and facilitated deeper characterization of cell states and subtypes. Co-
embedding of 1000-plex CosMx RNA and 67-plex CosMx protein profiles preserved tissue
structure and cellular identities, confirming the robustness of multimodal spatial profiling (**Fig.**
5a, Extended Data Fig. 7a-b). Canonical markers such as CD45 (PTPRC), E-cadherin (CDH1),
CD3D, and CD20 (MS4A1) exhibited concordant expression across RNA and protein modalities
(**Fig. 5b-d**). Protein-based annotations aligned closely (Adjusted Rand Index, ARI = 0.2719)

with RNA-integrated labels using Maxfuse [10], supporting consistent cell type classification
across modalities and batches (**Fig. 5c, Extended Data Fig. 7c-d**). Protein data also provided
clearer delineation of architectural features, such as the epithelial-stromal interface in the
stomach and lymphoid aggregates in the appendix (**Fig. 5d**), reflecting the stability, localization,
and abundance of structural proteins relative to transcriptomic signals [11]. Transcriptomic
profiles offered broader molecular coverage, enabling finer discrimination of epithelial subtypes
and detection of rare populations such as stem and transit-amplifying cells.
266
Beyond validating cell identity and tissue structure, protein datasets offer orthogonal
confirmation of receptor-ligand interactions inferred from RNA data (**Fig. 5e**) and provide direct
evidence of translational activity. While such multimodal comparisons reinforce biological
interpretation, it is important to note that RNA-protein correlations are not always linear.
Although this discordance has been widely hypothesized, systematic large-scale comparisons
across tissues have remained limited. Leveraging SAHA's multimodal resolution, we directly
compared matched RNA and protein measurements across multiple organs (**Fig. 5f**).
Housekeeping and structural genes such as VIM and FN1 showed strong concordance,
whereas key immune and signaling molecules—including RELA and IGHD—displayed
substantial discrepancies between RNA and protein levels (**Fig. 5f, Extended Data Fig. 7e-f**).
These differences were further highlighted in cross-organ analyses: VIM exhibited consistent
RNA-protein correlation across tissues (R^2 = 0.5781 with p-value = 0.0026 when compared
across all SAHA cells), whereas IL1B showed widespread RNA expression with minimal protein
detection. Conversely, CD276 demonstrated robust protein abundance despite low transcript
levels, consistent with post-transcriptional regulation [12]. These comparisons underscore the
necessity of multimodal profiling to accurately capture both transcriptional potential and protein-
level function.
284
The integration of histopathology-based annotations with spatial molecular profiles reinforced
biological interpretability and enabled precise mapping of anatomical features to molecular
states. Histological landmarks such as crypt architectures, lymphoid aggregates, and smooth
muscle layers corresponded closely to spatial clustering patterns observed in molecular
datasets (**Fig. 6**). This alignment between morphology and molecular identity validates the
technical robustness and biological fidelity of the SAHA datasets, supporting consistent cross-
tissue, cross-platform, and cross-modality analyses at subcellular resolution.
292
To further harness histological information, we applied a multi-modal vision-text foundation
model [13] that decodes tissue architecture from hematoxylin and eosin (H&E) stained whole-
slide images (**Fig. 6a-b**). The model extracts quantitative morphological features from image
tiles and learns spatial embeddings, which preserve anatomical continuity and discriminate
between distinct structural zones (See **Methods**, **Fig. 6c-d**). Unsupervised clustering of these
image-derived features reconstructed tissue organization, including epithelial boundaries,
stromal interfaces, and muscular layers (**Fig. 6e**). Visualization of representative clusters using
UMAP revealed discrete and biologically meaningful groupings that corresponded to
histologically distinct morphotypes, such as crypt bases, glandular epithelium, and lymphoid-rich

regions most of which aligned with unbiased, RNA-based spatial clustering results (**Fig. 4g-k,**
6f).

This approach generalizes across gastrointestinal tissues, as image-derived embeddings from
the colon, ileum, and appendix consistently show conserved morphological classes, including
crypt epithelium, extracellular matrix, and immune niches (**Fig. 6g**). Morphological clusters
overlapped with specific histopathological terms scored by the multi-modal vision-text model
(**Fig. 6h**) and displayed enrichment for specific tissue compartments such as epithelial,
connective, or immune components (**Fig. 6i**). Spatial adjacency patterns among clusters (**Fig.**
6j) and their correlations with RNA-derived cell type proportions (**Fig. 6k**) further confirmed that
image-derived features captured biologically relevant gradients in tissue organization. Finally,
direct comparison of H&E image and CosMx RNA projection revealed highly consistent
alignment shown by positive CCA correlation, between image-based morphology and
transcriptomically defined cell types (**Fig. 6l**), illustrating the power of histology-informed
molecular modeling for spatial atlas construction.

4. SAHA's utility and impact in biomedical research
A key motivation for constructing SAHA was to enable comparative spatial analysis of diseased
tissues, particularly in contexts where matched healthy controls are unavailable. To
demonstrate its translational utility, we applied SAHA to two clinically relevant diseased
conditions: colorectal cancer (CRC) and inflammatory bowel disease (IBD).

In CRC, we analyzed transcriptional and structural profiles in tumor adjacent crypt regions and
compared healthy tissue (SAHA COL, $n = 3$) to histologically normal crypts from CRC
resections ($n = 1$) (**Fig. 7a**). Crypt regions were stratified and annotated as 'top' and 'bottom'
crypts, based on gene expressions and their locations (**Fig. 7b, Extended Data Fig. 8a-b**).
Bottom crypts were identified by enrichment of genes associated with intestinal stem cells
(*OLFM4*) and differentiated secretory cells (*LEFTY1, STMN1*), whereas top regions expressed
mature epithelial markers (e.g., *KRT20, PLAC8, CEACAM1*). Notably, crypts adjacent to CRC
exhibited disruption in spatial expression patterns; markers normally restricted to bottom regions
(i.e., *H2AZ1, SPINK1*) were aberrantly expressed in the upper crypts, and vice versa for
CEACAM6 and *CEACAM1* (**Fig. 7c, Supplementary Table 3**). These patterns potentially
indicate crypt dedifferentiation and spatial disorganization often associated with tumor formation
[14], [15].

Tumor-adjacent crypts also harbored a distinct transcriptional program with genes significantly
more differentially expressed than in healthy crypts (**Fig. 7c**). Several upregulated genes,
including *REG1A*, have been previously associated with poor prognosis and crypt-inflammation
in CRC, while upregulation of *IL6* and *IL22* receptors are known to promote inflammation and
tumor development in CRC [16], [17]. CRC-associated crypts had a substantially higher
number of differentially expressed genes (DEGs) compared to COL crypts, indicating broader
transcriptional reprogramming (i.e., 96 vs 61 DEGs in bottom crypt and 116 vs. 72 DEGs in top
crypt; **Supplementary Table 3**). In terms of the tumor microenvironment, FOVs profiled from

CRC samples showed a significant increase in the number of mast cells (**Extended Fig. 9c**), a
known driver of angiogenesis and metastasis [18].
348
To further investigate spatial expression and organization of RNA transcripts, we performed
spatial autocorrelation analysis using Moran's I and clustered FOVs based on their top 200
spatially variable genes (SVGs) (**Fig. 7d**). This unsupervised approach effectively distinguished
crypts from CRC and COL samples, indicating their divergent spatial states. From matched top
and bottom crypt compartments across conditions, we visualized all SVGs and identified both
conserved spatial markers (e.g., *KRT20, CEACAM1, CEACAM6*) and spatially enriched
disease-associated genes (e.g., *REG1A, IL22RA1*) - both of which were also noted in our
differential expression analysis (**Fig. 7e-f, Extended Data Fig. 8e**).
357
Beyond structural comparisons, SAHA enabled exploration of intrapatient tumor heterogeneity
(**Fig. 7g-j**), including rare events such as perineural invasion (PNI) (**Fig. 7k-m**). Within a single
CRC sample, we identified two spatially and transcriptionally distinct tumor subtypes (labelled
as Type 1 and Type 2), based on dominant cell compositions and immunogenicity (**Fig. 7g**).
Type 1 tumor regions exhibited a higher lymphocyte to tumor ratio with increased proportions of
cytotoxic and naive CD8+ T cells, whereas Type 2 regions were enriched for macrophage
subsets (**Fig. 7g-j, Extended Data Fig. 8f-g**).
365
Among the analyzed FOVs, we captured a rare perineural invasion event, characterized by
dysplastic stroma surrounding a nerve and tumor cells encircling the perineurium (**Fig. 7k**).
Although perineural invasion is recognized as a hallmark of aggressive CRC, it remains difficult
to resolve with conventional histopathology and immunofluorescence, which lack
comprehensive transcriptomic resolution. While recent single-cell RNA-seq studies have
attempted to profile these events, they often fail to capture neural cells altogether—or, if
captured, cannot confidently localize them or distinguish whether they are part of a PNI
structure or simply present in unrelated regions of the tissue [19]–[21]. By leveraging the spatial
context of SAHA, we subclustered this PNI region and uncovered a rare subset of fibroblasts
(less than 100 cells in the PNI region) in close proximity to glial cells, characterized by a hybrid
transcriptional signature expressing both glial-associated genes (*CLU, NGFR, SLC2A1, APOD*)
and cancer-associated fibroblast markers (<2% of fibroblasts identified in the sample) such as
CXCL14 and *MMP2* (**Fig. 7k-l, Extended Data Fig. 8h**). Cell-cell communication analysis
further revealed unique signaling pathways within the PNI niche, including the ANGPTL
pathway originating from fibroblasts, previously associated with increased metastatic risk in
CRC [22]–[24], and the LIFR pathway, exclusively signaling from tumor cells to glial cells,
known to facilitate tumor-fibroblast crosstalk and activate pro-invasive programs [25]. These
findings offer new insights into the interactive dynamics within this rare microenvironment and
its ability to specify high resolution details of rare tumor types.
385
We also compared SAHA ILE with ileal samples from patients with inflammatory bowel disease
(IBD), including individuals stratified by clinical response to TNFα inhibitor therapy (**Fig. 8a**).
Integration of CosMx RNA profiles across healthy and IBD ileum samples revealed clear
separation by disease state, while preserving major cell type identities (**Fig. 8b, Extended Data**

Fig. 9a-b). Correlation analysis of average gene expression profiles demonstrated strong
consistency (e.g., spearman correlation of 0.74 for epithelial cells from 1K and 6K datasets)
across datasets generated with different plex sizes, validating the reproducibility of cell-type
transcriptional programs (**Extended Data Fig. 9a**). Spatial mapping of cell types demonstrated
increased immune cell infiltration into epithelial compartments in IBD samples relative to healthy
controls (**Fig. 8c, Extended Data Fig. 9c-d**).
396
Quantification of epithelial-proximal immune infiltration within 300 μm confirmed a significant
increase in immune cell density in IBD tissues ($p < 0.001$; **Fig. 8d**). Ligand–receptor interaction
analyses further revealed enhanced immune-epithelial signaling in IBD, particularly among non-
responders to TNFα therapy, who exhibited elevated spatial interaction scores relative to
responders (**Fig. 8e**). Mesenchymal-stromal populations in IBD tissues showed increased
expression of pro-inflammatory and extracellular matrix remodeling markers, including elevated
TNFRSF1A (TNF receptor 1) expression among non-responders compared to responders ($p <$
0.05; **Fig. 8f**). Spatial mapping of TNF-TNFRSF1A ligand–receptor interactions localized active
signaling to immune-mesenchymal interfaces, particularly in non-responder tissues (**Fig. 8g**).
406
These molecular level changes were consistent with the histological analyses. As performed for
the healthy GI tract (**Fig. 6**), we applied a multi-modal text-vision model to quantify and describe
the morphological patterns of H&E-stained ileum sections of healthy and IBD tissues (**Fig. 8h**).
IBD samples were distinguishable from healthy, with histopathological terms related to
metaplasia, immune cell infiltration, crypt branching, and apoptosis more frequent in IBD
samples compared to controls (**Fig. 8i**). Furthermore, we also found differentially enriched
morphological changes in response to TNFα treatment (**Fig. 8i**), which were aligned with
molecular findings, reinforcing the multi-layered disruption of epithelial and stromal architecture
in IBD [26]. These analyses demonstrate how SAHA enables spatially resolved comparisons of
healthy and diseased tissues at both molecular and histological levels, providing critical insights
into disease-associated microenvironmental changes and therapeutic response heterogeneity.
418
The integration of spatial transcriptomics and proteomics extends the analytical frameworks
traditionally applied to single-cell sequencing by adding critical layers of structural and
contextual information (**Fig. 9a**). Spatial multi-omics enables not only transcriptomic and
proteomic profiling at subcellular resolution, but also the characterization of extracellular matrix
(ECM) features, direct cell-cell interactions, and subcellular biomolecule localization, which are
dimensions inaccessible to dissociated single-cell methods. The addition of spatial context
enables direct measurement of cellular phenotypes, resolving the challenge of drawing
conclusions from disconnected data. This multidimensional mapping redefines efforts in building
biological atlases and opens new avenues for understanding tissue organization and function.
428
Recent technological advances now enable high-throughput, same-cell spatial multi-omics at
subcellular resolution—capturing whole-transcriptome RNA (~19,000 genes), high-plex protein
expression (~67 markers), and histological context within a single assay (**Fig. 9a**). These spatial
platforms also provide cell morphology, boundary, and segmentation data that are inaccessible
via traditional dissociated single-cell methods. To scale this approach across large tissue

volumes and billions of cells, we implemented computational workflows capable of integrating
transcriptomic, proteomic, and morphological data at population scale. Compared to
conventional scRNA-seq, spatial multi-omics approaches using Xenium and CosMx offer
orders-of-magnitude lower cost per cell and support efficient scaling to billions of cells with
fewer experimental runs—facilitating population-scale spatial atlasing (**Fig. 9b**).
439
To maximize accessibility and impact, SAHA adheres to the FAIR (Findable, Accessible,
Interoperable, and Reusable) data principles. The SAHA data portal provides an open-access
platform for exploring spatial multi-omics datasets, equipped with interactive visualization tools,
high-resolution tissue navigation, and download capabilities for downstream analysis (**Fig. 9c**).
Beyond serving as a reference, SAHA establishes standardized protocols and integration
frameworks that can be adopted across institutions to enable scalable, reproducible spatial
profiling.
447
448
DISCUSSION
The Spatial Atlas of Human Anatomy (SAHA) establishes the first multi-organ, multimodal, and
multi-platform reference map of healthy human gastrointestinal and immune systems at
subcellular resolution. By integrating histological, proteomic, and transcriptomic data across
tissues and spatial technologies, SAHA addresses major limitations of previous atlasing efforts,
which have predominantly relied on dissociated single-cell data and lacked spatial context. A
key strength of SAHA lies in its rigorously standardized experimental and computational
pipelines, which ensure reproducible, high-quality data generation and facilitate integrative
analyses across organs and modalities. To support cross-platform consistency, we implemented
systematic validation across CosMx, Xenium, GeoMx, and RNAscope platforms, alongside
robust quality control metrics. While SAHA initially leveraged 1000-plex RNA profiling, we also
introduce the field's first whole-transcriptome and paired protein multi-omics datasets at
subcellular resolution—expanding beyond mid-plex spatial transcriptomics to unlock deeper
biological insights and establish a future-ready reference for spatial biology.
463
A central innovation of SAHA lies in its multimodal design. The integration of RNA, protein, and
histopathological (H&E) data facilitates both cross-validation of molecular signals and nuanced
dissection of tissue structure and function. This multimodal synergy enabled identification of
conserved spatial niches, lineage-specific expression gradients, and regionally restricted cell
states, including rare and adaptive immune populations such as follicular B cells and
intraepithelial lymphocytes. We also uncovered fine-grained epithelial and stromal subtypes,
such as tuft and enterochromaffin-like cells, and delineated spatially encoded state transitions
within common lineages—patterns that would be obscured in single-modality or dissociated
datasets. Some segmentation ambiguity is inevitable in a subset of cells, in part due to the
physical limitations of tissue sectioning through three-dimensional structures. However, we
argue that such instances also encode meaningful biological information, as spatial proximity—
retained in intact tissue—offers a critical dimension absent in single-cell suspensions.
Dissociation-based workflows may also lose spatially tethered subpopulations during washing
and library preparation steps. While future pixel-level or nucleus-aware segmentation methods

may refine these boundaries, our current analysis emphasizes broader tissue-level organization,
highlighting reproducible spatial relationships that extend beyond cell-intrinsic expression.
480
Across gastrointestinal tissues—including appendix, ileum, colon, and stomach—SAHA
captures a conserved layered organization spanning mucosa, submucosa, muscularis externa,
and serosa. At the same time, it reveals organ-specific features, such as crypt-villus gradients,
lymphoid aggregates (e.g., Peyer's patches), and vascular-rich stromal zones. These features
were consistent across individuals and histologically validated, reinforcing the atlas's technical
fidelity. Notably, cross-organ analysis within gastrointestinal tissues uncovered conserved
principles of mucosal organization while highlighting unique adaptations in the appendix, ileum,
colon, and stomach. A comparative profiling of lymphoid aggregates across GI organs and
matched germinal centers in lymph nodes revealed both preserved core immune architectures
and distinct spatial intermingling with epithelial and stromal compartments, emphasizing the
utility of SAHA in decoding tissue-specific immunological microenvironments.
492
Importantly, SAHA serves not only as a reference but also as an actionable spatial comparator
for disease studies—especially in clinical contexts where matched healthy tissue is unavailable.
We demonstrate this translational potential by benchmarking SAHA against datasets from
colorectal cancer (CRC) and inflammatory bowel disease (IBD). This revealed distinct
alterations in crypt top–bottom zonation, immune composition, and spatial gene expression
programs. For instance, spatial transcriptomics exposed crypt dedifferentiation in CRC and
aberrant cytokine signaling, while perineural invasion—an aggressive metastatic route—was
resolved at high granularity, highlighting glia-associated fibroblast states and tumor-nerve
signaling circuits. These examples illustrate how SAHA enables spatial biomarker discovery and
mechanistic insight into rare or complex pathological events, areas that have been briefly
explored in previous studies.
504
While SAHA represents a significant advance, several limitations remain. First, despite including
over 100 donors, our cohort size is modest relative to the full scope of human biological
diversity. Expanding SAHA across age groups, ancestries, and environmental exposures is
essential for broader generalizability. Second, although cross-platform integration across
CosMx, Xenium, GeoMx, and RNAscope demonstrates strong concordance, inherent technical
differences—such as probe capture efficiencies and antibody variability—necessitate ongoing
methodological harmonization. Third, this initial phase focuses on gastrointestinal and lymphoid
organs; a complete human reference will require comprehensive coverage across additional
anatomical systems.
514
While the current datasets represent the initial phases of spatial atlas construction, the platform
is designed for scalability and community-driven expansion. Active efforts are underway to
expand its organ coverage (second phase to genitourinary systems and additional immune
organs), apply whole-transcriptome and high-plex proteomics protocols, increase demographic
representation, and integrate advanced computational frameworks, including deep learning for
tissue segmentation and spatial graph modeling. We also aim to leverage Common Coordinate
Framework (CCF)-based registration and analysis pipelines to enable 3D spatial integration

across tissue sections and donors. This will support more anatomically precise mapping,
facilitate multi-plane reconstruction of tissue architectures, and allow in-depth cross-sample
comparisons at scale.
525
By integrating spatial molecular data with histological and clinical metadata, SAHA provides the
infrastructure for building digital twins of human tissues—enabling in silico simulations of
disease progression, therapeutic response, and tissue regeneration. These capabilities position
SAHA as a translational bridge between spatial biology and precision medicine, advancing both
fundamental research and clinical diagnostics. For example, we can extend SAHA toward
perturbation-aware modeling, using spatial benchmarks to assess disease-associated changes
and therapeutic responses—analogous to recent large-scale single-cell initiatives such as
Tahoe-100M [27]. Together, these directions will transform SAHA into a foundational tool for
spatially resolved systems biology and experimental medicine.
535
The SAHA data portal, built on FAIR (Findable, Accessible, Interoperable, Reusable) principles,
will enable broad community access and invite collaborative expansion. We envision that SAHA
will not only support basic biological discovery and accelerate clinical translation by providing a
standardized framework for the development of spatially informed diagnostics, prognostics, and
therapies. As spatial omics technologies continue to evolve, SAHA serves as a dynamic
foundation to integrate emerging data types, drive systems-level insights into tissue function and
disease, and chart new frontiers in precision medicine. We invite researchers worldwide to
utilize SAHA datasets, develop new analytical tools, and contribute additional spatial multi-
omics data to collectively accelerate biological discovery and the construction of a
comprehensive human spatial atlas.
546
547
METHODS
Patient material, ethics approval, and consent for publication
Human tissue samples were obtained from archived formalin-fixed, paraffin-embedded (FFPE)
blocks curated by board-certified pathologists and collaborating investigators (**Supplementary
Table 1**). Each tissue sample was selected from a diverse ancestry (if possible) from healthy
adults (no cancer or organ wide immune issues, ideal age 20-40). If tissues were difficult to get,
age and ancestry parameters can be compromised, but no cancer patients or patients with
diseases that will cause an impact on not diseased tissues was considered. Representative
fields of view (FOVs) were selected following comprehensive histopathological evaluation of
hematoxylin and eosin (H&E)-stained slides. Tissue acquisition and research use were
conducted under protocols approved by the Institutional Review Board (IRB) at Weill Cornell
Medicine, specifically IRB protocols 1306014012 (Hissong), 1008011210 (Robinson), and
107004999 (Inghirami). All experiments adhered to ethical guidelines, and consent for research
use of de-identified samples was obtained in accordance with IRB regulations.
Tissue microarray (TMA) formalin-fixed, paraffin-embedded (FFPE) blocks were constructed
following standard procedures. SAHA FFPE blocks were selected based on histopathological
evaluation, and representative areas were identified on hematoxylin and eosin (H&E)-stained
slides. Tissue cores (variable in diameter depending on the organ) were extracted using a

manual or semi-automated tissue arrayer and transferred into a pre-designed recipient paraffin
block. The TMA block was re-embedded and cooled to ensure structural integrity. SAHA FFPE
blocks were stored in cardboard boxes at room temperature (RT) under ambient humidity until
the day of sectioning.
570
Tissue processing and histology
The FFPE blocks were sectioned at a thickness of 5 µm, and the sections were placed within
Bruker's 15mm x 20mm allowed scan area on Leica Bond Plus slides (Leica Biosystems,
S21.2113.A). Fresh ultrapure water was used to fill the water bath and new microtome blades
were used for each tissue type. The first few sections from the block face were discarded. The
number of serial sections to be collected was determined in advance according to the planned
assays. Slides were labeled with a serial number to keep track of the order, ensuring that
multiple assays were run on adjacent sections and facilitate subsequent data integration. If
needed, one section, usually from the middle of the series, was dedicated to H&E staining,
followed by full-slide image acquisition using the Aperio Digital Pathology Slide Scanner system
(Leica Biosystems), allowing pathologists to identify and mark the area of interest in
consideration of the available number of CosMx fields of view (FOVs). Most sections were also
H&E stained after data acquisition. **Supplementary Table 2** lists the serial section numbers for
each assay and for the corresponding H&E for each analyzed organ. Following sectioning, the
slides were dried upright at RT for 30 minutes (min), followed by a horizontal overnight (ON)
drying step at 37°C in an oven to improve tissue adherence. The slides were then stored in a
desiccator at RT until processing. Most slides dedicated to the RNA assay were used within two
weeks, as recommended by Bruker. Exact sectioning and run dates are reported in
Supplementary Table 2.
590
CosMx processing
Processing of the slides followed the guidelines outlined in the Bruker manuals MAN-10184 for
the RNA assay and MAN-10185 for the Protein assay, utilizing the equipment, materials, and
reagents recommended in the manuals. Protocol details may have varied over time, reflecting
updated manuals throughout the SAHA project. The specific manual version used for each
assay is reported in **Supplementary Table 2**.
For the CosMx RNA assay, slides were baked at 60°C ON in vertical position in a baking oven.
The following day, CitriSolv (VWR, 89426-268) was used for deparaffinization. Specifically,
slides were brought to RT for 3 min, then subjected to two sequential 5 min immersions in
CitriSolv, followed by two 5 min washes in 100% ethanol (Et-OH) (Sigma, E7023). The slides
were dried vertically for 5 min at 60°C in the same baking oven. Antigen retrieval was conducted
in a BioSB® TintoRetriever digital pressure cooker (BioSB, BSB 7015) at 100°C for 15 minutes.
After the 5-minute drying step, slides were transferred into the preheated 1X Target Retrieval
Solution (part of Bruker 121500006) as per the Bruker manual. The pressure cooker was then
closed, and the temperature was allowed to return to 100°C. At this point, a 15 min timer was
started, after which the slides were quickly rinsed up and down in DEPC water (Fisher Scientific,
BP561-1) at RT for 15 seconds, washed for 3 min in 100% Et-OH, and then dried for 30 min
horizontally on a clean surface at RT. 10 min before the end of the drying step, the incubation
frame (part of Bruker 121500006) was applied to the slide and a 3 µg/mL Proteinase K (part of
Bruker 121500006) working solution in 1XPBS (ThermoFisher Scientific, AM9625) was applied
to the sections. The slides were then incubated for 30 min at 40°C in a RapidFISH slide

hybridizer oven (Boekel Scientific, 240200). During the final 10 min of digestion, Bruker fiducial
beads (part of Bruker 121500006) were resuspended through alternating 1-min vortexing and 2-
min sonication steps (three vortexing steps and two sonication steps in total), then diluted in 2X
SCC-T (part of Bruker 121500006). Fiducials were diluted at a final working concentration of
0.001% for Appendix, Ileum, Colon, and Pancreas samples, 0.00075% for Stomach and Lymph
Nodes samples, and 0.0005% for inflammatory bowel disease ileum samples. Following
digestion, slides were washed twice in DEPC water per the manual and then incubated with the
prepared fiducials for 5 min, protected from light at RT. The slides were then washed in 1XPBS
for 1 min, fixed in 10% Neutral Buffered Formalin (NBF) (EMS Diasum, 15740) for 1 min,
washed twice in a 0.1M Tris-Glycine NBF buffer solution for 5 min, once in 1XPBS for 5 min,
and incubated with a freshly prepared 100 mM NHS-Acetate (Fisher Scientific, 26777) solution
in NHS-Acetate buffer (part of Bruker 121500006) for 15 min in the dark at RT. Finally, the
slides were washed twice in 2X SSC (ThermoFisher Scientific, AM9763) for 5 min. At this point,
slides were stored in 2X SSC, protected from light, until it was possible to proceed with the ON
hybridization step. The required amount of probe for the run was transferred to 0.2 mL tubes,
denatured for 2 min at 95°C in a standard thermocycler, and crash-cooled on ice for 1 min. A
probe hybridization mix containing RNase inhibitors (Bruker, 121500004) and Buffer R (part of
Bruker 121500006) was prepared according to the Bruker manual. Slides were laid horizontally
on a clean surface and the hybridization mix was applied to the sections. A plastic coverslip
(part of Bruker 121500006) was then placed over the frame. The slides were incubated ON for
17-18 hours (h) at 37°C in the RapidFISH slide hybridizer oven. The RNA probe panel used for
the SAHA project was the CosMx™ Human Universal Cell Characterization Panel (Bruker,
121500002), except for inflammatory bowel disease ileum samples, which were run using the
CosMx™ Human 6K Discovery Panel (Bruker, 121500041). On the following day, after
removing the coverslip, slides were quickly rinsed in 2X SCC, washed twice for 25 min at 37°C
in pre-warmed 50% 4X SSC/50% formamide (ThermoFisher, AM9342), and washed again in 2X
SCC twice for 2 min. The slides were then incubated in the dark at RT for 15 min with a 1:40
dilution of DAPI (part of Bruker, 121500020) in blocking buffer (part of Bruker, 121500006). After
a 5-min wash in 1XPBS, cell segmentation marker staining was performed using a cocktail of
CD298/B2M, PanCK, CD45, and CD68 antibodies. The antibodies are conjugated to a readout
domain that binds fluorophores on-instrument. Slides were incubated in the dark at RT for 1 h
with the morphology staining marker mix, prepared using the CosMx™ Human Universal Cell
Segmentation Kit, RNA (Bruker, 121500020) and the CosMx™ Human CD68 A La Carte
Marker, RNA (Bruker, 121500022), at the dilutions specified in the Bruker manual. Slides were
then washed three times in 1X PBS for 5 min and transferred to 2X SSC. For most samples,
the protocol spanned three days, meaning that at this stage, the adhesive frame was removed,
and slides were stored ON in 2X SSC at 4°C, protected from light. The only exceptions were
stomach and lymph node samples, which were fully processed and loaded on the instrument in
just two days. On the following day, the flow cells were assembled using the Flow Cell
Assembly Tool provided by Bruker and Bruker glass coverslips with ports (CosMx™ Flow Cell
Assembly Kit, Bruker, 122000061). They were then kept in 2X SSC, ensuring the solution filled
the flow cells to prevent section dehydration, and stored in the dark until loading onto the CosMx
instrument.
For the CosMx Protein assay, slides were baked for just 1 h at 65°C in a vertical position.
Similar to the RNA assay, after baking slides were equilibrated for 3 min at RT and then
immersed in CitriSolv twice for 5 minutes. In the case of the protein workflow, this was followed
by a series of washes: two 10 min washes in 100% Et-OH, two 5 min washes in 95% EtOH, one
5 min wash in 70% EtOH, and two 5 min washes in 1XPBS. The target retrieval step was

performed as described above for the RNA assay, with the only difference that after the 15 min retrieval, the solution containing the slides was left to cool at RT for 25 min. The slides were then transferred to 1XPBS and washed three times for 5 min. At this point, the incubation frame (part of Bruker 121500008) was applied to the slides. Sections were incubated with blocking Buffer W (part of Bruker 121500008) for 1 h at RT, followed by ON incubation at 4°C with a single mix containing both the morphology marker antibodies for segmentation (CosMx Human Universal Cell Segmentation Kit, IO PanCK/CD45 Supplemental Segmentation Kit, and CD3 A La Carte Marker, Bruker 121500026, 121500027, and 121500028) and the antibodies from the CosMx Human Immuno-oncology 64-plex panel (Bruker, 121500010). The sections were covered with a plastic coverslip (part of Bruker 121500008) during ON incubation. The following day, after removing the coverslip, the slides were washed three times for 10 min in 1XTBS-T (ThermoFisher, J77500.K2) and once for 2 min in 1XPBS. During the 1XTBS-T washes, fiducials were prepared as for the RNA assay. For the protein assay, the final fiducial working concentration was 0.00005%, requiring a two-step serial dilution in 1XTBS-T, as recommended by the manual. After a 5 min incubation at RT with fiducials, slides were washed once with 1XPBS for 5 min, fixed with 4% PFA (Electron Microscopy Sciences, 15710) for 15 min at RT in the dark, washed three times in 1XPBS for 5 min each, and incubated for 10 min at RT with a 1:40 dilution of DAPI (Bruker 121500026) in 1XPBS. The slides were then washed twice more in 1XPBS for 5 min, incubated with freshly prepared 100 mM NHS-Acetate for 15 min protected from light at RT, and subjected to a final 5 min wash in 1XPBS. As for the RNA assay, for most of the SAHA protein assays, the protocol extended over three days. Slides were transferred to fresh 1XPBS tubes and stored at 4°C in the dark ON. On the following day, flow cells were assembled as described above, kept in 1XPBS in the dark, and loaded onto the CosMx instrument.

To set up a new CosMx runs on the Bruker CosMx Spatial Molecular Imager (SMI), the Bruker manual MAN-10161 was used as a guide. Different versions of the manual were followed throughout the SAHA project, and various SMI instrument software versions were installed on the machine over time. **Supplementary Table 2** lists the instrument manual and software versions for each analyzed sample. The SMI Control Center webpage interface allows for the setup of all run specifications, as well as the assignment of FOVs. The setup begins with configuring a new flow cell, the first step of which is defining the primary information for the run, including the Pre-Bleaching Profile and the Cell Segmentation Profile. As these profiles are tissue-type specific, they were selected based on Bruker's recommendations in MAN-10161, but they ultimately ended up being Pre-Bleaching Profile C and Cell Segmentation Profile A for all our runs, both RNA and protein. The probe panel and cell segmentation kits used also needed to be specified at this stage. The sensor-based interactive SMI Control Center interface guided the loading of the instrument. The Imaging Tray (CosMx RNA Imaging Tray, Bruker 122000156, 122000157, or 122000158, CosMx Protein Imaging Tray, Bruker 122000162 or 122000163), flow cells, and buffers required for the run (CosMx RNA Instrument Buffer Kit, Bruker 100480 or 100481, CosMx Protein Instrument Buffer Kit, Bruker 100482) were loaded into their designated slots. For RNA assays, RNase inhibitors (Bruker, 121500004) were added to a specific well of the Imaging Tray. For all runs, two enzymes - Catalase and Pyranose Oxidase (part of Bruker 100480, 100481, and 100482) - needed to be resuspended, centrifuged, and added free of precipitate to Buffer 4 in advance. Once the assembled flow cell were loaded onto the instrument and all run parameters were approved, the system proceeded through a deck validation step, an instrument preparation step, and a preliminary morphology marker scan to allow the FOVs placement. RNA target readout on the CosMx SMI instrument was performed as described in He et al. [28]. Reporter Wash, Imaging, and Strip Wash Buffers all supplied by

Bruker Spatial Biology. At this point, the FOV Selection Workspace opened, and the FOV
Placement Tool enabled for the placement, movement, or deletion of individual FOVs or grids of
FOVs over the scanned image of the section. For large tissue areas, a serial H&E scan was
used in advance by pathologists to annotate the image and define the regions of interest to be
covered by the available FOVs. For TMAs, we aimed to allocate a predefined number of FOVs
to individual tissue cores. Once the FOVs were reviewed and approved, the machine began
cycling. RNA readout began by flowing 100 µl of Reporter Pool 1 into the flow cell and
incubating for 15 min. Reporter Wash Buffer (1 mL) was flowed to wash unbound reporter
probes, and Imaging Buffer was added to the flow cell for imaging. Eight Z-stack images (0.8
µm step size) for each FOV were acquired, and photocleavable linkers on the fluorophores of
the reporter probes were released by UV illumination and washed with Strip Wash buffer. The
fluidic and imaging procedure was repeated for all the reporter pools, and the different rounds of
reporter hybridization (based on assay plex) imaging were repeated multiple times to increase
target detection sensitivity. Cell morphology was imaged on-instrument prior to RNA or protein
readout by adding fluorophore-bound reporters to the flow cell and capturing eight z-stack
images in the channels 488 nm (CD298/B2M), 532 nm (PanCK), 594 nm (CD45), 647 nm
(CD68), and 385 nm (DAPI). Runs lasted anywhere from two to ten days, depending on the
panel type and plex, and the number of FOVs. CosMx data were automatically uploaded to
Bruker's cloud-based AtoMx Spatial Analysis Platform both during and after the run.
727
Single-nuclei sequencing (snPATHO-seq)
Single nucleus RNA sequencing (snRNA-seq) was performed following the snPATHO-Seq
protocol as previously described [29]. Two scrolls of 30–50 µm thickness were obtained from
each FFPE sample, adjacent to the sections used for CosMx, Xenium, and H&E staining.
Scrolls were collected into 1.5-mL Eppendorf tubes, stored at 4°C in Ziplock bags containing
desiccants to control humidity, and processed within one week. For nuclei isolation, scrolls were
dewaxed by three sequential washes in 1 mL of xylene (Millipore Sigma, 214736) for 10 minutes
each, with the first xylene wash performed at 55°C to enhance paraffin removal. Rehydration
was achieved through serial immersions in decreasing concentrations of ethanol (100% twice,
followed by 70%, 50%, and 30%; each for 1 minute) (Millipore Sigma, E7023), followed by a
single wash with 1 mL of RPMI 1640 medium (Gibco, 11875093).
Tissues were then digested in RPMI 1640 supplemented with 1 mg/mL Liberase TH (Millipore
Sigma, 5401151001) and 1 U/µL RNase inhibitor (Thermo Fisher Scientific, EO0382). The
scrolls were fragmented using a pellet pestle, and the digestion volume was adjusted to 1 mL.
Digestion proceeded at 800 rpm in a Thermomixer. Following digestion, 400 µL of EZ Lysis
Buffer (Sigma-Aldrich, NUC-101) was added, mixed, and centrifuged at 850 × g for 5 minutes at
4°C. The nuclei-containing pellet was resuspended in EZ Lysis Buffer with 2% BSA (Miltenyi,
130-091-376) and RNase inhibitor, then homogenized by pestle and pipetting. The suspension
was filtered through a 70-µm strainer (pluriSelect, 43-10070-50) and centrifuged. The nuclei
pellet was washed three times with 1× PBS containing 1% BSA (Gibco, 70011044), followed by
two washes with 0.5× PBS containing 0.02% BSA. Final resuspension was performed in 500–
1000 µL of 0.5× PBS with 0.02% BSA, depending on pellet size, and filtered through a 40-µm
strainer (pluriSelect, 43-10040-50).
Nuclei were quantified using the Luna-FX7 instrument with Acridine Orange/Propidium Iodide
staining (Logos). For immediate processing, nuclei suspensions were kept on wet ice. snRNA-
seq libraries were prepared using the Chromium X Controller (10x Genomics) and the

Chromium Fixed RNA Profiling Reagent Kits for multiplexed samples (Chromium Fixed RNA Kit,
PN-1000475; Chromium Next GEM Chip Q, PN-1000418; Dual Index Kit TS Set A, PN-
1000251), following the manufacturer's user guide (CG000527, Rev E). For cryopreservation,
nuclei suspensions were supplemented with 10% glycerol and 0.1× Enhancer (10x Genomics,
PN-2000482), rested on ice for 10 minutes, and stored at -80°C.
Hybridization was performed using 500,000–1.5 million nuclei per sample at 42°C for 20 hours.
Barcoding reagents BC1–BC4 were used. Post-hybridization washes were conducted in a
pooled setting according to the user guide with an additional wash step. Approximately 10,000
nuclei were targeted for capture per sample. An extra cycle was added during the pre-
amplification PCR step, and indexing PCR was optimized to achieve final library concentrations
between 50 and 200 nM. Library quality was assessed using a Fragment Analyzer 5200
(Agilent) and quantified using a Qubit Flex Fluorometer (Thermo Fisher Scientific). Libraries
were sequenced on an Illumina NovaSeq 6000 using a 28 × 10 × 10 × 90 bp read configuration,
targeting 20,000–30,000 read pairs per nucleus.
Transcript count matrices generated by Cell Ranger v8.0.0 (10x Genomics) were imported into
Seurat v5.0.1 for downstream analysis [30]–[34]. Doublet identification was performed using
scDblFinder v1.14.0 with a preset doublet rate of 10%, reflecting the expected frequency in
Single Cell Gene Expression Flex multiplexing experiments [35]. Quality control based on UMI
and gene count distributions identified a population of low-complexity droplets, which lacked
distinct transcriptomic signatures and were excluded from further analysis. Nuclei flagged as
doublets or containing fewer than 1,000 detected genes were removed, ensuring retention of
major cell populations as validated by comparison with the original dataset. Normalization and
highly variable feature selection were conducted using SCTransform with default parameters.
Dimensionality reduction was performed using principal component analysis (PCA), with the first
30 principal components guiding Uniform Manifold Approximation and Projection (UMAP)
embedding (RunUMAP) and Louvain clustering.
780
Xenium Slide Preparation and In Situ Gene Expression Assay
Sections for Xenium assays were prepared according to the *Xenium In Situ for FFPE—Tissue*
Preparation Guide (CG000578 Rev C, 10x Genomics) following the manufacturer's instructions.
Briefly, 5-µm-thick FFPE tissue sections were floated in an RNase-free water bath and mounted
onto the 10.45 mm × 22.45 mm sample area of Xenium slides (PN-3000941, 10x Genomics).
Slides were air-dried at room temperature for 30 minutes, followed by a 3-hour incubation at
42°C on a Xenium Thermocycler Adapter plate positioned atop a 96-well thermal cycler (C1000
Touch, Bio-Rad) with the lid open. Slides were further dried ON at room temperature in the
presence of desiccants.
Subsequent processing, including deparaffinization and decrosslinking, was performed following
the *Xenium In Situ for FFPE—Deparaffinization and Decrosslinking Protocol* (CG000580 Rev C,
10x Genomics). Slides were incubated at 60°C for 2 hours, equilibrated at room temperature,
then sequentially immersed in xylene (twice, 10 minutes each), 100% ethanol (twice, 3 minutes
each), 96% ethanol (twice, 3 minutes each), 70% ethanol (once, 3 minutes), and water (once,
20 seconds). Following rehydration, slides were assembled into Xenium cassettes and hydrated
with 1× PBS. Tissue sections were reverse crosslinked using a decrosslinking buffer containing

tissue enhancer, urea, and perm enzyme B at 80°C for 30 minutes, followed by multiple PBS-T
washes.
Slides were then processed according to the *Xenium In Situ Gene Expression User Guide*
(CG000582 Rev E, 10x Genomics). Gene expression profiling was performed using the Xenium
Human Multi-Tissue and Cancer Panel (10x-1000626), targeting 377 genes for prostate, ileum,
appendix, pancreas, and colon samples, and the Xenium Human Breast Panel (10x-1000463,)
targeting 380 genes for breast cancer samples. Probe hybridization involved probe denaturation
at 95°C for 2 minutes, crash-cooling on ice, equilibration to room temperature, and 17-hour
hybridization at 50°C. Post-hybridization, slides underwent standardized washes, ligation,
amplification, autofluorescence quenching, and nuclear staining following the manufacturer's
protocols.
Slides assembled into Xenium cassettes were processed using the Xenium Analyzer according
to the *Xenium Analyzer User Guide* (CG000584 Rev D/E, 10x Genomics). Instrument setup
included loading of the Xenium Decoding Consumables Kit (PN-1000487) with appropriate
wash buffers and reagents. Slides were scanned to image fluorescently labeled nuclei, and
fields of view (FOVs) were selected for sequencing. Data acquisition utilized the Xenium In Situ
software suite (v1.9 for standard samples; v2.0 for multimodal segmentation samples), with
simultaneous collection of gene expression and segmentation data for multimodal samples.
Post-run, slides were washed with PBS-T and stored at 4°C in the dark until further staining and
analysis.
817
RNAscope for quality assessment
Chromogenic in situ hybridization was performed on an automated stainer (Leica Bond RX,
Leica Biosystems, Deer Park, IL) with RNAscope 2.5 LS Assay Reagent Kit-Red (Cat. #
322150, Advanced Cell Diagnostics, Newark, CA) and Bond Polymer Refine Red Detection
(Cat. # DS9390, Leica Biosystems) following manufacturer's standard protocol. Positive control
probe detecting a housekeeping gene (human PPIB, cat # 313908, Advanced Cell Diagnostics)
and a negative control probe detecting bacterial (*Bacillus subtilis*) dapB gene (cat #312038,
Advanced Cell Diagnostics) were used to assess RNA preservation and detection, and absence
of non-specific signal, respectively. The chromogen was fast red and the counterstain
hematoxylin. Positive RNA hybridization was identified as discrete, punctate chromogenic red
dots under brightfield microscopy. Whole slide images were acquired with an Olympus VS200
slides scanner and VS200 ASW 4.1.1 software (Evident Scientific, Hamburg, Germany) using a
20X/0.80NA objective to generate whole slide images with a pixel size of 0.2738 μm. Image
analysis was performed with QuPath 0.5.1 software (Bankhead, P. et al. QuPath: Open source
software for digital pathology image analysis [36]. The positive cell detection and subcellular
detection algorithms were used to detect cells and hybridization dots (individual spots and
clusters), respectively. The mean number of estimated spots per cell was reported for each
sample. The analysis we performed and validated by a board-certified veterinary pathologist
(SM).
837
Data processing and annotation
For CosMx RNA and protein data, AtoMx Spatial Informatics Platform (SIP) was used for image
processing and preprocessing of raw data. A slight modification was applied to the foundational

pipeline available on the AtoMx platform (v1.3). For quality control, cells with the following

cutoffs were flagged as low-quality cells: total transcript counts less than 20, negative probe

counts more than 10%, count distribution 1, area outlier 0.01 (outlier p-value cutoff 0.01). FOV

QC was done with default settings (mean, FOV count cutoff 100). The negative control probe

quantile cutoff for target QC was set to 0.5, and detection was set to 0.01.

The datasets were normalized by scale factors and total counts, while 6k panels and combined

datasets were normalized using total count normalization for scaling. For organ-specific

analyses, count matrices were normalized using sc.pp.normalize_total() and log-transformed via

sc.pp.log1p() (Scanpy v1.9.3). Dimensionality reduction was performed using PCA, followed by

neighborhood graph construction (sc.pp.neighbors()) and UMAP embedding (sc.tl.umap()).

UMAP was generated with the following parameters: 50 PCs for clustering, 40 number of

neighbors, 0.01 minimum distance, 5 spread, Cosine distance metric, 0.25 data fraction. Spatial

networks were calculated based on the distance (50 um), and neighborhoods based on the

Jaccard cutoff of 0.099 and cosine distance. The datasets were exported for cell type annotation

and further downstream analysis. For initial automated cell type annotation of RNA data, we

used Insitutype [37] with organ-specific profile matrices derived from published single-cell and

spatial transcriptomics datasets (available via the SAHA GitHub repository; see **Code**

Availability). For protein data, cell phenotyping was performed using SCIMAP based on a

marker-guided gating strategy. Manual refinement was applied as needed following

unsupervised clustering to improve resolution of rare or ambiguous populations. Various

batches and organ types were integrated for analysis using Harmony [38].

Organ-specific analyses

To characterize lymphoid architecture in lymph nodes, we applied a multimodal spatial analysis

workflow. Spatial niche detection was performed using spaVAE [39], a dependency-aware

variational autoencoder model. Normalized lymph node datasets were input into spaVAE using

default parameters over 50 iterations. The resulting niche clusters were compared to Leiden-

based transcriptomic clusters for biological interpretability. Germinal centers (GCs) and peri-GC

regions were manually segmented using an interactive workflow implemented with

matplotlib.widgets.PolygonSelector and matplotlib.path.Path. Visualizations were generated in

Jupyter notebooks using %matplotlib tk, enabling real-time polygon annotation of GC

boundaries. Cells were classified based on spatial overlap as intra-GC, peri-GC, or non-GC.

For spatial cell–cell interaction analysis, neighborhood graphs were constructed using Squidpy's

sq.gr.spatial_neighbors() function. Cell–cell interaction matrices were generated with

sq.gr.interaction_matrix(), leveraging Leiden-inferred annotations to quantify contact frequencies

across annotated cell types within GCs and adjacent regions. Gene expression patterns were

visualized using Seaborn's clustermap() to highlight transcriptional diversity among niche

clusters. Spatial autocorrelation statistics were calculated using Moran's I

(sq.gr.spatial_autocorr()), excluding the top 1% of expression outliers. Genes with the highest

spatial variability were selected for downstream interpretation and figure display. To identify

differentially expressed genes (DEGs) between intra-GC and peri-GC compartments, we

applied Wilcoxon rank-sum tests (scipy.stats.ranksums()), followed by multiple hypothesis

correction using the Benjamini–Hochberg procedure (statsmodels.stats.multitest.multipletests()).

Genes with adjusted p < 0.05 and |log$_2$FC| > 0.5 were considered significant. Volcano plots
were generated to highlight key upregulated and downregulated genes across compartments.
886
Comparison and integration of CosMx Protein and RNA data
To annotate protein data, we used SCIMAP (v1.3.2), a gating-based hierarchical cell annotation
tool which takes a workflow with cell types noted by their marker genes to conduct cell typing
(**Extended Data Fig 7a**). Integration of CosMx RNA and protein data was performed using
MaxFuse (version 0.02). a Python (version 3.8.20) package that uses co-embedding, data
smoothing, and cell matching to integrate "weakly-linked" multi-modal omics data that have little
overlapping features [10]. Broad and granular annotations from RNA samples are integrated
into serial sectioned protein samples to generate protein cell type annotations. We generated
cell type mapping to analogous RNA and protein regions to create visual comparisons. To
conduct the concordance comparison between SCIMAP and Maxfuse annotations, we
examined how cells called by one modality are called by the other by calculating adjusted rand
index and mapping a matrix of cell counts for the cell types of both modalities as a heatmap.
For RNA-protein correlations and concordance comparisons, all analyses were conducted using
R version 4.4.3 and the following R packages: Seurat (v5.2.1), reticulate (v1.41.0), anndata
(v0.7.5.6), dplyr (v1.1.4), ggplot2 (v3.5.2), ggrepel (v0.9.6), reshape2 (v1.4.4), and
RColorBrewer (v1.1.3). No cell-wise or feature-wise filtering was applied to either the
transcriptomic or proteomic datasets for this analysis. We identified 53 features common to both
RNA and protein modalities across 15 matched tissue cross-sections from 9 different organs.
For each tissue pair, the mean RNA and protein expression levels for each common feature
were computed after excluding values equal to zero, to avoid bias from low-input
measurements. Mean values were also calculated at the organ level by averaging feature
expression across all tissue sections from the same organ. In addition, pooled averages were
computed by aggregating all features across all organs. To assess the relationship between
RNA and protein expression, we computed the log10-transformed mean expression values of
each feature and visualized them using scatter plots, with log10(RNA mean) on the y-axis and
log10(protein mean) on the x-axis. These plots were generated for tissue pairs, organs, and the
aggregated dataset. Linear regression models were fit to each of these comparisons using the
lm() function in base R, with the following formula: log10(RNA.mean) ~ log10(PRT.mean). We
also generated analogous plots and regressions to compare average RNA and protein
expression values across all common features within each tissue pair and organ. Spearman's
rank correlation coefficient was used to calculate RNA-protein correlations across organs and
genes.
919
Comparative analysis of H&E images with spatial transcriptomics datasets
Whole-slide H&E imaging data was processed using LazySlide [40]. Gastrointestinal tract
samples (appendix, colon, ileum) were segmented and tesselated with tiles of 512 pixels at 0.5
µm/pixel (~20X magnification). For each slide, we randomly sampled 250 tiles and extracted
high-dimensional morphological embeddings leveraging the CONCH multi-modal vision–
language model [13]. Subsampling was performed to ensure equal contribution of various
slides/tissue types. Tile embedding values were scaled, reduced with Principal Component
Analysis (PCA) and visualized using the Isomap (scikit-learn) or Uniform Manifold

Approximation and Projection (UMAP) algorithm [41] and clustered using the Leiden algorithm
[42] at resolution 0.5 with Scanpy [43]. For downstream interpretation, we assembled a panel of
79 histolopathological terms related to instestinal biology or pathology. We computed the
similarity between the text embedding of each term in each CONCH feature vector of each tile.
Text embedding term values were compared between morphological clusters using a Wilcoxon
test, and the top 3 most differential terms were used to describe each cluster.
In parallel, we applied CellViT [44] to the same whole-slide images to perform end-to-end
nuclear segmentation and classification. Within each sampled tile, we quantified cell-type
abundances (e.g., epithelial, immune, connective) and averaged these cellular proportions per
morphological cluster previously derived by the CONCH features. To link tile-based
morphological phenotypes with molecular cell-type composition, we performed canonical
correlation analysis (CCA) implemented in scikit-learn between the relative abundance of each
morphology cluster and the corresponding cell-type proportions inferred by the CosMx platform
per tissue slide.
942
Integrative analysis of SAHA COL and CRC CosMx data
Downstream analysis was performed using Seurat (v5.0.1) [30]–[34] throughout to assess cell
types and perform graph-based clustering. To harmonize gene probes across both datasets,
gene probes targeting closely related genes (previously combined in an earlier spatial probe set
but separated into individual genes in the current set) were merged. Specifically, genes
originally represented as single combined probes (e.g., *CCL3/L1/L3, CXCL1/2/3*) were merged
into single meta-probes by summing their expression values across individual gene
measurements. A complete list of merged genes is available in **Supplementary Table 3**. Genes
without common representation across both spatial datasets were excluded to ensure
consistent comparisons. After matching the probes, each FOV in the CRC sample was matched
to an FOV from the COL sample to achieve relatively equal cell numbers (Figure 7, Extended
Data Fig. 8).
Following this process, raw count matrices from each object were exported and combined using
the merge() function from Seurat. Normalization, scaling, and PCA calculation were performed
using standard Seurat functions. Objects were then integrated using Seurat's IntegrateLayers()
function with the Harmony algorithm [38] (parameters: theta = 2, lambda = 0.1). After
integration, Uniform Manifold Approximation and Projection (UMAP) embedding was performed
using 30 principal components (PCs). Clusters were defined using a resolution of 0.3, and low-
quality cells (Cluster 11) were excluded based on low feature counts. Prior to marker
identification, the JoinLayers() function was applied as recommended by Seurat developers.
Cluster markers were identified using Seurat's FindAllMarkers() function (parameters: min.pct =
0.2, logfc.threshold = 0.5, adjusted p-value < 0.05; Supplementary Table X).
Differentially expressed unique genes highlighted in Fig. 7c were identified with the
FindMarkers() function using the MAST test (min.pct = 0.1, logfc.threshold = 0.5, adjusted p-
value < 0.05; Supplementary Table X). Spatially differentially expressed genes (Fig. 7d-f) were
computed using the Voyager tool [45] to calculate Moran's I. Field of views (FOVs) were
clustered based on the top 200 spatially variable genes per field using principal component
analysis (25 PCs), followed by k-means clustering (k = 2) as visualized in Fig. 7d.
Visualizations were generated with Seurat's functions: UMAPs (Fig 7a, Extended Data Fig. f,h)
(DimPlot), Dotplots (Fig. 7h-2, Extended Data Fig. b,g) (DotPlot), violin plots (Fig 7c and e ,
Extended Data Fig. a) (VlnPlot), and pseudo-spatial images (Fig 7g-4, h-2, Extended Data Fig.

h-1) (ImageDimPlot). Bar plots (Fig 7g-3, Extended Data Fig. c,f) were created using
dittoBarPlot from dittoSeq [46], and spatial crypt visualizations were generated using ggplot2's
geom_point() (Fig. 7b,f, Extended Data Fig. d,e). Cell-cell interaction analysis (Fig. 7h-3) was
performed using CellChat software [47] (conversion factor = 0.18; interaction range = 250;
scale.distance = 4.5; contact range = 20).
979
Data explorer generation
All generated data were made available for interactive exploration using Vitessce [48]. For each
individual sample, cell segmentation polygons were converted into a JSON format in which each
cell is represented as an object containing a list of [x, y] coordinate pairs corresponding to its
boundary vertices. The visualization interface included a "Spatial" viewer, which displays the
sample in physical (x, y) space, allowing users to explore gene expression for any gene
alongside cell-level annotations derived from downstream analyses. Additional panels included
a "Quality Control" window showing violin plots for metrics such as transcripts per cell and
unique genes per cell, a "Features" window listing available genes, and a "Cell Labels" window
summarizing cell annotations. All panels were linked, enabling dynamic cross-filtering:
selections made in one viewer were automatically reflected across the others. Files generated
for the Vitessce viewers are hosted on Zenodo and the script used to generate these files was
deposited on github (See **Data and Code Availability**).
993
994
995 **Data and Code Availability**
996 Raw sequences, images, and processed datasets from single-nuclei sequencing and spatial
997 transcriptomics are available via Zenodo (link will be available upon acceptance), and code
998 blocks used to analyze the data can be accessed in the GitHub repository (link will be available
999 upon acceptance).
1000
1001

Acknowledgments
We thank the WorldQuant, GI Research Foundation, Bumrungrad International Hospital, the
National Institutes of Health (R01MH117406), and the LLS (MCL 7001-18, LLS 9238-16, 7029-
23). E.A., Y.Z., and A.F.R received funding from Angelini Ventures S.p.A. Rome, Italy. P.S.D.
thanks NIDDK (1U01DK134321, 5R01DK135620), NIAID (1U19AI181102), and Digestive
Health Foundation for the support.
We would like to acknowledge former SAHA members from NanoString Technologies, whose
contribution was crucial for establishing and generating the early phase of the project,
specifically Nathan Schurman, Joachim Schmid, Raymond Tecotzky, Sarah Church, and
Charlie Glaser. Most importantly, we thank the patients and their family members who provided
valuable samples for this research.

1013
1014

1015 **Author contributions**
1016 Conceptualization: J.P., C.E.M., R.E.S., S.L.H., A.A., R.D.G., E.H.
1017 Patient sample coordination: E.H., B.R., S.P., F.S.,

Experiment and data generation: A.A., R.D.G., H.C., S.M., P.S.D., Y.L., L.P., K.Y., A.H., M.K., D.M., L.W., A.W.

Investigation and data visualization: J.P., E.O., N.B., F.S.D., L.Z., M.M., Y.Z., E.A., J.K., M.R.A., E.M., J.R., Jac.P., A.A.A., P.D.,

Supervision: C.E.M., R.E.S., S.L.H. A.F.R., J.T.P., L.M., O.E., G.C., M.L., J.B.

Writing: J.P., R.D.G., F.S.D., L.Z., Y.Z., A.F.R., E.A., A.A.A., E.O., N.B., J.T.P., C.E.M., S.B., S.R., P.D., S.M.L.

All authors discussed the results and contributed to the final manuscript.

Competing interests

C.E.M. serves on the Scientific Advisory Board of Cellanome Inc. Competing interests for G.C. is detailed here: https://arep.med.harvard.edu/gmc/tech.html. E.M., S.R., P.D., J.R., Y.L., L.P., S.B., K.Y., A.H., M.K., D.M., L.W., A.W., and J.B. are current/previous Bruker Spatial Biology employees.

SUPPLEMENTARY TABLES

Supplementary Table 1. SAHA Cohort Demographics and Clinical Metadata.
Summary of demographic and clinical characteristics of the SAHA cohort, including patient age, sex, tissue of origin, comorbidity (if applicable), and sample designation.

Supplementary Table 2. SAHA Experimental Run Metadata.
Detailed metadata for all SAHA spatial transcriptomics runs, including informations such as slide/sample identifiers, tissue types, assay panels, staining protocols, quality metrics/statistics, platform used (e.g., CosMx, Xenium), and batch/run identifiers.

Supplementary Table 3. Additional Information related to CRC Analysis.
Additional files regarding the CRC and COL comparative analysis including list of probe substitutes, differential gene expression lists, cell marker lists, full result list of Moran's I results.

REFERENCES

1. Palla, G., Fischer, D. S., Regev, A. & Theis, F. J. Spatial components of molecular tissue biology. *Nat. Biotechnol.* **40**, 308–318 (2022).

2. Park, J. *et al.* Spatial omics technologies at multimodal and single cell/subcellular level. *Genome Biol.* **23**, 256 (2022).

3. Rood, J. E. *et al.* The Human Cell Atlas from a cell census to a unified foundation model. *Nature* **637**, 1065–1071 (2025).

4. Rozenblatt-Rosen, O. *et al.* The Human Tumor Atlas Network: Charting Tumor Transitions across Space and Time at Single-Cell Resolution. *Cell* **181**, 236–249 (2020).

5. Velten, B. & Stegle, O. Principles and challenges of modeling temporal and spatial omics data. *Nat. Methods* **20**, 1462–1474 (2023).

6. Cook, D. P. *et al.* A Comparative Analysis of Imaging-Based Spatial Transcriptomics Platforms. *BioRxiv* (2023) doi:10.1101/2023.12.13.571385.

7. Wilkinson, M. D. *et al.* The FAIR Guiding Principles for scientific data management and stewardship. *Sci. Data* **3**, 160018 (2016).

8. Liu, X. *et al.* Spatial multi-omics: deciphering technological landscape of integration of multi-omics and its applications. *J. Hematol. Oncol.* **17**, 72 (2024).

9. Yu, Y., Mai, Y., Zheng, Y. & Shi, L. Assessing and mitigating batch effects in large-scale omics studies. *Genome Biol.* **25**, 254 (2024).

10. Chen, S. *et al.* Integration of spatial and single-cell data across modalities with weak linkage. *BioRxiv* (2023) doi:10.1101/2023.01.12.523851.

11. Novoselsky, R. *et al.* Intracellular polarization of RNAs and proteins in the human small intestinal epithelium. *PLoS Biol.* **22**, e3002942 (2024).

12. Liu, S. *et al.* The role of CD276 in cancers. *Front. Oncol.* **11**, 654684 (2021).

13. Lu, M. Y. *et al.* A visual-language foundation model for computational pathology. *Nat. Med.* **30**, 863–874 (2024).

14. Liu, Y. & Chen, Y.-G. Intestinal epithelial plasticity and regeneration via cell

dedifferentiation. *Cell Regen (Lond)* **9**, 14 (2020).

15. Bala, P. *et al.* Aberrant cell state plasticity mediated by developmental reprogramming

precedes colorectal cancer initiation. *Sci. Adv.* **9**, eadf0927 (2023).

16. Koltsova, E. K. & Grivennikov, S. I. IL-22 gets to the stem of colorectal cancer. *Immunity*

40, 639–641 (2014).

17. Waldner, M. J., Foersch, S. & Neurath, M. F. Interleukin-6--a key regulator of colorectal

cancer development. *Int. J. Biol. Sci.* **8**, 1248–1253 (2012).

18. Liu, X., Li, X., Wei, H., Liu, Y. & Li, N. Mast cells in colorectal cancer tumour progression,

angiogenesis, and lymphangiogenesis. *Front. Immunol.* **14**, 1209056 (2023).

19. Xue, M. *et al.* Schwann cells regulate tumor cells and cancer-associated fibroblasts in the

pancreatic ductal adenocarcinoma microenvironment. *Nat. Commun.* **14**, 4600 (2023).

20. Zhang, B. *et al.* Single-cell RNA sequencing reveals intratumoral heterogeneity and

potential mechanisms of malignant progression in prostate cancer with perineural invasion.

Front. Genet. **13**, 1073232 (2022).

21. Zhang, Z. *et al.* Integrating clinical and genetic analysis of perineural invasion in head and

neck squamous cell carcinoma. *Front. Oncol.* **9**, 434 (2019).

22. Chen, H. *et al.* ANGPTL1 attenuates colorectal cancer metastasis by up-regulating

microRNA-138. *J. Exp. Clin. Cancer Res.* **36**, 78 (2017).

23. Jiang, K. *et al.* Exosomal ANGPTL1 attenuates colorectal cancer liver metastasis by

regulating Kupffer cell secretion pattern and impeding MMP9 induced vascular leakiness. *J.*

Exp. Clin. Cancer Res. **40**, 21 (2021).

24. Chang, T.-Y., Lan, K.-C., Chiu, C.-Y., Sheu, M.-L. & Liu, S.-H. ANGPTL1 attenuates cancer

migration, invasion, and stemness through regulating FOXO3a-mediated SOX2 expression

in colorectal cancer. *Clin. Sci.* **136**, 657–673 (2022).

25. Tang, W. *et al.* LIF/LIFR oncogenic signaling is a novel therapeutic target in endometrial

cancer. *Cell Death Discov.* **7**, 216 (2021).

26. Oliver, A. J. *et al.* Single-cell integration reveals metaplasia in inflammatory gut diseases.

Nature **635**, 699–707 (2024).

27. Zhang, J. *et al.* Tahoe-100M: A Giga-Scale Single-Cell Perturbation Atlas for Context-

Dependent Gene Function and Cellular Modeling. *BioRxiv* (2025)

doi:10.1101/2025.02.20.639398.

28. He, S. *et al.* High-Plex Multiomic Analysis in FFPE Tissue at Single-Cellular and Subcellular

Resolution by Spatial Molecular Imaging. *BioRxiv* (2021) doi:10.1101/2021.11.03.467020.

29. Wang, T. *et al.* snPATHO-seq, a versatile FFPE single-nucleus RNA sequencing method to

unlock pathology archives. *Commun. Biol.* **7**, 1340 (2024).

30. Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell

transcriptomic data across different conditions, technologies, and species. *Nat. Biotechnol.*

36, 411–420 (2018).

31. Satija, R., Farrell, J. A., Gennert, D., Schier, A. F. & Regev, A. Spatial reconstruction of

single-cell gene expression data. *Nat. Biotechnol.* **33**, 495–502 (2015).

32. Hao, Y. *et al.* Dictionary learning for integrative, multimodal and scalable single-cell

analysis. *Nat. Biotechnol.* **42**, 293–304 (2024).

33. Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902.e21

(2019).

34. Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.

(2021).

35. Germain, P.-L., Lun, A., Garcia Meixide, C., Macnair, W. & Robinson, M. D. Doublet

identification in single-cell sequencing data using scDblFinder. *F1000Res.* **10**, 979 (2021).

36. Bankhead, P. *et al.* QuPath: Open source software for digital pathology image analysis. *Sci.*

Rep. **7**, 16878 (2017).

37. Danaher, P. *et al.* Insitutype: likelihood-based cell typing for single cell spatial

transcriptomics. *BioRxiv* (2022) doi:10.1101/2022.10.19.512902.

38. Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with

Harmony. *Nat. Methods* **16**, 1289–1296 (2019).

39. Tian, T., Zhang, J., Lin, X., Wei, Z. & Hakonarson, H. Dependency-aware deep generative

models for multitasking analysis of spatial omics data. *Nat. Methods* **21**, 1501–1513 (2024).

40. Zheng, Y., Abila, E., Chrenková, E., Winkler, J. & Rendeiro, A. F. LazySlide: accessible and

interoperable whole slide image analysis. *BioRxiv* (2025) doi:10.1101/2025.05.28.656548.

41. McInnes, L., Healy, J. & Melville, J. UMAP: Uniform Manifold Approximation and Projection

for Dimension Reduction. *arXiv* (2018) doi:10.48550/arxiv.1802.03426.

42. Traag, V. A., Waltman, L. & van Eck, N. J. From Louvain to Leiden: guaranteeing well-

connected communities. *Sci. Rep.* **9**, 5233 (2019).

43. Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene expression

data analysis. *Genome Biol.* **19**, 15 (2018).

44. Hörst, F. *et al.* CellViT: Vision Transformers for precise cell segmentation and classification.

Med. Image Anal. **94**, 103143 (2024).

45. Moses, L. *et al.* Voyager: exploratory single-cell genomics data analysis with geospatial

statistics. *BioRxiv* (2023) doi:10.1101/2023.07.20.549945.

46. Bunis, D. G., Andrews, J., Fragiadakis, G. K., Burt, T. D. & Sirota, M. dittoSeq: universal

user-friendly single-cell and bulk RNA sequencing visualization toolkit. *Bioinformatics* **36**,

5535–5536 (2021).

47. Jin, S. *et al.* Inference and analysis of cell-cell communication using CellChat. *Nat.*

Commun. **12**, 1088 (2021).

48. Keller, M. S. *et al.* Vitessce: integrative visualization of multimodal and spatially resolved

single-cell data. *Nat. Methods* **22**, 63–67 (2025).

1152

FIGURES

Figure 1: Overview of SAHA Workflow and Data Scope.
a, Schematic overview of SAHA resource, summarizing sample collection and spatial profiles. The diagram illustrates the scale of (1) cohort, (2) organ types, (3) batches, (4) cells, and (5) biomolecules profiled across multiple platforms and modalities.
b, Heatmap showing the SAHA sample (healthy tissues) demographics across profiled organs, colored by age, sex, and ethnicity, with missing information labeled as unknown in gray.
c, Distribution of total cell numbers across healthy and diseased tissues stratified by technology and organ type; top right corner shows the summary of spatial technologies deployed (CosMx RNA and Protein, Xenium, GeoMx, RNAscope, H&E) with representative images from the same section in each category labeled with the same color.
d, Comparative overview of the published spatial atlases and SAHA by number of cells profiled, organ coverage, and assay modality (x axis). The dot size indicates the number of cells, and the colour indicates the maximum number of genes in the panel.

Figure 2: Key Structures, Cell Types, and Cellular Networks across SAHA Organs.
a, UMAP embeddings (n = 2,865,647 cells) colored by organ (top) or broad cell type (bottom).
b, Stacked violin plots showing canonical marker genes across major cell types, with color intensity reflecting median expression level.
c, Stacked bar chart summarizing the broad cell-type composition per section, illustrating the relative abundances of epithelial, immune, stromal, and neuronal cells.
d, Representative histological (H&E) sections, spatial RNA and protein images, and organ-specific UMAP embeddings from each tissue, demonstrating subcellular resolution and morphological context.

Figure 3: Spatial Neighborhoods and Cellular Niches of the Crypt.
a, Representative colon cross-section, showing spatially resolved cell types (colors represent cell types) across the mucosa, submucosa, and muscularis layers. Graphs to the right display the expression of key markers (e.g., *PIGR*, *MHC-I*, *ACTA2*) stratified by tissue depth.
b, Spatial enrichment scores of key biological pathways (keratinization, antigen processing, smooth muscle contraction) stratified by tissue depth.
c, Spatial mapping of proliferative and stress-response gene expressions (i.e., *CDKN1A*, *CCND1*, *TFEB*, and *SMARCB1*) within a magnified region at crypt boundaries.
d, Dot plot of enriched immune-epithelial crosstalk represented by ligand-receptor interactions from the crypt tip niche, sized by -$\log_2$ p-value.
e, Segmented overlays and heatmaps depicting cell type composition (e.g., strong T- and B-cell predominance near crypts, variable mesenchymal content deeper in the colon wall) across different tissue layers.
f, Cell type enrichment comparisons across all colon sections (left) and selected crypt regions, including colon cross-section (middle) and crypt tips (right).
g, Spatial projection of large lymphoid aggregates in SAHA APE, color-coded by cell type.
h, Force-directed graph showing adjacency networks among immune and stromal cells within lymphoid aggregates.

45   **i**, Spatial projections of cell types by both broad and detailed annotations, highlighting
46   specialized immune subtypes (e.g., T follicular helper) and their heterogeneity.
47   **j**, Chord diagram and heatmaps of global and niche-specific ligand-receptor interactions, The
48   left panels indicate overall interaction frequency among major cell types, while the right panels
49   focus on specialized niches (e.g., lymphoid structures identified from APE).
50
Figure 4: Comparison of Spatial Neighborhoods and Cellular Niches across Organs.
a, Representative spatial mapping of lymphoid structures in SAHA lymph node (LN) samples,
showing transcriptionally defined niches and cell type composition.
b, Bar plots comparing the proportion of B cells across "in GC (or lymphoid structure for
appendix)" and "around GC" niches in LN and appendix (APE).
c, Spatial expression plots of canonical B cell markers (*MS4A1*, *IGHG1*) in LN and APE,
illustrating organ-specific localization and expression intensity.
d, Cell-cell interaction matrices stratified by spatial niche showing overall interaction patterns
around germinal centers.
e, Pathway enrichment analysis of genes shared between LN and APE that are both
differentially expressed and spatially autocorrelated, showing enrichment for immune and
follicular activation processes.
f, Top enriched pathways among APE-specific spatially variable DEGs, highlighting epithelial
signaling and extracellular remodeling.
g, UMAP embedding of integrated gastrointestinal spatial datasets (APE, ILE, COL, STO),
colored by tissue of origin and spatial neighborhood clusters.
h, Representative spatial projections showing neighborhood clustering across organs. Colors
represent unbiased spatial clustering involving epithelial, immune, and stromal compartments.
i, Network graph illustrating relationships among organs, spatial neighborhood clusters, and
broad cell types. Nodes represent variables (organ, spatial cluster, and cell type), colors of the
edges represent edge weights (spatial connectivities), and the distance between nodes
represents expression-level correlation.
j, Stacked bar chart of organ-specific distributions of spatial neighborhoods, where colors
represent unbiased spatial clusters shown in **g-h**.
k, Relative frequencies of selected niche clusters, highlighting shared and organ-restricted
spatial patterns; colors represent cell types.
l, Expression level (top) and spatial autocorrelation (bottom) comparison of selected genes
across spatial neighborhoods, illustrating spatial heterogeneity and functional specialization
within gastrointestinal tissue clusters.
80
Figure 5: Integration of Spatial Proteomics Data for Multi-omics Map
a, UMAP projection of normal SAHA spatial proteomics datasets from normal tissues following
integration. Data are colored by tissue of origin (left), cell type labels using protein information
(middle), and labels done with both RNA and protein information (right).
b, Heatmap of protein expression of RNA-derived cell type labels, showing canonical and cross-
modality marker distributions.
c, Comparison of cell type assignments from RNA (top) and protein (bottom) datasets in
representative fields of view.

d, Representative spatial images from SAHA PROS, APE, and STO tissues, comparing RNA-
based (top) and protein-based (bottom) cell type assignments from the same fields of view.
e, Validation of ligand-receptor analysis using spatially resolved protein imaging, where protein
expression of key immune markers (e.g., CCR7, HLA-DRA, CD4) overlaid on tissue sections
(left) and ligand–receptor pair contributions based on spatially restricted analysis within selected
regions (right) were shown.
f, Correlation plot of average RNA and protein expressions across overlapping markers..
96
Figure 6: Integration of Matched Histopathological imaging with Transcriptomics Data
using Multimodal Foundation Models
a, Whole-slide H&E image of a representative SAHA slide (APE) showing tiled tissue regions
used for feature extraction.
b, Schematic overview of the computational workflow: tissue regions are segmented,
tesselated, and input to a multimodal foundation model that extracts quantitative morphological
features for each region.
c, Low-dimensional embedding of image-derived morphological features using Isomap, showing
tissue structure captured solely from histological patterns corresponding to morphologically-
driven clusters.
d, Clustered heatmap of morphological feature embeddings across tissue locations, revealing
spatial gradients and local heterogeneity.
e, Spatial reconstruction of tissue based on morphological clusters, demonstrating regional
organization inferred from morphological features alone.
f, UMAP embedding of morphological tile clusters from the gastrointestinal tract (ileum,
appendix, colon), with representative histological images from each cluster.
g, Organ-wise comparison of morphological clusters (top), annotated with similarity scores to
histopathology terms (bottom), revealing conserved structures such as crypts, lymphoid follicles,
and extracellular matrix regions.
h, Clustered heatmap showing enrichment of histopathological terms (rows) across morphology-
derived clusters (columns), highlighting distinct text-derived phenotypes.
i, Composition of broad cell types (epithelial, connective, immune), segmented and classified in
the histology images, within each morphology-derived cluster, revealing dominant biological
compartments.
j, Pairwise distance matrix showing spatial proximity between clusters, with warmer colors
indicating closer physical co-localization.
k, Canonical correlation analysis (CCA) comparing tile-level morphological cluster abundances
with RNA-defined cell type abundances, showing correspondence between structure and cell
states.
l, Example region from an H&E image (top) alongside matched CosMx cell-type annotation
(bottom) illustrating alignment between morphological features and transcriptomics.
128
Figure 7: Spatial Heterogeneity in Cellular and Molecular Profiles Across Healthy Crypts
from COL and CRC Samples.

a, UMAP projection of integrated healthy SAHA colon samples (SAHA COL, n=3) with colorectal
cancer (CRC, n=1) following Harmony integration, colored by cell types (left) and disease status
(right).
b, Representative spatial images from CRC tumor adjacent and SAHA COL tissues, colored by
cell types.
c, Gene expression profiles of top and bottom crypt regions in SAHA COL and CRC tumor
adjacent samples. Violin plots show normalized expression levels, colored by disease status.
Genes shared between SAHA COL and CRC samples (left) and differentially expressed genes
in CRC tumor adjacent crypts (right) are shown.
d, K-means clustering of crypt FOVs based on spatially variable genes identified by Moran's I.
Points are colored by sample status and shaded regions denote k-means clusters.
e, Comparison of spatially variable genes between CRC tumor adjacent and healthy crypts.
Moran's I correlation plot from a representative CRC vs. SAHA COL FOV comparison (left) are
shown, genes with Moran's I > 0.2 in either condition are highlighted in blue. Dashed red line
indicates equal spatial autocorrelation. Violin plots of condition-specific spatial genes are shown
(right).
f, Spatial expression of *IL22RA1* in healthy and CRC tumor adjacent crypts, representative of
differentially enriched crypt genes in panel **e,** shown across normalized assay values; insets
shown cell type annotations.
g, Intrapatient tumor heterogeneity in CRC sample, where H&E-stained section with annotated
regions showing Type 1 (purple) and Type 2 (blue) tumor areas.
h, Lymphocyte-to-tumor cell ratio across FOVs, grouped by dominant tumor type calculated by
the percentage of B cells and T cells divided by the tumor cell percentage in each FOV. Mixed
denotes the FOV samples that have a close percentage of type 1 and type 2 tumor cells. Rest
of the dots are colored based on the region of the tumor within the tissue.
i, Proportion of immune cell types in Type 1 vs. Type 2 tumor FOVs, colored by cell types that
are manually curated.
j, Representative FOVs from each tumor type, colored by the cell type.
k, Spatial transcriptomics reveals unique perineural invasion (PNI) structure with great detail,
demonstrated by H&E image of the corresponding PNI region (left) and spatial projection of cell
types, where bright green highlights the glial cells that are surrounded by fibroblasts and tumor
glands (right).
l, Dot plot of fibroblast markers highlighting the difference between close-contact fibroblasts to
the nerve versus rest of the fibroblasts in the data.
m, Unique signaling interactions from close fibroblasts (ANGPTL pathway, left) and glial cells
(LIFR pathway, right) toward immune and tumor compartments.
167
Figure 8: Application of the SAHA healthy ileum reference to spatially profile IBD tissues
and therapeutic response.
a, Schematic overview of comparative analysis using SAHA ILE reference and ileal samples
from IBD patients stratified by TNFα inhibitor response (Responder, Non-responder).
b, UMAP embeddings of CosMx RNA profiles colored by major cell types (left) and sample
identities (right) across healthy ILE and IBD tissues.

c, Spatial mapping of cell types in representative IBD and healthy ileum sections, showing
increased immune infiltration in IBD tissues.
d, Quantification of immune cell proximity to epithelial cells (within 300 µm), demonstrating
significantly higher immune infiltration scores in IBD compared to healthy ileum (***$p < 0.001$).
e, Ligand-receptor interaction scores between immune and epithelial cells across IBD
responders, non-responders, and healthy ileum, showing elevated spatial signaling in non-
responders.
f, Violin plot of *TNFRSF1A* (TNF receptor 1) expression in mesenchymal-stromal cells across
IBD responders, non-responders, and healthy ileum (**$p < 0.01$).
g, Spatial projection of TNF-TNFRSF1A ligand-receptor interactions in a representative IBD
non-responder sample, highlighting immune–stromal interface activation.
h, Represented images of IBD and SAHA ILE samples (left) with UMAP embedding of
unsupervised clustering of H&E-stained ileum sections, stratifying normal and IBD tissues
based on morphological features (right).
i, Combined scores of histological features associated with IBD pathologies in responders and
non-responders, including increased crypt branching, immune cell infiltration, and mucosal
edema.
191
Figure 9: Impact of SAHA, SAHA Data Portal and Accessibility.
A, (to be updated) Comparison between traditional single-cell transcriptomic profiling (left) and
spatial multi-omics profiling integrating morphological information (right), combining H&E
staining, cell boundary segmentation, and multiplexed RNA and protein expression profiling
(19k whole transcriptome probes + 67-plex protein).
b, Overview of cost and scalability considerations for spatial multi-omics at subcellular
resolution.
c, SAHA data portal architecture, enabling integrated visualization and analysis of spatial multi-
omics datasets. The portal supports interactive exploration of spatial transcriptomics and
proteomics through UMAP and PCA projections, cell-type annotations, spatial imaging data,
gene expression matrices, and downloadable formats including image files, flatfiles, AnnData,
and Seurat objects for analysis in Python and R.
204

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 7

Fig. 8

Fig. 9

EXTENDED DATA FIGURES AND TABLES

Extended Data Fig. 1: SAHA project workflow and data generation pipeline.
Overview of tissue collection, sample preparation, multimodal spatial profiling (CosMx, Xenium, GeoMx, RNAscope, snPATHO-seq), data integration, and open-access portal deployment.

Extended Data Fig. 2: Detailed Cell Types Identified from SAHA.
a, UMAP projections colored by detailed cell-type annotations derived from canonical marker gene expression.
b, Stacked bar charts showing detailed cell-type composition across sections and the relative abundance of epithelial, immune, stromal, and neuronal subtypes.
c, Organ-specific distribution of broad cell types (epithelial, immune, stromal, neuronal, etc.).

Extended Data Fig. 3: Overall Quality Assessment of SAHA Data.
a, Representative RNAscope segmentation and quantification results.
b, Comparison of RNAscope mean transcript counts of control probe PPIB (left) versus CosMx RNA mean counts for matched samples (right).
c, Correlation analyses (Spearman's ρ, top) and Bland-Altman plots (bottom) assessing agreement between RNAscope and CosMx measurements. CosMx RNA mean counts (left) and CosMx RNA feature counts (right) across samples were compared, and each dot represents a sample. Grey dashed lines represent the linear regression fit with shaded areas indicating the 95% confidence intervals. For Bland-Altman analysis, the x-axis represents the mean of the two measurements, and the y-axis shows the difference (CosMx – RNAscope), where each sample is labeled with its sample ID number, and the grey dashed line indicates the mean difference (bias), and red dashed lines indicate the 95% limits of agreement (mean ± 1.96 × SD).
d, Quality control metrics compared across gastrointestinal organ cohorts, demonstrating consistent assay performance.

Extended Data Fig. 4: Spatial mapping of lymphoid structures and cellular neighborhoods in the appendix.
a, Spatial expression patterns of highly variable genes (*IGHA1, IGKC, CD74, CCL21*) highlighting lymphoid structures in SAHA APE as an example.
b, Spatial projections of cellular neighborhoods and detailed cell types across representative fields of view (FOVs) in SAHA APE, colored by cell type and spatial neighborhoods (left); representative spatial projection of lymphoid structure showing distributions of spatial neighborhoods (middle) and broad and granular cell types (right) in a single FOV.

Extended Data Fig. 5: Cellular diversity and spatial neighborhood organization in SAHA lymph nodes.
a, UMAP embeddings of SAHA LN B cell niches (left) and transcriptionally defined cell types (right), including Lymphoid-B, Follicular B, Plasma subsets, and stromal components.
b, Heatmap showing normalized expression (z-score) of B cell and plasma cell marker genes across annotated LN cell types, highlighting class-switching and activation gradients.

c, Spatial projections of LN section showing annotated GC and peri-GC regions used for niche
comparison. Cells not used in GC analysis are shaded in gray.
d, Cell-cell interaction matrices stratified by cell types revealing distinct interaction patterns in
germinal center, relative to **Fig. 4d**.
e, Expression projections of selected differentially expressed genes in GC vs. peri-GC cells.
f, Volcano plot comparing GC vs. peri-GC B cells in LN, showing significantly upregulated (red)
and downregulated (blue) genes.
g, Spatial projections of colon (top) and LN (bottom) samples colored by broad cell type
annotations.
h, UpSet plot showing intersecting gene sets between differentially expressed genes (DEGs)
and spatially variable genes across LN and colon GC niches. Shared and unique signatures
highlight conserved and tissue-specific B cell programs.
56
Extended Data Fig. 6: Spatial network architecture and immune-epithelial interactions
across gastrointestinal tissues.
a, Spatial network statistics (average clustering coefficient, closeness centrality, degree
centrality) across SAHA APE, ILE, COL, and STO tissues.
b, Co-occurrence analysis of epithelial cells with other cell types within crypt-associated regions.
c, Top ligand–receptor interactions between immune and epithelial cells within lymphoid
structures.
d, Comparison of various spatial neighborhood clustering methods across APE, ILE, and COL
crypt structures.
e, Significant immune-epithelial ligand-receptor interactions across gastrointestinal organs.
67
Extended Data Fig. 7: Comparison of RNA and protein expression across tissues and
validation of annotation consistency.
a, Hierarchical cell type classification scheme used for protein-based phenotyping with SciMap.
b, Cell type proportions across major organs based on protein-based annotations.
c, Comparison of cell type assignments from RNA (top) and protein (bottom) data from **Fig. 5c**.
d, Label concordance between MaxFuse-integrated annotations and SciMap gating-based
annotations confirming consistent cell type classification across platforms.
e, Scatter plots of average RNA versus protein expression across matched genes for each
organ, showing tissue-specific correlation patterns. Each dot represents a gene with paired RNA
and protein measurements; red line indicates linear regression, and shaded area denotes 95%
confidence interval.
f, Zoomed-in views for representative genes with different RNA–protein concordance profiles.
80
Extended Data Fig. 8: Integrated spatial annotation and cell-type resolution in CRC and
SAHA COL samples.
a, Violin plots showing the raw distribution of detected features and counts across CRC and
SAHA COL samples.
b, Dot plot of cell-type specific marker genes used for annotation in the integrated dataset of
CRC and SAHA COL samples. Dot size corresponds to the percentage of cells within a cluster
expressing the gene, and the color represents the scaled gene expression level. Clusters

labeled "mixed" and "Epithelial_tcell" include expression patterns that may represent multiple
cell types likely due to segmentation artifacts.
c, Bar plots showing the overall percentage and the number of cells across CRC and SAHA
COL.
d, Representative FOVs from CRC and SAHA COL, colored by cell types from panel **c**.
e, Expression maps showing examples of CRC-enriched spatially variable genes identified in
Fig. 7e. Expression of *REG1A* and *MX1* is shown across CRC and SAHA COL FOVs.
f, UMAP projection of CRC sample (n=1), colored by the cell types.
g, Dot plot of marker genes used to identify CRC clusters. Dot size corresponds to the
percentage of cells expressing the gene within each cluster and color shows the scaled
expression value.
h, Perineural invasion (PNI) FOV clustered independently to resolve fibroblast subtypes. Close-
contact fibroblasts adjacent to the nerve are distinguished (purple) from other fibroblast
populations (red).
102
Extended Data Fig. 9: Cell-type level comparisons between SAHA healthy ileum and IBD
tissues.
a, Spearman correlation matrix of cell-type average expression profiles between 1000-plex
(lower-plex) and 6000-plex (higher-plex) CosMx RNA datasets, clustered by broad and detailed
cell types. Higher correlation values indicate consistent cell-type transcriptional programs across
datasets.
b, UMAP embedding of detailed cell-type annotations derived from CosMx RNA profiling,
colored by cell type.
c, Heatmap showing relative cell-type composition across individual samples from healthy ileum
(ILE) and IBD tissues, stratified by sample ID.
d, Stacked bar plots summarizing cell-type composition aggregated by clinical condition (healthy
ileum, IBD responder, IBD non-responder), highlighting changes in epithelial, immune, and
stromal populations between healthy and disease states.
116
117

 Extended Data Fig. 1

 Extended Data Fig. 2

 Extended Data Fig. 3

 Extended Data Fig. 4

 Extended Data Fig. 5

  Extended Data Fig. 6

Extended Data Fig. 7

 Extended Data Fig. 8