1

Evolutionary Genomics Guides Scalable Coral Probiotics for Climate Resilience

Mei Xie[1]†, Nan Xiang[1]†, Tianhua Liao[1]†, Chun Ting Cheung[1], Chi Hin Lee[1], Wai Yin Tse[1],

Kwong Kit Tsang[1], Kaitlyn E. Ho[1], Qian He[1], Melanie Dörr[2], Hannah Manns[2], Pan Li[1], Zhihao

Xian[1], Xiaojun Wang[1], Danli Luo[1], Róisín Hayden[3,4], Emily Chei[3,4], Zhongyue Wan[3,4], Philip

Thompson[3,4], Joseph Brennan[3,4], Raquel S. Peixoto[5], Guoxin Cui[5], Shelby E. McIlroy[1,3,4], Apple

Pui Yi Chui[1], Christian R. Voolstra[2], Haiwei Luo[1,6,7,8]*

[1]Simon F. S. Li Marine Science Laboratory, School of Life Sciences, The Chinese University of

Hong Kong; Shatin, Hong Kong SAR.

[2]Department of Biology, University of Konstanz; Konstanz, Germany.

[3]The Swire Institute of Marine Science, The University of Hong Kong; Cape D'Aguilar Road,

Shek O, Hong Kong SAR.

[4]School of Biological Sciences, The University of Hong Kong; Pok Fu Lam, Hong Kong SAR.

[5]Division of Biological and Environmental Science and Engineering (BESE), Red Sea Research

Center, King Abdullah University of Science and Technology (KAUST); Thuwal, Saudi Arabia.

[6]State Key Laboratory of Agrobiotechnology, The Chinese University of Hong Kong; Shatin,

Hong Kong SAR.

[7]Department of Earth and Environmental Sciences, The Chinese University of Hong Kong;

Shatin, Hong Kong SAR.

[8]Institute of Environment, Energy and Sustainability, The Chinese University of Hong Kong;

Shatin, Hong Kong SAR.

23   †These authors contributed equally to this work.

25   ***Corresponding author:**

26   E-mail: haiweiluo@cuhk.edu.hk

27
28

Abstract

Coral reefs sustain nearly 30% of marine biodiversity [1] and provide ecosystem services valued at $2.7

trillion USD annually [2], yet face catastrophic collapse under climate change [3], with 70-90% projected for

extinction by 2050 even if the 1.5°C Paris target is met [4]. While microbial probiotics could enhance

resilience, scalability is limited by transient colonization and uncertain long-term efficacy [5-9]. Here, we

introduce an evolutionary genomics framework to select next-generation probiotics targeting bacteria

undergoing irreversible transitions to host dependency. Through laboratory proof-of-principle, we show

the coral-sourced *Ruegeria* MC10, identified by genomic signatures of host adaptation (insertion

sequence proliferation and pseudogenization of essential metabolic genes), increases thermal tolerance

thresholds by 1°C in model cnidarians. Subsequent, real-world application of one-month nursery

inoculations enabled *Acropora pruinosa* corals to retain MC10 for the monitored period (eight months)

post-outplanting, including persistence through a natural bleaching event, with evidence for increased

resilience. This work establishes lifestyle-oriented probiotic selection as a scalable solution for microbial-

driven reef restoration. Critically, this framework addresses poor probiotic colonization and limited

scalability, which pose universal limitations across conservation and aquaculture that hinder microbial-

based interventions [10-13]. By merging evolutionary biology with conservation science, we provide a

generalizable blueprint applicable to diverse threatened wildlife.

Main Text

Coral reefs sustain nearly 30% of marine biodiversity [1] and support the livelihoods of over one

billion people through ecosystem services and coastal protection [14]. Yet these vital ecosystems

are collapsing under climate change: since the 1950s, half of global live coral cover has vanished,

and even under the 1.5 °C Paris target, 70-90% of reefs face extinction by 2050 [4]. Central to this

crisis is the breakdown of the coral-algal symbiosis during thermal stress, which triggers

bleaching and starvation [15]. The accelerating frequency of mass bleaching events, including the

catastrophic 2014-2017 [16] and 2023-2024 [17] global events, has left reefs with insufficient

recovery intervals, necessitating urgent interventions to enhance resilience.

The targeted manipulation of microbial communities is considered a promising strategy to

enhance the resilience of endangered wildlife [7], but uncertainty regarding establishment of long-

term relationships remains [11]. Microbial therapies, particularly probiotics (i.e., beneficial

microbes administered to improve host health), offer promise for mitigating coral bleaching by

modulating holobiont health. Laboratory studies demonstrate that probiotic inoculations can

enhance thermal tolerance, disease resistance, and calcification [8,18,19], while field trials confirm

their safe delivery with limited evidence for off-target effects [5].

To address the issue of long-term residency of bacterial inoculants, we introduce so-called

next-generation probiotics selected through evolutionary principles. By targeting beneficial

bacteria undergoing irreversible transitions from free-living to host-dependent symbiosis, we

identify microbes pre-adapted for stable colonization. We focus on evolutionary young

endosymbionts, which retain culturability but exhibit genomic adaptations (e.g.,

pseudogenization of core metabolic genes) indicative of host dependency, assumed to align with

more persistent association in the tissue. Through screening of 1,212 bacterial isolates from

corals, we identified a novel population taxonomically affiliated with *Ruegeria*, a genus ranked

among the most prevalent coral-associated taxa [20,21] with a coral-sourced isolate recently shown

to be a coral probiotic [22], that exhibits genomic hallmarks of host adaptation. Nursery

inoculations enable MC10 to persist in coral tissues throughout an 8-month field trial, including

through a natural bleaching event, with evidence for enhancing coral thermal resilience without

repeated applications. Unlike traditional probiotics, which are selected for transient traits,

MC10's efficacy stems from evolutionary compatibility with its host. By merging evolutionary

biology with wildlife conservation, we establish a novel framework for climate-resilient

restoration.

Results

Ruegeria population MC10 as a putative next-generation coral probiotic

To identify candidate probiotics, we cultured 419 *Ruegeria* isolates from five coral species

(*Acropora* cf. *samoensis*, *Acropora solitaryensis*, *Favites pentagona*, *Platygyra acuta*, and

Oulastrea crispata) across nine Hong Kong reef sites spanning diverse environmental gradients,

including dissolved inorganic nitrogen concentrations of 21-107 µM (**Fig. S1**). Most bacterial

isolates were obtained from coral mucus and skeleton (**Fig. 1A**).

87   Using PopCOGenT to delineate bacterial populations based on recombination barriers [23],

we resolved 116 genetically cohesive *Ruegeria* populations, 29 of which included ≥3 isolates

(**Fig. 1A**). Phylogenomic analysis confirmed monophyly for most populations, validating

PopCOGenT's accuracy and enabling targeted surveys of probiotic dynamics. Among these,

population MC10 emerged as a prime candidate, exhibiting genomic hallmarks of evolutionary

young endosymbionts: expansion of IS (**Fig. 1B**) and proliferation of pseudogenes (**Fig. 1C**), as

quantified from closed genome assemblies (**Fig. S2**). Observed frameshift mutations disrupted

key tricarboxylic acid (TCA) cycle genes, including aconitate hydratase (*ACO*), isocitrate

dehydrogenase (*IDH1*), succinyl-CoA synthetase (*sucC*), malate dehydrogenase (*mdh*), and

succinate dehydrogenase (*sdhA*) (**Fig. S3**). While 5′-terminal mutations likely abolished gene

function, 3′-terminal disruptions may permit residual activity (**Fig. S3** & **Fig. S4**). Notably,

pseudogenized loci lacked functional paralogs, suggesting heightened metabolic reliance on the

coral host. Strikingly, only 8 of 15 closed MC10 genomes retained a complete TCA cycle, with

similar heterogeneity observed in glycolysis, amino acid biosynthesis, and vitamin synthesis

pathways (**Fig. S5**). While population MC10 exemplifies the hallmark features for next-

generation probiotics, within population functional heterogeneity presents a potential source of

variation in the efficacy of probiotic candidate lineages.

Host dependency may come at the cost of free-living capacity. To test this, we compared

the growth patterns of the strain MC10-B4 (from MC10) in culture to its closely related *Ruegeria*

strain MC15-BG7 (from MC15) sourced from the same *O. crispata* colony and a free-living fast-

growing model strain *Ruegeria pomeroyi* DSS-3 isolated from coastal water [24] (**Fig. S6**).

Consistent with its host-adapted genomic architecture, MC10-B4 exhibited reduced maximum

growth rates (μ_{max} = 0.15 h^{-1}) in culture compared to its congenic relative MC15-BG7 (μ_{max} =

0.27 h^{-1}), but comparable to DSS-3 (μ_{max} = 0.12 h^{-1}). However, MC10-B4 achieved a 1.4-fold

higher stationary-phase biomass yield (7.4 × 10$\square$ CFU/mL) than MC15-BG7 (5.2 × 10$\square$

CFU/mL) and a 7.7-fold increase over the DSS-3 (9.6 × 10$\square$ CFU/mL) despite similar optical

densities.

Phylogenetically, MC10's closest relative, MC46 (**Fig. 1A**), shares 99.45% 16S rRNA

identity though diverges more genomically (average nucleotide identity or ANI: 86.88 ±

0.04%; **Fig. S7**), lacking IS expansion or pseudogenization (**Fig. 1B & 1C**). This contrast

underscores MC10's recent origin.

Ruegeria MC10 is ubiquitously associated with diverse corals

To assess *Ruegeria* MC10's ecological distribution and coral host associations, we performed

amplicon sequencing of two population-resolving marker genes, *parC* (encoding a

topoisomerase IV subunit) and *ATP5B* (encoding an ATP synthase subunit). These markers were

selected from over 700 conserved single-copy *Ruegeria* gene families and validated for their

ability to resolve PopCOGenT-defined populations into monophyletic groups (**Fig. S8**). Custom

primers achieved high recovery rates (64.8% for *parC*; 74.3% for *ATP5B*; **Table S1**), enabling

targeted surveys across corals.

Phylogenetic analysis of amplicon sequences revealed MC10's presence in the five

surveyed coral species (*O. crispata*, *Acropora solitaryensis*, *Plesiastrea* sp., *Porites* sp., and *P.*

acuta) sampled from multiple reef sites in Hong Kong, as well as in seawater and sediment (**Fig.**

2). MC10's occurrence varied among individuals of the same coral species, typically constituting

<2% of *Ruegeria* communities, where detected. Notably, *parC-* and *ATP5B*-based abundance

estimates showed no linear correlation, particularly at proportions below 0.5% (**Fig. S9**), likely

reflecting methodological challenges in quantifying low-abundance populations.

Attempts to quantify MC10's absolute abundance via qPCR were unsuccessful (**Fig. S10**),

though its recurrent detection across phylogenetically distant corals and abiotic habitats

underscores its ecological versatility. Despite methodological limitations for quantification,

MC10's broad distribution and transient environmental persistence support its classification as a

low-risk probiotic candidate, minimizing concerns of ecological disruption in restoration efforts.

Ruegeria MC10 increases thermal tolerance of a model cnidarian

To validate MC10's probiotic potential, we employed short-term acute thermal assays using the

Coral Bleaching Automated Stress System (CBASS) [25,26], exposing Aiptasia (sensu *Exaipatasia*

diaphna, a sea anemone model for coral-algal symbiosis) [27] to MC10-B4 at a pre-screened

concentration of 10^6 CFU/ml (**Fig. S11**) with subsequent subjection to thermal stress (**Fig. S12**).

The CBASS trial involved a 3-hour temperature ramp from 30 °C to target levels (34 °C, 36 °C,

or 39 °C) under light, followed by a 3-hour hold, a 1-hour ramp-down to 30 °C in darkness, and

an 11-hour overnight reprieve period [28]. Dark-acclimated photosynthetic efficiency (F_v/F_m) was

used to measure response to heat stress at two time points: T7 (7[th] hour, immediately post-stress)

to assess acute thermal tolerance, and T18 (18[th] hour, post-recovery) to evaluate thermal

recovery [28]. Thermal tolerance was quantified by means of the Effective Dose 50 (ED50), which

signifies the temperature at which a 50% decline in F_v/F_m relative to unstressed controls is

observed [29]. MC10-inoculated Aiptasia exhibited significantly higher ED50 standardized thermal

tolerance values than controls: a 1 °C increase at T7 ($P = 0.001$, one-way ANOVA) and a 0.6 °C

increase at T18 ($P = 0.013$), demonstrating enhanced acute stress tolerance and recovery capacity,

respectively (**Fig. 3A & 3B, Table S2**). Fluorescence in situ hybridization (FISH) assays with an

MC10-specific probe confirmed presence of MC10-B4 in inoculated Aiptasia tissues (**Fig. S13**).

These results, generated using Aiptasia's conserved symbiosis mechanisms, validate MC10's

cross-host efficacy.

Real-world application demonstrates MC10's probiotic capacity through sustained thermal tolerance in reef corals

To evaluate *Ruegeria* MC10's probiotic capacity in natural settings, we conducted an 8-month

field trial with *Acropora pruinosa*, initially designed to compare two evolutionarily distinct

strains: MC10-B4 (exhibiting host-adapted genomic signatures) and its phylogenetically closely

related (**Fig. 1**) congener MC15-BG7 (lacking such signatures), both isolated from the same *O.*

crispata colony. Following a one-month laboratory inoculation regime (January 2024, **t0**) using

genetically identical coral fragments to control for host variability, amplicon sequencing revealed

stark contrasts in colonization success. MC10-B4 achieved significantly higher abundance in

inoculated corals (median 18.331% of *Ruegeria* communities; 0.003% of total bacteria),

significantly higher than the placebo group treated with autoclaved 0.22 μm-filtered seawater

(AFSW) (2.410% of *Ruegeria* communities and < 0.001% of total bacteria; $P = 0.002$ and $P =$

0.007, respectively; one-way ANOVA; **Fig. 3C, Table S3**). In contrast, corals inoculated with

MC15-BG7 showed no significant increase in MC15 abundance (median 3.242% of *Ruegeria*

communities and < 0.001% of total bacteria) compared to the placebo group (0.251% of

Ruegeria communities and < 0.001% of total bacteria; $P = 0.136$ and 0.281, respectively; one-

way ANOVA; **Fig. 3D, Table S3**). Given MC10-B4's superior colonization, we prioritized these

colonies for outplanting to Knob Reef, Hong Kong (**Fig. S14**).

The outplanted corals were monitored through September 2024 without further probiotic

intervention, with measurements taken at five intervals: **t1** (1 month post-outplanting, February),

t3 (April), **t6** (July), **t7** (August), and **t8** (September). Crucially, a natural thermal stress event

occurred during **t6**-**t8**, with degree heating weeks (DHW) reaching 8.8-18 ºC-weeks (**Fig. S15**),

surpassing the 4-8 DHW bleaching threshold. During the 8-month monitoring period, MC10-B4-

inoculated corals exhibited enhanced thermal tolerance, with significantly reduced signs of visual

bleaching at the peak of thermal stress relative to placebo-treated group (CoralWatch Coral

Health Chart coloration scores at **t6**: 3.3 ± 0.5 vs. 2.5 ± 0.4, $P < 0.001$; **t7**: 3.2 ± 0.4 vs. 2.4 ± 0.4,

$P < 0.001$; two-way ANOVA, Tukey's HSD; **Fig. 3E, Table S4**). Although partial recovery

occurred in the subsequent month, neither MC10-B4-inoculated corals nor placebo-treated group

fully regained pre-bleaching coloration levels by **t8**, indicating lingering stress effects.

Concurrently, F_v/F_m' from corals inoculated with MC10-B4 varied by treatment at some

timepoints ($P < 0.001$, two-way ANOVA, interaction effect), with probiotic-inoculated corals

maintaining significantly higher F_v/F_m' (**t6**: $P < 0.001$; **t7**: $P < 0.001$; two-way ANOVA, Tukey's

HSD; **Fig. 3F, Table S5**).

FISH assays with an MC10-specific probe at **t0**, **t6**, and **t8** confirmed MC10's persistent

presence in coral tissues, particularly the gastrodermis (**Fig. 3G**), with additional nonsense probe

confirming the hybridization specificity (**Fig. 3G**). MC10's low abundance within *Ruegeria*

communities (median 0.23%) and *Ruegeria*'s minor representation in coral microbiomes

(median 0.02%) precluded precise quantification, falling below the 0.5% detection threshold for

reliable amplicon sequence variant (ASV) resolution [30].

To monitor microbiome dynamics in outplanted corals, we performed *parC* and 16S rRNA

gene amplicon sequencing at **t0, t1, t3, t6**, and **t8** (**t7** samples unavailable; see Supplemental

Methods). Analysis of β-diversity (compositional differences between communities) revealed

that bacterial communities in MC10-B4-inoculated corals became progressively more similar to

those in placebo-treated corals over time (**Fig. 4A**; ANOSIM). This convergence may reflect

environmental acclimatization or dispersal-based homogenization at the reef site (groups

separated by only 40 meters). Despite this community-level similarity, differential enrichment of

individual taxa could be observed. During the bleaching phase (**t6**), ASV-based genus-level

analysis identified significant enrichment of several Bacteroidetes genera (*Cryomorpha*,

Spirosoma, *Sunxiuqinia*, *Gelatiniphilus*, *Aureispira*, and *Flavobacterium*) and depletion of

multiple Roseobacteraceae genera (*Sulfitobacter*, *Roseovarius*, *Pseudoruegeria*, *Donghicola*,

Aliiroseovarius, and *Phaeobacter*) in MC10-B4-inoculated corals compared to the placebo group

(FDR-corrected $P < 0.05$; ANCOM-BC; **Fig. 4B**). Crucially, most of these taxa showed no

enrichment or depletion during pre-bleaching (**t3**) or post-bleaching (**t8**) phases (**Fig. S16**),

indicating that MC10 may exert a microbial community restructuring effect during natural

thermal stress.

This trial demonstrates that nursery-integrated probiotic treatments can enhance coral

resilience to thermal stress without the need for post-outplanting intervention. MC10's sustained

retention, even at low abundances, aligns with its evolutionary adaptations (e.g.,

pseudogenization-driven host dependency) and underscores its potential as a scalable tool for

reef restoration.

Discussion

The accelerating loss of biodiversity under climate change demands innovative, scalable

strategies to enhance resilience across ecosystems. While probiotics have shown promise in

diverse systems, their scalability is universally hampered by transient colonization. Here, we

advance a generalizable solution: probiotics selected through evolutionary principles, targeting

bacteria undergoing irreversible transitions from free-living to host-dependent symbiosis. Using

coral reefs as a model system, we identify *Ruegeria* MC10, a coral-associated bacterial

population exhibiting genomic hallmarks of host adaptation. An 8-month field trial demonstrates

that nursery MC10 inoculations confer sustained retention in corals through a natural bleaching

event, enhancing thermal resilience without the need for repeated applications.

Evolutionary innovation in probiotic selection

Traditional probiotics, for marine organisms like corals, farmed shellfish and fish, as well as for

terrestrial wildlife and farmed species, are selected for transient functional traits such as

pathogen inhibition or nutrient provisioning [7,12,31]. However, uncertainty regarding their ability to

persist long-term in host tissues limits scalability [32]. Here, we introduce a lifestyle-oriented

strategy rooted in evolutionary genomics, shifting the paradigm from what probiotics do to how

they evolve, their evolutionary trajectory. By targeting bacterial symbionts undergoing

irreversible transitions from free-living to host-dependent symbiosis, we exploit their inherent

genomic trajectory toward stable host association, arguably due to their putatively important role

in holobiont functioning. This adaptation comes at a measurable and detectable genomic cost, as

relaxed purifying selection during the lifestyle transition leads to the accumulation of mildly

deleterious mutations, including IS expansion and pseudogene proliferation [33]. These genomic

hallmarks, such as frameshift mutations or IS-driven disruptions in essential genes, reflect

reduced selective pressure to maintain free-living metabolic pathways, driving metabolic

dependency that favors sustained colonization [33,34]. *Ruegeria* MC10 exemplifies this novel

approach, exhibiting hallmark signatures of these evolutionary young endosymbionts.

This strategy draws parallels with terrestrial, recently host-dependent endosymbionts, such

as *Sodalis glossinidius* in Tsetse flies, which exhibit similar genomic erosion during early host

adaptation [33,35]. However, while such terrestrial symbionts have not yet been repurposed as

probiotics for economically or ecologically critical animals, our work demonstrates the

feasibility of this approach in marine systems. MC10's success, retained in coral tissues for eight

months post-inoculation, validates evolutionary principles as a blueprint for probiotic design

across ecosystems. For example, host-dependent symbionts in oysters, shrimp, or even terrestrial

livestock could be harnessed to improve stress tolerance. Notably, at current we are naïve

regarding the measurable host beneficial effect of such isolates. However, given our success here,

one may argue it might be a rather more frequent than less occurrence.

By prioritizing evolutionary compatibility over inferred functional trait screening, this

framework addresses a universal limitation in microbiome engineering: the uncertainty regarding

long-term establishment of administered microbes [10]. For corals, MC10's persistence reduces

reliance on repeated applications, a critical advance for large-scale restoration. Beyond reefs, this

approach could transform probiotics for aquaculture, agriculture, and wildlife restoration, where

long-term microbial colonization is essential. The integration of evolutionary theory into

conservation practice marks a paradigm shift, enabling scalable solutions that align microbial

interventions with natural host-microbe coevolution.

Facultative endosymbionts as host-generalist probiotics

Ruegeria MC10 is a facultative endosymbiont, given its variable presence among colonies of the

same coral species and its presence as a free-living bacterium. Facultative endosymbionts

undergoing early host adaptation exhibit a unique duality: genomic erosion drives metabolic

dependency while retaining ecological flexibility. This paradox enables broad host ranges, as

exemplified by *Sodalis glossinidius*, which persists in Tsetse flies and other insects despite

pseudogenized metabolic pathways. Similarly, *Ruegeria* MC10's host versatility may arise from

mechanisms that exploit universal host interfaces rather than lineage-specific adaptations. For

instance, MC10's pseudogenized TCA cycle necessitates scavenging host-derived metabolites

such as succinate or fumarate, pathways conserved across eukaryotes. This strategy aligns with

resource tracking, a mechanism highlighted by Lajoie and Parfrey [36], where microbial symbionts

exploit conserved host resources (e.g., vitamins, metabolites) to colonize diverse hosts without

specialization.

MC10's colonization of the coral gastrodermis likely involves eukaryotic receptors (e.g.,

integrins or lectins), akin to *Lactiplantibacillus plantarum*, which colonizes *Drosophila* spp. and

many metazoan species through a conserved colonization island, allowing for wide use as a

human and animal probiotic [37]. This touches upon the concept of ecological fitting, where pre-

existing traits enable symbionts to colonize novel hosts without coevolution, a key driver of host

generalization in facultative symbioses [36]. Additionally, MC10's transient environmental

persistence reflects weak host selection pressures. As emphasized by Lajoie and Parfrey [36], when

hosts impose minimal filtering (e.g., due to low symbiont abundance or metabolic byproduct

exploitation), symbionts face reduced costs of adaptation to novel hosts, enabling niche

expansion. MC10's low relative abundance in coral microbiomes likely minimizes its burden on

the host, weakening selection for specialization and favoring persistence across taxa.

Scalability hinges on this generality. Unlike obligate symbionts restricted to single hosts [35],

facultative endosymbionts like MC10 exploit conserved molecular gateways, enabling nursery-

based priming of diverse corals.

Ruegeria MC10's growth dynamics balance evolutionary trade-offs and scalable cultivation

Growth rate comparisons between MC10 and its conventional *Ruegeria* relatives reveal that

MC10 exhibits reduced proliferation under laboratory conditions, a phenotype potentially linked

to pseudogenization of core metabolic pathways. Disrupted TCA cycle genes likely constrain

ATP synthesis, while pseudogenized amino acid biosynthesis pathways increase reliance on

externally sourced metabolites, collectively imposing energetic inefficiencies. Despite these

constraints, MC10 retains culturability, potentially through mechanisms like ribosomal

frameshifting, where translational machinery bypasses frameshift mutations, or codon

reassignment, where alternative decoding of stop codons permits partial gene expression. For

example, *Salmonella enterica* translates 101 of 106 pseudogenes [38], though the specific

mechanisms driving translation in this species remain untested. In general, such processes often

produce misfolded proteins due to destabilized residue interactions, leading to energetic

frustration and reduced functionality [39]. These molecular inefficiencies compound physiological

deficits (e.g., reduced ATP yield), creating a dual burden that favors MC10's dependency on

coral hosts for metabolic compensation. Thus, while pseudogene translation may sustain

transient culturability, its inherent instability reinforces evolutionary trajectories toward host

reliance, aligning genomic decay with ecological dependency.

Critically, MC10-B4's growth kinetics ($\mu_{max} = 0.15\ \text{h}^{-1}$) align with established probiotics

like *Lactobacillus* spp. (0.12-0.23 h^{-1}) and marine probiotic *Phaeobacter inhibens* (0.16-0.18 h^{-1}).

Its stationary-phase yields ($7.4 \times 10^8\ \text{CFU/mL}$) approach commercial probiotic standards

(typically $\geq 10^9$ CFU/dose in pharmaceutical applications [40]), validating scalability for reef

restoration. Industrial fermentation processes achieve high yields through tailored nutrient-rich

media (e.g., whey-supplemented formulations for *Lactobacillus*) and advanced fermentation

engineering (e.g., fed-batch cultivation), overcoming growth limitations in fastidious strains [41].

For MC10-B4, tailored media formulations, such as coral-mimicking nutrient formulations,

could further enhance biomass production by compensating for pseudogenization-driven

metabolic deficiencies. This industrial compatibility, coupled with MC10-B4's high cell density

output, positions it as a tractable candidate for large-scale marine probiotic production.

Functional mechanisms underlying Ruegeria MC10's probiotic potential

Functional mechanisms underlying Ruegeria MC10's probiotic potential

Ruegeria MC10's benefits under thermal stress, including reduced algal photosynthetic

efficiency loss and reduced bleaching, align with functional roles observed in terrestrial

facultative endosymbionts. For example, *Serratia symbiotica* buffers its insect hosts against heat

stress via metabolite exchange [42]. Though MC10's exact mechanisms remain unresolved, its

retention in gastrodermis and pseudogenized metabolism suggest analogous stress-mitigating

roles, potentially through stabilizing algal symbiont function or modulating host physiology. A

key debate centers on whether probiotics act as nutritional sources to support host energetic

demands [7,32,43]. MC10's sustained colonization over eight months argues against digestion-driven

benefits, as degraded cells would not persist. Future experiments may leverage transcriptomics

and metabolomics to resolve whether MC10 stabilizes algal symbiont function, modulates host

immunity, or facilitates metabolite exchange during thermal stress.

Ecological flexibility and safety

While *Ruegeria* MC10's ecological flexibility, demonstrated by its cross-species efficacy in

enhancing thermal tolerance across Hexacorallia, could raise concerns about off-target

colonization, multiple factors converge that argue for its ecological safety. MC10 occurs

naturally at low abundances (0.5-2% of coral-associated *Ruegeria* communities, where detected)

across diverse hosts including faviid (*O. crispata*), acroporid (*A. solitaryensis*), and poritid

(*Porites* sp.) corals, as well as in seawater and sediment. This widespread but sparse distribution

may be due to constrained environmental fitness resulting from pseudogenized metabolism (e.g.,

disrupted TCA cycle), which may intrinsically limit competitive proliferation outside host tissues.

Our results also suggest that microbiome restructuring is fine-scale and restricted to context-

specific depletion of certain Roseobacteraceae genera during bleaching events, which may

suggest transient niche restructuring. Of note, MC10's evolutionary trajectory toward host

dependency manifests in tissue-specific localization (gastrodermis) and ultralow persistence

abundances (<0.005% total bacteria), reducing environmental shedding risks. Collectively, these

attributes, encompassing natural rarity, self-limiting metabolic constraints, and host-adapted

persistence, position MC10 as a coral "core microbiome" taxon: a low-abundance generalist that

supports holobiont function without ex hospite ecosystem disruption, making it ideally suited for

multi-species restoration [44].

Microbiome dynamics and stability

A central goal of probiotic intervention aligns with microbial transfer therapy (MTT) in human

medical settings: introducing beneficial taxa to reset dysbiotic microbiomes and enhance

resilience through community restructuring [21,45]. By leveraging *Ruegeria* MC10 as a keystone

inoculant, we achieved microbiome modulation without long-term destabilization, evidenced by

transient β-diversity differences that peaked post-inoculation but converged with placebo-treated

corals over eight months. Crucially, during bleaching stress (**t6**), MC10-B4 association co-

occurred with a functional microbiome reset characterized by enrichment of Bacteroidetes

polysaccharide degradation specialists (*Cryomorpha*, *Spirosoma*, *Sunxiuqinia*, *Gelatiniphilus*,

Aureispira, *Flavobacterium*) [46], ROS-tolerant taxa (*Deinococcus*, *Desertibacter*, *Spirosoma*) [47],

Vibrio-predatory specialist (*Aureispira*) [48], and known probiotics (*Veillonella*,

Limosilactobacillus) [49,50]. Concurrent depletion of multiple Roseobacteraceae genera

(*Sulfitobacter*, *Roseovarius*, *Pseudoruegeria*, *Donghicola*, *Aliiroseovarius*, *Phaeobacter*)

indicated competitive restructuring within this ecologically critical family, notably including

both MC10 itself and the established probiotics *Phaeobacter* [51]). This taxonomic shift, largely

absent pre- or post-bleaching, argues MC10 primed the microbiome for stress-responsive

optimization.

While algal symbiont switching (e.g., from sensitive *Cladocopium* to resilient *Durusdinium*)

represents a known thermal adaptation pathway in some corals [52], this mechanism is unlikely to

explain our results. Regional characterization confirms Hong Kong corals, including *Acropora*

spp., exclusively harbor *Cladocopium* spp. (except *O. crispata*) [53]. This consistency across host

genotypes and sites makes algal symbiont-mediated thermal tolerance gains as an alternative

explanation for MC10-B4's efficacy in inoculated *A. pruinosa* unlikely.

Next-generation probiotics as a scalable solution

Coral restoration typically involves growing colony fragments in nurseries before outplanting

[54,55], a framework highly compatible with integrating next-generation probiotics like Ruegeria

MC10 as a nursery-applied treatment. Our 8-month field trial provides a proof of concept that

evolutionarily selected probiotics can achieve sustained retention and enhance stress tolerance

evading the need for post-outplanting intervention. However, critical knowledge gaps remain for

evaluating long-term restoration success, an issue plaguing coral restoration initiatives globally [55].

Multi-year data on coral survival, reproductive fitness, and reef-scale outcomes (e.g.,

biodiversity recovery, structural complexity) are essential next steps to validate this approach.

Notwithstanding these open questions, the evolutionary design of next-generation probiotics

offers a uniquely promising pathway for scalability: host dependency minimizes post-outplanting

dispersal risks, while transient environmental viability and cross-species efficacy enable efficient

"probiotic priming" within existing nursery workflows. Given the accelerating pace of ocean

warming, which increasingly outstrips natural adaptation [56], such scalable interventions represent

one of the most viable options to bridge the resilience gap: the widening disparity between

organismal stress tolerance and environmental insult magnitude that exceeds compensatory

capacities of acclimation or adaptation.

By aligning mechanistic research with scalable delivery systems, the here-presented

approach offers a blueprint to explore microbial-driven resilience across ecosystems. Further

validation of long-term ecological impacts and industrial-scale production will be crucial, but the

urgency of the coral crisis demands parallel exploration of multiple strategies, among which

evolution-guided probiotics stand as a compelling, scalable candidate.

Acknowledgements: We thank all the colleagues who have supported experiments on outplant

fieldwork and FISH experiments, including Jingqi Gao and Kai Yan Lai. We would like to thank

Kristin Bär-Häge and Myriam Schmid for their support with Aiptasia animal husbandry for

CBASS acute thermal assays.

Funding: Hong Kong Research Grants Council (RGC) General Research Fund project #:

14114724 granted to H.L. Grand Challenges Scheme project #: 4720293 by the CUHK-Exeter

Joint Centre for Environmental Sustainability and Resilience (ENSURE) granted to H.L.

"Reviving Our Corals" initiative agreement #: TW2318074 by WWF Hong Kong and Coral

Academy, The Chinese University of Hong Kong granted to A.P.Y.C. German Research

Foundation (DFG) project #: 458901010 granted to C.R.V.

Author Contributions: Conceptualization: H.L. Methodology on *Ruegeria* isolation, genome

analysis, and population-level primer design: M.X., T.L., H.L. Methodology on CBASS and

coral outplant : N.X., C.R.V., C.T.C., M.X., A.P.Y.C., H.L. Code development: T.L., X.W.

Investigation on coral sampling and *Ruegeria* isolation: M.X., K.E.H., D.L., S.E.M., R.H., E.C.,

Z.W., P.T., J.B. Investigation on CBASS: N.X., M.D., H.M., C.R.V. Investigation on coral

outplant: C.T.C., M.X., N.X., A.P.Y.C., C.H.L, W.Y.T, K.K.T., Q.H., P.L. Methodology on

FISH: Z.X., G.C. Data visualization and statistical analysis: N.X., S.E.M. Funding acquisition:

H.L., A.P.Y.C. Project administration: H.L. Supervision: H.L. and C.R.V. Writing – original

draft: H.L. Writing – review and editing: H.L., C.R.V., N.X., M.X., S.E.M., R.S.P., G.C. Writing

– supplementary materials: M.X., N.X., T.L., H.L.

Competing interests: The authors declare that they have no competing interests.

Data and materials availability: Raw sequencing data are available on NCBI under several

BioProjects with the private access link for reviewers. The raw reads of 409 and assembled

genomes of 419 *Ruegeria* isolates (10 genomes missing raw data) sequenced on the second-

generation platform (DNBseq) and 34 *Ruegeria* isolates sequenced on the third-generation

platform (Nanopore) are available under NCBI BioProject accessions PRJNA1264799

(https://dataview.ncbi.nlm.nih.gov/object/PRJNA1264799?reviewer=mp7rulknapinfb95c9733nf

72r) and PRJNA1275854

(https://dataview.ncbi.nlm.nih.gov/object/PRJNA1275854?reviewer=g4mffnb8b965tq6b2662cpt

6i8), respectively. The *parC* and *ATP5B* amplicon sequencing data for natural coral samples are

available under PRJNA1275610

(https://dataview.ncbi.nlm.nih.gov/object/PRJNA1275610?reviewer=nst2vifa08hqr8otcjvv39ge4

l) and PRJNA1275576

(https://dataview.ncbi.nlm.nih.gov/object/PRJNA1275576?reviewer=g865emc830spbkko9b86qt

cc35), respectively. The 16S rRNA and *parC* amplicon sequencing data for outplanted corals are

available under PRJNA1275585

(https://dataview.ncbi.nlm.nih.gov/object/PRJNA1275585?reviewer=9ne88342f7at8ndur1klq05

mo8) and PRJNA1275617

(https://dataview.ncbi.nlm.nih.gov/object/PRJNA1275617?reviewer=e6g3aatk7pg8r6m6ao0riovj

nu), respectively. All data will be publicly accessible upon publication. The scripts used for

genome assembly and annotations, population genomic analysis, amplicon data analysis, and

pseudogene analysis have been deposited in an online repository

(https://github.com/444thLiao/CoralProbiotic).

446
References

1   Fisher, R. *et al.* Species richness on coral reefs and the pursuit of convergent global

estimates. *Current Biology* **25**, 500-505, doi:https://doi.org/10.1016/j.cub.2014.12.022

(2015).

2   Souter, D. *et al.* Status of coral reefs of the world: 2020. *Global Coral Reef Monitoring*

Network (GCRMN) and International Coral Reef Initiative (ICRI). (2021).

3   Hughes, T. P. *et al.* Global warming transforms coral reef assemblages. *Nature* **556**, 492-

496 (2018).

4   IPCC. Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and

III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change.

35–115 (Geneva, Switzerland, 2023).

5   Delgadillo-Ordoñez, N. *et al.* Probiotics reshape the coral microbiome in situ without

detectable off-target effects in the surrounding environment. *Communications Biology* **7**,

434, doi:10.1038/s42003-024-06135-3 (2024).

6   Doering, T., Maire, J., van Oppen, M. J. & Blackall, L. L. Advancing coral microbiome

manipulation to build long-term climate resilience. *Microbiology Australia* **44**, 36-40

(2023).

7   Garcias-Bonet, N. *et al.* Horizon scanning the application of probiotics for wildlife.

Trends in Microbiology **32**, 252-269, doi:https://doi.org/10.1016/j.tim.2023.08.012

(2023).

8   Santoro, E. P. *et al.* Coral microbiome manipulation elicits metabolic and genetic

restructuring to mitigate heat stress and evade mortality. *Science Advances* **7**, eabg3088,

doi:doi:10.1126/sciadv.abg3088 (2021).

9   Thatcher, C., Høj, L. & Bourne, D. G. Probiotics for coral aquaculture: challenges and

considerations. *Current Opinion in Biotechnology* **73**, 380-386 (2022).

10   Albright, M. B. N. *et al.* Solutions in microbiome engineering: prioritizing barriers to

organism establishment. *The ISME Journal* **16**, 331-338, doi:10.1038/s41396-021-01088-

5 (2022).

11   Čaušević, S. *et al.* Niche availability and competitive loss by facilitation control

proliferation of bacterial strains intended for soil microbiome interventions. *Nature*

Communications **15**, 2557, doi:10.1038/s41467-024-46933-1 (2024).

12   Chauhan, A. & Singh, R. Probiotics in aquaculture: a promising emerging alternative

approach. *Symbiosis* **77**, 99-113, doi:10.1007/s13199-018-0580-1 (2019).

13   Peixoto, R. S. *et al.* Coral probiotics: premise, promise, prospects. *Annual review of*

animal biosciences **9**, 265-288 (2021).

14   Kittinger, J. N., Finkbeiner, E. M., Glazier, E. W. & Crowder, L. B. Human dimensions

of coral reef social-ecological systems. *Ecology and Society* **17**, 17, doi:10.5751/ES-

05115-170417 (2012).

15   Rädecker, N. *et al.* Heat stress destabilizes symbiotic nutrient cycling in corals.

Proceedings of the National Academy of Sciences **118**, e2022653118,

doi:doi:10.1073/pnas.2022653118 (2021).

16   Eakin, C. M. *et al.* The 2014-17 global coral bleaching event: the most severe and

widespread coral reef destruction. *Preprint* (2022).

17   Reimer, J. D. *et al.* The fourth global coral bleaching event: Where do we go from here?

Coral Reefs **43**, 1121-1125, doi:10.1007/s00338-024-02504-w (2024).

18   Ushijima, B. *et al.* Chemical and genomic characterization of a potential probiotic

treatment for stony coral tissue loss disease. *Communications Biology* **6**, 248,

doi:10.1038/s42003-023-04590-y (2023).

19   Moradi, M. *et al.* Probiotics mitigate thermal stress- and pathogen-driven impacts on

coral skeleton. *Frontiers in Marine Science* **10**, doi:10.3389/fmars.2023.1212690 (2023).

20   Huggett, M. J. & Apprill, A. Coral microbiome database: Integration of sequences

reveals high diversity and relatedness of coral-associated microbes. *Environ. Microbiol.*

Rep. **11**, 372-385, doi:10.1111/1758-2229.12686 (2019).

21   Voolstra, C. R. *et al.* The coral microbiome in sickness, in health and in a changing world.

Nature Reviews Microbiology **22**, 460-475, doi:10.1038/s41579-024-01015-3 (2024).

22   Xu, M., Cai, Z., Cheng, K., Chen, G. & Zhou, J. Mitigation of *Vibrio coralliilyticus*-

induced coral bleaching through bacterial dysbiosis prevention by *Ruegeria profundi*.

Applied and Environmental Microbiology **90**, e02274-02223,

doi:doi:10.1128/aem.02274-23 (2024).

23   Arevalo, P., VanInsberghe, D., Elsherbini, J., Gore, J. & Polz, M. F. A reverse ecology

approach based on a biological definition of microbial populations. *Cell* **178**, 820-

834.e814 (2019).

24   Moran, M. A. *et al.* Genome sequence of *Silicibacter pomeroyi* reveals adaptations to the

marine environment. *Nature* **432**, 910-913, doi:10.1038/nature03170 (2004).

25   Evensen, N. R. *et al.* The Coral Bleaching Automated Stress System (CBASS): A low-

cost, portable system for standardized empirical assessments of coral thermal limits.

Limnology and Oceanography: Methods **21**, 421-434,

doi:https://doi.org/10.1002/lom3.10555 (2023).

26   Voolstra, C. R. *et al.* Standardized short-term acute heat stress assays resolve historical

516   differences in coral thermotolerance across microhabitat reef sites. *Global Change*

517   *Biology* **26**, 4328-4343, doi:https://doi.org/10.1111/gcb.15148 (2020).

518   27   Baumgarten, S. *et al.* The genome of *Aiptasia*, a sea anemone model for coral symbiosis.

519   *Proceedings of the National Academy of Sciences* **112**, 11893-11898,

520   doi:doi:10.1073/pnas.1513318112 (2015).

521   28   Dörr, M. *et al.* Short-term heat stress assays resolve effects of host strain, repeat stress,

522   and bacterial inoculation on Aiptasia thermal tolerance phenotypes. *Coral Reefs*,

523   doi:10.1007/s00338-023-02427-y (2023).

524   29   Voolstra, C. R. *et al.* Standardized methods to assess the impacts of thermal stress on

525   coral reef marine life. *Ann Rev Mar Sci* **17**, 193-226, doi:10.1146/annurev-marine-

526   032223-024511 (2025).

527   30   Callahan, B. J. *et al.* High-throughput amplicon sequencing of the full-length 16S rRNA

528   gene with single-nucleotide resolution. *Nucleic Acids Research* **47**, e103-e103,

529   doi:10.1093/nar/gkz569 (2019).

530   31   Peixoto, R. S. *et al.* Coral probiotics: premise, promise, prospects. *Annual Review of*

531   *Animal Biosciences* **9**, 265-288, doi:10.1146/annurev-animal-090120-115444 (2021).

532   32   Doering, T., Maire, J., van Oppen, M. J. H. & Blackall, L. L. Advancing coral

533   microbiome manipulation to build long-term climate resilience. *Microbiology Australia*

534   **44**, 36-40, doi:https://doi.org/10.1071/MA23009 (2023).

33 Moran, N. A. & Plague, G. R. Genomic changes following host restriction in bacteria. *Current Opinion in Genetics & Development* **14**, 627-633, doi:10.1016/j.gde.2004.09.003 (2004).

34 Toft, C. & Andersson, S. G. E. Evolutionary microbial genomics: insights into bacterial host adaptation. *Nature Reviews Genetics* **11**, 465-475, doi:10.1038/nrg2798 (2010).

35 Perreau, J. & Moran, N. A. Genetic innovations in animal–microbe symbioses. *Nature Reviews Genetics* **23**, 23-39, doi:10.1038/s41576-021-00395-z (2022).

36 Lajoie, G. & Parfrey, L. W. Beyond specialization: re-examining routes of host influence on symbiont evolution. *Trends in Ecology & Evolution* **37**, 590-598, doi:https://doi.org/10.1016/j.tree.2022.03.006 (2022).

37 Gutiérrez-García, K. *et al.* A conserved bacterial genetic basis for commensal-host specificity. *Science* **386**, 1117-1122, doi:doi:10.1126/science.adp7748 (2024).

38 Feng, Y. *et al.* "Pseudo-pseudogenes" in bacterial genomes: Proteogenomics reveals a wide but low protein expression of pseudogenes in *Salmonella enterica*. *Nucleic Acids Research* **50**, 5158-5170, doi:10.1093/nar/gkac302 (2022).

39 Jaafari, H. *et al.* The physical and evolutionary energy landscapes of devolved protein sequences corresponding to pseudogenes. *Proceedings of the National Academy of Sciences* **121**, e2322428121, doi:doi:10.1073/pnas.2322428121 (2024).

40 Bertazzoni, E., Gianfranco, D., Tore, M., Jacques, N. & and Sanz, Y. Probiotics and clinical effects: is the number what counts? *Journal of Chemotherapy* **25**, 193-212, doi:10.1179/1973947813Y.0000000078 (2013).

41 Aguirre-Ezkauriatza, E. J., Aguilar-Yáñez, J. M., Ramírez-Medrano, A. & Alvarez, M. M.

Production of probiotic biomass (*Lactobacillus casei*) in goat milk whey: Comparison of

batch, continuous and fed-batch cultures. *Bioresource Technology* **101**, 2837-2844,

doi:https://doi.org/10.1016/j.biortech.2009.10.047 (2010).

42 Burke, G., Fiehn, O. & Moran, N. Effects of facultative symbionts and heat stress on the

metabolome of pea aphids. *The ISME Journal* **4**, 242-252, doi:10.1038/ismej.2009.114

(2009).

43 Morgans, C. A., Hung, J. Y., Bourne, D. G. & Quigley, K. M. Symbiodiniaceae

probiotics for use in bleaching recovery. *Restoration Ecology* **28**, 282-288,

doi:https://doi.org/10.1111/rec.13069 (2020).

44 van Oppen, M. J. H. & Blackall, L. L. Coral microbiome dynamics, functions and design

in a changing world. *Nature Reviews Microbiology* **17**, 557-567, doi:10.1038/s41579-

019-0223-4 (2019).

45 Kang, D.-W. *et al.* Long-term benefit of Microbiota Transfer Therapy on autism

symptoms and gut microbiota. *Scientific Reports* **9**, 5821, doi:10.1038/s41598-019-

42183-0 (2019).

46 McKee, L. S. *et al.* Polysaccharide degradation by the Bacteroidetes: mechanisms and

nomenclature. *Environmental Microbiology Reports* **13**, 559-581,

doi:https://doi.org/10.1111/1758-2229.12980 (2021).

47 Pavlopoulou, A. *et al.* Unraveling the mechanisms of extreme radioresistance in

prokaryotes: Lessons from nature. *Mutation Research/Reviews in Mutation Research* **767**,

92-107, doi:https://doi.org/10.1016/j.mrrev.2015.10.001 (2016).

48 Lien, Y.-W. *et al.* Mechanism of bacterial predation via ixotrophy. *Science* **386**,

eadp0614, doi:doi:10.1126/science.adp0614 (2024).

49 Ksiezarek, M., Grosso, F., Ribeiro, T. G. & Peixe, L. Genomic diversity of genus

Limosilactobacillus. *Microb Genom* **8**, mgen000847, doi:10.1099/mgen.0.000847 (2022).

50 Zhang, S. M., Hung, J. H., Yen, T. N. & Huang, S. L. Mutualistic interactions of lactate-

producing lactobacilli and lactate-utilizing Veillonella dispar: Lactate and glutamate

cross-feeding for the enhanced growth and short-chain fatty acid production. *Microb*

Biotechnol **17**, e14484, doi:10.1111/1751-7915.14484 (2024).

51 Sonnenschein, E. C., Jimenez, G., Castex, M. & Gram, L. The *Roseobacter*-group

bacterium *Phaeobacter* as a safe probiotic solution for aquaculture. *Applied and*

Environmental Microbiology **87**, e02581-02520, doi:doi:10.1128/AEM.02581-20 (2021).

52 Quigley, K. M. *et al.* Symbioses are restructured by repeated mass coral bleaching.

Science Advances **8**, eabq8349, doi:doi:10.1126/sciadv.abq8349 (2022).

53 Saad, O. S. *et al.* Species richness and generalists–specialists mosaicism of

symbiodiniacean symbionts in corals from Hong Kong revealed by high-throughput ITS

sequencing. *Coral Reefs* **41**, 1-12, doi:10.1007/s00338-021-02196-6 (2022).

54   Suggett, D. J. & van Oppen, M. J. H. Horizon scan of rapidly advancing coral restoration

approaches for 21st century reef management. *Emerging Topics in Life Sciences* **6**, 125-

136, doi:10.1042/etls20210240 (2022).

55   Boström-Einarsson, L. *et al.* Coral restoration – A systematic review of current methods,

successes, failures and future directions. *PLOS ONE* **15**, e0226631,

doi:10.1371/journal.pone.0226631 (2020).

56   Lachs, L. *et al.* Natural selection could determine whether Acropora corals persist under

expected climate change. *Science* **386**, 1289-1294 (2024).

   Figure Legends

Fig. 1 *Ruegeria* MC10's phylogenetic position and genomic signatures showing recent host

dependency. (A) Rooted maximum-likelihood phylogenomic tree comprising 419 *Ruegeria*

isolates associated with five coral species sampled from nine Hong Kong reef sites, along with

26 *Ruegeria* genomes retrieved from public databases. This tree was constructed with IQ-TREE

based on concatenated single-copy orthologous genes at the amino acid level. PopCOGenT was

used to delineate the *Ruegeria* populations (i.e., main clusters or MCs). The 19 MCs that contain

representative members additionally sequenced by Nanopore were labelled with numbers in the

tree. From the inner to outer rings: (1) denotes 29 *Ruegeria* populations which consists of at least

three isolates as defined by PopCOGenT, along with MC46, the sister group of MC10; (2) - (5)

denote the source of *Ruegeria* isolates with each corresponding to coral species, sampling site,

coral compartment, and coral colony replicate, respectively. (**B)** Stacked bars showing the

number of IS and IS family composition in 34 newly closed *Ruegeria* genomes and the reference

genome *Ruegeria pomeroyi* DSS-3. (**C)** Number of pseudogenes in 34 newly closed *Ruegeria*

genomes and the reference genome *R. pomeroyi* DSS-3. Pseudogenes were grouped by the

mechanism of formation, namely "IS disruption", "Frameshift mutations", "Nonsense mutations",

and "Singletons". "Unclassified" pseudogenes are those that cannot be assigned to any of the

three defined mechanisms.

Fig. 2 Relative abundance of *Ruegeria* MC10 in coral holobionts and environmental

samples from Hong Kong reefs. (**A)** The relative abundance of MC10 in *Ruegeria* communities

was quantified based on population-resolving *ATP5B* and *parC* amplicon sequencing across

natural reefs in Hong Kong. Different bar plots under the same coral genus represented for

different coral colonies (i.e., 15 colonies for *Oulastrea* sp., 11 for *Platygyra* sp., 10 colonies for

Porites sp., 4 colonies for *Acropora* sp., and 4 colonies for *Plesiastrea* sp. across five sites). Each

bar plot was displayed with the mean value and standard deviation (SD) calculated from three

compartments (i.e., mucus, tissue and skeleton) for the same coral colony. (**B**) The relative

abundance of MC10 in seawater and sediment samples across different locations. The error bar

denotes standard deviation among three replicates. CI, Crescent Island; SI, Sharp Island; WP,

Wu Pai. Sd, sediment; Sw, seawater.

Fig. 3 Application of *Ruegeria* MC10-B4 through laboratory and field trials. (**A**) ED50

values of Aiptasia H2 inoculated with *Ruegeria* MC10-B4 or placebo FSW after heat stress (T7)

and post-stress recovery (T18). Data were presented with the mean values and standard errors. (**B**)

ED50 values modeled based on log-logistic regression curves of dark-adapted photosynthetic

efficiency (F_v/F_m) across experimental temperatures. F_v/F_m was measured at 7 h and 18 h of the

CBASS with one-hour dark acclimation. Treatment-specific ED50 values were denoted as

vertical lines. The relative abundance of (**C**) MC10 and (**D**) MC15 in the *Ruegeria* (left axis;

based on *parC* amplicon) and total bacteria community (right axis; combined 16S rRNA and

parC gene amplicon) following twice-weekly inoculations over a month. The central line of each

box indicates the median value with the upper and lower error bar showing maximum and

minimum potential variabilities, respectively. (**E**) Coral phenotype and coloration, and (**F**)

effective quantum yield (F_v/F_m') were assessed on MC10-B4-inoculated and placebo-treated

corals across five timepoints: 1, 3, 6, 7 and 8 months post-outplanting (**t1**, **t3**, **t6**, **t7**, **t8**,

respectively). Data were presented with the mean values and standard errors. (**G**) Localization of

MC10 in coral *A. pruinosa* after one-month inoculations (**t0**, before outplanting), six months

post-outplanting (**t6**, during bleaching), and eight months post-outplanting (**t8**, post-bleaching).

Representative confocal images depicting the hybridization of high-resolution *parC* probe

specifically targeting MC10 labeled with Cy3 (purple). The purple arrows pointed at the specific

location of MC10 in the coral gastrodermis. The blue arrows pointed at the non-specific binding

sites. Grey, DAPI-stained dsDNA; Green, autofluorescence of Symbiodiniaceae; Scale bars, 10

μm. All significant differences were indicated in bold and marked with asterisks (***: $P < 0.001$,

**: $P < 0.01$; *: $P < 0.05$; n.s.: non-significant).

Fig. 4 Coral microbiome dynamics during field trial. (**A**) Principal Coordinates Analysis

(PCoA) of coral holobiont-associated bacterial communities based on Bray-Curtis dissimilarity,

comparing placebo and MC10-B4-treated groups across five timepoints: baseline (**t0**, pre-

outplanting) and 1, 3, 6, and 8 months post-outplanting (**t1**, **t3**, **t6**, **t8**, respectively). Ellipses

represent 95% confidence intervals for group centroids. ANOSIM was used to assess the

separation of the placebo and MC10-B4-treated groups, with R values interpreted as: little to no

($R < 0.25$), weak ($0.25 \leq R < 0.5$), moderate ($0.5 \leq R < 0.75$), and strong ($R \geq 0.75$) separation.

(**B**) Differentially abundant taxa between placebo and MC10-B4-treated corals at t6 (during

bleaching) identified by ANCOM-BC analysis ($P < 0.05$, FDR corrected). A heatmap displaying

top 20 significantly upregulated (positive log2FC) and downregulated (negative log2FC) genera,

respectively. A log2-transformed read counts (log2[reads+1]) were shown to indicate the

absolute fold-change magnitudes of taxa. Pseudocounts were added to handle the case of zeros in

the calculation.

A From inner to outer ring

(1) *Ruegeria* MCs (in clockwise order)

MC14, MC27, MC9, MC0, MC49, MC18, MC10, MC50, MC15, MC1, MC11s3, MC5, MC13, MC26, MC21, MC2s2, MC19, MC25, MC61, MC3, MC22, MC56, MC6, MC16, MC45, MC46, MC23, MC44, MC17, MC7

(2) Coral species

Acropora cf. *samoensis*, *Acropora solitaryensis*, *Favites pentagona*, *Platygyra acuta*, *Oulastrea crispata*

(3) Sampling site

Kiu Tsui Chau (KTC), Ninepin Islands (NP), Wong Wan Chau (WWC), Chek Chau (CC), Ngo Mei Chau (NMC), Bluff Island (BI), Lo Chau (LC), Sham Wan (SW), Yam Tsai Wan (YTW)

(4) Coral compartment

mucus, tissue, skeleton

(5) Coral colony replicate

1, 2

tree scale: 0.1

B   **IS families**

IS200/IS605, Unknown, IS1634, IS6, IS4, ISNCY, IS630, IS110, IS30, IS66, IS21, IS5, IS1380, IS481, IS256, IS3, IS1595, IS91, IS1182

Number of IS

C   **Types of pseudogenes**

IS disruption, Frameshift mutations, Nonsense mutations, Singletons, Unclassified

Number of pseudogene

R. pomeroyi DSS-3 (MC1), AL3 (MC1), BG7 (MC15), AL7 (MC3), AF3 (MC7), AO1 (MC44), H4 (MC19), AJ12 (MC46), AK1 (MC46), AB1 (MC10), AA12 (MC10), AC11 (MC10), AJ1 (MC10), AJ9 (MC10), B4 (MC10), E4 (MC10), B2 (MC10), B7 (MC10), D9 (MC10), E12 (MC10), F6 (MC10), F7 (MC10), B1 (MC10), E9 (MC10), BB10 (MC2s2), AM7 (MC45), BK1 (MC21), AN11 (MC26), AP11 (MC13), AH3 (MC9), G7 (MC5), A5 (MC0), AB11 (MC6), BC8 (MC11s3), BJ3 (MC14)

MC10

A

Coral genus

Oulastrea (Oul)

Platygyra (Platy)

Porites (Pori)

Acropora (Acro)

Plesiastrea (Plesi)

Marker gene

ATP5B

parC

Wu Pai

Crescent Island

Port Island

Sharp Island

Ninepin Islands

Lo Chau

Sham Wan

B

Relative abundance of MC10 among *Ruegeria* community

A

After heat stress (T7h)

After post-stress recovery (T18h)

ED50 (°C)

Placebo   MC10   **

Placebo   MC10   *

B

Dark-adapted photosynthetic efficiency (Fv/Fm)

37.0   38.0

36.6   37.2

Treatment   ● Placebo   ▲ MC10

C

MC10 relative abundance within the *Ruegeria* community (%)

t0   **

MC10 relative abundance within the bacteria community (%)

t0   **

Treatment   ■ Placebo   ◻ MC10

D

MC15 relative abundance within the *Ruegeria* community (%)

t0   n.s.

MC15 relative abundance within the bacteria community (%)

t0   n.s.

Treatment   ● Placebo   ● MC15

E

Natural heat stress

Color score

n.s.   n.s.   ***   ***   n.s.

F

Fv/Fm'

n.s.   n.s.   ***   ***   n.s.

t1(Feb 22)   t3(Apr 26)   t6(Jul 17)   t7(Aug 9)   t8(Sep 2)

Treatment   ● Placebo   ● MC10

G

Treatment   Placebo   MC10   MC10

Probe   Sense   Sense   Nonsense

t0

t6

t8