Role of Atg1 in morphologic changes of the pathogenic fungus *Trichosporon asahii*

Mei Nakayama[1], Yasuhiko Matsumoto[1]*, Sanae Kurakado[1], Takashi Sugita[1]

[1]Department of Microbiology, Meiji Pharmaceutical University, 2-522-1, Noshio, Kiyose, Tokyo

204-8588, Japan.

*Corresponding author

E-mail: ymatsumoto@my-pharm.ac.jp (YM)

Short title: Atg1 and morphology of *T. asahii*

Abstract

Trichosporon asahii is a dimorphic fungus that causes severe invasive fungal infections, particularly in patients with neutropenia. Depending on nutrient availability, *T. asahii* exists in yeast, hyphae, or arthroconidia forms. Autophagy, a cellular degradation pathway responsible for removing old or damaged organelles, is essential for the survival of *Saccharomyces cerevisiae* under nutrient-limited conditions. Atg1 is a key regulator of the early phases of autophagy, especially under nitrogen starvation. The role of Atg1 in regulating morphology, stress resistance, or virulence in *T. asahii*, however, remains poorly understood. Here, we generated three *atg1* gene-deficient *T. asahii* mutants and investigated their phenotypic characteristics to reveal the role of Atg1 in *T. asahii*. The *atg1* gene-deficient mutants exhibited no growth defects under high-temperature or various chemical stress conditions, including antifungal drugs. Under a nutrient-rich culture condition, the mutants exhibited an increased proportion of hyphal cells. On the other hand, no morphologic differences were observed between the parent strain and the *atg1* gene-deficient mutants under a nitrogen-limited condition. The virulence of these *atg1* gene-deficient mutants was maintained in a silkworm infection model. Furthermore, all three generated *atg1* gene-deficient mutants exhibited consistent phenotypes. Our findings suggest that while Atg1 does not play a major role in stress tolerance or virulence in *T. asahii*, it plays a role in regulating its dimorphic morphologic changes.

Keywords: *Trichosporon asahii*, Atg1, morphology, virulence

Introduction

Trichosporon asahii is a dimorphic fungus that exists in yeast, hyphal, and arthroconidia forms [1]. *T. asahii* causes severe invasive fungal infections, particularly in immunocompromised hosts, such as those with neutropenia [2–4]. While the mortality rate of invasive fungal infections caused by the pathogenic fungus *Candida albicans* is approximately 40%, that caused by *T. asahii* is as high as 70%–80% [5]. Furthermore, *T. asahii* exhibits tolerance to echinocandin antifungal agents commonly used in clinical settings, often leading to breakthrough infections in patients receiving micafungin [6]. The ability of *T. asahii* to form biofilms on medical device surfaces, such as catheters, further complicates treatment by conferring resistance to various antifungal drugs and contributing to chronic, treatment-resistant infections [7]. These characteristics of *T. asahii* emphasize its clinical significance as a pathogenic fungus.

Autophagy is a conserved cellular process that maintains cellular homeostasis by degrading and recycling cytoplasmic components, including damaged organelles and misfolded proteins [8,9]. Autophagy is regulated by signaling pathways such as mTOR and AMPK and is activated under stress conditions such as nutrient deprivation [10,11]. In the model eukaryote *Saccharomyces cerevisiae*, autophagy plays a critical role in survival under nutrient-limited conditions [12,13]. The process is initiated in response to nutrient deprivation, with Atg1 acting as a key regulator in its early stages [8,14]. Atg1 and other autophagy-related proteins are essential for the progression of autophagy and are required for stress resistance and pathogenicity in *Cryptococcus neoformans*, a basidiomycete closely related to *T. asahii* [15,16]. In *Candida albicans*, another dimorphic fungus, Atg1 is required for biofilm formation [17,18]. The role of the autophagy-related protein Atg1 in morphologic transitions, stress resistance, and virulence in *T. asahii*, however, remains unclear.

57   In the present study, we independently generated three *atg1* gene-deficient mutants of *T.*

58 *asahii* and evaluated their morphology, stress sensitivity, and virulence. Our results revealed that

59 *atg1* gene deficiency promotes hyphal formation under nutrient-rich conditions but does not affect

60 growth under various stress conditions or virulence in the silkworm infection model. These

61 findings suggest that Atg1 is involved in regulating the morphologic changes of *T. asahii*.

Materials & Methods

T. asahii strains

65 *T. asahii* MPU129Δ*ku70* was used as the parental strain [19]. Three independent *T. asahii*

66 *atg1* gene-deficient mutants were generated in this study (Table 1).

Culture conditions for *T. asahii*

69 *T. asahii* strains were cultured according to previously described methods [19]. *T. asahii* strains

70 were grown on Sabouraud dextrose agar (SDA; 1% hipolypeptone, 4% dextrose, and 1.5% agar)

71 containing G418 (100 μg/mL) and cefotaxime (100 μg/mL) at 27°C for 1–2 days.

Reagents

Nourseothricin was purchased from JENA BIOSCIENCE GMBH (Dortmund, Germany).

Fluconazole was purchased from Tokyo Chemical Industry Co., Ltd. (Tokyo, Japan). Cefotaxime

sodium, D-glucose, agar, NaCl, sorbitol, H_2O_2, dithiothreitol (DTT), and Yeast Nitrogen Base

without Amino Acids and Ammonium Sulfate were obtained from FUJIFILM Wako Pure

Chemical Corporation (Osaka, Japan). Congo red was purchased from Sigma-Aldrich Co., LLC

(St. Louis, MO, USA). G418 was obtained from Enzo Life Science, Inc. (Farmingdale, NY, USA).

Hipolypeptone was purchased from NIHON PHARMACEUTICAL Co., LTD. (Tokyo, Japan).

Tunicamycin was purchased from Cayman Chemical Company (Ann Arbor, MI, USA).

Amphotericin B and KCl were obtained from Wako Pure Chemical Industries, Ltd. (Osaka, Japan).

Sodium dodecyl sulfate (SDS) was purchased from NIPPON GENE Co., LTD. (Tokyo, Japan).

Tween 80 was obtained from MP Biomedicals LLC (Santa Ana, CA, USA).

86 Conservation of Atg1 protein amino acid sequence in *T. asahii* and other fungi

Homology and domain analyses were performed as previously described [20]. The amino acid

sequence of the *T. asahii* Atg1 protein was obtained from the National Center for Biotechnology

Information (NCBI) (https://www.ncbi.nlm.nih.gov/). BLASTp

(https://blast.ncbi.nlm.nih.gov/Blast.cgi) was used to compare the sequence homology with the

Atg1 proteins of *C. albicans*, *A. fumigatus*, *C. neoformans*, and *S. cerevisiae*. A phylogenetic tree

was constructed using the maximum likelihood method with MEGA11 Software

(https://www.megasoftware.net/). The NCBI gene accession numbers of Atg1 proteins used for

phylogenetic analysis and alignment are listed in Table 2. The NCBI domain accession numbers

used for functional domain analysis and alignment are shown in Table 3.

97 Generation of *atg1* gene-deficient *T. asahii* mutants

Generation of the gene-deficient *T. asahii* mutants was performed according to previously

described methods [21]. The pUC19-*atg1* (5'UTR) -*NAT1*-*atg1* (3'UTR) plasmid was constructed

using the In-Fusion method. Approximately 1200 bp of the 5′-UTR and 3′-UTR regions of

the *atg1* gene were cloned upstream and downstream of the *NAT1* gene region in the pUC19-

NAT1 vector using the In-Fusion method (In-Fusion Snap Assembly Master Mix; Takara Bio Inc.,

Shiga, Japan). Using the pUC19-*atg1* (5'UTR) -*NAT1-atg1* (3'UTR) plasmid, polymerase chain

reaction (PCR) was performed to amplify a DNA fragment containing *atg1* (5'UTR) -*NAT1-atg1*

(3'UTR). The primers used for PCR are listed in Table 4. Gene transfer to *T. asahii* cells was

performed as described previously with modifications [22]. The parental strain was cultured on

SDA at 27°C for 1 day. Cells on the agar medium were suspended in 2 mL of saline and filtered

through a 40-µm cell strainer. The filtered suspension was centrifuged at 10,000 rpm for 5 min,

and the pellet was resuspended in 1 mL of distilled water and centrifuged again under the same

conditions. This washing step was repeated four times. The washed cells were then suspended in

1 mL of 1.2 M sorbitol solution and centrifuged at 10,000 rpm for 5 min. The harvested pellet was

resuspended in 40 µL of 1.2 M sorbitol to obtain competent cells. To perform electroporation, 4

µL of the PCR-amplified *atg1* (5'UTR) -*NAT1-atg1* (3'UTR) DNA fragment was added to 40 µL

of the competent cells and incubated on ice for 15 min. The mixture was transferred to a 2-mm

gap cuvette (Bio-Rad Laboratories, Inc., CA, USA), and electroporation was performed using a

Gene Pulser Xcell (Bio-Rad Laboratories, Inc., CA, USA) with the time constant protocol (1800

V, 5 ms). After electroporation, 500 µL of Yeast Polypeptone Dextrose (YPD) medium (2%

hipolypeptone, 1% yeast extract, 2% glucose) containing 0.6 M sorbitol was added, and the cells

were incubated at 27°C for 3 h. After incubation, the cells were plated on SA medium containing

nourseothricin (300 µg/mL) and incubated at 27°C for 3 days. Genomic DNA was extracted from

the nourseothricin-resistant transformants, and deletion of the *atg1* gene was confirmed by PCR.

Temperature sensitivity test

A temperature sensitivity test was performed as described previously with modifications [22]. *T.*

asahii strains were grown on SDA containing cefotaxime (100 μg/mL) at 27°C for 1 day. *T. asahii*

cells were suspended in saline, and the absorbance at 630 nm was adjusted to 1. A 10-fold serial

dilution of each suspension was prepared using saline. Five microliters of each dilution were

spotted onto SDA and incubated at 27°C, 37°C, or 40°C for 48 h. Photographs of the agar plates

were taken after incubation. For evaluation in liquid culture, Sabouraud dextrose broth (1%

hipolypeptone, 4% dextrose) was used. *T. asahii* cells were suspended in saline and filtered

through a 40-μm cell strainer. Absorbance at 630 nm was adjusted to 0.005. The suspensions were

incubated at 27°C, 37°C, or 40°C for 96 h, and absorbance at 630 nm was measured every 24 h

using a microplate reader (iMark™ microplate reader; Bio-Rad Laboratories Inc., Hercules, CA,

USA).

Drug sensitivity test

A drug sensitivity test was performed as previously reported with modifications [20]. *T. asahii* was

grown on SDA containing cefotaxime (100 μg/mL) at 27°C for 2 days. Cells were suspended in

saline and filtered through a 40-μm cell strainer. Absorbance of the suspension was adjusted to 1.0

at 630 nm. A 10-fold serial dilution of each suspension was prepared using saline. Five microliters

of each dilution were spotted onto SDA containing NaCl, KCl, Congo red, SDS, H_2O_2,

tunicamycin, DTT, amphotericin B, or fluconazole. The plates were incubated at 27°C or 37°C for

24 to 72 h, and images of the agar plates were captured after incubation.

The minimum inhibitory concentration (MIC) was determined using the broth

microdilution method according to CLSI guideline M27-A3, with a commercial antifungal

susceptibility testing plate for yeasts (DP Eiken; Eiken Chemical Co., Ltd., Tokyo, Japan) [20]. *T.*

asahii was grown on SDA containing cefotaxime (100 μg/mL) at 27°C for 2 days. The cells were

suspended in saline and adjusted to a turbidity equivalent to McFarland standard 1. The suspensions were then diluted 20-fold with saline, followed by a 100-fold dilution with RPMI 1640 medium (Thermo Fisher Scientific Inc., MA, USA) supplemented with 0.165 M 3-morpholinopropane-1-sulfonic acid (RPMI-MOPS, pH 7.0; FUJIFILM Wako Pure Chemical Corporation, Osaka, Japan). A 100-µL volume of the prepared cell suspension was added to each well of the plate containing one of the following eight antifungal agents: micafungin, caspofungin, amphotericin B, 5-fluorocytosine, fluconazole, itraconazole, voriconazole, or miconazole. The plates were incubated at 37°C for 48 h. MIC values were determined based on the results of two independent experiments.

Morphologic analysis of *T. asahii*

T. asahii strains were grown on SDA containing cefotaxime (100 µg/mL) at 27°C for 2 days. *T. asahii* cells were suspended in saline and filtered through a 40-µm cell strainer. Absorbance of the suspension was adjusted to 1.0 at 630 nm. A 20-µL aliquot of the suspension was added to 200 µL each of the Sabouraud dextrose broth (SB) and Synthetic Defined – Nitrogen (SD–N) medium. The cells were incubated at 37°C for 24 h. After incubation, culture samples were placed on microscope slides, covered with coverslips, and observed under a light microscope (BX51; OLYMPUS Corp., Tokyo, Japan). Photographs of the cells were randomly captured from multiple microscopic fields. The *T. asahii* morphology was classified as follows: yeast form, observed as single, spherical cells including budding yeasts; arthroconidia, observed as chains of three or more round or rectangular cells separated by septa; hyphal form, observed as elongated structures at least twice the length of yeast cells, including pseudohyphae, but lacking septa characteristic of

arthroconidia [20,23]. More than 400 cells were observed, and the number of each morphologic

type (yeast, arthroconidia, hyphae) was counted.

Silkworm infection experiments

Silkworm infection experiments were performed according to previously described methods with

modifications [22]. Silkworm eggs (KINSYU X SHOWA) were purchased from Ehime-Sanshu

Co., Ltd. (Ehime, Japan), disinfected, and incubated at 25–27°C until hatching. The hatched larvae

were reared on an artificial diet, SilkMate 2S (Nosan Corp., Kanagawa, Japan). Fifth-instar

silkworm larvae were fed SilkMate 2S overnight before use in the infection experiments. *T.*

asahii was grown for 1 day at 27°C on SDA containing cefotaxime (100 μg/mL). Cells were

suspended in saline and filtered through a 40-μm cell strainer. Absorbance of the resulting

suspension at 630 nm was adjusted to 1. The cell suspension was then diluted 250-fold with saline,

and 50 μL of the diluted suspension was injected into the silkworm hemolymph from the dorsal

side using a 1-mL tuberculin syringe (Terumo Medical Corporation, Tokyo, Japan). The silkworms

were fed SilkMate 2S and maintained in a 37°C incubator, where their survival was monitored.

Statistical analysis

All experiments were performed at least twice, and representative results are shown. For

morphologic analysis, differences between groups were analyzed by Dunnett's test. For silkworm

infection assays, differences between groups were analyzed using the log-rank test based on

Kaplan–Meier survival curves generated with EZR version 1.5.5 (https://www.jichi.ac.jp/saitama-

sct/SaitamaHP.files/download.html). A *P*-value of < 0.05 was considered statistically significant.

Results

Identification and conservation of the Atg1 Protein in *T. asahii*

We sought to identify a gene encoding a homologous Atg1 protein in the *T. asahii* genome. In *C. neoformans*, *atg1* was identified based on the amino acid sequence of the *S. cerevisiae* Atg1 protein [11,16]. *T. asahii* hypothetical protein A1Q1_02656 was identified as a candidate protein with 52% amino acid identity to *C. neoformans* Atg1 (Table 2). Next, we performed a homology analysis comparing the amino acid sequence of *T. asahii* hypothetical protein A1Q1_02656 with Atg1 proteins from representative fungi such as *S. cerevisiae*, *A. fumigatus,* and *C. albicans*. *T. asahii* hypothetical protein A1Q1_02656 showed 52%, 52%, 50%, and 47% amino acid identity with Atg1 proteins of *C. neoformans*, *A. fumigatus*, *C. albicans*, and *S. cerevisiae*, respectively (Figure 1A, Table 2). A phylogenetic tree generated from these amino acid sequences revealed that, among the fungi analyzed, *T. asahii* hypothetical protein A1Q1_02656 was most closely related to *C. neoformans* Atg1 (Fig. 1B). Functional domain analysis revealed that the *T. asahii* hypothetical protein A1Q1_02656 also contained a PKc_like superfamily domain, a catalytic domain of protein kinases (Fig. 1C, Table 3). Therefore, the *T. asahii* hypothetical protein A1Q1_02656 was estimated to be the Atg1 protein in *T. asahii,* and the corresponding gene was designated *atg1*.

Fig. 1. Conservation of *T. asahii* Atg1 among fungi.

(**A**) Phylogenetic tree of Atg1 proteins from *T. asahii*, *C. neoformans*, *A. fumigatus*, *C. albicans*, and *S. cerevisiae,* constructed using the maximum likelihood method. (**B**) Phylogenetic tree of Atg1 proteins from *T. asahii*, *C. neoformans*, *A. fumigatus*, *C. albicans*, and *S. cerevisiae,* constructed using the maximum likelihood method. Phylogenetic analysis was performed using

216   MEGA11 Software (https://www.ncbi.nlm.nih.gov/). (**C**) Schematic diagrams of domains present

217   in Atg1 proteins. PKc: PKc_like superfamily domain. STKc: STKc_ATG1_ULK_like domain.

218   HB: Herpes_BLLF1 superfamily domain. DUF: DUF3543. Functional domain prediction was

219   performed   using   the   National   Center   for   Biotechnology   Information   (NCBI)

220   (https://www.ncbi.nlm.nih.gov/).

Generation of *atg1* gene-deficient *T. asahii* mutants

The gene-targeting method used to generate gene-deficient *T. asahii* mutants was established

previously [19][24]. Using the method, we generated three *atg1* gene-deficient *T. asahii* mutants.

The DNA fragment used for *atg1* gene deletion contained the *NAT1* gene, which confers resistance

to nourseothricin (Fig. 2A). Homologous recombination replaced the *atg1* gene with

the *NAT1* gene, thereby conferring resistance to nourseothricin (Fig. 2B). Transformation was

performed by introducing the DNA fragment into the parental strain via electroporation, and

transformants resistant to nourseothricin were obtained (Fig. 2B). Genomic DNA was extracted

from the nourseothricin-resistant transformants, and *atg1* gene deficiency was confirmed by PCR

(Fig. 2C, D). These results suggest that the *atg1* gene-deficient *T. asahii* mutants were generated.

Fig. 2. Generation of the *atg1* gene-deficient *T. asahii* mutants.

(**A**) Schematic diagram of the gene recombination strategy used to construct the *atg1* gene-

deficient *T. asahii* mutant (Δ*atg1*), along with the predicted genomic structure of the *atg1* gene-

deficient mutant. (**B**) The parental strain (Parent) and the *atg1* gene-deficient mutants (Δ*atg1* #1-

3) were spread on SDA or SDA containing nourseothricin (100 µg/mL; Nou (+)) and incubated at

27°C for 2 days. (**C**) Schematic illustration of primer positions and expected amplified DNA

fragment sizes by PCR to verify the genomic structure of the *atg1* gene-deficient candidates using

extracted genomic DNA. (**D**) DNA electrophoresis of PCR products to confirm the *atg1* gene

deficiency (Δ*atg1*).

Effects of *atg1* gene deficiency on stress sensitivity

Next, we investigated whether growth of the *atg1* gene-deficient mutants was similar to that of the

parental strain. Under temperatures of 27°C, 37°C, or 40°C, no significant differences in growth

were observed between the parental strain and the *atg1* gene-deficient mutants (Fig. 3). In *C.*

neoformans, mutants lacking autophagy-related genes exhibit sensitivity to oxidative and osmotic

stress [15]. To evaluate whether the *T. asahii atg1* gene is involved in osmotic stress, cell

wall/membrane stress, endoplasmic reticulum stress, or oxidative stress, stress sensitivity assays

were performed. The growth of the *atg1* gene-deficient mutants was not altered under the tested

compounds (Fig. 4). The MICs of antifungal agents for the *atg1* gene-deficient strains were

comparable to those for the parental strain (Table 5). These results suggest that the *atg1* gene is

not involved in susceptibility to compound-induced stresses and antifungal agents in *T. asahii*.

Fig. 3. Growth of the *atg1* gene-deficient *T. asahii* mutants under high-temperature

conditions.

(**A**) The parental strain (Parent) and the *atg1* gene-deficient *T. asahii* mutants (Δ*atg1* #1-3) were

grown on SDA at 27°C for 2 days. Cells were suspended in saline, and 10-fold serial dilutions

were prepared. Each dilution (5 μl) was spotted onto SDA and incubated at 27°C, 37°C, or 40°C

for 48 h. (**B**) The parental strain (Parent) and the *atg1* gene-deficient *T. asahii* mutants (Δ*atg1* #1-

3) were grown in Sabouraud dextrose broth at 27°C, 37°C, or 40°C, and absorbance at 630 nm

was measured. Data are presented as mean ± standard error of the mean (SEM).

Fig. 4. Growth of the *atg1* gene-deficient *T. asahii* mutants under stress-inducing conditions.

The parental strain (Parent) and the *atg1* gene-deficient *T. asahii* mutants (Δ*atg1* #1-3) were

cultured on SB at 27°C for 2 days. Cells were suspended in saline, and 10-fold serial dilutions

were prepared. Each dilution (5 µL) was spotted onto SA medium containing NaCl (1 M), KCl

(1.5 M), Congo red (300 µg/mL), SDS (0.01%), tunicamycin (TM, 1 µg/mL), dithiothreitol (DTT,

15 mM), or hydrogen peroxide (H_2O_2, 4 mM), and incubated at 27°C for 48 h.

Role of the *atg1* gene in *T. asahii* morphology

T. asahii exhibits dimorphism, alternating between yeast forms and filamentous/arthroconidial

forms [1]. In *C. albicans*, a dimorphic fungus, hyphal formation is induced by various nutritional

conditions, such as growth at 37°C and exposure to GlcNAc [25]. To observe the morphologic

changes of *T. asahii* yeast cells under various nutritional conditions, we developed a method to

collect *T. asahii* yeast cells. More than 90% of *T. asahii* cells in suspension filtered through a 40-

µm filter were yeast cells (Fig. 5A, B). Using the *T. asahii* yeast cells, we tested the growth of the

cells in various culture media, such as SB and SD-N. The absorbance of *T. asahii* culture medium

at 630 nm was increased in the order SB and SD-N (Fig. 5C). The ratio of hyphae was increased

in SB by incubating at 37°C for 24 h (Fig. 5D, E). The ratio of arthroconidia was also increased

by incubation in SB and SD-N at 37°C for 24 h (Fig. 5D, E). We investigated whether the *atg1*

gene contributes to the morphologic changes of *T. asahii* in various culture media. The *atg1* gene-

deficient mutants exhibited a higher proportion of hyphal cells in SB (Fig. 6). No significant

morphologic differences were observed, however, between the parental strain and the *atg1*-

deficient *T. asahii* mutants in SD-N (Fig. 6). These results suggest that the *atg1* gene negatively

regulates hyphal formation in *T. asahii* in a nutrient-dependent manner.

Fig. 5 Effects of culture conditions on *T. asahii* morphology.

(**A**) Method for preparing the yeast form of *T. asahii*. The parental *T. asahii* strain (Parent) was

cultured on SDA at 27°C for 2 days. *T. asahii* cells were suspended in saline and filtered through

a 40-μm cell strainer. (**B**) The ratio of each cell shape in the filtered *T. asahii* cells was measured.

(**C**) Absorbance of the cell suspension at 630 nm was adjusted to 0.5, and 5 μL of the filtered cell

suspension was added to 200 μL of Sabouraud dextrose medium or SD-N liquid medium. Cultures

were incubated at 27°C for 24 h, and absorbance at 630 nm was measured for 24 h. (**D-F**)

Absorbance of the filtered cell suspension at 630 nm was adjusted to 1.0, and 20 μL of the

suspension was added to 200 μL of Sabouraud dextrose broth (SB) or synthetic defined-nitrogen

(SD–N) medium, followed by incubation at 27°C for 24 h. Cells were then observed under a light

microscope. Scale bars represent 40 μm. Images were captured randomly from multiple

microscopic fields. Cell numbers in three morphologic categories: yeast, arthroconidia, and hyphae,

were quantified. (**D**) Cell numbers in three morphologic categories were measured at 0 or 24 h

after incubation at 37°C. (**E**) Photographs of cells after incubation at 37°C for 24 h were obtained

under microscopic fields. *n* represents the total number of microscopic fields analyzed from two

independent experiments.

Fig. 6. Effect of *atg1* gene deficiency on *T. asahii* morphology.

The parental *T. asahii* strain (Parent) and three *atg1* gene-deficient mutants (Δ*atg1* #1-3) were

grown on SDA at 27°C for 2 days. Cells were suspended in saline and filtered through a 40-µm

cell strainer. Absorbance at 630 nm was adjusted to 1. A 20-µL aliquot of each suspension was

added to (**A**) Sabouraud dextrose broth (SB), or (**B**) synthetic defined-nitrogen (SD–N) medium

(200 µL). The samples were incubated at 37°C for 24 h. After incubation, the *T. asahii* cells were

dropped onto microscope slides, covered with coverslips, and observed under a light microscope.

Upper panels show representative microscopic images. Scale bars represent 20 µm. Lower panels

show quantification of the three morphologic forms—yeast, arthroconidia, and hyphae—based on

randomly selected microscopic fields. The total number of cells was obtained from two

independent experiments. (**C**) The relative ratio of the three morphologic forms were determined

compared to the ratio of parental strain (Parent) under a culture condition with SB at 37°C for 24

h. The data of the relative ratio were obtained from seven independent experiments. Error bars

indicate the SEM of the means (n = 7). Statistical differences between parental strain and Δ*atg1*

#1-3 were analyzed by Dunnett's test. (*: $P < 0.05$).

Virulence of the *atg1* gene-deficient mutants in a silkworm infection model

In *C. neoformans*, *atg1* gene deficiency did not affect its virulence in a *Galleria mellonella*

infection model [15]. We tested whether the *atg1* gene is required for *T. asahii* virulence in a

silkworm infection model. The survival time of silkworms infected with the *atg1*-deficient mutants

was not significantly different from that of silkworms infected with the parental strain (Fig. 7).

The result suggests that the *atg1* gene does not contribute to *T. asahii* virulence in the silkworm

model.

Fig. 7. Virulence of *T. asahii atg1* gene-deficient mutants in a silkworm infection model.

The parental strain (Parent; 530 cells/larva) and three independent *atg1* gene-deficient *T. asahii*

mutants (Δ*atg1* #1-3; 790 - 950 cells/larva) were injected into the hemocoel of silkworms, and

survival was monitored at 37°C for 96 h. Statistical significance between the parental and *atg1*-

deficient groups was determined by the log-rank test based on Kaplan–Meier survival curves. Ten

silkworms per group were used.

Discussion

The findings of the present study demonstrated that the *atg1* gene is involved in morphologic

transitions of *T. asahii* under nutrient-rich conditions. On the other hand, *atg1* gene deficiency did

not affect growth under various stress conditions or virulence of *T. asahii* in the silkworm model.

These findings suggest that Atg1 might play a role in regulating hyphal formation under specific

culture conditions but is not involved in stress resistance or virulence in *T. asahii*.

T. asahii Atg1 protein was estimated based on the amino acid sequence of other fungal

Atg1 proteins and their domain analysis. *T. asahii* hypothetical protein A1Q1_02656 has high

sequence similarity to the Atg1 proteins of other representative fungi, including *S. cerevisiae*, *A.*

fumigatus, and *C. albicans*. Domain analysis revealed that *T. asahii* hypothetical protein

A1Q1_02656 harbors a Pkc_like superfamily domain—a catalytic domain of protein kinases

conserved among the analyzed fungi. In *S. cerevisiae*, Atg1 kinase activity contributes to non-

selective autophagy [8]. Therefore, we determined that *T. asahii* hypothetical protein A1Q1_02656

is the Atg1 protein of *T. asahii*.

T. asahii atg1 gene deficiency does not significantly affect the stress response and

antifungal drug resistance of *T. asahii*. We generated *atg1* gene-deficient *T. asahii* mutants and

analyzed their phenotypes. When generating gene-deficient mutants, unintended secondary

mutations may be introduced in addition to the targeted gene disruption. To minimize the

possibility of this rare occurrence, we generated three independent *T. asahii atg1* gene-deficient

mutants in this study and examined whether they exhibited consistent phenotypes. The *atg1* gene

deficiency in *T. asahii* did not affect sensitivity to high temperature, NaCl, KCl, tunicamycin, DTT,

H_2O_2, or antifungal drugs. In *C. neoformans*, the *atg1* gene-deficient mutant also did not exhibit

high-temperature and H_2O_2-sensitive phenotypes [15,16]. Therefore, the *atg1* gene is not involved

in high temperature and oxidative stress tolerance in *T. asahii* or *C. neoformans*. On the other hand,

in *C. neoformans*, the growth of the *atg1* gene-deficient mutant was delayed by NaCl [15]. The

atg1 genes in *T. asahii* and *C. neoforman*s might differ in their tolerance to NaCl. The calcineurin

and Hog1 pathways are involved in high-temperature tolerance in *T. asahii* [19,20]. We therefore

concluded that the *T. asahii* Atg1 is regulated in a calcineurin and Hog1 pathway-independent

manner.

The *atg1* gene contributes to the morphologic changes of *T. asahii*. In SB, the ratio of

hyphae was increased by *atg1* gene deficiency after 24-h incubation. On the other hand, in SD-N,

a nitrogen-limited medium, the morphology in the *atg1* gene-deficient mutants was not different

from that in the parental strain. In fungi such as *S. cerevisiae*, *C. neoformans*, and *C. albicans*,

Atg1 is involved in nitrogen limitation-induced autophagy and required for survival under

nitrogen-limited conditions [26]. In the present study, we revealed the role of Atg1 in the

morphologic changes of *T. asahii* under nutrient-rich conditions. Further studies are needed to

elucidate the intracellular mechanisms regulated by Atg1 under nutrient-rich conditions.

The *T. asahii atg1* gene is not involved in *T. asahii* virulence in the silkworm infection

model. In *C. neoformans*, Atg1 does not influence virulence in the insect model *Galleria*

mellonella [15]. Therefore, we assume that Atg1 does not have a pivotal role in *T. asahii* and *C.*

neoformans virulence in insect models.

A limitation of this study is that the effects of *atg1* gene-deficiency of *T. asahii* on

autophagy were not evaluated. Autophagy validation methods have been established in several

fungi using fungal cells expressing GFP-Atg8 [27]. The development of a detection system for

autophagy in *T. asahii* using cells expressing GFP-Atg8–expressing cells remains a future

challenge.

Conclusion

384 The *atg1* gene contributes to the morphologic changes of *T. asahii* under nutrient-rich conditions.

385 Hyphal formation of *T. asahii* might be partially regulated by Atg1-dependent cellular events.

Acknowledgements

388 We thank Renta Endo, Momoka Matsumura, and Tomoya Sanbongi (Meiji Pharmaceutical

389 University) for technical assistance rearing the silkworms. We also thank the editors of

390 SciTechEdit International LLC (Highlands Ranch, CO, USA) for English-language editing.

Funding

393 This study was supported by JSPS KAKENHI grant number JP23K06141 (Scientific Research (C)

394 to Y.M.) and in part by the Research Program on Emerging and Re-emerging Infectious Diseases

395 of the Japan Agency for Medical Research and Development, AMED (Grant number

396 JP23fk0108679h0401 to T.S.).

References

1.   Colombo AL, Padovan ACB, Chaves GM. Current knowledge of trichosporon spp. and
trichosporonosis. Clinical Microbiology Reviews. 2011 Oct;24(4):682-700.

2.   Walsh TJ, Lee JW, Melcher GP, Navarro E, Bacher J, Callender D, et al. Experimental
Trichosporon infection in persistently granulocytopenic rabbits: implications for
pathogenesis, diagnosis, and treatment of an emerging opportunistic mycosis. J Infect Dis.
1992 Jul;166(1):121-33.

3.   Walsh TJ, Melcher GP, Lee JW, Pizzo PA. Infections due to Trichosporon species: new
concepts in mycology, pathogenesis, diagnosis and treatment. Curr Top Med Mycol.
1993;5:79-113.

4.   Duarte-Oliveira C, Rodrigues F, Gonçalves SM, Goldman GH, arvalho A, Cunha C. The
cell biology of the Trichosporon-host interaction. Front Cell Infect Microbiol. 2017 Apr
7;7:118.

5.   Krcmery V Jr, Mateicka F, Kunová A, Spánik S, Gyarfás J, Sycová Z, et al. Hematogenous
trichosporonosis in cancer patients: report of 12 cases including 5 during prophylaxis with
itraconazol. Support Care Cancer. 1999 Jan;7(1):39-43.

6.   Kimura M, Araoka H, Yamamoto H, Nakamura S, Nagi M, Yamagoe S, et al. Micafungin
Breakthrough Fungemia in Patients with Hematological Disorders. 2018 Apr
26;62(5):e02183-17.

7.   Iturrieta-González IA, Padovan ACB, Bizerra FC, Hahn RC, Colombo AL. Multiple species
of Trichosporon produce biofilms highly resistant to triazoles and amphotericin B. PLoS
One. 2014 Oct 31;9(10):e109553.

8.   Cai YY, Li L, Zhu XM, Lu JP, Liu XH, Lin FC. The crucial role of the regulatory
mechanism of the Atg1/ULK1 complex in fungi. Frontiers in Microbiology. 2022 Oct
26;13:1019543.

9.   Ohsumi Y. Molecular mechanism of autophagy in yeast, Saccharomyces cerevisiae. Philos
Trans R Soc Lond B Biol Sci. 1999 Sep 29;354(1389):1577-80; discussion 1580-1.

10.   Wong PM, Puente C, Ganley IG, Jiang X. The ULK1 complex sensing nutrient signals for
autophagy activation. Autophagy. 2013 Feb 1;9(2):124-37.

11.   Kamada Y, Funakoshi T, Shintani T, Nagano K, Ohsumi M, Ohsumi Y. JCB Report Tor-
mediated Induction of Autophagy Via an Apg1 Protein Kinase Complex. J Cell Biol. 2000
Sep 18;150(6):1507-13.

12.   Metur SP, Klionsky DJ. Nutrient-dependent signaling pathways that control autophagy in
yeast. FEBS Letters. 2024 Jan;598(1):32-47.

13.   Kitamoto K, Yoshizawa K, Ohsumi Y, Anraku2 Y. Dynamic Aspects of Vacuolar and
Cytosolic Amino Acid Pools of Saccharomyces cerevisiae. J Bacteriol. 1988
Jun;170(6):2683-6.

14.   Suzuki K, Kubota Y, Sekito T, Ohsumi Y. Hierarchy of Atg proteins in pre-autophagosomal
structure organization. Genes to Cells. 2007 Feb;12(2):209-18.

15.   Zhao X, Feng W, Zhu X, Li C, Ma X, Li X, et al. Conserved Autophagy Pathway
Contributes to Stress Tolerance and Virulence and Differentially Controls Autophagic Flux
Upon Nutrient Starvation in Cryptococcus neoformans. Front Microbiol. 2019 Nov
26;10:2690.

16.   Ding H, Caza M, Dong Y, Arif AA, Horianopoulos LC, Hu G, et al. ATG genes influence
the virulence of Cryptococcus neoformans through contributions beyond core autophagy
functions. Infect Immun. 2018 Aug 22;86(9):e00069-18.

17.   Siqi L, Liuliu J, Haochen M, Ying L, Qinqin Z, Ming M, et al. Autophagy regulation of
ATG13 and ATG27 on biofilm formation and antifungal resistance in Candida albicans.
Biofouling. 2022 Oct;38(9):926-939.

18.   Huang Y, Yu S, Liu S, Zhao X, Chen X, Wei X. Autophagy Activated by Atg1 Interacts
With Atg9 Promotes Biofilm Formation and Resistance of Candida albicans. J Basic
Microbiol. 2025 May;65(5):e2400603.

19.   Matsumoto Y, Yoshikawa A, Nagamachi T, Sugiyama Y, Yamada T, Sugita T. A critical
role of calcineurin in stress responses, hyphal formation, and virulence of the pathogenic
fungus Trichosporon asahii. Sci Rep. 2022 Sep 27;12(1):16126.

20.   Matsumoto Y, Sugiyama Y, Nagamachi T, Yoshikawa A, Sugita T. Hog1-mediated stress
tolerance in the pathogenic fungus Trichosporon asahii. Sci Rep. 2023 Aug 19;13(1):13539.

21.   Matsumoto Y, Nagamachi T, Yoshikawa A, Yamazaki H, Yamasaki Y, Yamada T, et al.
Development of an efficient gene-targeting system for elucidating infection mechanisms of
the fungal pathogen Trichosporon asahii. Sci Rep. 2021 Sep 14;11(1):18270.

22.   Matsumoto Y, Nagamachi T, Yoshikawa A, Yamada T, Sugita T. A joint PCR-based gene-
459   targeting method using electroporation in the pathogenic fungus Trichosporon asahii. AMB
460   Express. 2022 Jul 14;12(1):91.

461   23.   Kurakado S, Miyashita T, Chiba R, Sato C, Matsumoto Y, Sugita T. Role of arthroconidia
462   in biofilm formation by Trichosporon asahii. Mycoses. 2021 Jan;64(1):42-47.

463   24.   Matsumoto Y, Kurakado S, Yamada T, Sugita T. Strategy to Identify Virulence-Related
464   Genes of the Pathogenic Fungus Trichosporon asahii Using an Efficient Gene-Targeting
465   System. Microbiology and Immunology. 2025 Feb;69(2):77-84.

466   25.   Su C, Yu J, Lu Y. Hyphal development in Candida albicans from different cell states.
467   Current Genetics. 2018 Dec;64(6):1239-1243.

468   26.   Cheong H, Nair U, Geng J, Klionsky DJ. The Atg1 Kinase Complex Is Involved in the
469   Regulation of Protein Recruitment to Initiate Sequestering Vesicle Formation for
470   Nonspecific Autophagy in Saccharomyces cerevisiae. Mol Biol Cell. 2008 Feb;19(2):668-
471   81.

472   27.   Kirisako T, Baba M, Ishihara N, Miyazawa K, Ohsumi M, Yoshimori T, et al. Formation
473   Process of Autophagosome Is Traced with Apg8/Aut7p in Yeast. J Cell Biol. 1999 Oct
474   18;147(2):435-46.

Supporting information

S1 Dataset. Datasets included in this study.

480 **Table 1 *T. asahii* strains used in this study**

T. asahii strains	Relevant genotype	Background	Reference
MPU129 Δ*ku70* (Parent strain)	*ku70*::*nptII*	MPU129	Matsumoto Y., *et. al*. [21]
Δ*atg1* #1-#3	*ku70*::*nptII*, *atg1*::*NAT1*	MPU129 Δ*ku70*	This study

482   Table 2 NCBI gene accession numbers for the Atg1 protein used to create the phylogenetic
483   tree

Species	Definition	NCBI accession	Size (aa)	Query Cover (%)	Percent Identical (%)
Trichosporon asahii	hypothetical protein A1Q1_02656 (*Trichosporon asahii var. asahii* CBS 2479)	XP_014179392.1	995	100	100
Cryptococcus neoformans	ULK/ULK protein kinase (*Cryptococcus neoformans var. grubii* H99)	XP_012048654.1	985	98	51.87
Aspergillus fumigatus	serine/threonine protein kinase (Pdd7p), putative (*Aspergillus fumigatus* Af293).	XP_751920.1	973	77	52.33
Candida albicans	serine/threonine protein kinase (*Candida albicans* SC5314)	XP_717245.2	834	31	50.46
Saccharomyces cerevisiae	serine/threonine protein kinase ATG1 (*Saccharomyces cerevisiae* S288C)	NP_011335.1	897	61	47.45

485
486

487 **Table 3 NCBI domain accession number for the Atg1 protein.**
488

Domain name	NCBI accession	Description	Species	Interval
PKc_like super family	cl21453	Protein Kinases, catalytic domain	*Trichosporon asahii*	34-339
STKc_ATG1_ULK_like	cd14009	Catalytic domain of the Serine/Threonine kinases, Autophagy-related protein 1 and Unc-51-like kinases	*Cryptococcus neoformans* *Aspergillus fumigatus* *Candida albicans* *Saccharomyces cerevisiae*	33-336 29-327 64-350 30-324
DUF3543	pfam12063	Domain of unknown function (DUF3543)	*Trichosporon asahii* *Cryptococcus neoformans* *Aspergillus fumigatus* *Candida albicans* *Saccharomyces cerevisiae*	737-974 741-979 651-941 621-829 586-893
Herpes_BLLF1 super family	cl37540	Herpes virus major outer envelope glycoprotein	*Trichosporon asahii*	370-536

489
490

491   **Table 4 Primers used to generate *atg1* gene-deficient mutants.**
492

Primers	Nucleic acid sequence
pUC19-*NAT1* for cloning	
F *NAT1* (for 5UTR typing)	GAGCTGATACCGCTCGCCGCTGCGAGGATGTGAGCTGGAGAGC
R *NAT1* (for 3UTR typing)	GTTTCGGTGATGACGGTGGAAGAGATGTAGAAACTAGCTTCCTGGTTTCAGAG
F pUC19 (*NAT1* inf)	CACCGTCATCACCGAAACGCGC
R pUC19 (*NAT1* inf)	GGCGAGCGGTATCAGCTCACTCAAAGGCGGTAATACG
pUC19-*atg1* (5'UTR) -*NAT1*-*atg1* (3'UTR) for cloning	
pUC19-*atg1* (5'UTR) -*NAT1* (1st cloning)	
F *NAT1* (for 5'UTR typing)	GCTGCGAGGATGTGAGCTGGAGAGCGGCGC
R pUC19 (for 5'UTR typing)	GGCGAGCGGTATCAGCTCACTCAAAGGCGGTAATACG
F *atg1* (5'UTR)	GAGCTGATACCGCTCGCCGCCCATGTCTCGTCCGGTCCCAAACCTC
R *atg1* (5'UTR)	AGCTCACATCCTCGCAGCGCGTGTGGTGGGTGAGTGGGAGTGGATG
pUC19-*atg1* (5'UTR) -*NAT1*-*atg1* (3'UTR) (2nd cloning)	
F pUC19 (for 3'UTR typing)	CACCGTCATCACCGAAACGCGC
R *NAT1* (for 3'UTR typing)	GAAGAGATGTAGAAACTAGCTTCCTGGTTTCAGAGAC
F *atg1* (3'UTR)	TAGTTTCTACATCTCTTCCCCTCTGCTTCTCCTTCTTCGCCTGCGC
R *atg1* (3'UTR)	GTTTCGGTGATGACGGTGGCAAAGTCGCCTGCAAGTCGCCTGACCG
Primers-1 for *atg1* Genotyping	
F atg1 gene ORF	GGCAGAAGTTGACGAGCAAGCTGC
R atg1 gene ORF	CCGAAGCGAAGAATGCGTCGAAACC
Primers-2 for *atg1* genotyping	
F atg1 locus	GCCCATGTCTCGTCCGGTCCCAAACCTC
R atg1 locus	GCAAAGTCGCCTGCAAGTCGCCTGACCG
Primers-3 for *atg1* genotyping	
F atg1 gene outside	CGACCGACACAGAATTGAGCCACAAC
R atg1 gene outside	CTGGTGCGGTACCGGTAAGCCGTGTCGTCAAG

493
494

Table 5 MIC values of *atg1* gene-deficient mutants against antifungal agents.

MIC (μg/mL)	Parent	*Δatg1* #1	*Δatg1* #2	*Δatg1* #3
MCFG	>16	>16	>16	>16
CPFG	4	4	4	4
AMPH-B	0.5	0.5	0.5	0.5
5-FC	>64	>64	>64	>64
FLCZ	1	1	1	1
ITCZ	0.5	0.5	0.5	0.5
VRCZ	0.12	0.12	0.12	0.12
MCZ	0.5	0.5	0.5	0.5

The *T. asahii* parental strain (Parent) and three independent *atg1* gene-deficient mutant *T. asahii* strains (Δ*atg1* #1-3) were cultured in Sabouraud medium for 2 days at 27 °C. *T. asahii* cells were suspended in saline, and cell suspensions were prepared to a turbidity of 1 according to the McFarland turbidity method. Each bacterial solution was diluted 20-fold in saline and then 100-fold in RPMI MOPS. The prepared fungal solutions were then diluted with eight antifungal agents (Micafungin [MCFG], caspofungin [CPFG], amphotericin B [AMPH-B], 5-fluorocytosine [5-FC], fluconazole [FLCZ], itraconazole [ITCZ], voriconazole [VRCZ], miconazole [MCZ]) and inoculated on plates at 37°C for 48 h. The minimum inhibitory concentration (MIC) was determined from the results of two independent experiments.

Figure 1

Fig. 1. Conservation of *T. asahii* Atg1 among fungi.
(**A**) Phylogenetic tree of Atg1 proteins from *T. asahii*, *C. neoformans*, *A. fumigatus*, *C. albicans*, and *S. cerevisiae*, constructed using the maximum likelihood method. (**B**) Phylogenetic tree of Atg1 proteins from *T. asahii*, *C. neoformans*, *A. fumigatus*, *C. albicans*, and *S. cerevisiae*, constructed using the maximum likelihood method. Phylogenetic analysis was performed using MEGA11 Software (https://www.ncbi.nlm.nih.gov/). (**C**) Schematic diagrams of domains present in Atg1 proteins. PKc: PKc_like superfamily domain. STKc: STKc_ATG1_ULK_like domain. HB: Herpes_BLLF1 superfamily domain. DUF: DUF3543. Functional domain prediction was performed using the National Center for Biotechnology Information (NCBI) (https://www.ncbi.nlm.nih.gov/).

Figure 2

Fig. 2. Generation of the *atg1* gene-deficient *T. asahii* mutants.
(**A**) Schematic diagram of the gene recombination strategy used to construct the *atg1* gene-deficient *T. asahii* mutant (Δ*atg1*), along with the predicted genomic structure of the *atg1* gene-deficient mutant. (**B**) The parental strain (Parent) and the *atg1* gene-deficient mutants (Δ*atg1* #1-3) were spread on SDA or SDA containing nourseothricin (100 μg/mL; Nou (+)) and incubated at 27° C for 2 days. (**C**) Schematic illustration of primer positions and expected amplified DNA fragment sizes by PCR to verify the genomic structure of the *atg1* gene-deficient candidates using extracted genomic DNA. (**D**) DNA electrophoresis of PCR products to confirm the *atg1* gene deficiency (Δ*atg1*).

Figure 2

Figure 3

Fig. 3. **Growth of the _atg1_ gene-deficient _T. asahii_ mutants under high-temperature conditions.**
(**A**) The parental strain (Parent) and the _atg1_ gene-deficient _T. asahii_ mutants (Δ_atg1_ #1-3) were grown
on SDA at 27° C for 2 days. Cells were suspended in saline, and 10-fold serial dilutions were prepared.
Each dilution (5 μl) was spotted onto SDA and incubated at 27° C, 37° C, or 40° C for 48 h. (**B**) The
parental strain (Parent) and the _atg1_ gene-deficient _T. asahii_ mutants (Δ_atg1_ #1-3) were grown in
Sabouraud dextrose broth at 27° C, 37° C, or 40° C, and absorbance at 630 nm was measured. Data
are presented as mean ± standard error of the mean (SEM).

Figure 4

Fig. 4. Growth of the *atg1* gene-deficient *T. asahii* mutants under stress-inducing conditions.
The parental strain (Parent) and the *atg1* gene-deficient *T. asahii* mutants (Δ*atg1* #1-3) were cultured on
SB at 27° C for 2 days. Cells were suspended in saline, and 10-fold serial dilutions were prepared.
Each dilution (5 μL) was spotted onto SA medium containing NaCl (1 M), KCl (1.5 M), Congo red (300
μg/mL), SDS (0.01%), tunicamycin (TM, 1 μg/mL), dithiothreitol (DTT, 15 mM), or hydrogen peroxide
(H_2O_2, 4 mM), and incubated at 27° C for 48 h.

Figure 5

Fig. 5 Effects of culture conditions on *T. asahii* morphology.
(**A**) Method for preparing the yeast form of *T. asahii*. The parental *T. asahii* strain (Parent) was cultured on SDA at 27° C for 2 days. *T. asahii* cells were suspended in saline and filtered through a 40-μm cell strainer. (**B**) The ratio of each cell shape in the filtered *T. asahii* cells was measured. (**C**) Absorbance of the cell suspension at 630 nm was adjusted to 0.5, and 5 μL of the filtered cell suspension was added to 200 μL of Sabouraud dextrose medium or SD-N liquid medium. Cultures were incubated at 27° C for 24 h, and absorbance at 630 nm was measured for 24 h. (**D-F**) Absorbance of the filtered cell suspension at 630 nm was adjusted to 1.0, and 20 μL of the suspension was added to 200 μL of Sabouraud dextrose broth (SB) or synthetic defined-nitrogen (SD–N) medium, followed by incubation at 27° C for 24 h. Cells were then observed under a light microscope. Scale bars represent 40 μm. Images were captured randomly from multiple microscopic fields. Cell numbers in three morphologic categories,: yeast, arthroconidia, and hyphae, were quantified. (**D**) Cell numbers in three morphologic categories were measured at 0 or 24 h after incubation at 37° C. (**E**) Photographs of cells after incubation at 37° C for 24 h were obtained under microscopic fields. *n* represents the total number of microscopic fields analyzed from two independent experiments.

Figure 6

Figure 6-continued

Fig. 6. Effect of *atg1* gene deficiency on *T. asahii* morphology.
The parental *T. asahii* strain (Parent) and three *atg1* gene-deficient mutants (Δ*atg1* #1-3) were grown on SDA at 27° C for 2 days. Cells were suspended in saline and filtered through a 40-µm cell strainer. Absorbance at 630 nm was adjusted to 1. A 20-µL aliquot of each suspension was added to (**A**) Sabouraud dextrose broth (SB), or (**B**) synthetic defined-nitrogen (SD–N) medium (200 µL). The samples were incubated at 37° C for 24 h. After incubation, the *T. asahii* cells were dropped onto microscope slides, covered with coverslips, and observed under a light microscope. Upper panels show representative microscopic images. Scale bars represent 20 µm. Lower panels show quantification of the three morphologic forms—yeast, arthroconidia, and hyphae—based on randomly selected microscopic fields. The total number of cells was obtained from two independent experiments. (**C**) The relative ratio of the three morphologic forms were determined compared to the ratio of parental strain (Parent) under a culture condition with SB at 37° C for 24 h. The data of the relative ratio were obtained from seven independent experiments. Error bars indicate the SEM of the means (n = 7). Statistical differences between parental strain and Δ*atg1* #1-3 were analyzed by Dunnett's test. (*: $P < 0.05$).

Figure 7

Fig. 7. Virulence of *T. asahii atg1* gene-deficient mutants in a silkworm infection model.
The parental strain (Parent; 530 cells/larva) and three independent *atg1* gene-deficient *T. asahii* mutants
(Δ*atg1* #1-3; 790 - 950 cells/larva) were injected into the hemocoel of silkworms, and survival was
monitored at 37° C for 96 h. Statistical significance between the parental and *atg1*-deficient groups was
determined by the log-rank test based on Kaplan–Meier survival curves. Ten silkworms per group were
used.