1 **A chromosomal inversion accumulates adaptive and barrier loci in a cold-**
2 **adapted insect species**
3
4 **Running title: Inversion polymorphism in *Drosophila montana***
5

6 Poikela, N.[1,2], Hoikkala, V.[2], Ritchie, M. G.[1] & Kankare, M[2].
7
8 [1] Centre for Biological Diversity, University of St Andrews, United Kingdom
9 [2] Department of Biological and Environmental Science, University of Jyväskylä, Finland
10
Corresponding author:
FT Noora Poikela, np82@st-andrews.ac.uk, noora.p.poikela@gmail.com

Abstract

Chromosomal inversions can play a crucial role in population adaptation and divergence by
reducing gene flow and preserving adaptive allelic combinations between populations or
ecotypes that carry alternative arrangements. However, demonstrating this empirically remains
challenging because of the numerous interacting processes with similar genomic signatures
that can influence inversion evolution. In this study, we characterize a large (9.5Mb)
polymorphic inversion in the cold-adapted and widely distributed insect species, *Drosophila*
montana, using a combination of long- and short-read sequencing across several populations.
The origin of this inversion predates the divergence of North American (NA) and
Fennoscandian (North European) populations, indicating that it emerged in the ancestral *D.*
montana population in the Rocky Mountains of North America. Despite the species' expansion
across the northern hemisphere, this inversion has remained exclusive to the Rocky Mountain
populations, where it is fixed in the southernmost high-elevation population and appears at
lower frequencies in the more northern and lower-elevation populations. By independently
mapping SNPs associated with climate and cold adaptation and barrier regions that restrict gene
flow, we found that the inversion acts as a hotspot for both, with the barrier and adaptive
regions partially overlapping. Overall, the enrichment of adaptive and barrier loci, along with
its exclusive presence in Rocky Mountain populations, strongly suggests that chromosomal
inversions play a key role in environmental adaptation by restricting gene flow and facilitating
adaptive divergence between populations.

Keywords: barrier loci, climate adaptation, cold tolerance, demographic modelling, inversion
polymorphism, migration rate

Introduction

Barrier loci are defined as any loci that reduce gene exchange between populations, and they typically contribute to assortative mating and reduced fitness of hybrids (Butlin and Smadja 2018; Dopman et al. 2024). A central goal in evolutionary biology is to understand how barrier loci evolve. For example, do they begin as loci that promote local adaptation and become barrier loci, or are local adaptation and barrier loci independent? Can they become coupled by features of the genome, such as chromosomal inversions? Recent advances in genome sequencing of wild organisms have accelerated the development of computational tools to detect signatures of local adaptation, gene flow, and chromosomal rearrangements, enhancing our understanding of the selective, demographic, and molecular mechanisms driving population divergence despite ongoing gene flow.

Inversions, genomic regions with reversed gene order relative to the standard form, may play a major role in local adaptation and population divergence both with and without gene flow (reviewed in Hoffmann and Rieseberg 2008; Jackson 2011; Faria et al. 2019; Berdan et al. 2023). Inversions can gain a fitness advantage by significantly reducing recombination between individuals with different arrangements (Sturtevant 1921; Dobzhansky 1940), which allows for the accumulation and preservation of alleles with epistatic interactions (Hoffmann and Rieseberg 2008) and co-adapted gene complexes (Kirkpatrick and Barton 2006). The reduction in recombination is expected to be highest at the breakpoints, where it often extends beyond the inverted sequence (Kulathinal et al. 2009; Stevison et al. 2011), while gene flux that occurs via double crossovers and gene conversion can homogenize the regions towards the centre of the inversion (Navarro et al. 1997). Inversions may also be directly under selection if the breakpoints modify reading frames of genes, or if the new gene position, order or orientation, or changes in chromatin structure alter the expression of the genes (Villoutreix et al. 2021; Wright and Schaeffer 2022; Berdan et al. 2023). If there are no counteracting forces, these mechanisms can drive an inversion to high frequency, potentially leading to fixed differences between populations (Hoffmann and Rieseberg 2008; Berdan et al. 2023). On the other hand, inversion polymorphism within a species can be maintained through some form of balancing selection, such as heterozygote advantage (overdominance), or frequency-dependent or spatially or temporally varying selection. However, distinguishing the different mechanisms that act on inversions can be extremely difficult because multiple, sometimes interacting processes affect their evolution and because the different processes may result in similar genomic signatures (Berdan et al. 2023).

Showing empirically that alternatively fixed inversions between populations restrict gene flow and accumulate barrier loci remains challenging. Traditionally, outlier loci have been identified through summary statistics, like genetic divergence (d_{xy}) and genetic differentiation (F_{st}). However, these statistics can be misleading due to their susceptibility to different demographic, selective, and genetic factors, potentially resulting in false positives (Cruickshank and Hahn 2014; Ravinet et al. 2017; Laetsch et al. 2023). This limitation has led to the development of new methods that better account for these confounding factors and focus on detecting reduced migration as a signal of barrier regions (Fraïsse et al. 2021; Laetsch et al. 2023; Burban et al. 2024). While some studies have shown than mean migration rate (m_e) is lower within inverted compared to non-inverted (colinear) regions between populations connected by gene flow (e.g. Lohse et al. 2015; Poikela et al. 2024), studies have rarely investigated individual barrier regions and their association with rearrangements (Mackintosh et al. 2023). In light of recent studies, both inversions that have emerged at the onset of lineage divergence (Lohse et al. 2015) and inversions that predate lineage divergence and have existed already in the common ancestral population of the diverging lineages (Fuller et al. 2018; Poikela et al. 2024) can potentially drive divergence with gene flow.

To better understand the role of inversions in population divergence, it is essential to search for signs of local adaptation alongside barriers to gene flow. The presence of inversion frequency clines along environmental gradients, where balanced polymorphisms are maintained by spatially varying selection, was one of the first compelling pieces of evidence for the role of inversions in local adaptation (Dobzhansky and Levene 1948). Such clines have since been found in other systems (reviewed in Wellenreuther and Bernatchez 2018). Also, Quantitative Trait Loci (QTL) mapping and genome-wide association studies (GWAS) have proven effective in evaluating the role of inversions in adaptation. For example, Ayala et al. (2019) found that single nucleotide polymorphisms (SNPs) linked to desiccation resistance were located within the 2La and 2Rb inversions in *Anopheles gambiae*. It is also crucial to investigate the precise overlap between barrier loci and adaptive loci, as this may reveal whether adaptive loci act as barriers to gene flow as a result of ecologically-based divergent selection (Schluter 2001; Rundle and Nosil 2005). Finally, inversions may contribute to "coupling", where different barrier effects become linked through reduced recombination within inversions, ultimately promoting the overall barrier to gene flow (Butlin and Smadja 2018; Dopman et al. 2024).

The malt fly *Drosophila montana* is an ideal species for studying the role of chromosomal inversions in facilitating local adaptation and reducing gene flow between populations. It occupies a broad range of habitats, including extreme cold environments, shows reproductive barriers between populations, and possesses inversion polymorphisms. Spanning the northern hemisphere, its distribution ranges from Fennoscandia (Northern Europe) to Far East Asia near the Bering Strait, and North America (NA), with southern extensions into the Rocky Mountains (from elevations of 1400m to above 3000 m) and along the continents' western coast (Fig. 1A, Table S1; Patterson 1952; Throckmorton 1982; Hoikkala and Poikela 2022). The split between Fennoscandian and North American *D. montana* is estimated to have occurred ~1.75mya, soon after which the Crested Butte (NA) and other North American populations diverged (Garlovsky et al. 2020). Populations from these different regions show partial reproductive isolation (Jennings, Snook, et al. 2014). Additionally, *D. montana*, which has the highest recorded cold tolerance among *Drosophila* species (Kellermann et al. 2012; MacMillan et al. 2015) and can overwinter for up to six months under snow cover (Hoikkala and Poikela 2022), shows latitudinal variation in cold tolerance across its distribution (Poikela et al. 2021; Wiberg et al. 2021). While previous polytene chromosome studies suggest inversion polymorphism in *D. montana* (Stone et al. 1960; Throckmorton 1982), these inversions have not yet been characterized and studied at the genomic level.

Here, we use long- and short-read whole-genome sequencing (WGS) data, along with climate and cold tolerance data, GWAS and demographic modelling, to characterize polymorphic inversions and to investigate their role in population adaptation and divergence. First, we examine variation in climate conditions and cold tolerance and characterize polymorphic inversions and their frequencies across the species' distribution range. Next, we explore whether a large polymorphic inversion on chromosome 4 is enriched for loci associated with climate or cold adaptation. Finally, we investigate the origin of this large polymorphic inversion, assess whether it is enriched for barriers to gene flow, and determine whether barrier loci overlap with adaptive loci. We hypothesize that inversions are important drivers of population divergence by reducing recombination and facilitating local adaptation and the build-up of barriers to gene flow.

Result and Discussion

D. montana flies live in diverse geographic and climatic conditions

D. montana samples were collected from sites in the western coast and Rocky Mountains of North America (NA), Fennoscandia, and Kamchatka (Fig. 1A; Table S1). To characterize the climatic environments each population has adapted to, we conducted a principal component analysis (PCA) based on 19 bioclimatic variables of the geographic coordinates of each sampling site (Fig. 1B; Table S2). Three PCs had Eigenvalues >1, but the first two had the highest contribution and explained more than 81% of the total variation (Fig. 1B; Table S3). PC1 separated the western coast populations from the others (Fig. 1B). The coastal populations have warmer annual and winter temperatures (bio1, bio6, bio9 and bio11), lower seasonal variation in temperatures (bio4 and bio7) and receive higher precipitation throughout the year (bio12-13, bio15-16, bio19) compared to the other populations (Fig. 1B, Table S2, S4). PC2 separated the southern and northern coastal populations, as well as the Rocky Mountains populations from the Fennoscandian and Kamchatkan populations (Fig. 1B). These populations differed in daily temperature changes (bio2), isothermality (bio3, the ratio of day-to-night temperature variation to summer-to-winter variation) and variation in temperature and precipitation during warm months (bio5 and bio18; Table S4). In conclusion, *D. montana* has adapted to highly variable climates in its circumboreal distribution.

Figure 1. (A) *D. montana* sampling sites in the western coast and in the Rocky Mountains of North America, in Fennoscandia, and in Kamchatka (Asia). The Arctic circle is marked with a dashed line. (B) Climatic conditions across *D. montana* sampling sites summarized in a PCA using information on 19 bioclimatic variables (Table S2) of each fly collecting site.

D. montana cold tolerance traits show latitudinal variation and are partially correlated

An ability to cope with cold temperatures is one of the most important adaptations to high latitudes and altitudes. To examine clinal variation in cold tolerance, we studied cold tolerance traits in *D. montana* across its distribution range using nine different populations (Fig. 2, Table

S1). Cold tolerance was assessed using two ecologically relevant and widely used methods:
critical thermal minimum (CT_{min}) and chill-coma recovery time (CCRT). CT_{min} represents the
lowest temperature the fly can withstand without falling into chill-coma, while CCRT measures
the time required for flies to recover from a chill-coma. Lower values in both tests indicate
better cold tolerance.

The best-fit model for the traits was chosen based on Akaike's information criterion (AIC;
Table S5, Fig. 2A-B). Flies' ability to resist cold temperatures (CT_{min}) improves towards
northern (higher/colder) latitudes, and the positive interaction term between latitude and
altitude suggests that the effect of latitude on CT_{min} becomes more pronounced as altitude
increases (Fig. 2A). In other words, flies' ability to resist cold temperatures improves towards
northern latitudes and higher altitudes. Flies' ability to recover from chill coma (CCRT) also
improves towards northern (higher/colder) latitudes, but the negative interaction term suggests
that the relationship between latitude and CCRT becomes weaker as altitude increases (Fig.
2B). However, the significant interaction terms should be interpreted with caution since they
likely derive from one outlier population (Crested Butte) with an altitude of 2900m. Measured
on the same individuals, CT_{min} and CCRT were positively correlated (Fig. 2C, Table S6), but
this correlation was population-dependent and significant only in the northern western coast
populations (Seward and Terrace; Fig. 2C, Table S6).

Generally, populations living at higher latitudes or altitudes experience colder environments
than those at lower latitudes or altitudes, which is expected to lead to genetic differences and
clinal variation in cold tolerance traits (Addo-Bediako et al. 2000). The clinal variation
observed in *D. montana* supports previous findings in this species (Wiberg et al. 2021), and
contributes to the expanding evidence for clinal variation in cold tolerance across ectotherms
(e.g. Hoffmann et al. 2002; Overgaard et al. 2011; Zuther et al. 2012). While the discovery of
clinal variation in cold tolerance traits is expected, the partial correlation between CT_{min} and
CCRT is novel as it has not been observed before in *D. montana* (Wiberg et al. 2021) or other
species (e.g. Overgaard et al. 2011; Andersen et al. 2015). This correlation suggests that the
resistance to chill coma (CT_{min}) and the recovery from it (CCRT) may either share mechanisms
or coevolve in specific populations or environments, but not universally.

Figure 2. *D. montana* cold tolerance measured as (A) chill coma temperature (CT_{min}) and (B) chill coma recovery time (CCRT) across latitudes and altitudes. The points and error bars show the mean and standard errors, respectively. Dashed lines indicate the predicted values from the partial effect of latitude from the best-fit model (model comparison in Table S5). (C) Pearson correlation and regression lines between CT_{min} and CCRT across all nine populations (black dashed line), and separately for each population (all statistics in Table S6).

The Rocky Mountains populations harbour a large polymorphic inversion on chromosome 4

We characterized polymorphic inversions in *D. montana* using PacBio and Illumina re-sequence data and genome assemblies (Table S7-S10). We found one 9.5Mb polymorphic inversion located on chromosome 4 (Table S11, Fig. S1). This inversion was found only in the Rocky Mountains populations, being fixed in Crested Butte (100%) and present at lower frequencies in Jackson (17%) and McBride (22%) (Fig. 3, Table S11), though the inversion frequencies in Jackson and McBride are less reliable due to the low sample sizes (N = 3 and N = 9, respectively, Table S11). Overall, the inversion contains ~850 genes; those closest to the proximal breakpoint include *tRNA-specific adenosine deaminase* and *UPF0428 protein CG16865*, while those near the distal breakpoint include *sorting nexin-17* and *tRNA-dihydrouridine(20a/20b) synthase [NAD(P)+]-like* (see the exact locations in Fig. S1).

We also found a small inversion of approximately 1500bp at the proximal end of chromosome 4 (Table S12, Fig. S1). This inversion was present only in the Fennoscandian and Kamchatkan samples and is located in the intronic region of a gene coding for a leucine-rich repeat

transmembrane neuronal protein 3. Given its small size and the synonymous effects on the
protein product, it was not investigated further. Older polytene chromosome studies (Stone et
al. 1960; Throckmorton 1982) suggest that *D. montana* may harbor additional polymorphic
inversions not identified here, possibly due to the limited number of strains sequenced.

Tahami et al. (2024) found Gypsy and MITE transposable elements (TEs) at the breakpoints
of the large chromosome 4 inversion fixed in Crested Butte, which are absent in populations
without the inversion. Similarly, the small chromosome 4 inversion in Fennoscandian
populations contains unique Helitron and Gypsy TE insertions. These findings suggest that TE
insertions near breakpoints may have facilitated the origin of the inversions through ectopic
recombination, although further research is needed to confirm this. We explore the role of the
large chromosome 4 inversion in local adaptation and population divergence in the following
sections.

220
221 Figure 3. Frequencies of the large polymorphic inversion (9.5Mb) on chromosome 4. The inversion is
222 fixed in the southernmost population of the Rocky Mountains (Crested Butte) and occurs at lower
223 frequencies in the more northern Rocky Mountains populations (Jackson, McBride). It is absent in all
224 other studied North American, Asian and Fennoscandian populations (see Table S11 for details).

Geography and the polymorphic chromosome 4 inversion contribute to the population genetic structure of *D. montana*

To summarize the population genetic structure of *D. montana* and the role of geography and
the polymorphic inversion therein, we conducted PCAs for the whole genome and for the
inversion using the Illumina re-sequence data of 101 wild-caught females (Fig. 1A, Table S1,
S8). The first PCA included 1,986,534 filtered SNPs across the genome. The first three
principal components (PCs) explained 35%, 22% and 6% of the genetic variation (Fig. 4A;
Table S13). PC1 and PC2 revealed four major clusters corresponding to Crested Butte and
other North American (NA) populations, Fennoscandian populations, and Kamchatka (Asia).
PC3 showed a latitudinal cline within North American populations, which was primarily due
to genetic variation on chromosome 4 (Fig. S2E). The second PCA included 166,193 filtered
SNPs from the chromosome 4 inversion, revealing similar patterns to the genome-wide PCA
(Fig. 4B, Table S13), with some key differences. Here, Jackson and McBride samples that carry
the inversion in either homozygous or heterozygous states clustered more closely with Crested
Butte, where the inversion is fixed (Fig. 4B, Table S11). Notably, a homozygous inversion
carrier from McBride, clustered closest with Crested Butte, while heterozygous inversion
carriers were more distant. This pattern was not observed on colinear regions on chromosome
4 (Fig. S2, Table S13). Hence, genetic variants in the inversion are maintained across
populations.

We also measured genetic diversity (π) for each population, and divergence (d_{xy}) and
differentiation (F_{st}) between populations (Fig. 4C-E, Table S14-S15, Fig. S3-S5). North
American populations showed ~40% higher π than Fennoscandian populations, except for
Crested Butte (NA) which had the lowest diversity of all samples. Kamchatka had similarly
low diversity, although this may be unreliable because the samples were obtained from
isofemale strains, not directly from the wild. π was particularly high in the inverted region of
Jackson and McBride, reflecting their mixed inverted and non-inverted haplotypes (Fig. 4B-C,
Table S15). d_{xy} and F_{st} generally increased with geographic distance, following an isolation-
by-distance model, except for Kamchatka and Crested Butte. d_{xy} and F_{st} were highest between
North American and Fennoscandian populations, lower among North American populations,
and lowest within Fennoscandia. Despite its geographic proximity to North America,
Kamchatka was genetically closer to Fennoscandia, likely due to the isolation by the Bering
Strait. Crested Butte exhibited the highest genetic divergence from other populations,
regardless of geographic location, indicating that it is a relatively isolated population. Finally,
d_{xy} and F_{st} between Crested Butte, where the inversion is fixed, and populations without the
inversion were higher in the inverted regions compared to the colinear regions (Fig. 4D-G,
Table S15). Although d_{xy} and F_{st} were also elevated in the inverted regions between Crested
Butte and the Rocky Mountains populations (Jackson and McBride), the values were not as
high due to the presence of the inversion at lower frequencies in Jackson and McBride (Fig.
4D-G, Table S15).

Taken together, genetic variation in Crested Butte, other North American populations,
Fennoscandia, and Kamchatka contribute to large-scale genetic variation across *D. montana*'s
range, while the chromosome 4 inversion is one of the key drivers of finer-scale genetic
variation in *D. montana*. The phylogeographic history of the species has been uncertain, with
hypotheses suggesting either a North American origin, followed by a spread to Asia and Europe,
or an Asian origin, with subsequent spread to North America and Europe (Throckmorton 1982;
Mirol et al. 2007; Garlovsky et al. 2020). Given the highest genetic diversity observed in North
America, which gradually decreases towards Alaska (NA) and Northern Europe (Fennoscandia)
(Table S14-S15), along with the presence of all *montana* complex species in North America
(Throckmorton 1982; Hoikkala and Poikela 2022), it is likely that *D. montana* originated in
North America and spread into Asia and Europe from there. The low genetic diversity in
Crested Butte (NA) may indicate a founder effect or population bottleneck, which could
explain the fixation of the inversion in that population, while not in Jackson or McBride (NA).
This inversion may still play a key role in local adaptation, for example, by reducing
recombination and gene flow between populations, as indicated by the increased genetic
divergence within the inversion, a topic we will explore in the following sections.

Figure 4. Genetic population structure of *D. montana*. Genetic variation was summarized in principal component analyses (PCA) including SNPs (A) from all chromosomes and (B) from the large inversion on chromosome 4. The inversion is fixed in Crested Butte (marked with *), i.e. all samples are homozygous for the inversion (*INVhom). The Jackson and McBride samples that carry the inversion at homozygous (INVhom) or heterozygous (INVhet) state are marked separately; see Table S11). (C) Mean genetic diversity (π) was calculated for each population and for inverted (INV) and non-inverted (colinear, COL) regions. Mean (D) genetic divergence (d_{xy}) and (E) genetic differentiation (F_{st}) were calculated between Crested Butte (inversion fixed) and other populations (inversion at lower frequencies in Jackson and McBride and absent in the other populations) for INV and COL regions. (F) d_{xy} and (G) F_{st} were plotted in sliding windows of a fixed block size between Crested Butte and the other populations (125 blocks × blocks of 256bp in length, with a step change of 25 blocks × blocks of 256bp in length). Populations are highlighted with dashed lines whenever their divergence strongly deviates from that of the other populations of the same geographic region. The inversion breakpoints are marked with vertical solid lines. Since recombination is often suppressed beyond the inverted region into the colinear region, 1.5Mb buffer region was marked at both ends of the inversion with vertical

296   dashed lines. Kamchatka samples were excluded from the d_{xy} and F_{st} window-wise plots since they were
297   collected from inbred laboratory strains.

SNPs associated with climate variability and cold tolerance are enriched within the inversion

To explore the role of the large chromosome 4 inversion in accumulating and preserving loci
under local adaptation, we linked genomic data to climatic conditions and experimental cold
tolerance data. This was achieved through a GWAS using Illumina pool-sequencing data from
six populations (Table S9), combined with population-specific environmental variables (PC1,
PC2; Fig. 1B) and cold tolerance traits (CT_{min}, CCRT; Fig. 2A-B). The inversion is fixed in the
Crested Butte population but absent in the other three North American and two Fennoscandian
populations. The recombination-suppressing effects of inversions can extend up to 2-3Mb
beyond the inverted sequence into the colinear regions (Kulathinal et al. 2009; Stevison et al.
2011). Therefore, in addition to exploring SNPs strictly within the inversion breakpoints, we
extended the inversion by a conservative estimate of 1.5Mb at both ends since there is no prior
knowledge on the effect of inversions on recombination in *D. montana*.

Across different genomic regions, only 0.05-1.38% of SNPs were significantly associated with
environmental variables (PC1, PC2) or cold tolerance phenotypes (CT_{min}, CCRT; Fig. 5, Fig.
S6, Table S16). The inversion exhibited a significantly higher enrichment of SNPs associated
with CCRT compared to colinear regions, and this enrichment persisted regardless of whether
the region analyzed was limited to the inversion breakpoints (Chi Square: $X^2=5.1$, P=0.023) or
extended 1.5Mb beyond them into the colinear regions ($X^2=1228.7$, P<0.001; Fig. 5, Fig. S6D,
Table S17). The inversion was also enriched for SNPs associated with PC1 ($X^2=754.3$,
P<0.001), PC2 ($X^2=142.9$, P<0.001) and CT_{min} ($X^2=1058.7$, P<0.001), although these were
significant only when extending 1.5Mb beyond the inversion breakpoints (Fig. 5, Fig. S6D,
Table S17). Additionally, SNPs associated with PC2 showed two narrow peaks near the
telomeric end of chromosome 4, and the X chromosome displayed a strong enrichment of SNPs
associated with all four variables (Fig. 5, Fig. S6, Table S16).

323
Figure 5. Genome-wide association studies (GWAS; BayeScEnv). Points above the horizontal line
indicate significant SNPs, describing significant association between (A) genetic and environmental
(PC1=dark purple, PC2=light green) differentiation and (B) genetic differentiation and cold tolerance

(CT$_{min}$= dark purple, CCRT=light green; see separate plots in Fig. S7). q-values for the g parameter of
two chains (Fig. S7-S10) were used to control the false positive rate at 0.05, i.e. SNPs with q-values>1.3,
to obtain the final candidate SNPs. The solid vertical lines represent the inversion breakpoints, while
the dashed vertical lines mark a conservative estimate of 1.5Mb beyond the breakpoints, where
recombination is expected to remain reduced.

The significant SNPs inside the inversion were distributed across 104-239 genes, depending
on the analyzed dataset (Table S18). Based on the gene ontology (GO) enrichment analysis,
the genes linked to all four variables were significantly associated with calcium transport,
membrane and transmembrane functions, and PC2 and CCRT were additionally associated
with immunoglobulin (Ig)-like domain (Table S18, Supplementary file 1). Many of these
groups have been repeatedly linked to the ability of *D. montana* to withstand cold temperatures
(Kauranen et al. 2019; Parker et al. 2021; Wiberg et al. 2021), and genes connected to
membrane, transmembrane and immunoglobulins were among the fastest evolving when
comparing several cold-tolerant *Drosophila* species for their rates of molecular evolution
(Parker et al. 2018). Membrane proteins play a crucial role in membrane and cuticular functions,
which are especially important in cold temperatures and desiccation tolerance (Gibbs 2002;
Stinziano et al. 2015). Additionally, cell functioning at low temperatures is strongly dependent
on changes in ion transport and fluidity (Hazel 1995; Teets and Denlinger 2013).

Overall, the accumulation of SNPs linked to climate and cold adaptation within the inversion
and near its breakpoints (Fig. 5), along with its exclusive presence in the Rocky Mountains
populations (Fig. 3), suggests that the inversion plays a role in local adaptation. However, it
remains unclear whether this local adaptation has driven the fixation of the inversion in the
high-altitude Crested Butte population, or if the fixation is due to other factors not related to
selection, such as a population bottleneck or founder effect. In contrast, the inversion is
maintained in a polymorphic state in the more northern, lower-altitude Rocky Mountains
populations (Jackson and McBride), but the type of balancing selection responsible for this
remains unknown. For instance, the varying inversion frequencies may provide seasonal
advantages by adjusting stress resistance, thereby helping these mountainous populations better
cope with significant environmental fluctuations (Fig. 1B).

The inversion is old and is enriched with barrier loci

To investigate the evolutionary history of *D. montana* and the role of the large chromosome 4
inversion in population divergence, we used a demographic modelling approach (gIMble;
Laetsch et al. 2023). First, we compared and identified the best-fit demographic model between
three genetically, climatically, and geographically isolated populations: Crested Butte (North
America, inversion fixed), Vancouver (North America, inversion absent), and Storslett
(Fennoscandia, inversion absent; Fig. 1, 3). We tested models both without (strict divergence,
DIV) and with post-divergence gene flow in both directions (isolation with migration, IM).
This initial model selection focused on colinear regions, as the evolutionary history of
inversions may differ from other genomic regions. The best-fit demographic scenario for
Storslett-Crested Butte and Vancouver-Crested Butte comparisons suggests post-divergence
gene flow (IM model), with gene flow occurring from Storslett and Vancouver to Crested Butte
(forward in time; Table S19). Our parametric bootstrap analysis showed that the IM model
significantly improved the fit compared to the DIV model in these comparisons, indicating a
genuine signal of post-divergence gene flow (Fig. S11). However, while the IM model had
higher support than the DIV model in the Storslett-Vancouver comparison (Table S19), the
parametric bootstrap analysis showed that the improvement in fit was not significant (see Fig.
S11). Consequently, the DIV model could not be rejected for this comparison. Although *D.*
montana likely originated in North America and later spread to Asia and Europe (as discussed

earlier), post-divergence gene flow can occur in multiple directions, reflecting more recent gene exchange rather than the original migration patterns. The similar climate conditions in Northern Europe (Storslett) and the Rocky Mountains of North America (Crested Butte; see Fig. 1B) may explain the gene flow from Northern Europe to the Rocky Mountains. Conversely, the lack of gene flow from Storslett to Vancouver could be due to climate differences, as Vancouver is a low-latitude, low-altitude population (Fig. 1B). This is also supported by the observation that in Europe this species has not spread to southern locations.

Second, the best-fit model of each population pair (Table S19) was used to analyse the mean time of divergence (T) and migration rate (m_e) separately for inverted and colinear chromosomal regions to estimate the age of the inversion and its effects on migration rate (Fig. 6A-B, Table 1). Based on the colinear background, the North American (Crested Butte, Vancouver) and Fennoscandian (Storslett) populations of *D. montana* diverged approximately 700,000 to 1,000,000 years ago (Fig. 6A, Table 1). The estimated divergence time for the inverted region was approximately 1.4 million years between the Crested Butte population, where the inversion is fixed, and the Vancouver and Storslett populations, where the inversion is absent (Fig. 6A, Table 1), suggesting that the inversion was already present in the ancestral population before *D. montana* populations began to diverge. Notably, the colinear and inverted regions between Vancouver and Storslett – neither of which carry the inversion – diverged around the same time (Fig. 6A, Table 1). Additionally, the mean migration rate estimate was significantly lower within the inverted compared to colinear regions between Crested Butte and Vancouver, where gene flow has been strongest and may still be ongoing (Fig. 6B, Table 1). Migration rate estimates were very low for both inverted and colinear regions between Crested Butte and Storslett (Fig. 6B, Table 1), likely due to strong geographic isolation reducing gene flow across the genome, not solely within the inversion. Since the strict divergence (DIV) scenario between Vancouver and Storslett could not be rejected, m_e was estimated to be zero for both inverted and colinear regions (Fig. 6B, Table 1). Overall, the inversion likely has not played a major role in the divergence between North American and Fennoscandian populations, as none of the Fennoscandian populations and most North American populations do not carry the inversion, and the strong geographic isolation has reduced gene flow across the entire genome.

Table 1. Parameter for effective population sizes (N_e), divergence times (T in years/generations) and migration rates (m_e) for three *D. montana* populations pairs estimated from 256bp blocks under the best-fit IM model (Table S16), including intergenic non-repetitive regions to minimize the direct effects of selection. m_e estimates correspond to M ($=4N_e m_e$) individuals per generation (forwards in time). The chromosome 4 inversion is fixed in Crested Butte and absent in the other populations. The m_e estimates were inferred under the best-fit IM model, except for the Vancouver-Storslett comparison where the m_e estimates were based on the best-fit DIV model that assumes no post-divergence gene flow (m_e=0).

Population	Region	Ancestral N_e	Pop. A N_e	Pop. B N_e	T (years)	m_e (M)
Crested Butte (A) - Vancouver (B)	4 INV	646,563	179,131	1,137,026	1,347,905	2.06E-07 (0.15)
IM *Vancouver --> Crested Butte*	COL	438,089	147,816	1,070,603	958,812	4.95E-07 (0.29)
Crested Butte (A) - Storslett (B)	4 INV	647,591	203,253	670,724	1,407,776	1.43E-07 (0.12)
IM *Storslett --> Crested Butte*	COL	474,766	198,026	471,474	1,016,841	2.19E-07 (0.17)
Storslett (A) - Vancouver (B)	4 INV	808,211	561,850	1,159,870	727,619	0.0 (0.0)
IM *Vancouver --> Storslett*	COL	527,819	413,484	1,228,026	651,134	0.0 (0.0)

Figure 6. Evolutionary history of *D. montana* populations and the polymorphic inversion on chromosome 4. (A) Mean divergence time (T) and (B) mean migration rate (m_e) of inverted (4INV) and colinear (COL) regions between three *D. montana* populations. The m_e estimates were inferred under the best-fit IM model, except for Vancouver-Storslett comparison where the m_e estimates were based on the DIV model that assumes no post-divergence gene flow (m_e=0; see Table 1). Crested Butte, the population with the fixed inversion, is highlighted with * (absent in the other populations). Confidence intervals were estimated using the parametric bootstrap as +/- 2 SD across the 100 replicates under the best-fit IM model with recombination (see Methods). (C) Support for locally reduced m_e between Crested Butte and Vancouver was measured via positive Δ_{B0} (data points above the dashed horizontal line) and a false positive rate at 5% (more detailed plot in Fig. S11). Barrier regions ($\Delta_{B0} > 0$ and FPR ≤ 0.05) are marked with black segments. The solid vertical lines represent the inversion breakpoints, while the dashed vertical lines mark a conservative estimate of 1.5Mb beyond the breakpoints, where recombination is expected to remain reduced.

Finally, gIMble was used to assess window-wise m_e between Crested Butte (inversion fixed) and Vancouver (inversion is absent) to identify whether the inversion harbors barriers to gene flow, characterized by reduced m_e. As before, the barriers were explored strictly within the inversion breakpoints and by extending the inversion by 1.5Mb at both ends to account for the recombination-suppressing effects of inversions. Support for reduced m_e was indicated by positive Δ_{B0} values with a false positive rate (FPR) of 5%. Overall, 11.1% (1,050 out of 9,468) of the windows showed $\Delta_{B0}>0$, indicating potential barriers to gene flow (Fig. 6C). After accounting for an FPR of 5%, 44.4% (466 out of 1,050) of these windows passed the FPR criteria (Fig. 6C). By merging the remaining overlapping windows, a total of 10.7% (15.6Mb out of 145.5Mb) of the genome comprised barriers, distributed in 132 regions across the genome (Fig. 6C, Table S20). The barriers only partially overlapped with d_{xy} and F_{st} outliers (Fig. S12), a pattern also observed in *Heliconius melpomene* and *H. cydno* butterflies (Laetsch et al. 2023), highlighting the need for demographic modelling approaches to identify outlier barrier regions.

Notably, 62.9% (6.0 Mb out of 9.5 Mb) of the large inversion on chromosome 4 constituted
barriers, compared to 7.0% (9.6 Mb out of 135.9 Mb) of the colinear regions across
chromosomes. This difference was statistically significant, as determined by the number of
barrier and non-barrier windows in inverted and colinear regions (Chi-Square: $X^2 = 309.9$, P <
0.0001; Fig. 6C, Table S20-S21). This pattern remained consistent when extending the inverted
region beyond the breakpoints into the adjacent colinear regions (Chi-Square: $X^2 = 337.4$, P <
0.0001; Table S20-S21).

How closely associated are barrier and local adaptation loci? SNPs linked to PC1 and CCRT
overlapped with the barrier regions more than expected by chance, whereas SNPs linked to
PC2 and CT_{min} did not show significant overlap (Fig. 7). This suggests that loci underlying
local adaptation may at least partly act as barrier loci, reducing gene flow between populations.

452
Figure 7. The observed (vertical dashed lines) and simulated (distributions) overlap between barrier
windows and SNPs significantly associated with (A) PC1, (B) PC2, (C) CT_{min} and (D) CCRT. The
significance of the circular resampling is indicated in each plot.

Our results suggest that *D. montana* populations from Crested Butte (NA), Vancouver (NA)
and Storslett (Fennoscandia) diverged approximately 700,000-1,000,000 years ago. This
estimate aligns with a recent estimate of 1.75mya (Garlovsky et al. 2020) if we assume two
generations per year instead of one. The origin of the chromosome 4 inversion predates these
population split times based on the colinear background, indicating that the inversion originated
in the ancestral *D. montana* population before it spread across North America, Asia, and
Northern Europe. Similarly, species-specific inversions between sibling species pairs
Drosophila persimilis and *D. pseudoobscura* (Fuller et al. 2018) and *D. montana* and *D.*
flavomontana (Poikela et al. 2024) originated in the ancestral population before species
divergence. Interestingly, the polymorphic *D. montana* inversion, which has remained
restricted to the North American Rocky Mountain populations, is enriched with both adaptive
and barrier loci, potentially through coupling. The significant overlap between these loci
suggests that genetic regions associated with climate and cold adaptation may have reduced
gene flow between populations, acting as a barrier, potentially due to ecologically-based
divergent selection (Schluter 2001; Rundle and Nosil 2005). Other features of *D. montana* also
act as potential barriers to gene flow and may contribute to the barrier loci we have identified.
For instance, *D. montana* populations exhibit partial sexual and postmating-prezygotic (PMPZ)

isolation (Jennings, Snook, et al. 2014), likely driven by differences in courtship songs (Routtu
et al. 2007), cuticular hydrocarbons (CHCs; Jennings, Etges, et al. 2014), and reproductive
proteins (Garlovsky et al. 2020). While these findings suggest that the selective advantage of
the inversion arises explicitly from suppressed recombination, it could also result from changes
in gene expression. This opens the door to future research comparing inversion genotypes
(inverted and non-inverted homozygous, as well as heterozygous individuals) to identify
expression changes associated with the inversion and responses to environmental stressors such
as cold temperatures.

Conclusions

Chromosomal inversions have been suggested to influence population adaptation and
divergence by reducing gene flow and maintaining adaptive allelic combinations in linkage
between populations or ecotypes with different arrangements. However, determining whether
and how chromosomal inversions facilitate population adaptation and divergence is
challenging due to sparse empirical evidence (Faria and Navarro 2010; Fuller et al. 2019) and
the complexity of interacting processes that produce similar genomic signatures (Berdan et al.
2023). In this study, we characterize and investigate a large (~9.5Mb) polymorphic inversion
in *Drosophila montana*, a cold-adapted and widely distributed insect species.

The inversion's origin predates the divergence of North American and Fennoscandian
populations, suggesting it emerged in the ancestral *D. montana* population in the North
American Rocky Mountains. Despite the species' spread across the northern hemisphere, this
inversion remains exclusive to the Rocky Mountains populations, being fixed in the high-
altitude Crested Butte population and present at lower frequencies in the northern, lower-
altitude Jackson and McBride populations. We independently mapped SNPs associated with
climate and cold adaptation and barriers to gene flow, finding that the inversion serves as a
hotspot for both, with partial overlap between barrier and adaptive regions. This enrichment,
along with its exclusive presence in the Rocky Mountain populations, strongly supports the
role of chromosomal inversions in environmental adaptation and speciation by restricting gene
flow, facilitating adaptive divergence, and contributing to the coupling of different barriers to
gene flow.

Material and methods

D. montana sampling sites and overview of the data

D. montana samples were collected altogether from 13 geographically and climatically diverse sites in North America, Fennoscandia, and Asia in 2013−2020 (Fig. 1, Table S1). The newly collected females were brought to the fly laboratory in the University of Jyväskylä (Finland), and were kept in constant light, 19±1 ˚C and ~60% humidity. The females were allowed to lay eggs in malt vials for several days to produce the next generation, and to establish isofemale strains. The wild-caught females and their F_1 daughters were then stored in 70% EtOH at -20 ˚C. Furthermore, mass-bred populations were established from the F_4-progenies of 16-20 isofemale strains (320-400 flies) per population.

The mass-bred populations or isofemale strains were used in the experiments measuring fly cold tolerance (Table S1). PacBio (Pacific BioSciences) long-read sequencing data was generated from pools of females from isofemale strains (Table S1) and used to characterize chromosomal inversions. Finally, Illumina short-read sequencing data was generated from wild-caught females or their F_1-daughters collected and stored in 2013−2020 (Table S1). Illumina data were used to characterize chromosomal inversions and in genomic analyses. Details of the samples are described later in the text.

Climatic conditions of *D. montana* sampling sites

A principal component analysis (PCA) was used to summarize differences in climatic conditions between the 13 *D. montana* sampling sites. For this, 19 bioclimatic variables of each sampling site were extracted from WorldClim database v2.1 (2.5min spatial resolution, dataset 1970-2000, http://www.worldclim.org, Fick and Hijmans 2017) using latitudinal and longitudinal coordinates of each site (Table S1) and "raster" R package v2.8-19 (Hijmans et al. 2015; Table S2). The PCA was performed for the 19 bioclimatic variables of each population using the "FactoMineR" R package (Lê et al. 2008). All R analyses were conducted in R (v4.2.0) and R studio (v2023.03.0).

Assessment of *D. montana* cold tolerance

Variation in *D. montana* cold tolerance was assessed using two commonly used and ecologically relevant methods, chill coma temperature (CT_{min}, also called critical thermal minimum) and chill coma recovery time (CCRT) (MacMillan and Sinclair 2011; Overgaard et al. 2011). CT_{min} describes insects' ability to resist cold temperatures until falling into a reversible paralytic state, chill-coma, and CCRT their ability to recover from the chill-coma.

Fly cold tolerance was measured for nine *D. montana* populations: Seward, Terrace, Vancouver, Ashford, McBride, Crested Butte, Storslett, Oulanka and Korpilahti (Fig. 1, Table S1). Flies were collected from the mass-bred populations, apart from Oulanka for which the flies were collected from four isofemale strains. All flies had been maintained in the laboratory for 63−70 generations prior the experiments, except for Storslett and Oulanka which were maintained in the laboratory for 19 generations prior the experiments. To prevent the flies from entering reproductive diapause, they were maintained at constant light and 19±1°C prior and throughout the experiments. Flies were provided fresh malt-medium once a week. Only female cold tolerance was measured, but variation between *D. montana* females and males has been studied in Wiberg et al. (2021). Females were collected using light CO_2 anaesthesia within 72h after their emergence, placed in malt-vials, and used in cold tolerance experiments when sexually mature (18–23 days old, e.g. Jennings, Snook, et al. 2014).

A combined CT_{min}-CCRT test was performed by placing females of each population
individually in glass vials sealed with parafilm, which were then submerged into a 30% glycol
bath. The CT_{min} test was initiated by decreasing the bath temperature from 19 °C to -6 °C at
the rate of 0.5 °C/min, and the temperature at which a fly can no longer resist cold and loses
neurophysiological activity and coordination and falls into a chill coma was scored (MacMillan
and Sinclair 2011). After the CT_{min} test, the flies were left in -6 °C for 16 hours, after which
they were returned into room temperature and the time required for each fly to recover from
the chill coma and stand on its legs was measured (MacMillan and Sinclair 2011; Vesala et al.
2012). The tests were performed using Julabo F32-HL Refrigerated/Heating Circulator. The
assays were done in 13 batches, each consisting of 19-42 females that were split between
populations. For each population 40-44 females were tested, except for the Oulanka population
where 16-26 females from each of the four isofemale strain were tested (altogether 413
females).

In statistical tests, CT_{min} data (Celsius degrees + 10 ˚C because Gamma distribution does not
allow negative values) and CCRT data (seconds) were used as response variables. Since the
data was not normally distributed (Shapiro-Wilk test: CT_{min}; W = 0.974, P < 0.001, CCRT; W
= 0.851, P < 0.001), they were analysed with a generalized linear mixed model (GLMM) with
Gamma distribution, using *glmmTMB* function from glmmTMB R package (Brooks et al.
2017). To define the relevant fixed effects (latitude, altitude and their interaction) and random
effects (experimental batch and room temperature during the experiment), we performed model
selection first on the fixed effects, and then for the random effects using the best-fit fixed effects.
Th best-fit model for CT_{min} and CCRT was chosen based on Akaike information criterion (AIC)
value and Akaike weight using *aictab* function from AICcmodavg R package (Mazerolle and
Mazerolle 2020). The best-fit model for CT_{min} included latitude, altitude, their interaction as
fixed effects and experimental batch and room temperature as random effects, while the best-
fit model for CCRT included latitude, altitude, their interaction as fixed effects, with no random
effects.

Genomic data

PacBio long-read sequencing data from 11 *D. montana* populations (60 females from one
isofemale strain per population; Table S1, S7), Illumina short-read sequencing data of single
females from 13 *D. montana* populations (3-11 females per population, totalling 101 samples;
Table S1, S8) and Illumina pool-sequence data from six *D. montana* populations (each pool
containing 49-50 individuals; Table S1, S9) were used in the genomic analyses. Six of the 11
PacBio data sets were generated in the current study, but other PacBio and all Illumina data
sets were first published by Wiberg et al. (2021), Poikela et al. (2024), and Tahami et al. (2024).
Details of DNA extraction methods and sequencing technologies are given in these
publications, but the relevant information is summarized in Table S7-S9. For PacBio data,
average raw read coverage for each sample varied between 14-77X (Table S7). For Illumina
single female data, average coverage for each sample varied between 12-97X, except for
monSE13F37 sample which was sequenced to 435X (Table S8). For the Illumina pool-
sequence data the average coverage for each sample varied between 99-234X (Table S9).

D. montana chromosome-level reference genome (Seward) and the respective gene and repeat
annotations were obtained from Poikela et al. (2024). To characterize inversions, contig-level
assemblies of *D. montana* originating from Seward, Jackson, Crested Butte, Oulanka and
Kamchatka were obtained from Poikela et al. (2024) and Tahami et al. (2024). Assembly
statistics are given in Table S10.

Characterization of inversions and their frequencies

The presence of polymorphic *D. montana* inversions was investigated i) by mapping-based approaches and ii) by aligning assembled genomes of different populations. First, the PacBio raw reads of each of the 11 samples (one per population, except no data from Vancouver and Korpilahti; Table S7) were mapped against the contig-level Seward genome with the highest contiguity (Table S10) using ngmlr v0.2.7 (Sedlazeck et al. 2018). The resulting BAM-files were sorted with SAMtools v1.16.1 (Li et al. 2009) and structural variant (SV) candidates were obtained from Sniffles v2.0.7 (Sedlazeck et al. 2018). Also, Illumina paired-end reads of 101 single wild-caught samples (3-11 samples per population; Table S8) and six pool-sequence samples (Table S9) were quality-checked using FastQC v0.11.8 (Andrews 2010), trimmed for adapter contamination and low-quality bases using fastp v0.20.0 (Chen et al. 2018) and mapped against the contig-level Seward genome using BWA mem v0.7.17 with read group information (Li and Durbin 2009). The resulting BAM-files were sorted with SAMtools and PCR duplicates marked with sambamba v0.7.0 (Tarasov et al. 2015). The BAM-files were given to Delly v0.8.7 for SV identification (Rausch et al. 2012). SV signals from Sniffles (PacBio) and Delly (Illumina) were combined using SURVIVOR (Jeffares et al. 2017) to identify inversion candidates and their frequencies. The putative breakpoints of inversions were confirmed visually with IGV v2.8.0 (Thorvaldsdottir et al. 2013) using both long- and short-read data (Example view in Fig. S1). Second, alignments between *D. montana* contig-level assemblies (Table S10) were generated using nucmer of MUMmer package (Marçais et al. 2018) together with Dot plots (https://dot.sandbox.bio/). Finally, the breakpoints of reliable inversion candidates were mapped against the chromosome-level *D. montana* reference genome. Downstream analyses were focused on the large 9.5Mb inversion on chromosome 4 (Table S11), since the other characterized polymorphic inversion was tiny (~1500bp; Table S12).

Genetic diversity and divergence of *D. montana* populations

To define the genetic architecture of *D. montana*, the genetic variation of 101 Illumina single female samples was summarized using a principal component analysis (PCA). As explained in the previous paragraph, the Illumina raw reads were quality-checked, trimmed and mapped with read group information against the chromosome-level reference genome. The resulting BAM files were sorted, marked for PCR duplicates, and given to freebayes v1.3.6 for variant calling (Garrison and Marth 2012). The VCF file was filtered using preprocess function of gIMble with default settings (Laetsch et al. 2023). Variants with missing genotypes were excluded and the resulting VCF file was used to run the PCA i) for all chromosomes combined, ii) for each chromosome separately, iii) for the large chromosome 4 inversion (see Results), and iv) for the colinear regions of chromosome 4. All runs included coding and non-coding SNPs. The PCAs were run using PLINK's (v1.9) --pca function (Chang et al. 2015), by converting the VCF-file to PLINK's BED/BIM format and filtering SNPs and pruning them for linkage disequilibrium using PLINK's --indep-pairwise function.

Genetic diversity (π) within *D. montana* populations, and genetic divergence (d_{xy}) and differentiation (F_{st}) between pairwise comparisons of *D. montana* populations were investigated using the filtered VCF-file and info and windows modules of gIMble. π, d_{xy} and F_{st} were first analysed for coding, intronic and intergenic regions across the chromosomes to investigate the genetic architecture of *D. montana* in general, and then for the inverted and colinear intergenic regions to estimate the effects of the inversion on genetic diversity and differentiation/divergence. Repetitive regions were excluded from all data partitions as they may bias the results. Window-wise heterozygosity (H), d_{xy} and F_{st} (including coding, intronic and intergenic regions) were also plotted across the genome.

Signatures of local adaptation linked to climate conditions, cold tolerance and the chromosome 4 inversion

BayeScEnv is an F_{st} based genome-scan method that identifies loci linked to a given environmental variable or phenotype, while accounting for confounding effects of population structure (de Villemereuil and Gaggiotti 2015). We used the method to detect signatures of local adaptation linked to environmental conditions (PC1 and PC2 for each population) and cold phenotypes (mean CT_{min} and CCRT for each population) of *D. montana* populations. To maximize the power of the method, we used *D. montana* pool-seq data originating from four sites in North America (Seward, Terrace, Ashford and Crested Butte) and two sites in Fennoscandia (Oulanka and Korpilahti) (Table S1, S9) (Wiberg et al. 2021). The pool-sequence data were quality-checked and trimmed for adapter contamination and low-quality based as described before. Since coverage of the Seward pool was approximately twice as high as that of the other samples (Table S9), it was resampled to 94 million forward and reverse reads using seqtk (https://github.com/lh3/seqtk). Reads of each pool were mapped against the reference genome using BWA mem with read group information, and the alignments were sorted and marked for PCR duplicates, like before. The BAM-files of each population were filtered for mapping quality >30 using SAMtools. Allele counts for each population at each genomic position were obtained with SAMtools mpileup using options to keep reads with a mapping quality >30 and sites with a base quality >30. The resulting BAM-files were used for variant calling using a heuristic variant calling software PoolSNP (Kapun et al. 2020). PoolSNP settings included a minimum count, coverage and frequency of 15, 50 and 0.05, respectively, a missing fraction of 0.01, and 95% coverage percentile for a given population and chromosome was considered as a maximum coverage. Variant calling detected a total of 3,400,762 biallelic SNPs.

The VCF-file was given to BayeScEnv v1.1, which was run twice for each dataset. Each run consisted of five pilot runs of length of 1000, and 4000 iterations of which 2000 were excluded as burn-in. Because of the uneven number of males and the resulting variation in the number of X chromosomes in each pool (Table S9), BayeScEnv analysis was performed separately on autosomal and X chromosomal SNPs. Raw data were used for the autosomal SNPs. For X-linked SNPs, allele count data were scaled to the known number of X chromosomes in the pool (Table S9), following Wiberg et al. (2021). The MCMC chains were checked for convergence using coda R package v0.19-4.1 (Plummer et al. 2006), which was reached across the two chains for all analyses and parameters (potential scale reduction factors (PSRFs) of 1.00-1.02 in a Gelman-Rubin diagnostic test; Figures S7–S10). The union of significant SNPs, using q-values for the g parameter (FDR at 0.05, i.e. SNPs with q-values<0.05), were taken as the final candidate set. These SNPs describe the association between genetic differentiation and environmental or phenotypic differentiation.

The enrichment of significant SNPs was compared between inverted and colinear regions using a Chi-square test to identify whether adaptive loci have enriched within the inversion. The effect of inversions in suppressing recombination may extend up to 2-3Mb beyond the inverted sequence (Kulathinal et al. 2009; Stevison et al. 2011). Thus, the effect of the inversion on adaptive loci was tested with a) the inversion ending strictly at the breakpoints and b) the inversion extending from each breakpoint into the colinear regions by a conservative estimate of 1.5Mb, as the impact of inversions on recombination has not been studied in *D. montana*.

Finally, we performed GO-term enrichment analysis on the significant SNPs using DAVID 2021 (Huang, Sherman, Tan, Kir, et al. 2007; Huang, Sherman, Tan, Collins, et al. 2007). We extracted all genes containing significant SNPs within the chromosome 4 inversion and blasted them against *Drosophila virilis* RefSeq proteins (BLASTp 2.15.0+; Camacho et al. 2009),

687 because *D. montana* is more closely related to *D. virilis* (both belong to the *virilis* group; Yusuf
688 et al. 2022)) than to *D. melanogaster*. Functional annotation clustering of the best *D. virilis* hits
689 was carried out using DAVID.

Evolutionary history of *D. montana* and the inversion on chromosome 4

We used gIMble (Laetsch et al. 2023) to resolve the evolutionary history of *D. montana* and to
estimate when the large chromosome 4 inversion (see Results) originated and whether it has
reduced gene flow between populations. gIMble is a composite likelihood approach that uses
a joint distribution of mutation types in short sequence blocks, the blockwise site frequency
spectrum (bSFS), across pairs of individual genomes.

First, a "global" mode of gIMble was used to investigate the likely evolutionary history of three
genetically, geographically and climatically diverged *D. montana* populations, Vancouver (the
western coast of North America), Crested Butte (the Rocky Mountains of North America) and
Storslett (Fennoscandia), using the respective singly sequenced, wild-caught Illumina samples
(Fig. 1, Table S1, S8). For each population pair, three demographic scenarios were fitted: a
strict divergence without post-divergence gene flow (DIV, migration rate (m_e) = 0), and
isolation with migration with both gene flow directions between the populations ($IM_{monA \rightarrow monB}$,
$IM_{monB \rightarrow monA}$). This initial model selection was limited to intergenic, non-repetitive sequences
of colinear chromosomes by excluding sequences with gene and repeat annotations to minimize
the direct effects of selection. The large chromosome 4 inversion was excluded from the
analysis due to its potentially different evolutionary history compared to colinear regions
(Lohse et al. 2015; Fuller et al. 2018; Poikela et al. 2024), while colinear regions, ending at
breakpoints, were combined across the genome as these regions are expected to share the same
evolutionary history. Data was summarized by the bSFS for a block length of 256bp with kmax
values of 2, and blocks were grouped in sliding windows (--blocks 12,500, --steps 2,500). The
bSFS-based analyses assume no recombination within blocks and a constant mutation rate (μ)
across them. Based on an estimate of the spontaneous mutation rate in *Drosophila*
melanogaster (Keightley et al. 2014), $\mu = 2.8 \times 10^{-9}$ per base and generation was used. The
estimates of divergence time (*T*) were converted into absolute time using t = T×2N$_e$×g, where
N$_e$ = θ/(4μ) and g is generation time. One generation per year was assumed for each population
(Tyukmaeva et al. 2020). To evaluate the global support for an IM model versus a DIV model,
we conducted 100 simulations using msprime (Baumdicker et al. 2022) through gIMble
simulate, parameterised by the DIV history. Depending on the dataset, we simulated 8,343,
8,435 or 8,738 windows, matching the size of the real data, specifically 12,500 pair-blocks, for
10 diploid individuals per population, consistent with the empirical dataset. We assumed a
constant recombination rate, based on one crossover per female meiosis per chromosome
(recombination occurs only in females), resulting in map length of 25cM per chromosome. We
fitted both a DIV model and an IM model to each simulation and compared the null distribution
of lnCL to the observed empirical lnCL (see Fig. S11).

Second, a "global" mode of gIMble was used to estimate the time of divergence of the
polymorphic inversion on chromosome 4 and the mean migration rate inside and outside of the
inversion. For this, the best-fit model of each population pair was fitted separately for the
inverted and colinear regions using the same settings as before. Here, a parametric bootstrap
was performed to estimate the uncertainty of *T* and *M* parameters using gIMble simulate with
100 replicate datasets. These simulations used the same number of samples, block length, and
number of blocks per window as the real data, and assumed the same mutation and
recombination rates as before. Since the windows of the real data partially overlap and are in
linkage, whereas the simulated windows are assumed to be statistically independent, we
simulated datasets with 36 times fewer windows than the real data, resulting in approximately

500kb between windows, following Laetsch et al. (2023). Confidence intervals were estimated
using the parametric bootstrap as +/- 2 SD across the 100 replicates.

Third, a "local" mode of gIMble was used to identify regions of reduced gene flow, referred to
as barrier loci, across the genome between Crested Butte (inversion fixed) and Vancouver (no
inversion). gIMble accounts for background selection through heterogeneity in effective
population size (N_e) and barriers to gene flow through heterogeneity in migration rate (m_e). To
quantify the heterogeneity in m_e and N_e, a grid of parameters was fitted to sliding windows of
a fixed length of 12,500 pair-blocks with a step size of 2,500 pair-blocks along the genome.
This resulted in 9,468 windows with a minimum span of 32 kb. Since only intergenic blocks
(excluding repetitive sequences) were used, the window span was typically greater than 32 kb
(median window span 68.6 kb; Fig. S14). Variation in $N_e s$ and m_e across the windows were
explored by searching for parameter combinations in a 12×12×12×16 grid (N_e mon$_A$, N_e mon$_B$,
N_e ancestral, m_e; Fig. S15). Given that the inversion on chromosome 4 has diverged earlier, i.e.
is older than the colinear regions, the value for T was allowed to be either 958,812 or 1,347,905
(COL vs. 4INV T; see Table 1). The mutation rate was set to μ = 2.9 × 10^{-9} per base and
generation (Keightley et al. 2014). Windows were identified as barriers to gene flow when the
support for a DIV model (i.e. m_e = 0) was greater than the support for an IM model. The IM
model was parameterized with the most likely inferred effective migration rate under a null
model of m_e variation, indicated by $\Delta_{B0} > 0$. In other words, the local support for reduced gene
flow was measured via Δ_{B0}, where positive Δ_{B0} indicates strongly reduced gene flow. To
estimate the uncertainty in these local estimates, we used a parametric bootstrap approach.
Specifically, we simulated 100 datasets for each window under the best-fitting local
background history using the msprime software (Baumdicker et al. 2022) via the gIMble
simulate module, providing a null distribution of Δ_{B0} values. In these simulations, m_e was fixed
across windows at the global background level, while N_e parameters were allowed to vary
between windows based on the best composite log-likelihood N_e parameters inferred for each
window. Windows from the real data were labelled as barriers if their Δ_{B0} value was greater
than the highest Δ_{B0} value observed in the simulated null distribution, ensuring a false positive
rate (FPR) of ≤0.05. The same recombination rate was used in these simulations as before.
Finally, windows with significantly reduced gene flow ($\Delta_{B0}>0$ and FPR≤0.05) that overlapped
were merged to define barrier regions.

The number of the barrier and non-barrier windows (divided by five to obtain the number of
non-overlapping windows) was compared between the inverted and colinear regions using a
Chi-Square test to determine whether barriers are enriched within the inversion. Like in the
previous paragraph, the effect of the inversion on barrier loci was tested with a) the inversion
ending strictly at the breakpoints and b) the inversion extending from each breakpoint into the
colinear regions by a conservative estimate of 1.5Mb. Finally, to test whether SNPs
significantly associated with PC1, PC2, CT$_{min}$ or CCRT were enriched within barrier windows
(proportion of SNPs in barrier vs. non-barrier windows), we used a circular resampling
approach (https://github.com/LohseLab/circular_bootstrap/tree/main) by Ebdon et al. (2025),
where the observed estimate was compared to the distribution of means of 1000 resampled
datasets.

Acknowledgements

We want to thank Sara Lommi for her help with DNA extractions and cold tolerance experiments. This work was supported by an Academy of Finland project 322980 to M.K., and grants from Jenny and Antti Wihuri Foundation to N.P. Fig. 1A was created with Biorender.com. Fig. 3 was created with MapMixture (https://tomjenkins.shinyapps.io/mapmixture/).

Data Accessibility and Benefit-Sharing

Data accessibility statement

The new PacBio raw sequence data are available at SRA under BioProject PRJNA1237710. The published raw sequence data and genome assemblies were obtained from BioProject accessions PRJNA939085, PRJNA828433, and PRJNA588720. Cold tolerance data and all scripts will be made publicly available at https://github.com/noorlinnea upon publication.

Benefit-Sharing Statement

Not applicable.

Author contributions

MK and NP designed the study. NP and VH performed cold tolerance experiments. NP conducted statistical and genomic analyses, with input from VH and MGR. MK and MGR supervised, and MK funded the study. NP drafted the first version of the manuscript, and all authors edited and read the final draft.

ORCID

Noora Poikela – https://orcid.org/0000-0002-4627-9647
Ville Hoikkala – https://orcid.org/0000-0003-3995-3215
Michael G. Ritchie – https://orcid.org/0000-0001-7913-8675
Maaria Kankare – https://orcid.org/0000-0003-1541-9050

Ethics declaration

D. montana is not endangered, and the flies were collected along watersides on public lands outside National and State parks, where insect collecting does not require permits in the USA (The Wilderness Act of 1964, section 6302.15).

Conflicts of interest

None.

References

Addo-Bediako A, Chown SL, Gaston KJ. 2000. Thermal tolerance, climatic variability and latitude. *Proceedings of the Royal Society of London. Series B: Biological Sciences* 267:739–745.

Andersen JL, Manenti T, Sørensen JG, MacMillan HA, Loeschcke V, Overgaard J. 2015. How to assess *Drosophila* cold tolerance: chill coma temperature and lower lethal temperature are the best predictors of cold distribution limits. *Functional Ecology* 29:55–65.

Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data.

Ayala D, Zhang S, Chateau M, Fouet C, Morlais I, Costantini C, Hahn MW, Besansky NJ. 2019. Association mapping desiccation resistance within chromosomal inversions in the African malaria vector *Anopheles gambiae*. *Molecular Ecology* 28:1333–1342.

Baumdicker F, Bisschop G, Goldstein D, Gower G, Ragsdale AP, Tsambos G, Zhu S, Eldon B, Ellerman EC, Galloway JG, et al. 2022. Efficient ancestry and mutation simulation with msprime 1.0. *Genetics* 220:iyab229.

Berdan EL, Barton NH, Butlin R, Charlesworth B, Faria R, Fragata I, Gilbert KJ, Jay P, Kapun M, Lotterhos KE, et al. 2023. How chromosomal inversions reorient the evolutionary process. *Journal of Evolutionary Biology* 36:1761–1782.

Brooks ME, Kristensen K, Van Benthem KJ, Magnusson A, Berg CW, Nielsen A, Skaug HJ, Machler M, Bolker BM. 2017. glmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modeling. *The R journal* 9:378–400.

Burban E, Tenaillon MI, Glémin S. 2024. RIDGE, a tool tailored to detect gene flow barriers across species pairs. *Molecular Ecology Resources* 24:e13944.

Butlin RK, Smadja CM. 2018. Coupling, reinforcement, and speciation. *The American Naturalist* 191:155–172.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009. BLAST+: architecture and applications. *BMC Bioinformatics* 10:421.

Chang CC, Chow CC, Tellier LC, Vattikuti S, Purcell SM, Lee JJ. 2015. Second-generation PLINK: rising to the challenge of larger and richer datasets. *GigaSci* 4:7.

Chen S, Zhou Y, Chen Y, Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* 34:i884–i890.

Cruickshank TE, Hahn MW. 2014. Reanalysis suggests that genomic islands of speciation are due to reduced diversity, not reduced gene flow. *Molecular Ecology* 23:3133–3157.

Dobzhansky T. 1940. Speciation as a stage in evolutionary divergence. *The American Naturalist* 74:312–321.

Dobzhansky T, Levene H. 1948. Genetics of natural populations. Xvii. Proof of operation of natural selection in wild populations of *Drosophila pseudoobscura*. *Genetics* 33:537–547.

Dopman EB, Shaw KL, Servedio MR, Butlin RK, Smadja CM. 2024. Coupling of barriers to gene exchange: causes and consequences. *Cold Spring Harb Perspect Biol* 16:a041432.

Ebdon S, Laetsch DR, Vila R, Baird SJE, Lohse K. 2025. Genomic regions of current low hybridisation mark long-term barriers to gene flow in scarce swallowtail butterflies. *PLOS Genetics* 21:e1011655.

Faria R, Johannesson K, Butlin RK, Westram AM. 2019. Evolving inversions. *Trends in Ecology & Evolution* 34:239–248.

Faria R, Navarro A. 2010. Chromosomal speciation revisited: rearranging theory with pieces of evidence. *Trends in Ecology & Evolution* 25:660–669.

Fick S, Hijmans R. 2017. WorldClim 2: New 1-km spatial resolution climate surfaces for
global land areas. *International Journal of Climatology* 37.
Fraïsse C, Popovic I, Mazoyer C, Spataro B, Delmotte S, Romiguier J, Loire É, Simon A,
Galtier N, Duret L, et al. 2021. DILS: Demographic inferences with linked selection
by using ABC. *Molecular Ecology Resources* 21:2629–2644.
Fuller ZL, Koury SA, Phadnis N, Schaeffer SW. 2019. How chromosomal rearrangements
shape adaptation and speciation: case studies in *Drosophila pseudoobscura* and its
sibling species *Drosophila persimilis*. *Molecular Ecology* 28:1283–1301.
Fuller ZL, Leonard CJ, Young RE, Schaeffer SW, Phadnis N. 2018. Ancestral
polymorphisms explain the role of chromosomal inversions in speciation. *PLoS Genet*
14:e1007526.
Garlovsky MD, Yusuf LH, Ritchie MG, Snook RR. 2020. Within-population sperm
competition intensity does not predict asymmetry in conpopulation sperm precedence.
Philosophical Transactions of the Royal Society B: Biological Sciences
375:20200071.
Garrison E, Marth G. 2012. Haplotype-based variant detection from short-read sequencing.
arXiv preprint arXiv:1207.3907.
Gibbs AG. 2002. Lipid melting and cuticular permeability: new insights into an old problem.
Journal of Insect Physiology 48:391–400.
Hazel JR. 1995. Thermal adaptation in biological membranes: is homeoviscous adaptation
the explanation? *Annu. Rev. Physiol.* 57:19–42.
Hijmans RJ, Van Etten J, Cheng J, Mattiuzzi M, Sumner M, Greenberg JA, Lamigueiro OP,
Bevan A, Racine EB, Shortridge A, et al. 2015. Package 'raster.' *R package* 734:473.
Hoffmann AA, Anderson A, Hallas R. 2002. Opposing clines for high and low temperature
resistance in *Drosophila melanogaster*. *Ecology Letters* 5:614–618.
Hoffmann AA, Rieseberg LH. 2008. Revisiting the impact of inversions in evolution: from
population genetic markers to drivers of adaptive shifts and speciation? *Annual
Review of Ecology, Evolution, and Systematics* 39:21–42.
Hoikkala A, Poikela N. 2022. Adaptation and ecological speciation in seasonally varying
environments at high latitudes: *Drosophila virilis* group. *Fly* 16:85–104.
Huang DW, Sherman BT, Tan Q, Collins JR, Alvord WG, Roayaei J, Stephens R, Baseler
MW, Lane HC, Lempicki RA. 2007. The DAVID Gene Functional Classification
Tool: a novel biological module-centric algorithm to functionally analyze large gene
lists. *Genome Biol* 8:R183.
Huang DW, Sherman BT, Tan Q, Kir J, Liu D, Bryant D, Guo Y, Stephens R, Baseler MW,
Lane HC, et al. 2007. DAVID Bioinformatics Resources: expanded annotation
database and novel algorithms to better extract biology from large gene lists. *Nucleic
Acids Research* 35:W169–W175.
Jackson BC. 2011. Recombination-suppression: how many mechanisms for chromosomal
speciation? *Genetica* 139:393–402.
Jeffares DC, Jolly C, Hoti M, Speed D, Shaw L, Rallis C, Balloux F, Dessimoz C, Bähler J,
Sedlazeck FJ. 2017. Transient structural variations have strong effects on quantitative
traits and reproductive isolation in fission yeast. *Nat Commun* 8:14061.
Jennings JH, Etges WJ, Schmitt T, Hoikkala A. 2014. Cuticular hydrocarbons of *Drosophila
montana*: geographic variation, sexual dimorphism and potential roles as pheromones.
Journal of Insect Physiology 61:16–24.
Jennings JH, Snook RR, Hoikkala A. 2014. Reproductive isolation among allopatric
Drosophila montana populations. *Evolution* 68:3095–3108.
Kapun M, Barrón MG, Staubach F, Obbard DJ, Wiberg RAW, Vieira J, Goubert C, Rota-
Stabelli O, Kankare M, Bogaerts-Márquez M, et al. 2020. Genomic analysis of

European *Drosophila melanogaster* populations reveals longitudinal structure, continent-wide selection, and previously unknown DNA viruses. *Molecular Biology and Evolution* 37:2661–2678.

Kauranen H, Kinnunen J, Hiillos A-L, Lankinen P, Hopkins D, Wiberg RAW, Ritchie MG, Hoikkala A. 2019. Selection for reproduction under short photoperiods changes diapause-associated traits and induces widespread genomic divergence. *Journal of Experimental Biology* 222:jeb205831.

Keightley PD, Ness RW, Halligan DL, Haddrill PR. 2014. Estimation of the spontaneous mutation rate per nucleotide site in a *Drosophila melanogaster* full-sib family. *Genetics* 196:313–320.

Kellermann V, Loeschcke V, Hoffmann AA, Kristensen TN, Fløjgaard C, David JR, Svenning J-C, Overgaard J. 2012. Phylogenetic constraints in key functional traits behind species' climate niches: patterns of desiccation and cold resistance across 95 *Drosophila* species. *Evolution* 66:3377–3389.

Kirkpatrick M, Barton N. 2006. Chromosome inversions, local adaptation and speciation. *Genetics* 173:419–434.

Kulathinal RJ, Stevison LS, Noor MAF. 2009. The genomics of speciation in *Drosophila*: diversity, divergence, and Introgression estimated using low-coverage genome sequencing. *PLoS Genet* 5:e1000550.

Laetsch DR, Bisschop G, Martin SH, Aeschbacher S, Setter D, Lohse K. 2023. Demographically explicit scans for barriers to gene flow using gIMble. *PLoS Genet* 19:e1010999.

Lê S, Josse J, Husson F. 2008. FactoMineR: an R package for multivariate analysis. *Journal of statistical software* 25:1–18.

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics* 25:1754–1760.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000 Genome Project Data Processing Subgroup. 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* 25:2078–2079.

Lohse K, Clarke M, Ritchie MG, Etges WJ. 2015. Genome-wide tests for introgression between cactophilic *Drosophila* implicate a role of inversions during speciation. *Evolution* 69:1178–1190.

Mackintosh A, Vila R, Laetsch DR, Hayward A, Martin SH, Lohse K. 2023. Chromosome fissions and fusions act as barriers to gene flow between *Brenthis fritillary* butterflies. *Molecular Biology and Evolution* 40:msad043.

MacMillan HA, Andersen JL, Davies SA, Overgaard J. 2015. The capacity to maintain ion and water homeostasis underlies interspecific variation in *Drosophila* cold tolerance. *Sci Rep* 5:18607.

MacMillan HA, Sinclair BJ. 2011. Mechanisms underlying insect chill-coma. *Journal of Insect Physiology* 57:12–20.

Marçais G, Delcher AL, Phillippy AM, Coston R, Salzberg SL, Zimin A. 2018. MUMmer4: A fast and versatile genome alignment system. *PLoS Comput Biol* 14:e1005944.

Mazerolle MJ, Mazerolle MMJ. 2020. Package 'AICcmodavg.' *See https://cran.uib.no/web/packages/AICcmodavg/AICcmodavg.pdf*.

Mirol PM, Schäfer MA, Orsini L, Routtu J, Schlötterer C, Hoikkala A, Butlin RK. 2007. Phylogeographic patterns in *Drosophila montana*. *Molecular Ecology* 16:1085–1097.

Navarro A, Betrán E, Barbadilla A, Ruiz A. 1997. Recombination and gene flux caused by gene conversion and crossing over in inversion heterokaryotypes. *Genetics* 146:695–709.

Overgaard J, Hoffmann AA, Kristensen TN. 2011. Assessing population and environmental effects on thermal resistance in *Drosophila melanogaster* using ecologically relevant assays. *Journal of Thermal Biology* 36:409–416.

Parker DJ, Envall T, Ritchie MG, Kankare M. 2021. Sex-specific responses to cold in a very cold-tolerant, northern *Drosophila* species. *Heredity* 126:695–705.

Parker DJ, Wiberg RAW, Trivedi U, Tyukmaeva VI, Gharbi K, Butlin RK, Hoikkala A, Kankare M, Ritchie MG. 2018. Inter and intraspecific genomic divergence in *Drosophila montana* shows evidence for cold adaptation. *Genome Biology and Evolution* 10:2086–2101.

Patterson JT. 1952. Revision of the *montana* complex of the *virilis* species group. *Univerisity Texas Publ* 5204:20–34.

Poikela N, Laetsch DR, Hoikkala V, Lohse K, Kankare M. 2024. Chromosomal inversions and the demography of speciation in *Drosophila montana* and *Drosophila flavomontana*. *Genome Biology and Evolution* 16:evae024.

Poikela N, Tyukmaeva V, Hoikkala A, Kankare M. 2021. Multiple paths to cold tolerance: the role of environmental cues, morphological traits and the circadian clock gene *vrille*. *BMC Ecol Evo* 21:117.

Rausch T, Zichner T, Schlattl A, Stütz AM, Benes V, Korbel JO. 2012. DELLY: structural variant discovery by integrated paired-end and split-read analysis. *Bioinformatics* 28:i333–i339.

Ravinet M, Faria R, Butlin RK, Galindo J, Bierne N, Rafajlović M, Noor M a. F, Mehlig B, Westram AM. 2017. Interpreting the genomic landscape of speciation: a road map for finding barriers to gene flow. *Journal of Evolutionary Biology* 30:1450–1477.

Routtu J, Mazzi D, Van Der Linde K, Mirol P, Butlin RK, Hoikkala A. 2007. The extent of variation in male song, wing and genital characters among allopatric *Drosophila montana* populations. *Journal of Evolutionary Biology* 20:1591–1601.

Rundle HD, Nosil P. 2005. Ecological speciation. *Ecology Letters* 8:336–352.

Schluter D. 2001. Ecology and the origin of species. *Trends in Ecology & Evolution* 16:372–380.

Sedlazeck FJ, Rescheneder P, Smolka M, Fang H, Nattestad M, von Haeseler A, Schatz MC. 2018. Accurate detection of complex structural variations using single-molecule sequencing. *Nat Methods* 15:461–468.

Stevison LS, Hoehn KB, Noor MAF. 2011. Effects of inversions on within- and between-species recombination and divergence. *Genome Biology and Evolution* 3:830–841.

Stinziano JR, Sové RJ, Rundle HD, Sinclair BJ. 2015. Rapid desiccation hardening changes the cuticular hydrocarbon profile of *Drosophila melanogaster*. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology* 180:38–42.

Stone WS, Guest WC, Wilson FD. 1960. The evolutionary implications of the cytological polymorphism and phylogeny of the *virilis* group of *Drosophila*. *Proceedings of the National Academy of Sciences* 46:350–361.

Sturtevant AH. 1921. A case of rearrangement of genes in *Drosophila*. *Proc. Natl. Acad. Sci. U.S.A.* 7:235–237.

Tahami MS, Vargas-Chavez C, Poikela N, Coronado-Zamora M, González J, Kankare M. 2024. Transposable elements in *Drosophila montana* from harsh cold environments. *Mobile DNA* 15:18.

Tarasov A, Vilella AJ, Cuppen E, Nijman IJ, Prins P. 2015. Sambamba: fast processing of NGS alignment formats. *Bioinformatics* 31:2032–2034.

Teets NM, Denlinger DL. 2013. Physiological mechanisms of seasonal and rapid cold-hardening in insects. *Physiological Entomology* 38:105–116.

Thorvaldsdottir H, Robinson JT, Mesirov JP. 2013. Integrative Genomics Viewer (IGV):
high-performance genomics data visualization and exploration. *Briefings in*
Bioinformatics 14:178–192.
Throckmorton LH. 1982. The *virilis* species group. *Genetics and Bioogy of Drosophila*
3:227–296.
Tyukmaeva V, Lankinen P, Kinnunen J, Kauranen H, Hoikkala A. 2020. Latitudinal clines in
the timing and temperature-sensitivity of photoperiodic reproductive diapause in
Drosophila montana. Ecography 43:759–768.
Vesala L, Salminen TS, Kankare M, Hoikkala A. 2012. Photoperiodic regulation of cold
tolerance and expression levels of *regucalcin* gene in *Drosophila montana. Journal of*
Insect Physiology 58:704–709.
de Villemereuil P, Gaggiotti OE. 2015. A new FST-based method to uncover local adaptation
using environmental variables. *Methods in Ecology and Evolution* 6:1248–1258.
Villoutreix R, Ayala D, Joron M, Gompert Z, Feder JL, Nosil P. 2021. Inversion breakpoints
and the evolution of supergenes. *Molecular Ecology* 30:2738–2755.
Wellenreuther M, Bernatchez L. 2018. Eco-evolutionary genomics of chromosomal
inversions. *Trends in Ecology & Evolution* 33:427–440.
Wiberg R a. W, Tyukmaeva V, Hoikkala A, Ritchie MG, Kankare M. 2021. Cold adaptation
drives population genomic divergence in the ecological specialist, *Drosophila*
montana. Molecular Ecology 30:3783–3796.
Wright D, Schaeffer SW. 2022. The relevance of chromatin architecture to genome
rearrangements in *Drosophila. Phil. Trans. R. Soc. B* 377:20210206.
Yusuf LH, Tyukmaeva V, Hoikkala A, Ritchie MG. 2022. Divergence and introgression
among the *virilis* group of *Drosophila. Evolution Letters* 6:537–551.
Zuther E, Schulz E, Childs LH, Hincha DK. 2012. Clinal variation in the non-acclimated and
cold-acclimated freezing tolerance of *Arabidopsis thaliana* accessions. *Plant, Cell &*
Environment 35:1860–1878.
1035