Domain gain or loss in fungal chitinases drives ecological specialization toward antagonism or immune suppression

Ruben Eichfeld[1,2], Asmamaw B. Endeshaw[1], Margareta J. Hellmann[3], Bruno M.
Moerschbacher[3], Alga Zuccaro[1,2*]

[1]University of Cologne, Institute for Plant Sciences, Cologne, Germany

[2]Cluster of Excellence on Plant Sciences (CEPLAS), Cologne, Germany

[3]University of Münster, Institute of Biology and Biotechnology of Plants, Münster,
Germany

*Correspondence: azuccaro@uni-koeln.de

Abstract

The evolutionary origins of fungal effector proteins remain poorly understood, particularly how structural changes reprogram antimicrobial enzymes into host-adapted immune suppressors. Here, we show that domain modularity drives ecological specialization in chitinase effectors of the beneficial root endophyte *Serendipita indica*. The GH18 chitinase *Si*CHIT, which carries a C-terminal carbohydrate-binding module (CBM5), is expressed during fungal competition and antagonizes the fungal pathogen *Bipolaris sorokiniana* in the rhizosphere, thereby protecting plant roots. Deletion of the CBM5 abolishes this antifungal activity, while fusion of CBM5 to the CBM5-lacking paralog *Si*CHIT2 restores pathogen inhibition. In contrast, *Si*CHIT2 is induced during root colonization and suppresses chitin-triggered reactive oxygen species production, promoting immune evasion and host compatibility. These results identify CBM5 as a modular determinant of effector function, switching chitinase activity between microbial antagonism and host immune suppression. Our findings support an evolutionary scenario

26 in which effector function in planta arises through domain loss and transcriptional
27 divergence from an antimicrobial precursor, consistent with transitions along the
28 saprotrophy-to-symbiosis continuum.

Significance Statement

Effector proteins play key roles in shaping fungal interactions with both plant hosts and
microbial competitors, yet how these functions evolve remains unclear. Here, we show
that structural domain modularity enables ecological specialization of two paralogous
chitinases arising from gene duplication in the root endophyte *Serendipita indica*. Through
domain deletion and fusion, we demonstrate that a carbohydrate-binding module (CBM5)
determines whether a chitinase functions in microbial antagonism or immune evasion.
Our findings provide mechanistic evidence for effector diversification via domain loss and
transcriptional divergence, supporting an evolutionary trajectory from antimicrobial
activity to host adaptation. This work advances our understanding of how modular
architecture drives effector evolution and niche specialization in symbiotic fungi.

Highlights

• Gain or loss of a CBM5 binding domain drives effector specialization between
fungal antagonism and immune suppression

• CBM5 acts as a modular determinant enabling antifungal activity in the GH18
chitinase *Si*CHIT

• The CBM5-lacking paralog *Si*CHIT2 suppresses host immunity and promotes root
colonization

• Functional divergence following gene duplication illustrates evolutionary
repurposing of an antimicrobial enzyme into an immune-suppressive effector

Introduction

Mutualistic fungi are ecologically important members of the rhizosphere microbiome, where they support plant health and influence microbial community structure and function (Martin & Tan, 2025) This diverse group includes ectomycorrhizal and arbuscular mycorrhizal fungi, as well as beneficial root endophytes. Unlike mycorrhizal fungi, root endophytes often colonize plant roots without forming specialized symbiotic structures and, depending on the host and environmental context, can enhance plant performance by promoting nutrient uptake, water acquisition, and tolerance to environmental stress (Vandenkoornhuyse *et al.*, 2002; Dastogeer *et al.*, 2018; Mesny *et al.*, 2021; Mahdi *et al.*, 2022; Andreo-Jimenez *et al.*, 2023; Zhang *et al.*, 2024). These fungi exemplify the evolutionary plasticity of fungal lifestyles, flexibly occupying ecological niches along the saprotrophy-to-symbiosis continuum. To colonize plant roots and shape their ecological niche, beneficial root endophytes must overcome two key challenges: evasion of host immunity and competition with microbial antagonists and plant pathogens, a process recently shown to be supported by synergistic interactions with bacterial members of the core microbiota (Mahdi *et al.*, 2022). Both immune evasion and microbial competition are mediated by secreted effector proteins. Effector-mediated immune evasion often involves the degradation or masking of microbe-associated molecular patterns (MAMPs), such as β-1,6-glucan and chitin, to prevent recognition by host pattern recognition receptors (Lahrmann & Zuccaro, 2012; Fesel & Zuccaro, 2016; Wawra *et al.*, 2016). In parallel, effectors can alter the microbiota composition by suppressing microbial competitors, a strategy well documented in plant pathogens (Snelders *et al.*, 2020; Snelders *et al.*, 2021) and increasingly recognized in mutualistic fungi.

The beneficial root endophyte *Serendipita indica* (*Si*), a member of the widespread fungal order Sebacinales (Basidiomycota), colonizes a broad range of host plants and is known for its ability to promote growth and suppress plant immunity while protecting the host from pathogens. Colonization is confined to the rhizoplane, epidermis, and cortex, and depends on an intact host immune system (Lahrmann *et al.*, 2015). In addition to its host-associated functions, *Si* exhibits strong antifungal activity in the rhizosphere, particularly against the pathogen *Bipolaris sorokiniana (Bs)*, the causal agent of spot blotch and root

rot in cereals (Sarkar *et al.*, 2019; Mahdi *et al.*, 2022; Eichfeld *et al.*, 2024). This
antagonism is mediated by a GH18-CBM5 endochitinase (*Si*CHIT), whose catalytic
activity restricts *Bs* growth and recapitulates *the* protective effect of *Si* in planta (Eichfeld
et al., 2024).

GH18 chitinases, found across all fungal phyla, cleave β-1,4-glycosidic bonds in chitin, a
major structural component of fungal cell walls. These enzymes fulfill diverse roles in
nutrient scavenging, hyphal remodeling, and fungal antagonism (Seidl, 2008; Chen *et al.*,
2020). Their catalytic activity depends on a conserved DxxDxDxE motif, which defines
the substrate-assisted catalytic mechanism characteristic of this family (van Aalten *et al.*,
2001; Bußwinkel *et al.*, 2018). While some chitinase-derived effectors have lost enzymatic
activity and evolved to sequester MAMPs (Fiorin *et al.*, 2018), others retain catalytic
function and are functionally enhanced by accessory domains such as carbohydrate-
binding modules (CBMs) (Limón *et al.*, 2001; Limón *et al.*, 2004; Monge *et al.*, 2018).
These non-catalytic domains increase substrate affinity and specificity, particularly for
crystalline or polymeric forms of chitin (Itoh *et al.*, 2006; Matroodi *et al.*, 2013). *Si*CHIT
features a C-terminal CBM5 domain, a modular architecture found predominantly in
bacteria and Basidiomycota but absent in Ascomycota, suggesting a lineage-specific
evolutionary adaptation (Chen *et al.*, 2020; Eichfeld *et al.*, 2024). Here, we investigate the
functional significance of CBM5 in *Si*CHIT and examine how domain architecture
modulates effector activity, enabling a switch between microbial antagonism and immune
evasion.

We show that the CBM5 domain is essential for the antifungal and plant-protective
function of *Si*CHIT. This chitinase is expressed during fungal competition and antagonizes
the pathogen *Bs* in the rhizosphere, thereby protecting plant roots. In contrast, its closely
related paralog *Si*CHIT2 lacks CBM5 and is induced during root colonization. *Si*CHIT2
promotes evasion of chitin-triggered immunity, preventing the generation of host-derived
reactive oxygen species. Through domain-swapping experiments, we demonstrate that
CBM5 functions as a modular determinant that switches the ecological role of a chitinase
from microbial antagonism to host immune suppression. Together, our findings support a
model in which effector specialization emerges through gene duplication, domain loss,

and transcriptional divergence, reprogramming an antimicrobial enzyme into a host-
adapted immune effector.

Results

CBM5 enhances binding to crystalline chitin and fungal cell wall without altering product specificity
To investigate the role of the C-terminal CBM5 domain in *Si*CHIT (PIIN_03543), we
generated three recombinant variants: full-length *Si*CHIT, a catalytically inactive mutant
(*Si*CHIT^{E196Q}), and a truncated version lacking the CBM5 domain (*Si*CHIT^{-CBM5}), which
retained the proline-rich linker. The truncated construct was produced by excluding the
final 141 bp from the coding sequence. All recombinant proteins were purified via affinity
chromatography and verified by SDS-PAGE (Fig. S1).

We first assessed chitinase activity on insoluble crab shell chitin using thin-layer
chromatography (TLC) and by quantifying reducing sugars after 20 hours of incubation
(Fig. 1A, B). These assays, along with endpoint measurements on chitin azure (CA) (Fig.
1C), revealed no major differences between full-length *Si*CHIT and *Si*CHIT^{-CBM5}.
However, time-course experiments on CA showed that the presence of CBM5 enhanced
early hydrolysis (Fig. 1D), suggesting that the domain increases the reaction rate during
initial substrate engagement.

To evaluate whether CBM5 influences product outcome, we analyzed the hydrolysates of
colloidal chitin by mass spectrometry. Both *Si*CHIT and *Si*CHIT^{-CBM5} produced similar
oligomer profiles dominated by chitobiose (A2), indicating that CBM5 does not alter
product specificity (Fig. 1E). Notably, substantial amounts of partially deacetylated
products, such as A1D1 and A2D2, were also detected. This likely reflects the fact that
colloidal chitin is not fully acetylated; deacetylated regions may form stretches with higher
solubility and increased accessibility, leading to their preferential cleavage by the
enzymes. To assess whether CBM5 affects subsite preferences, we analyzed the
hydrolysis products of chitosans with defined degrees of acetylation (DA). *Si*CHIT^{-CBM5}
displayed the same requirement for an acetylated unit at the -1 subsite and no detectable
shifts at adjacent positions compared to *Si*CHIT, confirming that CBM5 does not influence
catalytic specificity (Fig. 1F; Eichfeld *et al*., 2024).

To determine the role of CBM5 in substrate interaction, we performed pull-down assays
using crab shell chitin, highly deacetylated chitosan, and lyophilized, protein-free *Bs*
mycelium. *Si*CHIT^{-CBM5} showed reduced binding to crab shell chitin compared to full-
length *Si*CHIT, while no difference was observed with chitosan, suggesting that CBM5
enhances binding specifically to acetylated, crystalline substrates (Fig. 1G). Binding to *Bs*
mycelium was also reduced in the absence of CBM5, indicating that the domain
contributes to cell wall targeting.

Together, these results show that while CBM5 is dispensable for full catalytic activity and
does not alter product specificity, it enhances early hydrolysis rates and substrate binding,
particularly to crystalline or fungal cell wall-like chitin. This suggests that CBM5 improves
functional efficiency under biologically relevant conditions, likely contributing to the
ecological role of *Si*CHIT in microbial competition.

CBM5 is required for antifungal activity and plant protection
We previously demonstrated that *Si*CHIT restricts the growth of *Bs* and reduces its
colonization of plant roots through its chitinolytic activity (Eichfeld *et al*., 2024). However,
the specific contribution of the CBM5 domain to this antifungal function remained
unresolved. To test whether CBM5 is required for *Bs* inhibition, we incubated *Bs* spores
with the full-length *Si*CHIT, the catalytically inactive mutant *Si*CHIT^{E196Q}, the truncated
variant lacking CBM5 *Si*CHIT^{-CBM5}, or the empty vector control Ev. As expected, *Bs* spore
germination was significantly reduced in the presence of active *Si*CHIT, but not with
*Si*CHIT^{E196Q}. *Si*CHIT^{-CBM5} showed a similar lack of inhibitory effect as *Si*CHIT^{E196Q} and
had only a minimal effect compared to the Ev control (Fig. 2A).

We next examined whether the loss of antifungal activity translates into reduced
protection of host roots. *Bs* spores treated with the same chitinase variants were used to
inoculate the roots of the *Bs* host *Hordeum vulgare* (*Hv*). As previously shown, *Si*CHIT-
treated spores led to a significant reduction in *Bs* colonization, as measured by fungal
transcript levels in roots. In contrast, *Si*CHIT^{E196Q} and *Si*CHIT^{-CBM5} had no effect (Fig. 2B).
Correspondingly, only *Si*CHIT mitigated the reduction in root biomass caused by *Bs*
infection (Fig. 2C). To test whether this effect is conserved in other hosts, we repeated

the experiment using *Arabidopsis thaliana* (*At*). *Si*CHIT reduced *Bs* biomass and
preserved root elongation, while *Si*CHIT^{E196Q} did not. *Si*CHIT^{-CBM5} had a weak protective
effect, albeit significantly lower than the full-length enzyme. (Fig. S2.).

Together, these results demonstrate that both chitinase activity and the CBM5 domain are
required for *Si*CHIT's antifungal and plant-protective functions. The loss of protection
observed in the CBM5-lacking variant suggests that this domain contributes to cell wall
binding and is critical for effective niche defense.

A CBM5-lacking paralog, *Si*CHIT2, is transcriptionally specialized for host colonization
GH18-CBM5 chitinases are absent in Ascomycota but occur frequently in the
Agaricomycetes (Basidiomycota), where gene copy number varies across species with
different ecological lifestyles (Eichfeld et al., 2024). To better define the structural
requirements for antifungal activity in *Si*, we compared the four GH18 chitinases encoded
in the genome with respect to phylogenetic relationships, domain architecture, and
transcriptional regulation. Of the four, only *Si*CHIT harbors a CBM5 domain, whereas the
other three chitinases contain no additional domains besides the GH18 catalytic domain
(Fig. 3A). *Si*CHIT and its closest paralog *Si*CHIT2 (PIIN_03542) share the highest
sequence identity and form a distinct phylogenetic pair among the *Si* GH18 chitinases
(Fig. 3A; Supplementary Fig. S3). The catalytic DxxDxDxE motif located at the -1 subsite
is conserved across all four chitinases, suggesting that each is catalytically active (Fig.
3B). Despite this shared motif, the chitinases exhibit distinct structural features. *Si*CHIT2
retains an 86-amino-acid C-terminal linker but lacks a CBM5 domain. *Si*CHIT3
(PIIN_11727) lacks a predicted signal peptide, while *Si*CHIT4 (PIIN_07603), features
extended surface-exposed loops within the GH18 domain (Supplementary Table S1).
Taken together, *Si*CHIT and *Si*CHIT2 are more similar to each other in sequence,
structure, and genomic context than to the other chitinases in *Si*. Notably, they are located
adjacent in the genome (Fig. 3C). Their tandem organization, high sequence identity, and
absence of CBM5 in *Si*CHIT2 support a gene duplication event followed by structural
divergence through domain loss.

To assess structural similarity, we generated AlphaFold models of both enzymes after
removing the predicted signal peptides. The models confirmed the overall structural
conservation of the GH18 catalytic domain and the presence of a C-terminal linker in
*Si*CHIT2, but no CBM5 domain (Fig. 3D & E)

To examine transcriptional regulation, we analyzed RNA-seq data from Eichfeld et al.
2024 and found that *Si*CHIT and *Si*CHIT2 were the only chitinases significantly
upregulated in biotic contexts. *Si*CHIT was specifically induced during fungal
confrontation, while *Si*CHIT2 was induced during colonization of barley, *Brachypodium*
and *Arabidopsis*. In contrast, *Si*CHIT3 and *Si*CHIT4 were not transcriptionally activated
under any tested biotic conditions, including interactions with pathogenic fungi, beneficial
bacteria, or different plant hosts (Fig. 3F & G). To extend the RNA-seq findings, we
measured the expression of *Si*CHIT and *Si*CHIT2 by qPCR following confrontation with
Bs and six additional filamentous fungi representing the major fungal lineages
Ascomycota, Basidiomycota, and Mucoromycota. *Si*CHIT was induced in several of these
interactions, whereas *Si*CHIT2 expression remained low under all tested microbial
conditions (Fig. 3H). Additionally, both *Si*CHIT and *Si*CHIT2 were induced when *S. indica*
was grown on crab shell chitin as the sole carbon source (Fig. 3I).

Together, these findings suggest that *Si*CHIT2 evolved a distinct ecological function from
*Si*CHIT following gene duplication. Their divergent domain architecture and expression
profiles indicate functional specialization, with *Si*CHIT adapted for fungal antagonism and
*Si*CHIT2 for host colonization.

Fusion of CBM5 reprograms *Si*CHIT2 for fungal antagonism
To test whether the CBM5 domain of *Si*CHIT is sufficient to confer antifungal activity, we
constructed a chimeric protein consisting of the GH18 catalytic domain of *Si*CHIT2 fused
to the proline-rich linker and CBM5 domain of *Si*CHIT (*Si*CHIT2^{+CBM5}). As a control, we
generated a second construct in which only the linker was fused to the GH18 domain of
*Si*CHIT2 (*Si*CHIT2^{+link}) (Fig. 4A). As observed for native *Si*CHIT and *Si*CHIT^{-CBM5}, the
presence of CBM5 did not affect endpoint chitinase activity on crab shell chitin or chitin
azure (Fig. 4B, C). However, only *Si*CHIT2^{+CBM5} suppressed *Bs* spore germination,

comparable to the effect of full-length SiCHIT. In contrast, SiCHIT2^{+link} showed no

inhibitory activity (Fig. 4D). Consistent with these results, both SiCHIT and SiCHIT2^{+CBM5}

reduced root growth inhibition in barley caused by Bs, and limited pathogen colonization.

In contrast, SiCHIT^{-CBM5} and SiCHIT2^{+link} had no measurable protective effect (Fig. 4E &

F).

These results demonstrate that fusing CBM5 to SiCHIT2 is sufficient to confer antifungal

activity, indicating that CBM5 acts as a modular domain enabling functional adaptation

toward microbial antagonism.

SiCHIT2 suppresses chitin-triggered immunity and facilitates root colonization by S. indica

Given the distinct plant-induced expression profile of SiCHIT2, we hypothesized that this

chitinase contributes to immune evasion during root colonization. Previous studies have

shown that *Podosphaera xanthii* uses chitinase-like effectors to degrade immunogenic

chitin oligomers and suppress host defense responses (Martínez-Cruz *et al.*, 2021). To

test whether Si chitinases function similarly, we incubated immunogenic chitohexaose

(A6) with recombinant SiCHIT, SiCHIT2, their respective variants, or the Ev control at a

1:1 molar ratio. The resulting hydrolysates were applied to barley roots to measure

reactive oxygen species (ROS) production. ROS accumulation was completely abolished

when chitohexaose was pre-treated with either SiCHIT2^{+link} or SiCHIT^{-CBM5}. In contrast,

treatment with SiCHIT2^{+CBM5} or full-length SiCHIT resulted in only partial or no

suppression of ROS accumulation (Fig. 5A, B). An intermediate reduction in ROS

accumulation was observed when chitohexaose was pre-treated with SiCHIT^{E196Q}, likely

due to the inactive catalytic cleft sequestering a fraction of the chitohexaose. None of the

recombinant chitinases induced ROS when applied in the absence of chitohexaose (Fig.

S4).

To assess whether this immune suppression contributes to host colonization, we co-

inoculated Si with 20 µM of each recombinant chitinase and quantified fungal biomass in

barley roots at three days post-inoculation. SiCHIT2^{+link} significantly enhanced root

colonization, whereas the other chitinases did not (Fig. 5C). SiCHIT^{-CBM5} showed a slight

but non-significant trend toward increased colonization.

These results suggest that *Si*CHIT2's ability to suppress ROS depends on the absence

of CBM5, which may otherwise restrict access to soluble substrates and favor binding to

insoluble chitin. This immune-suppressive function, together with its host-specific

transcriptional induction, underscores a distinct ecological role for SiCHIT2 compared to

its CBM5-containing paralog.

Discussion

Discussion

Effector proteins are central to the ecological success of root-colonizing fungi, mediating

interactions not only with host plants but also with competing microbes. While traditionally

associated with immune suppression (Stergiopoulos & de Wit, 2009), fungal effectors are

now also recognized for their roles in inter-microbial competition. They facilitate the

acquisition and maintenance of ecological niches in the rhizosphere and on root surfaces,

thereby supporting fungal persistence and competitive advantage (Snelders *et al.*, 2020;

Gómez-Pérez *et al.*, 2023; Eichfeld *et al.*, 2024). These dual roles are not restricted to

pathogens; beneficial fungi such as *Si* employ similar strategies to persist in complex

microbial communities, maintain host compatibility, and defend their ecological niche.

This functional versatility reflects broader evolutionary trends across the fungal kingdom,

where transitions from saprotrophy to symbiosis are accompanied by the repurposing of

ancestral traits to meet ecological demands.

Here, we show that *Si* encodes two closely related GH18 chitinases, *Si*CHIT and

*Si*CHIT2, that have diverged structurally and transcriptionally to perform distinct

ecological functions. *Si*CHIT, which carries a C-terminal CBM5 domain, is expressed

during fungal confrontation and contributes to pathogen growth inhibition and plant

protection. In contrast, *Si*CHIT2, is induced during root colonization, and suppresses

chitin-triggered immune responses in the host (Fig. 5). This divergence possibly reflects

an evolutionary trajectory driven by gene duplication and domain loss, enabling *Si* to

balance microbial antagonism and host compatibility through modular effector

specialization.

The importance of domain modularity in effector function is supported by previous studies

in *Trichoderma* spp., where the addition of substrate-binding domains such as CBM1 or

CBM18 increases their binding affinity for insoluble substrates and enhances antifungal activity by improving proximity and substrate targeting (Limón *et al.*, 2001; Limón *et al.*, 2004). However, such effects can be masked in vivo by genome-wide redundancy in cell wall-degrading enzymes and antimicrobial effectors. Using purified proteins, we demonstrate that the CBM5 domain is critical for the antifungal activity of *Si*CHIT. Its deletion abolishes fungal antagonism and its reintroduction into *Si*CHIT2 restores this function. This is consistent with studies in bacterial chitinases, where CBM5 loss reduces antifungal activity, likely due to impaired substrate binding (Tsujibo *et al.*, 2003; Itoh *et al.*, 2006). In our assays, CBM5 enhanced early hydrolysis rates and binding to crystalline chitin and fungal cell wall, but did not affect overall catalytic activity or product profile. This suggests a proximity-based enhancement mechanism, in which CBM5 anchors the enzyme at the fungal cell wall. Unlike antimicrobial or agglutinating lectins such as WSC3, a β-1,3-glucan-binding lectin from *S. indica* that agglutinates fungal cells independently of enzymatic activity (Wawra *et al.*, 2019), CBM5 alone does not provide antifungal activity which requires a catalytically active GH18 core. Some fungal pathogens employ catalytically inactive chitinase effectors (e.g., *Mp*Chi in *Moniliophthora perniciosa*; Fiorin *et al.*, 2018) or active chitinases (e.g., *Mo*Chia1 in *Magnaporthe oryzae*; Yang & Yu, 2019; Martínez-Cruz *et al.*, 2021), or EWCAs; Martínez-Cruz *et al.*, 2021) to suppress host immunity by sequestering or degrading free chitin oligomers. Similarly, we show that both *Si*CHIT2 and *Si*CHIT[-CBM5] can suppress chitohexaose-induced ROS production. These findings suggest that CBM5 may hinder activity on soluble substrates not by altering catalytic specificity, but by limiting access to short oligomers through spatial constraints. CBM5 domains are structurally optimized for engaging extended, crystalline chitin surfaces via aromatic stacking and hydrogen bonding (Kezuka *et al.*, 2006; Vaaje-Kolstad *et al.*, 2019). Accordingly, CBM5-containing chitinases like *Si*CHIT likely localize to insoluble polysaccharides in fungal cell walls, anchoring the enzyme to non-immunogenic material. Although such substrate targeting could in principle restrict access to soluble MAMPs, this is unlikely in the ROS assay, where only chitohexaose is present. A more plausible explanation is that CBM5 increases the rigidity or steric bulk of the protein, thereby reducing flexibility and limiting productive interaction with short ligands. In contrast, CBM5-lacking variants such as *Si*CHIT2 remain free to access and degrade free

immunogenic chitin oligomers. This suggests that suppression of host immunity may

require a CBM5-free configuration optimized for apoplastic MAMP degradation during

root colonization. Such structural neofunctionalization provides a plausible trajectory for

the evolution of fungal effector functions. Given the saprotrophic ancestry and facultative

symbiotic lifestyle of *S. indica*, *Si*CHIT likely evolved under selective pressure from fungal

competitors, whereas *Si*CHIT2 emerged as an immune-suppressive effector during the

transition to a root-associated niche. This represents a clear case of functional divergence

driven by domain loss and regulatory rewiring. By comparison, the *Verticillium dahliae*

effector *Vd*AMP3 illustrates a different evolutionary strategy: co-opted from an ancestral

antimicrobial protein, it retains both its antimicrobial activity and core structural features

while functioning in microbiota manipulation in planta. Rather than undergoing major

neofunctionalization, *Vd*AMP3 shows how effectors can be repurposed for new ecological

roles with minimal structural change (Snelders *et al.*, 2021). This highlights the diversity

of evolutionary pathways by which fungal effectors adapt to changing environments.

Together, our findings show that domain gain and loss can reprogram effector proteins for

distinct ecological roles. CBM5 serves as a modular determinant, switching GH18

chitinase activity between microbial antagonism and immune suppression. This highlights

how structural modularity enables fungi to adapt effectors to the dual pressures of

rhizosphere competition and host interaction in symbiotic fungi.

Material and Methods

Plant and fungal materials

Hordeum vulgare (*Hv*, L. cv Golden Promise) and *Arabidopsis thaliana* (*At*, Col-0) were

used as plant hosts. *Hv* seeds were surface-sterilized in 6 % sodium hypochlorite for 1 h

under constant shaking and subsequently washed five times for 30 min in sterile water.

Seeds were germinated for four days on wet filter paper at 21 °C in the dark. Seeds of *At*

were surface-sterilized in 70 % ethanol for 10 min and 100 % ethanol for 7 min. *At* seeds

were placed on ½ MS medium (Murashige-Skoog medium including vitamins, pH 5.6)

containing 1 % sucrose and stratified for three days in the dark prior to germination on a

day-night cycle of 8/16 h at 22/18 °C, 60 % humidity and 125 µmol/m^2 light intensity for

seven days. *Serendipita indica* (*Si*, DSM11827) and *Bipolaris sorokiniana* (*Bs*, ND90Pr)

were used as fungal species. *Si* was grown on CM medium (Complete Medium, 1.5 %

agar) at 28 °C for 30 days in the dark. *Si* spores were harvested by adding sterile water

and scraping the spores with a scalpel. The suspension was filtered through a sterile

miracloth filter and spores were centrifuged at 2,000 x g. The supernatant was discarded

and spores were washed three times in water. *Bs* was grown on modified CM medium

(Sarkar *et al.*, 2019) at 28 °C for 14 days in the dark. *Bs* spores were harvested like *Si*

spores but a Drigalski spatula was used to scrape the spores instead of a scalpel.

Preparation of fungal inoculants and plant colonization

For the preparation of *Bs* inoculants, spores were harvested from 21-day old plates and

incubated with the respective recombinant proteins (10 µM in 50 mM phosphate buffer)

for 16 h at 28 °C in the dark. Spore mixtures were diluted to 5000 spores/mL to yield the

final working inoculants. For inoculation of *At*, ten sterile seven-day old seedlings were

placed on solid 1/10 PNM (pH 5.6) plates (0.005 % KNO_3, 0.005 % KH_2PO_4, 0.0025 %

K_2HPO_4, 0.049 % $MgSO_4$, 0.00472 % $Ca(NO_3)_2$, 0.0025 % NaCl, 0.5 % (v/v) Fe-EDTA

stock solution, 1.2 % Gelrite, pH 5.6. After autoclaving, 10 mL 1 M MES pH 5.6 was

added. The EDTA stock solution contains 2.78 % $FeSO_4$ x 7 H_2O, 4.13 % Na_2EDTA x

$2H_2O$. Concentrations are indicated in w/v if not stated otherwise) and the roots were

treated with 1 mL of the respective inoculant. The root elongation was quantified by

measuring the difference root length at the start of the inoculation and the end of the

experiment at three days post inoculation (dpi). Roots were washed with water and snap

frozen in liquid nitrogen for further RNA extraction. For inoculation of *Hv*, four sterile

seedlings were placed in glass jars containing solid 1/10 PNM medium (pH 5.6) and

inoculated with three mL of the respective inoculant. The root weight was quantified at

three dpi. Roots were washed in water and snap frozen for further RNA extraction. For

the preparation of *Si* inoculants, spores were harvested from 28-day old plates and were

allowed to germinate overnight in sterile water at 28 °C. Prior to plant root inoculation, the

respective recombinant proteins were added to 20 µM in 50 mM phosphate buffer.

Subsequently, the roots of four day-old *Hv* seedlings were inoculated with three mL of

500,000 spores/mL. *Hv* seedlings were harvested at three dpi, washed in water and snap
frozen in liquid nitrogen.

Preparation of fungal confrontations

For the preparation of fungal inoculants, liquid CM medium was inoculated with 500,000
spores/mL. For all other fungi, mycelium was scraped from plates and transferred to liquid
CM. All fungi were grown for six days at 28 °C and 120 rpm. The mycelium was blended
and re-suspended in fresh CM and regenerated for another 24 h. Regenerated mycelium
was filtered through miracloth filters and washed thoroughly with sterile milliQ water.
Fungal mycelia were mixed in a 1:1 ratio and 1 g of mycelium was streaked on PNM
plates containing 15 % agar. After six days, the mycelia were harvested, dried with tissue
papers and snap frozen in liquid nitrogen.

Fungal spore germination

To quantify the *Bs* spore germination, isolated spores were resolved in liquid CM to a
concentration of 100,000 spores/mL and 100 µL were added to the wells of sterile Lab-
Tek 2 chamber slides with cover (VWR). 100 µL of the respective treatment (Ev control or
purified protein) were added to a final concentration of 10 µM. The germination rate of
spores was quantified 6 h after treatment.

RNA extraction from plant roots and RT-qPCR

Plant materials were ground in liquid nitrogen and 1 mL of TRIzol reagent (Thermo
Fischer scientific) was added. Samples were vortexed for 20 sec until all material was
resolved. 200 µL chloroform were added and samples vortexed again. Samples were
centrifuged at 17,000 x g and 4 °C for 30 min and 500 µL of the upper aqueous phase
were transferred to 500 µL isopropanol. The nucleic acids were precipitated at -20 °C for
16 h and pelleted by centrifugation at 17,000 x g and 4 °C for 30 min. Pellets were
washed three times with 70 % ethanol diluted in DEPC water. Remaining DNA was
removed by adding 1 µL DNase1 (1 U/µL) in 3 µL DNase1 buffer (10 x) (Thermo Fischer
scientific) and 26 µL RNase-free water. Samples were incubated at 37 °C for 30 min.
DNase digest was stopped by heating to 70 °C for 5 min and subsequent precipitation
with 2 volumes isopropanol and ½ volume of 7.5 M ammonium acetate for 16 h at -20 °C.
Samples were washed twice with 70 % ethanol in DEPC water and resolved in 30 µL

RNase-free water. Concentrations were determined using a NanoDrop 2000c
spectrophotometer (Thermo Fischer Scientific).

1 µg of RNA was used for reverse transcription into cDNA (Thermo Fischer Scientific).
First, 1 µL of random hexamer and of oligo-dt oligomers were added and incubated at
65 °C for 5 min and subsequently placed on ice. 4 µL 5 x reaction buffer, 1 µL Riboblock
RNase inhibitor (20 U/µL), 2 µL 10 mM dNTPs and 2 µL MMLV reverse transcriptase were
added and samples were incubated at 42 °C for 1 h. cDNA synthesis was stopped by
heating to 70 °C for 5 min. RT-qPCR was conducted using the primers listed in
Supplementary Table S2.

Cloning of chitinase expression vectors
The coding sequence (CDS) of the chitinase genes were amplified by PCR with Q5 High-
Fidelity Polymerase (New England Biolabs) using the primers listed in Supplementary
Table S2. PCR products were cloned using the Gibson assembly cloning technique into
the PQE80L expression vector (Qiagen, Hilden, Germany) after vector linearization with
Kpn1 (New England Biolabs).

Production and purification of recombinant protein in *E. coli*
Chemically competent *E. coli* BL21 cells were transformed with the vectors for protein
expression under control of a lac operon. 5 mL starter cultures were prepared in liquid LB
medium containing 100 µg/mL carbenicillin and grown for 16 h at 37 °C and 180 rpm.
Starter cultures were transferred to 300 mL of LB medium with carbenicillin and grown to
an OD_{600} of 0.5 at 28 °C. Gene expression was induced by adding IPTG to a final
concentration of 1 mM and cells were grown at 16 °C for 20 h. Cells were pelleted by
centrifugation at 5000 g for 15 min at 4 °C. Bacterial pellets were resolved in 5 mL lysis
buffer (50 mM NaH_2PO_4, 300 mM NaCl and 10 mM imidazole, pH 8) and sonicated three
times for 1 min with pulses on ice and centrifuged at 16,200 g for 30 min. 5 mL of the
supernatant were added to 1 mL of Nickel-NTA Agarose (Qiagen, Hilden, Germany) and
incubated under continuous shaking at 4 °C for 1 h. Samples were spun down at 500 g
for 10 secs, the flow through was removed, 4 mL wash buffer (50 mM NaH_2PO_4, 300 mM
NaCl and 20 mM imidazole, pH 8) were added and samples were incubated under

continuous shaking at 4 °C for 10 min. Washing steps were repeated five times. For

elution, 1 mL of elution buffer (50 mM NaH_2PO_4, 300 mM NaCl and 250 mM imidazole,

pH 8) was added and samples were incubated under continuous shaking at 4 °C for 10

min. Eluted fractions were dialyzed against 50 mM phosphate buffer (pH 6) over night

and protein concentrations were measured using Bradford reagent. To test the purity of

the protein samples, aliquots were boiled for 5 min in SDS-sample buffer, loaded on a 10

% SDS gel and visualized by Coomassie staining.

Chitinase activity assays

To quantify the enzymatic activity of the purified chitinases, 2 mg/mL chitin azure (Sigma

Aldrich) were incubated with 5 µM recombinant chitinase in a total volume of 200 µL.

Reactions were incubated at 28 °C for 20 h. Reactions were stopped by heating to 95 °C

for 5 min and samples were centrifuged 5 min at 17,000 x g and the absorbance of

supernatants was quantified at 560 nm. For quantifying reducing ends, 10 g crab shell

chitin were incubated with 5 µM enzyme in 200 µL 50 mM phosphate buffer for 20 h at

28 °C. Reactions were stopped by heating to 95 °C for 5 min. Samples were centrifuced

for 5 min at 17,000 x g and supernatant were mixed in a 1:1 ratio with DNSA (3,5-

dinitrosalicylic acid) reagent. DNSA reagent is prepared by mixing 1 % (w/v) 3,5-

dinitroslicylic acid, 1.6 % (w/v) sodium hydroxide, 30 % (w/v) potassium sodium tartrate

tetrahydrate. The samples were heated for 10 min at 95 °C and subsequently transferred

to ice. The absorbance of the samples was measured at 540 nm.

For TLC, 10 mg crab shell chitin (Sigma Aldrich) were weighed in 2 mL reaction tubes

and recombinant chitinases were added to a final concentration of 5 µM in 200 µL.

Reactions were incubated at 28 °C for 20 h and stopped by heating to 95 °C for 5 min.

Samples were centrifuged at 17,000 x g for 5 min. 10 µL of the supernatant were

separated on 60 F_{254} TLC (Sigma Aldrich) silica gel plates with running buffer (n-

butanol/methanol/28 % ammonium hydroxide/water, 5/4/2/1). Plates were dried and then

bathed in developer solution (1.6 g diphenylamine/1.6 mL aniline/12 mL 85 % phosphoric

acid/80 mL acetone). TLC plates were developed in an incubation oven at 100 °C for 20

to 30 min.

Chitinase subsite specificity

The subsite specificity of SiCHIT^{-CBM5} was determined as previously reported (Cord-Landwehr $et\ al.$, 2017; Eichfeld $et\ al.$, 2024).

Substrate binding assay

15 mg of crab shell chitin (Sigma Aldrich), Chitosan (Sigma Aldrich) or freeze-dried mycelium of Bs were weighed in a 2 mL reaction tube and incubated with 400 µL chitin-binding buffer (2 mM KH_2PO_4, 8 mM Na_2HPO_4, 2 mM KCL) and 100 µL of 20 µM recombinant protein shaking at 4 °C for 1 h. Samples were centrifuged at 17,000 x g for 5 min and 60 µL of the supernatant were mixed with 60 µL Bradford reagent. Protein amounts were quantified at 595 nm and compared to the input amount.

Quantification of reactive oxygen species (ROS)

For elicitor preparation, 10 µM of chitohexaose were incubated with 10 µM of recombinant chitinase or the buffer control for 16 h at 28 °C. Reactions were stopped by heating to 95 °C for 5 min and chitohexaose was diluted to 250 nM.

Four root pieces (0.5 cm) of four-day old barley seedlings were transferred to each well of a 96-well microtiter plate (white, flat bottom) containing 200 µL of 2.5 mM 2-(N-morpholino)ethanesulfonic acid (MES) buffer, pH 5.6. The plate was incubated for 16 h at 21°C for recovery. The buffer was replaced with 100 µL 2.5 mM MES buffer containing 20 µM LO-12 and 20 µg/mL Horseradish peroxidase (HRP). After 25-min incubation in the dark, 100 µL of 250 nM chitohexaose elicitor solution was added to each well. Chemiluminescence was measured using a TECAN SPARK 10 M microplate reader over all wells for 2 h with an integration time of 450 ms.

Data availability

All data supporting the findings of this study are available within the article and supporting information.

Acknowledgements

We acknowledge support from the Cluster of Excellence on Plant Sciences (CEPLAS) funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy-EXC 2048/1-Project ID: 390686111 and the ZU 263/11-2 (SPP 2125 DECRyPT).

Competing interest

The authors declare no competing interest.

Author contributions

AZ and RE conceptualized the study. RE and AE conducted most of the research. MJH, BMM and RE conducted mass spectrometric analysis of the product profile and the subsite specificity. AZ and RE wrote the manuscript with input from all authors.

References

Andreo-Jimenez B, Te Beest DE, Kruijer W, Vannier N, Kadam NN, Melandri G, Jagadish SVK, van der Linden G, Ruyter-Spira C, Vandenkoornhuyse P, et al. 2023. Genetic Mapping of the Root Mycobiota in Rice and its Role in Drought Tolerance. *Rice (N Y)* **16**(1): 26.

Bußwinkel F, Goñi O, Cord-Landwehr S, O'Connell S, Moerschbacher BM. 2018. Endochitinase 1 (Tv-ECH1) from Trichoderma virens has high subsite specificities for acetylated units when acting on chitosans. *International Journal of Biological Macromolecules* **114**: 453-461.

Chen W, Jiang X, Yang Q. 2020. Glycoside hydrolase family 18 chitinases: The known and the unknown. *Biotechnology Advances* **43**: 107553.

Cord-Landwehr S, Ihmor P, Niehues A, Luftmann H, Moerschbacher BM. 2017. Quantitative Mass-Spectrometric Sequencing of Chitosan Oligomers Revealing Cleavage Sites of Chitosan Hydrolases. **89**(5): 2893-2900.

Dastogeer KMG, Li H, Sivasithamparam K, Jones MGK, Wylie SJ. 2018. Fungal endophytes and a virus confer drought tolerance to Nicotiana benthamiana plants through modulating osmolytes, antioxidant enzymes and expression of host drought responsive genes. *Environmental and Experimental Botany* **149**: 95-108.

Eichfeld R, Mahdi LK, De Quattro C, Armbruster L, Endeshaw AB, Miyauchi S, Hellmann MJ, Cord-Landwehr S, Peterson D, Singan V, et al. 2024. Transcriptomics reveal a mechanism of niche defense: two beneficial root endophytes deploy an antimicrobial GH18-CBM5 chitinase to protect their hosts. *New Phytologist* **244**(3): 980-996.

Fesel PH, Zuccaro A. 2016. β-glucan: Crucial component of the fungal cell wall and elusive MAMP in plants. *Fungal Genetics and Biology* **90**: 53-60.

Fiorin GL, Sanchéz-Vallet A, Thomazella DPT, do Prado PFV, do Nascimento LC,
Figueira AVO, Thomma B, Pereira GAG, Teixeira P. 2018. Suppression of Plant
Immunity by Fungal Chitinase-like Effectors. *Curr Biol* **28**(18): 3023-3030.e3025.
Gómez-Pérez D, Schmid M, Chaudhry V, Hu Y, Velic A, Maček B, Ruhe J, Kemen A,
Kemen E. 2023. Proteins released into the plant apoplast by the obligate parasitic
protist Albugo selectively repress phyllosphere-associated bacteria. *New*
Phytologist **239**(6): 2320-2334.
Itoh Y, Watanabe J, Fukada H, Mizuno R, Kezuka Y, Nonaka T, Watanabe T. 2006.
Importance of Trp59 and Trp60 in chitin-binding, hydrolytic, and antifungal
activities of Streptomyces griseus chitinase C. *Appl Microbiol Biotechnol* **72**(6):
1176-1184.
Kezuka Y, Ohishi M, Itoh Y, Watanabe J, Mitsutomi M, Watanabe T, Nonaka T. 2006.
Structural Studies of a Two-domain Chitinase from Streptomyces griseus
HUT6037. *Journal of Molecular Biology* **358**(2): 472-484.
Lahrmann U, Strehmel N, Langen G, Frerigmann H, Leson L, Ding Y, Scheel D,
Herklotz S, Hilbert M, Zuccaro A. 2015. Mutualistic root endophytism is not
associated with the reduction of saprotrophic traits and requires a
noncompromised plant innate immunity. *New Phytologist* **207**(3): 841-857.
Lahrmann U, Zuccaro A. 2012. Opprimo ergo sum--evasion and suppression in the root
endophytic fungus Piriformospora indica. *Mol Plant Microbe Interact* **25**(6): 727-
737.
Limón MC, Chacón MR, Mejías R, Delgado-Jarana J, Rincón AM, Codón AC, Benítez
T. 2004. Increased antifungal and chitinase specific activities of Trichoderma
harzianum CECT 2413 by addition of a cellulose binding domain. *Applied*
Microbiology and Biotechnology **64**(5): 675-685.
Limón MC, Margolles-Clark E, Benítez T, Penttilä M. 2001. Addition of substrate-
binding domains increases substrate-binding capacity and specific activity of a
chitinase from Trichoderma harzianum. *FEMS Microbiology Letters* **198**(1): 57-63.
Mahdi LK, Miyauchi S, Uhlmann C, Garrido-Oter R, Langen G, Wawra S, Niu Y, Guan
R, Robertson-Albertyn S, Bulgarelli D, et al. 2022. The fungal root endophyte
Serendipita vermifera displays inter-kingdom synergistic beneficial effects with the
microbiota in Arabidopsis thaliana and barley. *The ISME Journal* **16**(3): 876-889.
Martin F, Tan H. 2025. Saprotrophy-to-symbiosis continuum in fungi. *Current Biology*
35(11): R469-R475.
Martínez-Cruz J, Romero D, Hierrezuelo J, Thon M, de Vicente A, Pérez-García A.
2021. Effectors with chitinase activity (EWCAs), a family of conserved, secreted
fungal chitinases that suppress chitin-triggered immunity. *The Plant Cell* **33**(4):
1319-1340.
Matroodi S, Motallebi M, Zamani M, Moradyar M. 2013. Designing a new chitinase with
more chitin binding and antifungal activity. *World J Microbiol Biotechnol* **29**(8):
1517-1523.
Mesny F, Miyauchi S, Thiergart T, Pickel B, Atanasova L, Karlsson M, Hüttel B, Barry
KW, Haridas S, Chen C, et al. 2021. Genetic determinants of endophytism in the
Arabidopsis root mycobiome. *Nature Communications* **12**(1): 7227.
Monge EC, Tuveng TR, Vaaje-Kolstad G, Eijsink VGH, Gardner JG. 2018. Systems
analysis of the glycoside hydrolase family 18 enzymes from Cellvibrio japonicus

characterizes essential chitin degradation functions. *J Biol Chem* **293**(10): 3849-
3859.
Sarkar D, Rovenich H, Jeena G, Nizam S, Tissier A, Balcke GU, Mahdi LK,
Bonkowski M, Langen G, Zuccaro A. 2019. The inconspicuous gatekeeper:
endophytic Serendipita vermifera acts as extended plant protection barrier in the
rhizosphere. *New Phytologist* **224**(2): 886-901.
Seidl V. 2008. Chitinases of filamentous fungi: a large group of diverse proteins with
multiple physiological functions. *Fungal Biology Reviews* **22**(1): 36-42.
Snelders NC, Petti GC, van den Berg GCM, Seidl MF, Thomma BPHJ. 2021. An
ancient antimicrobial protein co-opted by a fungal plant pathogen for in planta
mycobiome manipulation. *Proceedings of the National Academy of Sciences*
118(49): e2110968118.
Snelders NC, Rovenich H, Petti GC, Rocafort M, van den Berg GCM, Vorholt JA,
Mesters JR, Seidl MF, Nijland R, Thomma BPHJ. 2020. Microbiome
manipulation by a soil-borne fungal plant pathogen using effector proteins. *Nature*
Plants **6**(11): 1365-1374.
Stergiopoulos I, de Wit PJGM. 2009. Fungal Effector Proteins. *Annual Review of*
Phytopathology **47**(Volume 47, 2009): 233-263.
Tsujibo H, Kubota T, Yamamoto M, Miyamoto K, Inamori Y. 2003. Characterization of
Chitinase Genes from an Alkaliphilic Actinomycete, <i>Nocardiopsis prasina</i>
OPC-131. *Applied and Environmental Microbiology* **69**(2): 894-900.
Vaaje-Kolstad G, Tuveng TR, Mekasha S, Eijsink VGH 2019. Enzymes for Modification
of Chitin and Chitosan. *Chitin and Chitosan*, 189-228.
van Aalten DM, Komander D, Synstad B, Gåseidnes S, Peter MG, Eijsink VG. 2001.
Structural insights into the catalytic mechanism of a family 18 exo-chitinase. *Proc*
Natl Acad Sci U S A **98**(16): 8979-8984.
Vandenkoornhuyse P, Baldauf SL, Leyval C, Straczek J, Young JPW. 2002. Extensive
Fungal Diversity in Plant Roots. *Science* **295**(5562): 2051-2051.
Wawra S, Fesel P, Widmer H, Neumann U, Lahrmann U, Becker S, Hehemann J-H,
Langen G, Zuccaro A. 2019. FGB1 and WSC3 are in planta-induced β-glucan-
binding fungal lectins with different functions. *New Phytologist* **222**(3): 1493-1506.
Wawra S, Fesel P, Widmer H, Timm M, Seibel J, Leson L, Kesseler L, Nostadt R,
Hilbert M, Langen G, et al. 2016. The fungal-specific β-glucan-binding lectin
FGB1 alters cell-wall composition and suppresses glucan-triggered immunity in
plants. *Nature Communications* **7**(1): 13188.
Yang C, Yu Y. 2019. Binding of the Magnaporthe oryzae Chitinase MoChia1 by a Rice
Tetratricopeptide Repeat Protein Allows Free Chitin to Trigger Immune Responses.
31(1): 172-188.
Zhang Y, Yang Z, Yang Y, Han A, Rehneke L, Ding L, Wei Y, Liu Z, Meng Y, Schäfer
P, et al. 2024. A symbiont fungal effector relocalizes a plastidic oxidoreductase to
nuclei to induce resistance to pathogens and salt stress. *Current Biology* **34**(13):
2957-2971.e2958.

Figure 1: CBM5 enhances binding to fungal cell wall and crystalline chitin and promotes hydrolysis without affecting product specificity. A) Thin layer chromatography (TLC) of crab shell chitin incubated with the Ev control or 10 µM of *Si*CHIT, *Si*CHIT[E196Q], or *Si*CHIT[-CBM5] at 28 °C for 20 h. **B)** Chitinase activity of recombinant chitinases on crab shell chitin at 10 µM and after 20 h at 28 °C. Samples were spun down and supernatants mixed 1:1 with DNSA reagent. Samples were incubated at 100 °C for 10 min and placed on ice. Subsequently, the relative amount of reducing ends was measured at 540 nm (mean +/- SD, n = 3). **C)** Chitinase activity of recombinant chitinases at 10 µM after 20 h at 28 °C on chitin azure (CA). Samples were spun down and supernatants were quantified at 560 nm (mean +/- SD, n = 3). **D)** Chitinase activity of recombinant chitinases at 10 µM measured over 4 h at 28 °C and normalized to the value of t_0 (mean +/- SEM, n = 3). **E)** Product profile of *Si*CHIT and *Si*CHIT[-CBM] on colloidal chitin. Samples were incubated with

5 µM of chitinases for 24 h at 28 °C and hydrolysis products were identified and quantified via MS. A = acetylated, D = de-acetylated unit. **F)** Subsite preference of *Si*CHIT^-CBM5 measured by MS. Chitosan of three degrees of acetylation (DA) were hydrolyzed for 1 h or 24 h. Based on the sequenced products, the frequency of acetylated units at the -2 to +2 subsites of *Si*CHIT^-CBM5 was determined. The black arrow indicates the gylcosidic bond between the -1 and +1 subsite that is cleaved by the enzyme. **G)** Substrate-binding ability of *Si*CHIT and *Si*CHIT^-CBM5 on crab shell chitin, > 75 % de-acetylated chitosan or lyophilized *Bs* mycelium. Binding was assessed by measuring the protein amount in the supernatant of reactions compared to input amount (n = 6-8). Limits of the boxplots represent the 25th–75th percentile, the horizontal line represents the median and the whiskers the minimum/maximum values without outliers. Statistical analysis: **B** and **D)** Different letters indicate significant differences according to one-way ANOVA followed by Tukey's HSD test (adjusted p-value < 0.05). **G)** Student's t-test (P-value: * < 0.05; ** < 0.01).

Figure 2: The CBM5 domain is essential for the antifungal and plant-protective activity of SiCHIT.
A) Relative *Bs* spore germination six h post incubation in sterile chamber slides with the Ev control or 10 µM of *Si*CHIT, *Si*CHIT^E196Q or *Si*CHIT^-CBM5 (n = 6). **B)** Colonization of *Hv* roots by *Bs* at three dpi inferred from relative expression of the fungal housekeeping gene *BsTEF* compared to the *Hv* housekeeping gene *HvUBI* inferred by qPCR using the $2^{-\Delta CT}$ method. Colonization values were normalized to the Ev control. *Bs* spores were either pre-treated with the Ev control or 10 µM of recombinant chitinases for 20 h at 28 °C (mean +/- SD, n = 4). **C)** *Hv* root fresh weight at three dpi with *Bs* spores. *Bs* spores were either pre-treated with the Ev control or 10 µM of recombinant chitinases for 20 h at 28 °C (n = 5-7). Limits of the boxplots represent the 25th–75th percentile, the horizontal line represents the median and the whiskers the minimum/maximum values without outliers. Statistical analysis: Different letters indicate significant differences according to one-way ANOVA followed by Tukey's HSD test (adjusted p-value < 0.05).

A Phylogeny of GH18 chitinases in *Si*

Phylogeny · Domain structure

85 · *Si*CHIT
68 · *Si*CHIT2
*Si*CHIT3
*Si*CHIT4
*Pf*CHIT

0.20

B Conservation of the DxxDxDxE motif of GH18 chitinases in *Si*

```
                                    *      140                 160
SiCHIT  : EDNGFDGIDIDYEYPSSTEQGN--------GFAALIAELRT:156
SiCHIT2 : EDNGFDGIDIDYEYPSSVEQGV--------GFASLLAELRI:156
SiCHIT3 : EDYGFDGIDIDYEYPNNHEQAR--------GYVDLLRELRH:149
SiCHIT4 : EKFELDGIDIDWEYPNGDGAGNEKSSKDTANFLKFLQLLRK:141
```

C Relative genomic location of *SiCHIT* and *SiCHIT2* on contig 0064

1965 bp · 1390 bp · 2117 bp

SiCHIT2 · *SiCHIT*

D Predicted structure of *SiCHIT*

CBM5: 486-530

GH18: 61-405

E Predicted structure of *SiCHIT2*

GH18: 65-409

F Expression profile of *SiCHIT* and *SiCHIT2*

SynCom · *Bs* · *At* · *Bd* · *Hv*

SiCHIT
SiCHIT2

1 3 6 10 · 1 3 6 10 · 1 3 6 10 · 1 3 6 10 · 3 10

dpi

L2FC
-2 -1 0 1 2 3

H Expression of *SiCHIT* and *SiCHIT2* in fungal confrontation

SiCHIT · *SiCHIT2*

Rel. expr. (*Chitinase/SiTEF*)

Si, *Bs*, *Bcs*, *Pa*, *Po*, *St*, *Bj*, *Mo* + *Si*

G Expression of *SiCHIT* and *SiCHIT2* during *Hv* root colonization

SiCHIT · *SiCHIT2*

Rel. expr. (*Chitinase/SiTEF*)

Si, *SiHv* 3 dpi, *SiHv* 10 dpi, *SiHv* 21 dpi

I Expression of *SiCHIT* and *SiCHIT2* during growth on chitin

SiCHIT · *SiCHIT2*

Rel. expr. (*Chitinase/SiTEF*)

0 · 0.5 · 1 · 2 · 4

Crab shell chitin (mg/mL)

Figure 3: *SiCHIT2* encodes a GH18 chitinase without CBM5 and is expressed during host plant colonization. A) Maximum likelihood tree of all four GH18 chitinases encoded by *Si*. Signal peptides were predicted by SignalP 5.0 and removed prior to amino acid sequence alignment. Amino acid sequences were aligned using the MUSCLE algorithm. A maximum likelihood tree was constructed with 500 bootstraps. A GH18 chitinase from *Plasmodium falciparum* was used as outgroup. Symbols on the right indicate the domain architecture of the full length proteins: small blue box: signal peptide; grey line: inter-domain regions; larger boxes (red, green, blue, beige): GH18 domain; small green box in *Si*CHIT: CBM5. **B)** Amino acid sequence alignment of the catalytic DxxDxDxE motif. Blue: conserved aspartate (D); red: conserved glutamate (E). The asterisk indicates the glutamate that was mutated to glutamin (Q) in the catalytically inactive *Si*CHIT^{E196Q}. **C)** Scheme of the relative location of *SiCHIT* and *SiCHIT2* in the genome of *Si* on contig 0064. The grey bars illustrate introns. *SiCHIT* has 11, and *SiCHIT2* 10 introns. **D, E)** Structure prediction of *Si*CHIT and *Si*CHIT2 by AlphaFold. The sequences of predicted signal peptides were identified with SignalP 5.0 and removed prior to structure prediction. **F)** Expression profile of *SiCHIT* and *SiCHIT2* during confrontation with a bacterial community (SynCom), the pathogen *B. sorokiniana* (*Bs*), or during colonization of the host plants *A. thaliana* (*At*), *B. distachyon* (*Bd*) or *H. vulgare* (*Hv*). The data was taken from Eichfeld *et al.* 2024. **G)** Expression of *SiCHIT* or *SiCHIT2* during *Hv* root colonization at 3, 10 and 21 dpi. Four day old *Hv* roots were inoculated with *Si* and grown on 1/10 PNM medium. Expression was inferred by relative expression of the fungal housekeeping gene *SiTEF* compared to *SiCHIT* or *SiCHIT2* by qPCR using the $2^{-\Delta CT}$ method (mean +/- SD, n = 3-5). **H)** Expression of *SiCHIT* or *SiCHIT2* at six days of growth on 1/10 PNM medium with different fungal species (*Bs* = *Bipolaris sorokiniana*; *Bcs* = *Brunneochlamydosporium sp.*; *Pa* = *Papulaspora equi*; *Po* = *Podospora sp.*; *St* = *Stachybotrys sp.*; *Bj*: *Bjerkandera sp.*; *Mo* = *Mortierella sp.*. Fungal liquid cultures were grown in CM and washed with water, mixed in a 1:1 ratio and 1 g mycelium was streaked out on plates. Expression was inferred by relative expression of the fungal housekeeping gene *SiTEF* compared to *SiCHIT* or *SiCHIT2* by qPCR using the $2^{-\Delta CT}$ method (mean +/- SD, n = 4). **I)** Expression of *SiCHIT* or *SiCHIT2* at six days of growth on crab shell chitin-containing PNM medium at different concentrations. *Si* liquid culture grown in CM was washed with water and 1 g mycelium was streaked out on 1/10 PNM plates. Expression was inferred by relative expression of the fungal housekeeping gene *SiTEF* compared to *SiCHIT* or *SiCHIT2* by qPCR using the $2^{-\Delta CT}$ method (n = 4). Statistical analysis: Statistical analysis: Different letters indicate significant differences according to one-way ANOVA followed by Tukey's HSD test (adjusted p-value < 0.05).

Figure 4: Fusion of the CBM5 to *Si*CHIT2 activates its antifungal and plant-protective capacity. A) Domain architecture of recombinant proteins. *Si*CHIT2+link contains the GH18 domain of *Si*CHIT2 and the linker region of *Si*CHIT. *Si*CHIT2+CBM5 contains the GH18 domain of *Si*CHIT2 and the linker region with the CBM5 of *Si*CHIT. **B)** Chitinase activity of recombinant chitinases on crab shell chitin at 10 μM and after 20 h at 28 °C. Samples were spun down and supernatants mixed 1:1 with DNSA reagent. Samples were incubated at 100 °C for 10 min and placed on ice. Subsequently, the relative amount of reducing ends was measured at 540 nm (mean +/- SD, n = 3). **C)** Chitinase activity of recombinant chitinases at 10 μM after 20 h at 28 °C on chitin azure (CA). Samples were spun down and supernatants were quantified at 560 nm (mean +/- SD, n = 3). **D)** Relative *Bs* spore germination six h post incubation in sterile chamber slides with Ev control or 10 μM of *Si*CHIT, *Si*CHIT-CBM5, *Si*CHIT2+link or *Si*CHIT+CBM5. (n = 6). **E)** Colonization of *Hv* roots by *Bs* at three dpi inferred from relative expression of the fungal housekeeping gene *BsTEF* compared to the *Hv* housekeeping gene *HvUBI* by qPCR using the $2^{-\Delta CT}$ method. Colonization values were normalized to the Ev control. *Bs* spores were either pre-treated with the Ev control or 10 μM of recombinant chitinases for 20 h at 28 °C (n = 6). **F)** *Hv* root fresh weight at three dpi with *Bs* spores. *Bs* spores were either pre-

treated with the Ev control or 10 μM of recombinant chitinases for 20 h at 28 °C. Limits of the boxplots represent the 25th–75th percentile, the horizontal line represents the median and the whiskers the minimum/maximum values without outliers (n = 6). Statistical analysis: Different letters indicate significant differences according to one-way ANOVA followed by Tukey's HSD test (adjusted p-value < 0.05).

Figure 5:

***Si*CHIT2 suppresses chitin-triggered immunity and facilitates root colonization by *Si* in *Hv*. A)** ROS burst of 4 d *Hv* roots after treatment with 125 nM chitohexaose (A6). Chitohexaose was incubated with the indicated recombinant chitinases in an equimolar concentration (10 μM) 20 h prior to treatment of *Hv* roots. Values represent means +/- SD from five to six individual wells containing four root pieces of ~ 0.5 cm. **B)** Maximum values of ROS accumulation of *Hv* roots (data from **A**). **C)** Colonization of *Hv* roots by *Si* at three dpi inferred from relative expression of the fungal housekeeping gene *SiTEF* compared to the *Hv* housekeeping gene *HvUBI* by qPCR using the $2^{-\Delta CT}$ method. Colonization values were normalized to the Ev control. *Si* spores were germinated for 16 h at 28 °C and either incubated with the Ev control or 20 μM of recombinant chitinases immediately before *Hv* root inoculation (n = 4). **D)** Transcript accumulation of the *Hv* defense marker gene *Hv*PR10 at three dpi inferred by relative expression of *HvPR10* relative to the *Hv*

housekeeping gene *HvUBI* by qPCR using the $2^{-\Delta CT}$ method (n = 4). Limits of the boxplots represent the 25th–75th percentile, the horizontal line represents the median and the whiskers the minimum/maximum values without outliers. Statistical analysis: Different letters indicate significant differences according to one-way ANOVA followed by Tukey's HSD (adjusted p-value < 0.05).

Figure 6: Model displaying the divergent functions of *Si*CHIT and *Si*CHIT2. During confrontation with the pathogen *Bs*, *Si* secretes a GH18-CBM5 chitinase that efficiently inhibits the growth of *Bs*, enabling *Si* to secure its ecological niche. To effectively function in this context, *Si*CHIT relies on the presence of the CBM5. The paralogous *Si*CHIT2 has a catalytically active GH18 domain but no CBM5. It serves *Si* during host colonization by preventing the elicitation of chitin-triggered immunity at the fungus-plant interface, putatively by degrading immunogenic chitin oligomers that are released from the cell wall of *Si* by host chitinases. Figure was created using Biorender.