Circadian disruption beneath the surface: a screening approach to identify actigraphy-based markers of split shift work in submariners

Daniel Marques[a,b], Carina Fernandes[c,d], Nuno L. Barbosa-Morais[a,b,*], Cátia Reis[a,b,e,*]

[a]*Gulbenkian Institute for Molecular Medicine, Lisbon, Portugal;* [b]*Faculdade de Medicina, Universidade de Lisboa, Lisbon, Portugal;* [c]*Centro de Medicina Naval, Portuguese Navy, Lisbon, Portugal;* [d]*NOVA National School of Public Health, Universidade NOVA de Lisboa, Lisbon, Portugal;* [e]*Universidade Católica Portuguesa, Católica Research Centre for Psychological, Family and Social Wellbeing, Lisbon, Portugal*

*Corresponding authors: Daniel Marques – daniel.marques@gimm.pt, Nuno Morais – nuno.morais@gimm.pt, Cátia Reis – catia.reis@medicina.ulisboa.pt

Address: *Gulbenkian Institute for Molecular Medicine, Faculdade de Medicina,* Universidade de Lisboa, Av. Prof. Egas Moniz, 1649-028 Lisboa, Portugal

Circadian disruption beneath the surface: a screening approach to identify actigraphy-based markers of split shift work in submariners

Abstract

Submarines pose significant physiological and behavioral challenges due to their dimly lit, isolated, confined, and extreme (DICE) environment, which deprives individuals from exposure to a natural light-dark cycle and limits opportunities for physical activity. In this study, we developed a minimal yet robust exploratory feature-screening approach to identify candidate actigraphy markers of split shift work during a three-week military submarine mission. Actigraphy data was collected across the pre-mission, mission and post-mission periods. During the mission, crew members followed a 6-hours-on/6-hours-off split shift schedule, except for submariners with on-call duty or extended hours. Forty-one actigraphy features were screened for their ability to differentiate between study periods and shifts. A Monte Carlo permutation test was used for this initial screening, followed by receiver operating characteristic (ROC) curve analysis for pairwise comparisons. Light-related features, the slopes of regression lines fitted to the daily predicted dim light melatonin onset (DLMO) trajectories, and Pearson correlation coefficients between each recording day and its corresponding predicted DLMO demonstrated the greatest potential to distinguish between study periods. Among shifts, most high-performing features discriminated between split shift work schedules and on-call or extended-hours schedules. Only the relative amplitude (*RA*) and the average light intensity during the five least-illuminated hours of the day (*L5*) reliably distinguished between the shift performing the bulk of nighttime work and its counterpart. Given its robustness under conditions of low activity and light exposure, our methodology may be applicable to other operational or clinical populations with reduced activity levels, such as individuals with depression or dementia.

Keywords

Biological rhythms, circadian disruption, submarine mission, actigraphy.

Introduction

Circadian rhythms are near 24-hour endogenous oscillations that regulate behavior, physiology, and metabolism, allowing the organism to optimally manage bodily needs

in response to predictable changes in the environment (Boivin & Boudreau, 2014; Ca-

jochen & Schmidt, 2025). These are maintained by genetically encoded timekeeping sys-

tems, known as circadian clocks or oscillators, which can be entrained by environment

cues called *zeitgebers* (Aschoff, 1960). In humans and other diurnal primates, light is the

main *zeitgeber*, as it entrains the phase of the suprachiasmatic nucleus (SCN) with the

natural light-dark cycle (Cajochen & Schmidt, 2025). Additionally, peripheral clocks are

sensitive to non-photic *zeitgebers*, such as physical activity and meals (Peters et al., 2024).

Mistimed and irregular exposure to photic and non-photic *zeitgebers*, as is often the case

in modern societies, leads to circadian disruption (Meléndez-Fernández et al., 2023).

Submarine and submarine-like conditions seem to be remarkably disruptive in this

regard. Military submariners are required to operate in a dimly lit, isolated, confined and

extreme (DICE) environment, for periods that could last several months (Nieuwenhuys

et al., 2021; Zivi et al., 2020). Indeed, to maintain continuous operations, crews often

perform skilled and demanding tasks during their biological night, when the endogenous

circadian system promotes sleep (Marando et al., 2023; McHill et al., 2014). In addition,

light intensity levels are likely insufficient to promote effective entrainment, and oppor-

tunities for consolidated sleep are slim, with rest periods often diverted to meals, personal

care and recreational activities (Nieuwenhuys et al., 2021). Likewise, the prolonged social

isolation and reduced opportunities for physical activity may further contribute to worse

physical and mental health outcomes and well-being (Nieuwenhuys et al., 2021).

Shift work in DICE environments accentuates many characteristics of night shift

work, and it may also exacerbate its long-term health effects (Marando et al., 2023). Night

shift work has been associated with multiple health conditions, including cancer and car-

diometabolic disorders (Ansu Baidoo & Knutson, 2023; Erren et al., 2019). Proposed

mechanisms to explain this association point to circadian desynchrony, including circa-

dian misalignment, chronic partial sleep deprivation and melatonin suppression by noc-

turnal light exposure (Smith & Eastman, 2012). Night shifts uncouple *zeitgeber* exposure

from internal time, inducing a state of internal desynchrony among circadian clocks in

different tissues and organs (Heyde & Oster, 2022). Compared to night shift work, the

health effects of submarine shift work, particularly in the post-deployment period, remain

poorly characterized, limiting our ability to assess whether any alterations persist beyond

the mission (Marando et al., 2023).

Moreover, drawing generalizable conclusions from extreme operational environ-

ments, such as submarines, is constrained by small sample sizes and heterogenous

watchkeeping schedules, which vary not only in the distribution of work and rest periods but also in the length of the operational day (Marando et al., 2023). For instance, between 1960 and 2016, U.S. Navy submariners were subjected to an 18-hour period, which is outside the range of entrainment of the human circadian clock and results in rest-activity cycles shifting by 6 hours within each 24-hour period (Chabal et al., 2024). Additionally, under a 6-hours-on/6-hours-off schedule, a biphasic sleep pattern was observed, consisting of a primary sleep episode lasting 4–5 hours and a secondary sleep episode lasting 1.5 hours per each 24-hour cycle (Van Puyvelde et al., 2022).

Furthermore, submarines present a difficult environment for actigraphy data analysis due to the restricted range of movement and persistently low light levels. Given all these constraints, we wondered whether it would be possible to extract a putative signature of actigraphy features that both reflected the conditions of living inside a submarine and the physiological impact of split shift work in such an extreme environment. To this end, we aimed to develop a minimal yet robust exploratory feature-screening approach to identify candidate actigraphy markers of split shift work during a military submarine mission aboard a diesel-electric vessel. These candidate markers may serve as potential signatures of circadian desynchrony and misalignment in submarine and submarine-like conditions and, if validated, could help inform the development of effective mitigation strategies.

Materials and methods

Participants and actigraphy data collection

Actigraphy data from a group of twenty-nine Portuguese military submariners (mean age 35.6 ± 4.9 years) was collected to validate the feature-screening methodology proposed in this study. Participants were mostly male (26/29) and quite experienced both in active duty (15.5 ± 5.1 years) and in the submarine fleet (6.0 ± 5.6 years). Data collection was conducted as part of a broader investigation approved by the Ethics Committee of the NOVA National Public Health School (CE-ENSP nº12/2022). All participants provided informed consent prior to enrollment. Psychological variables and metadata from the same cohort have been previously published (Fernandes et al., 2025). Inclusion was limited to individuals actively serving as crew members during the mission, and those with a previous diagnosis of a sleep disorder or taking chronic medication affecting sleep or wakefulness were excluded.

Figure 1 Study periods and 6-hours-on/6-hours-off shift schedules during a typical submarine mission day. Shift schedules were implemented only during the mission period. Shift 0 is shown in grey, representing personnel assigned to on-call or extended hours.

Data acquisition window spanned three consecutive periods: pre-mission (P1),

mission (P2), and post-mission (P3) **[Figure 1]**. Pre-mission refers to the period of two

weeks before the submarine mission and serves as a baseline condition, as no operational

deployments occurred in the preceding months. During this period, participants kept their

usual routines, including standard working hours at the naval base. Mission refers to a

three-week submarine mission, in which submariners adhered to a 6-hours on/6-hours off

split work schedule. To ensure continuous operations during the mission, submariners

were assigned to three different shifts. Shift 1 (n = 13) operated from 07:00 to 13:00 and

from 19:00 to 01:00, while shift 2 (n = 12) worked from 01:00 to 07:00, and from 13:00

to 19:00. Shift 0 (n = 5) included personnel with on-call duty or extended working hours.

Finally, post-mission corresponds to a three-week recovery period, in which submariners

returned to a standard working schedule and/or off-duty days.

Feature-screening methodology

Acquisition and preprocessing. Throughout the eight-week acquisition window,

submariners wore an ActTrust2 actigraphy monitoring device (Condor Instruments™,

Brazil) on their non-dominant wrist. Activity (PIM), temperature, and light were recorded

at a sampling rate of sixty seconds. Invalid sequences were manually removed at the start

and end of the actigraphy recordings using the ActStudio® software (version 2.2.2). Off-

wrist periods of the actigraphy trace were identified through visual inspection and marked

as missing values.

Panel construction. A panel of forty-one actigraphy features was calculated with

circStudio (v1.0-*alpha*.2)*,* a package for actigraphy data analysis built on top of *pyActig-*

raphy (Hammad et al., 2021, 2024). This custom version was modified for compatibility

with Python 3.12+ and extended with a module for analyzing light recordings using math-

ematical models of circadian rhythms. Specifically, twelve features were derived from

two families of light-informed mathematical models of circadian rhythms (Forger et al.,
1999; Hannay et al., 2019): Kronauer-based (Forger, Jewett) and Hannay (Single Popula-
tion [HannaySP], Two Populations [HannayTP]). Prior to numerical integration, missing
values from off-wrist periods of the actigraphy trace were imputed by averaging the light
intensity values recorded at the same time on different days, after which the signal was
resampled at a 10-minute rate. For a given light intensity trajectory in *lux*, $L(t)$, these
models – systems of first-order ordinary differential equations – were integrated using the
SciPy's *odeint* function, which calls the LSODA integrator from the FORTRAN library
ODEPACK. Dim light melatonin onset (DLMO), a marker of circadian phase, was esti-
mated by applying a seven-hour offset to the core body temperature minimum estimates
from the four models. *scipy.stats.linregress* was used to calculate the Pearson's correlation
coefficient (Pearson's *r*) and the slope of the regression line for the daily predicted DLMO
in P1, P2 and P3. All actigraphy features are detailed in **Table S1**.

Initial feature screening. Feature relationships were first explored by computing a
Sperman correlation matrix and visualizing it using *Seaborn's clustermap* (average link-
age, Euclidean distance). The main axes of variance were identified via principal compo-
nent analysis (PCA) on z-scored data (*sklearn.preprocessing.scale*) using *sklearn.decom-
position.PCA*. To evaluate if any actigraphy features from the panel could discriminate
between at least one study period or shift, a null distribution of F values was generated
using a Monte Carlo permutation test (10^4 permutations). From this distribution, a false
positive rate (FPR) was estimated and controlled for the false discovery rate (FDR) asso-
ciated with multiple hypothesis testing via the Benjamini-Hochberg procedure. Study pe-
riod labels (P1, P2, P3) were randomly permuted within each subject using the *pandas
sample* method, preserving the repeated-measures structure of this factor. Shift labels (S0,
S1, S2) were also randomly permuted within each study period block. To select the most
informative features, a dual filtering criterion was applied based on statistical significance
(FDR < 0.05) and effect size (Cohen's f > 0.63). This effect size threshold corresponds to
the minimum detectable effect size for a repeated measures design with two factors, given
our total sample size (n = 29) and approximately 83% power, as estimated using *G*Power*
(version 3.1.97).

Pairwise feature analysis. To assess the ability of the previously selected features
to discriminate across pairwise comparisons of period and shift levels, receiver operating
characteristic (ROC) curves were drawn using the *roc_curve* function, and the area under
the curve (AUC) was computed with the *auc* function, both from the *scikit-learn* package.

An AUC null distribution was constructed using a Monte Carlo permutation test, as in the

initial feature screening. Plots were generated using the *seaborn* and *matplotlib* libraries.

All analyses were performed in Python 3.12 using the PyCharm 2024.3.3 (Professional

Edition) integrated development environment (IDE). Complete code and documentation

can be found in the **Supplementary Notebook** and in *submarine_notebook.ipynb,* located

in the *notebooks/submarine_project* directory of the *circStudio* GitHub repository.

Results

Identifying candidate actigraphy submarine mission markers

To assess the consequences of the submarine mission on activity-inactivity cycles,

41 features were extracted from actigraphy data. Potential relationships among those

Figure 2 Hierarchical cluster map of Spearman's rank correlation coefficients among the 41 actigraphy features. The dendrogram reveals three major correlation clusters. Clustering was performed using average linkage and Euclidean distance on the correlation matrix.

Figure 3 Principal component analysis of actigraphy features. (A) PCA plot displaying the first two principal components (PC1 and PC2), with samples colored by study period. PC1 effectively separates P2 (the mission period) from P1 and P3 (pre- and post-mission periods, respectively), indicating that study period is a major source of variance. (B) The same PCA projection with samples colored by shift, showing no clear separation, suggesting limited shift-related structure in the two dominant components.

features were explored by computing a Spearman correlation matrix and visualizing the

results as a hierarchically clustered heatmap **[Figure 2]**. This analysis revealed three main

clusters. The first consisted of positively correlated features associated to mathematical

models of circadian rhythms, namely the slope and the Pearson's correlation coefficient

(Pearson's *r*) of the linear models fitted to the daily DLMO prediction according to the

Forger, Jewett, HannaySP and HannayTP formalisms. This cluster also included the du-

ration of minor sleep bouts (*minor_sleep_bouts*), sleep midpoint (*sleep_midpoint*), mean

activity and hourly light exposure levels during the five least active or least-illuminated

hours of the day, respectively (*L5*, *L5_light*), and intradaily variability (*IV*).

A second cluster of positively correlated features, characterized by negative cor-

relations with the first cluster, aggregated several light metrics (e.g., light exposure level

[*exp_level*], light interdaily stability [*IS_light*], time above light intensity thresholds

[*tat_10-500lux*]), some cosinor parameters (*mesor* and *amplitude*), the mean activity and

light intensity levels during the ten most active or brightest hours of the day, respectively

(*M10, M10_light*), relative amplitude (*RA*), and the median duration of all sleep bouts

(*all_sleep_bouts*) and of the main sleep bout (*main_sleep_bout*). A third cluster incorpo-

rated features showing weak to no correlation with each other nor with variables in the

other two clusters (e.g., *acrophase, L5_light_onset*).

To identify the dominant sources of variance in the dataset, we performed princi-

pal component analysis (PCA). Over a third of the total variance observed was concen-

trated within the first principal component (PC1), which distinctly separated the mission

period (P2) from the pre- and post-mission periods (P1 and P3), respectively **[Figures**

Figure 4 Ability of features to discriminate across study periods. (A) Scatter plot of module of effect size (Cohen's f – X-axis) and significance (-log₁₀(FDR), Y-axis) of features with potential to distinguish between study periods. The five features with the largest effect sizes are shown in the inset. Light-related variables ranked the highest, alongside with the Pearson's *r* for several mathematical models. (B) Jittered violin plots of AUC values across features for each pairwise contrast. Discrimination was strongest when comparing the mission period (P2) to either pre- (P1) or post-mission (P3), while pre- (P1) vs. post-mission (P3) periods were markedly less distinguishable, indicating a transient effect of submarine confinement.

3A, S1]. However, none of the two first principal components differentiated between

shifts **[Figure 3B]**. The top ten contributors to PC1, by loading coefficient, comprised

variables from both the first and second correlation clusters **[Supplementary Notebook**

1]. From the first cluster, we found the Pearson's *r* of all four mathematical models

(*forger_r, jewett_r, hannaytp_r, hannaysp_r*) and the slope of the Jewett model

(*jewett_slope*). From the second cluster, the key contributors included light-related fea-

tures such as the time spent per day above 100 and 500 lux (*tat_100lux, tat_500lux*) and

M10_light, but also *RA*, a metric that measures the contrast between periods of highest

(M10) and lowest activity (L5) hours. Overall, this seems to indicate that the primary axis

of variation in the dataset reflects both the kinetics (captured by the slope) and the strength

and direction of temporal shifts in the predicted daily DLMO (captured by Pearson's *r*),

alongside with light intensity metrics and *RA*.

Given these observations, we sought to identify which features provided the great-

est discriminatory ability across study periods. Features were initially filtered based on a

Monte Carlo permutation approach to estimate the Cohen's f, as an effect size metric, and

the FPR, which was subsequently adjusted for multiple comparisons via Benjamini–

Hochberg FDR procedure. Among the five highest-ranking features by effect size were

several variables mentioned before, with *tat_100lux* having the highest effect size (Co-

hen's f =1.91, FDR $\leq 1.64 \times 10^{-4}$), followed by *forger_r* (Cohen's f = 1.69, FDR $\leq 1.64 \times$

10^{-4}), *tat_500lux* (Cohen's f = 1.66, FDR $\leq 1.64 \times 10^{-4}$), *jewett_r* (Cohen's f = 1.62, FDR

$\leq 1.64 \times 10^{-4}$), and *hannaytp_r* (Cohen's f = 1.38, FDR $\leq 1.64 \times 10^{-4}$) **[Figure 4A, Table**

S2]. Similarly, *hannaysp_r* (Cohen's f = 1.12, FDR $\leq 1.64 \times 10^{-4}$) exhibited comparable

discriminative power across at least one study period. Taken together, these findings point
to a high degree of agreement among the four mathematical models, supporting the notion
that the stability and/or temporal dynamics of the predicted DLMO trajectory signifi-
cantly differed in at least one study period.
234

Table 1 Mean and standard deviation in each study period of features with FPR < 0.05 AUC ≥ 0.99 in at least
one pairwise comparison between study periods. Variable descriptions: *vat_10lux* – median daily light exposure
above 10 lux; *forger r, jewett_r* – Pearson's correlation coefficient between each day in the recording and the corre-
sponding DLMO predicted by the Forger and Jewett model, respectively; *tat_500lux, tat_100lux* – median daily light
exposure time above 500 and 100 lux, respectively; *M10_light* – mean light intensity during the ten brightest hours of
the day.

Feature	P1 (mean ± SD)	P2 (mean ± SD)	P3 (mean ± SD)
vat_10lux (lux)	115.17 ± 58.52	27.17 ± 4.01	90.32 ± 26.25
forger_r	-0.36 ± 0.64	0.95 ± 0.04	-0.81 ± 0.41
jewett_r	-0.18 ± 0.62	0.93 ± 0.07	-0.77 ± 0.41
tat_500lux (minutes)	156.1 ± 51.96	0.57 ± 1.93	133.54 ± 55.28
M10_light (lux)	1600.58 ± 490.09	264.44 ±177.83	1219.72±605.44
tat_100lux (minutes)	284.17 ± 76.28	11.11 ± 14.33	241.17 ± 85.01

To determine whether specific study period contrasts were associated with greater
discriminability across features, we examined the AUC distributions derived from ROC
curves for all pairwise comparisons (P1 vs. P2, P2 vs. P3, and P1 vs. P3) **[Figure 4B]**.
The AUC distribution for the pre- vs. post-mission contrast (P1 vs. P3) differed signifi-
cantly from the pre- vs. mission (P1 vs. P2*; p = 7.53 ×10^{-7}, AUC=0.85*) and mission vs.
post-mission (P2 vs. P3*; p = 4.98 ×10^{-5}, AUC = 0.7*) contrasts. Yet, no significant

Figure 5 Pairwise feature discrimination across study periods. (A) Heatmap showing the performance of features
in pairwise period comparisons, ranked by average AUC. (B) Illustrative jittered violin plots and (C) ROC curves for
jewett_r (Pearson's correlation coefficient between each day in the recording and the respective DLMO predicted by
the Jewett model), showing a strong ability to distinguish P2 from P1 and P3, but not between P1 and P3.

difference was observed between the P1 vs. P2 and P2 vs. P3 distributions (*p = 0.22,*

AUC=0.58). These results point to greater similarity between the pre- and post-mission

periods, with a higher concentration of discriminative features in comparisons involving

the mission period.

Building on these findings, we used the AUC to select features with the strongest

ability to discriminate between pairs of study periods **[Table S3]**. Features meeting the

criteria of an FDR < 0.05 and AUC ≥ 0.99 in at least one contrast were deemed highly

discriminative and thereby chosen. The mean and standard deviation of the features meet-

ing these criteria are shown in **Table 1** and include both light-related metrics and the

Pearson's *r* for the Forger and Jewett models **[Figure 5A]**. Summary statistics for addi-

tional features are provided in **Table S4**. Illustrative ROC curves and violin plots for

jewett_r are shown in **Figure 5B,C,** with the remaining plots available in **Figures S2 and**

S4.

During the submarine mission, light-related metrics providing the greatest dis-

crimination between shifts consistently exhibited lower intensity values, and decreased

time, per day, above 100 and 500 lux (*tat_100lux*: 11.11 ± 14.33 min; *tat_500lux*: 0.57 ±

1.93), with lower light intensity values during the ten brightest hours of the day

Figure 6 Predicted daily circadian phase trajectory based on the Jewett model. Predicted dim light melatonin onset (DLMO) for each individual across the whole study, separated by shifts. The submarine mission period (P2) is shaded in light grey.

(*M10_light*: 264.44 ± 177.83). Moreover, in P1, *forger_r* and *jewett_r* showed minimal

correlation of the predicted daily DLMO with time (-0.36 ± 0.64 and -0.18 ± 0.62, respec-

tively). However, during P2, both coefficients increased markedly (0.95 ± 0.04 and 0.93

± 0.07), indicating a strong positive association consistent with a phase delay. In contrast,

P3 was characterized by strongly negative correlations (-0.81 ± 0.41 and -0.77 ± 0.41),

suggestive of a phase advance. A similar pattern was followed by the daily DLMO pre-

dicted by HannaySP model (P1: -0.24 ± 0.60; P2: 0.80 ± 0.46 ; P3: -0.56 ± 0.50) and

HannayTP model (P1: -0.24 ± 0.61; P2: 0.90 ± 0.37; P3: -0.70 ± 0.45) **[Figures 6, S6–8]**.

Taken together, all four models seem to consistently predict a pronounced phase delay in

response to the dim-light conditions of the submarine, followed by a phase advance in the

post-mission period.

Identifying shift-specific actigraphy markers during the submarine mission.

While submarine missions represent a strong circadian perturbation, we investi-

gated whether their physiological impact might also vary depending on shift assignment.

If so, we would expect distinct actigraphy-based signatures to emerge across shifts. Alt-

hough the distribution of effect sizes was significantly lower when grouping by shift ra-

ther than by study period (*Mann Whitney's U = 3747.00, p = 3.00×10⁻⁶*) **[Figure 7A]**, we

wondered if it would be possible to identify features capable of distinguishing between

shifts. In this context, P1 served as a negative control, as participants had not yet been

assigned to shifts, while P3 provided insights into post-mission recovery upon return to

baseline conditions. No features differentiated shifts at a 5% significance level in either

P1 or P3. However, during the mission period (P2), seventeen features met the threshold

Figure 7 Discriminative power of features across shifts during the mission period (P2). (A) Jittered violin plots of Cohen's f distributions across features, by period and shift, showing that larger effect sizes are mainly associated with distinctions between study periods, while shift comparisons within P2 are characterized by smaller, yet non-negligible, effect sizes. (B) Scatter plot of module of effect size (Cohen's F – X axis) and significance (-log10(FDR), Y-axis) of features that significantly distinguish at least one shift in P2. The inset on the right highlights the top five features ranked by effect size (Cohen's F).

for statistical significance and were ranked by effect size (Cohen's f) **[Figure 7B]**. Among

the top five were several sleep-related variables, namely the *main_sleep_bout* (Cohen's f

= 1.59, FDR ≤ 8.19 × 10⁻⁴), *sleep_midpoint* (Cohen's f =1.41, FDR ≤ 8.19 × 10⁻⁴) and

the sleep regularity index (*SRI*) (Cohen's f = 1.27, FDR ≤ 8.19 × 10⁻⁴). The highest effect

size was observed for the *RA* (Cohen's f = 1.68, FDR ≤ 8.19 × 10⁻⁴), and the lowest among

the top 5 was observed in *exp_level* (Cohen's f = 1.06; FDR = 0.0023).

Variables derived from mathematical models of circadian rhythms were notably

absent from the set of features that robustly differentiated between at least one shift. While

forger_slope yielded a statistically significant result (Cohen's f = 0.93, FDR = 0.0033),

the effect size was small and it was not verified in the other models (*jewett_slope*: Cohen's

f = 0.58, FDR = 0.054, *hannaysp_slope*: Cohen's f = 0.17, FDR = 0.79; *hannaytp_slope*:

Cohen's f = 0.20, FDR = 0.53). Similarly, a small yet significant effect was observed for

jewett_r (Cohen's f = 0.75, FDR = 0.015), but not for the Pearson's *r* of the other models

(*forger_slope*: Cohen's f = 0.53, FDR = 0.082; *hannaysp_r*: Cohen's f = 0.18, FDR =

0.79; *hannaytp_r*: Cohen's f = 0.21, FDR = 0.79). Thus, even though all models consist-

ently predicted a phase delay during P2 when contrasting study periods, the lack of con-

sensus when grouping by shift leads us to conclude that there is no conclusive evidence

Figure 8 Pairwise feature discrimination across shifts. (A) Heatmap of pairwise shift comparisons, with features ordered by average AUC across the three shift-pair contrasts. Empty cells denote non-significant (FDR > 0.05) results. (B) Jittered violin plots and (C) ROC curves for *SRI* (sleep regularity index). Most features exhibiting near-perfect (AUC > 0.99) discriminative performance differentiated between the split shift schedules (shifts 1 and 2) and the on-call schedule (shift 0), with *RA* and *L5* being notable exceptions.

305 that either the linear dependence of the predicted DLMO on time or the rate of change in
306 predicted DLMO with respect to time varied across shifts.

Table 2 Mean and standard deviation in each shift of features with FDR < 0.05 and AUC ≥ 0.99 in at least one pairwise comparison between shifts. Variable descriptions: *SRI* – probability that any two time points exactly 24 hours apart are in the same state (either sleep or wake), averaged across all days; *main_sleep_bout* – duration of the primary sleep episode each day, defined as the longest consolidated period of sleep within a 24-hour interval; *exp_level* – median light exposure level; *L5* – mean light intensity during the five least active hours of the day; *RA* – relative amplitude between the ten most active hours (M10) and the five least active hours of the day (L5).

Feature	Shift 0 (mean ± SD)	Shift 1 (mean ± SD)	Shift 2 (mean ± SD)
SRI (%)	51.55 ± 8.26	73.03 ± 6.03	65.02 ± 5.16
main_sleep_bout (minutes)	377.62 ± 52.23	259.15 ± 29.99	299.91 ± 25.63
exp_level (lux)	4.13 ± 3.27	0.70 ± 0.63	0.53± 0.32
tat_10lux (minutes)	527.75 ± 108.62	309.15 ± 93.77	292.91 ± 74.81
L5 (PIM)	846.84 ± 420.88	472.22 ± 116.59	1159.07 ± 428.97
RA (%)	60 ± 17	71 ± 6	45 ± 4

The discriminative capacity of the seventeen features was further evaluated across
pairwise shift comparisons (S0 vs. S1, S0 vs. S2, S1 vs. S2) **[Table S3]**. As previously,
discrimination ability was measured using ROC curves, with the AUC serving as a per-
formance metric. Shift-discriminating features meeting the criteria of an FDR < 0.05 and
AUC ≥ 0.99 in at least one comparison are presented in **Figure 8**, with the mean and
standard deviation shown in **Table 2**, and the remaining descriptives on **Table S5**. Illus-
trative ROC curves and violin plots are presented for *SRI* in **Figure8B,C**, with the other
violin plots and ROC curves shown in **Figures S3 and S5**. Notably, two sleep features,
SRI and *main_sleep_bout*, were significant across all shift comparisons. Interestingly, *L5*
and *RA* were found to exclusively differentiate between shift 1 .and 2, with a lower *RA*
and higher *L5* in shift 2 (*L5*: 1159.07 ± 428.97 [PIM]; *RA*: 45% ± 4%) in comparison with
shift 1 (*L5*: 472.22 ± 116.59 [PIM]; *RA*: 71% ± 6%), suggestive of increased nighttime
movement during the five hours of lower activity.

Discussion

This study aimed to identify actigraphy signatures associated with living aboard a
submarine for an extended period of time and with split shift work. The rationale for
comparing signatures between different shifts stemmed from a considerable body of lit-
erature linking night shift work with a wide spectrum of detrimental health outcomes
(Dutheil et al., 2020; Hansen & Pedersen, 2024; Jahn et al., 2024; Manohar et al., 2017;
Moon et al., 2024; Wang et al., 2014). However, some meta-analyses emphasized the need
for well-designed and adequately powered studies using standardized methods to

facilitate comparisons (Jahn et al., 2024; Viramgami et al., 2025). The dual filtering cri-

terion adopted in this study addresses this concern by retaining only those actigraphy

features with an effect size above Cohen's f = 0.63 – corresponding to the minimum de-

tectable effect size for our total sample size (n = 29) at a statistical power of approximately

82%.

As expected, light-related variables emerged as predominant sources of variance

in our dataset. Interestingly, the Pearson's *r*, which measured the correlation between the

predicted DLMO trajectory and time, was effective in distinguishing the mission period

(P2) from the pre- and post-mission periods (P1 and P3, respectively). Remarkably, no

conclusive evidence was found of a shift-specific impact on either the Pearson's *r* or the

slope of the regression line fitted to the DLMO model predictions, seemingly contradict-

ing a previous study on a nuclear submarine in which an adaptation to the watchkeeping

schedule seems to occur (Chabal et al., 2024). Nonetheless, the absence of shift-related

differences in the predicted DLMO is consistent with results from a prior study involving

submariners also on a 6-hours-on/6-hours-off routine in a diesel-electric submarine,

where melatonin secretion failed to adjust to the biphasic work-sleep cycle (Van Puyvelde

et al., 2022). These results lead us to hypothesize that adaptation and recovery dynamics

may vary depending on the heterogeneity of submarine environments, highlighting the

importance of reporting vessel type and characteristics in future research. Another plau-

sible explanation involves the constant and very low light levels onboard, which may

have been insufficient for effective entrainment. Consequently, the observed circadian

phase delay could reflect a non-entrained rhythm pattern, as previously documented in

underground isolation studies (Aschoff et al., 1971) and during the restrictive measures

of the COVID-19 lockdown conditions (Korman et al., 2020).

Although all four mathematical models demonstrated general agreement in detect-

ing a phase delay during the mission and a subsequent phase advance in the post-mission

period, the corresponding effect sizes and discriminatory performance were not identical.

Future research should compare the performance of different mathematical models of cir-

cadian rhythms in DICE environments to determine under which conditions their predic-

tions diverge the most. A potential explanation for the residual variability across models

may lie in how each processes the light input, $L(t)$ (Hannay et al., 2019). It may be worth

exploring if these differences capture a plausible range of inter-individual variability in

human light sensitivity. Regarding the four mathematical models used in this study, a

previous report (Huang et al., 2021) found larger errors in circadian phase predictions for

night shift workers compared to non-shift workers when default parameters were used. Strikingly, only 30% of the DLMO predictions for night workers fell within one hour of the actual DLMO, underlining the importance of improving model accuracy (Huang et al., 2021).

Establishing a comparative framework to identify parameter sets that bring model predictions into closer agreement – or alternatively, conducting population-specific parameter calibrations – could enhance predictive power and help identify which parameters are most informative for characterizing circadian dynamics across human groups. Additionally, using a fixed seven-hour offset to estimate DLMO, a common practice in the literature (Lim et al., 2025; Murray et al., 2021; Revell et al., 2006), may only provide a gross approximation of this biological night marker and, thus, contribute to suboptimal prediction performance. Indeed, considerable interindividual variability in sensitivity to evening light has been reported, with most sensitive individuals showing over 50% melatonin suppression at just 10 lux, whereas the least sensitive required exposure to 400 lux to show an equivalent melatonin suppression level. This variability may also be reflected in the phase angle between DLMO and sleep onset, which can range from approximately 5.0 hours before to 1.5 hours after sleep onset in patients with sleep-wake phase disorder, a range comparable to the observed in healthy individuals (Reis et al., 2025). Future studies should take into consideration this variability, and determine whether incorporating additional *zeitgebers*, such as meal timing, and/or explicitly modelling melatonin secretion and degradation dynamics in response to SCN output, may enhance the predictive power of current mathematical frameworks (St Hilaire et al., 2007).

An interesting pattern observed in our study was that individuals assigned to split shift schedules differed consistently from those assigned to on-call/extended hours, that is, increased similarity between shifts 1 and 2, and a higher dissimilarity between these two and shift 0. A noteworthy exception to this pattern was the reduction in *RA* and an elevation in *L5* observed in shift 2, with a working bout entirely during the biological night. This signature likely reflects increased nocturnal activity and, thus, points to impaired sleep quality and/or increased circadian disruption. Future studies in diesel-electric submariners should attempt to include some modality of portable PSG, as it would give further insights into the differential impact of being assigned to a shift during the biological night, in the absence of a robust photic *zeitgeber*.

Ultimately, these findings are consistent with established literature showing that night shift workers experience poor sleep due to misalignment between the sleep-wake

cycle and circadian phase (Costa, 2015). Remarkably, this disruptive pattern was detectable in shift 2, the one most closely resembling a night shift, despite all submariners being exposed to the same environmental conditions. These results indicate that, even in the absence of natural light-dark cues, working during the biological night may contribute to sleep impairment or, more likely, that sleeping during the biological day leads to a more disrupted sleep, as observed in nurses (Martin et al., 2006). Given the parallels between submarine and night shift work, with the latter being linked to increased cardiometabolic risk and certain cancers (Su et al., 2021), it would be particularly valuable to investigate whether molecular, namely metabolomic, alterations are observed in submariners and if we can observe an overlap with those documented in night shift workers.

A major limitation of our data was the small representation of female participants, reflecting the reality in most submarine crews, but leaving open the question of whether the impact of submarine and submarine-like environments would be comparable in women, as several studies point to sex-related differences in circadian rhythms (Walton et al., 2022). This study was also limited by the fact that only data from a single submarine mission was available. Since the post-mission recovery period has scarcely been evaluated and the importance of ensuring adequate recuperation between deployments, future studies should attempt to evaluate the long-term physiological effects of repeated submarine missions. Similarly, it would be valuable to investigate the differential impacts of shorter versus longer deployments, as well as the influence of vessel type, particularly in relation to the degree of spatial confinement.

Finally, while some results may align with prior expectations, they show the ability of our feature-screening approach to select candidate actigraphy biomarkers of split shift work in a submarine environment. Future refinements upon this methodology should focus on enhancing the F-statistic estimation, for instance, by introducing mixed-effect models. Nevertheless, the ability to extract biologically meaningful signal in such constrained conditions highlights the robustness and broad applicability of this effect-size-centered screening strategy to other challenging operational and clinical settings. For example, it could be used to monitor activity-inactivity cycles in bed-ridden populations, including older adults, patients with dementia (Tolea et al., 2016) or depression (Elkjær et al., 2022). In addition, this approach is sufficiently flexible to incorporate other types of measurements, including PSG, metabolomics, transcriptomics, and better mathematical models of circadian rhythms. Ultimately, by integrating diverse data types, it may help to uncover distinct physiological programs associated with DICE environments.

References

Ansu Baidoo, V., & Knutson, K. L. (2023). Associations between circadian disruption and

cardiometabolic disease risk: A review. *Obesity*, *31*(3), 615–624.

https://doi.org/10.1002/oby.23666

Aschoff, J. (1960). Exogenous and endogenous components in circadian rhythms. *Cold*

Spring Harbor Symposia on Quantitative Biology, *25*, 11–28.

https://doi.org/10.1101/sqb.1960.025.01.004

Aschoff, J., Fatranská, M., Giedke, H., Doerr, P., Stamm, D., & Wisser, H. (1971). Human

circadian rhythms in continuous darkness: Entrainment by social cues. *Science*

(New York, N.Y.), *171*(3967), 213–215. https://doi.org/10.1126/sci-

ence.171.3967.213

Boivin, D. B., & Boudreau, P. (2014). Impacts of shift work on sleep and circadian

rhythms. *Pathologie Biologie*, *62*(5), 292–301. https://doi.org/10.1016/j.pat-

bio.2014.08.001

Cajochen, C., & Schmidt, C. (2025). The Circadian Brain and Cognition. *Annual Review*

of Psychology, *76*(Volume 76, 2025), 115–141. https://doi.org/10.1146/annurev-

psych-022824-043825

Chabal, S. A., Markwald, R. R., & Chinoy, E. D. (2024). Life onboard a submarine: Sleep,

fatigue, and lifestyle behaviors of sailors on a circadian-aligned watchstanding

schedule. *Applied Ergonomics*, *119*, 104321.

https://doi.org/10.1016/j.apergo.2024.104321

Costa, G. (2015). Chapter 24—Sleep deprivation due to shift work. In M. Lotti & M. L.

Bleecker (Eds.), *Handbook of Clinical Neurology* (Vol. 131, pp. 437–446). Else-

vier. https://doi.org/10.1016/B978-0-444-62627-1.00023-8

Dutheil, F., Baker, J. S., Mermillod, M., De Cesare, M., Vidal, A., Moustafa, F., Pereira, B., & Navel, V. (2020). Shift work, and particularly permanent night shifts, promote dyslipidaemia: A systematic review and meta-analysis. *Atherosclerosis*, *313*, 156–169. https://doi.org/10.1016/j.atherosclerosis.2020.08.015

Elkjær, E., Mikkelsen, M. B., Michalak, J., Mennin, D. S., & O'Toole, M. S. (2022). Motor alterations in depression and anxiety disorders: A systematic review and meta-analysis. *Journal of Affective Disorders*, *317*, 373–387. https://doi.org/10.1016/j.jad.2022.08.060

Erren, T. C., Morfeld, P., Groß, J. V., Wild, U., & Lewis, P. (2019). IARC 2019: "Night shift work" is probably carcinogenic: What about disturbed chronobiology in all walks of life? *Journal of Occupational Medicine and Toxicology (London, England)*, *14*, 29. https://doi.org/10.1186/s12995-019-0249-6

Fernandes, C., Maruta, C., Marques-Dias, C. M., & Reis, C. (2025). Sleep in Military Submariners: Exploring its Dynamics in Relation to a Submarine Mission and Interaction with Psychological Factors. *Behavioral Sleep Medicine*, *0*(0), 1–14. https://doi.org/10.1080/15402002.2025.2493650

Forger, D. B., Jewett, M. E., & Kronauer, R. E. (1999). A Simpler Model of the Human Circadian Pacemaker. *Journal of Biological Rhythms*, *14*(6), 533–538. https://doi.org/10.1177/074873099129000867

Hammad, G., Reyt, M., Beliy, N., Baillet, M., Deantoni, M., Lesoinne, A., Muto, V., & Schmidt, C. (2021). pyActigraphy: Open-source python package for actigraphy data visualization and analysis. *PLOS Computational Biology*, *17*(10), e1009514. https://doi.org/10.1371/journal.pcbi.1009514

Hammad, G., Wulff ,Katharina, Skene ,Debra J., Münch ,Mirjam, & and Spitschan, M. (2024). Open-Source Python Module for the Analysis of Personalized Light

Exposure Data from Wearable Light Loggers and Dosimeters. *LEUKOS*, *20*(4), 380–389. https://doi.org/10.1080/15502724.2023.2296863

Hannay, K. M., Booth, V., & Forger, D. B. (2019). Macroscopic Models for Human Circadian Rhythms. *Journal of Biological Rhythms*, *34*(6), 658–671. https://doi.org/10.1177/0748730419878298

Hansen, J., & Pedersen, J. E. (2024). Night shift work and breast cancer risk – 2023 update of epidemiologic evidence. *Journal of the National Cancer Center*, *5*(1), 94–103. https://doi.org/10.1016/j.jncc.2024.07.004

Heyde, I., & Oster, H. (2022). Induction of internal circadian desynchrony by misaligning zeitgebers. *Scientific Reports*, *12*(1), 1601. https://doi.org/10.1038/s41598-022-05624-x

Huang, Y., Mayer, C., Cheng, P., Siddula, A., Burgess, H. J., Drake, C., Goldstein, C., Walch, O., & Forger, D. B. (2021). Predicting circadian phase across populations: A comparison of mathematical models and wearable devices. *Sleep*, *44*(10), zsab126. https://doi.org/10.1093/sleep/zsab126

Jahn, A., Nielsen, M. L., Kyndi, M., & Dalbøge, A. (2024). Association between night work and prostate cancer: A systematic review and meta-analysis. *International Archives of Occupational and Environmental Health*, *97*(2), 207–215. https://doi.org/10.1007/s00420-023-02037-9

Korman, M., Tkachev, V., Reis, C., Komada, Y., Kitamura, S., Gubin, D., Kumar, V., & Roenneberg, T. (2020). COVID-19-mandated social restrictions unveil the impact of social time pressure on sleep and body clock. *Scientific Reports*, *10*(1), 22225. https://doi.org/10.1038/s41598-020-79299-7

Lim, D., Choi, S. J., Song, Y. M., Park, H. R., Joo, E. Y., & Kim, J. K. (2025). Enhanced Circadian Phase Tracking: A 5-h DLMO Sampling Protocol Using Wearable Data.

Journal of Biological Rhythms, *40*(3), 249–261.
https://doi.org/10.1177/07487304251317577

Manohar, S., Thongprayoon, C., Cheungpasitporn, W., Mao, M. A., & Herrmann, S. M.
(2017). Associations of rotational shift work and night shift status with hyperten-
sion: A systematic review and meta-analysis. *Journal of Hypertension*, *35*(10),
1929. https://doi.org/10.1097/HJH.0000000000001442

Marando, I., Lushington, K., Owen, M., Matthews, R. W., & Banks, S. (2023). The sleep,
circadian, and cognitive performance consequences of watchkeeping schedules in
submariners: A scoping review. *Sleep Medicine Reviews*, *72*, 101845.
https://doi.org/10.1016/j.smrv.2023.101845

Martin, J. L., Webber, A. P., Alam, T., Harker, J. O., Josephson, K. R., & Alessi, C. A.
(2006). Daytime Sleeping, Sleep Disturbance, and Circadian Rhythms in the
Nursing Home. *The American Journal of Geriatric Psychiatry*, *14*(2), 121–129.
https://doi.org/10.1097/01.JGP.0000192483.35555.a3

McHill, A. W., Melanson, E. L., Higgins, J., Connick, E., Moehlman, T. M., Stothard, E.
R., & Wright, K. P. (2014). Impact of circadian misalignment on energy metabo-
lism during simulated nightshift work. *Proceedings of the National Academy of
Sciences of the United States of America*, *111*(48), 17302–17307.
https://doi.org/10.1073/pnas.1412021111

Meléndez-Fernández, O. H., Liu, J. A., & Nelson, R. J. (2023). Circadian Rhythms Dis-
rupted by Light at Night and Mistimed Food Intake Alter Hormonal Rhythms and
Metabolism. *International Journal of Molecular Sciences*, *24*(4), 3392.
https://doi.org/10.3390/ijms24043392

Moon, J., Holzhausen, E. A., & Mun, Y. (2024). Risk of prostate cancer with increasing
years of night shift work: A two-stage dose-response meta-analysis with duration

of night shift work as exposure dose. *Heliyon*, *10*(8). https://doi.org/10.1016/j.heliyon.2024.e29080

Murray, J. M., Magee, M., Sletten, T. L., Gordon, C., Lovato, N., Ambani, K., Bartlett, D. J., Kennaway, D. J., Lack, L. C., Grunstein, R. R., Lockley, S. W., Rajaratnam, S. M. W., & Phillips, A. J. K. (2021). Light-based methods for predicting circadian phase in delayed sleep–wake phase disorder. *Scientific Reports*, *11*(1), 10878. https://doi.org/10.1038/s41598-021-89924-8

Nieuwenhuys, A., Dora, J., Knufinke-Meyfroyt, M., Beckers, D., Rietjens, G., & Helmhout, P. (2021). "20,000 leagues under the sea": Sleep, cognitive performance, and self-reported recovery status during a 67-day military submarine mission. *Applied Ergonomics*, *91*, 103295. https://doi.org/10.1016/j.apergo.2020.103295

Peters, B., Vahlhaus, J., & Pivovarova-Ramich, O. (2024). Meal timing and its role in obesity and associated diseases. *Frontiers in Endocrinology*, *15*, 1359772. https://doi.org/10.3389/fendo.2024.1359772

Reis, C., Pilz, L. K., Paiva, T., Hidalgo, M. P., & Wright Jr, K. P. (2025). Sleep and circadian rhythms in delayed sleep–wake phase disorder: Phenotypic differences between patients with and without comorbid depression. *Journal of Sleep Research*, *n/a*(n/a), e14437. https://doi.org/10.1111/jsr.14437

Revell, V. L., Burgess, H. J., Gazda, C. J., Smith, M. R., Fogg, L. F., & Eastman, C. I. (2006). Advancing Human Circadian Rhythms with Afternoon Melatonin and Morning Intermittent Bright Light. *The Journal of Clinical Endocrinology and Metabolism*, *91*(1), 10.1210/jc.2005-1009. https://doi.org/10.1210/jc.2005-1009

Smith, M. R., & Eastman, C. I. (2012). Shift work: Health, performance and safety problems, traditional countermeasures, and innovative management strategies to

reduce circadian misalignment. *Nature and Science of Sleep*, *4*, 111–132.

https://doi.org/10.2147/NSS.S10372

St Hilaire, M. A., Gronfier, C., Zeitzer, J. M., & Klerman, E. B. (2007). A physiologically

based mathematical model of melatonin including ocular light suppression and

interactions with the circadian pacemaker. *Journal of Pineal Research*, *43*(3),

294–304. https://doi.org/10.1111/j.1600-079X.2007.00477.x

Su, F., Huang, D., Wang, H., & Yang, Z. (2021). Associations of shift work and night

work with risk of all-cause, cardiovascular and cancer mortality: A meta-analysis

of cohort studies. *Sleep Medicine*, *86*, 90–98.

https://doi.org/10.1016/j.sleep.2021.08.017

Tolea, M. I., Morris, J. C., & Galvin, J. E. (2016). Trajectory of Mobility Decline by Type

of Dementia. *Alzheimer Disease & Associated Disorders*, *30*(1), 60.

https://doi.org/10.1097/WAD.0000000000000091

Van Puyvelde, M., Rietjens, G., Helmhout, P., Mairesse, O., Van Cutsem, J., & Pattyn, N.

(2022). The submariners' sleep study: A field investigation of sleep and circadian

hormones during a 67-day submarine mission with a strict 6-h-on/6-h-off watch

routine. *Journal of Applied Physiology*, *132*(4), 1069–1079.

https://doi.org/10.1152/japplphysiol.00130.2021

Viramgami, A., Balachandar, R., Bagepally, B. S., & Sheth, A. (2025). Study on the as-

sociation between night shift work and reproductive functions among male work-

ers: A systematic review and meta-analysis. *Endocrine*, *88*(2), 410–419.

https://doi.org/10.1007/s12020-025-04166-2

Walton, J. C., Bumgarner, J. R., & Nelson, R. J. (2022). Sex Differences in Circadian

Rhythms. *Cold Spring Harbor Perspectives in Biology*, *14*(7), a039107.

https://doi.org/10.1101/cshperspect.a039107

Wang, F., Zhang, L., Zhang, Y., Zhang, B., He, Y., Xie, S., Li, M., Miao, X., Chan, E. Y.
Y., Tang, J. L., Wong, M. C. S., Li, Z., Yu, I. T. S., & Tse, L. A. (2014). Meta-
analysis on night shift work and risk of metabolic syndrome. *Obesity Reviews*,
15(9), 709–720. https://doi.org/10.1111/obr.12194

Zivi, P., De Gennaro, L., & Ferlazzo, F. (2020). Sleep in Isolated, Confined, and Extreme
(ICE): A Review on the Different Factors Affecting Human Sleep in ICE. *Fronti-*
ers in Neuroscience, *14*. https://doi.org/10.3389/fnins.2020.00851