2 Major transitions in early coral development: novel insights enabled by visualisation
3 of a comprehensive transcriptomic dataset for *Acropora millepora*

5 Ramona Brunner[1,2]*, Mila Grinblat[1,2]*, Aurelie Moya[1,2]*, Sylvain Foret[3‡], David C
6 Hayward[3], Bruno Lapeyre[1,4], **Eldon E Ball**[3], Ira Cooke[2,5], **David J Miller**[1,2]
7

[1]ARC Centre of Excellence for Coral Reef Studies, James Cook University,
Townsville, QLD 4811 Australia
[2]Molecular Genetics Unit, James Cook University, Townsville, QLD 4811 Australia
[3]Research School of Biology, Australian National University, Acton, ACT 2601
Australia
[4]Laboratoire d'excellence CORAIL, Centre de Recherches Insulaires et Observatoire
de l'Environnement (CRIOBE), Moorea, B.P. 1013, Papeete, French Polynesia
[5]Department of Marine Biology and Aquaculture, James Cook University, Townsville,
QLD 4811 Australia
17

18 ‡Deceased
19 Email addresses: RB (ramona.brunner@my.jcu.edu.au), MG
20 (mila.grinblat@my.jcu.edu.au), AM (aurelie.moya@gmail.com), DH
21 (david.hayward@anu.edu.au), BL (viso3841@gmail.com), EB
22 (eldon.ball@anu.edu.au), IC (ira.cooke@jcu.edu.au), DM (david.miller@jcu.edu.au)
23

Corresponding authors: EB and DM

Keywords: Maternal-zygotic transition, juvenile-adult transition, metamorphosis, coral
development, larval settlement

Ethics and integrity statement: Our work does not require ethics approval, but all
work involving the biota of the Great Barrier Reef requires approval by The Great
Barrier Reef Marine Park Authority (GBRMPA) which rigorously assesses research
plans before issuing appropriate permits and carries out routine surveillance. The
work described here was conducted under GBRMPA permit G13/36196.1. The
authors were all directly involved in the work and all have been consulted on draft
and final versions of the manuscript. All analyses have been subject to scrutiny by
other authors and can be confirmed using the web-based tool that we have released
(see below).

Data availability: The transcriptome data and web-based browser reported here are
available at https://amil2dev.mmb.group/. The source code is available at
https://github.com/iracooke/amil2_dev.

Funding statement: This work was supported by grant numbers DP170104734,
FS110200046 and CE14100020 from the Australian Research Council
45
46   The authors declare no conflicts of interest
47
48   Ethics approval, patient consent and clinical trial registration are not required for this
49   work
50
Permission to reproduce material from other sources: Permission has been obtained
to reproduce parts of Figure 5 from a previous publication by our group (Hayward et
al. 2004). Smaller parts of other figures from our previous publications have been
included in other figures, as cited in their captions.

Abstract

Background

Given the ecological importance of reef-building corals (Scleractinia), it is perhaps

surprising that the molecular mechanisms underlying many of the morphological and

metabolic changes during their development remain unclear. In part, this is due to

the lack of a comprehensive transcriptomic dataset for any coral. A second challenge

in the analysis of such non-model developmental datasets is that the volume of data

often complicates its interpretation.

Results

To overcome these limitations, we profiled gene expression in *Acropora millepora*

across 26 life stages from unfertilised eggs to juvenile polyps and developed an

interactive online tool based on the R-application Shiny to simultaneously visualise

changes in the expression of large numbers of genes. As expected, major

transcriptomic changes (transitions) occurred during gastrulation and the acquisition

of competence. Surprisingly, however, settlement triggered by using the natural

inducer CCA did not immediately lead to major changes in gene expression, but a

major transition involving many genes was observed 3 - 6 hours after settlement

induction.

Conclusions

We hope that providing access to this extensive developmental transcriptome

dataset and software to facilitate its analysis will expedite a better understanding of

the changes that occur during coral development. The online tool is available at

https://amil-deview.mmb.guide.

Background

Broadcast spawning corals, such as *Acropora*, release buoyant eggs and sperm and

fertilisation occurs in the water column. The early developmental stages are

planktonic, ultimately reaching a motile stage known as a planula larva which

becomes "competent" – that is, it becomes capable of settlement and

metamorphosis - after a period of development of variable duration (Figure 1).

Competence is marked by a change in behaviour, from horizontal swimming to

swimming into the substratum and testing it for chemical cues. In nature, the

settlement of competent larvae generally involves selection of appropriate sites,

based on perception of a range of cues that include small molecules, at least some

of which are produced by bacteria associated with substrate biofilms and algae (e.g.

Tebben et al. 2015; Siboni et al. 2020; Quinlan et al. 2023; Turnlund et al. 2025).

The quality and nature of available light also plays a part in settlement site selection

in the case of many reef-building corals (Mason and Cohen, 2012; Strader et al.

2015; Mason et al. 2023).

Various aspects of early coral development have previously been studied (Table 1),

with the initial focus being characterisation of individual genes homologous to those

playing key developmental roles in bilaterians (Ball et al. 2002; Hayward et al. 2002;

Ball et al. 2004; Hayward et al. 2004). The first larger scale analyses were based on

microarrays (Grasso et al. 2008; Reyes-Bermudez et al. 2009; Grasso et al. 2011),

454 GSFlx sequencing (Meyer et al. 2009), SOLiD sequencing (Meyer et al. 2011) or

subtractive hybridisation (Hayward et al. 2011). More recent studies have used high-

throughput DNA sequencing technologies to characterise specific developmental

stages (Summarised in Table 1). Much of this work has focussed on members of the

Acroporidae (genera Acropora and Montipora), with sampling regimes designed to

address specific questions. For example, Chille et al. (2021) concentrated on gene

expression changes *in Montipora digitata* during the maternal/zygotic transition

(MZT). However, two recent developmental studies on *M. digitata* have gone beyond

transcriptomics and are particularly relevant to the work presented here. He et al.

(2024) combined spatial gene expression data with a comprehensive temporal

dataset to study the segmentation of the gastric cavity, and Huffmeyer et al. (2025)

supplemented transcriptomic data with metabolomic analyses to investigate the

nutritional shift from maternal lipids to symbiont photosynthates in this vertically

transmitting coral. Much higher cellular resolution expression datasets are becoming

available for scleractinians (Levy et al. 2021; Gilbert et al. 2024; Ramon-Mateu et al.

2025), but so far these have been limited to a few developmental stages.

To address the need for a comprehensive transcriptomic dataset, we profiled gene

expression of *Acropora millepora* across 26 life stages from fertilised eggs to juvenile

polyps and developed an interactive online tool based on the Shiny web framework

to visualise changes in the expression of groups of genes. We present an overview

of the data, focussing on major transitions in gene expression and on processes not

previously investigated, but hope that further exploration of the dataset via the

website facilitates progress towards solutions for the crises currently faced by coral

reefs worldwide. Our study also reveals the importance of combining spatial

expression data, as revealed by in situ hybridisation, with the data obtained by

RNAseq, in order to better understand gene function.

Materials and Methods

Colonies of *Acropora millepora* were collected at Orpheus Island, Queensland,

Australia, (18.6141° S, 146.4892° E) under GBRMPA permit G13/36196.1 shortly

before they spawned at approximately 9pm on 21 November 2013. The samples

sequenced were the result of crosses between two colonies and were raised in three

separate flow-through tanks to control for tank effects.

Material was collected from 26 different time points of which 14 were sampled before

and 12 after settlement induction. Pre-settlement, samples were collected from

unfertilised eggs, at 5 hours post-fertilisation (hpf; 16-64 cell stages), 13hpf

(prawnchip), 18hpf, 23hpf, 29hpf, 35hpf, 40hpf (sphere, some swimming), 45hpf

(spheres, all swimming), 57hpf, 69hpf, 81hpf (planulae prior to the searching phase),

93hpf (searching planulae) and at 133hpf. At these stages, approx. 500 embryos per

replicate (n=3) and cross (n = 2) were immersed in 2 mL ethanol (100%). This

caused them to sink to the tube bottom, so that the overlying liquid was removed

before tubes were snap-frozen in liquid nitrogen.

To assess settlement competency, between 69 and 93hpf, aliquots of planulae were

exposed (independently) to either ethanolic extracts of the crustose coralline alga

(CCA) *Porolithon onkodes*, (kindly provided by Mikhail V. Matz) or the neuropeptide

Hym-248, both of which are known to induce settlement of *Acropora* planulae. At the

93 hpf stage 100% of planulae rapidly responded to the CCA extract by undergoing

settlement followed by metamorphosis, hence the settlement-induced stages used

for RNAseq were derived from planulae exposed to this inducer. Post-settlement

samples were collected by stocking 144 petri dishes (2 replicates x 3 larval tanks x 2

crosses x 12 time points) with respectively 100 larvae at the age of 133 h. Settlement

was induced with ethanolic CCA extract in all petri dishes at the same time. At the

respective sampling times of 15 minutes after induction (mpi), 30mpi, 1, 2, 3, 6, 9,

12, 18, 24, 48 and 72 hours post induction (hpi), unattached larvae were transferred

into tubes, and attached settlers were snap-frozen on petri-dishes using liquid

nitrogen after excess seawater was decanted. Sample tubes and petri dishes were

taken to James Cook University, (Townsville, Australia) in dry-ice, and stored at -

80°C until further processing.

Frozen samples in the plates were submerged in RNAlater-ICE and allowed to thaw

overnight at -20°C. Once thawed, RNAlater-ICE was removed and replaced by 2ml

of *mir*Vana™ homogenisation buffer. Polyps were gently scraped off the plate with a

sterilized scalpel in the homogenisation buffer and transferred into a 2ml tube for

immediate processing.

For both pre- and post-settlement samples, total RNA was extracted using

the *mir*Vana™ miRNA Isolation Kit (Ambion) according to the manufacturer's

instructions. RNA quantity and quality were assessed using a NanoDrop ND-1000

spectrometer and denaturing gel electrophoresis using standard methods (Russell &

Sambrook 2001) before being shipped on dry-ice to AGRF in Brisbane for Hi-Seq

paired-end sequencing.

RNA sequencing data for all samples was first checked using a Nextflow (Di

Tommaso et al. 2017) pipeline https://github.com/marine-omics/moqc. This pipeline

includes fastqc (v0.11.5; https://www.bioinformatics.babraham.ac.uk/projects/fastqc/)

to check raw read quality, and read-level taxonomic classification via krakenuniq

(v1.0.2 ; Breitwieser, Baker and Salzberg, 2018) to determine the dominant symbiont

genus and check for bacterial or eukaryotic contaminants. All samples were

dominated by reads attributable to the coral host but also contained a small fraction

(0.5-2%) from Symbiodiniaceae, of which *Durusdinium* was the dominant genus. No

samples were flagged as having low quality base calls, excessive adapter

contamination or high levels of potential contaminants (eg bacteria).

After confirming the overall quality of read data for all samples, reads were trimmed

using fastp (Chen, Zhou, Chen and Gu, 2018) to remove adapters and low-quality

sequences. Trimmed reads were then aligned to a combined reference database

consisting of all *Acropora millepora* transcripts predicted from gene models along

with the transcriptome of an unnamed *Durusdinium* species (Shoguchi et al. 2021).

Alignment was performed using Bowtie2 (v2.5.0; Langmead and Salzberg, 2012) as

part of RSEM (v1.3.3; Li, B., Dewey, C.N., 2011) and used to produce transcript level

counts for each sample. All of these steps were run using a Nextflow pipeline

https://github.com/marine-omics/morp.

Transcript counts produced by RSEM were imported into R and processed using the

variance stabilising transform function in DESeq2 (vv1.38.3; Love, Huber and

Anders, 2014) to produce values on a log2 scale for visualisation.

Functional annotations for *Acropora millepora* transcripts were obtained using a

nextflow pipeline (https://github.com/marine-omics/moat) that is primarily based

around identifying homologous proteins in Swissprot (downloaded December 2022)

via blastp and blastx (NCBI Blast+ v 2.13.0). In this process, any protein with an

evalue less than 1e-5 is considered a hit and the best hit is used to lookup Gene

Ontology ID's and a protein name from the Swissprot database. In addition,

transcripts were annotated with conserved domains, gene family membership and an

alternative Gene Ontology ID inferred using Interproscan (v5.59-91.0; Jones et al.

2014).

Principal component analysis (PCA) was conducted using the prcomp function of the

stats package in R with vst normalised counts.

Weighted gene co-expression network analysis (WGCNA) was conducted with

normalised data that were filtered for genes with excessive number of missing values

or zero-variance using the goodSamplesGenes function of the WGCNA package

(Langfelder & Horvath, 2008). The samples were clustered using the function hclust

of the stats package (R Core Team, 2024) to detect outliers. One sample collected 3

hpi was removed from the analysis. To determine groups of genes +(modules) with

correlated developmental gene expression profiles, a weighted gene co-expression

network was constructed blockwise using the function blockwiseModules with a

maximum block size of 18000 genes, a minimal module size of 18 and a

mergeCutHeight set to 0.25. A signed, scale-free topological overlap matrix for each

block was created by choosing a soft threshold power of 12 and a biweight

midcorrelation.

To facilitate reuse and rapid exploration of the *Acropora millepora* developmental

transcriptome we created an interactive web application using the shiny framework

(https://shiny.posit.co/) . This application is hosted on servers provided by the

Australian Research Data Commons Nectar Research Cloud to ensure that it is

scalable and highly available for long-term use. Users can access the web

230 application directly at https://amil2dev.mmb.group/ or obtain its source code and the

231 underlying data from our github repository (https://github.com/iracooke/amil2_dev).

232 The application is designed to facilitate access to developmental expression profiles

233 for individual genes or groups of genes. Users can define genes of interest in two

234 ways by either conducting a BLAST search of a sequence or by directly searching

235 for genes with metadata matching a set of gene identifiers, Pfam domains, GO term

236 IDs or text as part of the protein name. Once a list of genes has been identified, the

237 corresponding normalised gene expression data is rapidly retrieved from an sqlite

238 relational database and used to generate an interactive display. When the number

239 of genes is small (default <10) each gene is displayed using a separate line plot

240 generated by the R package ggplot2 (Wickham, 2016) so that detailed changes in

241 the expression of each gene can be carefully inspected. For larger numbers of

242 genes (default>=10) we assume that the user is more interested in broad patterns

243 among genes and the display changes to an interactive heatmap generated with the

244 R package InteractiveComplexHeatmap (Gu and Hübschmann, 2022). In both

245 cases a list of genes with annotation information is also displayed and users can

246 select rows in this table to adjust which genes are displayed in the main display.

247 Comprehensive data and code exporting functions are also provided so that users

248 can easily export both data and R code to reproduce the plots seen in the app to

249 allow further customisation by the user.

Results

Major transcriptional transitions visualised using DEView

During the development of *A. millepora*, the period immediately following fertilisation

(0 – 5h) was characterised by rapid rounds of cell division (Figure. 1) accompanied

by relatively limited changes in gene expression (Figure. 2), implying that this phase

is driven largely by maternal mRNAs and proteins. After that time more extensive

changes in gene expression occurred resulting in resolvable PCA clusters. After

completion of gastrulation at 35hpf, embryos were spherical and became motile at

45hpf. Elongation from pear- (57hpf) to spindle-shaped larvae (69hpf) resulted in

planulae that were not searching at 81hpf but were probing the environment for

appropriate settlement sites at 93hpf, by which time they were competent to settle

(Figure. 1). At 133hpf, settlement was induced with ethanolic CCA extract but the

133hpf transcriptome data represent the untreated state. Settlement induction

caused larvae to further elongate at 15mpi before forming disc-like shapes by

contracting along the oral-aboral axis. Most larvae were metamorphosed at 2hpi and

by 3hpi 100% of larvae had initiated metamorphosis characterised by a six-fold

symmetry around an oral pore in the centre. Whereas transcriptome data for the 0-

3hpi stages clustered with those for late planula stages, data for 6 – 72hpi stages

formed a discrete and well-resolved cluster (Figure. 2) during which metamorphosing

polyps developed a 12-fold symmetry and formed tentacles around the oral opening.

The polyp initiates calcification at the former aboral part, which becomes the

calicoblastic ectoderm.

Differential gene expression analysis (Figure 3) clearly highlights that, in addition to

the expected early (13-29hpf) transcriptional events corresponding to zygotic

genome activation, major transitions in gene expression also occurred at 81-93hpf

and 3-6hpi. The construction of weighted gene co-expression networks (WGCNA;

Langfelder & Horvath (2008), which group modules of genes according to correlation

in their expression patterns, also support a 3-transition hypothesis of early coral

development (Supplementary Figure 1). Whilst such analyses are informative with

respect to the major transitions in development, they are less useful for exploration

of the expression patterns of subsets of genes in terms of structure or function

according to Pfam or Gene Ontology (GO) IDs. To facilitate these latter kinds of

analyses, a software tool based on the Shiny web framework (https://shiny.posit.co/)

was developed (https://amil2dev.mmb.group/).

One powerful feature of the new software tool is that it allows the simultaneous

visualisation of developmental gene expression patterns from a subset of genes

defined by gene identifiers, Pfam domains or GO terms. These subsets can be

selected for further analyses and comparison based on morphology, behaviour, or

gene expression patterns, such as occur during gastrulation and settlement.

Whilst early zygotic gene expression accompanied gastrulation, the major transition

at 81-93hpf is likely to correspond to larval competence which is the preparation for

settlement and metamorphosis. Both of these transitions have previously received

considerable attention (e.g. Hayward et al. 2011; Grasso et al. 2011; Strader et al.

2018). Metamorphosis was underway by 3hpi and was followed by changes in the

expression of large numbers of genes (Figure 2, Figure 3). This post-settlement

change in the expression patterns of many coral genes has previously received only

limited attention (Ishii et al. 2022) and therefore merited closer examination. In what

follows we have proceeded chronologically, with a particular focus on processes and

classes of genes not discussed in previous studies.

Maternal transcripts and the earliest transcriptional events

As in most animals, early post-fertilisation development in corals consists of rapid

cell division during which the maternal cytoplasm is serially subdivided (reviewed in

Ball et al. 2021), and the transcripts required to facilitate these events are largely

provided maternally. Extensive zygotic gene expression appears to be initiated

leading into gastrulation (i.e. around 13h post-fertilisation; Figure 1, Figure 2)

corresponding to the "prawn-chip" morphology.

Rapid rounds of cell division during early development require that replication occurs

quickly, generating a need for core histone proteins in large amounts to form the

H2A/H2B and H3/H4 dimers that constitute the nucleosomes responsible for

packaging the DNA into chromatin. As in *Hydractinia* (*Klyxum*) *echinata* (Ayers et al.

2023), histone mRNAs are amongst both the most abundant maternal transcripts

and earliest expressed zygotic genes in *Acropora* (Figure 4A). Note, however, that

the early zygotic (canonical) histone transcript profile is likely to be dominated by the

replication-coupled types.

Transcripts encoding the core histones H2B, H3 and H4 are particularly abundant in

the *Acropora* maternal mRNA complement (Figure 4A), and the zygotic expression

of these histone types began early at the onset of gastrulation with a peak at 13-

18hpf for H3, H4 and H2B (XP_044167100.1, XP_044175889.1) (Figure 4A).

Curiously, however, canonical H2A mRNA was not maternally provisioned at

significant levels, instead, one of two H2A.X homologs (H2A.X2; XP_029204486.1)

and, to a lesser extent, H2A.V (H2A.F/Z; XP_029205888.1) transcripts were

maternal. Levels of mRNA encoding maternal H2A.X in the unfertilised eggs were

very high but fell dramatically after 13hpf as zygotic expression of canonical H2A

began (e.g. XP_044167969.1), and H2A.X transcripts were essentially absent by

29hpf (Figure 4A).

Gastrulation is characterised by large scale activation of zygotic genes, requiring

extensive chromatin modification. Consistent with this, high levels of mRNAs

encoding homologs of the histone acetyltransferase (HAT) KAT6A

(XP_029189526.2) and SETD2 (XP_029190384.2) were provided maternally (Figure

4B). The latter is the only histone methyltransferase capable of generating the

H3K36me3 modification essential for transcriptional activation (Michail et al. 2025).

Note that both of these genes were reactivated during the post-settlement transition

discussed below.

The MZT occurring at gastrulation marks the first major developmental

transition

To date, the process of gastrulation has been the most extensively studied aspect of

developmental gene expression in both corals and *Nematostella* (Technau, 2020). In

situ hybridisation analyses have shown that orthologs of several key developmental

regulators from bilaterians are temporally and spatially expressed in *A. millepora*

consistent with conservation of function. Examples of these include *snail* (Hayward

et al. 2004), *brachyury* (Hayward et al. 2015; Yasuoka et al. 2016), *forkhead*

(Hayward et al. 2015) and *dpp/bmp4* (Hayward et al. 2002) (Supplementary Figure

2).

With the sole exception of *Pavona,* all complex corals examined to date go through a

"prawn chip" stage of early development that corresponds to the early gastrula

(Okubo et al. 2013 and references cited there; Okubo et al. 2016). At this stage, a

homolog of *Drosophila snail* (XP_029185944.1) was expressed in the presumptive

mesendoderm (Hayward et al. 2004). As *snail*-expressing cells are undergoing an

epithelial-mesenchymal transition (EMT), one of several *Acropora* genes related to

Drosophila aristaless (XP_044163657.1/XP_029193169.1) was expressed

throughout the presumptive ectoderm (Figure 5). Hence these two genes effectively

demarcate the two primary tissue layers of *Acropora* during gastrulation.

The late planula transition corresponds to the acquisition of larval competence

During the period 81-93hpf, many genes show strong down-regulation, but this is

immediately followed by the up-regulation of an even larger number of genes during

the 93-133hpf time window (Figure 3).

These timepoints correspond to the acquisition of settlement competence and when

Acropora larvae are searching for cues that identify appropriate settlement sites (eg.

Tebben et al. 2015, Siboni et al. 2020, Quinlan et al. 2023, Turnlund et al. 2025) by

swimming along the substratum to periodically sample it with the aboral end.

Settlement involves the aboral larval ectoderm making contact with substratum,

which is likely coated with attractive and potentially hostile bacteria, and among the

upregulated genes a number encode proteins with known or suspected antimicrobial

properties. Among them are for example AmAMP1 (Mason et al. 2021;

XP_029188806.1), the MAC-Perforin protein apextrin (Miller et al. 2007; Hayward et

al. 2011; XP_044181104.1) and the quorum quenching enzyme AmNtNH1 (Mason

et al. 2024; XP_029207486.1). Figure 6 summarises developmental expression data

for these three genes.

This study also provides the most direct evidence to date in support of Hadfield's

hypothesis (Hadfield, 2000) that coral larvae are primed for settlement and

metamorphosis via prior gene expression. Whilst several thousands of genes are up

or down regulated during the onset of competency (81- 133hpf), settlement and the

initiation of metamorphosis largely happen without changes in gene expression

(Figure 3). With the caveat that datasets and technologies were much less

sophisticated than those presently available, Grasso et al. (2011) came to the same

conclusion for *A millepora* based on microarrays, as did Strader et al. (2018)

indirectly by correlating expression data for competent larvae with settlement-

induced larvae from Meyer et al. (2011).

The post-settlement gene expression transition resembles an MZT in terms of

scale and involvement of some specific components

The scale of the post-settlement transition between 3-6hpi resembles that typical of

an MZT. In *Acropora*, we estimate that approximately 20% of mRNAs undergo at

least a two-fold change in expression level (i.e. are either up- or down-regulated over

the 3-6hpi time window). By comparison, in the zebrafish approximately 25% of

maternal transcripts are cleared during the MZT (Mishima and Tomari, 2017) and the

corresponding number for *Caenorhabditis* is around 33% (Baugh et al. 2003). Many

of the mRNAs up-regulated over the 3-6hpi time window in *Acropora* coded for

transcription factors or other regulatory proteins involved in chromatin modifications

and mRNA processing.

Chromatin modification during early development and at the post-settlement

transcriptional transition

Major changes in gene expression patterns also generally involve chromatin

modification events, so histone modification and DNA methylation changes are to be

expected. Homologs of a number of key chromatin modifiers are differentially

expressed at the 3-6hpi transition.

In many animals, homologs of TET (ten-eleven translocation) are the primary agents

responsible for demethylation of 5-methylcytosine residues (MeCpG) in DNA and are

required for gene activation (Mohr et al. 2011; Wu and Zhang 2017; Jessop et al.

2018). As well as a transient earlier (5hpf) window of expression leading into the

early MZT, the *Acropora TET* homolog (XP_029179484.2) was strongly upregulated

at the 6hpi time point (Figure 7A), suggesting a role for the *Acropora* TET homolog in

relieving repression by erasing pre-existing MeCpG marks.

DNA methylation is generally associated with repression of transcription, the

enzymes associated with maintenance and *de novo* DNA methylation being the DNA

methyltransferases DNMT1 and DNMT3, respectively.

In the case of *Acropora*, mRNAs encoding the DNA methylases DNMT1

(XP_029197815.2) and DNMT3a (XP_029213858.1) were maternally provided,

zygotic expression of DNMT1 being detected between 29hpf and 12hpi, despite

having faded by 81hpf. Both DNMTs were upregulated 12 to 18hpi (Figure 7A)

expression of DNMT3a appeared to be stronger than that of DNMT1 (Figure.7A)

mRNA degradation during major transitions in development

Perhaps the major characteristic of MZTs is the elimination of a subset of maternally

provided mRNAs (often a major proportion) to facilitate the transition to the zygotic

developmental program. The mRNA elimination process employs a range of

mechanisms to destabilise or degrade transcripts, but some conserved players have

been identified during major transitions, including homologs of the *Drosophila* genes

Pumilio (Gerber et al. 2006; Rabani et al. 2017) and *Smaug* (Chen et al. 2014) as

well as the *Caenorhabditis* protein TRIM71/LIN41 (Spike et al. 2022).

Transcripts annotated as *Pumilio* (XP_029180703.2 and XP_029180704.2) were

maternally provided but were essentially cleared after 29hpf, whereas a distinct

Pumilio homolog (XP_044164192.1) was not represented in the maternal mRNA

complement but was expressed from 18hpf (Figure 7B). Each of these Pumilio

homologs was up-regulated over the 3-6hpi time window after earlier periods of

activity.

Members of the Smaug/SAMD4 family of proteins are highly conserved (yeast to

man) regulators of mRNA stability and transcriptional repressors that act in

development by binding to specific RNA sequence motifs. Smaug proteins have

been implicated in erasing maternal transcripts during the MZT in the early

development of diverse animals (Chen et al. 2014; Chille et al. 2021) implying that

the Smaug homolog fulfills a similar function in the early development (around 4hpf)

of the coral *Montipora capitata*. Although an early "flutter" of Smaug expression,

possibly corresponding to that reported in *Montipora* (Chille et al. 2021), was

observed in *Acropora* at 5hpf, Smaug (XP_029197445.1) was predominantly

expressed at 15mpi - 6hpi (peak) (Figure 7B). A second transcript annotated as

Smaug (XP_044163536.1) was mildly up-regulated at 13-18hpf.

Members of the TRIM71/LIN41 family of E-3 ubiquitin ligases are highly conserved

RNA-binding proteins (Aviv et al. 2003) that facilitate major developmental transitions

in *Caenorhabditis* (Spike et al. 2022). At least one member of this gene family

(XP_029198843.2/XP_029198844.2) is up-regulated at the 3-6hpi time point during

Acropora development, suggesting that they may also be involved in protein

clearance at the 3-6hpi transition.

Small RNA processing pathway components

Small RNAs (miRNAs, siRNAs and piRNAs) have important regulatory functions

during the development of many bilaterians (Bartel, 2004) and this is likely to also be

the case in *Acropora* given that at least some components of the miRNA processing

pathway are known to be differentially expressed during the development of

Nematostella (Praher et al. 2017). In cnidarians, the biogenesis of miRNAs follows a

plant-like pathway (Moran et al. 2017), but homologs of some of the enzymes in the

animal miRNA processing system are also present and may also participate in

miRNA synthesis.

To address the possibility of miRNAs participating in gene regulation during early

coral development, the expression of components of both the plant and animal

processing pathways was investigated (Figure. 8).

Some of the small RNA processing pathway components were provided maternally,

levels of transcripts annotated as Dicer-like (XP_029197160.2) and the HEN1

methyltransferase (HEN1MT; XP_029185006.2) being present at high levels. In

cnidarians, HEN1 is responsible for methylation of the 3'-ends of both miRNAs and

piRNAs, resulting in their stabilisation. The observed upregulation of HEN1 at 93hpf

(Figure. 8) is consistent with a role in metamorphosis of *Acropora* larvae, as in

Nematostella (Modepalli et al. 2018).

The maternal mRNA repertoire also included an Argonaute homolog

(XP_044181068.1) and a *Acropora* Piwi-like protein (XP_029187700.2/

XP_029187701.2) (Figure. 8) that likely function together in transposon silencing

during early development, as in *Nematostella* (Fridrich et al. 2020; Bamberger et al.

2022). The high abundance of mRNA encoding a homolog of the Maelstrom protein

(XP_044166539.1) also attests to the importance of the piRNA-mediated transposon

silencing pathway (Matsumoto et al. 2015; Sato and Siomi, 2015) during the earliest

stages of coral development.

Although not maternal, other miRNA pathway components were expressed very

early in development (Figure 8). Transcripts annotated as DGCR8/Pasha

(XP_029194207.2) and SRRT/Serrate (XP_044181962.1) were first detected as the

early embryo was going into gastrulation at about 13hpf. For both of these genes,

other transcripts carrying the same annotation appeared to be expressed during

different stages of development. In the case of SRRT/Serrate, XP_044181962.1 is

expressed in the early gastrula but most heavily after the initiation of metamorphosis

(6hpi). A second isoform/splice variant (XP_044181963.1) was most highly

expressed very early in the settlement process (15mpi - 2hpi) – i.e. prior to what Ishii

et al. (2022) referred to as the "point of no return".

Similarly, the DGCR8/Pasha paralogs or splice variants XP_029194207.2 and

XP_029194208.2 seem to have distinct expression profiles during development

(data not shown).

In addition to the maternal Piwi-like mRNA (XP_029187700.2/ XP_029187701.2), a

second (non-maternal) Piwi (XP_044168784.1/ XP_044168785.1/ XP_044168786.1)

was expressed from the early gastrula stage, peaking in expression as the maternal

type was fading.

Discussion

The study of early coral (*A. millepora*) development presented here was based on

gene expression profiles obtained for 26 developmental stages spanning the period

between unfertilised eggs and 3-day-old primary polyps. Downstream analyses via a

DEView app revealed three major transcriptomic transitions - the maternal-zygotic

transition, the acquisition of competence by planulae (93-133hpf) and a post-

settlement transition (3-6hpi) that has not previously received significant attention.

This latter novel transition in gene expression resembled an MZT in terms of scale

and the involvement of some key genes. In addition to regulatory genes involved in

chromatin modification and RNA processing, we focused here on genes likely to

have protective roles, specifically those potentially involved in photoprotection, anti-

microbial activity and transposon silencing.

In many animals, mRNAs encoding canonical histones (including H2A) as well as the

corresponding proteins are provided maternally at sufficient levels to support the

earliest cell division events. To our knowledge, finding that in *Acropora*, an mRNA

encoding one H2A.X variant (H2A.XII) replaced that for canonical H2A in the eggs is

novel. H2A.X is a histone variant whose best-defined role is in DNA repair (reviewed

in Lorkovic and Berger, 2017; Herchenröther et al. 2023). It is closely related to

canonical H2A but distinguished from it by the presence of a characteristic C-

terminal sequence extension that includes a serine residue subject to

phosphorylation generating the gammaH2A.X form which is required for binding at

double-stranded DNA breaks (Osakabe and Molaro, 2023). The enzymes

responsible for this phosphorylation of H2A.X are Phosphoinositide-3-Kinase-related

protein kinases (PIKKs), and at least one PIKK-related protein kinase

(XP_029200091.2) is provided maternally in *Acropora*, though it is unclear whether

H2A.X is a substrate for this activity.

Whilst most coral spawning occurs during the night, the gamete bundles (or eggs)

released are buoyant so that the earliest developmental events occur near the

water's surface. In the present study, *Acropora* larvae became motile 40-45hpf by

which time they would have been exposed to two days of direct sunlight. Early

developmental stages of corals therefore face exposure to UV radiation at levels that

can cause DNA damage, and H2A.X2 mRNA may be maternally provided at high

levels to facilitate maintenance of DNA integrity through the earliest developmental

stages. Chille et al. (2021) noted high representation of GO terms associated with

DNA repair in the maternal mRNA complement of *Montipora* but did not identify

specific genes/proteins. GFP-related proteins are also thought to be involved in DNA

protection in corals (Salih et al. 2000; Palmer et al. 2009; Bou-Abdallah et al. 2006;

Clarke et al. 2024) and, consistent with the study of Strader et al. (2018), at least four

distinct types are expressed during the larval and early polyp stages. Although

mRNAs encoding GFPs did not appear to be maternally provided at significant

levels, the pigmentation of the eggs of many coral species is consistent with the idea

of maternal provision of photoprotective agents that likely include GFP-related

proteins, which are known to be long-lived.

Note that in *Acropora* there is a second H2A.X-like gene (H2A.X1; XP_029200990.1

- annotated as H2A-like) that is expressed only at relatively low levels during early

development. This second H2A.X isoform (XP_029200990.1) has an N-terminal

tyrosine residue replacing the phenylalanine in XP_029204486.1 and has close

homologs in *Hydra* (CDG71330.1; 86.47% protein identity to *Acropora*) and

Hydractinia (now *Klyxum*; AOF43153.1; 87.02% identity) and, like those, is

presumed to be the main isoform expressed during post-larval life. In *Hydractinia*

(now *Klyxum*) *echinata*, the H2A.X variant AOF43152.1 is expressed specifically in

female germ cells along with canonical histone H2A (Torok et al. 2016) and may be

functionally equivalent to the maternally provided *Acropora* H2A.X homolog

XP_029204486.1 (protein sequence identity of 68.15%).

Comparisons with published studies of coral development

The most interesting observation made here was the extent and nature of the post-

settlement transcriptional transition observed at 3-6hpi. This major change in gene

expression is more apparent than in previous studies which either did not include

post-settlement stages (Strader et al. 2018; Chille et al. 2021) or sampled sparsely

pre-settlement (Huffmeyer et al. 2025) or after settlement induction (Ishii et al. 2022,

Meyer et al. 2011) (Table 1). Furthermore, comparisons between the work presented

here and previously published papers in this field (Table 1) are complicated due to

the diversity of species studied, differing temperatures during development, differing

sampling points and differing inducers used to trigger settlement and

metamorphosis.

Development has been studied primarily in members of the three coral genera

Acropora, Montipora and *Porites*. Whilst these are all members of the "superfamily"

Complexa, and the first two belong to the same coral family (Acroporidae), both

Montipora and *Porites* spp. maternally transmit Symbiodiniaceae whereas *Acropora*

spp. acquire their symbionts from the environment during early life.

Chille et al. (2021) focussed on MZT and therefore studied gene expression changes

during development of *Montipora capitata* from unfertilised egg to planula in addition

to adults. Some of their MZT results are consistent with those presented here for

Acropora, for example the expression of Smaug (Figure 7B) and Brachyury

(Supplementary Figure 2) in prawn chip and early gastrula.

However, the rate of development of *M. capitata* observed was apparently slower

than that of *Acropora*. *Montipora capitata* reached the late gastrula stage after 1.5

days, whereas *Acropora* generally reaches the corresponding stage earlier (around

30hpf). This demonstrates the complication of relating developmental stages across

datasets where interspecific differences can be compounded by the

presence/absence of maternal photosymbionts and different rearing temperatures,

as higher temperatures accelerate development. Using GO-term data, Huffmeyer et

al. (2025) recognised three distinct (but overlapping) phases of gene expression

during the early development of *Montipora*.

Since morphology reflects changes in gene expression, we have found that

comparisons of gene expression based on morphology are more meaningful than

those based on time, even in the single species *A millepora*. The post-settlement

transition remained unexplored in the study of Chille et al. (2021) since the focus on

MZT did not require the sampling of post-settlement stages. The 3-6hpi gene

expression transition observed here may correspond to what Ishii et al. (2022) refer

to in their study as "the point of no return" (PNR), a point in post-settlement

development after which reversibility by aborting settlement and returning to the

motile phase is lost. Whilst the Ishii et al. (2022) paper focussed on gene expression

changes between three time points during metamorphosis in *Acropora*, comparison

between their results and those presented here is complicated by the differing

triggers used to drive the process. For *Acropora* spp., external exposure of

competent larvae to the *Hydra* GLWamide (Hym-248), directly triggers

metamorphosis, essentially overriding the need for external settlement cues (Iwao et

al. 2002; Grasso et al. 2011; Attenborough et al. 2019). Whilst Ishii et al. (2022)

induced metamorphosis using Hym-248, the study described here triggered

settlement by addition of CCA extract, which induces searching behaviour rather

than directly driving metamorphosis (Grasso et al. 2011). Hence it is not possible to

directly compare developmental timing between these two studies. The 1hpi and

6hpi time points in the Ishii paper are likely to correspond to later time points in the

present study, as the duration of the settlement period must be several hours longer

when using CCA extract compared to Hym-248. The PNR inferred by Ishii as

occurring at 4hpi may correspond to the 3-6h transition identified in the present work

but note that rates of larval development can be strongly influenced by temperature.

The differences in gene expression between untreated (0 hr) and 1 hour after

induction described in the Ishii et al. paper are therefore likely to encompass a

number of distinct time points in our study, so subtle transcriptional changes

occurring during the searching and very early settlement phases were presumably

not noted by Ishii et al. (2022).

The post-settlement transition – a cnidarian equivalent of the *Caenorhabditis*

"juvenile to adult" transition?

In *Acropora*, the transition from larvae to recruits at 3-6hpi has several parallels to

maternal-zygotic transitions (Vastenhouw et al. 2019) regarding the removal of

earlier transcripts and genome activation appropriate for stages following the

transition. A precedent for such a later transition is provided by *Caenorhabditis*

elegans, where it is recognised as the juvenile to adult transition (Spike et al. 2022).

Above, the involvement of homologs of both *Drosophila* Smaug and Pumilio in the

mRNA clearance processes during MZT and the juvenile to adult transition at 3-6hpi

were highlighted. In addition, two transcripts with the same sequence

(XP_029198843.2 and XP_029198844.2) annotated as a member of the

TRIM71/LIN41 family of RNA-binding proteins were upregulated during the 3-6hpi

transition. Other members of this protein family have important roles in the regulation

of both oocyte to embryo (Spike et al. 2022) and juvenile to adult (Aeschimann et al.

2019) developmental transitions in *Caenorhabditis*. As in *Caenorhabditis* (Spike et al.

2022), this *Acropora* TRIM71/LIN41-related protein is likely to be involved in

regulatory changes at the 3-6hpi transition, and on this basis deserves further

exploration.

In addition, peaks in the expression of the DNA demethylase TET (Figure 7A) and of

SETD2 (XP_029190384.2) as well as homologs of the histone acetyltransferases

HAC5 (XP_029203404.2), KAT5 (XP_029179982.2) and KAT6A (XP_029189526.2)

(Figure 4B) around 3-6hpi imply extensive chromatin modification at that time.

Presumably, the corresponding proteins function in facilitating transcription of genes

required for early polyp development (e.g. He et al. 2024). Transcription factors (TF)

strongly up-regulated at the 3-6hpi time point include genes annotated as homologs

of FoxB1 (XP029196672.1) and FoxO (XP_029208531.2), Aristaless

(XP_029193170.1), Rax (XP_029193171.1), Meis2 (XP_029187327.1) and Otx

(XP_029190339.1), whilst homologs of Emx (XP_029211983.2), Pitx2

(XP_029179676.1) and Brn3C (XP_029213135.1) were simultaneously down-

regulated.

Limitations: Heatmaps are useful but tell only part of the story

A caveat in making inferences from heatmaps across developmental stages, and

more generally from whole-larvae overall expression levels, is that important subtle

changes may be missed. Particularly with transcription factors or other regulatory

molecules, low overall levels of expression may have major effects, especially when

restricted to subsets of cells. Consequently, interpreting developmental gene

expression requires not only temporal data, but also spatial information from the

application of in situ hybridisation or other techniques. This is clearly illustrated in

Supplementary Figure 2, which provides expression data for genes implicated in

conserved functions during gastrulation.

Whilst in situ hybridisation data show expression of these four genes in smaller

subsets of genes across all developmental stages between gastrulation and

metamorphosed recruit, the heatmaps show that the expression maxima occur post-

settlement rather than at gastrulation around 13 – 18hpf. Only a weaker peak in

expression of Brachyury at 13hpf corresponds to the early gastrulation window of

expression visualised via in situ hybridisation (Hayward et al. 2015). Consequently,

visualisations of spatial expression via in situ hybridisation are required to study the

function of selected genes. For example, whilst gene expression data presented in

the heatmap suggested functions post-settlement, ISH visualisations revealed that

temporal expression of an *Acropora* arx-related gene

(XP_044163657.1/XP_029193169.1) during gastrulation closely complements that of

snail (Figure 5). This *Acropora* sequence is referred to here as homeobrain, based

on its orthology with *Nematostella* XP_048583705.1 (Mazza et al. 2010). Snail and

homeobrain respectively demarcate the early mesendoderm and presumptive

ectoderm during gastrulation (Figure 5), but both are also expressed at higher levels

after settlement-induction where snail expression reaches its expression peak earlier

than in the case of homeobrain. Assuming these two genes fulfill similar function

during gastrulation and post-settlement, this suggests that a cell migration such as in

an epithelial-mesenchymal transition occurs during the initiation of metamorphosis.

The upper limit of expression in a heatmap is determined by the genes included in

that map so important changes in gene expression may not be as apparent as if the

gene was plotted against itself.

Expression of the small RNA processing machinery during early development

Despite deep evolutionary divergence between the sea anemones (Actiniaria) and

Despite deep evolutionary divergence between the sea anemones (Actiniaria) and

the coral/corallimorpharian lineage, many of the molecular processes underlying

early development in *Acropora* closely parallel those described in *Nematostella*

(Hayward et al. 2015), and this appears to extend to the machinery for processing

small RNAs (miRNAs and piRNAs). However, some differences were observed in

the timing of expression of several of genes implicated in small RNA processing.

Whilst mRNAs for Vasa1, Maelstrom and Piwi are provided maternally in *Acropora*

and *Nematostella*, Vasa2 (aka Vasa-like or DDX4) is also maternal in the coral

whereas the corresponding mRNA was not detected until around gastrulation in

Nematostella (Praher et al. 2017). Whether Dicer/DCL is maternal is unclear in the

case of *Nematostella*, whereas mRNA of the *Acropora* homolog of *Nematostella*

Dicer 2 (AGW15596.1), encoding XP_029197160.2, is abundant in *Acropora* eggs.

In summary, the preliminary data presented here indicate that all or most of the small

RNA processing machinery is either maternal or expressed very early in *Acropora*

development. It is therefore likely that, in addition to roles for some of these

components in transposon silencing, miRNAs are likely to have important regulatory

roles in the early development of *Acropora*.

Heatmaps enable identification of candidate genes for key roles in early development

721 A major strength of the DEView app is that it provides a means of scanning classes

722 of genes individually or collectively for their expression patterns across early coral

723 development, allowing identification of candidates for roles in specific processes,

724 including settlement induction and the initiation of calcification. Whilst significant

Expression of the small RNA processing machinery during early development

Despite deep evolutionary divergence between the sea anemones (Actiniaria) and

the coral/corallimorpharian lineage, many of the molecular processes underlying

early development in *Acropora* closely parallel those described in *Nematostella*

(Hayward et al. 2015), and this appears to extend to the machinery for processing

small RNAs (miRNAs and piRNAs). However, some differences were observed in

the timing of expression of several of genes implicated in small RNA processing.

Whilst mRNAs for Vasa1, Maelstrom and Piwi are provided maternally in *Acropora*

and *Nematostella*, Vasa2 (aka Vasa-like or DDX4) is also maternal in the coral

whereas the corresponding mRNA was not detected until around gastrulation in

Nematostella (Praher et al. 2017). Whether Dicer/DCL is maternal is unclear in the

case of *Nematostella*, whereas mRNA of the *Acropora* homolog of *Nematostella*

Dicer 2 (AGW15596.1), encoding XP_029197160.2, is abundant in *Acropora* eggs.

In summary, the preliminary data presented here indicate that all or most of the small

RNA processing machinery is either maternal or expressed very early in *Acropora*

development. It is therefore likely that, in addition to roles for some of these

components in transposon silencing, miRNAs are likely to have important regulatory

roles in the early development of *Acropora*.

Heatmaps enable identification of candidate genes for key roles in early development

A major strength of the DEView app is that it provides a means of scanning classes of genes individually or collectively for their expression patterns across early coral development, allowing identification of candidates for roles in specific processes, including settlement induction and the initiation of calcification. Whilst significant

725 progress has recently been made, we have as yet a relatively limited understanding

726 of the nature and recognition of settlement cues (Hadfield, 2011; Tran and Hadfield,

727 2012; Ball et al. 2021; Turnlund et al. 2025). The larvae of many corals select

728 appropriate settlement sites by detection of small molecules that either favour or

729 discourage settlement (e.g. Morse et al. 1988; Negri et al. 2001; Kitamura et al.

730 2007; Birrell et al. 2008; Kitamura et al. 2009; Tebben et al. 2015, Quinlan et al.

731 2023). Most likely the outcome reflects a combination of molecules acting as ligands

732 for specific receptors that are expressed in competent larvae and downregulated

733 after settlement and metamorphosis. Receptors that potentially could be involved in

734 settlement cue reception and downstream signalling pathways, respectively, are G-

735 Protein coupled receptors (GPCRs) and nuclear receptors. With the help of the

736 DEView app, search terms such as "G-protein coupled receptor" or specific Pfam

737 domains can be used as filters to search the developmental transcriptome to identify

738 genes that were upregulated at competency (i.e. 93hpf) consistent with their

739 potential involvement in settlement cue perception. Searching the *Acropora* data

740 using the terms PF00104 Ligand-binding domain of nuclear hormone receptor and

741 PF00105 Zinc finger, C4 type (two domains) yielded 21 hits corresponding to 14 loci

742 (Supplementary Figure 3). Nuclear receptors play key roles in metamorphosis in

743 both insects and some vertebrates, potentially acting in the internal transmission of

744 metamorphosis signals rather than their external reception. Of the 14, seven were

745 upregulated at 133hpf and all of these mRNAs appear to be cleared at the 3-6hpi

746 transition so could have roles in the cue recognition pathway and/ or metamorphosis.

748 The *Acropora* developmental transcriptome data, together with the DEView app,

749 have provided novel insights into early coral development and allowed identification

of candidate genes for key roles. We hope that the availability of this app via the web

will lead to more extensive exploration of the data and ultimately facilitate much

better understanding of the molecular processes underlying the development of the

coral body plan.

For a range of organisms, including some corals, considerably more advanced

genomic and transcriptomic resources are now available than could have been

imagined even a few years ago. However, despite the developmental transcriptomic

data provided here being to our knowledge the most comprehensive available for

any coral, users need to be aware that the quality of the *Acropora millepora* dataset

falls short of those available for the model invertebrates *Drosophila* and

Caenorhabditis. Nevertheless, the developmental novelties uncovered here provide

clear indications of how useful the dataset can be in facilitating coral developmental

research.

<u>Acknowledgements</u>

The authors gratefully acknowledge the support of the Australian Research

Council under Discovery Grant DP170104734, SuperScience Grant (round 2)

FS110200046 and Centres of Excellence Grant CE14100020. We also thank

Gergely Torda and the staff of the Orpheus Island Research Centre for fieldwork

assistance. Some of the in situ images presented here were produced by Lauretta

Grasso and Nikki Hislop.

References

Aeschimann F, Neagu A, Rausch M, Großhans H. *let-7* coordinates the transition to adulthood through a single primary and four secondary targets. Life Sci Alliance. 2019; 2(2):e201900335.

Attenborough RMF, Hayward DC, Wiedemann U, Foret S, Miller DJ, Ball EE. Expression of the neuropeptides RFamide and LWamide during development of the coral Acropora millepora in relation to settlement and metamorphosis. Dev Biol. 2019; 446: 56-67.

Aviv T, Lin Z, Lau S, Rendl LM, Sicheri F, Smibert CA. The RNA-binding SAM domain of Smaug defines a new family of post-transcriptional regulators. Nat Struct Biol. 2003;10(8):614-21.

Ayers TN, Nicotra ML, Lee MT. Parallels and contrasts between the cnidarian and bilaterian maternal-to-zygotic transition are revealed in Hydractinia embryos. PLoS Genet. 2023;19(7):e1010845.

Ball EE, Hayward DC, Bridge TCL, Miller DJ. *Acropora* – the most studied coral genus. In: Boutet A, Schierwater B (eds). Handbook of Marine Model Organisms in Experimental Biology - Established and Emerging. 2021. Boca Raton (FL): CRC Press. p. 173-193. DOI: 10.1201/9781003217503-10.

Ball E, Hayward D, Reece-Hoyes J, Hislop N, Samuel G, Saint R, Harrison P and Miller, D. Coral development: from classical embryology to molecular control. Int J Dev Biol. 2002;46:671-678.

Ball EE, Hayward DC, Saint R, Miller, DJ. A simple plan—cnidarians and the origins of developmental mechanisms. Nat Rev Genetics. 2004;*5*(8):567-577.

Bamberger C, Pankow S, Yates JR 3rd. Nvp63 and nvPIWIL1 suppress retrotransposon activation in the sea anemone Nematostella vectensis. J Proteome Res. 2022; 21(11):2586-2595.

Bartel DP. MicroRNAs: genomics, biogenesis, mechanism, and function. Cell. 2022; 116(2):281-97.

Baugh LR, Hill AA, Slonim DK, Brown EL, Hunter CP. Composition and dynamics of the Caenorhabditis elegans early embryonic transcriptome. Development. 2003; 130(5):889-900.

820 Birrell CL, Mccook LJ, Willis BL, Harrington L. Chemical effects of macroalgae on
821 larval settlement of the broadcast spawning coral Acropora millepora. Mar Ecol Prog
822 Ser. 2008;362:129–137.
823
824 Bou-Abdullah F, Chasteen ND, Lesser MP. Quenching of superoxide radicals by
825 green fluorescent protein. Biochim Biophys Acta. 2006;1760(11):1690-5.
826
827 Breitwieser FP, Baker DN, Salzberg SL. KrakenUniq: confident and fast
828 metagenomics classification using unique k-mer counts. Genome Biol. 2018;19:198.
829
830 Chen L, Dumelie JG, Li X, Cheng MH, Yang Z, Laver JD, Siddiqui NU, Westwood
831 JT, Morris Q, Lipshitz HD, Smibert CA. Global regulation of mRNA translation and
832 stability in the early Drosophila embryo by the Smaug RNA-binding protein. Genome
833 Biol. 2014;15(1):R4.
834
835 Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor.
836 Bioinformatics. 2018;34:i884-i890.
837
838 Chille E, Strand E, Neder M, Schmidt V, Sherman M, Mass T, Putnam H.
839 Developmental series of gene expression clarifies maternal mRNA provisioning and
840 maternal-to-zygotic transition in a reef-building coral. BMC Genomics. 2021;
841 22(1):815.
842
843 Clarke DN, Rose NH, De Meulenaere E, Rosental B, Pearse JS, Pearse VB, Deheyn
844 DD Fluorescent proteins generate a genetic color polymorphism and counteract
845 oxidative stress in intertidal sea anemones. Proc Natl Acad Sci USA. 2024;
846 121(11):e2317017121.
847
848 Di Tommaso P, Chatzou M, Floden EW, Barja PP, Palumbo E, Notredame C.
849 Nextflow enables reproducible computational workflows. Nat Biotechnol. 2017;
850 35:316-319.
851
852 Fridrich A, Modepalli V, Lewandowska M, Aharoni R, Moran Y.) Unravelling the
853 developmental and functional significance of an ancient Argonaute duplication. Nat
854 Commun. 2020;11(1):6187.
855
856 Gerber AP, Luschnig S, Krasnow MA, Brown PO, Herschlag D. Genome-wide
857 identification of mRNAs associated with the translational regulator PUMILIO in
858 Drosophila melanogaster. Proc Natl Acad Sci USA. 2006;103(12):4487-92.
859
860 Gilbert E, Craggs J, Modepalli V. Gene Regulatory Network that Shaped the
861 Evolution of Larval Apical Organ in Cnidaria. Mol Biol Evol. 2024;41: msad285.
862

Grasso L, Maindonald J, Rudd S, Hayward D, Saint R, Miller DJ, Ball EE. Microarray
analysis provides candidate genes for key roles in coral development. BMC
Genomics. 2008;9:540.

Grasso LC, Negri AP, Foret S, Hayward DC, Miller DJ, Ball EE. The biology of coral
metamorphosis: molecular responses of coral larvae to inducers of settlement and
metamorphosis. Dev Biol. 2011;353:411-419.

Gu Z, Hübschmann D. Make Interactive Complex Heatmaps in R. Bioinformatics.
2022;38:1460–1462. https://doi.org/10.1093/bioinformatics/btab806.

Hadfield MG. Biofilms and Marine Invertebrate Larvae: What Bacteria Produce That
Larvae Use to Choose Settlement Sites. Ann Rev Marine Sci. 2011;3:453-470.

Hadfield MG. Why and how marine-invertebrate larvae metamorphose so fast. Sem
Cell Dev Biol. 2000;11(6):437-443.

Hayward DC, Miller DJ, Ball EE. Snail expression during embryonic development of
the coral *Acropora*: Blurring the diploblast / triploblast divide? Development Genes
and Evolution. 2004;214:257-260.

Hayward DC, Samuel G, Pontynen PC, Catmull J, Saint RB, Miller DJ, Ball EE.
Localized expression of a DPP/BMP2-4 ortholog in a coral embryo. Proc Natl Acad
Sci USA. 2002;99:8106-8111.

Hayward DC, Hetherington S, Behm CA, Grasso L, Foret S, Miller DJ, Ball EE.
Differential gene expression at coral settlement and metamorphosis – a subtractive
hybridization study. PLoS One. 2011;6: e26411.

Hayward DC, Grasso LC, Saint RB, Miller DJ, Ball EE. The organizer in evolution:
Gastrulation and organizer gene expression highlight the importance of Brachyury
during development of the coral *Acropora millepora*. Dev Biol. 2015;399:227-247.

He S, Rangel-Huerta E, Hill E, Ellington L, Chen S, Robb S, Majerová E, Drury C,
Gibson MC. An evolutionarily conserved Hox-Gbx segmentation code in the rice
coral, *Montipora capitata*. bioRxiv preprint. 2024; doi:
https://doi.org/10.1101/2024.09.29.615694.

Herchenröther A, Wunderlich TM, Lan J, Hake SB. Spotlight on histone H2A
variants: From B to X to Z. Sem Cell Dev Biol. 2023;135:3-12.

Huffmyer AS, Wong KH, Becker DM, Strand E, Mass T, Putnam HM. Shifts and
critical periods in coral metabolism reveal energetic vulnerability during development.
Curr Biol. 2025;35:1-14;doi https://doi.org/10.1016/j.cub.2025.05.013.

Ishii Y, Hatta M, Deguchi R, Kawata M, Maruyama S. Gene expression alterations
from reversible to irreversible stages during coral metamorphosis. Zool Lett. 2022;
8(1):4.

Iwao K, Fujisawa T, Hatta M. A cnidarian neuropeptide of the GLWamide family
induces metamorphosis of reef-building corals in the genus Acropora. Coral Reefs.
2002;21:127-129.

Jessop P, Ruzov A, Gering M. Developmental Functions of the Dynamic DNA
Methylome and Hydroxymethylome in the Mouse and Zebrafish: Similarities and
Differences. Front Cell Dev Biol. 2018;20(6):27.

Jones P, Binns D, Chang HY, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J,
Mitchell A, Nuka G, Pesseat S. InterProScan 5: genome-scale protein function
classification. Bioinformatics. 2014;30(9):1236-40.

Kitamura M, Koyama T, Nakano Y, Uemura D. Characterization of a natural inducer
of coral larval metamorphosis. J Exp Mar Biol Ecol. 2007;340:96-102.

Kitamura M, Schupp PJ, Nakano Y, Uemura D. Luminaolide, a novel
metamorphosis-enhancing macrodiolide for scleractinian coral larvae from crustose
coralline algae. Tetrahedron Letts. 2009;50:6606-6609.

Langfelder P, Horvath S. WGCNA: an R package for weighted correlation network
analysis. BMC Bioinformatics. 2008;9:559. https://doi.org/10.1186/1471-2105-9-559.

Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods.
2012;9:357-359.

Levy S, Elek A, Grau-Bové X, Menéndez-Bravo S, Iglesias M, Tanay A, Mass T,
Sebé-Pedrós A. A stony coral cell atlas illuminates the molecular and cellular basis
of coral symbiosis, calcification, and immunity. Cell. 2021;184(11):2973-2987.

Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with
or without a reference genome. BMC Bioinformatics. 2011;12:323.

Lorković ZJ, Berger F. Heterochromatin and DNA damage repair: Use different
histone variants and relax. Nucleus. 2017;8(6):583-588.

Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. Genome Biol. 2014;15:550.

Mansour TA, Rosenthal JJ, Brown CT, Roberson LM. Transcriptome of the
Caribbean stony coral Porites astreoides from three developmental stages.
Gigascience. 2016;5(1):33.
953
954 Mason BM, Cohen JH. Long-wavelength photosensitivity in coral planula larvae. Biol
955 Bull. 2012;222(2):88-92.
956
957 Mason B, Cooke I, Moya A, Augustin R, Lin M-F, Satoh N, Bosch TCG, Bourne DG,
958 Hayward DC, Andrade N, Forêt S, Ying H, Ball EE, Miller DJ. AmAMP1 from
959 *Acropora millepora* and damicornin define a family of coral-specific antimicrobial
960 peptides related to the Shk toxins of sea anemones. Dev Comp Immunol. 2021;114:
961 103866.
962
963 Mason B, Hayward DC, Moya A, Cooke I, Sorenson A, Brunner R, Andrade N,
964 Huerlimann R, Bourne DG, Schaeffer P, Grinblat M, Ravasi T, Ueda N, Tang SL,
965 Ball EE, Miller DJ. Microbiome manipulation by corals and other Cnidaria via quorum
966 quenching. Curr Biol. 2024;34(14):3226-3232.e5.
967
968 Mason BM, Koyanagi M, Sugihara T, Iwasaki M, Slepak V, Miller DJ, Sakai Y,
969 Terakita A. Multiple opsins in a reef-building coral, *Acropora millepora*. Scientific
970 Reports 2023;23:1628.
971
972 Matsumoto N, Sato K, Nishimasu H, Namba Y, Miyakubi K, Dohmae N, Ishitani R,
973 Siomi H, Siomi MC, Nureki O. Crystal structure and activity of the endoribonuclease
974 domain of the piRNA pathway factor Maelstrom. Cell Reps. 2015;11:366-375.
975
976 Mazza ME, Pang K, Reitzel AM, Martindale MQ, Finnerty JR. A conserved cluster of
977 three PRD-class homeobox genes (homeobrain, rx and orthopedia) in the Cnidaria
978 and Protostomia. EvoDevo. 2010;1:1-15.
979
980 Meyer E, Aglyamova GV, Matz MV. Profiling gene expression responses of coral
981 larvae (Acropora millepora) to elevated temperature and settlement inducers using a
982 novel RNA-Seq procedure. Mol Ecol. 2011;20:3599-616.
983
984 Meyer E, Aglyamova GV, Wang S, Buchanan-Carter J, Abrego D, Colbourne JK,
985 Willis BL, Matz MV. Sequencing and de novo analysis of a coral larval transcriptome
986 using 454 GSFlx. BMC Genomics. 2009;10:1-8.
987
988 Michail C, Rodrigues Lima F, Viguier M, Deshayes F. Structure and function of the
989 lysine methyltransferase SETD2 in cancer: From histones to cytoskeleton.
990 Neoplasia. 2025;59:101090.
991

992 Miller DJ, Hemmrich G, Ball EE, Hayward DC, Khalturin K, Funayama N, Agata K,
993 Bosch TCG.The innate immune repertoire in Cnidaria – ancestral complexity and
994 stochastic gene loss. Genome Biol. 2007;8:R59.
995
996 Mishima Y, Tomari Y. Pervasive yet nonuniform contributions of Dcp2 and Cnot7 to
997 maternal mRNA clearance in zebrafish. Genes Cells. 2017;22(7):670-678.
998
999 Modepalli V, Fridrich A, Agron M, Moran Y. The methyltransferase HEN1 is required
1000 in Nematostella vectensis for microRNA and piRNA stability as well as larval
1001 metamorphosis. PLoS Genet. 2018;14(8):e1007590.
1002
1003 Mohr F, Döhner K, Buske C, Rawat VP. TET genes: new players in DNA
1004 demethylation and important determinants for stemness. Exp Hematol. 2011;
1005 39(3):272-81.
1006
1007 Moran Y, Agron M, Praher D, Technau U. The evolutionary origin of plant and animal
1008 microRNAs. Nat Ecol Evol. 2017;1(3):27.
1009
1010 Morse DE, Hooker N, Morse AN, Jensen RA. Control of larval metamorphosis and
1011 recruitment in sympatric agariciid corals. J Exp Mar Biol Ecol. 1988;116(3):193-217.
1012
1013 Negri AP, Webster NS, Hill RT, Heyward AJ. Metamorphosis of broadcast spawning
1014 corals in response to bacteria isolated from crustose algae. Mar Ecol Prog Ser.
1015 2001;*223*:121-131.
1016
1017 Okubo N, Hayward DC, Forêt S, Ball EE. A comparative view of early development
1018 in the corals *Favia lizardensis*, *Ctenactis echinata*, and *Acropora millepora* -
1019 morphology, transcriptome, and developmental gene expression. BMC Evol Biol.
1020 2016;16:1-12.
1021
1022 Okubo N, Mezaki T, Nozawa Y, Nakano Y, Lien YT, Fukami H, Hayward DC, Ball EE
1023 Comparative embryology of eleven species of stony corals (Scleractinia). PLoS One.
1024 2013;8(12): e84115.
1025
1026 Osakabe A, Molaro A Histone renegades: Unusual H2A histone variants in plants
1027 and animals. Sem Cell Dev Biol. 2023;135:35-42.
1028
1029 Palmer CV, Modi CK, Mydlarz LD. Coral fluorescent proteins as antioxidants. PLoS
1030 One. 2009;4(10):e7298.
1031
1032 Praher D, Zimmermann B, Genikhovich G, Columbus-Shenkar Y, Modepalli V,
1033 Aharoni R, Moran Y, Technau U. Characterization of the piRNA pathway during
1034 development of the sea anemone Nematostella vectensis. RNA Biol. 2017;
1035 14(12):1727-1741.

1036

Quinlan ZA, Bennett MJ, Arts MGI, Levenstein M, Flores D, Tholen HM, Tichy L,
Juarez G, Haas AF, Chamberland VF, Latijnhouwers KRW, Vermeij MJA, Johnson
AW, Marhaver KL, Kelly LW. Coral larval settlement induction using tissue-
associated and exuded coralline algae metabolites and the identification of putative
chemical cues. Proc Roy Soc B. 2023;290:20231476.
https://doi.org/10.1098/rspb.2023.1476.

1043

1044 Rabani M, Pieper L, Chew G-L, Schier AH. A Massively Parallel Reporter Assay of 3'
1045 UTR Sequences Identifies In Vivo Rules for mRNA Degradation. Mol Cell. 2017;
1046 68(6):1083-1094.e5.

1047

1048 Reyes-Bermudez A, Desalvo MK, Voolstra CR, Sunagawa S, Szmant AM, Iglesias-
1049 Prieto R, Medina M. Gene expression microarray analysis encompassing
1050 metamorphosis and the onset of calcification in the scleractinian coral Montastraea
1051 faveolata. Mar Genomics 2009;2:149-159.

1052

1053 Ramon-Mateu J, Ferraioli A, Teixidó N, Domart-Coulon I, Houliston E, Copley RR.
1054 Aboral cell types of Clytia and coral larvae have shared features and link taurine to
1055 the regulation of settlement. Science Adv. 2025;16;11(20):eadv1159.

1056

1057 Reyes-Bermudez A, Villar-Briones A, Ramirez-Portilla C, Hidaka M, Mikheyev AS
1058 Developmental Progression in the Coral *Acropora digitifera* Is Controlled by
1059 Differential Expression of Distinct Regulatory Gene Networks. Genome Biol Evol.
1060 2016;8(3):851-70.

1061

1062 Salih A, Larkum A, Cox G, Kuhl M, Hoegh-Guldberg Fluorescent pigments in corals
1063 are photoprotective. Nature.2000;408:850-853.

1064

1065 Shoguchi E, Beedessee G, Hisata K, Tada I, Narisoko H, Satoh N, Kawachi M,
1066 Shinzato C. A New Dinoflagellate Genome Illuminates a Conserved Gene Cluster
1067 Involved in Sunscreen Biosynthesis. Genome Biol Evol. 2021;13:evaa235.

1068

1069 Sato K, Siomi MC. Functional and structural insights into the piRNA factor
1070 Maelstrom. FEBS Lett. 2015;589:1688-1693.

1071

1072 Siboni N, Abrego D, Puill-Stephan E, King WL, Bourne DG, Raina J-B, Seymour JR,
1073 Harder T. Crustose coralline algae that promote coral larval settlement harbor
1074 distinct surface bacterial communities. Coral Reefs 2020;39:1703–1713.

1075

1076 Spike CA, Tsukamoto T, Greenstein D. Ubiquitin ligases and a processive
1077 proteasome facilitate protein clearance during the oocyte-to-embryo transition in
1078 *Caenorhabditis elegans*. Genetics. 2022;221(1):iyac051.

1079

Strader ME, Davies SW, Matz MV (2015) Differential responses of coral larvae to the
colour of ambient light guide them to suitable settlement microhabitat. R Soc Open
Sci 2(10):150358

Strader ME, Aglyamova GV, Matz MV. Molecular characterization of larval
development from fertilization to metamorphosis in a reef-building coral. BMC
Genomics. 2018;19(1):1-17.

Takeuchi T, Yamada L, Shinzato C, Sawada H, Satoh N. Stepwise Evolution of
Coral Biomineralization Revealed with Genome-Wide Proteomics and
Transcriptomics. PLoS One 2016;11(6):e0156424.

Tebben J, Motti CA, Siboni N, Tapiolas DM, Negri AP, Schupp PJ, Kitamura M,
Hatta M, Steinberg PD, Harder T. Chemical mediation of coral larval settlement by
crustose coralline algae. Sci Rep. 2015;5:10803.

Technau U. Gastrulation and germ layer formation in the sea anemone Nematostella
vectensis and other cnidarians. Mech Dev. 2020;163:103628.

Török A, Schiffer PH, Schnitzler CE, Ford K, Mullikin JC, Baxevanis AD, Bacic A,
Frank U, Gornik SG. The cnidarian Hydractinia echinata employs canonical and
highly adapted histones to pack its DNA. Epigenetics Chromatin. 2016;9(1):1-17.

Tran C, Hadfield MG. Are G-protein-coupled receptors involved in mediating larval
settlement and metamorphosis of coral planulae? Biol Bull 2012;222(2):128-36.

Turnlund AC, Vanwonterghem I, Botté ES, Randall CJ, Giuliano C, Kam L, Bell S,
O'Brien P, Negri AP, Webster NS, Lurgi M. Linking differences in microbial network
structure with changes in coral larval settlement. ISME Comm. 2023;3:114.

Vastenhouw NL, Cao WX, Lipshitz HD. The maternal-to-zygotic transition revisited.
Development. 2019;146(11): dev161471

Walker NS, Fernández R, Sneed JM, Paul VJ, Giribet G, Combosch DJ. Differential
gene expression during substrate probing in larvae of the Caribbean coral *Porites
astreoides*. Mol Ecol. 2019;28(22):4899-4913.

Wickham H. Data Analysis, in: Wickham, H. (Ed.), Ggplot2: Elegant Graphics for
Data Analysis. Springer International Publishing, Cham, 2016; pp. 189–201.
https://doi.org/10.1007/978-3-319-24277-4_9.

Wu X, Zhang Y. TET-mediated active DNA demethylation: mechanism, function and
beyond. Nat Rev Genet. 2017;18(9):517-534.

1123

1124 Yasuoka Y, Shinzato C, Satoh N. The Mesoderm-Forming Gene brachyury
1125 Regulates Ectoderm-Endoderm Demarcation in the Coral Acropora digitifera. Curr
1126 Biol. 2016;26(21):2885-2892.

1127

1128

1129 **Availability of data and materials**

1130

Data accessibility statement: To facilitate use and rapid exploration of the

Acropora millepora developmental transcriptome we created an interactive web

application that is hosted on servers provided by the Australian Research Data

Commons Nectar Research Cloud to ensure that it is scalable and highly available

for long-term use. Users can access the web application directly at

https://amil2dev.mmb.group/ or obtain its source code and the underlying data from

our github repository (https://github.com/iracooke/amil2_dev).

1138

1139

1140 **Author Contributions**

1141

Designed research: AM, DH, BL, EB, DM

Carried out field work: AM, DH, BL, EB

Laboratory work: AM, BL

Performed bioinformatic analyses: SF, AM, DH, IC

Contributed new analytical tools: IC

Analysed the data: AM, RB, MG, DH, DM

Wrote the paper: DH, RB, EB, IC, DM (lead)

1149

1150

1151 Figure legends

1152

Figure 1. Images showing the morphologies recorded at the time of collection at the

timepoints indicated. Times are given in hours post fertilisation (hpf) and hours post

induction (hpi). The places in the developmental sequence of two major events,

gastrulation and induction with a settlement cue, are indicated. Some of the SEM

images have been previously published in Grasso et al. 2008, Hayward et al. 2011,

and Ball et al. 2021.

1159

Figure 2. PCA analysis of the developmental transcriptomic data. Principal

Components Analysis (PCA) of the developmental transcriptome data with ellipses

representing 95% confidence levels. At 93 hours post fertilisation (hpf), larvae were

competent to settle in response to ethanolic CCA extract and settlement was

induced at the age of 133hpf. Settlement induction caused larvae to elongate at 15

minutes post induction (mpi) and 100% of larvae had initiated metamorphosis at 3

hours post induction (hpi). A major change in gene expression occurred between 3

and 6hpi.

1168

1169 Figure 3. Serial up-down expression plots showing the number of genes changing in

1170 expression between consecutive time points during development. There are three

1171 periods that show major changes in expression: early development leading up to

1172 gastrulation (5-29hpf), 81-133hpf as competence to settle is developing, and 3-6hpi

1173 as polyp development is getting under way after settlement.

1174

Figure 4. Expression of histone genes and histone modifying enzymes during the

early development of *Acropora millepora*. (A) Heatmap showing expression of

selected histone genes during development. mRNAs encoding canonical histones

H3, H4 and H2B are present at high levels in eggs (rows 3, 4 and 5 from the top), as

are H2A.X and H2A.V (but not canonical H2A) messages. Zygotic expression of the

four core canonical histones begins at the start of gastrulation (i.e. the 13hpf time

point) note that the set of histone genes shown in the bottom four rows are likely to

represent a replication-dependent subset that may be required at high levels only

early in development. Consistent with this, the top row shows data for another H2A

gene, expression of which begins later in development. (B) Heatmap showing

expression of histone modifying enzymes during the early development of *Acropora*.

mRNAs encoding the histone acetyltransferase KAT6A the histone H3K36

methylase SETD2 are maternal (rows 2 and 3, respectively). Two other histone

acetyltransferases (HAC5 and KAT5; top and bottom rows) are most highly

expressed at the 6hpi time point, reflecting their possible role in activating chromatin

during the post-settlement transition. Note that a second wave of expression of both

KAT6A and SETD2 also occurs at that time.

1192

Figure 5. Heatmaps and in situ hybridisation provide complementary data, as

illustrated by data for coral homologs of the *Drosophila* genes, snail and homeobrain.

During gastrulation, these genes have complementary morphological expression

patterns, *Am-snail* marking the presumptive mesendoderm (Hayward et al. 2004)

and *Am-homeobrain* the presumptive ectoderm. (A) Diagrammatic vertical sections

of embryos aged from early prawn chip to sphere. (B-D) Embryos aged from prawn

chip to sphere (blastopore closure) viewed in three different ways. (B) Critical-point

dried embryos viewed with scanning electron microscopy reveal the external

morphology, (C) Whole mount in situ hybridisation reveals the spatial expression of

Am-snail. (D) Vertical sections of in situ preparations reveal the internal distribution

of the expressing tissue in the developing mesendoderm. (E) Expression of *Am-*

homeobrain in the developing ectoderm revealed using whole-mount in situ

hybridisation. Note that the two genes are complimentary in their expression. (C', E')

Heatmap showing the relative expression of the two corresponding genes over the

course of development. Parts of the figure were published in Hayward et al. (2004)

Development Genes & Evolution 214:257-260, Springer Nature.

1209

Figure 6. In situ hybridisation and heatmaps provide complementary data in support

of roles for known and suspected anti-bacterial activities during settlement and

metamorphosis in *Acropora*. The upper three rows show in situ hybridisation data for

developmental stages leading into and immediately following settlement for (A) the

anti-microbial protein AmAMP1 (Mason et al. (2021), (B) the quorum quenching

activity AmNtNH1 (Mason et al. 2024), and (C) the MAC-Perforin protein Apextrin

Miller et al. 2007; Hayward et al. 2011). A', B' and C' are heatmaps showing the

corresponding temporal expression levels of the three genes. The arrowheads

indicate the time of settlement induction.

1219

Figure 7. Heatmaps for genes implicated in (A) DNA methylation /demethylation and

(B) clearance of mRNAs and proteins are consistent with major changes in gene

activity occurring at 3-6hpi. The upper panel (A) illustrates relative expression levels

of the *Acropora* homologs of ten-eleven-translocase (tet), the primary role of which is

removal of CpG methylation marks, the de novo DNA methyltransferase, DNMT3,

and the maintenance DNA methyltransferase, DNMT1. Although all three activities

are expressed during gastrulation, TET is most heavily expressed at the 6hpi time

point. The lower panel (B) shows relative expression levels of *Acropora* homologs of

the conserved mRNA clearance activities, Smaug and Pumilio, and the TRIM-NHL

ubiquitin ligase, TRIM71/LIN41. Smaug and TRIM71/LIN41are most heavily

expressed at the 6hpi time point; three gene predictions annotated as Pumilio

homologs (corresponding to at least two distinct gene products) were also up-

regulated at that time.

1233

Figure 8. Expression of genes involved in small RNA (miRNA and piRNA)

processing during early development. mRNAs encoding HEN1 and Dicer homologs

(rows 6 and 7) were present at high levels in eggs, as were those encoding

homologs of Maelstrom, Vasa/DDX4.1 and .2, and Piwi2, which function primarily in

piRNA-mediated transposon silencing. Other miRNA processing activities, including

homologs of Serrate/SRRT and DGCR8/Pasha, were also expressed in embryos,

larvae and during post settlement life, implying extensive involvement of miRNAs in

regulating gene activities during early coral development.

1242

Supplementary Figure 1.

Weighted gene co-expression network analysis (WGCNA) clustered the

developmental gene expression data into 39 modules based on correlated

expression patterns. The number of genes in each module range from 18 (plum

module) to 5851 (turquoise module) and the grey module contains all genes that did

not fit into any other module. The WGCNA heatmap indicated that there is a major

change in gene expression between 3 and 6 hours post induction (hpi) and also

showed that settlement induction at the age of 133 hours post fertilisation (hpf) did

not trigger an immediate strong change in gene expression 15 minutes post

induction (mpi).

1253

Supplementary Figure 2.

Genes previously characterised in the context of gastrulation illustrate the need for

spatial as well as temporal expression data. In situ data for the *Acropora* homologs

of several genes imply conservation of function during gastrulation with Bilateria. (A)

BMP2/4 (BMP2B), (B) Forkhead (FOXA2), (C) Goosecoid (GSC) and (D) Brachyury

(TBXT). However, heatmaps show that, for most of these genes, rather than at

gastrulation (around 13 – 18 hpf) the expression maxima occur post-settlement. Only

one of the genes characterised in the context of gastrulation (snail) has a major peak

of expression at the late "prawn chip" (gastrula) stage (Figure 5) – which is

consistent with in situ data (Hayward et al. 2004) suggesting expression throughout

the presumptive mesendoderm. The in situs were previously published in Hayward

et al. (2015) Developmental Biology 399:337-347 (Elsevier).

1266

1267
Supplementary Figure 3.

Differential expression of genes encoding nuclear receptors during coral

development. Searching the *Acropora* data using the terms PF00104 Ligand-binding

domain of nuclear hormone receptor and PF00105 Zinc finger, C4 type (two

domains) yielded hits corresponding to 14 loci. Two of these loci code for predicted

protein isoforms with distinct expression profiles (XP_029208454.1 and

XP_029208455.1 are protein isoforms from the locus LOC114972094;

XP_044184859.1 and XP_044184860.1 are isoforms from the locus

LOC114969161). Consistent with possible roles in cue recognition and/ or

metamorphosis, seven of the 14 genes were upregulated at 133hpf. All of these

mRNAs were cleared at the 3-6hpi transition.

1279

egg

5hpf

13hpf

18hpf

gastrulation

23hpf

29hpf

40hpf

57hpf

Induction

24hpi

9hpi

133hpf

Small RNAs

XP_044168784.1_PIWL1
XP_044181962.1_SRRT
XP_029194207.2_DGCR8
XP_044181963.1_SRRT
XP_044181068.1_AGO2
XP_029185006.2_HENMT
XP_029197160.2_DICER
XP_044164157.1_DDX4
XP_044166539.1_MAEL
XP_044168427.1_DDX4
XP_029187700.2_PIWL2

0hpf 5hpf 13hpf 18hpf 23hpf 29hpf 35hpf 40hpf 45hpf 57hpf 69hpf 81hpf 93hpf 133hpf 15mpi 30mpi 1hpi 2hpi 3hpi 6hpi 9hpi 12hpi 18hpi 24hpi 48hpi 72hpi

Table 1. Previous studies on the molecular bases of early development in hard corals

Coral species	Reference	Stages sampled						Focus of study	Methodology*
Acropora millepora									
	Grasso et al., 2008	prawn chip 8 hpf	planula 83 hpf	primary polyp 130 hpf	adult			Early survey of developmental changes	ESTs
	Grasso et al., 2011	egg (unfertilised)	prawn chip 8 hpf	planula 83 hpf	primary polyp 130 hpf	adult		Pre- vs post-settlement	ESTs
	Hayward et al., 2011			planula	primary polyp			Pre vs post settlement	SSH
	Strader et al., 2018	22 hpf	46 hpf	73 hpf	89 hpf	24 hour intervals		Competence and fluorescence	
Acropora digitifera									
	Takeuchi et al., 2016	egg	blastula	gastrula	larva	meta	adult	Calcification genes	
	Reyes-Bermudez et al., 2016	prawn chip 12 hpf	gastrula 24 hpf	sphere 48 hpf	planula 96 hpf	adult		Stage specific transcript profiles	
Acropora tenuis									
	Ishii et al. 2022	0, 1 and 6 hr post induction						Commitment to settlement	
Montipora capitata									
	Chillie et al., 2021	0 hpf (egg) (Unfertilised)	1 hpf	4hpf	9hpf	14hpf	22hpf	Maternal-zygotic transition	
	He et al., 2024	0 hpf (egg) (unfertilised?)	prawn chip 14hpf	sphere 38hpf	pear 62hpf	86hpf	planula 110hpf	Developmental roles of Hox-Gbx genes	
	Huffmyer et al., 2025	Egg (fertilised) 1hpf	cleavage 5hpf	late gastrula 38hpf	late gastrula 65hpf	93hpf	183hpf		
Porites asteroides									
	Walker et al., 2019	Compared pre-competent and competent planulae						Compared non-probing and probing planulae	
	Mansour et al., 2016	Compared pre-settlement, post-settlement and adult						Calcification, settlement and symbiosis	
Montastrea faveolata	Reyes-Bermudez et al., 2009	Planula, primary polyp and adult						Metamorphosis and onset of calcification	Arrays
Stylophora pistllata	Levy et al., 2021	Planula, primary polyp and adult						Characterisation of single cell types	

*With the exception of those where the methodology is specified, these studies were based on high-throughput sequencing data

Nuclear Receptors