Spleen-derived Small Extracellular Vesicles Protect against Myocardial Infarction via Mediating Spleen-heart Crosstalk

Keyu Liu[1,*], MD, Yue Luo[1,*], MD, Weifeng Liao[2,3,*], MD, Riken Chen[4,*], MD, PhD, Tugang Chen[5], MD, PhD, Haitao Huang[5], MD, Jinrong Xu[5], MD, Fang Fang[6], PhD, Qinghua Chen[2], Dongling Lin[2], Ting Gu[2], Burton B Yang[7,#], MD, PhD, Wenliang Chen[2,3,#], MD, PhD, Liangqing Zhang[1,#], MD, PhD.

Department of Anesthesiology, the Second Affiliated Hospital of Guangdong Medical University, Guangdong Medical University, Zhanjiang, Guangdong, China

Translational Medicine Center, the Second Affiliated Hospital of Guangdong Medical University, Guangdong Medical University, Zhanjiang, Guangdong, China

Medical Interdisciplinary Science Research Center of Western Guangdong, the Second Affiliated Hospital of Guangdong Medical University, Guangdong Medical University, Zhanjiang, Guangdong, China

Respiratory and Critical Care Medicine, the Second Affiliated Hospital of Guangdong Medical University, Zhanjiang, Guangdong, China

Department of Cardiovascular Medicine, the Second Affiliated Hospital of Guangdong Medical University, Guangdong Medical University, Zhanjiang, Guangdong, China

Office of Academic Research, the Second Affiliated Hospital of Guangdong Medical University, Guangdong Medical University, Zhanjiang, Guangdong, China

Department of Laboratory Medicine and Pathobiology, University of Toronto, 2075 Bayview Ave, Toronto, Ontario, M4N 3M5 Canada

Short title: A novel mediator in cardiosplenic axis

* These authors contribute equally.

 # Correspondence:

Liangqing   Zhang,   email: zhangliangqing@gdmu.edu.cn,   Department   of

Anesthesiology, the Second Affiliated Hospital of Guangdong Medical University,

Guangdong Medical University, Zhanjiang, Guangdong, China

Wenliang Chen, email: chenwl@gdmu.edu.cn, Translational Medicine Center, the

Second Affiliated Hospital of Guangdong Medical University, Guangdong Medical

University, Zhanjiang, Guangdong, China

Burton B Yang, email: byang@sri.utoronto.ca, Department of Laboratory Medicine

and Pathobiology, University of Toronto, 2075 Bayview Ave, Toronto, Ontario, M4N

3M5 Canada

Words count： 10238 words

Abstract

Background: The cardiosplenic axis has recently begun to be recognized as an important mediator in ischemic heart disease, but its mechanistic role remains incompletely understood. This study aimed to investigate novel molecular mechanisms underlying spleen-heart crosstalk in acute myocardial infarction (AMI), focusing on the role of spleen-derived small extracellular vesicles (sEVs).

Methods: We used both a murine AMI model and an oxygen-glucose deprivation (OGD) cell model. Orthogonal approaches, including splenectomy, TTC staining, Masson's trichrome staining, wheat agglutinin staining, echocardiography, TUNEL staining, RNA sequencing, transmission electron microscope imaging, GW4869 administration as a sEVs inhibitor, and functional assays, was employed to evaluate the role of spleen-derived sEVs in AMI-induced cardiac injuries and dysfunction. 4D-proteomics was conducted to profile the protein cargo of sEVs. The function of mitochondrial pyruvate carrier 1 (MPC1) was evaluated with specific antibody blockade and pharmacological inhibition (UK5099, GW604714x). Plasma levels of sEVs-capsulated MPC1 were compared between patients with and without coronary heart disease (CHD), and AMI using ELISA.

Results: Spleen immune activity was upregulated in AMI mice. Splenectomy worsened AMI outcomes by reducing survival, impairing cardiac function, and increasing fibrosis. Spleen-derived sEVs levels increased with a temporal manner following AMI and was taken up by ischemic myocardium, peaking at day 3 post AMI (defined as M3D-sEVs). Administration of M3D-sEVs improved survival, reduced

infarct size, and restored mitochondrial respiration. Conversely, sEVs depletion by

GW4869 aggravated cardiac ischemic damage and left ventricle dysfunction.

Proteomic analysis revealed that cardioprotective M3D-sEVs were enriched for

MPC1, which promoted pyruvate metabolism and increased ATP/ADP ratio in

recipient cardiomyocytes. Neutralization of MPC1 within M3D-sEVs abrogated these

beneficial effects. Clinically, plasma sEVs-MPC1 levels were significantly elevated in

AMI patients compared to CHD patients without AMI, and in CHD patients compared

to non-CHD controls.

Conclusions: The spleen exerts cardioprotection against AMI by releasing

MPC1-enriched sEVs that preserve mitochondrial energetics in the injured

myocardium. Augmenting this endogenous sEVs-mediated pathway may represent a

promising therapeutic strategy for ischemic cardiomyopathy.

Keywords: Acute myocardial infarction; Spleen; Small extracellular vesicles; MPC1;

Cardiosplenic axis

Clinical Perspective

What Is New?

- Acute myocardial infarction (AMI) induces transcriptional activation of splenic extracellular vesicle-related genes, resulting in elevated vesicle secretion.

- Following AMI, splenic sEVs enriched with mitochondrial pyruvate carrier 1 (MPC1) increase in response to cardiac injury. These sEVs enhance mitochondrial metabolic energy production in the heart, critically influencing post-MI outcomes, displaying as reducing mortality, promoting wound healing and repair, and attenuating long-term adverse cardiac remodeling.

- Levels of MPC1 in circulating sEVs are elevated in patients with coronary heart disease (CHD) and show a further significant increase in those with AMI patients.

What Are the Clinical Implication?

- Our findings reveal a novel heart-spleen communication axis mediated by sEVs following AMI. These vesicular messengers enhance cardiac mitochondrial energy metabolism in the injured myocardium, ultimately improving functional recovery and clinical outcomes post-MI.

- Circulating sEVs MPC1 is a potential biomarker for the diagnosis of myocardial injury.

Abbreviations

CVD, Cardiovascular Disease. **CHD,** Coronary Heart Disease. **AMI,** Acute myocardial

infarction. **OGD,** Oxygen-glucose deprivation. **sEVs,** Small extracellular vesicles.

P-sEVs, Plasma small extracellular vesicles. **AARIVE,** Animal Research: Reporting

of In vivo Experiments. **Echo,** Echocardiograph. **EF,** ejection fraction. **FS,** fractional

shortening. **MPC1,** Mitochondrial pyruvate carrier 1. **PCK2,** Phosphoenolpyruvate

carboxykinase 2. **α-SMA,** α-smooth muscle actin. **Bax,** Bcl-2-associated X protein.

TUNEL, terminal deoxynucleotidyl transferase dUTP nick end labeling. **C-Cas-3,**

cleaved caspase 3. **CK-MB,** creatine kinase MB isoenzyme. **ELISA,** Enzyme-Linked

Immunosorbent Assay. **WGA,** Wheat germ agglutinin. **nFCM,** Nano-flow cytometer.

OCR, Oxygen Consumption Rate. **LC-MS**, Liquid Chromatograph Mass

Spectrometer.

RNA-seq, RNA sequencing. **LV,** left ventricular. ETC, Electron Transport Chain.

TCA,

Tricarboxylic Acid. **4D-DIA,** Four-dimensional data-independent acquisition.

Introduction

Acute myocardial infarction (AMI) results from sudden interruption of coronary blood flow, leading to ischemia and subsequent injury to the myocardium. Despite major advancements in early reperfusion therapy, AMI remains a leading cause of morbidity and mortality worldwide[1]. A deeper understanding of the molecular mechanisms involved in AMI pathogenesis and an identification of novel therapeutic targets are still urgently needed.

The concept of crosstalk between the spleen and heart, termed the cardiosplenic axis, was first described in 1949 that splenic nerves activation by electric stimulation improves ventricle function in dogs[2]. Since then, the spleen has been implicated cardioprotection, potentially through its regulation of inflammatory processes after myocardial injury by mobilizing immune cell populations such as monocytes and by releasing signaling factors that influence downstream effects within the heart[3,4]. For example, stimulation of splenic activity vie the vagus nerve during remote ischemic conditioning was shown to reduce cardiac infarct size in a rat model of AMI[5], suggesting that spleen-derived factors may play a critical role in modulating ischemic cardiac injury. However, the precise molecular mechanisms underlying heart-spleen communication remain to be fully elucidated.

Small extracellular vesicles (sEVs), including exosomes, have be proposed as critical mediators of intercellular and inter-organ signaling[6], particularly in the context of cardiovascular injury[7]. sEVs are nanosized vesicles (30-150nm) that carry a diverse range of bioactive molecular cargo, including proteins, lipids, and nucleic acids. The

cargos are shuttled systemically to recipient cells to regulate their function. The roles

of sEVs in mediating cellular and organ-level communications make it promising

candidates for use as disease biomarkers, therapeutic agents, or drug delivery

vehivles[8]. Recent studies have shown that plasma sEVs derived from AMI patients

promote angiogenesis, regulate ferroptosis and autophagy, and thus to alleviate

hypoxic myocardial injuries[9-11]. It has also been reported that the majority of plasma

sEVs in AMI patients originate from the spleen[12], indicating that spleen-derived sEVs

play critical roles in the pathophysiology of AMI.

Ischemic injury triggers a cascade of pathological responses, including inflammation,

metabolic dysfunction, and cell death, which ultimately contribute to adverse cardiac

remodeling and progression to heart failure[13]. Among these processes, mitochondrial

dysfunction plays a central role in AMI, as the heart's high energy demands are

heavily dependent on mitochondrial oxidative metabolism[14]. Ischemia-induced

impairment of mitochondrial function compromises ATP production generates

excessive reactive oxygen species (ROS), and promotes cardiomyocyte apoptosis,

further exacerbating myocardial injury and impairing functional recovery[15].

Mitochondrial Pyruvate Carrier 1 (MPC1), a key subunit of the mitochondrial pyruvate

carrier (MPC) complex, has gained attention as a critical regulator of mitochondrial

bioenergetics. The MPC complex, composed of MPC1 and MPC2, facilitates

transport of pyruvate from the cytosol into the mitochondrial matrix, enabling its

subsequent entry into the tricarboxylic acid (TCA) cycle and oxidative

phosphorylation[16]. Pyruvate metabolism is essential for efficient energy production,

particularly in cardiomyocytes, which has a limited capacity to adapt to metabolic

stress[17]. Dysregulation of MPC1 expression or function in ischemic conditions

disrupts this process, leading to cytosolic pyruvate accumulation, diminished TCA

cycle activity, and a shift toward less efficient anaerobic glycolysis[18]. Restoration of

MPC1 expression during ischemic injury has been shown to enhance mitochondrial

function, reduce oxidative stress, and facilitate cardiomyocyte survival, making MPC1

as a promising therapeutic target for metabolic management of AMI[19].

In this study, we investigated the crosstalk between spleen and heart during AMI,

focusing on the potential role of spleen-derived sEVs in delivering MPC1 to the

ischemic heart and preserving mitochondrial function in cardiomyocytes. Elucidating

these mechanisms may offer valuable insights into systemic regulation of

mitochondrial bioenergetics and identify novel therapeutic strategies to enhance

myocardial repair and recovery following ischemic injury.

Methods

Detailed methods

Comprehensive descriptions of experimental procedures, protocols, and reagents are

provided in the Supplemental Material.

Data Availability Statement

The data, analytic methods, and study materials supporting the findings of this study

are available to other researchers from the corresponding author upon reasonable

request. RNA sequencing (RNA-seq) raw data have been deposited at the National

Center for Biotechnology Information (BioProject: PRJNA1262715). Mass

spectrometry proteomics data have been deposited to the Proteome Xchange

Consortium (https://proteomecentral.proteomexchange.org) with the dataset identifier

PXD064000 and PXD063946.

Study Participants

A total of 128 subjects undergoing evaluation for coronary artery disease were

enrolled in this study. Among these, 104 patients scheduled for coronary angiography

were included, being composed of 21 patients with coronary heart disease (CHD) and

myocardial infarction (MI), 48 patients with CHD without MI, and 35 patients without

CHD. Plasma and plasma sEVs MPC1 levels were measured in all participants.

Written informed consent was obtained from all subjects, and the study protocol was

approved by the Human Ethics Committee of the Second Affiliated Hospital of

Guangdong Medical University (No. PJKT2024-163), in accordance with the ethical

guidelines of the 1975 Declaration of Helsinki.

Animal Experiments

Male C57BL/6 mice (6-8 weeks old, 20-25 g) were purchased from Guangzhou

Tengke Biomedical Technology Co., Ltd (Guangzhou, China). Mice were housed

under specific pathogen free conditions at 25 ± 2 °C with a 12 h light/dark cycle and

ad libitum access to food and water. MI surgery was performed in 8- to 10-week-old

mice by permanent ligation of the left anterior descending coronary artery (LAD). All

animal protocols were approved by the Guangdong Medical University Animal Care

and Use Committee (GDMU-2024-000043) and conducted in accordance with the

ARRIVE 2.0 (Animal Research Reporting of In Vivo Experiments) guidelines.

Statistical Analysis

For in vitro and in vivo experiments, cells or animals were randomly assigned to

experimental groups. At least five biologically independent replicates were performed

per group. Data are presented as mean ± standard error of the mean (SEM) for

normally distributed variables, or as median (interquartile range) as appropriate.

Normality was assessed using the Shapiro-Wilk test. For normally distributed data,

comparisons between two groups were performed using unpaired two-tailed

Student's t test, and comparisons among more than two groups were performed

using one-way ANOVA with Tukey's or Dunn's multiple comparisons test as

appropriate. For non-normally distributed data, the Mann-Whitney U test (two groups)

or Kruskal-Wallis test with Dunn's post-hoc test (>2 groups) was used. When two

independent factors were analyzed, two-way ANOVA with Tukey's multiple

comparisons test was applied. Mixed-effects ANOVA was used for serial

echocardiographic measurements. Homogeneity of variance was tested with the

Brown-Forsythe test; if violated, the Welch test was used. Survival analyses were

performed using the Kaplan-Meier method and log-rank test. Statistical analyses

were conducted using GraphPad Prism (version 9.0). Significance was defined as

$P < 0.05$.

For clinical data, categorical variables are reported as counts and percentages, with

comparisons performed using Chi-square or Fisher's exact tests as appropriate. All

statistical tests were two-sided, with significance set at $P < 0.05$. Clinical statistical

analyses were performed using SPSS (version 27.0, IBM). Detailed statistical

approaches and replicate numbers are provided in the figure legends.

Results

Splenectomy aggravates cardiac injury in AMI mice

We first assessed the splenic immune response post myocardial ischemia in AMI

mice. As expected, both heart and spleen weights were significantly increased in AMI

mice compared to sham-operated controls on day 3 post AMI (Figure S1A-S1C). The

white pulp, which is critical for the initiation of adaptive immune responses in

spleen[20,21], was significantly enlarged in AMI mice, as demonstrated by H&E staining;

the marginal zone was also expanded relative to sham mice (Figure S1D-S1F). In the

meantime, mRNA expression of inflammatory cytokines, such as IL-6, IFN-γ, TNF-α,

IL-10, was significantly upregulated in the spleens of AMI mice, with all cytokines

peaking at day 3 post AMI (Figure S1G-S1J). These findings indicate that splenic

biological activity is markedly enhanced in response to myocardial ischemic injury.

Previous studies have reported variable effects of splenectomy on cardiac injury[22-24].

To clarify the impact of spleen removal, we performed splenectomy immediately

following ligation of left anterior descending (LAD) of coronary artery in AMI mice and

evaluated cardiac outcomes. Compared with AMI mice with intact spleens,

splenectomized AMI mice exhibited significantly poorer survival rates, reduced left

ventricular ejection fraction and fractional shortening (Figure 1A-1D), larger cardiac

infarct size (Figure 1E-1F), and higher serum creatine kinase-MB (CK-MB) levels

(Figure S2A). Splenectomy also led to a dramatical increase in cardiac collagen

deposition, as determined by Masson's trichrome staining (Figure 1G-1H), and

elevated expression of fibrosis-related proteins, such as α smooth muscle actin

(α-SMA), collagen I (Col), and fibronectin (FN), in MI hearts (Figure 1I-1J), indicating

that splenectomy increased cardiac fibrosis post AMI. Furthermore, splenectomized

mice also exhibited a higher proportion of TUNEL positive cardiomyocytes (Figure

1K-1L), increased expression of pro-apoptotic markers Bax and cleaved caspase-3

(C-Cas3) and decreased expression of anti-apoptotic protein Bcl2 expression in heart

at day 3 post AMI (Figure 1M-1N). Myocardial hypertrophic remodeling, as measured

by wheat germ agglutinin (WGA) staining at 28 days post MI, was also significantly

aggravated in splenectomized AMI mice compared to controls (Figure 1O-1P).

Histological analyses revealed that splenectomy with AMI mice worsened myocardial

structural disarray, as shown by H&E staining (Figure S2B). These results together

demonstrate that splenectomy significantly worsens cardiac dysfunction and adverse

remodeling following AMI, indicating the existence of spleen-heart crosstalk and

supporting an endogenously protective role of the spleen against ischemic cardiac

injury.

Extracellular vesicles-related gene expression is upregulated in the spleen of

AMI mice

To investigate transcriptomics alteration in the spleen post AMI, we performed

RNA-Seq analysis of spleen tissues (Figure 2A). A total of 2621 differentially

expressed genes (DEGs) were found in spleens from AMI mice versus

sham-operated controls, with 1129 genes upregulated and 1492 genes

downregulated (Figure 2B and Figure S3A). Gene Ontology (GO) analysis revealed

that these DEGs were predominantly enriched in biological function associated with

extracellular vesicles (Figure S3B-S3E). Gene set enrichment analysis (GSEA)

further demonstrated a significant increase in gene signatures related to extracellular

exosome and vesicle regulation in the MI group compared with the sham group

(Figure S3F-S3G). The biogenesis processes of EVs are intricate and can be

affected by multiple pathways[25], such as soluble N-ethylmaleimide-sensitive fusion

attachment protein receptor (SNARE), Neutral sphingomyelin phosphodiesterase 3

(Smpd3), and Rab GTPases (e.g., Rab27a/Rab27b, Rab11). Therefore, we sought to

validate RNA-Seq findings by assessing the expression levels of key sEVs

biogenesis-related genes via quantitative PCR. Several genes involved in sEVs

formation and trafficking, including Rab27a, Rab31, Rab11, Pdcd6ip, Rab22a, and

Smpd3, were found to be significantly upregulated in the spleens of AMI mice

compared to sham controls (Figure 2C), which is consistent with RNA-Seq results.

These findings indicate that myocardial ischemia induces the upregulation of

extracellular vesicles biogenesis in the spleen, prompting further investigation into the

role and function of spleen-derived sEVs in AMI.

Spleen-derived small extracellular vesicles biogenesis is enhanced in response

to AMI

sEVs were isolated from mouse spleen using established protocols from our previous

work[26] (Figure 2D). The physical characteristics of the isolated sEVs were validated

by transmission electron microscopy (TEM), western blotting, and nano-flow

cytometry (nFCM). The isolated particles were positive for conventional sEVs

markers (TSG101, CD63, and CD9), while the negative markers GM130 and TBP

were undetectable, confirming the purity of the preparations. We observed that both

the protein contents and total particle number of spleen-derived sEVs were

significantly increased in response to myocardial infarction (MI), peaking at day 3 post

MI (Figure 2E-2F, Figure S4A). TEM and nFCM analysis showed that vesicles

exhibited the expected cup-shaped, double-membrane morphology with diameters

primarily ranging from 30 to 200 nm (Figure 2G-2H). These results confirm that the

isolated particles are sEVs and indicate that splenic sEVs secretion is significantly

increased during AMI, particularly peaking at day 3 post MI.

To determine if spleen-derived sEVs could reside in cardiac tissue in vivo, we labeled

isolated spleen-derived sEVs with Dimethyl red dye (DIR dye) and administered them

intramyocardially to mice. Bioluminescence imaging results showed that DIR-labeled

sEVs isolated from the spleens of MI mice predominantly accumulated in the hearts

of recipient AMI mice post injection (Figure 2I-2J), while markedly weak

bioluminescence signals were observed in the liver, spleen, and lungs (Figure S4B),

suggesting that spleen-derived sEVs preferentially target the ischemic myocardium.

These findings demonstrate that biogenesis and release of spleen-derived sEVs are

significantly upregulated during the acute phase of MI and that these vesicles are

capable of trafficking to the injured myocardium and accumulating within it.

331　Spleen-derived sEVs alleviate ischemic injuries in the heart and

332　cardiomyocytes

We next evaluated the effects of spleen-derived sEVs on ischemic injury induced by

AMI. Recipient AMI mice received intramyocardial injections of spleen-derived sEVs

isolated from donor sham or AMI mice. Comparative analyses revealed that

M3D-sEVs exerted the most predominant beneficial effects on survival in recipient

AMI mice, significantly increasing survival rates compared to other groups (Figure 3A).

In addition, M3D-sEVs but not M1D-sEVs or M7D-sEVs administration markedly

improve cardiac function, as evidenced by significantly increased left ventricular

ejection fraction and fractional shortening (Figure 3B-3D), reduced infarct area

(Figure 3E-3F) and decreased collagen accumulation (Figure 3G and 3H) in recipient

AMI mice. Furthermore, M3D-sEVs significantly attenuated the expression of

fibrosis-related proteins (α-SMA, Col, and FN) (Figure 3I and 3J) in hearts of recipient

AMI mice. M3D-sEVs treatment led to decreased myocardial structural disarray,

TUNEL-positive cardiomyocytes, and serum CK-MB levels (Figure S5A-S5D). It also

modulated apoptosis-related protein expression, with downregulation of Bax and

cleaved caspase-3 (C-Cas-3), and upregulation of Bcl2 (Figure 3K-3L). Additionally,

M3D-sEVs alleviated cardiac hypertrophy in ischemic mouse hearts (Figure 3M-3N).

To further investigate the direct effects on cardiomyocytes, H9C2 cells were

incubated with PKH26-labeled M3D-sEVs. Fluorescent signal was readily observed

within H9C2 cells at 12 and 24 hours post treatment (Figure S6A), suggesting

efficient sEVs uptake by these cells. Consistent with the in vivo findings, M3D-sEVs

treatment affected expressions of apoptosis-related proteins in H9C2 cells subjected

to OGD, resulting in upregulation of Bcl2 and downregulation of C-Cas-3 and Bax

(Figure S6B-S6C).

Taken together, these results demonstrate that spleen-derived sEVs, specifically

those collected at day 3 post AMI, exert significant cardioprotective effects by

attenuating ischemic injury and apoptosis both in vivo and in vitro. Accordingly,

M3D-sEVs were used in subsequent experiments.

Blockage of sEVs biogenesis aggravated cardiac ischemic injury in mice

To further assess the functional importance of spleen-derived sEVs in AMI, we used

GW4869, a well-characterized pharmacological inhibitor that suppresses

ceramide-mediated sEVs formation and release by targeting neutral

sphingomyelinase (N-Smase). Initial safety experiments demonstrated that

intraperitoneal administration of GW4869 (2.5 mg/kg/day for consecutive three days)

did not induce cardiac toxicity, in terms of CK-MB levels, TNF-α and IL-10 mRNA

expression, cardiac function, cardiac fibrosis, and myocardial structure, as evaluated

by qPCR, echocardiography, TTC staining, H&E, and Masson's trichrome staining

(Figure S7A-S7I). A single dose of GW4869 injection reduced spleen-derived sEVs

protein content for one day, with levels comparable to the vehicle group thereafter

(Figure S8A), suggesting its inhibitory effects on sEVs biogenesis in the spleen are

short-lived. Thus, daily injections were used for subsequent studies (Figure 4A).

Under AMI conditions, GW4869 administration significantly decreased survival,

reduced left ventricular ejection fraction and fractional shortening, while increased

infarct area compared with the vehicle group (Figure 4B-4G). Additionally, GW4869

treatment exacerbated adverse remodeling and injury, evidenced by elevated

collagen deposition, increased protein expression of Col, FN, α-SMA, worse

myocardial structural disarray, and higher serum CK-MB levels (Figure 4H-4K, Figure

S8B and S8C). Pro-apoptotic markers Bax and C-Cas-3 were also upregulated,

whereas anti-apoptotic Bcl2 was downregulated in ischemic hearts following GW4869

treatment (Figure 4L-M). Furthermore, GW4869 aggravated MI-induced cardiac

hypertrophy (Figure 4N-O). Importantly, the deleterious effects of GW4869 were

rescued by the administration of M3D-sEVs, but not by sEVs from sham mice

(S-sEVs) (Figure 4A-4O, Figure S8B and S8C). These findings demonstrate that the

release of spleen-derived sEVs during AMI plays an endogenously protective role

against cardiac damage and adverse remodeling.

Spleen-derived sEVs improve myocardial mitochondria energy metabolism

To explore the mechanisms underlying the cardioprotective effects of M3D-sEVs, we

performed 4D-proteomic profiling to compare the protein cargo of spleen-derived

sEVs from sham and AMI mice (Figure 5A). A total of 1068 differentially expressed

proteins (DEPs) were identified (M3D-sEVs vs S-sEVs), with 389 up-regulated and

679 down-regulated proteins in M3D-sEVs relative to S-sEVs (Figure 5B). GO

enrichment analysis showed that these DEPs are associated with metabolic process,

transport, intracellular organelle, electron transfer activity, and structural molecular

activity (Figure 5C-5E). KEGG pathway analysis demonstrated that they were mainly

enriched in oxidative phosphorylation, metabolic pathways, and cardiac muscle

contraction (Figure 5F). These results suggest that mitochondrial metabolism-related

oxidative phosphorylation (OXPHOS) may serve as a key mechanism by

which M3D-sEVs alleviate injury in acute myocardial infarction (AMI). We next

examined mitochondrial energy metabolism, a critical determinant of myocardial

response to ischemic injury. As shown in Figure 6A-6B, M3D-sEVs treatment

alleviated MI-induced abnormalities in cardiac mitochondrial morphology, including

restoration of organized mitochondrial cristae, preservation of myofibril integrity and

recovery of mitochondrial size. These protective effects were not observed in mice

treated with spleen-derived sEVs from Sham controls.

To further elucidate the impact of spleen-derived sEVs in mitochondrial function, we

assessed the expression of genes and proteins associated with the mitochondrial

electron transport chain (ETC) complexes in cardiac tissue. Immunoblot analysis

demonstrated that protein levels of mitochondrial complex I (NDUFB8), complex II

(SDHB), complex IV (MTCO1), and complex V (ATP5A) were significantly reduced in

hearts from MI mice, while they were restored by M3D-sEVs treatment. In contrast,

expression of complex III (UQCRC2) remained unchanged across experimental

groups (Figure 6C-6F). Supporting these observations, qPCR analysis revealed

significant downregulation of ETC subunit mRNA expression, including complex I

(Ndufs1 and Ndufs2), complex II (Sdhc and Sdhd), complex III (Uqcrfs1 and Uqcrc1),

complex IV (Cox4il) and complex V (Atp5po and Atp5f1b) in ischemic myocardial

tissue and OGD-insulted H9C2 cells, which was significantly rescued by M3D-sEVs

but not S-sEVs administration (Figure 6G-6Q). These findings suggest that

spleen-derived M3D-sEVs prevent ischemia-induced disruption of the cardiac

mitochondrial ETC.

We next extended our investigation of energy metabolism in cardiomyocytes using

H9C2 cells. M3D-sEVs but not S-sEVs treatment restored ATP/ADP ratio in

OGD-insulted cells compared with those treated with vehicle (Figure 6R). In addition,

M3D-sEVs treatment improved the oxygen consumption rate (OCR) in H9C2 cells

subjected to OGD and attenuated abnormal mitochondrial superoxide production, as

examined by mitoSOX staining (Figure 6S-6U). Collectively, these data suggest that

the cardioprotective effects of M3D-sEVs in AMI are mediated, at least in part, by

mitigation of mitochondrial injury and restoration of mitochondrial energy metabolism

when hearts suffered from ischemic stress.

MPC1 protein expression is increased in M3D-sEVs from AMI mouse and in

circulating plasma sEVs from AMI patients.

To identify critical cardioprotective factors inside M3D-sEVs, we performed proteomic

profiling to compare the protein of plasma from sham and AMI mice (Figure S9).

When cross-referenced with DEPs between plasma and sEVs proteomics, 21

upregulated and 12 downregulated overlapping proteins were found by using Venn

diagrams (Figure 7A, Table S4). We then focused on mitochondrial energy

metabolism-related proteins, and selected MPC1 and PCK2 accordingly for further

validation as their changes were most significant among these proteins. As shown in

Figure 7B-7C, MPC1 protein expression was significantly increased in M3D-sEVs

relative to S-sEVs, while PCK2 levels remain unchanged. Furthermore, when

comparing sEVs from multiple organs post-AMI, increased MPC1 protein was

uniquely observed in spleen-derived sEVs but not in sEVs from other tissues (Figure

7D). Notably, MPC1 was even decreased in sEVs from the heart and lung of AMI

mice. Moreover, to assess the potential clinical relevance of MPC1 in ischemic heart

disease, we determined MPC1 expression in plasma-derived sEVs from human

subjects (Figure 7E). The baseline parameters of recruited patients are summarized

in Table S5. TEM images and nFCM analysis confirmed the successful isolation of

plasma sEVs (P-sEVs), which showed a diameter in the range of 30-200nm (Figure

7F-7G). Additionally, P-sEVs preparations were positive for marker proteins (CD63,

CD9, and TSG101) and negative for GM130 and TBP (Figure 7H), confirming sample

purity. As shown in Figure 7I-7J, quantification analysis exhibited that plasma

sEVs-MPC1 protein levels were significantly higher in AMI patients than that in

patients with CHD without MI, as well as in patients without CHD. Plasma MPC1

protein was significantly increased in AMI patients but not in CHD without AMI when

compared with that in these without CHD. These clinical findings imply that

sEVs-encapsulated MPC1 elevates in response to ischemic conditions in hearts.

Increased MPC1 expression accounts for the regulatory effects of M3D-sEVs

on myocardial mitochondrial energy metabolism

We neutralized MPC1 in M3D-sEVs using a specific antibody (sEVs+Ab) and

administered it via intramyocardial injection in AMI mice. This neutralization

significantly reduced the beneficial effects of M3D-sEVs, shown by decreased

survival rates (Figure 8A), worsened cardiac function (Figure 8B-8D), increased

infarct size (Figure 8E-8F), and elevated cardiac fibrosis and levels of fibrosis-related

proteins (Col, FN, and α-SMA) (Figure 8G-8J). Additionally, MI-induced increase in

cardiomyocyte area was more pronounced with MPC1 antagonism compared to

M3D-sEVs treatment alone or M3D-sEVs incubated with control IgG (Figure 8K-8L).

To further validate the role of MPC1, we employed MPC1 inhibitors, including UK5099

and GW604714x (GWx), to explore whether MPC1 contributes to the therapeutic

outcomes of M3D-sEVs. As expected, both MPC1 inhibitors reduced mitochondrial

pyruvate levels in H9C2 cells (Figure S10A-S10B) and were used at concentrations

of 2.5 μM and 60 nM, respectively, for subsequent investigations. The inhibition of

MPC1 with either compound abolished the beneficial effects of M3D-sEVs on ETC

complex subunit mRNA expression in AMI hearts (Figure 8M-8Q). Specifically, both

inhibitors significantly blunted the beneficial outcomes of M3D-sEVs in mitochondrial

morphological characteristics as shown by TEM (Figure 8R-8S), ATP/ADP ratio, and

oxygen consumption rate under OGD conditions (Figure 8T-8U). These results

demonstrate that the overall cardioprotective effects of M3D-sEVs were substantially

suppressed by MPC1 inhibition.

Discussion

In this study, we uncovered a novel mechanism accounting for spleen-heart crosstalk, providing compelling evidence that spleen-derived sEVs exert endogenous cardioprotective effects in AMI through enhancing delivery of MPC1. Our findings demonstrate that splenic sEVs biogenesis is dramatically upregulated following AMI, and these vesicles significantly attenuate ischemic cardiac damage in recipient mice. Moreover, MPC1 cargo is significantly increased in spleen-derived sEVs from AMI mice and is critical in the protective effects of these sEVs on ischemic myocardium. These results advance our understanding of the spleen's protective role in the early phase of AMI, highlighting a pathway whereby increasing sEVs biogenesis facilitates the delivery of MPC1 to injured myocardial tissue, and thus to improve mitochondrial energy metabolism. To our knowledge, this is the first study to identify spleen-derived sEVs as a mediator of spleen-heart crosstalk under ischemic conditions.

The concept of a cardio-splenic axis as a critical regulator of ischemic heart disease is well supported by previous studies.[3,27]. Increased metabolic activity of spleen was observed patients with acute coronary syndrome, which was measured by [18]F-fluorodeoxyglucose-positron emission tomography imaging, and it was associated with systemic proinflammatory status and higher risks in cardiovascular disease events[28]. In rodent models, myocardial ischemia leads to splenic hypertrophy[29], and myocardial ischemia/reperfusion (I/R) results in a rapid upregulation of splenic leukocyte mobilization-related gene expression (4h post reperfusion)[30]. Consistent with these reports, we observed acute splenic activation

post AMI in mice, characterized by increased spleen size, splenic weight, and

expansion of the white pulp area and marginal zone, as well as elevated expression

of inflammatory cytokines (IL-6, IFN-γ, TNF-α, IL-10) in the spleen, which is

consistent with Huizhen Lv and colleagues' study[31]. These findings support the notion

that the spleen responds dynamically to myocardial injury, possibly through

recognition of damage-associated molecular patterns derived from the ischemic

heart[27]. It was reported that indigenous Bajau people who could dive to 70 meters

underneath sea and were highly tolerant to diving-induced hypoxia, having increased

spleen size compared to their neighbor population, the Saluan[32]. This indicates that

enlarging spleen size post MI in mice might increase hypoxia tolerance to cardiac

ischemia.

The role of spleen in ischemic heart disease remains controversial. Recently, Qiang

Long and colleagues reported that splenectomy inhibited the infiltration of phagocytic

monocytes into the injured myocardium at the early phases and eventually improved

cardiac functions in I/R mice with ischemia for 60 min[30]. Similarly, splenectomy

significantly reduced cardiac infarct size in mice with ischemia for 40/50 min and

subsequent reperfusion for 60 min by lowering neutrophil infiltration in ischemic

myocardium. Moreover, the investigation in mice also showed that AMI enhanced

platelet releasing from spleen to facilitate myocardial inflammation, while

splenectomy blocked platelets and leucocytes accumulation in heart, reduced infarct

size, and improved cardiac functions[33]. In contrast, the epidemiological data from

World War II veterans suggest that splenectomy increases mortality rates following

myocardial infarction[34]. In consistent with this reports, a recent study showed that

splenectomy suppressed inflammation and exerted beneficial outcomes in long term

cardiac repair in AMI mice[23]. Our data demonstrate that splenectomy immediately

prior to MI surgery exacerbates cardiac injury and fibrosis, while reduced the survival

rates in mice, supporting that spleen works as a cardioprotective function in the early

phase of AMI. This is further supported by studies showing that remote ischemic

conditioning or neuromodulation in rats and pigs enhances the release splenic

cardioprotective factors, where splenectomy abrogates the deploying of these splenic

factors and blunts the cardiac protective effects of remote ischemic condition[3,35].

However, the precise nature of these protective factors has remained elusive.

Previous studies have focused predominantly on the role of splenic immune cells in

modulating post ischemic injury[30]. A set of studies demonstrated that splenic

neutrophils, monocytes, and lymphocytes could be rapidly released into circulation

post ischemic insults in hearts[30,33,36]. The splenic immune cells exhibit dual roles in

ischemic cardiac injury. For example, $Ly6C^{hi}$ monocytes deployed from subcapsular

red pulp of the spleen infiltrate to the infarct region of ischemic hearts, while inhibiting

Ly6C high monocyte infiltration promotes cardiac repair and improves cardiac

function after AMI[36]. In contrast, the injection of regulatory T cells (Tregs) enhanced

cardiac repair post AMI, showing as improved ejection fraction and decreased cardiac

fibrosis[37]. Recently, Mohamed Ameen Ismahil and colleagues found splenic

$CD169^+Tim4^+$ marginal metalophulid macrophages from marginal zone of spleen

increased in blood 24h post AMI, and these cells alleviated inflammation and restored

cardiac function after AMI in mice[23]. The dual and context-dependent roles of these

cells have complicated the development of targeted therapies for ischemic heart

disease. This complexity prompted us to investigate alternative mechanisms of

spleen-heart communication, focusing on sEVs as potential mediators.

sEVs have emerged as critical factors mediating intercellular and interorgan

communication [38,39], modulating the pathophysiology of ischemic heart disease by

delivering diverse biomolecules, including proteins, microRNAs, and lipids[40,41]. Our

transcriptomic analysis revealed that genes associated with extracellular vesicle

biology are upregulated in the spleen following AMI, prompting us to characterize the

temporal dynamics and functional properties of spleen-derived sEVs. We found that

splenic sEVs release peaks at day 3 post-AMI (M3D-sEVs), and that the

administration of these vesicles to recipient AMI mice markedly attenuates cardiac

injury, improves systolic function, and increases survival. These findings are

consistent with emerging evidence that organ-specific sEVs can modulate

inflammation, promote tissue repair, and limit ischemic injury[42]. Specifically, our

results suggest that splenic sEVs may deliver reparative cargo biomolecules, such as

proteins or regulatory RNAs, to the injured myocardium, coordinating beneficial

immune and metabolic responses.

In this study, proteomic profiling of M3D-sEVs identified significant enrichment of

proteins involved in oxidative phosphorylation, with MPC1 showing as a key

candidate. Both specific antibody-mediated neutralization and pharmacological

inhibition of MPC1 abolished the cardioprotective effects of M3D-sEVs, implicating

this protein as a critical effector. Importantly, increased MPC1 expression was unique

to spleen-derived sEVs post AMI in mice and not observed in sEVs from other main

organs. Functionally, M3D-sEVs enhanced MPC1 expression in recipient

cardiomyocytes, improved mitochondrial ATP/ADP ratios, while reduced

mitochondrial ROS generation, which are the hallmarks of optimized mitochondrial

bioenergetics and reduced oxidative stress. These findings were further supported by

our clinical data, which revealed significantly higher MPC1 levels in plasma sEVs

from AMI patients compared to non-AMI CHD patients, underscoring the translational

relevance of this mechanism.

MPC1 is responsible for mitochondrial pyruvate import, a critical step in maintaining

intracellular pyruvate and lactate balance and limiting lactate accumulation[43,44].

Previous studies have shown that impaired pyruvate metabolism and mitochondrial

dysfunction exacerbate ischemic injury and that MPC1 downregulation is associated

with heart failure and increased susceptibility to cell death[16,45,46]. Besides, Yue Liu

and colleges demonstrated that MPC1 downregulation enhances susceptibility to

ischemic injury by increasing mitochondria disruption, oxidative stress, calcium

overload, and autophagy in cortical neurons[47]. Conversely, augment MPC1

expression or activity have been shown to restore mitochondrial metabolism and

ameliorate cardiomyocyte injury in models of ischemia-reperfusion. Our findings

extend this paradigm by demonstrating that M3D-sEVs-mediated delivery of MPC1

constitutes a novel, endogenous rescue pathway during AMI. The observed

mitochondrial benefits, such as preserved ATP generation and reduced ROS

accumulation, highlight the importance of maintaining efficient mitochondrial pyruvate

flux for cardioprotection during acute ischemic stress. Thus, splenic sEVs-mediated

delivery of MPC1 represents a promising therapeutic axis for preserving

mitochondrial function during acute ischemia, warranting further mechanistic study

and consideration for translational application in AMI.

Clinically, we observed that plasma MPC1 levels are elevated not only in AMI patients

but also in those with non-AMI CHD compared to non-CHD controls, suggesting a

close association between sEVs-MPC1 and the severity of coronary heart disease.

Importantly, sEVs-associated biomarkers offer distinct advantages over traditional

markers such as troponins. They are stable in circulation, reflect dynamic

pathophysiological changes, and may provide organ-specific information[48]. Recent

work indicates that most plasma sEVs were derived from spleen in AMI patients[12],

supporting the feasibility of clinical investigations into splenic sEVs. The upregulation

of MPC1 in plasma sEVs in patients with ischemic heart disease may reflect acute

myocardial metabolic stress and mitochondrial adaptation, and sEVs-encapsulated

MPC1 could serve as a potential novel circulating biomarker for early AMI diagnosis.

In addition, sEVs-based platforms have demonstrated feasibility for early, minimally

invasive detection of various cardiac and non-cardiac conditions[49]. Our findings

suggest that plasma sEVs-MPC1, alone or in combination with established markers

such as troponins and natriuretic peptides, potentially enhance the diagnostic

accuracy for AMI, especially in challenging clinical scenarios or during the early

phase of myocardial injury. Larger prospective studies are warranted to validate the

diagnostic performance of sEVs-MPC1 and to establish standardized quantification

protocols and assess its utility in risk stratification and monitoring therapeutic

response in patients with suspected acute coronary syndromes.

Despite these promising findings, several limitations should be acknowledged. First,

our murine data and initial clinical data, while highly suggestive, may not fully capture

the complexity of sEVs biology and MPC1 regulation in humans. The clinical cohort

was modest in size and from a single center, potentially limiting generalizability.

Second, although we provide convincible evidence for the role of MPC1-enriched

spleen-derived sEVs in enhancing mitochondrial function and cardioprotection, the

precise molecular mechanisms underlying sEVs-MPC1 delivery, cardiac uptake, and

downstream signaling pathways require further elucidation. Finally, the temporal

kinetics of splenic sEVs-MPC1 elevation following AMI was not fully characterized in

this study.

In summary, our study reveals that spleen-derived sEVs carrying MPC1 exerting

significant cardioprotective effects by enhancing mitochondrial energy metabolism

and reducing oxidative stress in ischemic myocardium. We further demonstrate that

plasma sEVs-MPC1 levels are elevated in AMI patients, suggesting their potential as

a novel, clinically relevant biomarker for acute myocardial injury. Together, these

findings uncover a previously unrecognized role for spleen–heart crosstalk mediated

by splenic sEVs and highlight sEVs-MPC1 as both a mechanistic effector and a

translational target in ischemic heart disease.

Acknowledgements

None.

Source of Funding

This work was supported by the National Natural Science Foundation of China

(82370281 to L.Q.Z.), the Science and Technology Development Special Fund

Competitive Allocation Project of Zhanjiang City (2021A05086 to W.L.C.), the

High-level Talent Startup Fund (23H03, W.L.C.), Special Project for Clinical and Basic

Sci&Tech Innovation of Guangdong Medical University (GDMULCJC2024048,

W.L.C.),   Guangdong   Basic   and   Applied   Basic   Research   Foundation

(2025A1515010473, W.L.C.).

Disclosures

None.

Supplemental Material

Supplemental Methods

Tables S1-S5

Figure S1-S10

References 50-61

References

1.   Saito Y, Oyama K, Tsujita K, Yasuda S, Kobayashi Y. Treatment strategies of acute

myocardial infarction: updates on revascularization, pharmacological therapy, and beyond. *J*

Cardiol. 2023;81:168-178. doi: 10.1016/j.jjcc.2022.07.003

2.   Rein H. [The role of the spleen and liver in coronary or hypoxic myocardial insufficiency].

Pflugers Arch Gesamte Physiol Menschen Tiere. 1951;253:435-458. doi:

10.1007/BF00370032

3.   Heusch G, Kleinbongard P. The spleen in ischaemic heart disease. *Nat Rev Cardiol*. 2025. doi:

10.1038/s41569-024-01114-x

4.   Tian Y, Miao B, Charles EJ, Wu D, Kron IL, French BA, Yang Z. Stimulation of the Beta2

Adrenergic Receptor at Reperfusion Limits Myocardial Reperfusion Injury via an

Interleukin-10-Dependent Anti-Inflammatory Pathway in the Spleen. *Circ J*.

2018;82:2829-2836. doi: 10.1253/circj.CJ-18-0061

5.   Lieder HR, Paket U, Skyschally A, Rink AD, Baars T, Neuhauser M, Kleinbongard P, Heusch

G. Vago-splenic signal transduction of cardioprotection in humans. *Eur Heart J*.

2024;45:3164-3177. doi: 10.1093/eurheartj/ehae250

6.   Rohm TV, Cunha ERK, Olefsky JM. Metabolic Messengers: small extracellular vesicles. *Nat

Metab*. 2025;7:253-262. doi: 10.1038/s42255-024-01214-5

7.   Xiao J, Sluijter JPG. Extracellular vesicles in cardiovascular homeostasis and disease:

potential role in diagnosis and therapy. *Nat Rev Cardiol*. 2025. doi:

10.1038/s41569-025-01141-2

8.   Kumar MA, Baba SK, Sadida HQ, Marzooqi SA, Jerobin J, Altemani FH, Algehainy N, Alanazi

MA, Abou-Samra AB, Kumar R, et al. Extracellular vesicles as tools and targets in therapy for

diseases. *Signal Transduct Target Ther*. 2024;9:27. doi: 10.1038/s41392-024-01735-1

9.   Li H, Ding J, Liu W, Wang X, Feng Y, Guan H, Chen Z. Plasma exosomes from patients with

683   acute myocardial infarction alleviate myocardial injury by inhibiting ferroptosis through

684   miR-26b-5p/SLC7A11 axis. *Life Sci*. 2023;322:121649. doi: 10.1016/j.lfs.2023.121649

685 10. Duan S, Wang C, Xu X, Zhang X, Su G, Li Y, Fu S, Sun P, Tian J. Peripheral Serum

686   Exosomes Isolated from Patients with Acute Myocardial Infarction Promote Endothelial Cell

687   Angiogenesis via the miR-126-3p/TSC1/mTORC1/HIF-1alpha Pathway. *Int J Nanomedicine*.

688   2022;17:1577-1592. doi: 10.2147/IJN.S338937

689 11. Wang B, Cao C, Han D, Bai J, Guo J, Guo Q, Li D, Zhang J, Zhang Z, Wang Y, et al.

690   Dysregulation of miR-342-3p in plasma exosomes derived from convalescent AMI patients

691   and its consequences on cardiac repair. *Biomed Pharmacother*. 2021;142:112056. doi:

692   10.1016/j.biopha.2021.112056

693 12. Jin X, Xu W, Wu Q, Huang C, Song Y, Lian J. Detecting early-warning biomarkers associated

694   with heart-exosome genetic-signature for acute myocardial infarction: A source-tracking study

695   of exosome. *J Cell Mol Med*. 2024;28:e18334. doi: 10.1111/jcmm.18334

696 13. Thygesen K, Alpert JS, Jaffe AS, Chaitman BR, Bax JJ, Morrow DA, White HD, Mickley H,

697   Crea F, Van de Werf F, et al. [Fourth universal definition of myocardial infarction (2018)].

698   *Kardiol Pol*. 2018;76:1383-1415. doi: 10.5603/KP.2018.0203

699 14. Yang R, Ruan B, Wang R, Zhang X, Xing P, Li C, Zhang Y, Chang X, Song H, Zhang S, et al.

700   Cardiomyocyte betaII spectrin plays a critical role in maintaining cardiac function by regulating

701   mitochondrial respiratory function. *Cardiovasc Res*. 2024;120:1312-1326. doi:

702   10.1093/cvr/cvae116

703 15. Ong SB, Samangouei P, Kalkhoran SB, Hausenloy DJ. The mitochondrial permeability

704   transition pore and its role in myocardial ischemia reperfusion injury. *J Mol Cell Cardiol*.

2015;78:23-34. doi: 10.1016/j.yjmcc.2014.11.005

16.   Visker JR, Cluntun AA, Velasco-Silva JN, Eberhardt DR, Cedeno-Rosario L, Shankar TS,

Hamouche R, Ling J, Kwak H, Hillas JY, et al. Enhancing mitochondrial pyruvate metabolism

ameliorates ischemic reperfusion injury in the heart. *JCI Insight*. 2024;9. doi:

10.1172/jci.insight.180906

17.   Gray LR, Tompkins SC, Taylor EB. Regulation of pyruvate metabolism and human disease.

Cell Mol Life Sci. 2014;71:2577-2604. doi: 10.1007/s00018-013-1539-2

18.   Fernandez-Caggiano M, Prysyazhna O, Barallobre-Barreiro J, CalvinoSantos R, Aldama

Lopez G, Generosa Crespo-Leiro M, Eaton P, Domenech N. Analysis of Mitochondrial

Proteins in the Surviving Myocardium after Ischemia Identifies Mitochondrial Pyruvate Carrier

Expression as Possible Mediator of Tissue Viability. *Mol Cell Proteomics*. 2016;15:246-255.

doi: 10.1074/mcp.M115.051862

19.   Zhao S, Zhang Y, Bao S, Jiang L, Li Q, Kong Y, Cao J. A novel HMGA2/MPC-1/mTOR

signaling pathway promotes cell growth via facilitating Cr (VI)-induced glycolysis. *Chem Biol*

Interact. 2024;399:111141. doi: 10.1016/j.cbi.2024.111141

20.   Lewis SM, Williams A, Eisenbarth SC. Structure and function of the immune system in the

spleen. *Sci Immunol*. 2019;4. doi: 10.1126/sciimmunol.aau6085

21.   Mebius RE, Kraal G. Structure and function of the spleen. *Nat Rev Immunol*. 2005;5:606-616.

doi: 10.1038/nri1669

22.   Zhang X, Fang Y, Qin X, Zhang Y, Kang B, Zhong L, Liu B, Luo J, Wei Y. The Role of MCPIP1

in Macrophage Polarization and Cardiac Function Post-Myocardial Infarction. *Adv Sci (Weinh)*.

2025:e2500747. doi: 10.1002/advs.202500747

23.   Ismahil MA, Zhou G, Rajasekar S, Gao M, Bansal SS, Patel B, Limdi N, Xie M, Antipenko S,

Rokosh G, et al. Splenic CD169(+)Tim4(+) Marginal Metallophilic Macrophages Are Essential

for Wound Healing After Myocardial Infarction. *Circulation*. 2025. doi:

10.1161/CIRCULATIONAHA.124.071772

24.   Rolland TJ, Hudson ER, Graser LA, Zahra S, Cucinotta D, Weil BR. Splenic Modulation of the

Early Inflammatory Response to Regional and Global Ischemia/Reperfusion Injury in Swine.

Am J Physiol Heart Circ Physiol. 2025. doi: 10.1152/ajpheart.00714.2024

25.   Yu J, Sane S, Kim JE, Yun S, Kim HJ, Jo KB, Wright JP, Khoshdoozmasouleh N, Lee K, Oh

HT, et al. Biogenesis and delivery of extracellular vesicles: harnessing the power of EVs for

diagnostics and therapeutics. *Front Mol Biosci*. 2023;10:1330400. doi:

10.3389/fmolb.2023.1330400

26.   Luo Y, Liu K, Ling D, Gu T, Zhang L, Chen W. Isolation and Characterization of Exosomes

Derived from Mouse Spleen Tissues. *J Vis Exp*. 2024. doi: 10.3791/67234

27.   Tian Y, Pan D, Chordia MD, French BA, Kron IL, Yang Z. The spleen contributes importantly to

myocardial infarct exacerbation during post-ischemic reperfusion in mice via signaling

between cardiac HMGB1 and splenic RAGE. *Basic Res Cardiol*. 2016;111:62. doi:

10.1007/s00395-016-0583-0

28.   Emami H, Singh P, MacNabb M, Vucic E, Lavender Z, Rudd JH, Fayad ZA, Lehrer-Graiwer J,

Korsgren M, Figueroa AL, et al. Splenic metabolic activity predicts risk of future cardiovascular

events: demonstration of a cardiosplenic axis in humans. *JACC Cardiovasc Imaging*.

2015;8:121-130. doi: 10.1016/j.jcmg.2014.10.009

29.   Ruggiero SA, Huber JS, Murrant CL, Brunt KR, Simpson JA. Splenic blood-flow response

following myocardial infarction in rat. *Can J Physiol Pharmacol*. 2018;96:1060-1068. doi:

10.1139/cjpp-2018-0134

30.   Long Q, Rabi K, Cai Y, Li L, Huang S, Qian B, Zhong Y, Qi Z, Zhang Y, Huang K, et al.

Identification of splenic IRF7 as a nanotherapy target for tele-conditioning myocardial

reperfusion injury. *Nat Commun*. 2025;16:1909. doi: 10.1038/s41467-025-57048-6

31.   Lv H, Wang C, Liu Z, Quan M, Li K, Gou F, Shi X, Liu Q, Yu Y, Zhu P, et al. Suppression of the

Prostaglandin I2-Type 1 Interferon Axis Induces Extramedullary Hematopoiesis to Promote

Cardiac Repair After Myocardial Infarction. *Circulation*. 2025. doi:

10.1161/CIRCULATIONAHA.124.069420

32.   Ilardo MA, Moltke I, Korneliussen TS, Cheng J, Stern AJ, Racimo F, de Barros Damgaard P,

Sikora M, Seguin-Orlando A, Rasmussen S, et al. Physiological and Genetic Adaptations to

Diving in Sea Nomads. *Cell*. 2018;173:569-580 e515. doi: 10.1016/j.cell.2018.03.054

33.   Gao XM, Moore XL, Liu Y, Wang XY, Han LP, Su Y, Tsai A, Xu Q, Zhang M, Lambert GW, et

al. Splenic release of platelets contributes to increased circulating platelet size and

inflammation after myocardial infarction. *Clin Sci (Lond)*. 2016;130:1089-1104. doi:

10.1042/CS20160234

34.   Robinette CD, Fraumeni JF, Jr. Splenectomy and subsequent mortality in veterans of the

1939-45 war. *Lancet*. 1977;2:127-129. doi: 10.1016/s0140-6736(77)90132-5

35.   Lieder HR, Kleinbongard P, Skyschally A, Hagelschuer H, Chilian WM, Heusch G.

Vago-Splenic Axis in Signal Transduction of Remote Ischemic Preconditioning in Pigs and

Rats. *Circ Res*. 2018;123:1152-1163. doi: 10.1161/CIRCRESAHA.118.313859

36.   Zhang J, Hao W, Zhang J, Li T, Ma Y, Wang Y, Li X, Cui W, Du J. CXCL16 Promotes

Ly6Chigh Monocyte Infiltration and Impairs Heart Function after Acute Myocardial Infarction. *J*

Immunol. 2023;210:820-831. doi: 10.4049/jimmunol.2200249

37.   Alshoubaki YK, Nayer B, Lu YZ, Salimova E, Lau SN, Tan JL, Amann-Zalcenstein D, Hickey

PF, Del Monte-Nieto G, Vasanthakumar A, et al. Tregs delivered post-myocardial infarction

adopt an injury-specific phenotype promoting cardiac repair via macrophages in mice. *Nat*

Commun. 2024;15:6480. doi: 10.1038/s41467-024-50806-y

38.   Huang-Doran I, Zhang CY, Vidal-Puig A. Extracellular Vesicles: Novel Mediators of Cell

Communication In Metabolic Disease. *Trends Endocrinol Metab*. 2017;28:3-18. doi:

10.1016/j.tem.2016.10.003

39.   Hu S, Hu Y, Yan W. Extracellular vesicle-mediated interorgan communication in metabolic

diseases. *Trends Endocrinol Metab*. 2023;34:571-582. doi: 10.1016/j.tem.2023.06.002

40.   Tian C, Gao L, Rudebush TL, Yu L, Zucker IH. Extracellular Vesicles Regulate

Sympatho-Excitation by Nrf2 in Heart Failure. *Circ Res*. 2022;131:687-700. doi:

10.1161/CIRCRESAHA.122.320916

41.   Fu S, Zhang Y, Li Y, Luo L, Zhao Y, Yao Y. Extracellular vesicles in cardiovascular diseases.

Cell Death Discov. 2020;6:68. doi: 10.1038/s41420-020-00305-y

42.   Zheng X, Hermann DM, Bahr M, Doeppner TR. The role of small extracellular vesicles in

cerebral and myocardial ischemia-Molecular signals, treatment targets, and future clinical

translation. *Stem Cells*. 2021;39:403-413. doi: 10.1002/stem.3329

43.   Sichrovsky M, Lacabanne D, Ruprecht JJ, Rana JJ, Stanik K, Dionysopoulou M, Sowton AP,

King MS, Jones SA, Cooper L, et al. Molecular basis of pyruvate transport and inhibition of the

human   mitochondrial   pyruvate   carrier.   *Sci   Adv*.   2025;11:eadw1489.   doi:

10.1126/sciadv.adw1489

44.   He Z, Zhang J, Xu Y, Fine EJ, Suomivuori CM, Dror RO, Feng L. Structure of mitochondrial

pyruvate carrier and its inhibition mechanism. *Nature*. 2025;641:250-257. doi:

10.1038/s41586-025-08667-y

45.   Lesnefsky EJ, Chen Q, Tandler B, Hoppel CL. Mitochondrial Dysfunction and Myocardial

Ischemia-Reperfusion: Implications for Novel Therapies. *Annu Rev Pharmacol Toxicol*.

2017;57:535-565. doi: 10.1146/annurev-pharmtox-010715-103335

46.   Lopez-Vazquez P, Fernandez-Caggiano M, Barge-Caballero E, Barge-Caballero G,

Couto-Mallon D, Grille-Cancela Z, Blanco-Canosa P, Paniagua-Martin MJ, Enriquez-Vazquez

D, Vazquez-Rodriguez JM, et al. Reduced mitochondrial pyruvate carrier expression in hearts

with heart failure and reduced ejection fraction patients: ischemic vs. non-ischemic origin.

Front Cardiovasc Med. 2024;11:1349417. doi: 10.3389/fcvm.2024.1349417

47.   Liu Y, Yuan Y, Yan Y, Wang R, Wang Z, Liu X, Zhang Y, Hua J, Wang Y, Zhao L.

Mitochondrial pyruvate carrier 1 alleviates hypoxic-ischemic brain injury in rats. *Life Sci*.

2023;325:121686. doi: 10.1016/j.lfs.2023.121686

48.   Garcia-Martin R, Brandao BB, Thomou T, Altindis E, Kahn CR. Tissue differences in the

exosomal/small extracellular vesicle proteome and their potential as indicators of altered

tissue metabolism. *Cell Rep*. 2022;38:110277. doi: 10.1016/j.celrep.2021.110277

49.   Das D, Jothimani G, Banerjee A, Dey A, Duttaroy AK, Pathak S. A brief review on recent

advances in diagnostic and therapeutic applications of extracellular vesicles in cardiovascular

disease. *Int J Biochem Cell Biol*. 2024;173:106616. doi: 10.1016/j.biocel.2024.106616

50.   Li S, Shen S, Xu H, Cai S, Yuan X, Wang C, Zhang X, Chen S, Chen J, Shi DL, et al. IGF2BP3

promotes adult myocardial regeneration by stabilizing MMP3 mRNA through interaction with

m6A modification. *Cell Death Discov*. 2023;9:164. doi: 10.1038/s41420-023-01457-3

51.   Tian Y, French BA, Kron IL, Yang Z. Splenic leukocytes mediate the hyperglycemic

exacerbation of myocardial infarct size in mice. *Basic Res Cardiol*. 2015;110:39. doi:

10.1007/s00395-015-0496-3

52.   Ge X, Meng Q, Wei L, Liu J, Li M, Liang X, Lin F, Zhang Y, Li Y, Liu Z, et al. Myocardial

ischemia-reperfusion induced cardiac extracellular vesicles harbour proinflammatory features

and aggravate heart injury. *J Extracell Vesicles*. 2021;10:e12072. doi: 10.1002/jev2.12072

53.   Ge X, Meng Q, Zhuang R, Yuan D, Liu J, Lin F, Fan H, Zhou X. Circular RNA expression

alterations in extracellular vesicles isolated from murine heart post ischemia/reperfusion injury.

Int J Cardiol. 2019;296:136-140. doi: 10.1016/j.ijcard.2019.08.024

54.   Zuo R, Ye LF, Huang Y, Song ZQ, Wang L, Zhi H, Zhang MY, Li JY, Zhu L, Xiao WJ, et al.

Hepatic small extracellular vesicles promote microvascular endothelial hyperpermeability

during NAFLD via novel-miRNA-7. *J Nanobiotechnology*. 2021;19:396. doi:

10.1186/s12951-021-01137-3

55.   Lasser C, Kishino Y, Park KS, Shelke GV, Karimi N, Suzuki S, Hovhannisyan L, Radinger M,

Lotvall J. Immune-Associated Proteins Are Enriched in Lung Tissue-Derived Extracellular

Vesicles during Allergen-Induced Eosinophilic Airway Inflammation. *Int J Mol Sci*. 2021;22. doi:

10.3390/ijms22094718

56.   Crescitelli R, Lasser C, Lotvall J. Isolation and characterization of extracellular vesicle

subpopulations from tissues. *Nat Protoc*. 2021;16:1548-1580. doi:

10.1038/s41596-020-00466-1

57. Jingushi K, Uemura M, Ohnishi N, Nakata W, Fujita K, Naito T, Fujii R, Saichi N, Nonomura N,

Tsujikawa K, et al. Extracellular vesicles isolated from human renal cell carcinoma tissues

disrupt vascular endothelial cell morphology via azurocidin. *Int J Cancer*. 2018;142:607-617.

doi: 10.1002/ijc.31080

58. Chen L, Li S, Zhu J, You A, Huang X, Yi X, Xue M. Mangiferin prevents myocardial

infarction-induced apoptosis and heart failure in mice by activating the Sirt1/FoxO3a pathway.

J Cell Mol Med. 2021;25:2944-2955. doi: 10.1111/jcmm.16329

59. Zielonka J, Kalyanaraman B. Hydroethidine- and MitoSOX-derived red fluorescence is not a

reliable indicator of intracellular superoxide formation: another inconvenient truth. *Free Radic*

Biol Med. 2010;48:983-1001. doi: 10.1016/j.freeradbiomed.2010.01.028

60. Mol EA, Lei Z, Roefs MT, Bakker MH, Goumans MJ, Doevendans PA, Dankers PYW, Vader P,

Sluijter JPG. Injectable Supramolecular Ureidopyrimidinone Hydrogels Provide Sustained

Release of Extracellular Vesicle Therapeutics. *Adv Healthc Mater*. 2019;8:e1900847. doi:

10.1002/adhm.201900847

61. Wisniewski JR, Zougman A, Nagaraj N, Mann M. Universal sample preparation method for

proteome analysis. *Nat Methods*. 2009;6:359-362. doi: 10.1038/nmeth.1322

Figures and Figure Legends

Figure 1. Splenectomy aggravates cardiac injury in AMI mice

A, Twenty-eight-day Kaplan-Meier survival curves for sham operation (Sham) or MI

mice, with or without splenectomy (n=30 mice/group). **B**, Representative echocardiograms from sham, splenectomized sham (Sham+Spl), MI, and splenectomized MI (MI+Spl) mice 7 days post MI (n=6/group). **C** and **D**, Summary of left ventricle EF (LVEF%) and FS (LVFS%) in sham, Sham+Spl, MI, and MI+Spl mice at indicated time-points (n=6/group). **E** and **F**, Representative TTC images of cross-sectional cardiac slices and Quantitation of infarct size in hearts from sham, Sham+Spl, MI, and MI+Spl mice 1 day post MI (n=6 mice per group). **G** and **H,** Representative images of Masson's trichrome staining and quantitation of the fibrosis area in hearts from sham, Sham+Spl, MI, and MI+Spl mice 14 days after MI (n=6/group). **I-J**, Representative western blots quantitation of collagen I, Fibronectin, and α-SMA expression in mice hearts 14 days post MI (n=6/group). **K-L,** Representative TUNEL images and quantitation of TUNEL-positive rates in myocardial tissues from sham, Sham+Spl, MI, and MI+Spl mice 3 days post MI (n=6/group). **M-N,** Representative western blots and quantitation of C-Cas-3, Bax, and Bcl2 in mice hearts 3 days post MI (n=6/group). **O-P**, Representative WGA staining images of cardiac sections and quantitation of cardiomyocyte size in mice hearts 28 days post MI (n=6/group). *, vs Sham; [#], vs MI, $P<0.05$. Statistical analyses were performed using mixed-effects ANOVA followed by the Sidak multiple comparisons test (**C** and **D**), the one-way ANOVA with the Dunnett multiple comparisons test (**H**, **P** and **L**), the Brown-Forsythe and Welch ANOVA test with the Dunnett T3 multiple comparisons test (**F**, **J**; Collagen I and N; Bcl2, C-Cas-3), the one-way ANOVA with the Tukey multiple comparisons test (**J**; α-SMA and N; Bax), the

880   Kruskal Wallis with the Dunn multiple comparisons test (**J**; Fibronectin). Survival

881   curve rate was analyzed by the Kaplan-Meier method and compared by the log-rank

882   test in **A**. Spl indicates splenectomy; EF, ejection fraction; FS, fractional shortening;

883   LV, left ventricular; FN, Fibronectin; Col, Collagen I (COL1A1); and α-TUB, α-Tubulin.

Figure 2. Spleen-derived sEVs biogenesis is enhanced in response to AMI and

these sEVs were enriched in ischemic hearts in mice.

A, Schematic diagram of RNA-Seq for spleens from Sham and MI mice. **B**, Volcano

plot of differentially expressed genes (DEGs) of spleens from sham and MI mice. **C**,

sEVs biogenesis-related DEGs expression in spleen tissues from sham and MI mice

were validated using qPCR 1 day post MI. Data were normalized to GAPDH and

expressed as folds of Sham group (n=6/group). **D,** The flowchart of spleen-derived

sEVs extraction. **E**, Representative western blot images of sEVs markers. **F**, Protein

contents of sEVs (n=6/group). **G**, Representative TEM images of spleen-derived

sEVs from various mice groups as indicated. **H**, Size distributions of Spleen-derived

sEVs measured by nFCM. **I-J**, Detection of the biodistribution of DIR-labeled

spleen-derived sEVs in vivo. Representative heart and whole-body images and were

shown. *, vs Sham, *P*<0.05. Statistical analyses were performed by unpaired t test (**C**;

Rab27a, Rab11, Pdcd6ip and Smpd3), the Mann-Whitney U test (**C**; Rab31 and

Rab22a) and the one-way ANOVA with the Dunnett multiple comparisons test (**F**).

M1D indicates day 1 after MI; M3D, day 3 after MI; M7D, day 7 after MI.

909 **Figure 3. Spleen-derived sEVs from MI mice alleviate ischemia injuries in hearts**

910 **and cardiomyocytes.**

A, Twenty-eight-day Kaplan-Meier survival curves for sham or MI mice treated with indicated sEVs or vehicle (n=30 mice per group). **B-D**, Representative echocardiograms of sham or MI mice treated with indicated sEVs or vehicle 7 days after MI (n=6/group). C-D. Quantitation of LVEF% and LVFS%. **E-F**, Representative images of TTC staining and quantitation of infarct area in mice hearts 1 day after MI (n=6/group). **G-H**, Representative images of Masson's trichrome and quantitation of the scar area in mice hearts 14 days after MI (n=6/group). **I-J**, Representative western blot images and quantitation of cardiac collagen I, Fibronectin, and α-SMA 14 days after MI (n=6 mice per group). **K-L**, Representative western blot images and quantitation of cardiac C-Cas-3, Bax, and Bcl2 3 days after MI (n=6/group). **M-N**, Representative WGA staining images of cardiac sections and quantitation of cardiomyocytes (CMs) 28 days after MI (n=6/group). *, vs Sham; #, vs MI, $P<0.05$. Statistical analyses were performed using mixed-effects ANOVA followed by the Sidak multiple comparisons test (**C** and **D**), the one-way ANOVA with the Dunnett multiple comparisons test (**F** and **H**), the one-way ANOVA with the Tukey multiple comparisons test (**K**; α-SMA, Fibronectin and **L**; Bcl2, C-Cas-3), the Kruskal-Wallis with the Dunn multiple comparisons test (**K**; Collagen I and **L**; Bax), and the Brown-Forsythe and Welch-ANOVA test with the Dunnett T3 multiple comparisons test (**N**). Survival curve rate was analyzed by the Kaplan-Meier method and compared by the log-rank test in **A**. S-sEVs indicates small extracellular vesicles derived from spleens of sham mice; M1D-sEVs, sEVs derived from spleens of MI mice 1 day post MI; M3D-sEVs, sEVs derived from spleens of MI mice 3 day post MI;

933   and M7D-sEVs, sEVs derived from spleens of MI mice 7 day post MI. EF, ejection

934   fraction; FS, fractional shortening; LV, left ventricular; FN, Fibronectin; Col, Collagen I

935   (COL1A1); and α-TUB, α-Tubulin.

Figure 4. sEVs depletion aggravates cardiac ischemic injury in AMI mice.

A, Schematic diagram of experimental design for sEVs depletion. **B**,

Twenty-eight-day Kaplan-Meier survival curves for MI mice treated with indicated

conditions (n=30 mice per group). Log-rank test. **C**, Representative echocardiograms

of AMI mice treated with GW4869 plus S-sEVs or M3D-sEVs, or corresponding

vehicles 7 days after MI (n=6 mice per group). **D-E**, Quantitation of LVEF% and

LVFS%. **F-G**, Representative images of TTC staining of cardiac slices from AMI mice

treated with GW4869 plus S-sEVs or M3D-sEVs, or corresponding vehicles, and

quantitation of infarct area in mice hearts 1 day after MI (n=6/group). **J-K**,

Representative western blot images and quantitation of cardiac collagen I,

Fibronectin, and α-SMA 14 days after MI in mice treated as indicated conditions

(n=6/group). **L-M**, Representative western blot images and quantitation of cardiac

C-Cas-3, Bax, and Bcl2 of hearts 3 days after MI in mice (n=6/group). **N-P**,

Representative WGA staining images of cardiac sections and quantitation of

cardiomyocytes (CMs) 28 days after MI (n=6/group). *, vs MI; #, vs MI+GW4869,

$P<0.05$. Statistical analyses were performed using mixed-effects ANOVA followed by

the Sidak multiple comparisons test (**D** and **E**), the one-way ANOVA with the Dunnett

multiple comparisons test (**G**), the Brown-Forsythe and Welch-ANOVA test with the

Dunnett T3 multiple comparisons test (**I** and **M**; Bax), the Kruskal-Wallis with the

Dunn multiple comparisons test (**M**; C-Cas-3 and **P**), and the one-way ANOVA with

the Tukey multiple comparisons test (**K** and **M**; Bcl2). Survival curve rate was

analyzed by the Kaplan-Meier method and compared by the log-rank test in B. EF,

ejection fraction; FS, fractional shortening; LV, left ventricular; FN, Fibronectin; Col,

Collagen I (COL1A1); and α-TUB, α-Tubulin.

Figure 5. The proteomic profiling of spleen-derived sEVs between sham and

AMI mice 3 days post ischemia.

A, The flowchart of proteomics experiment for spleens from sham and AMI mouse. **B**,

Volcano plot showing the differentially expressed proteins in spleens between sham

and AMI mouse. **C-E**, GO enrichment analysis of differentially expressed proteins. **F**,

KEGG enrichment analysis of differentially expressed proteins. BP, Biological

Process. CC, Cellular Component. MF, Molecular Function.

Figure 6. Spleen-derived sEVs from MI mice improve myocardial mitochondrial

energy metabolism.

A, Representative TEM images showing mitochondrial ultrastructure of heart tissues from sham and MI mice treated with S-sEVs, M3D-sEVs, or vehicles. **B**, Quantitation of mitochondrial size in these TEM images as indicated groups, being presented as relative value to the sham group (n=6/group). **C-F**, Representative western blotting images and quantitation of ETC complex subunits expression in heart tissues from sham and MI mice treated with S-sEVs, M3D-sEVs, or vehicles for 7 days, and in normal control cells (NC) and OGD-insulted H9C2 cells treated with S-sEVs, M3D-sEVs, or vehicles for 12h (n=6/group). **G-K**, mRNA expressions of ETC-coded genes in the heart tissues from mice treated with as above indicated conditions (n=6/group). **L-Q**, mRNA expressions of ETC-coded genes in H9C2 cells treated as above indicated conditions (n=6/group). **R**, ATP/ADP ratios of NC and OGD-insulted H9C2 cells treated S-sEVs, M3D-sEVs, or vehicles for 24h (n=6/group). **S**, Real-time monitoring OCR in H9C2 cells treated NC and OGD-insulted H9C2 cells treated S-sEVs, M3D-sEVs, or vehicles for 12h (n=6/group). **T-U**, Representative MitoSOX staining images and quantitation of the MitoSOX fluorescent intensity in NC and OGD-insulted H9C2 cells treated S-sEVs, M3D-sEVs, or vehicles for 12h(n=6/group). *, vs Sham or NC, $P<0.05$; #, vs MI or OGD, $P<0.05$. Statistical analyses were performed by the one-way ANOVA with the Dunnett multiple comparisons test (**B** and **R**), the one-way ANOVA with the Tukey multiple comparisons test (**E**; ATP5A, MTCO1, NDUFB8, UQCRC2, **F**;NDUFB8, **G**; Sdhb, **H**, **I**; Uqcrc1, **J**, **K**; Atp5po, **L**; Sdhb, **M**; Ndufs2, **N** and **P**; Cox4il), the Brown-Forsythe and Welch-ANOVA test with the Dunnett T3 multiple comparisons test (**E**; SDHB, **L**; Sdhc, **P**; Cox7b, **Q** and **U**),

1001 and the Kruskal-Wallis with the Dunn multiple comparisons test (**F**; ATP5A, MTCO1,

1002 SDHB, UQCRC2, **G**; Sdhb, **I**; Uqcrc1, **K**; Atp5f1b and **M**; Ndufs1).

1003

1004

Figure 7. MPC1 protein expression is increased in M3D-sEVs from AMI mouse

and in circulating plasma sEVs from AMI patients.

A, Veen diagrams of differentially expressed proteins in spleen-derived sEVs and

plasma between sham and AMI mouse. **B-C**, Representative western blotting images

and quantitation of MPC1 and PCK2 expression inside sEVs from sham and AMI

mouse. Unpaired two-tailed Welch t-test. **D**, Quantitation results of ELISA analysis

determining MPC1 expression in organ tissues-derived sEVs after MI. unpaired t test.

E, Schematic diagram of plasma sEVs (P-sEVs) isolation from patients. **F**,

Representative TEM images of P-sEVs. **G**, Size distributions of P-sEVs measured by

nFCM. **H**, Immunoblot determination of sEVs markers. **I**, ELISA assay results of the

plasma MPC1 expression from patients. **J**, ELISA assay results of MPC1 expression

in P-sEVs. *, vs S-sEVs or No CHD, $P<0.05$. #, vs CHD no AMI, $P<0.05$. Statistical

analyses were performed by the unpaired two-tailed Welch t-test (**C**; MPC1), the

unpaired t test (**C**; PCK2 and **D**), the Kruskal-Wallis with the Dunn multiple

comparisons test (**I**) and the one-way ANOVA with the Dunnett multiple comparisons

test (**J**). Ti-sEVs indicates tissue sEVs.

1021

1022

Figure 8. Blockage of MPC1 blunts the protective effects of spleen-derived M3D-sEVs against ischemic cardiac damage and disruption of myocardial mitochondrial energy metabolism.

A, Twenty-eight-day Kaplan-Meier survival curves for AMI mouse treated with M3D-sEVs (sEVs), MPC1 antibody-neutralized M3D-sEVs (sEVs+Ab), or Ig G-incubated M3D-sEVs (sEVs+IgG), n=30/group. Log-rank test. **B**, Representative echocardiograms of AMI mice treated with indicated conditions 7 days after MI (n=6/group). **C-D**, Quantitation of LVEF% and LVFS%. **E-F**, Representative TTC staining images of cross-sectional cardiac slices from AMI mice treated as above indicated 1 day post MI, and quantitation of infarct area percentage (n=6/group). **G-H**, Representative western blotting images and quantitation of cardiac collagen I, Fibronectin, and α-SMA expression in AMI mice treated as above indicated 14 days after MI (n=6/group). **I-K**, Representative images of Masson's trichrome (top) and WGA staining (bottom) and quantitation of the hearts from AMI mice treated as above indicated 14- or 28-days post MI, respectively (n=6 mice per group). **L-P**, mRNA expression of ETC-coded genes in the OGD-insulted H9C2 cells treated with M3D-sEVs or M3D-sEVs plus MPC1 inhibitors (n=6/group). **Q**, Representative TEM images showing mitochondrial ultrastructure of heart tissues from AMI mice treated as above indicated. **R**, Quantitation of mitochondrial size in TEM images as indicated groups, being presented as relative value to the M3D-sEVs group (n=6/group). **S**, Quantitation of ATP/ADP ratio of OGD-H9C2 cells treated as indicated (n=6/group). **T**, Results of OCR assay in H9C2 cells treated as indicated. Statistical analyses were performed using mixed-effects ANOVA followed by the Sidak multiple comparisons test (**C** and **D**), the one-way ANOVA with the Dunnett multiple comparisons test (**F**, **J** and **S**), the one-way ANOVA with the Tukey multiple comparisons test (**H**; Collagen I,

L; Ndufs2, **M**, **N**, **O** and **P**), the Brown-Forsythe and Welch-ANOVA test with the

Dunnett T3 multiple comparisons test (**R**), and the Kruskal-Wallis with the Dunn

multiple comparisons test (**H**; α-SMA, Fibronectin, **K**, **L**; Ndufs1). The survival curve

rate was analyzed by the Kaplan-Meier method and compared by the log-rank test in

A. UK, UK5099 (a MPC1 inhibitor). GWx, GW604714x (an MPC1 inhibitor). *, vs

sEVs or M3D-sEVs; [#], vs sEVs+IgG, *P*<0.05. UK indicates UK5099; GWx,

GW604714x. EF, ejection fraction; FS, fractional shortening; LV, left ventricular; FN,

Fibronectin; Col, collagen I (COL1A1); and α-TUB, α-Tubulin.