Orco regulates the circadian activity of pheromone-sensitive olfactory

receptor neurons in hawkmoths

Abbreviated title: Circadian regulation by Orco

Author names and affiliations:

* Aditi Vijayan, Animal Physiology and Center for Interdisciplinary Nanostructure Science and

Technology, FB 10, University of Kassel, 34132 Kassel, Germany

* Mauro Forlino, Theoretical Physics and Center for Interdisciplinary Nanostructure Science and

Technology, FB10, University of Kassel, 34132 Kassel, Germany

Yajun Chang, Animal Physiology and Center for Interdisciplinary Nanostructure Science and

Technology, FB 10, University of Kassel, 34132 Kassel, Germany

Pablo Rojas, Theoretical Physics and Center for Interdisciplinary Nanostructure Science and

Technology, FB10, University of Kassel, 34132 Kassel, Germany

Katrin Schröder, Animal Physiology and Center for Interdisciplinary Nanostructure Science and

Technology, FB 10, University of Kassel, 34132 Kassel, Germany

Anna C. Schneider, Animal Physiology and Center for Interdisciplinary Nanostructure Science and

Technology, FB 10, University of Kassel, 34132 Kassel, Germany

Martin E. Garcia, Theoretical Physics and Center for Interdisciplinary Nanostructure Science and

Technology, FB10, University of Kassel, 34132 Kassel, Germany

Monika Stengl, Animal Physiology and Center for Interdisciplinary Nanostructure Science and

Technology, FB 10, University of Kassel, 34132 Kassel, Germany

*Both authors contributed equally

Corresponding author email address: anna.c.schneider@uni-kassel.de

Conflict of interest: Authors report no conflict of interests.

Acknowledgements: We thank Dr. Dieter Wicher (Max Planck Institute for Chemical Ecology, Jena,

Germany) for generously providing OLC15, and Rishaban Radhakrishnan for help with Python

codes. We thank Prof. Hülya Altuntaş for helpful discussions.

Funding: AV, MF, YC, ACS, MEG, and MS were supported by Deutsche Forschungsgemeinschaft

RTG 2749/1: "Biological Clocks on Multiple Time Scales"; KS and MS were supported by Deutsche

Forschungsgemeinschaft STE531/20-2.

Author contributions:

Conceptualization: AV, MF, MEG, MS; data curation: AV, MF, YC, PR, KS, ACS, MEG, MS; formal

analysis: AV, MF, YC, PD, ACS; funding acquisition: MEG, MS; investigation: AV, MF, YC, KS;

methodology: MF, PR, MEG, MS; project administration: ACS, MEG, MS; resources: MEG, MS;

software: AV, MF, PR, MEG; supervision: MEG, MS; validation: AV, MF, YC, PR, KS, ACS, MEG, MS;

visualization: AV, MF, YC, ACS, MS; writing – original draft: AV, MF, ACS, MEG, MS; writing –

reviewing & editing: AV, MF, YC, ACS, MEG, MS.

1 Abstract

The mating behavior of nocturnal *Manduca sexta* hawkmoths is under strict temporal control. It is orchestrated via circadian- and ultradian oscillations in sex-pheromone stimuli as social zeitgeber. The extremely sensitive pheromone-detecting olfactory receptor neurons (ORNs) that innervate the long trichoid sensilla on the male's antennae are peripheral circadian clocks. They express the transcriptional-translational feedback loop (TTFL) circadian clockwork, best characterized in *Drosophila melanogaster*. In hawkmoths, it is still unknown whether or how the ORN's TTFL clockwork regulates the daily rhythms in pheromone-sensitivity and in temporal resolution of ultradian pheromone pulses as prerequisites to the temporal regulation of hawkmoth mating behavior.

We hypothesize that, rather than the slow TTFL clock, a more rapidly adaptive post-translational feedback loop (PTFL) clockwork, associated with the ORN's plasma membrane, allows for temporal control of pheromone detection via generation of multiscale endogenous membrane potential oscillations. The potential oscillations of the PTFL clock could rapidly synchronize to oscillations of pheromone stimuli at different timescales, thus enable the prediction of stimulus patterns as a mechanism for active sensing. With *in vivo* long-term tip recordings of long trichoid sensilla of male hawkmoths, we analyzed the spontaneous spiking activity indicative of the ORNs' endogenous membrane potential oscillations. Consistent with our hypothesis of a multiscale PTFL clock in hawkmoth ORNs, spontaneous spiking was modulated on ultradian and circadian timescales, with maximum activity at night. When we blocked the evolutionarily conserved olfactory receptor coreceptor (Orco), the circadian modulation was abolished but the ultradian frequency modulation of the spontaneous activity remained. Consistent with PTFL control, Orco was not under the circadian control of the TTFL clock. We could replicate the experimental data in a conductance-based computational model of Orco. In this model, Orco conductivity changed as a function of fluctuating 2^{nd} messenger levels. This study demonstrates that a PTFL is sufficient to impose a circadian pattern on ORN sensitivity.

2 Significance statement

It is generally assumed that all circadian rhythms in an organism are driven by a transcriptional-translational feedback loop (TTFL) clock. In this study, we demonstrate with *in vivo* recordings of hawkmoth pheromone-sensitive olfactory receptor neurons (ORNs) that the olfactory

receptor coreceptor (Orco) is the key pacemaker channel for controlling circadian but not ultradian

rhythms in spontaneous spiking activity. Since Orco expression is not driven by the TTFL clock, its

conductance is rather controlled by a post-translational feedback loop (PTFL) membrane clock.

Accordingly, our computational model suggests that periodic changes in the conductivity of an

Orco ion channel, which is mediated by cycling levels of cyclic nucleotides, tune ORN sensitivity.

This highlights the role of the contribution of posttranslational modifications to the generation of

circadian rhythmicity.

3 Introduction

Biological clocks generate endogenous physiological and behavioral oscillations on multiple

timescales that persist in the absence of any external timing cues under constant conditions

(Lamont and Amir, 2010). Such clocks are essential to generate internal rhythms and to permit

their synchronization to external zeitgebers at timescales from milliseconds to years. These

synchronized endogenous multiscale clocks thus provide both a continuous time frame for

physiology and behavior - the organism's "present time" - and enable organisms to predict and

respond to both regular, periodic environmental changes (Stengl and Schneider, 2024). Even

though endogenous clocks are widespread in prokaryotes and eukaryotes (Géron et al., 2024; Hall

and Rosbash, 1993; Millar, 2016; Thaiss et al., 2014), current knowledge about how multiscale

clocks are linked over widely diverging timescales remains sparse.

Insect circadian clock neurons, which are tightly connected to visual brain circuits, are

amongst the best studied biological clocks (Hall and Rosbash, 1993; Nishiitsutsuji-Uwo and

Pittendrigh, 1968; Reinhard et al., 2024). They regulate physiological and behavioral oscillations

that underly sleep-wake rhythms tied to the 24 h light-dark cycles on Earth (Dubowy and Sehgal,

2017). Circadian clock neurons possess a molecular clockwork of transcriptional-translational

feedback loops (TTFLs), which generate endogenous oscillations of clock gene mRNA and protein

levels with a cycle period of ~24 h (Hardin, 2011). It is assumed for both insects and mammals that

this TTFL comprises the master clockwork that regulates all physiological and behavioral rhythms

in the circadian time range through genetically controlled output pathways (Hastings et al., 2020).

Molecular TTFL clockworks were not only found in central brain clock neurons but also in

sensory neurons, such as olfactory receptor neurons (ORNs) in insect antennae, where they play a

role in odor-dependent circadian behaviors like feeding, pollination, reproduction, and navigation

(Flecke et al., 2010; Flecke and Stengl, 2009; Krishnan et al., 1999; Merlin et al., 2009, 2006; Page

and Koelling, 2003; Plautz et al., 1997; Sauman and Reppert, 1998; Schendzielorz et al., 2015;

Schuckel et al., 2007; Tanoue et al., 2004; Zhou et al., 2005). Little is known about how the TTFL

clockwork of insect ORNs regulates circadian rhythms in chemosensory sensitivity that underlie

the circadian control of these different behaviors.

At night, male nocturnal *Manduca sexta* hawkmoths are very sensitive to the sex pheromone

blend that their conspecific females release in an intermittent pattern (Itagaki and Conner, 1988).

The intermittency is a prerequisite to maintain the male's search flight, which is suspended at

pulse frequencies >30 Hz (Baker et al., 1985; Baker and Vogt, 1988; Stengl, 2010). The pheromone

pulses are resolved of up to 3 Hz by pheromone-specific ORNs in the antenna's long trichoid

sensilla (Lee and Strausfeld, 1990; Marion-Poll and Tobin, 1992). Pheromone-specificity is provided

by odor receptor (OR) subunits in the ciliary membranes of ORNs that heteromerize with the

conserved olfactory receptor co-receptor Orco (Nakagawa and Vosshall, 2009; Stengl and Funk,

2013; Wicher and Miazzi, 2021). While the various roles of Orco are still under debate, it is known

that in heterologous expression systems Orco homomers form a slow, leaky, non-specific cation

channel, which depolarizes the cells (Jones et al., 2011; Sargsyan et al., 2011; Sato et al., 2008;

Wicher et al., 2008). *In vivo* tip recordings of long trichoid sensilla in *M. sexta* confirmed that Orco

acts as spontaneously opening cation channel, acting as a pacemaker channel, because it

promotes ultradian membrane potential oscillations that underly the spontaneous action potential

activity (Nolte et al., 2013). These potential oscillations turn the ORNs into ultradian temporal

filters that are sensitive to the frequency of the pheromone pulses they encounter. Pheromone

pulses that arrive at a similar frequency as the endogenous sub-threshold membrane potential

oscillations would synchronize these oscillations and would be more likely to elicit action potential

trains. Therefore, this pacemaker channel-dependent mechanism would allow for active sensing if

endogenous membrane potential oscillations are in the physiological range necessary for tracking

intermittent pheromone pulses.

It is not known whether the Orco-dependent membrane potential oscillations also occur at

circadian timescales, and whether or how they are linked to ultradian potential oscillations. One

hypothesis is that the molecular TTFL clockwork of ORNs exclusively controls endogenous

circadian rhythms of the membrane potential via circadian expression of the pacemaker channel

Orco. Alternatively, the plasma membrane of ORNs could by itself constitute an autonomous clock

that is entrained by but does not rely on the TTFL clock. Constant levels of ion channels that do not

require daily degradation and daily transcription via the TTFL clock, but with modulation of open-

time probability by posttranslational modifications, would constitute a more economical and

faster adjustable mechanism to generate endogenous membrane potential oscillations (Stengl and

Schneider, 2024). This would facilitate active sensing, the rapid tuning to detect behaviorally

relevant zeitgeber signals in the insect's environment.

Here, we performed a combination of experimental and computational modeling studies to

investigate whether Orco plays a role in active sensing at the circadian and/or ultradian timescale

via generation of endogenous membrane potential oscillations. We hypothesize that Orco is part of

a post-transcriptional feedback loop (PTFL) clockwork in the plasma membrane, rather than being

TTFL controlled. To capture naturally occurring, physiologically relevant processes in insect ORNs

we performed *in vivo* long-term tip recordings of the spontaneous spiking activity of pheromone-

sensitive trichoid sensilla of intact but restrained hawkmoth males.

We corroborated the experimental data with a Hodgkin-Huxley (HH) based computational

ORN model. In general, the widely used deterministic HH models do not include circadian control

of ion channels. Moreover, the highly dynamic, bursting firing patterns of ORNs cannot be

described by the traditional deterministic modeling of neuronal activity. This constitutes a

drawback because stochasticity plays a crucial role in shaping firing patterns which are influenced

not only by external stimuli but also by the inherent noise involved in channel gating. Therefore, to

address the above points realistically, we developed a Langevin formulation of the HH model,

which explicitly accounts for the noise introduced by the random opening and closing of ion

channels and circadian modulation of the Orco channel conductivity. By integrating channel noise,

we could explore how circadian regulation of ionic currents contributes to the variability of firing

patterns and how this noise interacts with external stimuli to influence olfactory signal processing.

Our model aims to highlight the advantages of using a stochastic framework for modeling the

circadian activity of ORNs. The combined experimental and modeling efforts generate a more

complete picture and predictions concerning these autonomous processes in sensory neurons,

which are required for future exploration of physiologically relevant anticipatory behaviors that are

driven by biological clocks.

4 Methods

4.1 Animals

Nocturnal *M. sexta* hawkmoths were bred and raised from eggs in the rearing facility at the

University of Kassel. Males and females were separated during pupal stages. Males were housed

isolated from the females to avoid exposure to female pheromones. All experiments were

performed with adult males reared in a long-day photoperiod (L:D 17:7 h to prevent diapause) at

25 °C with relative humidity of about 55%. Larvae were fed with an artificial diet modified after Bell

and Joachim (1976); adult moths could feed on sugar solution with added vitamins (Roth) *ad*

libitum (Riffell et al., 2008).

4.2 Solutions

Ringer compositions were taken from (Pézier et al., 2007) and contained (in mM): 6.4 KCl, 12

$MgCl_2$, 1 $CaCl_2$, 12 NaCl, 10 HEPES, 340 glucose for hemolymph ringer (HLR) at 450 mosmol/kg,

and 172 KCl, 3 $MgCl_2$, 1 $CaCl_2$, 25 NaCl, 10 HEPES, 22.5 glucose for sensillum lymph ringer (SLR) at

475 mosmol/kg, both at pH 6.5. The Orco antagonist OLC15 (N-(4-butylphenyl)-2-((4-ethyl-5-(2-

pyridinyl)-4H-1,2,4-triazol-3-yl)thio)acetamide (Chen and Luetje, 2012); kindly provided by Dr. D.

Wicher, Max Planck Institute for Chemical Ecology, Jena, Germany) was dissolved in DMSO to a

concentration of 100 mM (stock solution stored at 4-8 °C) before being diluted in SLR to a final

concentration of 50 μM immediately before the start of the recording. DMSO percentage in the final

solution was 0.05%.

4.3 Electrophysiology

The spontaneous electrical activity of the unstimulated pheromone-sensitive ORNs in a

single long trichoid sensillum was recorded extracellularly as *in vivo* tip recording (Kaissling et al.,

1987). 1-2-d-old male *M. sexta* that were raised in isolation from females and pheromone exposure

were fixed in a custom-made Teflon holder with adhesive tape. The right antenna used for

recordings was oriented with its dorsal surface facing up and immobilized near the base with

dental wax (Boxing wax strips, KerrHawe SA). Electrodes (Ag/AgCl wire in a glass pipette filled with

SLR for the recording electrode and HLR for the reference electrode) were pulled (DMZ-Universal

Puller, Zeitz Instruments) from borosilicate capillaries (OD:1.56mm, ID: 1.17mm, without filament;

Science Products) to a tip diameter of about 2 μm. After cutting off 20 distal annuli of the antenna

with micro scissors, the reference electrode was inserted into the antennal lumen up to the second

annulus from the cut end. The cut was covered with electrode gel (GE Medical Systems Information

Technology) to prevent desiccation. Electrodes were moved with manual micromanipulators (MMJ,

Märzhäuser Wetzlar; and Leitz, Leica Microsystems) or with motorized micromanipulators

(Microstar, Scientifica) with a joystick controller (Scientifica). Under microscopic control, some

tips of the long trichoid sensilla of the distal second annulus were cut off with sharpened Dumont

forceps, and the recording electrode slipped over a single sensillum of an upper sensillar row.

ORN activity was amplified 200-fold (custom built amplifier with $10^{12}\,\Omega$ input impedance, or

ELC-01 MX in combination with DPA-2FX, npi, or EXT 10-2F, npi), low pass filtered at 1.3 kHz,

digitized at 20 kHz with a Digidata (1550A or 1550B, Molecular Devices), and saved for offline

analysis with Clampex 10 (pCLAMP suite, Molecular Devices).

All tip recording experiments were performed at room temperature (21-27 °C) and a relative

humidity of 35-60%. We recorded ORN activity either in long-day conditions (17:7 LD) or constant

darkness (DD) to examine the free-running peripheral ORN clock. For LD experiments, the light

regime continued as in the rearing facility during the experiment. For DD and OLC15 experiments,

the lights were turned off at the beginning of the recording, i.e. all animals were exposed to the

same zeitgeber until the beginning of the experiment. To examine the effect of Orco on the

spontaneous action potential (AP) activity pattern, we used the recording electrode to infuse

OLC15 in DD conditions.

4.4 Data analysis

To characterize the spontaneous ORN spiking activity, we first detected spikes with a self-

written program in Python. Using the scipy.signal package and find_peaks function, we applied a

Bessel filter to the raw voltage trace to facilitate spike detection within the following conditions:

spike full width (3-15 ms), threshold search for amplitude of spike (0.5-2.5 mV) and threshold

prominence (0.1). Two or more consecutive spikes with intervals ≤ 50 ms were considered a burst

(Dolzer et al., 2001). We then calculated the following attributes in bins of either 10 min or 1 h for

further analysis: instantaneous spike frequency (1 / inter-spike interval (ISI)), inter-burst interval

(IBI, time between two consecutive bursts with no single spikes in between), inter-event interval

(IEI, time between two events, i.e. two consecutive single spikes, two consecutive bursts, a burst

and a consecutive spike, and vice versa), burst duration (time from first to last spike in a burst),

percentage of all spikes in the experiment that are members of burst, and mean number of spikes

per burst (Figure 1). All statistics were done with those binned values.

Hawkmoth behavioral and physiological rhythms desynchronize across the population in the

absence of zeitgebers like pheromone stimulation and light-dark cycles. Therefore, it was

necessary to phase align the recordings obtained in DD and OLC15. To achieve this, we used an

optimization procedure to calculate the optimal temporal lags for each dataset, aiming to

maximize the overall cross-correlation between the calculated binned spike frequency series. The

optimization was performed using simulated annealing, which provided a set of time shifts (lags)

for each series. These lags were applied to shift the frequency data in a manner that maximized the

synchronization between the different subjects' data, resulting in a "subjective ZT" with subjective

ZT 0 at the first maximum of the binned ISI for each individual. The objective function used for

optimization was the negative sum of pairwise correlations between all series after applying the

shifts. The resulting alignment allowed us to compare the frequency patterns across the different

conditions.

Once the data were aligned, we computed the mean frequency across all subjects for each

bin and each condition. Then, the first high activity period was identified by finding the maximum in

the mean frequency series. Subsequently, we identified the second high activity period as the time

24 hours after the first one and the first low activity period was found by locating the minimum in

the mean frequency series between the two high activity periods.

Besides the mean frequency, we also searched for significant oscillations in IBI, IEI, burst

duration, percentage of spikes that are part of burst, and mean number of spikes per burst. For this

we found the maximum value of these statistics within a 3 h range centered in the first low activity

period and compared it with the values at the second high activity period.

Continuous wavelet and Fourier analysis were performed using PyBoat (Mönke et al., 2020)

to examine the time series of neuronal firing frequency. The continuous wavelet method allows us

to detect and characterize rhythmic patterns within the data by identifying frequency components

that change over time. Specifically, we used wavelet analysis to determine whether the neuron

exhibits circadian and ultradian rhythms. Unlike traditional Fourier analysis, which provides an

overall frequency composition but assumes stationarity, wavelet analysis is particularly useful for

biological signals because it can reveal how these frequencies evolve dynamically. By applying this

approach, we could assess not only the presence of ultradian rhythms but also whether their

strength or occurrence fluctuated in a circadian manner, indicating modulation by the 24-hour

cycle.

4.5 Real-time quantitative polymerase chain reaction (qPCR)

We followed the qPCR methodology detailed in (Schneider et al., 2025). Briefly, male moths

were isolated and cultured until the second day after eclosion, and the antennae from four

individuals were collected at each zeitgeber time (ZT) 1, ZT9, and ZT17 using liquid nitrogen, with

three biological replicates per time point. Total RNA was extracted using TRIzol® Reagent

(Invitrogen) and quantified for concentration and purity with the NanoDrop ND-1000

spectrophotometer (Thermo Fisher Scientific). First-strand complementary DNA (cDNA) was

synthesized using the PrimeScript™ RT reagent Kit with gDNA Erase (Takara Bio Inc.). Primers (F:

AAGCACGTGGTCAGATTGGT, R: GTAAGCCAGGAGCGTCAGAG) for *Orco* (Gene ID: LOC115452348)

were designed using the Primer-BLAST online program of NCBI. The qPCR was conducted following

the protocol of TB Green® Premix Ex Taq™ II (Tli RNase H Plus) (Takara Bio Inc.), with the program:

95 °C for 30 s, followed by 40 cycles of 95 °C for 5 s, and 60 °C for 30 s. Relative expression levels

of *Orco* were calculated using the $2^{-\Delta\Delta Ct}$ method (Livak and Schmittgen, 2001).

4.6 Statistics

All statistics were done with self-written python codes, SigmaPlot 12 (Systat Software), or

JASP (version 0.19.03, University of Amsterdam) with a significance level α = 0.05. Unless noted

otherwise, data were checked for normal distribution with Anderson-Darling or Shapiro-Wilk tests

and for equal variance with Levene's test. If one of these assumptions failed, we used non-

parametric tests, otherwise we used parametric tests.

To assess the statistical significance of observed differences in neuronal activity of the

aligned data between ZTs, we applied the Wilcoxon signed-rank test to pairwise comparisons

between the first low activity period and the subsequent high activity period and discarded the

transient high activity at the beginning of most recordings.

For other data, we used an ANOVA on ranks with either Tukey's or Dunn's post-hoc test for

multiple comparisons as indicated.

4.7 ORN Model

In this study, we introduce a single-compartment Hodgkin-Huxley (HH) type model of the

spontaneous activity of *M. sexta* ORNs. The model contains 5 sets of ion channels that were

previously identified in electrophysiological recordings (Dolzer et al., 2021): the fast sodium (Na^+)

and potassium (K^+) channels responsible for the generation of spikes, low voltage-activated

calcium channels (LVA) responsible for the initiation of bursts, calcium-gated potassium channels

(BK) responsible for the termination of bursts, and the Orco channel, which is a voltage-

independent non-specific, cAMP-gated channel. A schematic of the model is provided in Figure 2A.

We modeled Orco as a ZT-dependent conductance to represent the effect of the oscillations in

cAMP concentration that were identified in these cells (Schendzielorz et al., 2014) (Figure 2B, C).

For simplicity we chose to use a sine function to directly represent the oscillations in Orco

conductivity, but similar results can be achieved by modeling it as an output of a Goodwin model of

the circadian clock via cAMP concentration.

Each type of ion channel possesses one or more gates that can open and close individually

with transition rates α and β for opening and closing, respectively, of activation gates (n, m) and

inactivation gates (h). As a result, each possible combination of open and closed gates defines a

possible state in which each ion channel can be found. A schematic of the Markov chain depicting

all possible states and transitions for each ion channel is shown in Figure 2D. Each ion channel can

conduct current only in the state where all its gates are open.

First, we constructed a deterministic 22-dimensional model to serve as the mean field of the

Langevin model following the methodology proposed by Pu and Thomas (2020). Here, one

dimension tracks the membrane potential, another the intracellular Ca^{2+} concentration, and the

remaining 20 dimensions track the number of channels that are in a specific state of the Markov

chains. Considering the conservation of the total number of ion channels, each type of ion channel

required one less dimension than the total possible states.

The complete Langevin model adds to the mean field an independent noise source for each

transition in the Markov chain. These noise sources are biophysically motivated since the transition

between ion channel states is a stochastic process, with the probability of transition determined by

the kinetics of each kind of ion channel. The variability in the spiking behavior of a neuron emerges

from this non-deterministic gating of individual ion channels. The detailed mathematical

description of the model can be found in the Supplementary Material.

5 Results

We tested the hypothesis that sensory neurons, such as insect ORNs, perform active,

anticipatory sensing based upon an endogenous, plastic plasma membrane clock with the

pacemaker channel Orco as the core element. We focused on pheromone-sensitive long trichoid

sensilla of male *M. sexta* hawkmoth antennae which exhibit daily rhythms in pheromone sensitivity

and temporal resolution that are controlled by a circadian clock. We performed minimally invasive

in vivo tip-recordings over the course of several days to search for daily rhythms in spontaneous

spiking activity as a measure of endogenous membrane potential oscillations.

5.1 Pheromone-sensitive hawkmoth ORNs exhibit Orco-dependent circadian rhythms in spontaneous spiking activity

The ORNs that innervate the pheromone-sensitive long trichoid sensilla on male *M. sexta*

antennae display spontaneous spiking patterns of irregular bursts with interspersed single action

potentials in short-term recordings (Dolzer et al., 2001). In the present study we documented the

same pattern in long-term recordings over the course of several days. This enabled us to search for

daily and circadian rhythmicity in spontaneous ORN spiking activity in the absence of pheromone

stimulation.

The spontaneous spiking patterns revealed clear daily rhythms in 5 of 11 animals under long-

day conditions (17:7 h LD) (Figure 3A). The maxima in spontaneous activity occurred mostly during

the dark phase when nocturnal hawkmoths are active (Figure 3A (top), Bi light blue arrow), and

minima occurred during the light phase when hawkmoths rest/sleep (Figure 3A (bottom), Bi dark

blue arrow). High spiking activity in the dark phase had a mean maximum frequency of ~3 Hz (theta

range: 0.5 – 4 Hz) (Figure 3Bii). In contrast, the minimum spiking frequencies during the

hawkmoth's resting phase were below 0.5 Hz (Figure 3A (bottom)). Since ORN firing patterns had a

clear daily rhythm in LD both in individual animals (Figure 3Bi) and averaged across animals (Figure

3Bii, Biii), we examined whether these rhythms depended on cycling light cues or whether they

persisted in constant darkness (DD), as expected for an endogenous, circadian clock-driven

rhythm.

In DD, the spontaneous ORN firing pattern remained rhythmic (7 of 10 animals), albeit with

less robustness: The single peak of high spiking activity that was observed at the same ZT across

several days in LD divided into multiple bouts of high spiking activity in the activity phase in 7 of 10

animals (Figure 3Ci). Also, in DD conditions, the high spiking activity phase shifted into the

subjective day due to the individual endogenous circadian periods of about 23.5 h +/- 2.8h (21.27 h

in Figure 3Ci). Thus, averaging the instantaneous spiking activity across all DD animals did not

depict clear circadian rhythms (Figure 3Cii). Therefore, we phase-aligned the mean frequency of

spontaneous activity (see Methods). After alignment, circadian rhythms were exposed across DD

animals (Figure 3Ciii), comparable to the daily rhythms in LD (cf. Figure 3 Biii and Ciii).

Next, we examined whether Orco, a leaky, non-specific cation channel, is the dominant

depolarizing pacemaker current of ORNs that drives circadian rhythms in spontaneous spiking

activity. Infusion of the Orco antagonist OLC15 into the sensillum lymph obliterated circadian

rhythms and attenuated the spontaneous activity in several, but not all experiments (N = 8 of 12)

(Figure 3Di). This attenuation resulted in a linear decrease in spiking activity over several days

(Figure 3Dii, Diii). Even after phase-aligning the individual experiments, no circadian rhythm was

present in the average across all OLC15-treated animals (Figure 3Diii).

Because OLC15 is not membrane-permeable on its own it was infused with low

concentrations (0.05%) of DMSO through the recording electrode into the sensillum lymph and,

therefore, effectiveness increased over time. Hence, we fit the spiking activity of each animal and

in each condition with linear regression lines over the whole recording time to compare the slopes

of the fits (Figure 3E). In LD, the mean ± SD slope was 0.004 ± 0.006 Hz/h, indicating that the

average spiking activity remained similar throughout the duration of the recording. In DD, the slope

was -0.003 ± 0.018 Hz/h and was not different from LD. In contrast, the mean slope during OLC15

treatment was -0.007 ± 0.022 Hz/h, significantly smaller than in LD (ANOVA on ranks with Dunn's

post-hoc test; $H(2) = 7.681$, $p = 0.021$). The negative slope corroborated the finding that spiking

activity decreased over time in the presence of the Orco blocker.

Activity rhythms between hawkmoths desynchronize and have only weak phase coupling in

the absence of pheromone stimulation, even in LD cycles (Ghosh et al., 2024). Therefore, instead

of comparing fixed ZT or circadian time (CT) intervals, we compared different attributes of

spontaneous activity between periods of high and low activity, for DD conditions after phase

alignment (see Methods; Figure 4). Despite the high variability in ORN spiking between individuals,

the mean frequency of spontaneous ORN activity was significantly higher during the high activity

phase than during low activity in both LD and DD recordings (Figure 4Ai, Aii). In LD but not in DD,

the frequency of bursting increased significantly during the high activity phase (Figure 4Bi, Bii). The

percentage of spikes that are part of bursts was less variable and lower during the high activity

phase in LD but not significantly different in DD (Figure 4Ci, Cii). The mean burst duration was less

variable during the high than the low activity phase in LD, but not significantly shorter in any

condition (Figure 4Di, Dii). Furthermore, both in LD and DD, the mean inter-event frequencies were

significantly higher during the high activity phase (Figure 4Ei,Eii), indicating a significantly higher

overall spontaneous spiking activity during the hawkmoth's activity phase. The mean number of

spikes per burst, and intra-burst frequencies did not differ significantly between high or low activity

periods in both LD and DD conditions (Figure 4Fi, Fii, Gi, Gii). The addition of Orco antagonist

OLC15 deleted any significant differences of spontaneous activity found in LD and DD (Figure 4Aiii-

Giii). In conclusion, the mean frequency of spontaneous spiking and the frequency of bursting

expressed daily modulation, and are both most likely controlled via a circadian clock that targets

the leak channel Orco.

5.2 Orco imposes circadian modulation on the ultradian rhythms in spontaneous

ORN activity

After establishing that Orco influences the circadian changes in ORN spiking activity between

the animal's resting and activity phase we examined if Orco also influences ORN spiking rhythms

on the largely different ultradian timescales in the frequency range between 0.01-1000 Hz. In all

conditions, two broad bands of ultradian instantaneous frequencies were evident: one upper band

of >10 to ~100 Hz and a lower band between 0.1 to ≤10 Hz (Figure 5). The upper band primarily

represents the frequencies of spikes within a burst, while the lower band represents single spikes

and frequencies between bursts.

In LD, two types of daily/circadian modulation of the ultradian frequencies occurred (Figure

5A). The upper higher frequency band was mostly modulated in prevalence (its frequency of

occurrence) but not in its frequency range. The lower band was modulated both in prevalence and

frequency range, which resulted in a sinusoidal change in the mean of the frequency band over 24

h. The low and high frequency bands appeared to merge during the activity phase around ZT 0 in the

animals that showed clear circadian rhythms (N = 5 of 11 in LD), when both bands had maximum

prevalence. During the hawkmoth's activity phase, higher frequencies of up to ~10 Hz occurred,

while frequencies dropped below 1 Hz during the resting phase.

The circadian modulation of both frequency bands was maintained in constant darkness

(Figure 5B) but phase-shifted with respect to the projected ZT due to the endogenous circadian

period of each individual. Again, this indicates that a clock is modulating the spontaneous ORN

spiking activity over the course of a day.

The addition of the Orco antagonist OLC15 deleted any circadian modulation of either

frequency band. Interestingly, the instantaneous frequencies in both bands still showed correlated

changes in prevalence and merging, but now at ultradian periods only.

Additional Fourier analysis (see Methods) of spontaneous spiking activity over the course of

several days revealed faster ultradian and slower infradian rhythms, ranging from 0.6 h to 38 h, in

addition to the circadian (24 ± 4 h) rhythm (Figure 6Ai, Bi, Ci). Some rhythms (factorials of 24 h, like

12 h, 6 h, etc.) appeared to be harmonics of the circadian rhythm, which were not excluded in our

analysis. Most animals expressed ultradian rhythms of 2-4 h in LD as well as DD conditions. With

OLC15 the occurrence of circadian rhythmicity decreased from 70% to 33.3 %, while the

prevalence of infradian periods increased from 0% to 33.3 % and ultradian periods remained the

same. These results were corroborated by continuous wavelet analysis, which provided time-

resolved spectral power across different rhythmic periods (Figure 6Aii, Bii, Cii).In control LD and DD

conditions, wavelet power for ultradian periods increased during times of elevated firing rate and

decreased during periods of low activity (Figure 6Aii and Bii, middle and bottom panels). This

dynamic pattern indicates that ultradian rhythms were not constant over time but were instead

modulated by the circadian rhythm, becoming more prominent during circadian peaks in neuronal

activity. Such circadian gating of ultradian power was clearly visible in the wavelet spectrograms as

alternating bands of higher and lower ultradian power aligned with daily activity cycles. Following

Orco blockade with OLC15, this modulation was disrupted (Figure 6Cii middle panel): although

ultradian components persisted (Figure 6Cii bottom panel), they no longer showed systematic

variation with the circadian cycle, suggesting a decoupling of ultradian rhythms from circadian

control.

5.3   Orco expression is not under the control of the molecular TTFL clockwork

So far, our results indicated a prominent role of Orco in the circadian modulation of ORN

spiking activity. Since it is the general opinion that the master circadian clock that dominates all

physiological and behavioral circadian rhythms functions via transcriptional control we examined

whether Orco transcription shows a circadian rhythm, as previously found for clock genes in

hawkmoth antennae (Schneider et al., 2025; Schuckel et al., 2007). However, qPCR of whole male

hawkmoth antennae revealed no significant circadian rhythm of Orco expression in DD (Figure 7;

ANOVA on ranks, $H(2) = 5.60$, $p = 0.061$). Thus, it is possible that Orco is controlled via a

posttranslational circadian clockwork mechanism that is associated directly with the plasma

membrane (Stengl and Schneider, 2024; Stengl and Schröder, 2021).

5.4   Circadian changes in Orco conductance are sufficient to modulate spiking activity in a computational model of a hawkmoth ORN

Although Orco is a leaky ion channel, its maximum conductivity is modulated by cAMP

(Getahun et al., 2013; Martín et al., 2001; Sargsyan et al., 2011; Wicher et al., 2008). In hawkmoth

antennae, cAMP levels oscillate throughout the day (Dolzer et al., 2021; Flecke et al., 2010; Flecke

and Stengl, 2009; Schendzielorz et al., 2014, 2015), which makes cAMP modulation of Orco a

putative mechanism for the Orco-dependent circadian modulation of ORN spiking activity,

integrating Orco in a circadian PTFL based membrane clockwork (Stengl and Schneider, 2024).

To test the hypothesis that Orco is a cAMP-gated, leaky ion channel, we constructed an ORN

model (Methods and Supplementary Material) and tuned the parameters to provide a quantitative

match with the biological data. The developed model provides a unified description of the

experimental observations. For instance, the ORN model yields irregular spiking activity with a

bimodal distribution, consisting of both bursts and isolated spikes, successfully reproducing the

experimental recordings (Figure 8A). This suggests a unified mechanism underlying the initiation of

both phenomena: If the pacemaker-driven depolarization reaches the Na^+ threshold a firing

process begins, and at this point randomness determines if other slower ion channels responsible

for bursting are sufficiently activated to sustain a burst of several spikes.

Additionally, the ORN model replicates the distribution of the number of spikes per burst,

which follows a decreasing exponential trend in both experimental and simulated results (Figure

8B).The model also successfully reproduces the circadian modulation of spiking activity: the

maximum firing rate occurs when the Orco conductance is maximal, whereas the maximum

percentage of spikes that are part of bursts correlates with the lowest Orco conductance, resulting

in antiphase oscillations, as were observed in the experimental recordings (Figure 9).

The frequency prevalences in the model over several days clustered in two distinct frequency

bands: a top band that corresponds to the intra-burst frequencies and a bottom band,

corresponding to inter-burst frequencies (Figure 10), in agreement with the biological experiments

(Figure 5A, B). When these circadian oscillations in Orco conductance were included, inter-burst

frequencies showed significant circadian variations, whereas intra-burst frequencies remained

relatively constant, reminiscent of the biological data. Notably, the simulation captured the

experimentally observed "merging" effect, where the low frequency band approaches the top band

during high-activity periods (Figure 10).

This merging of spiking and bursting frequencies suggests that the system approached a

bifurcation, transitioning from a bursting regime to tonic spiking activity. Further evidence for the

near bifurcation condition of the neurons is the distribution of number of spikes per burst that

followed a decreasing exponential trend in contrast to a constant value, or a bounded range of

spikes per burst, which would be expected from a standard bursting neuron model (Figure 8B). The

observation that the circadian modulation of spiking activity requires only one parameter, namely

changes in the Orco conductivity, to describe the circadian effects on the ORN spiking pattern

indicates the crucial role of Orco channel in circadian control.

6 Discussion

We searched for a membrane-bound clockwork component in insect ORNs that allows for

sensing of regular pheromone patterns to guide mating flights. For this, we recorded interlinked

circadian and ultradian frequencies in the spontaneous spiking activity of the pheromone-sensitive

long trichoid sensilla of male *Manduca sexta* hawkmoths with *in vivo* long-term tip recordings. We

found that Orco, which serves several important functions in insect olfaction (Stengl and Funk,

2013; Wicher and Miazzi, 2021), acts as a pacemaker channel that controls the circadian

modulation of the spiking activity without affecting the frequency ranges that are relevant for the

primary fast, phasic pheromone response (Dolzer et al., 2003; Nolte et al., 2016; Schneider et al.,

2025). While circadian rhythmicity disappeared in the presence of the Orco blocker OLC15,

ultradian activity rhythms remained. We confirmed with computational modeling that the circadian

modulation of open-time probability of Orco as a leaky pacemaker channel was sufficient to

account for our experimental results. Since we observed no evidence for circadian transcriptional

control of Orco, our findings are consistent with the hypothesis that the plasma membrane

constitutes an endogenous post-translational feedback loop (PTFL) clockwork. In this clockwork,

circadian changes in cyclic nucleotide levels (Schendzielorz et al., 2015) modulate Orco

conductance rather than transcription and insertion of additional Orco channels in the membrane.

The expression of a pacemaker channel that does not require daily degradation and

transcription would constitute an economic and rapidly adjustable mechanism to generate

endogenous membrane potential oscillations that are tuned to detect behaviorally relevant

zeitgeber signals in the insect's environment (Stengl and Schneider, 2024; Stengl and Schröder,

2021). The present study provides the first evidence for a systems view of timing that involves a

PTFL membrane clock in hawkmoth ORNs, instead of the prevailing hierarchical view that assumes

all endogenous circadian oscillations are outputs of the genetic TTFL clock (Stengl and Schneider,

2024; Stengl and Schröder, 2021).

6.1 Orco acts as a pacemaker channel to drive the circadian rhythm in spontaneous ORN spiking activity

A common motif of spontaneously active neurons is the expression of pacemaker channels

that open at hyperpolarized potentials and thus depolarize the neuron (Bose et al., 2014; Cochet-

Bissuel et al., 2014; Das et al., 2016; Golowasch et al., 2017; Lüthi and McCormick, 1998; Ratliff et

al., 2021; Robinson and Siegelbaum, 2003; Sharma et al., 2023). The resting membrane potential of

most neurons lies close to the negative equilibrium potential for potassium ions due to potassium

leak channels (Goldstein et al., 2001; Patel and Honoré, 2001; Talley et al., 2001). When the

pacemaker-driven depolarization reaches spike thresholds, voltage-gated Na^+, K^+ and Ca^{2+}

channels promote spiking. Voltage- and Ca^{2+}-gated K^+ channels then repolarize the neuron to

negative membrane potentials, re-starting the cycle

The core element of the different frequencies of endogenous oscillations that needs to be

identified is the respective pacemaker channel (Stengl and Schneider, 2024). The finding of distinct

rhythms in spontaneous spiking activity under constant conditions are indicative of underlying

endogenous membrane potential oscillations. Therefore, different sets of voltage- and Ca^{2+}-

dependent ion channels with antagonistic effects on the membrane potential appear to underlie

different ultradian oscillation frequencies of spontaneous activity that still need to be further

elucidated (Dolzer et al., 2021; Stengl, 1994, 1993; Zufall et al., 1991). While this study did not aim

to unravel the overall generation of spontaneous activity in the ORNs, we could demonstrate that

Orco acts as a circadian pacemaker channel that drives the circadian modulation of the spiking

activity.

6.2 The constitutively expressed Orco is a core element of a circadian PTFL clockwork in hawkmoth ORNs

In hawkmoth ORNs, at least two types of pacemaker channels exist: the highly conserved,

slow, leaky Orco, which we focus on in this study, and an HCN type channel (Butterwick et al.,

2018; Dolzer et al., 2021; Nolte et al., 2016, 2013; Sato et al., 2008; Stengl and Funk, 2013; Stengl

and Schneider, 2024; Stengl and Schröder, 2021; Wicher et al., 2008; Wicher and Miazzi, 2021).

When expressed *in vitro* in heterologous expression systems, *Drosophila* Orco homomers

assemble into leaky, non-specific cation channels with a reversal potential around 0 mV,

constitutively leaking Na^+, K^+, and Ca^{2+} (Sato et al., 2008; Wicher et al., 2008; Wicher and Miazzi,

2021). In *Drosophila*, Orco is regulated by Ca^{2+}/calmodulin and phosphorylation/de-

phosphorylation at multiple sites (Cao et al., 2016; Getahun et al., 2016; Guo and Smith, 2017;

Mukunda et al., 2016, 2014; Sargsyan et al., 2011). In *Drosophila*, the Ca^{2+}/calmodulin binding site

controls the localization of the OR-Orco complex in the ciliary membrane of ORNs and depends on

previous odor stimulation (Cao et al., 2016; Guo and Smith, 2022; Jain et al., 2021). After

phosphorylation of Orco by protein kinase C (PKC), a cyclic nucleotide binding site becomes

available, and subsequent binding of cAMP or cGMP increases the open-time probability (Getahun

et al., 2013; Wicher et al., 2008). This indicates two complementary mechanisms by which Orco

conductance can be regulated: by controlling the number of Orco channels in the membrane or by

modulating the open-time probability of existing channels through posttranslational modifications.

Since blocking Orco in *M. sexta* decreased the spontaneous activity and deleted its circadian

modulation, Orco is apparently the main target for modulation of circadian activity in hawkmoth

ORNs. Since Orco transcription levels were constant, posttranslational modifications of Orco that

modulate the open-time probability provide a mechanistic explanation. Increases in cAMP and

cGMP levels sensitize or adapt the ORNs to pheromone, respectively (Dolzer et al., 2021, 2008;

Flecke et al., 2010, 2006; Redkozubov, 2000; Schendzielorz et al., 2015, 2012; Ziegelberger et al.,

1990). While cAMP levels are under circadian control and peak during the activity phase, cGMP

levels oscillate in a daily rhythm and peak during the resting phase in insects (Schendzielorz et al.,

2014, 2015). We propose that cAMP and cGMP binding modulate Orco with opposing effects,

consistent with the physiological needs of the hawkmoth during its sleep-wake cycle: cAMP

binding increases Orco conductance while cGMP binding decreases it.

Our hypothesis that a circadian PTFL membrane clock in hawkmoth ORNs controls circadian

activity rhythms predicts that the pacemaker channel Orco is part of a membrane signalosome.

The protein complexes of this signalosome would comprise the pacemaker channel that generates

the specific circadian frequency of endogenous potential oscillations, linked to oscillations of

second messenger cascades that determine physiological setpoints of the circadian sleep-wake

cycles. Thus, the circadian cAMP oscillations could result from a PTFL membrane clock comprising

of a signalosome that minimally consists of voltage-gated Ca^{2+}-channels, Orco, and membrane-

anchored adenylyl cyclases (ACs). Since the membrane potential directly influences intracellular

Ca^{2+} levels, the AC activity could be periodically upregulated by the circadian oscillations of the

endogenous membrane potential oscillations. Alternatively, AC expression could be under the

control of the molecular TTFL clock, which would put Orco under indirect TTFL control. This

possibility should be further explored in hawkmoth antennae. Instead of, or in addition to, the

voltage-gated Ca^{2+} channels the signalosome could also work if it contained octopamine receptors

that cause circadian rises in cAMP levels through activation of $G_{\alpha s}$, as found in hawkmoths (Dacks

et al., 2006; Flecke et al., 2010). Since octopamine levels in the hemolymph exhibit circadian

oscillations and octopamine sets a physiological state of heightened alertness in insects, the

circadian PTFL clock in ORNs would synchronize with endogenous physiological rhythms via

octopamine receptor activation (Flecke and Stengl, 2009; Schendzielorz et al., 2015).

6.3   Circadian and ultradian rhythms in spontaneous ORN activity resemble the

physiologically relevant timescales for active sensing

Moths are more sensitive to pheromone stimulation during their activity phase than at rest,

apparently due to antennal circadian clocks (Flecke et al., 2010; Flecke and Stengl, 2009; Merlin et

al., 2007, 2006; Schendzielorz et al., 2015; Schuckel et al., 2007; Silvegren et al., 2005). The greater

the distance from the female, the lower the pheromone concentration. The increased pheromone

sensitivity during the activity phase enhances the male's chance to detect a calling female.

Pheromone concentration is encoded in the frequency of the first, phasic burst of spikes of

the characteristic triphasic spiking sequence of ORNs in response to brief pheromone pulses,

which is independent of Orco (Dolzer et al., 2003; Nolte et al., 2013; Schneider et al., 2025).

However, through the cAMP-dependent increase in Orco conductance during the activity phase,

the ORN membrane potential is closer to spike threshold and fewer pheromone molecules are

necessary to produce a receptor current that brings the membrane voltage over that threshold.

Thus, larger Orco conductance increases the likelihood of an ORN response to pheromone

stimulation without changing the typical triphasic spiking pattern of the response.

In the odor transduction of *Drosophila*, OR-Orco activation drives the initial receptor current

response. In contrast, Orco is not part of an OR-Orco odor-gated receptor ion channel complex in

hawkmoths but controls the spontaneous ORN activity and the late, long-lasting pheromone

response (LLPR). The LLPR begins several hundred milliseconds after the initiation of the

pheromone-dependent transduction current and correlates with the time course of pheromone-
induced rises in cGMP levels (Boekhoff et al., 1993; Nolte et al., 2016, 2013; Stengl, 2010; Stengl et
al., 2001; Wicher and Miazzi, 2021; Ziegelberger et al., 1990). Thus, in hawkmoths, Orco is not part
of the primary transduction channel but rather controls membrane potential oscillations and
thereby the temporal resolution and general pheromone sensitivity of ORNs. The higher
spontaneous burst frequency during the hawkmoth's activity phase improves the resolution of the
anticipated pheromone pulses during its mating flight in search for females.

In the mammalian suprachiasmatic nucleus (SCN), the clock that controls the sleep-wake
cycle, cAMP levels oscillate at an ultradian frequency, and cAMP and cGMP induce opposing phase
shifts in SCN spiking activity (Prosser et al., 1989; Prosser and Gillette, 1991). In *M. sexta* antennae,
cAMP levels express daily oscillations and peak during the activity phase. In contrast, cGMP levels
do not oscillate but are elevated by pheromone exposure, which adapts the spiking activity
(Boekhoff et al., 1993; Dolzer et al., 2021; Flecke et al., 2006; Schendzielorz et al., 2015;
Ziegelberger et al., 1990). This implies that cAMP and cGMP may have opposing effects on Orco
conductivity, with results similar to what is seen in the SCN.

The upregulation of neural activity during wakefulness/alertness seems to be ubiquitous
among animals. For example, the many synchronously recorded neurons in human EEG show high
frequency oscillations in the beta and gamma ranges during alertness and low frequency
oscillations in the delta and theta ranges during rest and sleep. Similarly, the spontaneous spiking
activity of hawkmoth ORNs is upregulated during their activity phase, albeit only from below delta
to the delta/theta range. The increased spontaneous spiking activity in the theta range during the
activity phase of the hawkmoth matches the maximal temporal resolution of pheromone pulses of
up to 10 Hz (Marion-Poll and Tobin, 1992), which supports a role for Orco as part of a membrane-
bound PTFL clockwork that tunes the ORNs for active sensing. Even through oscillations in the field
potential of electroantennogram responses to odor stimulation can reach 100 Hz in different
insects (Szyszka et al., 2014), these frequencies most likely do not encode physiological odor
responses. Apparently, they reflect the astounding adaptive ability of antennal ORNs to
synchronize their endogenous membrane potential oscillations across the antenna to a broad
range of stimulation frequencies. Pheromone pulse resolution of about 100 Hz was never reported
in single olfactory sensilla recordings, and in behavioral experiments male moths stopped their
upwind mate search if pheromone pulses exceeded 30 Hz (Baker et al., 1985; Bau et al., 2005,
2002; Tripathy et al., 2010).

Circadian oscillations in the ORN membrane potential would be ideally suited to tune

sensitivity and temporal resolution of the pheromone-sensitive ORNs to the hawkmoth's rest-

activity cycle, synchronizing male and female mating behavior. Further experiments are needed to

test our hypothesis that ORN signalosomes are highly plastic endogenous PTFL clocks comprising

receptors for circadian and ultradian Zeitgebers that allow to tune into internal physiological and

external environmental rhythms as basis for active sensing.

6.4   Stochastic ion channel dynamics and circadian modulation of Orco conductance explain modeled ORN firing patterns

Our modeling results provide mechanistic insight into the irregular spontaneous activity of

pheromone-sensitive ORNs in *M. sexta*, highlighting how stochastic ion channel dynamics and

circadian modulation of ion conductance together shape spiking patterns. By employing a Langevin

approximation of the stochastic HH formalism, we demonstrate that the intrinsic noise arising from

the random gating of ion channels is sufficient to generate both isolated spikes and bursts of

spikes, as observed experimentally. Our model effectively recapitulates the spike distributions

found *in vivo* and accounts for the role of channel noise as a critical determinant of ORN firing

patterns.

A key contribution of our model is the incorporation of the Orco ion channel as a pacemaker

component, with a conductance that varies linearly with cAMP concentration. We show that

circadian modulation of Orco conductance alone is sufficient to produce the observed time-of-day

dependent changes in spike distribution. This finding suggests that the rhythmic expression of

cAMP levels across the circadian cycle may act as a primary driver of diurnal changes in ORN

excitability, without requiring additional external inputs or modulation of other ion channels.

Importantly, our results reveal that the same set of ion channels can support the emergence

of both single action potentials and complex bursting behaviors purely through the stochastic

variability of their gating kinetics. This unifying explanation reduces the need for invoking separate

biophysical mechanisms or circuit-level inputs to explain these distinct firing modes.

For simplicity, the circadian variation in cAMP concentration was modeled using a sinusoidal

function. While this approximation is sufficient for capturing the qualitative features of the

rhythmic modulation, future work could enhance the biological realism of the model by coupling

Orco conductance dynamics to a molecular circadian oscillator, such as a Goodwin-type

transcriptional-translational feedback loop. Incorporating this layer would allow for a more

detailed exploration of how molecular clock components regulate cAMP synthesis and

degradation. Additionally, further extensions could include downstream regulatory mechanisms

such as phosphorylation and dephosphorylation events mediated by kinases and phosphatases

under circadian control, potentially contributing to fine-tuned modulation of ion channel function

over the daily cycle.

Together, these findings demonstrate the value of a biophysically grounded, noise-inclusive

modeling framework for understanding complex temporal patterns in sensory neuron activity and

open the door for future integrative models linking molecular circadian clocks to cellular

excitability.

7   References

Baker TC, Vogt RG. 1988. Measured Behavioural Latency in Response to Sex-Pheromone Loss in
the Large Silk Moth *Antheraea Polyphemus*. *J Exp Biol* **137**:29–38. doi:10.1242/jeb.137.1.29
Baker TC, Willis MA, Haynes KF, Phelan PL. 1985. A pulsed cloud of sex pheromone elicits upwind
flight in male moths. *Physiol Entomol* **10**:257–265. doi:10.1111/j.1365-3032.1985.tb00045.x
Bau J, Justus KA, Cardé RT. 2002. Antennal resolution of pulsed pheromone plumes in three moth
species. *J Insect Physiol* **48**:433–442. doi:10.1016/S0022-1910(02)00062-8
Bau J, Justus KA, Loudon C, Cardé RT. 2005. Electroantennographic Resolution of Pulsed
Pheromone Plumes in Two Species of Moths with Bipectinate Antennae. *Chem Senses*
30:771–780. doi:10.1093/chemse/bji069
Bell RA, Joachim FG. 1976. Techniques for Rearing Laboratory Colonies of Tobacco Hornworms
and Pink Bollworms1. *Ann Entomol Soc Am* **69**:365–373. doi:10.1093/aesa/69.2.365
Boekhoff I, Seifert E, Göggerle S, Lindemann M, Krüger B-W, Breer H. 1993. Pheromone-induced
second-messenger signaling in insect antennae. *Insect Biochem Mol Biol* **23**:757–762.
doi:10.1016/0965-1748(93)90063-X
Bose A, Golowasch J, Guan Y, Nadim F. 2014. The role of linear and voltage-dependent ionic
currents in the generation of slow wave oscillations. *J Comput Neurosci* **37**:229–242.
doi:10.1007/s10827-014-0498-4
Butterwick JA, del Mármol J, Kim KH, Kahlson MA, Rogow JA, Walz T, Ruta V. 2018. Cryo-EM
structure of the insect olfactory receptor Orco. *Nature* **560**:447–452. doi:10.1038/s41586-
018-0420-8
Cao L-H, Jing B-Y, Yang D, Zeng X, Shen Y, Tu Y, Luo D-G. 2016. Distinct signaling of *Drosophila*
chemoreceptors in olfactory sensory neurons. *Proc Natl Acad Sci* **113**:E902–E911.
doi:10.1073/pnas.1518329113
Chen S, Luetje CW. 2012. Identification of New Agonists and Antagonists of the Insect Odorant
Receptor Co-Receptor Subunit. *PLOS ONE* **7**:e36784. doi:10.1371/journal.pone.0036784
Cochet-Bissuel M, Lory P, Monteil A. 2014. The sodium leak channel, NALCN, in health and
disease. *Front Cell Neurosci* **8**:132. doi:10.3389/fncel.2014.00132
Dacks AM, Dacks JB, Christensen TA, Nighorn AJ. 2006. The cloning of one putative octopamine
receptor and two putative serotonin receptors from the tobacco hawkmoth, *Manduca*
sexta. *Insect Biochem Mol Biol* **36**:741–747. doi:10.1016/j.ibmb.2006.07.002
Das A, Holmes TC, Sheeba V. 2016. dTRPA1 in Non-circadian Neurons Modulates Temperature-
dependent Rhythmic Activity in *Drosophila melanogaster*. *J Biol Rhythms* **31**:272–288.
doi:10.1177/0748730415627037
Dolzer J, Fischer K, Stengl M. 2003. Adaptation in pheromone-sensitive trichoid sensilla of the
hawkmoth *Manduca sexta*. *J Exp Biol* **206**:1575–1588. doi:10.1242/jeb.00302
Dolzer J, Krannich S, Fischer K, Stengl M. 2001. Oscillations of the Transepithelial Potential of Moth
Olfactory Sensilla Are Influenced by Octopamine and Serotonin. *J Exp Biol* **204**:2781–2794.
doi:10.1242/jeb.204.16.2781
Dolzer J, Krannich S, Stengl M. 2008. Pharmacological Investigation of Protein Kinase C- and cGMP-
Dependent Ion Channels in Cultured Olfactory Receptor Neurons of the Hawkmoth
Manduca sexta. *Chem Senses* **33**:803–813. doi:10.1093/chemse/bjn043
Dolzer J, Schröder K, Stengl M. 2021. Cyclic nucleotide-dependent ionic currents in olfactory
receptor neurons of the hawkmoth *Manduca sexta* suggest pull–push sensitivity
modulation. *Eur J Neurosci* **54**:4804–4826. doi:10.1111/ejn.15346

Dubowy C, Sehgal A. 2017. Circadian Rhythms and Sleep in *Drosophila melanogaster*. *Genetics*
205:1373–1397. doi:10.1534/genetics.115.185157
Flecke C, Dolzer J, Krannich S, Stengl M. 2006. Perfusion with cGMP analogue adapts the action
potential response of pheromone-sensitive sensilla trichoidea of the hawkmoth *Manduca*
sexta in a daytime-dependent manner. *J Exp Biol* **209**:3898–3912. doi:10.1242/jeb.02432
Flecke C, Nolte A, Stengl M. 2010. Perfusion with cAMP analogue affects pheromone-sensitive
trichoid sensilla of the hawkmoth *Manduca sexta* in a time-dependent manner. *J Exp Biol*
213:842–852. doi:10.1242/jeb.032839
Flecke C, Stengl M. 2009. Octopamine and tyramine modulate pheromone-sensitive olfactory
sensilla of the hawkmoth *Manduca sexta* in a time-dependent manner. *J Comp Physiol A*
195:529–545. doi:10.1007/s00359-009-0429-4
Fox RF, Lu Y. 1994. Emergent collective behavior in large numbers of globally coupled
independently stochastic ion channels. *Phys Rev E* **49**:3421–3431.
doi:10.1103/PhysRevE.49.3421
Géron A, Werner ,Johannes, Wattiez ,Ruddy, and Matallana-Surget S. 2024. Towards the discovery
of novel molecular clocks in Prokaryotes. *Crit Rev Microbiol* **50**:491–503.
doi:10.1080/1040841X.2023.2220789
Getahun MN, Olsson SB, Lavista-Llanos S, Hansson BS, Wicher D. 2013. Insect Odorant Response
Sensitivity Is Tuned by Metabotropically Autoregulated Olfactory Receptors. *PLOS ONE*
8:e58889. doi:10.1371/journal.pone.0058889
Getahun MN, Thoma M, Lavista-Llanos S, Keesey I, Fandino RA, Knaden M, Wicher D, Olsson SB,
Hansson BS. 2016. Intracellular regulation of the insect chemoreceptor complex impacts
odour localization in flying insects. *J Exp Biol* **219**:3428–3438. doi:10.1242/jeb.143396
Ghosh S, Suray C, Bozzolan F, Palazzo A, Monsempès C, Lecouvreur F, Chatterjee A. 2024.
Pheromone-mediated command from the female to male clock induces and synchronizes
circadian rhythms of the moth *Spodoptera littoralis*. *Curr Biol* **34**:1414-1425.e5.
doi:10.1016/j.cub.2024.02.042
Goldstein SAN, Bockenhauer D, O'Kelly I, Zilberberg N. 2001. Potassium leak channels and the
KCNK family of two-p-domain subunits. *Nat Rev Neurosci* **2**:175–184.
doi:10.1038/35058574
Golowasch J, Bose A, Guan Y, Salloum D, Roeser A, Nadim F. 2017. A balance of outward and
linear inward ionic currents is required for generation of slow-wave oscillations. *J*
Neurophysiol **118**:1092–1104. doi:10.1152/jn.00240.2017
Guo H, Smith DP. 2022. Time-Dependent Odorant Sensitivity Modulation in Insects. *Insects* **13**:354.
doi:10.3390/insects13040354
Guo H, Smith DP. 2017. Odorant Receptor Desensitization in Insects. *J Exp Neurosci*
11:1179069517748600. doi:10.1177/1179069517748600
Hall JC, Rosbash M. 1993. Oscillating molecules and how they move circadian clocks across
evolutionary boundaries. *Proc Natl Acad Sci U S A* **90**:5382–5383.
Hardin PE. 2011. Molecular Genetic Analysis of Circadian Timekeeping in *Drosophila* In: Brody S,
editor. Advances in Genetics, The Genetics of Circadian Rhythms. Academic Press. pp.
141–173. doi:10.1016/B978-0-12-387690-4.00005-2
Hastings MH, Smyllie NJ, Patton AP. 2020. Molecular-genetic Manipulation of the Suprachiasmatic
Nucleus Circadian Clock. *J Mol Biol* **432**:3639–3660. doi:10.1016/j.jmb.2020.01.019
Itagaki H, Conner WE. 1988. Calling Behavior of *Manduca sexta* (L.) (Lepidoptera: Sphingidae) with
Notes on the Morphology of the Female Sex Pheromone Gland. *Ann Entomol Soc Am*
81:798–807. doi:10.1093/aesa/81.5.798

Jain K, Lavista-Llanos S, Grabe V, Hansson BS, Wicher D. 2021. Calmodulin regulates the olfactory
performance in Drosophila melanogaster. *Sci Rep* **11**:3747. doi:10.1038/s41598-021-
83296-9
Jones PL, Pask GM, Rinker DC, Zwiebel LJ. 2011. Functional agonism of insect odorant receptor ion
channels. *Proc Natl Acad Sci* **108**:8821–8825. doi:10.1073/pnas.1102425108
Kaissling KE, Zack Strausfeld C, Rumbo ER. 1987. Adaptation processes in insect olfactory
receptors. Mechanisms and behavioral significance. *Ann N Y Acad Sci* **510**:104–112.
doi:10.1111/j.1749-6632.1987.tb43475.x
Krishnan B, Dryer SE, Hardin PE. 1999. Circadian rhythms in olfactory responses of *Drosophila
melanogaster*. *Nature* **400**:375–378. doi:10.1038/22566
Lamont EW, Amir S. 2010. Circadian and Ultradian Clocks/Rhythms In: Koob GF, Moal ML,
Thompson RF, editors. Encyclopedia of Behavioral Neuroscience. Oxford: Academic Press.
pp. 257–261. doi:10.1016/B978-0-08-045396-5.00223-2
Lee JK, Strausfeld NJ. 1990. Structure, distribution and number of surface sensilla and their
receptor cells on the olfactory appendage of the male mothManduca sexta. *J Neurocytol*
19:519–538. doi:10.1007/BF01257241
Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time
quantitative PCR and the $2^{-\Delta\Delta C(T)}$ Method. *Methods San Diego Calif* **25**:402–408.
doi:10.1006/meth.2001.1262
Lüthi A, McCormick DA. 1998. H-Current: Properties of a Neuronal and Network Pacemaker.
Neuron **21**:9–12. doi:10.1016/S0896-6273(00)80509-7
Marion-Poll F, Tobin TR. 1992. Temporal coding of pheromone pulses and trains in *Manduca sexta*.
J Comp Physiol A **171**:505–512. doi:10.1007/BF00194583
Martín F, Charro M, Alcorta E. 2001. Mutations affecting the cAMP transduction pathway modify
olfaction in *Drosophila*. *J Comp Physiol A* **187**:359–370. doi:10.1007/s003590100208
Merlin C, François M-C, Queguiner I, Maïbèche-Coisné M, Jacquin-Joly E. 2006. Evidence for a
putative antennal clock in *Mamestra brassicae*: Molecular cloning and characterization of
two clock genes –*period* and *cryptochrome*– in antennae. *Insect Mol Biol* **15**:137–145.
doi:10.1111/j.1365-2583.2006.00617.x
Merlin C, Gegear RJ, Reppert SM. 2009. Antennal Circadian Clocks Coordinate Sun Compass
Orientation in Migratory Monarch Butterflies. *Science* **325**:1700–1704.
doi:10.1126/science.1176221
Merlin C, Lucas P, Rochat D, François M-C, Maïbèche-Coisne M, Jacquin-Joly E. 2007. An Antennal
Circadian Clock and Circadian Rhythms in Peripheral Pheromone Reception in the Moth
Spodoptera littoralis. *J Biol Rhythms* **22**:502–514. doi:10.1177/0748730407307737
Millar AJ. 2016. The Intracellular Dynamics of Circadian Clocks Reach for the Light of Ecology and
Evolution. *Annu Rev Plant Biol* **67**:595–618. doi:10.1146/annurev-arplant-043014-115619
Mönke G, Sorgenfrei FA, Schmal C, Granada AE. 2020. Optimal time frequency analysis for
biological data - pyBOAT. doi:10.1101/2020.04.29.067744
Mukunda L, Miazzi F, Kaltofen S, Hansson BS, Wicher D. 2014. Calmodulin modulates insect
odorant receptor function. *Cell Calcium* **55**:191–199. doi:10.1016/j.ceca.2014.02.013
Mukunda L, Miazzi F, Sargsyan V, Hansson BS, Wicher D. 2016. Calmodulin Affects Sensitization of
Drosophila melanogaster Odorant Receptors. *Front Cell Neurosci* **10**.
doi:10.3389/fncel.2016.00028
Nakagawa T, Vosshall LB. 2009. Controversy and consensus: noncanonical signaling mechanisms
in the insect olfactory system. *Curr Opin Neurobiol*, Signalling mechanisms **19**:284–292.
doi:10.1016/j.conb.2009.07.015

Nishiitsutsuji-Uwo J, Pittendrigh CS. 1968. Central nervous system control of circadian rhythmicity
in the cockroach II. The pathway of light signals that entrain the rhythm. *Z Für Vgl Physiol*
58:1–13. doi:10.1007/BF00302433

Nolte A, Funk NW, Mukunda L, Gawalek P, Werckenthin A, Hansson BS, Wicher D, Stengl M. 2013.
In situ Tip-Recordings Found No Evidence for an Orco-Based Ionotropic Mechanism of
Pheromone-Transduction in *Manduca sexta*. *PLOS ONE* **8**:e62648.
doi:10.1371/journal.pone.0062648

Nolte A, Gawalek P, Koerte S, Wei H, Schumann R, Werckenthin A, Krieger J, Stengl M. 2016. No
Evidence for Ionotropic Pheromone Transduction in the Hawkmoth *Manduca sexta*. *PLOS*
ONE **11**:e0166060. doi:10.1371/journal.pone.0166060

Page TL, Koelling E. 2003. Circadian rhythm in olfactory response in the antennae controlled by the
optic lobe in the cockroach. *J Insect Physiol* **49**:697–707. doi:10.1016/s0022-
1910(03)00071-4

Patel AJ, Honoré E. 2001. Properties and modulation of mammalian 2P domain K+ channels.
Trends Neurosci **24**:339–346. doi:10.1016/S0166-2236(00)01810-5

Pézier A, Acquistapace A, Renou M, Rospars J-P, Lucas P. 2007. Ca^{2+} Stabilizes the Membrane
Potential of Moth Olfactory Receptor Neurons at Rest and Is Essential for Their Fast
Repolarization. *Chem Senses* **32**:305–317. doi:10.1093/chemse/bjl059

Plautz JD, Kaneko M, Hall JC, Kay SA. 1997. Independent Photoreceptive Circadian Clocks
Throughout *Drosophila*. *Science* **278**:1632–1635. doi:10.1126/science.278.5343.1632

Prosser RA, Gillette MU. 1991. Cyclic changes in cAMP concentration and phosphodiesterase
activity in a mammalian circadian clock studied *in vitro*. *Brain Res* **568**:185–192.
doi:10.1016/0006-8993(91)91396-I

Prosser RA, McArthur AJ, Gillette MU. 1989. cGMP induces phase shifts of a mammalian circadian
pacemaker at night, in antiphase to cAMP effects. *Proc Natl Acad Sci U S A* **86**:6812–6815.

Pu S, Thomas PJ. 2020. Fast and Accurate Langevin Simulations of Stochastic Hodgkin-Huxley
Dynamics. *Neural Comput* **32**:1775–1835. doi:10.1162/neco_a_01312

Ratliff J, Franci A, Marder E, O'Leary T. 2021. Neuronal oscillator robustness to multiple global
perturbations. *Biophys J* **120**:1454–1468. doi:10.1016/j.bpj.2021.01.038

Redkozubov A. 2000. Guanosine 3',5'-Cyclic Monophosphate Reduces the Response of the Moth's
Olfactory Receptor Neuron to Pheromone. *Chem Senses* **25**:381–385.
doi:10.1093/chemse/25.4.381

Reinhard N, Fukuda A, Manoli G, Derksen E, Saito A, Möller G, Sekiguchi M, Rieger D, Helfrich-
Förster C, Yoshii T, Zandawala M. 2024. Synaptic connectome of the *Drosophila* circadian
clock. *Nat Commun* **15**:10392. doi:10.1038/s41467-024-54694-0

Riffell JA, Abrell L, Hildebrand JG. 2008. Physical processes and real-time chemical measurement
of the insect olfactory environment. *J Chem Ecol* **34**:837–853. doi:10.1007/s10886-008-
9490-7

Robinson RB, Siegelbaum SA. 2003. Hyperpolarization-activated cation currents: from molecules
to physiological function. *Annu Rev Physiol* **65**:453–480.
doi:10.1146/annurev.physiol.65.092101.142734

Sargsyan V, Getahun MN, Lavista Llanos S, Olsson SB, Hansson BS, Wicher D. 2011.
Phosphorylation via PKC Regulates the Function of the *Drosophila* Odorant Co-Receptor.
Front Cell Neurosci **5**. doi:10.3389/fncel.2011.00005

Sato K, Pellegrino M, Nakagawa Takao, Nakagawa Tatsuro, Vosshall LB, Touhara K. 2008. Insect
olfactory receptors are heteromeric ligand-gated ion channels. *Nature* **452**:1002–1006.
doi:10.1038/nature06850

Sauman I, Reppert SM. 1998. Brain Control of Embryonic Circadian Rhythms in the Silkmoth
Antheraea pernyi. *Neuron* **20**:741–748. doi:10.1016/S0896-6273(00)81012-0
Schendzielorz J, Schendzielorz T, Arendt A, Stengl M. 2014. Bimodal Oscillations of Cyclic
Nucleotide Concentrations in the Circadian System of the Madeira Cockroach *Rhyparobia*
maderae. *J Biol Rhythms* **29**:318–331. doi:10.1177/0748730414546133
Schendzielorz T, Peters W, Boekhoff I, Stengl M. 2012. Time of Day Changes in Cyclic Nucleotides
Are Modified via Octopamine and Pheromone in Antennae of the Madeira Cockroach. *J Biol*
Rhythms **27**:388–397. doi:10.1177/0748730412456265
Schendzielorz T, Schirmer K, Stolte P, Stengl M. 2015. Octopamine Regulates Antennal Sensory
Neurons via Daytime-Dependent Changes in cAMP and IP_3 Levels in the Hawkmoth
Manduca sexta. *PLOS ONE* **10**:e0121230. doi:10.1371/journal.pone.0121230
Schmidt DR, Galán RF, Thomas PJ. 2018. Stochastic shielding and edge importance for Markov
chains with timescale separation. *PLOS Comput Biol* **14**:e1006206.
doi:10.1371/journal.pcbi.1006206
Schmidt DR, Thomas PJ. 2014. Measuring Edge Importance: A Quantitative Analysis of the
Stochastic Shielding Approximation for Random Processes on Graphs. *J Math Neurosci* **4**:6.
doi:10.1186/2190-8567-4-6
Schneider AC, Schröder K, Chang Y, Nolte A, Gawalek P, Stengl M. 2025. Hawkmoth Pheromone
Transduction Involves G-Protein–Dependent Phospholipase Cβ Signaling. *eNeuro*
12:ENEURO.0376-24.2024. doi:10.1523/ENEURO.0376-24.2024
Schuckel J, Siwicki KK, Stengl M. 2007. Putative circadian pacemaker cells in the antenna of the
hawkmoth *Manduca sexta*. *Cell Tissue Res* **330**:271–278. doi:10.1007/s00441-007-0471-x
Sharma A, Rahman G, Gorelik J, Bhargava A. 2023. Voltage-Gated T-Type Calcium Channel
Modulation by Kinases and Phosphatases: The Old Ones, the New Ones, and the Missing
Ones. *Cells* **12**:461. doi:10.3390/cells12030461
Silvegren G, Löfstedt C, Qi Rosén W. 2005. Circadian mating activity and effect of pheromone pre-
exposure on pheromone response rhythms in the moth *Spodoptera littoralis*. *J Insect*
Physiol **51**:277–286. doi:10.1016/j.jinsphys.2004.11.013
Stengl M. 2010. Pheromone Transduction in Moths. *Front Cell Neurosci* **4**:133.
doi:10.3389/fncel.2010.00133
Stengl M. 1994. Inositol-trisphosphate-dependent calcium currents precede cation currents in
insect olfactory receptor neurons *in vitro*. *J Comp Physiol A* **174**:187–194.
doi:10.1007/BF00193785
Stengl M. 1993. Intracellular-Messenger-Mediated Cation Channels in Cultured Olfactory Receptor
Neurons. *J Exp Biol* **178**:125–147. doi:10.1242/jeb.178.1.125
Stengl M, Funk NW. 2013. The role of the coreceptor Orco in insect olfactory transduction. *J Comp*
Physiol A **199**:897–909. doi:10.1007/s00359-013-0837-3
Stengl M, Schneider AC. 2024. Contribution of membrane-associated oscillators to biological
timing at different timescales. *Front Physiol* **14**:1243455. doi:10.3389/fphys.2023.1243455
Stengl M, Schröder K. 2021. 14 - Multiscale timing of pheromone transduction in hawkmoth
olfactory receptor neurons. In: Blomquist GJ, Vogt RG, editors. Insect Pheromone
Biochemistry and Molecular Biology (Second Edition). London: Academic Press. pp. 435–
468. doi:10.1016/B978-0-12-819628-1.00014-6
Stengl M, Zintl R, De Vente J, Nighorn A. 2001. Localization of cGMP immunoreactivity and of
soluble guanylyl cyclase in antennal sensilla of the hawkmoth Manduca sexta. *Cell Tissue*
Res **304**:409–421. doi:10.1007/s004410000336

Szyszka P, Gerkin RC, Galizia CG, Smith BH. 2014. High-speed odor transduction and pulse
tracking by insect olfactory receptor neurons. *Proc Natl Acad Sci* **111**:16925–16930.
doi:10.1073/pnas.1412051111
Talley EM, Solórzano G, Lei Q, Kim D, Bayliss DA. 2001. CNS Distribution of Members of the Two-
Pore-Domain (KCNK) Potassium Channel Family. *J Neurosci* **21**:7491–7505.
doi:10.1523/JNEUROSCI.21-19-07491.2001
Tanoue S, Krishnan P, Krishnan B, Dryer SE, Hardin PE. 2004. Circadian Clocks in Antennal
Neurons Are Necessary and Sufficient for Olfaction Rhythms in *Drosophila*. *Curr Biol*
14:638–649. doi:10.1016/j.cub.2004.04.009
Thaiss CA, Zeevi D, Levy M, Zilberman-Schapira G, Suez J, Tengeler AC, Abramson L, Katz MN,
Korem T, Zmora N, Kuperman Y, Biton I, Gilad S, Harmelin A, Shapiro H, Halpern Z, Segal E,
Elinav E. 2014. Transkingdom Control of Microbiota Diurnal Oscillations Promotes
Metabolic Homeostasis. *Cell* **159**:514–529. doi:10.1016/j.cell.2014.09.048
Tripathy S, Staudacher EM, Peters O, Kalwar F, Hatfield M, Daly K. 2010. Odors pulsed at wing beat
frequencies are tracked by primary olfactory networks and enhance odor detection. *Front*
Cell Neurosci **4**. doi:10.3389/neuro.03.001.2010
Viertel R, Borisyuk A. 2019. A Computational model of the mammalian external tufted cell. *J Theor*
Biol **462**:109–121. doi:10.1016/j.jtbi.2018.10.003
Wicher D, Miazzi F. 2021. Functional properties of insect olfactory receptors: ionotropic receptors
and odorant receptors. *Cell Tissue Res* **383**:7–19. doi:10.1007/s00441-020-03363-x
Wicher D, Schäfer R, Bauernfeind R, Stensmyr MC, Heller R, Heinemann SH, Hansson BS. 2008.
Drosophila odorant receptors are both ligand-gated and cyclic-nucleotide-activated cation
channels. *Nature* **452**:1007–1011. doi:10.1038/nature06861
Zhou X, Yuan C, Guo A. 2005. *Drosophila* Olfactory Response Rhythms Require Clock Genes but
Not Pigment Dispersing Factor or Lateral Neurons. *J Biol Rhythms* **20**:237–244.
doi:10.1177/0748730405274451
Ziegelberger G, Berg M van den, Kaissling KE, Klumpp S, Schultz JE. 1990. Cyclic GMP levels and
guanylate cyclase activity in pheromone-sensitive antennae of the silkmoths *Antheraea*
polyphemus and *Bombyx mori*. *J Neurosci* **10**:1217–1225. doi:10.1523/JNEUROSCI.10-04-
01217.1990
Zufall F, Stengl M, Franke C, Hildebrand JG, Hatt H. 1991. Ionic currents of cultured olfactory
receptor neurons from antennae of male Manduca sexta. *J Neurosci* **11**:956–965.
doi:10.1523/JNEUROSCI.11-04-00956.1991
923

924 # 8 Figures and Figure legends

Figure 1: Attributes of pheromone-sensitive ORN spiking patterns.

Spikes were identified by their prominence in the raw ORN recording. The inter-spike interval (ISI) was defined as the time from one spike to the next. A train with ≥2 spikes with ISI ≤ 50 ms was considered a burst. The intra-burst interval was the ISIs within a burst, the inter-burst interval (IBI) the ISIs between two bursts, omitting any individual spikes in between. The inter-event interval (IEI) included the ISIs between bursts and individual spikes.

Figure 2: The conductance-based model of a simplified hawkmoth ORN.

(A) Schematic of the model containing a voltage gated sodium channel (Na⁺), a voltage gated potassium channel (K⁺), a low voltage activated calcium channel (LVA), a calcium and voltage activated potassium channel (BK), a leak channel, and a cAMP-gated leaky non-specific cation channel (Orco). The Orco channel is represented as a linear conductance whose value oscillates with a sinusoidal shape as a function of the ZT. The circadian rhythm in the Orco conductance is hypothesized to result from a cAMP-dependent increase in channel open-time probability (as found in *Drosophila* Orco: (Getahun et al., 2013)) via a circadian oscillation of antennal cAMP concentrations (Schendzielorz et al., 2015). This oscillation is driven by the circadian clock and serves as an input; the internal mechanism of the clock is not modeled explicitly. Arrows indicate the flow of ions through the channels. **(B)** Parametrized circadian oscillation in Orco conductivity,

introduced as an external input to our model, representing the gating of Orco by cAMP whose

concentration oscillates on a circadian timescale. The times of minimum and maximum cAMP

concentration are indicated by colored arrows. **(C)** Parametrized IV curve of the Orco channel

showing a linear behavior with the slope depending on the cAMP concentration. Line colors

correspond to the arrows in panel B. **(D)** Markov-chain representing all possible states and

transitions for each type of ion channel included in the model. The per capita transition rates (α and

β) depend on membrane potential and in the case of the BK channel also on the intracellular Ca^{2+}-

concentration. The "N" state vectors contain the population of channels in each state "n".

Subscripts of "n" indicate how many activation and inactivation gates are in the open state.

Directed edges are numbered in red. The only conducting state of each ion channel, representing

the condition where all its gates are open, is shown in green. Panel **A** was drawn using pictures

from Servier Medical Art. Servier Medical Art by Servier is licensed under a Creative Commons

Attribution 3.0 Unported License.

Figure 3: Spontaneous activity of pheromone-sensitive olfactory receptor neurons (ORNs) of male *M. sexta* shows Orco-dependent circadian modulation.

(A) Example tip recordings of the spontaneous activity of one long trichoid sensillum during a period of high (top, average frequency ~ 3 Hz) and low (bottom, average frequency ~ 0.2 Hz) activity. **(Bi)** Spontaneous spike frequency of one long-term ORN recording during 17:7 light-dark cycles (same experiment as in **A,** colored arrows indicate respective zeitgeber times (ZT)), indicated by the black-white bar at the top, increased during each activity phase and was low during the resting phases. Spike frequencies were averaged for each 1 h bin. **(Bii)** Mean spontaneous ORN spike frequency across all LD animals in 1 h bins. Data from the individual in Bi is highlighted in red. Recordings had different lengths, therefore the number of recordings used for the element-wise mean for each 1 h bin decreased with the time since the start of the recording, indicated in the top

panel. **(Biii)** Mean spontaneous ORN spike frequency across all LD animals in 1 h bins during the
first 48 hours revealed circadian activity. **(Ci)** The spontaneous spike frequency of one long-term
ORN recording in constant darkness (DD, the expected times of lights on are represented by grey
bars above) exhibited a circadian pattern. The peak activity shifted in DD due to the animal's
endogenous, free-running circadian period τ = 21.27 h. **(Cii)** Mean spontaneous ORN spike
frequency across all DD animals in 1 h bins. Data from the individual in Ci is highlighted in red. **(Ciii)**
Mean spontaneous ORN spike frequency across all DD animals in 1 h bins during the first 48 hours
after aligning the time to the first maximum in spontaneous activity (see Methods) revealed
circadian activity. **(Di)** The spontaneous spike frequency of one long-term ORN recording in
constant darkness with infusion of the Orco antagonist OLC15 (orange frame) dissipated the
circadian rhythm of spontaneous activity. **(Dii)** Mean spontaneous ORN spike frequency across all
OLC15 animals in 1 h bins. Data from the individual in Di is highlighted in red. In contrast to LD and
DD conditions, the spike frequency decreases over time with prolonged exposure to OLC15. **(Diii)**
Mean spontaneous ORN spike frequency across all OLC15 animals in 1 h bins (mean ± SD) during
the first 48 hours after aligning the time to the first maximum in spontaneous activity (see
Methods). In contrast to LD and DD conditions, the circadian change in spontaneous spiking
frequency disappeared. **(E)** The slopes of the linear fits to the binned spiking activity of each
individual animal in the three different conditions (see Methods). Each dot indicates the slope for
one experiment. The line indicates the mean. The slope for OLC15 is significantly different from
control LD.

Figure 4: Blocking Orco removed circadian modulation of spontaneous ORN spiking patterns.
Some attributes of the spontaneous ORN activity were significantly different (Wilcoxon signed-rank
test, α = 0.05) between the time windows of low vs. high activity, mostly in LD (white boxes) and DD
(gray boxes) conditions, but never when Orco was blocked with OLC15 (orange boxes). In both LD
and DD, the mean spiking frequency was increased during high activity (**Ai, Aii**). Only in LD but not
in DD, the mean inter-burst intervals (IBI; **Bi, Bii**), the relative number of spikes belonging to a burst

(**Ci, Cii**) and the mean inter-event interval (IEI; **Di, Dii**) decreased significantly during the high
activity period. Both in LD and DD the mean burst duration (**Ei,Eii**), mean number of spikes per bust
(**Fi,Fii**), and mean intra-burst intervals (**Gi, Gii**) did not differ between low vs. high spiking activity.
Also, with Orco blocking, the attributes for low vs. high activity did not differ for any of the attributes
tested (**Aiii-Giii**). In DD and OLC15conditions, activity phases were aligned as described in the
Methods; thus, low activity in DD was at subjective ZT 19 and in OLC15 at subjective ZT 13, high
activity was in both cases at subjective ZT 24. ZTs for low and high activity in LD were 10 and 0,
respectively.

1007

1008

Figure 5: Blocking Orco removed circadian regulation of ultradian rhythms in the spontaneous ORN spiking pattern.

Heat map of the instantaneous frequencies (1/ISI) over two consecutive days of long-term tip-recordings of one pheromone-sensitive long trichoid sensillum under different conditions (LD **(A)**, DD **(B)**, OLC15 **(C)**) in each panel. Pixel color indicates the counts of instantaneous frequencies in that respective bin. The points fall mostly in a band of high frequencies (>10 – ~100 Hz) and low frequencies (0.01 – ~10 Hz), with the high-frequency band representing the instantaneous frequencies of spikes within a burst and the low-frequency band the instantaneous frequency between bursts. **(A, B)** The high frequency band indicates daily and circadian modulation of frequency prevalence. In addition, the low frequency band (<0.01 Hz to ~ 10 Hz) also displayed circadian modulation of the frequency composition. **(C)** Infusion of OLC15 deleted the circadian modulation in both frequency bands and enhanced the ultradian rhythm of frequency prevalence.

1021

Figure 6: Blocking Orco affected ultradian and infradian frequencies in the spontaneous ORN firing patterns.

The Fourier analysis of spontaneous spiking activity revealed rhythms with ultradian (<20 h; dark grey), circadian (20-28 h; white), and infradian periods (>28 h; light grey) in LD (**Ai**), DD (**Bi**), and OLC15 (**Ci**). Each line represents one animal where each triangular marker is at the local maximum in the frequency spectra obtained for that animal. The histograms above illustrate how often

specific periods of spiking activity rhythms occurred averaged over all animals. Circadian rhythms
were detected in LD (N = 5 of 11), DD (N = 7 of 10), and OLC15 (N = 4 of 12). Wavelet analysis of LD
(**Aii**), DD (**Bii**), and OLC15 (**Cii**) confirmed the occurrence of multiscale periods. Example plots for
one animal each, the same animals as in Figure 5. For each panel, the top plot depicts the mean
firing frequency, the middle plot the wavelet power in the same period range as panels **i,** and the
bottom plot the wavelet power for infradian periods up to 7 h on a log scale to highlight infradian
time scales.

1036

1037

Figure 7: Orco is not under control of the TTFL-based molecular circadian clock.

1039 Relative expression levels of *Orco* mRNA from male hawkmoth antennae raised under LD
1040 conditions did not differ significantly (ANOVA on ranks, $H(2) = 5.60$, $p = 0.061$) between ZT 1 (end of
1041 activity phase), ZT 9 (resting phase), and ZT 17 (beginning of the activity phase). *timeless* mRNA
1042 levels from male hawkmoth antenna served as control and were published in Schneider et al.
1043 (2025).

1044

Figure 8: The model adequately reproduces the firing pattern of the biological ORN.

(A) The model activity (bottom trace) displays both isolated spikes and bursts of variable length as in the original recording (top trace). The model represents an intracellular recording whereas the biological recordings were done with extracellular electrodes. **(B)** Semi-logarithmic plot of burst length distribution of an experimental LD recording (top) and simulated model (bottom) of spontaneous action potential activity of pheromone-sensitive neurons. Both the model and the recordings display exponential decrease of the count's density as a function of the burst length, shown as a linear decrease in the logarithmic scale.

1055

Figure 9: Comparison of the circadian modulation of model and ORN spiking attributes.

Mean spike frequency and percentage of spikes that belong to bursts show anti-phasic circadian oscillations both in the simulated model (left) and in the biological recording (right).

1059

1060

Figure 10: The model captures the Orco-dependent circadian modulation of ultradian spiking

rhythms.

Heat map of the simulated model, pixel color indicates the density of instantaneous frequencies in

that respective bin. The points can be roughly divided in a band of high frequencies (top, ~50 –

80 Hz) and low frequencies (bottom, ~0.1 – 5 Hz), with the high-frequency band representing the

instantaneous frequencies of spikes within a burst and the low-frequency band the instantaneous

frequency between bursts. **(A)** Simulated spikes with circadian regulation by the Orco channel; the

inter-burst frequencies varied widely with circadian rhythmicity while the frequencies within burst

remained approximately constant. The simulated results also reproduced the "merging" effect

between the two bands when the bottom one approaches the top one. **(B)** Simulated spikes

without circadian regulation of Orco had no circadian oscillations.

1072

9 Supplementary Material

9.1 Mathematical Model

The mathematical model is a 22-dimensional system of stochastic differential equations designed to describe the behavior of ion channels and their effects on the membrane potential. I_{LVA} represents a low-voltage-activated calcium current, while I_{Na} and I_K are the transient sodium and potassium currents responsible for action potential generation. I_{BK} is a large-conductance potassium channel that is activated by both a depolarized membrane potential and an increased intracellular calcium concentration, and I_L is the leak current. Additionally, the Orco channel is a leaky, non-specific ion channel gated by cAMP, with the cAMP concentration oscillating in a circadian rhythm. To model the Orco conductance, we used a sinusoidal function of Zeitgeber time (ZT), independent of the membrane potential, as shown in Figure 2B, C. Each of these ion channels has been reported in vitro in previous studies (Dolzer et al., 2021).

The deterministic part of the 22D model is given by the mean field of the channel-based Langevin model. This approach was introduced by Fox and Lu's (1994) for a 14D HH model. In their framework, one dimension tracks the membrane potential, while the other 13 dimensions correspond to the fraction of channels in each of the five states in the potassium (K^+) Markov chain and the eight states in the sodium (Na^+) Markov chain. Pu and Thomas (2020) demonstrated that the deterministic 14D model and the original 4D HH model are dynamically equivalent, meaning that each solution of the 4D model corresponds to a solution of the 14D model.

Expanding upon Pu and Tomas' methodology to represent out system, we added one dimension to track the calcium concentration, six dimensions for each possible state of the LVA channel, and two dimensions for each possible state of the BK channel. The state vectors for each ion channel are therefore defined as follows: a five-component vector NK for the K^+ gates, an eight-component vector NNa for the Na^+ gates, a six-component vector NLVA for the LVA gates, and a two-component vector NBK for the BK gates. The state vectors are represented as:

$$N^K = [n_{K0}, n_{K1}, n_{K2}, n_{K3}, n_{K4}]^T \in [0,1]^5 \tag{1}$$

$$N^{Na} = [n_{Na00}, n_{Na10}, n_{Na20}, n_{Na30}, n_{Na01}, n_{Na11}, n_{Na21}, n_{Na31}]^T \in [0,1]^8 \tag{2}$$

$$N^{LVA} = [n_{LVA00}, n_{LVA10}, n_{LVA20}, n_{LVA01}, n_{LVA11}, n_{LVA21}]^T \in [0,1]^6 \tag{3}$$

$$N^{BK} = [n_{BK0}, n_{BK1}]^T \in [0,1]^2 \tag{4}$$

The K$^+$ channel consists of four independent n (activation) gates, forming a five-vertex

channel-state diagram with eight directed edges. The channel conducts a current only when it is in

the rightmost state (Figure 2D). Similarly, the Na$^+$ channel is composed of three identical n gates

and one h (inactivation) gate, forming an 8-vertex diagram with 20 directed edges, one of which is

conducting. The LVA channel includes two identical n gates and one h gate, resulting in a 6-vertex

diagram with 14 directed edges, one of which is conducting. The BK channel consists of a single n

gate, forming a 2-vertex diagram with two directed edges, one of which is conducting. The Markov

chain for every channel is shown in Figure 2B.

For a sufficiently large number of channels, Schmidt and Thomas (2014) and Schmidt et al.

(2018) demonstrated that, under voltage clamp, the equations governing the system can be

approximated by a multidimensional Ornstein-Uhlenbeck (OU) process, or Langevin equation, as

follows:

$$dN^{ion} = \sum_{k=1}^{k_{max}} \zeta^{ion} \left\{ \alpha_k^{ion}(V) N_{i(k)}^{ion} dt + \sqrt{\epsilon^{ion} \alpha_k^{ion} \, N_{i(k)}^{ion} \, dW_k^{ion}} \right\}$$

(5)

1112

1113

Here ζ^{ion} is the stoichiometry vector for the kth directed edge. If we write i(k) for the source

node and j(k) for the destination node of edge k, then $\zeta^{ion} = e_{j(k)}^{ion} - e_{i(k)}^{ion}$. The voltage-dependent

per capita transition rate along the kth edge is $\alpha_k^{ion}(V)$ and $N_{i(k)}^{ion}$ is the fractional occupancy of the

source node for the kth transition. The finite-time increment in the Poisson process is approximated

by a gaussian process, namely, the increment dW_k^{ion} in a Wiener (Brownian motion) process

associated with each directed edge. These independent noise terms are scaled by $\epsilon^{ion} = \frac{1}{N_{Tot}^{ion}}$,

where $\frac{1}{N_{Tot}^{ion}}$ represents the total number of that specific ion channel.

The membrane potential is governed by the current balance equation:

$$C \frac{dV}{dt} = \sum_{\chi} I_{\chi}$$

(6)

1122

1123

1124   where the set χ contains the six ionic currents included in the model:

$$\chi \in \{LVA, Na, K, BK, L, ORCO\}$$

(7)

1125

1126

1127   Each of the six ionic currents included in the model are described as:

1128

$$I_{Na} = -g_{Na}N_8^{Na}(V - V_{Na}) \qquad (8)$$

$$I_K = -g_K N_5^K (V - V_K) \qquad (9)$$

$$I_{LVA} = -g_{LVA}N_6^{LVA}(V - V_{LVA}) \qquad (10)$$

$$I_{BK} = -g_{BK}N_2^{BK}(V - V_{BK}) \qquad (11)$$

$$I_{Orco} = -g_{Orco}\left(0.8 + 0.2\,sin(\frac{2\pi t}{24h})\right)(V - V_{Orco}) \qquad (12)$$

$$I_L = -g_L(V_L - V) \qquad (13)$$

1129
1130   Here, g_χ is the maximal conductance, V is the membrane potential, and V_χ is the reversal

1131   potential for each ion current. The parameter values for each current can be found in Table 1.

1132   Each transition rate $\alpha_k^{ion}(V)$ is calculated in the following way:

1133

$$\alpha_k^{ion}(V) = \frac{p_{j(k)\infty}^{ion}(V)}{\tau_p(V)} \qquad (14)$$

1134
1135

where $p_{j(k)\infty}^{ion}(V)$ is the steady state value of the specific gate that switches states in the

destination state and $\tau_p(V)$ is the time constant of said gate. The steady state activation values and

time constants for the LVA, Na, K, and BK channels were based on Viertel and Borisyuk (2019). The

steady state functions for the activation gates are denoted with m_∞^{ion}, and those for the inactivation

gates with h_∞^{ion}.

Each gate's steady state activation function, except for the BK channel, takes the form:

1142
1143

$$p_{j\infty}^{ion}(V) = \frac{1.0}{1.0 + exp\left(\frac{(V - \theta_p)}{\beta_p}\right)} \qquad (15)$$

1144

1145   The I_{BK} current is dependent on the intracellular calcium concentration and membrane

1146   potential as represented by the following equations:

1147
1148

$$m_\infty^{BK}(V) = \frac{1.0}{1.0 + exp\left(\frac{\left(V - V_{\frac{1}{2}}\right)}{(-15.6)}\right)} \tag{16}$$

$$V_{\frac{1}{2}} = -20 + 59.2e^{-90[Ca]} + 96.7e^{-470[Ca]} \tag{17}$$

$$\tau_\alpha = -(q - 1.0)\frac{(f - 0.2)}{0.8} + 170 \tag{18}$$

$$q = 2.9 + 6.3e^{-360[Ca]} \tag{19}$$

$$s = -25.3 + 107.5e^{-120[Ca]} \tag{20}$$

$$f = \frac{1.0}{10.0\left(e^{\frac{-(V+100.0-s)}{63.6}} + e^{\frac{(-150.0+(V+100.0-s))}{63.6}}\right)} \tag{21}$$

1149
1150

1151   The intracellular concentration of calcium ions is described by the following equation:

1152

$$\frac{d[Ca]}{dt} = -\gamma\frac{I_{LVA}}{zFd} + \frac{[Ca_0] - [Ca]}{\tau_{Ca}} \tag{22}$$

1153

1154   where Ca_0 is the steady state calcium concentration, γ describes calcium buffering, z is the

1155   ionic valence of calcium, F is Faraday's constant and d is the depth in microns at which calcium

1156   ions concentrate near the membrane.

1157   The parameters of each steady state activation function as well as the time constant for each

1158   gating value can be found in Table 2.

1159

1160   **Table 1: Parameter values for model currents.**

Current	g_X (mS/cm²)	V_X (mV)
I_L	4.3	-62.5
I_{LVA}	10.0	120
I_{NA}	29.17	45.0
I_K	12.96	-105.0
I_{BK}	9.0	-105.0
I_{ORCO}	5.0	0.0

1161

1162   **Table 2: Gating variable parameter values.**

Variable	θ	β	τ(ms)

m^{LVA}	-30	-4.8916	$\dfrac{40.0}{cosh\left(\dfrac{(V+37.1)}{9.7832}\right)}$
h^{LVA}	-59.2	11.2326	$\dfrac{350.0}{cosh\left(\dfrac{(V+59.2)}{22.4652}\right)}$
m^{Na}	-25.0	-6.5	0.2
h^{Na}	-26.0	9.0	$\dfrac{10.0}{cosh\left(\dfrac{(V+26.0)}{18.0}\right)}$
m^{K}	-26.0	-9.0	$\dfrac{10.0}{cosh\left(\dfrac{(V+26.0)}{18.0}\right)}$

1163