Orco regulates the circadian activity of pheromone-sensitive olfactory

receptor neurons in hawkmoths

Abbreviated title: Circadian regulation by Orco

Author names and affiliations:

* Aditi Vijayan, Animal Physiology and Center for Interdisciplinary Nanostructure Science and

Technology, FB 10, University of Kassel, 34132 Kassel, Germany

* Mauro Forlino, Theoretical Physics and Center for Interdisciplinary Nanostructure Science and

Technology, FB10, University of Kassel, 34132 Kassel, Germany

Yajun Chang, Animal Physiology and Center for Interdisciplinary Nanostructure Science and

Technology, FB 10, University of Kassel, 34132 Kassel, Germany

Pablo Rojas, Theoretical Physics and Center for Interdisciplinary Nanostructure Science and

Technology, FB10, University of Kassel, 34132 Kassel, Germany

Katrin Schröder, Animal Physiology and Center for Interdisciplinary Nanostructure Science and

Technology, FB 10, University of Kassel, 34132 Kassel, Germany

Anna C. Schneider, Animal Physiology and Center for Interdisciplinary Nanostructure Science and

Technology, FB 10, University of Kassel, 34132 Kassel, Germany

Martin E. Garcia, Theoretical Physics and Center for Interdisciplinary Nanostructure Science and

Technology, FB10, University of Kassel, 34132 Kassel, Germany

Monika Stengl, Animal Physiology and Center for Interdisciplinary Nanostructure Science and

Technology, FB 10, University of Kassel, 34132 Kassel, Germany

*Both authors contributed equally

Corresponding author email address: anna.c.schneider@uni-kassel.de

Conflict of interest: Authors report no conflict of interests.

Acknowledgements: We thank Dr. Dieter Wicher (Max Planck Institute for Chemical Ecology, Jena,

Germany) for generously providing OLC15, and Rishaban Radhakrishnan for help with Python

codes. We thank Prof. Hülya Altuntaş for helpful discussions.

Funding: AV, MF, YC, ACS, MEG, and MS were supported by Deutsche Forschungsgemeinschaft

RTG 2749/1: "Biological Clocks on Multiple Time Scales"; KS and MS were supported by Deutsche

Author contributions:

Conceptualization: AV, MF, MEG, MS; data curation: AV, MF, YC, PR, KS, ACS, MEG, MS; formal

analysis: AV, MF, YC, PD, ACS; funding acquisition: MEG, MS; investigation: AV, MF, YC, KS;

methodology: MF, PR, MEG, MS; project administration: ACS, MEG, MS; resources: MEG, MS;

software: AV, MF, PR, MEG; supervision: MEG, MS; validation: AV, MF, YC, PR, KS, ACS, MEG, MS;

visualization: AV, MF, YC, ACS, MS; writing – original draft: AV, MF, ACS, MEG, MS; writing –

reviewing & editing: AV, MF, YC, ACS, MEG, MS.

41   1   Abstract

The mating behavior of nocturnal *Manduca sexta* hawkmoths is under strict temporal

control. It is orchestrated via circadian and ultradian oscillations in sex-pheromone stimuli as

social zeitgeber. The extremely sensitive pheromone-detecting olfactory receptor neurons (ORNs)

that innervate the long trichoid sensilla on the male's antennae are peripheral circadian clocks.

They express the transcriptional-translational feedback loop (TTFL) circadian clockwork, best

characterized in *Drosophila melanogaster*. In hawkmoths, it is still unknown whether or how the

ORN TTFL clockwork regulates the daily rhythms in pheromone sensitivity and in temporal

resolution of ultradian pheromone pulses as prerequisites to the temporal regulation of hawkmoth

mating behavior.

We hypothesize that, rather than the slow TTFL clock, a more rapidly adaptive post-

translational feedback loop (PTFL) clockwork, associated with the ORN plasma membrane, allows

for temporal control of pheromone detection via generation of multiscale endogenous membrane

potential oscillations. The potential oscillations of the PTFL clock could rapidly synchronize to

oscillations of pheromone stimuli at different timescales, thus enabling the prediction of stimulus

patterns as a mechanism for active sensing. With *in vivo* long-term tip recordings of long trichoid

sensilla of male hawkmoths, we analyzed the spontaneous spiking activity indicative of the ORNs'

endogenous membrane potential oscillations. Consistent with our hypothesis of a multiscale PTFL

clock in hawkmoth ORNs, spontaneous spiking was modulated on ultradian and circadian

timescales, with maximum activity at night. When we blocked the evolutionarily conserved

olfactory receptor coreceptor (Orco), the circadian modulation was abolished but the ultradian

frequency modulation of the spontaneous activity remained. Consistent with PTFL control, Orco

was not under the transcriptional control of the TTFL clock, but its modulation was dependent on

cAMP. We could replicate the experimental data in a conductance-based computational model of

an ORN. In this model, Orco conductivity changed as a function of fluctuating 2^{nd} messenger levels.

This study demonstrates that a PTFL clock is sufficient to impose a circadian pattern on ORN

sensitivity.

68   2   Significance statement

It is generally assumed that all circadian rhythms in an organism are driven by a

transcriptional-translational feedback loop (TTFL) clock. In this study, we demonstrate with *in vivo*

recordings of hawkmoth pheromone-sensitive olfactory receptor neurons (ORNs) that the olfactory

receptor coreceptor (Orco) is the key pacemaker channel for controlling circadian, but not all

ultradian, rhythms in spontaneous spiking activity. Since Orco expression is not driven by the TTFL

clock, its conductance appears to be controlled by a post-translational feedback loop (PTFL)

membrane clock via 2^{nd} messengers. Accordingly, our computational model suggests that ORN

sensitivity is tuned by periodic changes in the conductivity of an Orco ion channel, which is

mediated by cycling levels of cyclic nucleotides. This highlights the role of the contribution of

posttranslational modifications to the generation of circadian rhythmicity.

3 Introduction

Biological clocks generate endogenous physiological and behavioral oscillations on multiple

timescales that persist in the absence of any external timing cues under constant conditions

(Lamont and Amir, 2010). Such clocks are essential to generate internal rhythms and to permit

their synchronization to external zeitgebers at timescales from milliseconds to years. These

synchronized endogenous multiscale clocks thus provide both a continuous time frame for

physiology and behavior - the organism's "present time" - and enable organisms to predict and

respond to regular, periodic environmental changes (Stengl and Schneider, 2024). Even though

endogenous clocks are widespread in prokaryotes and eukaryotes (Géron et al., 2024; Hall and

Rosbash, 1993; Millar, 2016; Thaiss et al., 2014), current knowledge about how multiscale clocks

are linked over widely diverging timescales remains sparse.

Insect circadian clock neurons, which are tightly connected to visual brain circuits, are

amongst the best studied biological clocks (Hall and Rosbash, 1993; Nishiitsutsuji-Uwo and

Pittendrigh, 1968; Reinhard et al., 2024). They regulate physiological and behavioral oscillations

that underly sleep-wake rhythms tied to the 24 h light-dark cycles on Earth (Dubowy and Sehgal,

2017). Circadian clock neurons possess a molecular clockwork of transcriptional-translational

feedback loops (TTFLs). This TTFL clock generates endogenous oscillations of clock gene mRNA

and protein levels with a cycle period of ~24 h via negative feedback by proteins that inhibit their

own transcription (Hardin, 2011). It is assumed for both insects and mammals that this TTFL

comprises the master clockwork that regulates all physiological and behavioral rhythms in the

circadian time range through genetically controlled output pathways (Hastings et al., 2020).

Molecular TTFL clockworks were not only found in central brain clock neurons but also in

sensory neurons, such as olfactory receptor neurons (ORNs) in insect antennae, where they play a

role in odor-dependent circadian behaviors like feeding, pollination, reproduction, and navigation

(Flecke et al., 2010; Flecke and Stengl, 2009; Krishnan et al., 1999; Merlin et al., 2009, 2006; Page

and Koelling, 2003; Plautz et al., 1997; Sauman and Reppert, 1998; Schendzielorz et al., 2015;

Schuckel et al., 2007; Tanoue et al., 2004; Zhou et al., 2005). Little is known about how the TTFL

clockwork of insect ORNs regulates circadian rhythms in chemosensory sensitivity that underlie

the circadian control of these different behaviors.

At night, male nocturnal *Manduca sexta* hawkmoths are very sensitive to the sex pheromone

blend that their conspecific females release in an intermittent pattern (Itagaki and Conner, 1988).

The intermittency is a prerequisite to maintain the male's search flight, which is suspended at

pulse frequencies >30 Hz (Baker et al., 1985; Baker and Vogt, 1988; Stengl, 2010). The pheromone

pulses are resolved of up to 3 Hz by pheromone-specific ORNs in the antenna's long trichoid

sensilla (Lee and Strausfeld, 1990; Marion-Poll and Tobin, 1992). Pheromone-specificity is provided

by odor receptor (OR) subunits in the ciliary membranes of ORNs that heteromerize with the

conserved olfactory receptor co-receptor Orco (Nakagawa and Vosshall, 2009; Stengl and Funk,

2013; Wicher and Miazzi, 2021). The insect odor transduction cascade is still under debate and

seems to involve both ionotropic and metabotropic pathways (Sato et al., 2008; Schneider et al.,

2025; Wicher et al., 2008). While there is no general agreement on the role of Orco, it is known that

in heterologous expression systems Orco homo- and heteromers form a slow, leaky, non-specific

cation channel, which depolarizes the cells (Jones et al., 2011; Sargsyan et al., 2011; Sato et al.,

2008; Wicher, 2018; Wicher et al., 2008). *In vivo* tip recordings of long trichoid sensilla in *M. sexta*

confirmed that Orco acts as leaky cation channel, acting as a pacemaker channel, because it

promotes ultradian membrane potential oscillations that underly the spontaneous action potential

activity (Nolte et al., 2013). These potential oscillations turn the ORNs into ultradian temporal

filters that are sensitive to the frequency of the pheromone pulses they encounter. Pheromone

pulses that arrive at a similar frequency as the endogenous sub-threshold membrane potential

oscillations would synchronize these oscillations and would be more likely to elicit action potential

trains. Therefore, this pacemaker channel-dependent mechanism would allow for active sensing if

endogenous membrane potential oscillations are in the physiological range that is necessary for

tracking intermittent pheromone pulses.

It is not known whether the Orco-dependent membrane potential oscillations also occur at

circadian timescales, and whether or how they are linked to ultradian potential oscillations. One

hypothesis is that the molecular TTFL clockwork of ORNs exclusively controls endogenous

circadian rhythms of the membrane potential via circadian expression of pacemaker channels like

Orco. Alternatively, the plasma membrane of ORNs could by itself constitute an autonomous clock

that is entrained by but does not rely on the TTFL clock. Constant levels of ion channels that do not

require daily degradation and daily transcription via the TTFL clock, but with modulation of open-

time probability by posttranslational modifications, would constitute a more economical and

faster adjustable mechanism to generate endogenous membrane potential oscillations (Stengl and

Schneider, 2024). This would facilitate active sensing by rapidly tuning odorant/pheromone

detection to behaviorally relevant zeitgeber signals in the insect's environment.

Here, we performed a combination of experimental and computational modeling studies to

investigate whether Orco plays a role in active sensing at the circadian and/or ultradian timescale

via generation of endogenous membrane potential oscillations. We hypothesize that Orco is part of

a post-transcriptional feedback loop (PTFL) clockwork in the plasma membrane, rather than being

TTFL controlled. To capture naturally occurring, physiologically relevant processes in insect ORNs

we performed *in vivo* long-term tip recordings of the spontaneous spiking activity of pheromone-

sensitive trichoid sensilla of intact but restrained hawkmoth males.

We corroborated the experimental data with a Hodgkin-Huxley (HH) based computational

ORN model. In general, the widely used deterministic HH models do not include circadian control

of ion channels. Moreover, the highly dynamic, bursting firing patterns of ORNs cannot be

described by the traditional deterministic modeling of neuronal activity. This constitutes a

drawback because stochasticity plays a crucial role in shaping firing patterns which are influenced

not only by external stimuli but also by the inherent noise involved in channel gating. Therefore, to

address the above points realistically, we developed a Langevin formulation of the HH model,

which explicitly accounts for the noise introduced by the random opening and closing of ion

channels and circadian modulation of the Orco channel conductivity. By integrating channel noise,

we could explore how circadian regulation of ionic currents contributes to the variability of firing

patterns and how this noise interacts with external stimuli to influence olfactory signal processing.

Our model aims to highlight the advantages of using a stochastic framework for modeling the

circadian activity of ORNs. The combined experimental and modeling efforts generate a more

complete picture and predictions concerning these autonomous processes in sensory neurons,

which are required for future exploration of physiologically relevant anticipatory behaviors that are

driven by biological clocks.

4 Methods

4.1 Animals

Nocturnal *M. sexta* hawkmoths were bred and raised from eggs in the rearing facility at the

University of Kassel. Males and females were separated during pupal stages. Males were housed

isolated from the females to avoid exposure to female pheromones. All experiments were

performed with adult males reared in a long-day photoperiod (L:D 17:7 h to prevent diapause) at

25 °C with relative humidity of about 55%. Larvae were fed with an artificial diet modified after Bell

and Joachim (1976); adult moths could feed on sugar solution with added vitamins (Roth) *ad*

libitum (Riffell et al., 2008).

While all animals were exposed to the same lighting regime as zeitgeber to entrain them up

until the start of the experiments, we also attempted pheromone exposure as additional circadian

zeitgeber to phase-align spontaneous ORN activity in a subset of males (Ghosh et al., 2024).

Without allowing direct contact, virgin males were exposed at zeitgeber time (ZT) 16 to females and

naturally occurring pheromone blends in closed mating cages for 30 mins. Males were then moved

back into isolation before tip recordings (see below) were started at ZT 17 on the same day. Since

we focused on spontaneous spiking activity rather than on pheromone-dependent responses and

because repetitive or long pheromone exposure changes the composition of 2^{nd} messenger

modulated ion channels in ORNs (Gawalek and Stengl, 2018), we did not employ any further

pheromone synchronization protocols.

4.2 Solutions

Ringer compositions were taken from (Pézier et al., 2007) and contained (in mM): 6.4 KCl, 12

$MgCl_2$, 1 $CaCl_2$, 12 NaCl, 10 HEPES, 340 glucose for hemolymph ringer (HLR) at 450 mosmol/kg,

and 172 KCl, 3 $MgCl_2$, 1 $CaCl_2$, 25 NaCl, 10 HEPES, 22.5 glucose for sensillum lymph ringer (SLR) at

475 mosmol/kg, both at pH 6.5.

The Orco antagonist OLC15 (N-(4-butylphenyl)-2-((4-ethyl-5-(2-pyridinyl)-4H-1,2,4-triazol-3-

yl)thio)acetamide (Chen and Luetje, 2012); kindly provided by Dr. D. Wicher, Max Planck Institute

for Chemical Ecology, Jena, Germany) was dissolved in DMSO to a concentration of 100 mM (stock

solution stored at 4-8 °C) before being diluted in SLR to a final concentration of 50 µM to be used

within one week (stored at 4 °C). We determined in previous studies that, compared to other Orco

antagonists, OLC15 is the most specific (Chen and Luetje, 2012; Nolte et al., 2016, 2013; Pask et

al., 2013). It is effective at 50 µM, according to dose-response curves, and dose-dependently

antagonizes the action of the Orco agonist VUAA1 in intact hawkmoths (Nolte et al., 2016, 2013).

DMSO percentage was 0.05% in the final solution.

The cAMP analog 8-bromo-cAMP acetoxymethyl ester (8-Br-cAMP-AM; Biolog Life Science

Institute GmbH & Co. KG) was dissolved in DMSO to a concentration of 10 mM (stock solution

stored at -20°C) and diluted to a final concentration of 100 mM in either SLR or SLR + 50 µM OLC15.

4.3   Electrophysiology

The spontaneous electrical activity of the unstimulated pheromone-sensitive ORNs in a

single long trichoid sensillum was recorded extracellularly as *in vivo* tip recording (Kaissling et al.,

1987). 1-2-d-old male *M. sexta* that were raised and entrained to zeitgebers as described above

were fixed in a custom-made Teflon holder with adhesive tape. The right antenna used for

recordings was oriented with its dorsal surface facing up and immobilized near the base with

dental wax (Boxing wax strips, KerrHawe SA). Electrodes (Ag/AgCl wire in a glass pipette filled with

SLR for the recording electrode and HLR for the reference electrode) were pulled (DMZ-Universal

Puller, Zeitz Instruments) from borosilicate capillaries (OD:1.56mm, ID: 1.17mm, without filament;

Science Products) to a tip diameter of about 2 µm. All recordings were made at a distance of 2/3 to

3/4 of the antenna length towards its distal end because pheromone-sensitive long-trichoid

sensilla (sensilla trichodea type 1) occur in a characteristic, homogenous pattern along the

hawkmoth antenna (Kaissling et al., 1989; Kalinová et al., 2001; Lee and Strausfeld, 1990).

Furthermore, all ORNs in long trichoid sensilla express the same Orco staining pattern (Nolte et al.,

2016), share largely overlapping ion channel populations, and obvious differences in the activity of

ORNs from different annuli have not been reported (c.f., Dolzer et al., 2003; Gawalek and Stengl,

2018; Schneider et al., 2025). Thus, it is very likely that pheromone encoding mechanisms are not

locally restricted along the flagellum. Recording and reference electrodes were located on the

same annulus: After cutting off the distal 20 annuli of the antenna with micro scissors, the

reference electrode was inserted to a depth of 2 annuli into the antennal lumen from the cut end.

The cut was covered with electrode gel (GE Medical Systems Information Technology) to prevent

desiccation. Under microscopic control, some tips of the long trichoid sensilla of the annulus were

cut off with sharpened Dumont forceps, and the recording electrode was slipped over a single

sensillum of an upper sensillar row. Electrodes were moved with manual micromanipulators (MMJ,

Märzhäuser Wetzlar; and Leitz, Leica Microsystems) or with motorized micromanipulators

(Microstar, Scientifica) with a joystick controller (Scientifica).

ORN activity was amplified 200-fold (custom built amplifier with $10^{12}\,\Omega$ input impedance, or

ELC-01 MX in combination with DPA-2FX, npi, or EXT 10-2F, npi), low pass filtered at 1.3 kHz,

digitized at 20 kHz with a Digidata (1550A or 1550B, Molecular Devices), and saved for offline

analysis with Clampex 10 (pCLAMP suite, Molecular Devices).

All tip recording experiments were performed at room temperature (21-27 °C) and a relative

humidity of 35-60%. These environmental fluctuations across experiments were based on

seasonal influences. Throughout each experiment, conditions were stable. To ensure that neither

electrode reattachment to the same trichoid sensillum nor application of DMSO influenced spiking

activity, we performed two sets of paired experiments at ZT 1-3 in which we recorded the

spontaneous activity before and after removing and reattaching the SLR-filled recording electrode

to the same sensillum, or recorded the spontaneous activity of the same sensillum in SLR and SLR

+ 0.15% DMSO. Both treatments did not affect spontaneous spiking activity (Figure 1B).

We recorded ORN activity either in long-day conditions (17:7 LD) or constant darkness (DD)

to examine the free-running peripheral ORN clock. For LD experiments, the light regime continued

as in the rearing facility during the experiment. For DD and OLC15 experiments, the lights were

turned off at the beginning of the recording, i.e., all animals were exposed to the same light

zeitgeber until the beginning of the experiment. To examine the effect of Orco on the spontaneous

action potential (AP) activity pattern, we used the recording electrode to infuse OLC15 in DD

conditions. To examine the effect of cAMP on Orco gating, we recorded the spontaneous ORN

activity at the end of the moth's activity phase (ZT 1-3) in SLR as control, and either 8-Br-cAMP-AM,

or 8-Br-cAMP-AM + OLC15, infused through the recording electrode.

ORNs generate spontaneous APs. However, high spiking activity during control recordings is

indicative of stress, whereas low/no activity could be indicative of damage. Therefore, we only

considered experiments in which the average spiking frequency was between 2 Hz and 0.008 Hz in

control conditions.

4.4   Data analysis

To characterize the spontaneous ORN spiking activity, we first detected spikes with a self-

written program in Python. Using the scipy.signal package and find_peaks function, we used a

moving mean with 100 samples (5 ms) window size for baseline correction, followed by applying a

8th order low-pass Bessel filter (2000 Hz cutoff frequency) to the raw voltage trace to facilitate spike

detection within the following conditions: spike full width (3-15 ms), threshold search for amplitude

of spike (0.5-2.5 mV) and threshold prominence (0.1). Two or more consecutive spikes with

intervals ≤ 50 ms were considered a burst (Dolzer et al., 2001). We then calculated the following

attributes in bins of either 10 min or 1 h for further analysis: instantaneous spike frequency (ISI,

1 / inter-spike interval), inter-burst interval (IBI, time between two consecutive bursts with no single

spikes in between), inter-event interval (IEI, time between two events, i.e. two consecutive single

spikes, two consecutive bursts, a burst and a consecutive spike, and vice versa), burst duration

(time from first to last spike in a burst), percentage of all spikes in the experiment that are members

of burst, and mean number of spikes per burst (Figure 1A). All statistics were done with those

binned values.

Hawkmoth behavioral and physiological rhythms desynchronize across the population in the

absence of zeitgebers like circadian rhythms of pheromone stimulation and daily light-dark cycles.

Therefore, we first used RAIN analysis (Thaben and Westermark, 2014) to identify circadian

rhythms (24 ± 4 h) in the attributes above for each animal and condition (LD, DD, OLC15) in the 1 h-

binned data. Data was downsampled to 2 h resolution and thus included every other 1 h bin to

speed up computational time. To further quantify the circadian changes in attributes across

animals, it was then necessary to phase align the recordings obtained in DD and OLC15. To

achieve this, we used an optimization procedure to calculate the optimal temporal lags for each

dataset, aiming to maximize the overall cross-correlation between the calculated binned spike

frequency series. The optimization was performed using simulated annealing, which provided a set

of time shifts (lags) for each series. These lags were applied to shift the frequency data in a manner

that maximized the synchronization between the different subjects' data, resulting in a "subjective

ZT" with subjective ZT 0 at the first maximum of the binned ISI for each individual. The objective

function used for optimization was the negative sum of pairwise correlations between all series

after applying the shifts. The resulting alignment allowed us to compare the frequency patterns

across the different conditions.

Once the data were aligned, we computed the mean frequency across all subjects for each

bin and each condition. Then, the first high activity period was identified by finding the maximum in

the mean frequency series. Subsequently, we identified the second high activity period as the time

24 hours after the first one and the first low activity period was found by locating the minimum in

the mean frequency series between the two high activity periods. Besides the mean frequency, we

also searched for significant oscillations in IBI, IEI, burst duration, percentage of spikes that are

part of bursts, and mean number of spikes per burst. For this we found the maximum value of these

statistics within a 3 h range centered in the first low activity period and compared it with the values

at the second high activity period.

Continuous wavelet and Fourier analysis were performed using PyBoat (version 0.9.12)

(Mönke et al., 2020) to examine the time series of neuronal firing frequency. The continuous

wavelet method allows us to detect and characterize rhythmic patterns within the data by

identifying frequency components that change over time. Specifically, we used wavelet analysis to

determine whether a neuron exhibited circadian and ultradian rhythms. Unlike traditional Fourier

analysis, which provides an overall frequency composition but assumes stationarity, wavelet

analysis is particularly useful for biological signals because it can reveal how these frequencies

evolve dynamically. By applying this approach, we could assess not only the presence of ultradian

rhythms but also whether their strength or occurrence fluctuated in a circadian manner, indicating

modulation by the 24-hour cycle. For this analysis, we used the 10 min-binned data to have

sufficiently high resolution to detect ultradian rhythms. Periods were determined as local maxima

in the Fourier power spectra.

4.5   Real-time quantitative polymerase chain reaction (qPCR)

We followed the qPCR methodology detailed in (Schneider et al., 2025). Briefly, male moths

were isolated and cultured until the second day after eclosion, and the antennae from four

individuals were collected every 4 hours for one day, starting at ZT 0, using liquid nitrogen, with four

biological replicates per time point. Total RNA was extracted using TRIzol® Reagent (Invitrogen) and

quantified for concentration and purity with the NanoDrop ND-1000 spectrophotometer (Thermo

Fisher Scientific). First-strand complementary DNA (cDNA) was synthesized using the PrimeScript™

RT reagent Kit with gDNA Erase (Takara Bio Inc.). Primers for *Orco* (F: GCTCGCTACCACCAAATTGC,

R:TCGTGACCCAACTGACAACA; Gene ID: LOC115452348) and for *timeless* (*tim*) (F:

TTAAGCCGACCGTAGTGCTG, R: CGTCTTCCGTCCATGTGTCT; Gene ID: LOC115446009) were

designed using the Primer-BLAST online program of NCBI. The qPCR was conducted following the

protocol of TB Green® Premix Ex Taq™ II (Tli RNase H Plus) (Takara Bio Inc.), with the program: 95 °C

for 30 s, followed by 40 cycles of 95 °C for 5 s, and 60 °C for 30 s. Relative expression levels of *Orco*

were calculated using the $2^{-\Delta\Delta Ct}$ method (Livak and Schmittgen, 2001).

4.6 Statistics

All statistics were done with self-written python codes, SigmaPlot 12 (Systat Software), or

JASP (versions 0.19.03 and 0.95.4, University of Amsterdam) with a significance level α = 0.05.

Unless noted otherwise, data were checked for normal distribution with Anderson-Darling or

Shapiro-Wilk tests and for equal variance with Levene's test. If one of these assumptions failed, we

used non-parametric tests, otherwise we used parametric tests.

To assess the statistical significance of observed differences in neuronal activity of the

aligned data between ZTs, we applied the Wilcoxon signed-rank test to pairwise comparisons

between the first low activity period and the subsequent high activity period and discarded the

transient high activity at the beginning of most recordings.

For paired recordings (electrode reattachment, addition of DMSO, or cAMP/cAMP+OLC15 in

the same animal, we used either paired student-t test or Wilcoxon signed-rank test, depending on

normal distribution of data.

For other data, we used an ANOVA on ranks with either Tukey's or Dunn's post-hoc test for

multiple comparisons as indicated.

4.7 ORN Model

In this study, we introduce a single-compartment Hodgkin-Huxley (HH) type model of the

spontaneous activity of *M. sexta* ORNs (available on GitHub (Forlino et al., 2025)). The model

contains 5 sets of ion channels that were previously identified in electrophysiological recordings

(Dolzer et al., 2021): the fast sodium (Na^+) and potassium (K^+) channels responsible for the

generation of spikes, low voltage-activated calcium channels (LVA) responsible for the initiation of

bursts, calcium-gated potassium channels (BK) responsible for the termination of bursts, and the

Orco channel, which is a voltage-independent non-specific, cAMP-gated channel. A schematic of

the model is provided in Figure 2A. We modeled Orco as a ZT-dependent conductance to represent

the effect of the oscillations in cAMP concentration that were identified in both hawkmoth and

cockroach antennae (Schendzielorz et al., 2014, 2015, 2012) (Figure 2B, C). For simplicity we chose

to use a sine function to directly represent the oscillations in Orco conductivity, but similar results

can be achieved by modeling it as an output of a Goodwin model of the circadian clock via cAMP

concentration.

Each type of ion channel possesses one or more gates that can open and close individually

with transition rates α and β for opening and closing, respectively, of activation gates (n, m) and

inactivation gates (h). As a result, each possible combination of open and closed gates defines a

possible state in which each ion channel can be found. A schematic of the Markov chain depicting

all possible states and transitions for each ion channel is shown in Figure 2D. Each ion channel can

conduct current only in the state where all its gates are open.

First, we constructed a deterministic 22-dimensional model to serve as the mean field of the

Langevin model following the methodology proposed by Pu and Thomas (2020). Here, one

dimension tracks the membrane potential, another the intracellular Ca^{2+} concentration, and the

remaining 20 dimensions track the number of channels that are in a specific state of the Markov

chains. Considering the conservation of the total number of ion channels, each type of ion channel

required one less dimension than the total possible states.

The complete Langevin model adds to the mean field an independent noise source for each

transition in the Markov chain. These noise sources are biophysically motivated since the transition

between ion channel states is a stochastic process, with the probability of transition determined by

the kinetics of each kind of ion channel. The variability in the spiking behavior of a neuron emerges

from this non-deterministic gating of individual ion channels. The detailed mathematical

description of the model can be found in the Supplementary Material.

5   Results

We tested the hypothesis that sensory neurons, such as insect ORNs, perform active,

anticipatory sensing based upon an endogenous plasma membrane clock with the pacemaker

channel Orco as the core element. We focused on pheromone-sensitive long trichoid sensilla of

male *M. sexta* hawkmoth antennae which express daily rhythms in pheromone sensitivity and

temporal resolution that are controlled by a circadian clock. We performed minimally invasive *in

vivo* tip recordings over the course of several days to search for daily rhythms in spontaneous

spiking activity as a measure of endogenous membrane potential oscillations.

5.1 Pheromone-sensitive hawkmoth ORNs exhibit Orco-dependent circadian rhythms in spontaneous spiking activity

The ORNs that innervate the pheromone-sensitive long trichoid sensilla on male *M. sexta* antennae display spontaneous spiking patterns of irregular bursts with interspersed single action potentials in short-term recordings (Dolzer et al., 2001). In the present study we documented the same pattern in long-term recordings over the course of 2-7 days. This enabled us to search for daily and circadian rhythmicity in spontaneous ORN spiking activity in the absence of pheromone stimulation.

RAIN analysis of the spontaneous spiking patterns revealed significant daily rhythms in 7 of 11 animals under long-day conditions (17:7 h LD) (Figure 3A). The maxima in spontaneous activity occurred mostly during the dark phase when nocturnal hawkmoths are active (Figure 3A (top), Bi light blue arrow), and minima occurred during the light phase when hawkmoths rest/sleep (Figure 3A (bottom), Bi dark blue arrow). High spiking activity in the dark phase had a mean maximum frequency of ~3 Hz (delta range: 0.5 – 4 Hz) (Figure 3Bii). In contrast, the minimum spiking frequencies during the hawkmoth's resting phase were below 0.5 Hz (Figure 3A (bottom)).

Since ORN firing patterns had a clear daily rhythm in LD both in individual animals (Figure 3Bi) and averaged across animals (Figure 3Bii, Biii), we examined whether these rhythms depended on cycling light cues or whether they persisted in constant darkness (DD), as expected for an endogenous, circadian clock-driven rhythm. In DD, the spontaneous ORN firing pattern remained rhythmic (RAIN: 8 of 12 animals), albeit with less robustness: The single peak of high spiking activity that was observed at the same ZT across several days in LD divided into multiple bouts of high spiking activity in the activity phase in 8 of 12 animals (Figure 3Ci). Also, in DD conditions, the high spiking activity phase shifted into the subjective day due to the individual endogenous circadian periods of about 23.5 +/- 2.8 h (21.27 h in Figure 3Ci). Using pheromone exposure as additional zeitgeber also did not synchronize the animals (Figure 3 – Supplement). Thus, averaging the instantaneous spiking activity across all DD animals did not depict clear circadian rhythms (Figure 3Cii). Therefore, we phase-aligned the mean frequency of spontaneous activity (see Methods). After alignment, circadian rhythms were visible across DD animals (Figure 3Ciii), comparable to the daily rhythms in LD (cf. Figure 3 Biii and Ciii).

Next, we examined whether Orco, a leaky, non-specific cation channel, is the dominant depolarizing pacemaker current of ORNs that drives circadian rhythms in spontaneous spiking activity. Infusion of the Orco antagonist OLC15 into the sensillum lymph obliterated circadian

rhythms and attenuated the spontaneous activity in several, but not all experiments (RAIN: 2 of 12

animals expressed circadian rhythms) (Figure 3Di). This attenuation resulted in a linear decrease in

spiking activity over several days (Figure 3Dii, Diii). Even after phase-aligning the individual

experiments, no circadian rhythm was present in the average across all OLC15-treated animals

(Figure 3Diii).

Because OLC15 is not membrane-permeable on its own it was infused with low

concentrations (0.05%) of DMSO through the recording electrode into the sensillum lymph and,

therefore, effectiveness increased over time. Hence, we fit the spiking activity of each animal and

in each condition with linear regression lines over the whole recording time to compare the slopes

of the fits (Figure 3E). In LD, the mean ± SD slope was 0.004 ± 0.006 Hz/h, indicating that the

average spiking activity remained similar throughout the duration of the recording. In DD, the slope

was -0.003 ± 0.018 Hz/h and was not different from LD. In contrast, the mean slope during OLC15

treatment was -0.007 ± 0.022 Hz/h, significantly smaller than in LD (ANOVA on ranks with Dunn's

post-hoc test; $H(2) = 7.681$, $p = 0.021$). The negative slope corroborated the finding that spiking

activity decreased over time in the presence of the Orco blocker.

Activity rhythms between hawkmoths desynchronize and have only weak phase coupling in

the absence of pheromone stimulation, even in LD cycles (Ghosh et al., 2024). Therefore, we first

determined the percentage of animals that expressed circadian rhythmicity (period of 24 ± 4 h) for

each attribute with RAIN (Figure 4A). In LD, all attributes except #AP per burst and intra-burst

frequency showed daily modulation in most animals. Circadian rhythms were still present in more

than half of the animals in DD for mean spiking frequency, mean burst frequency, and mean event

frequency. In contrast, the attributes of only a few animals remained rhythmic with the Orco

channel blocker OLC15.

To further quantify the circadian differences, especially of the desynchronized animals, we

compared different attributes of spontaneous activity between periods of high and low activity, for

DD conditions after phase alignment (see Methods) instead of comparing fixed ZT or circadian time

(CT) intervals. Despite the high variability in ORN spiking between individuals, the mean frequency

of spontaneous ORN activity was significantly higher during the high activity phase than during low

activity in both LD and DD recordings (Figure 4Bi, Bii). In LD, but not in DD, the frequency of bursting

increased significantly during the high activity phase (Figure 4Ci, Cii). The percentage of spikes that

are part of bursts was less variable and lower during the high activity phase in LD but not

significantly different in DD (Figure 4Di, Dii). The mean burst duration was less variable during the

high than the low activity phase in LD, but not significantly shorter in any condition (Figure 4Ei, Eii).

Furthermore, both in LD and DD, the mean inter-event frequencies were significantly higher during

the high activity phase (Figure 4Fi,Fii), indicating a significantly higher overall spontaneous spiking

activity during the hawkmoth's activity phase. The mean number of spikes per burst, and intra-

burst frequencies did not differ significantly between high or low activity periods in both LD and DD

conditions (Figure 4Gi, Gii, Hi, Hii). The addition of Orco antagonist OLC15 deleted any significant

differences of spontaneous activity found in LD and DD (Figure 4Biii-Hiii). In conclusion, the mean

frequency of spontaneous spiking and the frequency of bursting expressed circadian modulation,

and are both most likely controlled via a circadian clock that involves Orco.

5.2   Orco imposes circadian modulation on the ultradian rhythms in spontaneous ORN activity

After establishing that Orco influences the circadian changes in ORN spiking activity between

the animal's resting and activity phase we examined if Orco also influences ORN spiking rhythms

on the largely different ultradian timescales in the frequency range between 0.01-1000 Hz. In all

conditions, two broad bands of ultradian instantaneous frequencies were evident: one upper band

of >10 to ~100 Hz (beta/gamma range) and a lower band between 0.1 to ≤10 Hz (delta/theta/alpha

range) (Figure 5). The upper band primarily represents the frequencies of spikes within a burst,

while the lower band represents single spikes and frequencies between bursts.

In LD, two types of daily/circadian modulation of the ultradian frequencies occurred (Figure

5A). The upper frequency band was mostly modulated in prevalence (its frequency of occurrence)

but not in its frequency range. The lower band was modulated both in prevalence and frequency

range, which resulted in a sinusoidal change in the mean of the frequency band over 24 h. During

the hawkmoth's activity phase, higher frequencies of up to ~10 Hz occurred, while frequencies

dropped below 1 Hz during the resting phase. Because of this circadian increase in the lower

frequency band, it merged with the high frequency band when both bands had maximum

prevalence during the activity phase around ZT 0.

The circadian modulation of both frequency bands was maintained in constant darkness

(Figure 5B) but phase-shifted with respect to the projected ZT due to the endogenous circadian

period of each individual. Again, this indicates that a clock is modulating the spontaneous ORN

spiking activity over the course of a day. The addition of the Orco antagonist OLC15 deleted any

circadian modulation of either frequency band. Interestingly, the instantaneous frequencies in

both bands still showed correlated changes in prevalence and merging, but now at ultradian

periods only.

Additional Fourier analysis (see Methods) of spontaneous spiking activity over the course of

several days revealed faster ultradian and slower infradian rhythms, ranging from 0.6 h to 38 h, in

addition to the circadian (24 ± 4 h) rhythm (Figure 6Ai, Bi, Ci). Some rhythms (factorials of 24 h, like

12 h, 6 h, etc.) appeared to be harmonics of the circadian rhythm, which were not excluded in our

analysis. Most animals expressed ultradian rhythms of 2-4 h in LD as well as DD conditions. Fourier

confirmed the decrease of animals with circadian spiking activity in OLC15 (58.3% in DD, 33.3% in

OLC15). Please note that discrepancies in the number of animals with circadian rhythms between

RAIN and Fourier/wavelet analyses are due to the fundamentally different sensitivities in

methodology. In brief, Fourier analysis assumes a stationary sinusoidal component, whereas RAIN

as a rank-based umbrella test detects consistent raise-fall patterns, including asymmetric

rhythms. Nonetheless, both methods confirmed a dramatic decrease in animals with circadian

spiking patterns in OLC15. In addition, 33.3% of animals expressed infradian spiking patterns (up

from 0% in DD) with OLC15. Ultradian periods could be detected in all animals in all conditions but

their distribution did not differ between conditions (1-way ANOVA;$F_{(2, 117)} = 0.87$, $p = 0.421$).

These results were corroborated by continuous wavelet analysis, which provided time-resolved

spectral power across different rhythmic periods (Figure 6Aii, Bii, Cii).In control LD and DD

conditions, wavelet power for ultradian periods increased during times of elevated firing rate and

decreased during periods of low activity (Figure 6Aii and Bii, middle and bottom panels). This

dynamic pattern indicates that ultradian rhythms were not constant over time but were instead

modulated with a circadian rhythm, becoming more prominent during circadian peaks in neuronal

activity. Such circadian gating of ultradian power was clearly visible in the wavelet spectrograms as

alternating bands of higher and lower ultradian power aligned with daily activity cycles. Following

Orco blockade with OLC15, this modulation was disrupted (Figure 6Cii middle panel): although

ultradian components persisted (Figure 6Cii bottom panel), they no longer showed systematic

variation with the circadian cycle, suggesting a decoupling of ultradian rhythms from circadian

control.

5.3   Orco expression is not under the control of the molecular TTFL clockwork

So far, our results indicated a prominent role of Orco in the circadian modulation of ORN

spiking activity. Since it is the general opinion that the master circadian clock is the TTFL clock that

dominates all physiological and behavioral circadian rhythms via transcriptional control, we

examined whether Orco transcript abundance shows a daily rhythm, as previously found for clock

genes in hawkmoth antennae (Schneider et al., 2025; Schuckel et al., 2007). However, in contrast

to the circadian clock gene *tim*, qPCR of whole male hawkmoth antennae across different ZTs

revealed no significant daily rhythm of Orco transcript abundance (Figure 7; ANOVA on ranks,

H(5) = 6.91, p = 0.227). Thus, we suggest that Orco function is controlled via a posttranslational

circadian clockwork mechanism that is associated directly with the plasma membrane (Stengl and

Schneider, 2024; Stengl and Schröder, 2021).

5.4 Circadian changes in Orco conductance are sufficient to modulate spiking activity in a computational model of a hawkmoth ORN

The maximum conductivity of *Drosophila* Orco as a leaky ion channel is modulated by cAMP

(Getahun et al., 2013; Martín et al., 2001; Sargsyan et al., 2011; Wicher et al., 2008). In hawkmoth

antennae, cAMP levels oscillate throughout the day and are involved in sensitizing the ORNs to

pheromone (Dolzer et al., 2021; Flecke et al., 2010; Flecke and Stengl, 2009; Schendzielorz et al.,

2014, 2015). Hence, cAMP modulation of Orco conductance could be a putative mechanism for

the Orco-dependent circadian modulation of ORN spiking activity, integrating Orco in a circadian

PTFL based membrane clockwork (Stengl and Schneider, 2024).

To test the hypothesis that Orco is a cAMP-dependent, leaky ion channel, we constructed an

ORN model (Methods and Supplementary Material) and tuned the parameters to provide a

quantitative match with the biological data. The developed model provides a unified description of

experimental observations. For instance, the ORN model yields irregular spiking activity with a

bimodal distribution, consisting of both bursts and isolated spikes, successfully reproducing the

experimental recordings (Figure 8A). This suggests a unified mechanism underlying the initiation of

both phenomena: If the pacemaker-driven depolarization reaches the Na^+ threshold a firing

process begins, and at this point randomness determines if other slower ion channels responsible

for bursting are sufficiently activated to sustain a burst of several spikes.

Additionally, the ORN model replicates the distribution of the number of spikes per burst,

which follows a decreasing exponential trend in both experimental and simulated results (Figure

8B).The model also successfully reproduces the circadian modulation of spiking activity: the

maximum firing rate occurs when the Orco conductance is maximal, whereas the maximum

percentage of spikes that are part of bursts correlates with the lowest Orco conductance, resulting

in antiphase oscillations, as were observed in the experimental recordings (Figure 8C).

The frequency prevalences in the model over several days clustered in two distinct frequency

bands: an upper band that corresponds to the intra-burst frequencies and a lower band,

corresponding to inter-burst frequencies (Figure 8D), in agreement with the biological experiments

(Figure 5A, B). When circadian oscillations in Orco conductance were included, inter-burst

frequencies showed significant circadian variations, whereas intra-burst frequencies remained

relatively constant, reminiscent of the biological data. Notably, the simulation captured the

experimentally observed "merging" effect, where the low frequency band approaches the top band

during high-activity periods (Figure 8D).

This merging of spiking and bursting frequencies suggests that the system approached a

bifurcation, transitioning from a bursting regime to tonic spiking activity. Further evidence for the

near bifurcation condition of the neurons is the distribution of number of spikes per burst that

followed a decreasing exponential trend in contrast to a constant value, or a bounded range of

spikes per burst, which would be expected from a standard bursting neuron model (Figure 8B). The

observation that the circadian modulation of spiking activity requires only one parameter, namely

changes in the Orco conductivity, to describe the circadian effects on the ORN spiking pattern

indicates the crucial role of the Orco channel in circadian control.

5.5   cAMP regulates the circadian function of Orco

Our computational model predicted that the daily/circadian oscillations in cAMP levels

(Schendzielorz et al., 2015) control the circadian modulation of spontaneous spiking activity of

hawkmoth ORNs. To test this prediction in hawkmoth ORNs, we increased cAMP levels by infusing

8-Br-cAMP-AM through the recording electrode into the sensillar lymph (Figure 9) at ZT 1-3. This

significantly increased the mean frequency of the spontaneous spiking activity (Figure 9A;

Wilcoxon signed-rank test Z = -2.701; p = 0.004). To check whether this increase in mean frequency

was caused by direct or indirect interaction of cAMP with the Orco channel, we infused the Orco

antagonist OLC15 together with 8-Br-cAMP-AM through the same recording electrode. Blocking

Orco prevented a significant change in the spontaneous spiking frequency via the cAMP analogue

(Figure 9B; paired t-test t(9) = -1.446; p = 0.182). This further corroborates cAMP-dependent

modulation of the circadian pacemaker channel Orco.

6   Discussion

With *in vivo* long-term tip recordings of pheromone-sensitive long trichoid sensilla of male *M.*

sexta hawkmoth antennae, we searched for a membrane-associated PTFL clockwork component

that orchestrates membrane potential oscillations to facilitate active sensing of pheromone

patterns during mating flights. Indeed, we detected interlinked circadian and ultradian frequencies

in the spontaneous spiking activity, indicative of underlying membrane potential oscillations in

ORNs. We found that the olfactory receptor co-receptor (Orco) acts as a pacemaker channel that

controls the circadian modulation of the spiking activity without affecting the ultradian frequency

ranges that are relevant for the primary fast, phasic pheromone response (Dolzer et al., 2003; Nolte

et al., 2016; Schneider et al., 2025). Our computational model confirmed that the circadian

modulation of open-time probability of Orco as a leaky pacemaker channel was sufficient to

account for our experimental results. Since Orco was not under circadian transcriptional control of

the TTFL present in hawkmoth ORNs, our findings are consistent with our novel hypothesis that the

ORN's plasma membrane constitutes an endogenous, plastic post-translational feedback loop

(PTFL) clockwork. Consistent with experimental data (Schendzielorz et al., 2015), we propose that

PTFL-generated endogenous multiscale membrane potential oscillations comprising interlinked

oscillations in Ca^{2+} and cAMP levels provide circadian modulation of Orco conductance rather than

TTFL-commanded circadian Orco transcription and turnover.

The expression of a 2^{nd} messenger-dependent pacemaker/leak channel in ORNs that does

not require daily transcription would constitute an economic and rapidly adjustable mechanism of

active sensing. It could generate endogenous membrane potential oscillations that are tuned to

detect and resonate with behaviorally relevant temporal patterns of zeitgeber signals, such as the

circadian and ultradian pheromone fluctuations in the insect's environment (Stengl and Schneider,

2024; Stengl and Schröder, 2021). The present study provides the first evidence for a systems view

of biological timing that involves a PTFL membrane clock in insect sensory neurons, instead of the

prevailing hierarchical view that assumes all endogenous circadian oscillations are outputs of the

genetic TTFL clock (Stengl and Schneider, 2024; Stengl and Schröder, 2021).

6.1   Orco acts as a pacemaker channel that controls membrane potential

oscillations underlying spontaneous ORN spiking activity

The resting membrane potential of most silent neurons lies close to the negative equilibrium

potential for potassium ions, mainly due to potassium leak channels (Goldstein et al., 2001; Hille,

2001; Patel and Honoré, 2001; Talley et al., 2001). Spontaneously active neurons require the

expression of leak or pacemaker channels that open at the hyperpolarized resting membrane

potential, thus depolarizing the neuron(Bose et al., 2014; Cochet-Bissuel et al., 2014; Das et al.,

2016; Golowasch et al., 2017; Lüthi and McCormick, 1998; Ratliff et al., 2021; Robinson and

Siegelbaum, 2003; Sharma et al., 2023). When the pacemaker-driven depolarization reaches spike

thresholds, voltage-gated Na^+, K^+ and Ca^{2+} channels promote Ca^{2+} influx and spiking. Voltage- and

Ca^{2+}-gated hyperpolarizing K^+ channels then repolarize the neuron to negative membrane

potentials, re-starting the cycle. Thus, membrane potential oscillations with interlinked oscillations

in intracellular Ca^{2+} levels drive spontaneous spiking activity which depends on the type of

pacemaker channel and antagonistic ion channels present.

In insect ORNs, at least two types of pacemaker channels exist: the highly conserved, slow,

leaky Orco, which we focus on in this study, and an inversely voltage-dependent HCN type channel

(Butterwick et al., 2018; Dolzer et al., 2021; Kodirov, 2022; Lee and MacKinnon, 2017; Mandala et

al., 2025; Nolte et al., 2016, 2013; Sato et al., 2008; Stengl and Funk, 2013; Stengl and Schneider,

2024; Stengl and Schröder, 2021; Wicher et al., 2008; Wicher and Miazzi, 2021). When expressed *in*

vitro in heterologous expression systems, *Drosophila* as well as *Manduca* Orco homo- and

heteromers assemble into leaky, non-specific cation channels with a reversal potential around 0

mV, constitutively leaking Na^+, K^+, and Ca^{2+} into ORNs (Nolte et al., 2016, 2013; Sato et al., 2008;

Stengl and Funk, 2013; Wicher et al., 2008; Wicher and Miazzi, 2021). Thus, while this study did not

aim to unravel the overall generation of spontaneous activity in ORNs, here, we examined whether

the leak channel Orco in hawkmoth ORNs is the core pacemaker channel responsible for

membrane potential oscillations that drive spontaneous circadian activity (Stengl, 2010; Stengl

and Schneider, 2024).

Differential location and function of Orco homo-or heteromers in the distinct membrane

compartments of ORNs that face different ionic environments appear likely (Stengl and Funk, 2013;

Stengl and Schröder, 2021). We propose that homomeric *Manduca* Orco pacemaker channels, as

core element of a PTFL membrane clock, are restricted to the soma/inner dendrite compartment of

ORNs (Stengl and Schneider, 2024; Stengl and Schröder, 2021). In contrast, we expect *Manduca*

Orco heteromers to serve as obligatory chaperons to maintain ORs in the ciliary membranes, as

proven in *Drosophila* (Benton et al., 2006). Since the ciliary compartment faces very high

concentrations of K^+ in the sensillum lymph, matching intracellular K^+ levels (Kaissling and

Thorson, 1980; Steinbrecht, 1992), we propose that Orco in ciliary membranes remains closed via

unknown mechanisms, while Ca^{2+} channels of the TRP ion channel family serve the primary events

of pheromone transduction (Gawalek and Stengl, 2018; Stengl, 1994, 1993). Otherwise, Orco

channel opening would clamp the ciliary membranes at 0 mV, strongly reducing the driving force of

pheromone-dependent receptor currents, not allowing for the ORN's extreme pheromone

sensitivity (Stengl and Funk, 2013).

In support of this hypothesis, in both *Drosophila* (Benton et al., 2006) and *Manduca* (Nolte et

al., 2016), Orco antisera labeled Orco in two of the three sealed off ORN membrane

compartments: in the membranes of soma/ inner dendrite compartment, as well as in the

membrane of outer cilia, but not in the axons of ORNs. In contrast, antisera against an OR in

Drosophila detected ORs only in ciliary membranes but not in the other compartments of ORNs.

Thus, the colocalization of OR-Orco heteromers only in cilia confirm that Orco is an obligatory

coreceptor and chaperone of ORs localizing and maintaining the OR-Orco complexes in the ciliary

membrane (Benton et al., 2006). Since antisera against *Manduca* ORs are not available, we do not

known whether the pheromone-sensitive OR-Orco heteromers are solely located to ciliary

membranes also in the hawkmoth while Orco homomers are found in the soma/inner dendrite

compartment (Fandino et al., 2019). However, we know that Orco exists as homomeric complexes

in wild-type hawkmoth ORNs because Orco expression starts during development well before ORs

are present and because Orco agonist VUAA1 activates Orco in primary cell cultures of not yet

pheromone-sensitive maturing ORNs (Nolte et al., 2016; Schweitzer et al., 1976).

In summary, we propose Orco to be a 2nd messenger cascade-dependent core element of a

PTFL membrane clock that is located in the soma/inner dendrite compartment of ORNs, which

controls the ORN's membrane potential and is coupled to - but not dominated by - the TTFL clock

in the hawkmoth ORNs (Schuckel et al., 2007; Stengl and Schneider, 2024).

6.2   Orco is suggested to be a second messenger-dependent control hub of a PTFL

membrane clock that regulates circadian membrane potential oscillations in

hawkmoth ORNs

Different 2nd messenger cascades regulate *Drosophila* Orco via Ca^{2+}/calmodulin and via

phosphorylation/de-phosphorylation at multiple sites (Cao et al., 2016; Getahun et al., 2016; Guo

and Smith, 2017; Mukunda et al., 2016, 2014; Sargsyan et al., 2011). The Ca^{2+}/calmodulin binding

site controls the localization of the OR-Orco complex in the ciliary membrane of ORNs, provides for

negative feedback control of Orco conductance, and depends on previous odor stimulation (Bahk

and Jones, 2016; Cao et al., 2016; Guo and Smith, 2022; Jain et al., 2021). After phosphorylation of

Orco by protein kinase C (PKC), Orco directly or indirectly becomes cyclic nucleotide-dependent,

with cGMP and cAMP increasing its open-time probability (Getahun et al., 2013; Wicher et al.,

2008). While we do not know the second messenger-dependent modulation of hawkmoth Orco,

due to Orco's highly conserved structure we assume similar Ca^{2+}/calmodulin-, PKC-, and cyclic

nucleotide-dependent control of Orco's localization and open-time probability.

Since blocking Orco in *M. sexta* decreased the spontaneous activity and deleted its circadian

but not ultradian modulation, Orco is apparently the main target for modulation of circadian

activity in hawkmoth ORNs. The ultradian rhythms in spontaneous activity are presumably

controlled by additional pacemaker channels. Since Orco transcript abundance was constant

throughout the day, Orco is not controlled directly by the circadian TTFL clock. Rather, a self-

assembling PTFL membrane clock controls open-time probability and/or membrane localization of

Orco via posttranslational modifications in a circadian manner. Because circadian rhythms in

cAMP levels and phase-leading circadian rhythms in IP_3 were found in hawkmoth antennae

(Schendzielorz et al., 2015), it is likely that Orco's phosphorylation state regulates its cAMP

dependence in a daily rhythm. Indeed, our results showed that the increase of spontaneous activity

caused by 8-Br-cAMP application is canceled with co-application of the Orco blocker OLC15. Thus,

we assume that a PTFL membrane clock determines the circadian oscillations in the pheromone

sensitivity and temporal resolution of ORNs via Ca^{2+}, PKC, and cAMP-dependent membrane

potential control (Dolzer et al., 2021, 2008; Flecke et al., 2010, 2006; Redkozubov, 2000;

Schendzielorz et al., 2015, 2012; Ziegelberger et al., 1990).

Our hypothesis that a circadian PTFL membrane clock in hawkmoth ORNs controls circadian

rhythms in spontaneous spiking activity predicts that this PTFL clock comprises a membrane

signalosome with the pacemaker channel Orco as central control hub (Stengl and Schneider,

2024). In the soma/inner dendrite compartment of the ORNs the protein complexes of this PTFL

signalosome would generate the specific ultradian and circadian frequencies of endogenous

potential oscillations, linked to oscillations of Ca^{2+} and cAMP that determine physiological

setpoints of the ORN during circadian sleep-wake cycles. Thus, the PTFL membrane clock that

generates endogenous membrane potential and Ca^{2+} level oscillations minimally consists of

depolarizing, Ca^{2+} permeable Orco, voltage-gated Ca^{2+}, and hyperpolarizing K^+ channels. In the

same PTFL soma clock, membrane-anchored adenylyl cyclases (ACs) and phosphodiesterases

(PDE), that are antagonistically controlled by intracellular Ca^{2+} levels, are proposed to generate

circadian cAMP oscillations (Halls and Cooper, 2011). Kinases, such as PKC and PKA, could couple

potential oscillations with cAMP oscillations that target Orco, phosphatases, and PDEs via

negative feedback. Very likely, this membrane PTFL clock is interlinked with metabolic clocks such

as PTFL clocks that generate endogenous rhythms of redox elements or ATP availability (Edgar et

al., 2012), as evident in, e.g., human red blood cells, which lack a nucleus (O'Neill and Reddy,

2011).

Furthermore, this PTFL-clock could contain octopamine receptors that couple to

endogenous oscillations of intracellular Ca^{2+} and cAMP levels through activation of $G_{\alpha o}$ and $G_{\alpha s}$, as

found in hawkmoths (Dacks et al., 2006; Flecke et al., 2010). Since octopamine levels in the

hemolymph exhibit circadian oscillations and octopamine sets a physiological state of heightened

alertness in insects, the circadian PTFL clock in ORNs would synchronize with endogenous

physiological rhythms via octopamine receptor activation (Flecke and Stengl, 2009; Schendzielorz

et al., 2015). This also constitutes a possible pathway to synchronize the ORN PTFL clock with

circadian TTFL clocks in both ORNs and the brain. In horseshoe crab, the master clock sends

multiscale patterned spike trains through retinal efferents at night, which carries information about

the circadian phase and enhances visual sensitivity based on octopamine release and cAMP

signaling (Battelle, 2013; Liu and Passaglia, 2011). In *M. sexta*, two biogenic amine-immunoreactive

neurons project into the sensory cell layer of the antenna, being suited to phase-shift endogenous

cAMP oscillations via stress signals such as octopamine (Flecke and Stengl, 2009; Schendzielorz et

al., 2015). Comparably, the circadian pacemaker center controls endogenous rhythms in the

electrical network activity in rodent brains via endogenous cAMP oscillations that are phase-

controlled through G-protein coupled receptor signaling (Ono et al., 2023; Prosser et al., 1989;

Prosser and Gillette, 1991). Future studies will explore the links between PTFL and TTFL clocks in

ORNs.

6.3   Circadian and ultradian rhythms in spontaneous ORN activity resemble the

physiologically relevant timescales for active sensing

Moths are more sensitive to pheromone stimulation during their activity phase than at rest,

apparently due to antennal circadian clocks (Flecke et al., 2010; Flecke and Stengl, 2009; Merlin et

al., 2007, 2006; Schendzielorz et al., 2015; Schuckel et al., 2007; Silvegren et al., 2005). The greater

the distance from the female, the lower the frequency of pheromone filaments in the air. The

increased pheromone sensitivity during the activity phase enhances the male's chance to detect

and locate a calling female. Pheromone concentration is encoded in the frequency of the first,

phasic burst of spikes of the characteristic triphasic spiking sequence of ORNs in response to brief

pheromone pulses, which is independent of Orco (Dolzer et al., 2003; Nolte et al., 2013; Schneider

et al., 2025). Consistent with these results, we found here that blocking Orco does not affect the

upper frequency band of spontaneous activity that correlates with the frequency ranges of the

pheromone-encoding primary fast pheromone response(Dolzer et al., 2003; Nolte et al., 2016,

2013). However, through the cAMP-dependent increase in Orco conductance during the activity

phase, the ORN membrane potential is closer to spike threshold and fewer pheromone molecules

are necessary to produce a receptor current that brings the membrane voltage over that threshold.

Thus, larger Orco conductance increases the likelihood of an ORN response to pheromone

stimulation without changing the typical triphasic spiking pattern of the response.

In the odor transduction of *Drosophila*, OR-Orco activation drives the initial receptor current

response (Sato et al., 2008; Wicher et al., 2008). In contrast, Orco is not part of an OR-Orco odor-

gated receptor ion channel complex in hawkmoths but controls the spontaneous ORN activity and

the late, long-lasting pheromone response (LLPR). The LLPR begins several hundred milliseconds

after the initiation of the pheromone-dependent transduction current and correlates with the time

course of pheromone-induced rises in cGMP levels, possibly playing a role in odor memory or long-

term gain control (Boekhoff et al., 1993; Nolte et al., 2016, 2013; Stengl, 2010; Stengl et al., 2001;

Wicher and Miazzi, 2021; Ziegelberger et al., 1990). Thus, in hawkmoths, Orco is not part of the

primary transduction channel but rather controls membrane potential oscillations and thereby the

temporal resolution and general pheromone sensitivity of ORNs. In addition, the increased

spontaneous burst frequency during the hawkmoth's activity phase improves the resolution of the

anticipated pheromone pulses during its mating flight in search for females.

In the mammalian suprachiasmatic nucleus (SCN), the clock that controls the sleep-wake

cycle, cAMP levels oscillate at an circadian frequency, and cAMP and cGMP induce ZT-dependent

opposing phase shifts in SCN spiking activity (Prosser et al., 1989; Prosser and Gillette, 1991). In

antennae of both *M. sexta* (Schendzielorz et al., 2015) and the cockroach *Rhyparobia maderae*

(Schendzielorz et al., 2012), cAMP levels express daily and circadian oscillations and peak during

the activity phase. In contrast, cGMP levels do not oscillate under constant conditions. They are in

antiphase to the daily oscillations in cAMP levels in LD and are elevated by strong or long

pheromone exposure, which adapts the sensitivity and temporal resolution of ORNs (Boekhoff et

al., 1993; Dolzer et al., 2021; Flecke et al., 2006; Schendzielorz et al., 2015; Ziegelberger et al.,

1990). This implies that cAMP and cGMP may have ZT-dependent effects on Orco conductivity,

thereby affecting pheromone sensitivity via control of the membrane potential and intracellular

Ca^{2+} concentrations.

The upregulation of neural activity during wakefulness/alertness seems to be ubiquitous

among different species. For example, the many synchronously recorded neurons in human EEG

show fast oscillations in the alpha, beta, and gamma frequency ranges during alertness and slow

oscillations in the delta and theta ranges during rest and sleep. Similarly, in the lower frequency

band, the spontaneous spiking activity of hawkmoth ORNs changes from below delta during rest to

theta/alpha ranges during their activity phase. The upper frequency band in the gamma range,

which did not depend on Orco regulation, comprises the spike frequency within a burst,

reminiscent of the first phasic pheromone response that encodes pheromone concentration

changes (Dolzer et al., 2003). The increased spontaneous spiking activity in the theta/alpha range

during the activity phase of the hawkmoth matches the maximal temporal resolution of pheromone

pulses of up to 10 Hz (Marion-Poll and Tobin, 1992), which supports a role for Orco as part of a

membrane-bound PTFL clockwork that tunes the ORNs for active sensing. Even through

oscillations in the field potential of electroantennogram responses to odor stimulation can reach

100 Hz in different insects (Szyszka et al., 2014), these frequencies most likely do not encode the

temporal resolution of physiological odor responses. Apparently, they reflect the astounding

adaptive ability of antennal ORNs to synchronize their endogenous membrane potential

oscillations across the antenna to a broad range of stimulation frequencies. Pheromone pulse

resolution of about 100 Hz was never reported in single olfactory sensilla recordings, and in

behavioral experiments male moths stopped their upwind mate search if pheromone pulses

exceeded 30 Hz, apparently due to adaptation (Baker et al., 1985; Bau et al., 2005, 2002; Tripathy et

al., 2010).

Circadian oscillations in the ORN membrane potential would be ideally suited to tune

sensitivity and temporal resolution of the pheromone-sensitive ORNs to the hawkmoth's rest-

activity cycle, synchronizing male and female mating behavior. Here, we have demonstrated how

modulation of a leaky pacemaker channel can achieve these circadian oscillations. Whether cAMP

levels are under control of the TTFL clock or oscillate based on Ca^{2+} concentration oscillations, and

would therefore be part of a stand-alone PTFL clock, remains to be studied.

6.4 Stochastic ion channel dynamics and circadian modulation of Orco conductance explain modeled ORN firing patterns

Our modeling results provide mechanistic insight into the irregular spontaneous activity of pheromone-sensitive ORNs in *M. sexta*, highlighting how stochastic ion channel dynamics and circadian modulation of ion conductance together shape spiking patterns. By employing a Langevin approximation of the stochastic HH formalism, we demonstrate that the intrinsic noise arising from the random gating of ion channels is sufficient to generate both isolated spikes and bursts of spikes, as observed experimentally. Our model effectively recapitulates the spike distributions found *in vivo* and accounts for the role of channel noise as a critical determinant of ORN firing patterns.

A key contribution of our model is the incorporation of the Orco ion channel as a pacemaker component, with a conductance that varies linearly with cAMP concentration. We show that circadian modulation of Orco conductance alone is sufficient to produce the observed time-of-day dependent changes in spike distribution. This finding suggests that the rhythmic expression of cAMP levels across the circadian cycle may act as a primary driver of diurnal changes in ORN excitability, without requiring additional external inputs or modulation of other ion channels.

Importantly, our results reveal that the same set of ion channels can support the emergence of both single action potentials and complex bursting behaviors purely through the stochastic variability of their gating kinetics. This unifying explanation reduces the need for invoking separate biophysical mechanisms or circuit-level inputs to explain these distinct firing modes.

For simplicity, the circadian variation in cAMP concentration was modeled using a sinusoidal function. While this approximation is sufficient for capturing the qualitative features of the rhythmic modulation, future work could enhance the biological realism of the model by coupling Orco conductance dynamics to a molecular circadian oscillator, such as a Goodwin-type TTFL. Incorporating this layer would allow for a more detailed exploration of how molecular clock components regulate cAMP synthesis and degradation. Additionally, further extensions could include downstream regulatory mechanisms such as phosphorylation and dephosphorylation events mediated by kinases and phosphatases under circadian control, potentially contributing to fine-tuned modulation of ion channel function over the daily cycle.

Together, these findings demonstrate the value of a biophysically grounded, noise-inclusive modeling framework for understanding complex temporal patterns in sensory neuron activity and

809   open the door for future integrative models linking molecular circadian clocks to cellular

810   excitability.

7 References

Bahk S, Jones WD. 2016. Insect odorant receptor trafficking requires calmodulin. BMC Biology **14**:83. DOI: https://doi.org/10.1186/s12915-016-0306-x

Baker TC, Vogt RG. 1988. Measured Behavioural Latency in Response to Sex-Pheromone Loss in the Large Silk Moth *Antheraea Polyphemus*. Journal of Experimental Biology **137**:29–38. DOI: https://doi.org/10.1242/jeb.137.1.29

Baker TC, Willis MA, Haynes KF, Phelan PL. 1985. A pulsed cloud of sex pheromone elicits upwind flight in male moths. Physiological Entomology **10**:257–265. DOI: https://doi.org/10.1111/j.1365-3032.1985.tb00045.x

Battelle B-A. 2013. What the Clock Tells the Eye: Lessons from an Ancient Arthropod. Integrative and Comparative Biology **53**:144–153. DOI: https://doi.org/10.1093/icb/ict020, PMID: 23639718

Bau J, Justus KA, Cardé RT. 2002. Antennal resolution of pulsed pheromone plumes in three moth species. Journal of Insect Physiology **48**:433–442. DOI: https://doi.org/10.1016/S0022-1910(02)00062-8

Bau J, Justus KA, Loudon C, Cardé RT. 2005. Electroantennographic Resolution of Pulsed Pheromone Plumes in Two Species of Moths with Bipectinate Antennae. Chemical Senses **30**:771–780. DOI: https://doi.org/10.1093/chemse/bji069

Bell RA, Joachim FG. 1976. Techniques for Rearing Laboratory Colonies of Tobacco Hornworms and Pink Bollworms1. Annals of the Entomological Society of America **69**:365–373. DOI: https://doi.org/10.1093/aesa/69.2.365

Benton R, Sachse S, Michnick SW, Vosshall LB. 2006. Atypical Membrane Topology and Heteromeric Function of Drosophila Odorant Receptors In Vivo. PLOS Biology **4**:e20. DOI: https://doi.org/10.1371/journal.pbio.0040020

Boekhoff I, Seifert E, Göggerle S, Lindemann M, Krüger B-W, Breer H. 1993. Pheromone-induced second-messenger signaling in insect antennae. Insect Biochemistry and Molecular Biology **23**:757–762. DOI: https://doi.org/10.1016/0965-1748(93)90063-X

Bose A, Golowasch J, Guan Y, Nadim F. 2014. The role of linear and voltage-dependent ionic currents in the generation of slow wave oscillations. Journal of Computational Neuroscience **37**:229–242. DOI: https://doi.org/10.1007/s10827-014-0498-4

Butterwick JA, del Mármol J, Kim KH, Kahlson MA, Rogow JA, Walz T, Ruta V. 2018. Cryo-EM structure of the insect olfactory receptor Orco. Nature **560**:447–452. DOI: https://doi.org/10.1038/s41586-018-0420-8

Cao L-H, Jing B-Y, Yang D, Zeng X, Shen Y, Tu Y, Luo D-G. 2016. Distinct signaling of *Drosophila* chemoreceptors in olfactory sensory neurons. Proceedings of the National Academy of Sciences **113**:E902–E911. DOI: https://doi.org/10.1073/pnas.1518329113

Chen S, Luetje CW. 2012. Identification of New Agonists and Antagonists of the Insect Odorant Receptor Co-Receptor Subunit. PLOS ONE **7**:e36784. DOI: https://doi.org/10.1371/journal.pone.0036784

Cochet-Bissuel M, Lory P, Monteil A. 2014. The sodium leak channel, NALCN, in health and disease. Frontiers in Cellular Neuroscience **8**:132. DOI: https://doi.org/10.3389/fncel.2014.00132, PMID: 24904279

Dacks AM, Dacks JB, Christensen TA, Nighorn AJ. 2006. The cloning of one putative octopamine receptor and two putative serotonin receptors from the tobacco hawkmoth, *Manduca sexta*. Insect Biochemistry and Molecular Biology **36**:741–747. DOI: https://doi.org/10.1016/j.ibmb.2006.07.002

Das A, Holmes TC, Sheeba V. 2016. dTRPA1 in Non-circadian Neurons Modulates Temperature-
dependent Rhythmic Activity in *Drosophila melanogaster*. Journal of Biological Rhythms
31:272–288. DOI: https://doi.org/10.1177/0748730415627037
Dolzer J, Fischer K, Stengl M. 2003. Adaptation in pheromone-sensitive trichoid sensilla of the
hawkmoth *Manduca sexta*. Journal of Experimental Biology **206**:1575–1588. DOI:
https://doi.org/10.1242/jeb.00302
Dolzer J, Krannich S, Fischer K, Stengl M. 2001. Oscillations of the Transepithelial Potential of Moth
Olfactory Sensilla Are Influenced by Octopamine and Serotonin. Journal of Experimental
Biology **204**:2781–2794. DOI: https://doi.org/10.1242/jeb.204.16.2781
Dolzer J, Krannich S, Stengl M. 2008. Pharmacological Investigation of Protein Kinase C- and cGMP-
Dependent Ion Channels in Cultured Olfactory Receptor Neurons of the Hawkmoth
Manduca sexta. Chemical Senses **33**:803–813. DOI:
https://doi.org/10.1093/chemse/bjn043
Dolzer J, Schröder K, Stengl M. 2021. Cyclic nucleotide-dependent ionic currents in olfactory
receptor neurons of the hawkmoth *Manduca sexta* suggest pull–push sensitivity
modulation. European Journal of Neuroscience **54**:4804–4826. DOI:
https://doi.org/10.1111/ejn.15346
Dubowy C, Sehgal A. 2017. Circadian Rhythms and Sleep in *Drosophila melanogaster*. Genetics
205:1373–1397. DOI: https://doi.org/10.1534/genetics.115.185157
Edgar RS, Green EW, Zhao Y, van Ooijen G, Olmedo M, Qin X, Xu Y, Pan M, Valekunja UK, Feeney
KA, Maywood ES, Hastings MH, Baliga NS, Merrow M, Millar AJ, Johnson CH, Kyriacou CP,
O'Neill JS, Reddy AB. 2012. Peroxiredoxins are conserved markers of circadian rhythms.
Nature **485**:459–464. DOI: https://doi.org/10.1038/nature11088, PMID: 22622569
Fandino RA, Haverkamp A, Bisch-Knaden S, Zhang J, Bucks S, Nguyen TAT, Schröder K,
Werckenthin A, Rybak J, Stengl M, Knaden M, Hansson BS, Große-Wilde E. 2019.
Mutagenesis of odorant coreceptor Orco fully disrupts foraging but not oviposition
behaviors in the hawkmoth *Manduca sexta*. Proceedings of the National Academy of
Sciences **116**:15677–15685. DOI: https://doi.org/10.1073/pnas.1902089116
Flecke C, Dolzer J, Krannich S, Stengl M. 2006. Perfusion with cGMP analogue adapts the action
potential response of pheromone-sensitive sensilla trichoidea of the hawkmoth *Manduca
sexta* in a daytime-dependent manner. Journal of Experimental Biology **209**:3898–3912.
DOI: https://doi.org/10.1242/jeb.02432
Flecke C, Nolte A, Stengl M. 2010. Perfusion with cAMP analogue affects pheromone-sensitive
trichoid sensilla of the hawkmoth *Manduca sexta* in a time-dependent manner. Journal of
Experimental Biology **213**:842–852. DOI: https://doi.org/10.1242/jeb.032839
Flecke C, Stengl M. 2009. Octopamine and tyramine modulate pheromone-sensitive olfactory
sensilla of the hawkmoth *Manduca sexta* in a time-dependent manner. Journal of
Comparative Physiology A **195**:529–545. DOI: https://doi.org/10.1007/s00359-009-0429-4
Forlino M, Vijayan A, Schneider AC, Stengl M, Garcia ME. 2025. ORN-Orco-Model.
https://github.com/a-c-schneider/VijayanForlinoEtAl2025_Model
Fox RF, Lu Y. 1994. Emergent collective behavior in large numbers of globally coupled
independently stochastic ion channels. Physical Review E **49**:3421–3431. DOI:
https://doi.org/10.1103/PhysRevE.49.3421
Gawalek P, Stengl M. 2018. The Diacylglycerol Analogs OAG and DOG Differentially Affect Primary
Events of Pheromone Transduction in the Hawkmoth *Manduca sexta* in a Zeitgebertime-
Dependent Manner Apparently Targeting TRP Channels. Frontiers in Cellular Neuroscience
12:218. DOI: https://doi.org/10.3389/fncel.2018.00218

Géron A, Werner ,Johannes, Wattiez ,Ruddy, and Matallana-Surget S. 2024. Towards the discovery
of novel molecular clocks in Prokaryotes. Critical Reviews in Microbiology **50**:491–503. DOI:
https://doi.org/10.1080/1040841X.2023.2220789, PMID: 37330701
Getahun MN, Olsson SB, Lavista-Llanos S, Hansson BS, Wicher D. 2013. Insect Odorant Response
Sensitivity Is Tuned by Metabotropically Autoregulated Olfactory Receptors. PLOS ONE
8:e58889. DOI: https://doi.org/10.1371/journal.pone.0058889
Getahun MN, Thoma M, Lavista-Llanos S, Keesey I, Fandino RA, Knaden M, Wicher D, Olsson SB,
Hansson BS. 2016. Intracellular regulation of the insect chemoreceptor complex impacts
odour localization in flying insects. Journal of Experimental Biology **219**:3428–3438. DOI:
https://doi.org/10.1242/jeb.143396
Ghosh S, Suray C, Bozzolan F, Palazzo A, Monsempès C, Lecouvreur F, Chatterjee A. 2024.
Pheromone-mediated command from the female to male clock induces and synchronizes
circadian rhythms of the moth *Spodoptera littoralis*. Current biology **34**:1414-1425.e5. DOI:
https://doi.org/10.1016/j.cub.2024.02.042, PMID: 38479388
Goldstein SAN, Bockenhauer D, O'Kelly I, Zilberberg N. 2001. Potassium leak channels and the
KCNK family of two-p-domain subunits. Nature Reviews Neuroscience **2**:175–184. DOI:
https://doi.org/10.1038/35058574
Golowasch J, Bose A, Guan Y, Salloum D, Roeser A, Nadim F. 2017. A balance of outward and
linear inward ionic currents is required for generation of slow-wave oscillations. Journal of
Neurophysiology **118**:1092–1104. DOI: https://doi.org/10.1152/jn.00240.2017
Guo H, Smith DP. 2022. Time-Dependent Odorant Sensitivity Modulation in Insects. Insects **13**:354.
DOI: https://doi.org/10.3390/insects13040354
Guo H, Smith DP. 2017. Odorant Receptor Desensitization in Insects. Journal of Experimental
Neuroscience **11**:1179069517748600. DOI: https://doi.org/10.1177/1179069517748600
Hall JC, Rosbash M. 1993. Oscillating molecules and how they move circadian clocks across
evolutionary boundaries. Proceedings of the National Academy of Sciences of the United
States of America **90**:5382–5383. PMID: 8516281
Halls ML, Cooper DMF. 2011. Regulation by Ca2+-Signaling Pathways of Adenylyl Cyclases. Cold
Spring Harbor Perspectives in Biology **3**:a004143. DOI:
https://doi.org/10.1101/cshperspect.a004143, PMID: 21123395
Hardin PE. 2011. Molecular Genetic Analysis of Circadian Timekeeping in *Drosophila*. In: Brody S
(Ed). *Advances in Genetics*, The Genetics of Circadian Rhythms. Academic Press. p. 141–
173. DOI: https://doi.org/10.1016/B978-0-12-387690-4.00005-2
Hastings MH, Smyllie NJ, Patton AP. 2020. Molecular-genetic Manipulation of the Suprachiasmatic
Nucleus Circadian Clock. Journal of Molecular Biology **432**:3639–3660. DOI:
https://doi.org/10.1016/j.jmb.2020.01.019, PMID: 31996314
Hille B. 2001. Ion Channels of Excitable Membranes, Third Edition, Third Edition. Oxford University
Press.
Itagaki H, Conner WE. 1988. Calling Behavior of *Manduca sexta* (L.) (Lepidoptera: Sphingidae) with
Notes on the Morphology of the Female Sex Pheromone Gland. Annals of the Entomological
Society of America **81**:798–807. DOI: https://doi.org/10.1093/aesa/81.5.798
Jain K, Lavista-Llanos S, Grabe V, Hansson BS, Wicher D. 2021. Calmodulin regulates the olfactory
performance in Drosophila melanogaster. Scientific Reports **11**:3747. DOI:
https://doi.org/10.1038/s41598-021-83296-9
Jones PL, Pask GM, Rinker DC, Zwiebel LJ. 2011. Functional agonism of insect odorant receptor ion
channels. Proceedings of the National Academy of Sciences **108**:8821–8825. DOI:
https://doi.org/10.1073/pnas.1102425108

Kaissling KE, Hildebrand JG, Tumlinson JH. 1989. Pheromone receptor cells in the male moth
Manduca sexta. Archives of Insect Biochemistry and Physiology **10**:273–279. DOI:
https://doi.org/10.1002/arch.940100403

Kaissling KE, Thorson J. 1980. Insect Olfactory Sensilla: Structural, Chemical and Electrical
Aspects of the Functional Organisation. In: Sattelle DB, Hall LM, Hildebrand JG (Eds).
Receptors for Neurotransmitters, Hormones and Pheromones in Insects. Elsevier/North-
Holland Biomedical Press. p. 261–282.

Kaissling KE, Zack Strausfeld C, Rumbo ER. 1987. Adaptation processes in insect olfactory
receptors. Mechanisms and behavioral significance. Annals of the New York Academy of
Sciences **510**:104–112. DOI: https://doi.org/10.1111/j.1749-6632.1987.tb43475.x, PMID:
3324874

Kalinová B, Hoskovec M, Liblikas I, Unelius CR, Hansson BS. 2001. Detection of Sex Pheromone
Components in *Manduca sexta* (L.). Chemical Senses **26**:1175–1186. DOI:
https://doi.org/10.1093/chemse/26.9.1175

Kodirov SA. 2022. Probability that there is a mammalian counterpart of cardiac clock in insects.
Archives of Insect Biochemistry and Physiology **110**:e21867. DOI:
https://doi.org/10.1002/arch.21867

Krishnan B, Dryer SE, Hardin PE. 1999. Circadian rhythms in olfactory responses of *Drosophila*
melanogaster. Nature **400**:375–378. DOI: https://doi.org/10.1038/22566

Lamont EW, Amir S. 2010. Circadian and Ultradian Clocks/Rhythms. In: Koob GF, Moal ML,
Thompson RF (Eds). *Encyclopedia of Behavioral Neuroscience*. Academic Press. p. 257–
261. DOI: https://doi.org/10.1016/B978-0-08-045396-5.00223-2

Lee C-H, MacKinnon R. 2017. Structures of the Human HCN1 Hyperpolarization-Activated
Channel. Cell **168**:111-120.e11. DOI: https://doi.org/10.1016/j.cell.2016.12.023

Lee JK, Strausfeld NJ. 1990. Structure, distribution and number of surface sensilla and their
receptor cells on the olfactory appendage of the male mothManduca sexta. Journal of
Neurocytology **19**:519–538. DOI: https://doi.org/10.1007/BF01257241

Liu JS, Passaglia CL. 2011. Spike firing pattern of output neurons of the Limulus circadian clock.
Journal of Biological Rhythms **26**:335–344. DOI:
https://doi.org/10.1177/0748730411409712, PMID: 21775292

Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time
quantitative PCR and the $2^{-\Delta\Delta C(T)}$ Method. Methods (San Diego, Calif.) **25**:402–408. DOI:
https://doi.org/10.1006/meth.2001.1262, PMID: 11846609

Lüthi A, McCormick DA. 1998. H-Current: Properties of a Neuronal and Network Pacemaker.
Neuron **21**:9–12. DOI: https://doi.org/10.1016/S0896-6273(00)80509-7

Mandala VS, Fu Z, MacKinnon R. 2025. Molecular contacts in self-assembling clusters of
membrane proteins. Proceedings of the National Academy of Sciences **122**:e2507112122.
DOI: https://doi.org/10.1073/pnas.2507112122

Marion-Poll F, Tobin TR. 1992. Temporal coding of pheromone pulses and trains in *Manduca sexta*.
Journal of Comparative Physiology A **171**:505–512. DOI:
https://doi.org/10.1007/BF00194583

Martín F, Charro M, Alcorta E. 2001. Mutations affecting the cAMP transduction pathway modify
olfaction in *Drosophila*. Journal of Comparative Physiology A **187**:359–370. DOI:
https://doi.org/10.1007/s003590100208

Merlin C, François M-C, Queguiner I, Maïbèche-Coisné M, Jacquin-Joly E. 2006. Evidence for a
putative antennal clock in *Mamestra brassicae*: Molecular cloning and characterization of
two clock genes –*period* and *cryptochrome*– in antennae. Insect Molecular Biology **15**:137–
145. DOI: https://doi.org/10.1111/j.1365-2583.2006.00617.x

Merlin C, Gegear RJ, Reppert SM. 2009. Antennal Circadian Clocks Coordinate Sun Compass
Orientation in Migratory Monarch Butterflies. Science **325**:1700–1704. DOI:
https://doi.org/10.1126/science.1176221

Merlin C, Lucas P, Rochat D, François M-C, Maïbèche-Coisne M, Jacquin-Joly E. 2007. An Antennal
Circadian Clock and Circadian Rhythms in Peripheral Pheromone Reception in the Moth
Spodoptera littoralis. Journal of Biological Rhythms **22**:502–514. DOI:
https://doi.org/10.1177/0748730407307737

Millar AJ. 2016. The Intracellular Dynamics of Circadian Clocks Reach for the Light of Ecology and
Evolution. Annual Review of Plant Biology **67**:595–618. DOI:
https://doi.org/10.1146/annurev-arplant-043014-115619

Mönke G, Sorgenfrei FA, Schmal C, Granada AE. 2020. Optimal time frequency analysis for
biological data - pyBOAT. DOI: https://doi.org/10.1101/2020.04.29.067744

Mukunda L, Miazzi F, Kaltofen S, Hansson BS, Wicher D. 2014. Calmodulin modulates insect
odorant receptor function. Cell Calcium **55**:191–199. DOI:
https://doi.org/10.1016/j.ceca.2014.02.013

Mukunda L, Miazzi F, Sargsyan V, Hansson BS, Wicher D. 2016. Calmodulin Affects Sensitization of
Drosophila melanogaster Odorant Receptors. Frontiers in Cellular Neuroscience **10**. DOI:
https://doi.org/10.3389/fncel.2016.00028

Nakagawa T, Vosshall LB. 2009. Controversy and consensus: noncanonical signaling mechanisms
in the insect olfactory system. Current Opinion in Neurobiology **19**:284–292. DOI:
https://doi.org/10.1016/j.conb.2009.07.015

Nishiitsutsuji-Uwo J, Pittendrigh CS. 1968. Central nervous system control of circadian rhythmicity
in the cockroach II. The pathway of light signals that entrain the rhythm. Zeitschrift für
vergleichende Physiologie **58**:1–13. DOI: https://doi.org/10.1007/BF00302433

Nolte A, Funk NW, Mukunda L, Gawalek P, Werckenthin A, Hansson BS, Wicher D, Stengl M. 2013.
In situ Tip-Recordings Found No Evidence for an Orco-Based Ionotropic Mechanism of
Pheromone-Transduction in *Manduca sexta*. PLOS ONE **8**:e62648. DOI:
https://doi.org/10.1371/journal.pone.0062648

Nolte A, Gawalek P, Koerte S, Wei H, Schumann R, Werckenthin A, Krieger J, Stengl M. 2016. No
Evidence for Ionotropic Pheromone Transduction in the Hawkmoth *Manduca sexta*. PLOS
ONE **11**:e0166060. DOI: https://doi.org/10.1371/journal.pone.0166060

O'Neill JS, Reddy AB. 2011. Circadian clocks in human red blood cells. Nature **469**:498–503. DOI:
https://doi.org/10.1038/nature09702

Ono D, Wang Huan, Hung CJ, Wang Hsin-tzu, Kon N, Yamanaka A, Li Y, Sugiyama T. 2023. Network-
driven intracellular cAMP coordinates circadian rhythm in the suprachiasmatic nucleus.
Science Advances **9**:eabq7032. DOI: https://doi.org/10.1126/sciadv.abq7032

Page TL, Koelling E. 2003. Circadian rhythm in olfactory response in the antennae controlled by the
optic lobe in the cockroach. Journal of Insect Physiology **49**:697–707. DOI:
https://doi.org/10.1016/s0022-1910(03)00071-4, PMID: 12837322

Pask GM, Bobkov YV, Corey EA, Ache BW, Zwiebel LJ. 2013. Blockade of Insect Odorant Receptor
Currents by Amiloride Derivatives. Chemical Senses **38**:221–229. DOI:
https://doi.org/10.1093/chemse/bjs100

Patel AJ, Honoré E. 2001. Properties and modulation of mammalian 2P domain K+ channels.
Trends in Neurosciences **24**:339–346. DOI: https://doi.org/10.1016/S0166-2236(00)01810-
5

Pézier A, Acquistapace A, Renou M, Rospars J-P, Lucas P. 2007. Ca^{2+} Stabilizes the Membrane
Potential of Moth Olfactory Receptor Neurons at Rest and Is Essential for Their Fast
Repolarization. Chemical Senses **32**:305–317. DOI: https://doi.org/10.1093/chemse/bjl059

Plautz JD, Kaneko M, Hall JC, Kay SA. 1997. Independent Photoreceptive Circadian Clocks
Throughout *Drosophila*. Science **278**:1632–1635. DOI:
https://doi.org/10.1126/science.278.5343.1632
Prosser RA, Gillette MU. 1991. Cyclic changes in cAMP concentration and phosphodiesterase
activity in a mammalian circadian clock studied *in vitro*. Brain Research **568**:185–192. DOI:
https://doi.org/10.1016/0006-8993(91)91396-I
Prosser RA, McArthur AJ, Gillette MU. 1989. cGMP induces phase shifts of a mammalian circadian
pacemaker at night, in antiphase to cAMP effects. Proceedings of the National Academy of
Sciences of the United States of America **86**:6812–6815. PMID: 2549549
Pu S, Thomas PJ. 2020. Fast and Accurate Langevin Simulations of Stochastic Hodgkin-Huxley
Dynamics. Neural Computation **32**:1775–1835. DOI:
https://doi.org/10.1162/neco_a_01312
Ratliff J, Franci A, Marder E, O'Leary T. 2021. Neuronal oscillator robustness to multiple global
perturbations. Biophysical Journal **120**:1454–1468. DOI:
https://doi.org/10.1016/j.bpj.2021.01.038
Redkozubov A. 2000. Guanosine 3',5'-Cyclic Monophosphate Reduces the Response of the Moth's
Olfactory Receptor Neuron to Pheromone. Chemical Senses **25**:381–385. DOI:
https://doi.org/10.1093/chemse/25.4.381
Reinhard N, Fukuda A, Manoli G, Derksen E, Saito A, Möller G, Sekiguchi M, Rieger D, Helfrich-
Förster C, Yoshii T, Zandawala M. 2024. Synaptic connectome of the *Drosophila* circadian
clock. Nature Communications **15**:10392. DOI: https://doi.org/10.1038/s41467-024-54694-
0
Riffell JA, Abrell L, Hildebrand JG. 2008. Physical processes and real-time chemical measurement
of the insect olfactory environment. Journal of Chemical Ecology **34**:837–853. DOI:
https://doi.org/10.1007/s10886-008-9490-7, PMID: 18548311
Robinson RB, Siegelbaum SA. 2003. Hyperpolarization-activated cation currents: from molecules
to physiological function. Annual Review of Physiology **65**:453–480. DOI:
https://doi.org/10.1146/annurev.physiol.65.092101.142734, PMID: 12471170
Sargsyan V, Getahun MN, Lavista Llanos S, Olsson SB, Hansson BS, Wicher D. 2011.
Phosphorylation via PKC Regulates the Function of the *Drosophila* Odorant Co-Receptor.
Frontiers in Cellular Neuroscience **5**. DOI: https://doi.org/10.3389/fncel.2011.00005
Sato K, Pellegrino M, Nakagawa Takao, Nakagawa Tatsuro, Vosshall LB, Touhara K. 2008. Insect
olfactory receptors are heteromeric ligand-gated ion channels. Nature **452**:1002–1006.
DOI: https://doi.org/10.1038/nature06850
Sauman I, Reppert SM. 1998. Brain Control of Embryonic Circadian Rhythms in the Silkmoth
Antheraea pernyi. Neuron **20**:741–748. DOI: https://doi.org/10.1016/S0896-6273(00)81012-
0
Schendzielorz J, Schendzielorz T, Arendt A, Stengl M. 2014. Bimodal Oscillations of Cyclic
Nucleotide Concentrations in the Circadian System of the Madeira Cockroach *Rhyparobia*
maderae. Journal of Biological Rhythms **29**:318–331. DOI:
https://doi.org/10.1177/0748730414546133
Schendzielorz T, Peters W, Boekhoff I, Stengl M. 2012. Time of Day Changes in Cyclic Nucleotides
Are Modified via Octopamine and Pheromone in Antennae of the Madeira Cockroach.
Journal of Biological Rhythms **27**:388–397. DOI:
https://doi.org/10.1177/0748730412456265
Schendzielorz T, Schirmer K, Stolte P, Stengl M. 2015. Octopamine Regulates Antennal Sensory
Neurons via Daytime-Dependent Changes in cAMP and IP_3 Levels in the Hawkmoth

Manduca sexta. PLOS ONE **10**:e0121230. DOI:
https://doi.org/10.1371/journal.pone.0121230
Schmidt DR, Galán RF, Thomas PJ. 2018. Stochastic shielding and edge importance for Markov
chains with timescale separation. PLOS Computational Biology **14**:e1006206. DOI:
https://doi.org/10.1371/journal.pcbi.1006206
Schmidt DR, Thomas PJ. 2014. Measuring Edge Importance: A Quantitative Analysis of the
Stochastic Shielding Approximation for Random Processes on Graphs. The Journal of
Mathematical Neuroscience **4**:6. DOI: https://doi.org/10.1186/2190-8567-4-6
Schneider AC, Schröder K, Chang Y, Nolte A, Gawalek P, Stengl M. 2025. Hawkmoth Pheromone
Transduction Involves G-Protein–Dependent Phospholipase Cβ Signaling. eNeuro
12:ENEURO.0376-24.2024. DOI: https://doi.org/10.1523/ENEURO.0376-24.2024, PMID:
39880675
Schuckel J, Siwicki KK, Stengl M. 2007. Putative circadian pacemaker cells in the antenna of the
hawkmoth *Manduca sexta*. Cell and Tissue Research **330**:271–278. DOI:
https://doi.org/10.1007/s00441-007-0471-x
Schweitzer ES, Sanes JR, Hildebrand JG. 1976. Ontogeny of electroantennogram responses in the
moth, *Manduca sexta*. Journal of Insect Physiology **22**:955–960. DOI:
https://doi.org/10.1016/0022-1910(76)90078-0
Sharma A, Rahman G, Gorelik J, Bhargava A. 2023. Voltage-Gated T-Type Calcium Channel
Modulation by Kinases and Phosphatases: The Old Ones, the New Ones, and the Missing
Ones. Cells **12**:461. DOI: https://doi.org/10.3390/cells12030461, PMID: 36766802
Silvegren G, Löfstedt C, Qi Rosén W. 2005. Circadian mating activity and effect of pheromone pre-
exposure on pheromone response rhythms in the moth *Spodoptera littoralis*. Journal of
Insect Physiology **51**:277–286. DOI: https://doi.org/10.1016/j.jinsphys.2004.11.013
Steinbrecht RA. 1992. Experimental morphology of insect olfaction: Tracer studies, x-ray
microanalysis, autoradiography, and immunocytochemistry with silkmoth antennae.
Microscopy Research and Technique **22**:336–350. DOI:
https://doi.org/10.1002/jemt.1070220404
Stengl M. 2010. Pheromone Transduction in Moths. Frontiers in Cellular Neuroscience **4**:133. DOI:
https://doi.org/10.3389/fncel.2010.00133
Stengl M. 1994. Inositol-trisphosphate-dependent calcium currents precede cation currents in
insect olfactory receptor neurons *in vitro*. Journal of Comparative Physiology A **174**:187–
194. DOI: https://doi.org/10.1007/BF00193785
Stengl M. 1993. Intracellular-Messenger-Mediated Cation Channels in Cultured Olfactory Receptor
Neurons. Journal of Experimental Biology **178**:125–147. DOI:
https://doi.org/10.1242/jeb.178.1.125
Stengl M, Funk NW. 2013. The role of the coreceptor Orco in insect olfactory transduction. Journal
of Comparative Physiology A **199**:897–909. DOI: https://doi.org/10.1007/s00359-013-0837-
3
Stengl M, Schneider AC. 2024. Contribution of membrane-associated oscillators to biological
timing at different timescales. Frontiers in Physiology **14**:1243455. DOI:
https://doi.org/10.3389/fphys.2023.1243455
Stengl M, Schröder K. 2021. 14 - Multiscale timing of pheromone transduction in hawkmoth
olfactory receptor neurons. In: Blomquist GJ, Vogt RG (Eds). *Insect Pheromone
Biochemistry and Molecular Biology (Second Edition)*. Academic Press. p. 435–468. DOI:
https://doi.org/10.1016/B978-0-12-819628-1.00014-6

Stengl M, Zintl R, De Vente J, Nighorn A. 2001. Localization of cGMP immunoreactivity and of
soluble guanylyl cyclase in antennal sensilla of the hawkmoth Manduca sexta. Cell and
Tissue Research **304**:409–421. DOI: https://doi.org/10.1007/s004410000336
Szyszka P, Gerkin RC, Galizia CG, Smith BH. 2014. High-speed odor transduction and pulse
tracking by insect olfactory receptor neurons. Proceedings of the National Academy of
Sciences **111**:16925–16930. DOI: https://doi.org/10.1073/pnas.1412051111
Talley EM, Solórzano G, Lei Q, Kim D, Bayliss DA. 2001. CNS Distribution of Members of the Two-
Pore-Domain (KCNK) Potassium Channel Family. Journal of Neuroscience **21**:7491–7505.
DOI: https://doi.org/10.1523/JNEUROSCI.21-19-07491.2001, PMID: 11567039
Tanoue S, Krishnan P, Krishnan B, Dryer SE, Hardin PE. 2004. Circadian Clocks in Antennal
Neurons Are Necessary and Sufficient for Olfaction Rhythms in *Drosophila*. Current Biology
14:638–649. DOI: https://doi.org/10.1016/j.cub.2004.04.009, PMID: 15084278
Thaben PF, Westermark PO. 2014. Detecting Rhythms in Time Series with RAIN. Journal of
Biological Rhythms **29**:391–400. DOI: https://doi.org/10.1177/0748730414553029
Thaiss CA, Zeevi D, Levy M, Zilberman-Schapira G, Suez J, Tengeler AC, Abramson L, Katz MN,
Korem T, Zmora N, Kuperman Y, Biton I, Gilad S, Harmelin A, Shapiro H, Halpern Z, Segal E,
Elinav E. 2014. Transkingdom Control of Microbiota Diurnal Oscillations Promotes
Metabolic Homeostasis. Cell **159**:514–529. DOI: https://doi.org/10.1016/j.cell.2014.09.048
Tripathy S, Staudacher EM, Peters O, Kalwar F, Hatfield M, Daly K. 2010. Odors pulsed at wing beat
frequencies are tracked by primary olfactory networks and enhance odor detection.
Frontiers in Cellular Neuroscience **4**. DOI: https://doi.org/10.3389/neuro.03.001.2010
Viertel R, Borisyuk A. 2019. A Computational model of the mammalian external tufted cell. Journal
of Theoretical Biology **462**:109–121. DOI: https://doi.org/10.1016/j.jtbi.2018.10.003
Wicher D. 2018. Tuning Insect Odorant Receptors. Frontiers in Cellular Neuroscience **12**:94. DOI:
https://doi.org/10.3389/fncel.2018.00094, PMID: 29674957
Wicher D, Miazzi F. 2021. Functional properties of insect olfactory receptors: ionotropic receptors
and odorant receptors. Cell and Tissue Research **383**:7–19. DOI:
https://doi.org/10.1007/s00441-020-03363-x
Wicher D, Schäfer R, Bauernfeind R, Stensmyr MC, Heller R, Heinemann SH, Hansson BS. 2008.
Drosophila odorant receptors are both ligand-gated and cyclic-nucleotide-activated cation
channels. Nature **452**:1007–1011. DOI: https://doi.org/10.1038/nature06861
Zhou X, Yuan C, Guo A. 2005. *Drosophila* Olfactory Response Rhythms Require Clock Genes but
Not Pigment Dispersing Factor or Lateral Neurons. Journal of Biological Rhythms **20**:237–
244. DOI: https://doi.org/10.1177/0748730405274451
Ziegelberger G, Berg M van den, Kaissling KE, Klumpp S, Schultz JE. 1990. Cyclic GMP levels and
guanylate cyclase activity in pheromone-sensitive antennae of the silkmoths *Antheraea*
polyphemus and *Bombyx mori*. Journal of Neuroscience **10**:1217–1225. DOI:
https://doi.org/10.1523/JNEUROSCI.10-04-01217.1990, PMID: 1970356
1178

1179 # 8 Figures and Figure legends

1180

Figure 1: Attributes of pheromone-sensitive ORN spiking patterns.

(**A**) Spikes were identified by their prominence in the raw ORN recording. The inter-spike interval
(ISI) was defined as the time from one spike to the next. A train with ≥2 spikes with ISI ≤ 50 ms was
considered a burst. The intra-burst interval was the ISIs within a burst, the inter-burst interval (IBI)
the ISIs between two bursts, omitting any individual spikes in between. The inter-event interval (IEI)
included the ISIs between bursts and individual spikes. Detaching and reattaching the recording
electrode to a long trichoid sensillum (**Bi**; paired t-test $t(9)$ = -2.184; p = 0.057; N = 10) or adding
DMSO to the SLR (**Bii**; Wilcoxon signed-rank test Z = 0.153; p = 0.922; N = 10) did not affect the
mean spiking frequency.

1190

1191

Figure 2: The conductance-based model of a simplified hawkmoth ORN.

1192

(A) Schematic of the model containing a voltage gated sodium channel (Na$^+$), a voltage gated
potassium channel (K$^+$), a low voltage activated calcium channel (LVA), a calcium and voltage
activated potassium channel (BK), a leak channel, and a cAMP-gated leaky non-specific cation
channel (Orco). The Orco channel is represented as a linear conductance whose value oscillates
with a sinusoidal shape as a function of the ZT. The circadian rhythm in the Orco conductance is
hypothesized to result from a cAMP-dependent increase in channel open-time probability (as
found in *Drosophila* Orco: (Getahun et al., 2013)) via a circadian oscillation of antennal cAMP
concentrations (Schendzielorz et al., 2015). This oscillation is driven by the circadian clock and
serves as an input; the internal mechanism of the clock is not modeled explicitly. Arrows indicate
the flow of ions through the channels. **(B)** Parametrized circadian oscillation in Orco conductivity,

introduced as an external input to our model, representing the gating of Orco by cAMP whose
concentration oscillates on a circadian timescale. The times of minimum and maximum cAMP
concentration are indicated by colored arrows. **(C)** Parametrized IV curve of the Orco channel
showing a linear behavior with the slope depending on the cAMP concentration. Line colors
correspond to the arrows in **B**. **(D)** Markov-chain representing all possible states and transitions for
each type of ion channel included in the model. The per capita transition rates (α and β) depend on
membrane potential and in the case of the BK channel also on the intracellular Ca^{2+}-concentration.
The "N" state vectors contain the population of channels in each state "n". Directed edges are
numbered in red. The only conducting state of each ion channel, representing the condition where
all its gates are open, is shown in green.

1213

1214

Figure 3: Spontaneous activity of pheromone-sensitive olfactory receptor neurons (ORNs) of male *M. sexta* shows Orco-dependent circadian modulation.
(A) Example tip recordings of the spontaneous activity of one long trichoid sensillum during a period of high (top, average frequency ~ 3 Hz) and low (bottom, average frequency ~ 0.2 Hz) activity. **(Bi)** Spontaneous spike frequency of one long-term ORN recording during 17:7 light-dark cycles (same experiment as in **A,** colored arrows indicate respective zeitgeber times (ZT)), indicated by the black-white bar at the top, increased during each activity phase and was low during the resting phases. Spike frequencies were averaged for each 1 h bin. **(Bii)** Mean spontaneous ORN spike frequency across all LD animals in 1 h bins. Data from the individual in **Bi** is highlighted in red. Recordings had different lengths, therefore the number of recordings used for the element-wise mean for each 1 h bin decreased with the time since the start of the recording, indicated in the top

panel. **(Biii)** Mean spontaneous ORN spike frequency across all LD animals in 1 h bins during the
first 48 hours revealed circadian activity. **(Ci)** The spontaneous spike frequency of one long-term
ORN recording in constant darkness (DD, the expected times of lights on are represented by grey
bars above) exhibited a circadian pattern. The peak activity shifted in DD due to the animal's
endogenous, free-running circadian period τ = 21.27 h. **(Cii)** Mean spontaneous ORN spike
frequency across all DD animals in 1 h bins. Data from the individual in Ci is highlighted in red. **(Ciii)**
Mean spontaneous ORN spike frequency across all DD animals in 1 h bins during the first 48 hours
after aligning the time to the first maximum in spontaneous activity (see Methods) revealed
circadian activity. **(Di)** The spontaneous spike frequency of one long-term ORN recording in
constant darkness with infusion of the Orco antagonist OLC15 (orange frame) dissipated the
circadian rhythm of spontaneous activity. **(Dii)** Mean spontaneous ORN spike frequency across all
OLC15 animals in 1 h bins. Data from the individual in **Di** is highlighted in red. In contrast to LD and
DD conditions, the spike frequency decreases over time with prolonged exposure to OLC15. **(Diii)**
Mean spontaneous ORN spike frequency across all OLC15 animals in 1 h bins (mean ± SD) during
the first 48 hours after aligning the time to the first maximum in spontaneous activity (see
Methods). In contrast to LD and DD conditions, the circadian change in spontaneous spiking
frequency disappeared. **(E)** The slopes of the linear fits to the binned spiking activity of each
individual animal in the three different conditions (see Methods). Each dot indicates the slope for
one experiment. The line indicates the mean. The slope for OLC15 is significantly different from
control LD.

1246

Figure 4: Blocking Orco removed circadian modulation of spontaneous ORN spiking patterns.
RAIN analyses demonstrated significant circadian rhythmicity for most animals in most attributes of the spontaneous spiking activity in LD and DD conditions but not in OLC15 (**A**). Values and background colors indicate the percentage of animals that expressed significant circadian rhythmicity. Circadian differences of attributes were further quantified between the time windows of low vs. high activity (Wilcoxon signed-rank test, α = 0.05; **B-H**). Significant differences occurred mostly in LD (white boxes) and DD (gray boxes) conditions, but never when Orco was blocked with OLC15 (orange boxes). In both LD and DD, the mean spiking frequency was increased during high activity (**Bi, Bii**). Only in LD but not in DD, the mean burst frequency (**Ci, Cii**) increased, whereas the relative number of spikes belonging to a burst (**Di, Dii**) decreased significantly during the high activity period. Both in LD and DD the mean event frequency (**Fi,Fii**) increased significantly. Mean burst duration (**Ei, Eii**), mean number of spikes per bust (**Gi,Gii**), and mean intra-burst intervals (**Hi, Hii**) did not differ between low vs. high spiking activity in LD and DD. When blocking Orco, the attributes for low vs. high activity did not differ for any of the attributes tested (**Biii-Hiii**). In DD and OLC15conditions, activity phases were aligned as described in the Methods; thus, low activity in

1263   DD was at subjective ZT 19 and in OLC15 at subjective ZT 13, high activity was in both cases at
1264   subjective ZT 24. ZTs for low and high activity in LD were 10 and 0, respectively.

1265

1266

Figure 5: Blocking Orco removed circadian regulation of ultradian rhythms in the spontaneous ORN spiking pattern.

Heat map of the instantaneous frequencies (1/ISI) over two consecutive days of long-term tip recordings of one pheromone-sensitive long trichoid sensillum under different conditions (LD **(A)**, DD **(B)**, OLC15 **(C)**) in each panel. Representative recordings from one animal in each panel. Pixel color indicates the counts of instantaneous frequencies in that respective bin. The points fall mostly in a band of high frequencies (>10 – ~100 Hz) and low frequencies (0.01 – ~10 Hz), with the high-frequency band representing the instantaneous frequencies of spikes within a burst and the low-frequency band the instantaneous frequency between bursts. **(A, B)** The high frequency band indicates daily and circadian modulation of frequency prevalence. In addition, the low frequency band (<0.01 Hz to ~ 10 Hz) also displayed circadian modulation of the frequency composition. **(C)** Infusion of OLC15 deleted the circadian modulation in both frequency bands but not the ultradian rhythm of frequency prevalence.

1280

Figure 6: Blocking Orco affected ultradian and infradian frequencies in the spontaneous ORN firing patterns.

The Fourier analysis of spontaneous spiking activity revealed rhythms with ultradian (<20 h; dark grey), circadian (20-28 h; white), and infradian periods (>28 h; light grey) in LD (**Ai**), DD (**Bi**), and OLC15 (**Ci**). Each line represents one animal where each triangular marker is at the local maximum in the frequency spectra obtained for that animal. The histograms above illustrate how often

specific periods of spiking activity rhythms occurred averaged over all animals. Circadian rhythms
were detected in LD (N = 5 of 11), DD (N = 7 of 10), and OLC15 (N = 4 of 12). Wavelet analysis of LD
(**Aii**), DD (**Bii**), and OLC15 (**Cii**) confirmed the occurrence of multiscale periods. Example plots for
one animal each, the same animals as in Figure 5. For each panel, the top plot depicts the mean
firing frequency, the middle plot the wavelet power in the same period range as panels **i,** and the
bottom plot the wavelet power for infradian periods up to 7 h on a log scale to highlight infradian
time scales. Gray dashed lines indicate cone of uncertainty.

1295

1296

Figure 7: Orco is not under control of the TTFL-based molecular circadian clock.

Relative expression levels of *Orco* and *tim* mRNA from male hawkmoth antennae raised under LD conditions. *Orco* expression levels did not differ significantly between ZTs (ANOVA on ranks, $H(5) = 6.91$, $p = 0.227$: N = 4 per ZT). *tim* expression levels changed significantly throughout the day (1-way ANOVA, $F(5, 18) = 6.215$, $p = 0.002$) Red lines indicate mean, *tim* served as positive control.

1302

1303

Figure 8: The model adequately reproduces the firing pattern of the biological ORN.

(A) The model activity (bottom trace) displays both isolated spikes and bursts of variable length as in the original recording (top trace). The model represents an intracellular recording whereas the biological recordings were done with extracellular electrodes. **(B)** Semi-logarithmic plot of burst length distribution of an experimental LD recording (top) and simulated model (bottom) of spontaneous action potential activity of pheromone-sensitive neurons. Both the model and the recordings display exponential decrease of the count's density as a function of the burst length, shown as a linear decrease in the logarithmic scale. **(C)** Comparison of the circadian modulationof model and ORN spiking attributes. Top panels (**Ci, Cii**) show mean spike frequency, whereas bottom panels (**Ciii, Civ**) show the percentage of spikes belonging to bursts. The simulated model reproduced the circadian dynamics observed in the experimental recordings, demonstrating good agreement between model and biological data. **(D)** Heat map of the simulated model, pixel color indicates the density of instantaneous frequencies in that respective bin. The points can be roughly divided in a band of high frequencies (top, ~50 – 80 Hz) and low frequencies (bottom, ~0.1 – 5 Hz), with the high-frequency band representing the instantaneous frequencies of spikes within a burst and the low-frequency band the instantaneous frequency between bursts. **(Di)** Simulated spikes with circadian regulation by the Orco channel; the inter-burst frequencies varied widely with circadian rhythmicity while the frequencies within burst remained approximately constant. The simulated results also reproduced the "merging" effect between the two bands when the bottom

1323   one approaches the top one. **(Dii)** Simulated spikes without circadian regulation of Orco had no
1324   circadian oscillations.

1325

1326

Figure 9: cAMP modulates open-time probability of Orco in M. sexta.

Perfusion of 8-Br-cAMP-AM significantly increased the spontaneous spiking activity at the end of the activity phase (ZT 1-3; **A**; Wilcoxon signed-rank test Z = -2.701; p = 0.004). However, activity did not increase when 8-Br-cAMP-AM was added together with the Orco blocker OLC15 (**B**; paired t-test t(9) = -1.446; p = 0.182). Electrodes were filled with SLR containing 0.1% and 0.15% DMSO as control in **A** and **B**, respectively. Data from individual animals are connected by lines.

1333

9 Supplementary Material

9.1 Mathematical Model

The mathematical model is a 22-dimensional system of stochastic differential equations designed to describe the behavior of ion channels and their effects on the membrane potential. I_{LVA} represents a low-voltage-activated calcium current, while I_{Na} and I_K are the transient sodium and potassium currents responsible for action potential generation. I_{BK} is a large-conductance potassium channel that is activated by both a depolarized membrane potential and an increased intracellular calcium concentration, and I_L is the leak current. Additionally, the Orco channel is a leaky, non-specific ion channel gated by cAMP, with the cAMP concentration oscillating in a circadian rhythm. To model the Orco conductance, we used a sinusoidal function of Zeitgeber time (ZT), independent of the membrane potential, as shown in Figure 2B, C. Each of these ion channels has been reported in vitro in previous studies (Dolzer et al., 2021).

The deterministic part of the 22D model is given by the mean field of the channel-based Langevin model. This approach was introduced by Fox and Lu's (1994) for a 14D HH model. In their framework, one dimension tracks the membrane potential, while the other 13 dimensions correspond to the fraction of channels in each of the five states in the potassium (K^+) Markov chain and the eight states in the sodium (Na^+) Markov chain. Pu and Thomas (2020) demonstrated that the deterministic 14D model and the original 4D HH model are dynamically equivalent, meaning that each solution of the 4D model corresponds to a solution of the 14D model.

Expanding upon Pu and Tomas' methodology to represent out system, we added one dimension to track the calcium concentration, six dimensions for each possible state of the LVA channel, and two dimensions for each possible state of the BK channel. The state vectors for each ion channel are therefore defined as follows: a five-component vector NK for the K^+ gates, an eight-component vector NNa for the Na^+ gates, a six-component vector NLVA for the LVA gates, and a two-component vector NBK for the BK gates. The state vectors are represented as:

$$N^K = [n_{K0}, n_{K1}, n_{K2}, n_{K3}, n_{K4}]^T \in [0,1]^5 \tag{1}$$

$$N^{Na} = [n_{Na00}, n_{Na10}, n_{Na20}, n_{Na30}, n_{Na01}, n_{Na11}, n_{Na21}, n_{Na31}]^T \in [0,1]^8 \tag{2}$$

$$N^{LVA} = [n_{LVA00}, n_{LVA10}, n_{LVA20}, n_{LVA01}, n_{LVA11}, n_{LVA21}]^T \in [0,1]^6 \tag{3}$$

$$N^{BK} = [n_{BK0}, n_{BK1}]^T \in [0,1]^2 \tag{4}$$

The K⁺ channel consists of four independent n (activation) gates, forming a five-vertex

channel-state diagram with eight directed edges. The channel conducts a current only when it is in

the rightmost state (Figure 2D). Similarly, the Na⁺ channel is composed of three identical n gates

and one h (inactivation) gate, forming an 8-vertex diagram with 20 directed edges, one of which is

conducting. The LVA channel includes two identical n gates and one h gate, resulting in a 6-vertex

diagram with 14 directed edges, one of which is conducting. The BK channel consists of a single n

gate, forming a 2-vertex diagram with two directed edges, one of which is conducting. The Markov

chain for every channel is shown in Figure 2B.

For a sufficiently large number of channels, Schmidt and Thomas (2014) and Schmidt et al.

(2018) demonstrated that, under voltage clamp, the equations governing the system can be

approximated by a multidimensional Ornstein-Uhlenbeck (OU) process, or Langevin equation, as

follows:

$$dN^{ion} = \sum_{k=1}^{k_{max}} \zeta^{ion} \left\{ \alpha_k^{ion}(V) N_{i(k)}^{ion} dt + \sqrt{\epsilon^{ion} \alpha_k^{ion} N_{i(k)}^{ion} dW_k^{ion}} \right\}$$ (5)

1373

1374

Here ζ^{ion} is the stoichiometry vector for the kth directed edge. If we write i(k) for the source

node and j(k) for the destination node of edge k, then $\zeta^{ion} = e_{j(k)}^{ion} - e_{i(k)}^{ion}$. The voltage-dependent

per capita transition rate along the kth edge is $\alpha_k^{ion}(V)$ and $N_{i(k)}^{ion}$ is the fractional occupancy of the

source node for the kth transition. The finite-time increment in the Poisson process is approximated

by a gaussian process, namely, the increment dW_k^{ion} in a Wiener (Brownian motion) process

associated with each directed edge. These independent noise terms are scaled by $\epsilon^{ion} = \frac{1}{N_{Tot}^{ion}}$,

where $\frac{1}{N_{Tot}^{ion}}$ represents the total number of that specific ion channel.

The membrane potential is governed by the current balance equation:

$$C \frac{dV}{dt} = \sum_{\chi} I_\chi$$ (6)

1383

1384

1385   where the set χ contains the six ionic currents included in the model:

$$\chi \in \{LVA, Na, K, BK, L, ORCO\}$$ (7)

1386

1387

1388   Each of the six ionic currents included in the model are described as:

1389

$$I_{Na} = -g_{Na}N_8^{Na}(V - V_{Na}) \qquad (8)$$

$$I_K = -g_K N_5^K(V - V_K) \qquad (9)$$

$$I_{LVA} = -g_{LVA}N_6^{LVA}(V - V_{LVA}) \qquad (10)$$

$$I_{BK} = -g_{BK}N_2^{BK}(V - V_{BK}) \qquad (11)$$

$$I_{Orco} = -g_{Orco}\left(0.8 + 0.2\,sin(\frac{2\pi t}{24h})\right)(V - V_{Orco}) \qquad (12)$$

$$I_L = -g_L(V_L - V) \qquad (13)$$

1390
1391   Here, g_χ is the maximal conductance, V is the membrane potential, and V_χ is the reversal

1392   potential for each ion current. The parameter values for each current can be found in Table 1.

1393   Each transition rate $\alpha_k^{ion}(V)$ is calculated in the following way:

1394

$$\alpha_k^{ion}(V) = \frac{p_{j(k)\infty}^{ion}(V)}{\tau_p(V)} \qquad (14)$$

1395
1396

where $p_{j(k)\infty}^{ion}(V)$ is the steady state value of the specific gate that switches states in the

destination state and $\tau_p(V)$ is the time constant of said gate. The steady state activation values and

time constants for the LVA, Na, K, and BK channels were based on Viertel and Borisyuk (2019). The

steady state functions for the activation gates are denoted with m_∞^{ion}, and those for the inactivation

gates with h_∞^{ion}.

Each gate's steady state activation function, except for the BK channel, takes the form:

1403
1404

$$p_{j\infty}^{ion}(V) = \frac{1.0}{1.0 + exp\left(\frac{(V - \theta_p)}{\beta_p}\right)} \qquad (15)$$

1405

1406   The I_{BK} current is dependent on the intracellular calcium concentration and membrane

1407   potential as represented by the following equations:

1408
1409

$$m_\infty^{BK}(V) = \frac{1.0}{1.0 + exp\left(\frac{\left(V - V_{\frac{1}{2}}\right)}{(-15.6)}\right)} \qquad (16)$$

$$V_{\frac{1}{2}} = -20 + 59.2e^{-90[Ca]} + 96.7e^{-470[Ca]} \qquad (17)$$

$$\tau_\alpha = -(q - 1.0)\frac{(f - 0.2)}{0.8} + 170 \qquad (18)$$

$$q = 2.9 + 6.3e^{-360[Ca]} \qquad (19)$$

$$s = -25.3 + 107.5e^{-120[Ca]} \qquad (20)$$

$$f = \frac{1.0}{10.0\left(e^{\frac{-(V+100.0-s)}{63.6}} + e^{\frac{(-150.0+(V+100.0-s))}{63.6}}\right)} \qquad (21)$$

1410
1411

1412   The intracellular concentration of calcium ions is described by the following equation:

1413

$$\frac{d[Ca]}{dt} = -\gamma\frac{I_{LVA}}{zFd} + \frac{[Ca_0] - [Ca]}{\tau_{Ca}} \qquad (22)$$

1414

1415   where Ca_0 is the steady state calcium concentration, γ describes calcium buffering, z is the

1416   ionic valence of calcium, F is Faraday's constant and d is the depth in microns at which calcium

1417   ions concentrate near the membrane.

1418   The parameters of each steady state activation function as well as the time constant for each

1419   gating value can be found in Table 2.

1420

1421   **Table 1: Parameter values for model currents.**

Current	g_X (mS/cm^2)	V_X (mV)
I_L	4.3	-62.5
I_{LVA}	10.0	120
I_{NA}	29.17	45.0
I_K	12.96	-105.0
I_{BK}	9.0	-105.0
I_{ORCO}	5.0	0.0

1422

1423   **Table 2: Gating variable parameter values.**

Variable	θ	β	τ(ms)

m^{LVA}	-30	-4.8916	$\dfrac{40.0}{cosh\left(\dfrac{(V + 37.1)}{9.7832}\right)}$
h^{LVA}	-59.2	11.2326	$\dfrac{350.0}{cosh\left(\dfrac{(V + 59.2)}{22.4652}\right)}$
m^{Na}	-25.0	-6.5	0.2
h^{Na}	-26.0	9.0	$\dfrac{10.0}{cosh\left(\dfrac{(V + 26.0)}{18.0}\right)}$
m^{K}	-26.0	-9.0	$\dfrac{10.0}{cosh\left(\dfrac{(V + 26.0)}{18.0}\right)}$

1424

1425

1426

**Figure 3 – Supplement: Spontaneous spiking activity of ORNs of pheromone-exposed male *M.*

sexta.**

To enhance synchronization, males were exposed to females and pheromones as additional

zeitgeber for 30 mins at ZT 16 before the start of the experiment. Recordings were obtained over

multiple days in DD. **(A)** The spontaneous spike frequency of one long-term ORN recording in

constant darkness (DD, the expected times of lights on are represented by grey bars above) **(B)**

Mean spontaneous ORN spike frequency across all DD animals in 1 h bins. Data from the individual

in **A** is highlighted in red. **(C)** Mean spontaneous ORN spike frequency across all DD animals in 1 h

bins during the first 48 hours after aligning the time to the first maximum in spontaneous activity

(see Methods). No circadian activity was evident in these pheromone-exposed males.