Preclinical evaluation of [11C]CHDI-009R for quantification of mutant huntingtin aggregates

Authors

Franziska Zajicek[1,2], Liesbeth Everix[1,2], Annemie Van Eetveldt[1,2], Jeroen Verhaeghe[1,2], Stef De Lombaerde[1,3], Alan Miranda[1,2], Jordy Akkermans[1,2], Celia Dominguez[4], Vinod Khetarpal[4], Jonathan Bard[4], Longbin Liu[4], Steven Staelens[1,2], Daniele Bertoglio[2,5]

[1]*Molecular Imaging Center Antwerp (MICA), University of Antwerp, Antwerp, Belgium*

[2]*µNeuro Research Center of Excellence, University of Antwerp, Antwerp, Belgium*

[3]*Department of Nuclear Medicine, Antwerp University Hospital, Antwerp, Belgium*

[4]*CHDI Management, Inc., the company that manages the scientific activities of CHDI Foundation, Inc., Los Angeles, CA, USA*

[5]*Bio-Imaging Lab, University of Antwerp, Antwerp, Belgium*

Corresponding author:

Dr. Daniele Bertoglio

Bio-Imaging Lab, Faculty of Pharmaceutical, Biomedical and Veterinary Sciences, University of Antwerp, Universiteitsplein 1, Wilrijk, Belgium

Tel. +32 3 265 8736

Email: daniele.bertoglio@uantwerpen.be

ORCID

Franziska Zajicek: 0000-0001-5410-687X

Liesbeth Everix: 0000-0001-5985-1609

Annemie Van Eetveldt: 0000-0002-6488-5456

Jeroen Verhaeghe: 0000-0002-4493-1902

Stef De Lombaerde: 0000-0002-5307-3575

Alan Miranda: 0000-0002-5381-015X

Jordy Akkermans: 0000-0002-1533-9504

Vinod Khetarpal: 0000-0002-2175-5117

Jonathan Bard: 0000-0002-5097-1361

Longbin Liu: 0000-0003-3995-0196

Steven Staelens: 0000-0003-3376-0519

Daniele Bertoglio: 0000-0003-4205-5432

37   **Keywords**

Abstract

Huntington's disease (HD) is a neurodegenerative disorder caused by an expanded trinucleotide repeat in the huntingtin gene (*HTT*) that subsequently leads to aggregation of the mutant huntingtin (mHTT) protein. Thus, lowering mHTT is a key therapeutic approach used by several candidate therapeutics currently under investigation. Visualization of the efficiency of these therapeutics through *in vivo* mHTT quantification rises in importance. For positron emission tomography (PET) imaging of mHTT aggregates, it is critical to characterize the *in vivo* kinetic profile of newly identified mHTT binders to assess their translational application. Here, we report the evaluation of [¹¹C]CHDI-009R, a PET imaging radioligand with higher affinity and selectivity for mHTT aggregates than previously reported radioligands, in the heterozygous (HET) zQ175DN mouse model of HD and wild-type (WT) littermates at 9 and 3 months of age.

[¹¹C]CHDI-009R displayed high stability in plasma and brain, which was reflected in brain kinetics as demonstrated by rapid uptake followed by relatively slow elimination. Kinetic modeling and volume of distribution $V_{T\ (IDIF)}$ indicated the radioligand's ability to quantify mHTT aggregation at 9 months of age with clear genotype differentiation ($p<0.0001$). [¹¹C]CHDI-009R showed an excellent test-retest reliability in 9-month-old mice (intraclass correlation coefficient: 0.62 - 0.79). A phenotypic difference in mHTT aggregates was also observed in 3-month-old mice in several brain structures ($p<0.05$) and was confirmed with [³H]CHDI-009R autoradiography.

Overall, this study suggests [¹¹C]CHDI-009R is a promising radioligand for the detection of cerebral mHTT aggregates in a mouse model of HD and supports its advance to clinical evaluation.

Introduction

Huntington's disease (HD) is a progressive neurodegenerative disorder caused by an expanded trinucleotide (CAG) repeat in exon 1 of the huntingtin (*HTT*) gene. The transcription and translation of mutant *HTT* leads to the aggregation of mutant huntingtin (mHTT) protein [1, 2], which, alongside mHTT fragments, is involved in cellular dysfunction and loss of neurons [3, 4]. With several disease-modifying therapeutic interventions aimed to lower mHTT expression under clinical investigation [5-7] (e.g. ClinicalTrials.gov identifier (ID): NCT06024265; NCT05032196), there is an increasing interest in accurately quantifying their ameliorative effect. Currently, the estimate of the central nervous system (CNS) target engagement of HTT-lowering therapeutics is indirect, relying on extrapolations of soluble HTT levels in non-brain compartments, such as cerebrospinal fluid, both in the preclinical [8] and clinical setting [9] (www.clinicaltrials.gov, identifier NCT02519036). Still, fluid biomarkers lack brain region-specific information that positron emission tomography (PET) imaging of mHTT aggregates could provide. Thus, PET imaging holds great potential as a complementary tool to assess brain region-specific target engagement, as previously demonstrated preclinically [10]. Thus, enabling accurate spatial and temporal evaluation of efficacy with potential therapeutic interventions.

We recently described a number of novel radiotracers for mHTT aggregate imaging in preclinical imaging studies, namely [11C]CHDI-626 [11-13], [11C]CHDI-180R [10, 12, 14, 15] and [18F]CHDI-650 [16, 17] with varying levels of applicability in mice. Additionally, [11C]CHDI-626 [18] was advanced to the clinic but later discontinued due to an undesirable brain uptake and decay profile. On the other hand, [11C]CHDI-180R detected a significant difference between people with HD (PwHD) and healthy controls; however, it also displayed high intersubject variability [19]. Thus, a more suited clinical candidate would be preferred, and the identification of novel radioligands targeting mHTT aggregates with superior PET radiotracer properties remains a priority. Especially regarding clinical importance, enhanced sensitivity of

mHTT aggregate detection in 3-month-old zQ175DN heterozygous (HET) mice is desired, as

mHTT aggregation is expected to be lower in abundance in humans than in mouse models.

Therefore, we report on the investigation of a novel radioligand, [^{11}C]CHDI-009R, that has

been shown to have an almost 3-fold higher maximum binding capacity than CHDI-180R

based on autoradiography studies in homozygous zQ175DN brains [20]. Here, we

characterize the *in vivo* performance of this tracer in wildtype (WT) and HET zQ175DN knock-

in mice through radiometabolite analysis, test-retest variability estimation, quantitative

phenotypic comparison at two different ages, and post-mortem autoradiography.

Methods

Animals

A total of 63 nine-month-old male HET (*n*=33) and WT (*n*=30) as well as 44 three-month-old male HET (*n*=24) and WT (*n*=20) zQ175DN mice (B6.129S1-*Htt*$^{tm1.1Mfc}$/190ChdiJ JAX stock #029928), were obtained from Jackson Laboratories (Bar Harbour, Maine, USA) [21, 22]. Due to sporadic congenital portosystemic shunt occurring in C57BL/6J mice [23], all animals were screened at Jackson Laboratories before shipment, and hence, all animals used in this study were shunt-free. Study design, group size, body weight, and age of the animals are reported in Fig. 1 and Tables S1-S3. Animals were single-housed at 9 months of age and group-housed at 3 months of age in ventilated cages under a 12 h light/dark cycle. While group-housing is generally considered to achieve better social well-being for mice, minimising aggression and potential injuries caused by dominant-submissive behaviour was prioritised for animals undergoing several imaging sessions. The animals were kept in a temperature- and humidity-controlled environment with food and water *ad libitum*. The animals were given at least one week for acclimatization after arrival before the start of any procedure. Experiments followed the European Committee (decree 2010/63/CEE) and were approved by the Ethical Committee for Animal Testing (ECD 2019-39) at the University of Antwerp (Belgium).

Radioligand synthesis

[^{11}C]CHDI-009R was prepared using an automated synthesis module (Carbonsynthon I, Comecer, The Netherlands). The radioligand was prepared via single-step carbon-11 labeling at the pyridazinone-NH site, starting with a solution of the precursor (0.3 mg) in DMF (0.5 ml), which was reacted with excess of [^{11}C]CH$_3$OTf in the presence of NaOH (1 M, 5 μl) for 3 min at room temperature. To terminate the reaction and to ensure good retention of the compound of interest on the semi-preparative HPLC column, acetonitrile/water (35:65, v/v, 2 ml) was introduced to the reaction mixture, and the resulting crude product was purified on HPLC using

an ACE 5 C18-HL 250x10 mm column (ACE, UK), eluted with a mixture of ammonium formate

(0.1 M) in acetonitrile (35/65, v/v) as mobile phase, at a flow rate of 6 ml/min. The collected

fraction was diluted with water (20 ml) before being loaded on a 1 cc 10 mg Oasis HLB

cartridge (Waters, MA, USA). The cartridge was washed with water for injection (3 ml) and

eluted through a sterile filter with ethanol (0.5 ml) and saline (4.5 ml) to afford the final product

solution.

The radiochemical purity of the produced [^{11}C]CHDI-009R was determined using a Kinetex

EVO C18, 5μm, 150×4.6 mm (Phenomenex, USA) HPLC column, with a solution of sodium

acetate (NaOAc) (0.05 M, pH 5.5) in acetonitrile/ (30:70, v/v) as mobile phase, at a flow of 1

ml/min, with UV absorbance set at 289 nm. Based on 37 productions, radiochemical purity

was greater than 99%, a molar activity (A_m) of 93.59±27.61 GBq/μmol, and a radiochemical

yield of 7.83±0.03% was achieved.

Radiometabolite analysis

Evaluation of *in vivo* plasma and brain radiometabolite profiles was performed at 5, 20, and

40 min post-injection (p.i.). Mice were injected via the lateral tail vein with [^{11}C]CHDI-009R

(WT: 3.78±0.87 MBq; HET: 3.75±0.62 MBq in 200 µl), and blood was withdrawn via cardiac

puncture at designated times, and brains were immediately excised. The homogenization and

extraction were conducted as previously described [10, 17]. The extraction efficiency from

plasma samples (percentage of recovery of radioactivity) was 98.4±2.6%, whereas in brain

samples, an extraction efficiency of 100.6±4.7% was achieved. Plasma or brain supernatant

was loaded onto a pre-conditioned reverse-phase (RP)-HPLC system (Luna C18(2), 5 µm

HPLC column (250x4.6 mm) + Phenomenex security guard pre-column) and eluted with

NaOAc 0.05M pH 5.5 and acetonitrile: (87:13 for 10 min v/v) buffer at a flow rate of 1 ml/min.

RP-HPLC fractions were collected at 0.5 min intervals for 10 min. Pure tracer (5μCi) was

loaded onto the same HPLC setup (Fig. S1). Control experiments confirmed that no apparent

degradation of the radiotracer occurred during procedural work-up in both plasma

(99.91±0.00% of intact tracer) and brain (92.14±0.07% of intact tracer).

Dynamic PET imaging acquisition

Dynamic microPET/Computed tomography (CT) images were acquired on Siemens Inveon

PET-CT scanners (Siemens Preclinical Solution, Knoxville, USA). Animal preparation was

performed as previously described [10, 24]. Animal and dosing information for both WT and

HET mice of all paradigms are given in the supplementary Tables S1-S3.

PET data were acquired in list mode format. Following the microPET scan, a 10 min 80 kV/500

µA CT scan was performed for attenuation and scatter correction. The same microPET scan,

was also used to estimate the blood radioactivity through a non-invasive image-derived input

function (IDIF) to estimate the volume of distribution ($V_{\text{T (IDIF)}}$) as a surrogate of V_{T}.

Acquired PET data were reconstructed into 39 frames of increasing length (12x10s, 3x20s,

3x30s, 3x60s, 3x150s, and 15x300s). Images were reconstructed using a list-mode iterative

reconstruction proprietary spatially variant resolution modeling in 8 iterations and 16 subsets

of the 3D ordered subset expectation maximization (OSEM 3D) algorithm [25]. Normalization,

dead time, and CT-based attenuation corrections were applied, and PET image frames were

reconstructed on a 128x128x159 grid with 0.776x0.776x0.796 mm^3 voxels.

Image processing and analysis

Image processing and analysis were performed with PMOD 3.6 software (PMOD

Technologies, Zurich, Switzerland) as previously described [11]. For spatial normalization of

the PET/CT images, brain normalization of the CT image to a CT template was performed by

adapting the previously described protocol [26]. We applied the volumes-of-interest (VOIs)

described in previous studies [10, 11], adapted from the Waxholm atlas [27]. Using those VOIs,

time activity curves (TACs) were then extracted for individual brain regions. The IDIFs were

extracted from the PET images by generating a VOI in the lumen of the left ventricle at an

early time frame exhibiting maximal activity in the heart and applying a threshold of 50% of the

max signal as previously described [24, 28-30]. Standardized uptake value (SUV) was

calculated as activity concentration [kBq/cc] divided by injected dose [kBq] per body weight

[g].

These regional TACs and IDIFs were used to estimate $V_{T\,(IDIF)}$ values through kinetic modeling.

$V_{T\,(IDIF)}$ was estimated by fitting the extracted TACs with standard two-tissue compartmental

model (2TCM), with blood volume fraction (V_B) fixed at 3.6% [10, 11] and Logan linear model

to calculate $V_{T\,(IDIF)}$. Pixelwise kinetic modeling was performed using the Logan graphical

analysis in PXMOD (PMOD 4.2). The averaged parametric maps were subsequently overlaid

onto a Waxholm MRI template.

Tissue collection for Autoradiography

Animals were euthanized by decapitation while under anaesthesia. Brains were dissected and

fresh-frozen in 2-metylbuthane at $-35°C$ for 2 min and preserved at $-80°C$. Coronal sections

of 10 μm thickness were collected at 1.10 mm from the bregma according to Paxinos and

Franklin [31] in triplicate using a cryostat (Leica, Germany) on Superfrost Plus slides (Thermo

Fischer Scientific, USA).

Autoradiography

[³H]CHDI-009R was provided by Novandi Chemistry AB (Sweden) with a A_m of 82 Ci/mmol.

Sections were air-dried at room temperature, pre-incubated for 20 min with binding buffer

(50mM Tris-HCl + 120mM NaCl + 5mM KCl + 2mM $CaCl_2$ + 1mM $MgCl_2$, pH 7,4) and air-dried

using warm airflow. Dried sections were subsequently incubated with total binding (TB)

solution (0.5 nM of [³H]CHDI-009R in binding buffer) and non-specific binding (NSB) solution

(0.5 nM of [³H]CHDI-009R + 10 μM of cold ligand in binding buffer) for 1h at room temperature.

Subsequently, sections were washed three times for 10 min in ice-cold washing buffer (50mM

Tris-HCl, pH 7,4), followed by a brief wash in ice-cold distilled water, and dried for 2h at room

temperature. Finally, all sections were exposed on imaging plates (BAS-TR2025, Fujifilm,

Japan) for 120 min. Radioactivity was detected with a phosphor imager (Fuji FLA-7000 image

reader) and quantified based on intensity values calculated using commercial tritium

standards (American Radiolabelled Chemicals Inc., USA). For all samples, the entire surface

of the coronal brain slice was used for quantification. The specific binding (SB) was determined

by subtracting NSB from TB, which were both measured in triplicate and subsequently

averaged. The SB averages were used for statistical analysis. SB was set to 0 if the values of

NSB and TB were below the limit of detectability of the calibration curve. Analysis was

performed blind to conditions in Fiji, ImageJ (National Institute of Health, USA, version: 2.1).

Statistical analysis

The Shapiro-Wilk test revealed that *in vivo* data were normally distributed, while *in vitro* data

demonstrated a non-Gaussian distribution; therefore, non-parametric testing was performed

on *in vitro* data. All data was visually inspected for outliers, followed, if necessary, by statistical

outlier testing with ROUT (Q=0.5%). Two outliers in motor cortex for 9-month-old HET mice

were identified and excluded. A Pearson's correlation with simple linear regression was

applied to $V_{T\,(IDIF)}$ values from 2TCM and Logan kinetic modeling for model verification. A

Spearman's correlation with interpolation (sigmoidal, 4PL, X is concentration) was applied to

injected masses and $V_{T\,(IDIF)}$ values from Logan for visualization of a potential mass dose

effect. Regular two-way ANOVA with *post hoc* Bonferroni correction for multiple comparisons

was applied to analyze $V_{T\,(IDIF)}$ value differences between genotypes.

The individual relative test-retest variability (rTRV) was calculated as follows:

228
$$\frac{test - retest}{average\ (test, retest)}$$

229   The individual absolute test-retest variability (aTRV) was calculated as follows:

230
$$\left| \frac{test - retest}{average\ (test, retest)} \right|$$

In addition, a Bland-Altman analysis was performed to assess the agreement between two

measures on an individual level. A Mann-Whitney test was performed to investigate

phenotypic differences in SB in autoradiography.

All statistical analyses above were performed with GraphPad Prism (version 9.2.0) statistical

software. The Intraclass correlation coefficient (ICC) for the test-retest variability study was

calculated with the software JMP Pro 17 (v17.0.0). Group and timepoint interaction was set

as fixed effect. Animal ID, timepoint, interaction of animal ID and group, and interaction of

animal ID and timepoint, were set as random effects. No effect was identified for either

interaction. The ICC was reported as the ratio of the animal variance component for the effect

to the sum of positive variance components. One-tailed power analyses were performed in

G*Power (v3.1.9.6) with significance level (α) at 0.05, confidence (β) set to 80%. Data are

represented as mean ± standard deviation (SD) or as mean ± standard error of the mean

(SEM). All tests were two-tailed unless stated otherwise. Statistical significance was set at

p<0.05 with the following standard abbreviations used to reference p values: ns, not

significant; *p<0.05; **p<0.01; ***p<0.001; ****p<0.0001.

Results

[¹¹C]CHDI-009R demonstrated high metabolic stability

[¹¹C]CHDI-009R radiometabolite analysis revealed a high stability over time in both brain (e.g. 40 min: WT=90.9±8.6%; HET=93.8±2.0%; Fig. 2a) and plasma (e.g. 40 min: WT=95.1±1.7%; HET=96.4±1.2%; Fig. 2b) and with no observable peak other than the intact radioligand (Fig. S1). In addition, the plasma-to-whole-blood ratio was stable over time with no difference observed between genotypes (WT=1.8±0.1; HET=1.7±0.2; Fig. 2c). IDIF profiles of both genotypes displayed a comparable peak within 2 min of radioligand injection and a similar stable signal after 5 min p.i. until the end of acquisition (Fig. 2d). Given the high stability and no difference between genotypes, no corrections were applied to IDIFs for accurate $V_{T\,(IDIF)}$ estimations.

[¹¹C]CHDI-009R kinetic modeling with 90 min scan duration

The averaged striatal SUV TACs of [¹¹C]CHDI-009R demonstrated rapid penetration into the brain in both genotypes, but a subsequent slower wash-out was observed in the brain in HET animals compared to WTs (Fig. 3a). We further observed an excellent agreement between 2TCM quantification and Logan graphical analysis (striatum: r^2=0.996; p<0.0001) with minimal deviation from the identity line (slope = 0.998) (Fig. 3b); thus, supporting the use of Logan graphical analysis to quantify [¹¹C]CHDI-009R in mice.

Next, we performed a $V_{T\,(IDIF)}$ time stability analysis to estimate the shortest recommended acquisition time. Although we observed no impact on $V_{T\,(IDIF)}$ in WT animals (Fig. 3c), $V_{T\,(IDIF)}$ in HET mice was susceptible to a steady underestimation, which passed the 90% threshold with a 60 min acquisition (Fig. 3d). Thus, shortening the scan acquisition time from 90 min was not recommended to avoid even minimal underestimation of a phenotypic effect. Further, we investigated the impact of injected mass on [¹¹C]CHDI-009R $V_{T\,(IDIF)}$. As shown in Fig. S2,

a definite negative association was observed; thus, the recommended upper limit for injectable

mass was set to 1.7 μg/kg.

[¹¹C]CHDI-009R quantification revealed phenotypic differences

The averaged parametric $V_{\text{T (IDIF)}}$ maps of WT and HET mice shown in Fig. 4a visualized a

clear genotype effect. The two-way ANOVA test revealed a significant genotype effect ($F_{(1, 183)}$

= 987.3; p<0.0001) and the Bonferroni *post hoc* analysis demonstrated statistically significant

increased signal in HET mice compared to their WT littermates in all tested brain regions

(p<0.0001) (Fig. 4b). Phenotypic $V_{\text{T (IDIF)}}$ differences ranged from 55.5±5.4% in the cerebellum

to 105.8±6.2% in the thalamus (Table 1).

[¹¹C]CHDI-009R displayed excellent test-retest reproducibility

To measure the reliability of [¹¹C]CHDI-009R, a test-retest study was performed in WT and

HET zQ175DN mice at 9 months. Parametric $V_{\text{T (IDIF)}}$ maps did not reveal any apparent

differences between scans (Fig. 5a). The aTRV was overall low, indicating a high reliability,

with a range from 5.1±4.0% to 7.7±4.2% in striatum of WT and HET animals (Fig. 5b; Table

2). The Bland-Altman plot displayed close-ranged levels of agreement (-16.0% to 14.7%) (Fig.

5c). In addition, the regional rTRV, with values ranging from -3.1±8.5% to 4.3±8.3% in WT and

from -4.8±6.2% to 1.2±8.3% in HET, demonstrated limited difference across brain regions

(Table 2). Overall, according to the ICC values, [¹¹C]CHDI-009R displayed a good to excellent

test-retest agreement (Table 2).

[¹¹C]CHDI-009R differentiated genotypes at 3 months

In addition to the comprehensive assessment of [¹¹C]CHDI-009R properties at 9 months, a

study was performed at 3 months, when less mHTT aggregates are present in HET zQ175DN

mice, providing insights into the degree of sensitivity for this radioligand. Based on parametric

$V_{T\ (IDIF)}$ maps, an increased signal in HET mice could be detected (Fig. 6a). Accordingly, the

two-way ANOVA test revealed a significant genotype effect ($F_{(1,\ 175)}$ = 62.3; p<0.0001). The

Bonferroni *post hoc* analysis displayed statistically significant phenotypic differences between

genotypes, ranging from 8.3±2.7% to 16.3±2.9% in test regions, with the exception of motor

cortex (5.0±2.8%; not significant) (Fig. 6b; Table 3). The mHTT aggregate binding capacity of

CHDI-009R was also demonstrated through a post-mortem autoradiography study with the

tritiated version of the ligand. [^{3}H]CHDI-009R displayed a significantly different SB in HET

mice compared to WT littermates (Fig. 6c, p<0.0001).

[^{11}C]CHDI-009R is suitable for evaluating therapeutic efficacy in zQ175DN mice

We performed power analyses to estimate group sizes needed for evaluating therapeutic

efficacy (α=0.05, β=80%) in zQ175DN mice. As shown in Table S4, statistical power was

good, as revealed by the range of Cohen's d of 0.59 – 1.70 and 3.52 – 5.15 at 3 and 9 months,

respectively. This indicated that at 9 months, [^{11}C]CHDI-009R could significantly display a 20%

difference in mHTT aggregates with acceptable group arms (e.g. striatum: n=21). At 3 months,

however, only in striatum (n=23) and thalamus (n=26) could a larger effect (50% difference)

between groups be detected with a similar sample size.

Discussion

Here we report the *in vivo* characterization of a novel mHTT aggregate PET radioligand, [¹¹C]CHDI-009R, with higher selectivity [20] than previously investigated radiotracers [12, 15, 16]. As in previous evaluation studies of mHTT aggregate radiotracers, we included only male mice for better comparison, as no sex differences have been observed in HET mice before [32]. Since the current study focused on evaluating the capability of [¹¹C]CHDI-009R in detecting different levels of mHTT aggregates in the brain *in vivo*, only one sex was used. While research on sex differences is of significant importance, the inclusion of both sexes in this study was considered secondary.

We show that [¹¹C]CHDI-009R rapidly enters the brain, followed by wash-out, indicating its reversible binding. The slower rate of radioligand wash-out from the brain of the HET animals was likely due to the presence of mHTT aggregates that are absent in WT. [¹¹C]CHDI-009R demonstrated high stability in mouse plasma, and no metabolite correction was deemed necessary for IDIFs, although we could only obtain data until 40 min p.i. due to the loss of radioactive signal at later time points in the limited blood sample that can be collected as a consequence of the radioisotope's half-life.

The cerebral retention time for both genotypes was suitable for estimating the $V_{T\,(IDIF)}$, for which both 2TCM and Logan graphical analysis were shown as adequate models. The time stability kinetic analysis revealed that a reduction in scan acquisition time was not feasible without introducing a genotype-specific underestimation of the $V_{T\,(IDIF)}$ values, which would result in an artificial decrease of phenotypic differences. The current scan duration of 90 min, however, borders the capability of obtaining data with limited noise due to the short half-life of the isotope (20.4 min), which requires a prompt scan acquisition and keeping the injected mass within the recommended limit of 1.7 μg/kg.

With those optimized PET imaging conditions, quantification with Logan indicated that [¹¹C]CHDI-009R could significantly distinguish genotypes at both 9 and 3 months of age, with

comparable magnitude as observed with previous mHTT radiotracers [10, 17]. Noteworthy,

there was no change in $V_{T (IDIF)}$ values of WT animals across brain regions, indicating that

[11C]CHDI-009R can reliably and uniformly show the lack of a target in WT animals. Moreover,

the post-mortem [3H]CHDI-009R autoradiography confirmed the lack of mHTT aggregates in

WT mice, corroborating the previously described findings [20]. This represents an important

indication of minimum off-target binding that could otherwise potentially diminish the utility of

a radioligand, as shown for several Alzheimer's disease-related radioligands [33].

Besides general affinity and selectivity, intra-subject variability was also an important factor to

be considered. The reliability of [11C]CHDI-009R was demonstrated by the excellent test-retest

performance as determined via Bland-Altman plot and ICC values [34]. This provides a

significant benefit for a more accurate application, especially in longitudinal settings, and

represents a favorable attribute compared to the poor reliability of [18F]CHDI-650 (ICC: 0.00 –

0.27) [17].

The capability of detecting early changes at the molecular level represents another important

utility in the context of monitoring (early) therapeutic intervention, as described in the studies

of HD mouse models [10, 32]. Our current work confirmed the *ex vivo* findings that [3H]-

radiolabeled CHDI-009R [20] can detect statistically significant phenotypic differences at 3

months of age with both *in vivo* PET imaging and post-mortem autoradiography.

We have described several mHTT radiotracers that could identify a significant difference in

striatal binding in 3-month-old zQ175DN HET mice [11, 17]. Especially [11C]CHDI-180R

exhibited high metabolic stability and appropriate brain wash-out for imaging studies in both

HD mouse models as well as in a large non-human primate (NHP) model of HD [35],

suggesting [11C]CHDI-180R as a well-performing mHTT aggregate-directed PET radioligand

for animal models [10, 14]. However, [11C]CHDI-180R demonstrated high intersubject

variability and relatively low signal-to-background ratio in the first-in-patient PET study [19].

The increased SB and binding potential of [3H]CHDI-009R (SB=205.3±10.3 fmol/mg; B_{max}=448

fmol/mg) compared to [^{3}H]CHDI-180 (SB=70.9±11.0 fmol/mg; B_{max}=163 fmol/mg) in HOM

zQ175DN brains likely points towards higher sensitivity to detect mHTT aggregates with

CHDI-009R, as shown in post-mortem HD brains [20]. Further, radio-binding assays revealed

increased binding to mHTT aggregates with CHDI-009R (IC_{50}=0.4±0.2 nM) in contrast to

CHDI-180R (IC_{50}=1.3±1.0 nM), while additionally minimizing the affinity for abeta- and/or tau-

associated aggregates [20]. Thus, [^{11}C]CHDI-009R demonstrates high potential for successful

clinical translation given its improved characteristics over CHDI-180R. The utility of CHDI-

009R as an imaging biomarker will be assessed in a clinical setting.

Conclusions

[^{11}C]CHDI-009R is a highly stable radioligand capable of detecting significant phenotypic

differences between HET and WT zQ175DN mice at different ages in a reproducible manner.

These findings suggest [^{11}C]CHDI-009R to be a suitable radioligand to image mHTT

aggregates in mice. Notably, a clinical investigation is underway to determine its suitability for

assessing mHTT aggregates in humans.

Declarations

Ethics approval

Availability of data and material

The PET datasets generated or analysed during the current study are available in the OpenNeuro repository, https://doi.org/10.18112/openneuro.ds006613.v1.0.1

Acknowledgements

The authors thank Philippe Joye, Caroline Berghmans, Romy Raeymakers and Eleni Van der Hallen of the Molecular Imaging Center Antwerp (MICA) for their valuable assistance. FZ, LE, AVE, JA, AM, JV, SS, and DB are members of the μNeuro Research Centre of Excellence at the University of Antwerp. The authors acknowledge the contribution of the following CHDI staff: Drs Mette Skinbjerg, Robert Doot and Yuchuan Wang for helpful discussions; Drs Simon Noble and Vahri Beaumont for reviewing and providing constructive feedback; Brenda Lager for coordinating animal supply and Braden Boone for project management.

Authors' contributions

Conceptualization: JV, VK, JB, LL, SS, DB

Methodology: JV, AM, SS, DB

Data acquisition: FZ, LE, AVE, SDL

Formal analysis and investigation: FZ, LE, AVE

Data interpretation: FZ, JV, JA, VK, JB, LL, SS, DB

Writing – original draft preparation: FZ, SS, DB

Writing – review and editing: FZ, JV, SDL, JA, CD, VK, JB, LL, SS, DB

Resources: CD, VK, JB, LL, SS, DB

Funding acquisition: SS, DB

Supervision: JV, SS, DB

Competing interests

This work was funded by CHDI Foundation, Inc., a privately-funded nonprofit biomedical

research organization exclusively dedicated to collaboratively developing therapeutics that

improve the lives of those affected by Huntington's disease. CD, VK, JB, and LL are employed

by CHDI Management, Inc., the company that manages the scientific activities of CHDI

Foundation. No other potential conflicts of interest relevant to this article exist.

Funding

This work was funded by CHDI Foundation, Inc. under agreement number A-11627. DB

acknowledges support by the Research Foundation Flanders (FWO) (ID: 1229721N).

University of Antwerp supported the work through professorships for JV, SS, and DB.

References

1. Saudou, F. and Humbert, S., *The Biology of Huntingtin.* Neuron, 2016. **89**(5): p. 910-26.

2. The-Huntington's-Disease-Collaborative-Research-Group, *A novel gene containing a trinucleotide repeat that is expanded and unstable on Huntington's disease chromosomes.* Cell, 1993. **72**(6): p. 971-983.

3. McColgan, P. and Tabrizi, S.J., *Huntington's disease: a clinical review.* Eur J Neurol, 2018. **25**(1): p. 24-34.

4. Waldvogel, H.J., Kim, E.H., Tippett, L.J., Vonsattel, J.P., and Faull, R.L., *The Neuropathology of Huntington's Disease.* Curr Top Behav Neurosci, 2015. **22**: p. 33-80.

5. Tabrizi, S.J., Ghosh, R., and Leavitt, B.R., *Huntingtin Lowering Strategies for Disease Modification in Huntington's Disease.* Neuron, 2019. **101**(5): p. 801-819.

6. Wild, E.J. and Tabrizi, S.J., *Therapies targeting DNA and RNA in Huntington's disease.* The Lancet Neurology, 2017. **16**(10): p. 837-847.

7. Saade, J. and Mestre, T.A., *Huntington's Disease: Latest Frontiers in Therapeutics.* Curr Neurol Neurosci Rep, 2024. **24**(8): p. 255-264.

8. Caron, N.S., Banos, R., Aly, A.E., Xie, Y., Ko, S., Potluri, N., Anderson, C., Black, H.F., Anderson, L.M., Gordon, B., Southwell, A.L., and Hayden, M.R., *Cerebrospinal fluid mutant huntingtin is a biomarker for huntingtin lowering in the striatum of Huntington disease mice.* Neurobiol Dis, 2022. **166**: p. 105652.

9. Kordasiewicz, H.B., Stanek, L.M., Wancewicz, E.V., Mazur, C., McAlonis, M.M., Pytel, K.A., Artates, J.W., Weiss, A., Cheng, S.H., Shihabuddin, L.S., Hung, G., Bennett, C.F., and Cleveland, D.W., *Sustained therapeutic reversal of Huntington's disease by transient repression of huntingtin synthesis.* Neuron, 2012. **74**(6): p. 1031-44.

10. Bertoglio, D., Bard, J., Hessmann, M., Liu, L., Gartner, A., De Lombaerde, S., Huscher, B., Zajicek, F., Miranda, A., Peters, F., Herrmann, F., Schaertl, S., Vasilkovska, T., Brown, C.J., Johnson, P.D., Prime, M.E., Mills, M.R., Van der Linden, A., Mrzljak, L., Khetarpal, V., Wang, Y., Marchionini, D.M., Skinbjerg, M., Verhaeghe, J., Dominguez, C., Staelens, S., and Munoz-Sanjuan, I., *Development of a ligand for in vivo imaging of mutant huntingtin in Huntington's disease.* Sci Transl Med, 2022. **14**(630): p. eabm3682.

11. Bertoglio, D., Verhaeghe, J., Miranda, A., Wyffels, L., Stroobants, S., Mrzljak, L., Khetarpal, V., Skinbjerg, M., Liu, L., Dominguez, C., Munoz-Sanjuan, I., Bard, J., and Staelens, S., *Longitudinal preclinical evaluation of the novel radioligand [11C]CHDI-626 for PET imaging of mutant huntingtin aggregates in Huntington's disease.* Eur J Nucl Med Mol Imaging, 2022. **49**(4): p. 1166-1175.

12. Herrmann, F., Hessmann, M., Schaertl, S., Berg-Rosseburg, K., Brown, C.J., Bursow, G., Chiki, A., Ebneth, A., Gehrmann, M., Hoeschen, N., Hotze, M., Jahn, S., Johnson, P.D., Khetarpal, V., Kiselyov, A., Kottig, K., Ladewig, S., Lashuel, H., Letschert, S., Mills, M.R., Petersen, K., Prime, M.E., Scheich, C., Schmiedel, G., Wityak, J., Liu, L., Dominguez, C., Munoz-Sanjuan, I., and Bard, J.A., *Pharmacological characterization of mutant huntingtin aggregate-directed PET imaging tracer candidates.* Sci Rep, 2021. **11**(1): p. 17977.

13. Liu, L., Johnson, P.D., Prime, M.E., Khetarpal, V., Lee, M.R., Brown, C.J., Chen, X., Clark-Frew, D., Coe, S., Conlon, M., Davis, R., Ensor, S., Esposito, S., Moren, A.F., Gai, X., Green, S., Greenaway, C., Haber, J., Halldin, C., Hayes, S., Herbst, T., Herrmann, F., Hessmann, M., Hsai, M.M., Kotey, A., Mangette, J.E., Mills, M.R., Monteagudo, E., Nag, S., Nibbio, M., Orsatti, L., Schaertl, S., Scheich, C., Sproston, J., Stepanov, V., Varnas, K., Varrone, A., Wityak, J., Mrzljak, L., Munoz-Sanjuan, I., Bard, J.A., and Dominguez, C., *[(11)C]CHDI-626, a PET Tracer Candidate for Imaging Mutant Huntingtin Aggregates with Reduced Binding to AD Pathological Proteins.* J Med Chem, 2021. **64**(16): p. 12003-12021.

14. Bertoglio, D., Weiss, A.R., Liguore, W., Martin, L.D., Hobbs, T., Templon, J., Srinivasan, S., Dominguez, C., Munoz-Sanjuan, I., Khetarpal, V., Verhaeghe, J., Staelens, S., Link, J., Liu, L., Bard, J.A., and McBride, J.L., *In Vivo Cerebral Imaging of Mutant Huntingtin Aggregates Using (11)C-CHDI-180R PET in a Nonhuman Primate Model of Huntington Disease.* J Nucl Med, 2023. **64**(10): p. 1581-1587.

15. Liu, L., Prime, M.E., Lee, M.R., Khetarpal, V., Brown, C.J., Johnson, P.D., Miranda-Azpiazu, P., Chen, X., Clark-Frew, D., Coe, S., Davis, R., Dickie, A., Ebneth, A., Esposito, S., Gadouleau, E., Gai, X., Galan, S., Green, S., Greenaway, C., Giles, P., Halldin, C., Hayes, S., Herbst, T., Herrmann, F., Hessmann, M., Jia, Z., Kiselyov, A., Kotey, A., Krulle, T., Mangette, J.E., Marston, R.W., Menta, S., Mills, M.R., Monteagudo, E., Nag, S., Nibbio, M., Orsatti, L., Schaertl, S., Scheich, C., Sproston, J., Stepanov, V., Svedberg, M., Takano, A., Taylor, M., Thomas, W., Toth, M., Vaidya, D., Vanras, K., Weddell, D., Wigginton, I., Wityak, J., Mrzljak, L., Munoz-Sanjuan, I., Bard, J.A., and Dominguez, C., *Imaging Mutant Huntingtin Aggregates: Development of a Potential PET Ligand.* J Med Chem, 2020. **63**(15): p. 8608-8633.

16. Liu, L., Johnson, P.D., Prime, M.E., Khetarpal, V., Brown, C.J., Anzillotti, L., Bertoglio, D., Chen, X., Coe, S., Davis, R., Dickie, A.P., Esposito, S., Gadouleau, E., Giles, P.R., Greenaway, C., Haber, J., Halldin, C., Haller, S., Hayes, S., Herbst, T., Herrmann, F., Hessmann, M., Hsai, M.M., Khani, Y., Kotey, A., Lembo, A., Mangette, J.E., Marriner, G.A., Marston, R.W., Mills, M.R., Monteagudo, E., Forsberg-Moren, A., Nag, S., Orsatti, L., Sandiego, C., Schaertl, S., Sproston, J., Staelens, S., Tookey, J., Turner, P.A., Vecchi, A., Veneziano, M., Munoz-Sanjuan, I., Bard, J., and Dominguez, C., *Design and Evaluation of [(18)F]CHDI-650 as a Positron Emission Tomography Ligand to Image Mutant Huntingtin Aggregates.* J Med Chem, 2023. **66**(1): p. 641-656.

17. Zajicek, F., Verhaeghe, J., De Lombaerde, S., Van Eetveldt, A., Miranda, A., Munoz-Sanjuan, I., Dominguez, C., Khetarpal, V., Bard, J., Liu, L., Staelens, S., and Bertoglio, D., *Preclinical evaluation of the novel [(18)F]CHDI-650 PET ligand for non-invasive quantification of mutant huntingtin aggregates in Huntington's disease.* Eur J Nucl Med Mol Imaging, 2024. **52**(1): p. 122-133.

18. Delva, A., Koole, M., Serdons, K., Bormans, G., Liu, L., Bard, J., Khetarpal, V., Dominguez, C., Munoz-Sanjuan, I., Wood, A., Skinbjerg, M., Wang, Y., Vandenberghe, W., and Van Laere, K., *Biodistribution and dosimetry in human healthy volunteers of the PET radioligands [(11)C]CHDI-00485180-R and [(11)C]CHDI-00485626, designed for quantification of cerebral aggregated mutant huntingtin.* Eur J Nucl Med Mol Imaging, 2022. **50**(1): p. 48-60.

19. Delva, A., Koole, M., Serdons, K., Bormans, G., Skinbjerg, M., Khetarpal, V.,
Liu, L., Bard, J., Doot, R., Warner, J.H., Sathe, S., Sampaio, C., Wood, A., Van
Laere, K., and Vandenberghe, W., *PET imaging with [(1)(1)C]CHDI-00485180-
R, designed as radioligand for aggregated mutant huntingtin, in people with
Huntington's disease.* Eur J Nucl Med Mol Imaging, 2025.
20. Liu, L., Johnson, P.D., Mills, M.R., Turner, P.A., Prime, M.E., Brown, C.J.,
Kotey, A., Coe, S., Giles, P.R., Lloyd, C., Hayes, S., Marlin, F.J., Rota, F.,
Herrmann, F., Hessmann, M., Schaertl, S., Zajicek, F., De Lombaerde, S.,
Elvas, F., Verhaeghe, J., Staelens, S., Bertoglio, D., Chen, X., Conlon, M.P.,
Davis, R., Ensor, S.F., Haber, J., Haber, K.C., Hsai, M.M., Mangette, J.E.,
Penniman, W.F., Anzillotti, L., Esposito, S., Lembo, A., Orsatti, L., Ventre, D.,
Veneziano, M., Sandiego, C., Haller, S., Marriner, G.A., Munoz-Sanjuan, I.,
Khetarpal, V., Bard, J.A., and Dominguez, C., *Isoindolinone-Based PET
Tracers for Imaging Mutant Huntingtin Aggregates.* J Med Chem, 2025. **68**(14):
p. 14818-14842.
21. Menalled, L.B., Kudwa, A.E., Miller, S., Fitzpatrick, J., Watson-Johnson, J.,
Keating, N., Ruiz, M., Mushlin, R., Alosio, W., McConnell, K., Connor, D.,
Murphy, C., Oakeshott, S., Kwan, M., Beltran, J., Ghavami, A., Brunner, D.,
Park, L.C., Ramboz, S., and Howland, D., *Comprehensive behavioral and
molecular characterization of a new knock-in mouse model of Huntington's
disease: zQ175.* PLoS One, 2012. **7**(12): p. e49838.
22. Heikkinen, T., Lehtimaki, K., Vartiainen, N., Puolivali, J., Hendricks, S.J.,
Glaser, J.R., Bradaia, A., Wadel, K., Touller, C., Kontkanen, O., Yrjanheikki,
J.M., Buisson, B., Howland, D., Beaumont, V., Munoz-Sanjuan, I., and Park,
L.C., *Characterization of neurophysiological and behavioral changes, MRI brain
volumetry and 1H MRS in zQ175 knock-in mouse model of Huntington's
disease.* PLoS One, 2012. **7**(12): p. e50717.
23. Cudalbu, C., McLin, V.A., Lei, H., Duarte, J.M., Rougemont, A.L., Oldani, G.,
Terraz, S., Toso, C., and Gruetter, R., *The C57BL/6J mouse exhibits sporadic
congenital portosystemic shunts.* PLoS One, 2013. **8**(7): p. e69782.
24. Bertoglio, D., Verhaeghe, J., Miranda, A., Kertesz, I., Cybulska, K., Korat, S.,
Wyffels, L., Stroobants, S., Mrzljak, L., Dominguez, C., Liu, L., Skinbjerg, M.,
Munoz-Sanjuan, I., and Staelens, S., *Validation and noninvasive kinetic
modeling of [(11)C]UCB-J PET imaging in mice.* J Cereb Blood Flow Metab,
2020. **40**(6): p. 1351-1362.
25. Miranda, A., Bertoglio, D., Glorie, D., Stroobants, S., Staelens, S., and
Verhaeghe, J., *Validation of a spatially variant resolution model for small animal
brain PET studies.* Biomed Phys Eng Express, 2020. **6**(4): p. 045001.
26. Bertoglio, D., Verhaeghe, J., Kosten, L., Thomae, D., Van der Linden, A.,
Stroobants, S., Wityak, J., Dominguez, C., Mrzljak, L., and Staelens, S., *MR-
based spatial normalization improves [18F]MNI-659 PET regional
quantification and detectability of disease effect in the Q175 mouse model of
Huntington's disease.* PLoS One, 2018. **13**(10): p. e0206613.
27. Johnson, G.A., Badea, A., Brandenburg, J., Cofer, G., Fubara, B., Liu, S., and
Nissanov, J., *Waxholm space: an image-based reference for coordinating
mouse brain research.* Neuroimage, 2010. **53**(2): p. 365-72.
28. Bertoglio, D., Verhaeghe, J., Korat, S., Miranda, A., Wyffels, L., Stroobants, S.,
Mrzljak, L., Dominguez, C., Liu, L., Skinbjerg, M., Munoz-Sanjuan, I., and

Staelens, S., *In vitro and In vivo Assessment of Suitable Reference Region and Kinetic Modelling for the mGluR1 Radioligand [(11)C]ITDM in Mice.* Mol Imaging Biol, 2020. **22**(4): p. 854-863.

29. Bertoglio, D., Zajicek, F., Lombaerde, S., Miranda, A., Stroobants, S., Wang, Y., Dominguez, C., Munoz-Sanjuan, I., Bard, J., Liu, L., Verhaeghe, J., and Staelens, S., *Validation, kinetic modeling, and test-retest reproducibility of [(18)F]SynVesT-1 for PET imaging of synaptic vesicle glycoprotein 2A in mice.* J Cereb Blood Flow Metab, 2022. **42**(10): p. 1867-1878.

30. Verhaeghe, J., Bertoglio, D., Kosten, L., Thomae, D., Verhoye, M., Van Der Linden, A., Wyffels, L., Stroobants, S., Wityak, J., Dominguez, C., Mrzljak, L., and Staelens, S., *Noninvasive Relative Quantification of [(11)C]ABP688 PET Imaging in Mice Versus an Input Function Measured Over an Arteriovenous Shunt.* Front Neurol, 2018. **9**: p. 516.

31. Paxinos, G. and Franklin, K.B.J., *The mouse brain in stereotaxic coordinates: Academic Press.* 2 edition ed. 2003.

32. Marchionini, D.M., De Lombaerde, S., van Rijswijk, J., Zajicek, F., Everix, L., Miranda, A., Aaltonen, M.J., Kluger, C.M., Wild, T., Kakoulidou, A., Gundelach, J., Fieblinger, T., Fentz, J., Rosinski, J., Obenauer, J., Greene, J.R., Liu, L., Munoz-Sanjuan, I., Verhoye, M., Verhaeghe, J., Bard, J., Staelens, S., and Bertoglio, D., *Pharmacodynamic biomarkers responsive to mutant huntingtin lowering in a Huntington's disease mouse model.* Neurobiol Dis, 2025. **209**: p. 106906.

33. Cassinelli Petersen, G., Roytman, M., Chiang, G.C., Li, Y., Gordon, M.L., and Franceschi, A.M., *Overview of tau PET molecular imaging.* Curr Opin Neurol, 2022. **35**(2): p. 230-239.

34. Cicchetti, D., *Guidelines, Criteria, and Rules of Thumb for Evaluating Normed and Standardized Assessment Instruments in Psychology.* Psychological Assessment, 1994. **6**(4): p. 284-290.

35. Weiss, A.R., Liguore, W.A., Brandon, K., Wang, X., Liu, Z., Domire, J.S., Button, D., Srinivasan, S., Kroenke, C.D., and McBride, J.L., *A novel rhesus macaque model of Huntington's disease recapitulates key neuropathological changes along with motor and cognitive decline.* Elife, 2022. **11**.

Tables

Table 1 [^{11}C]CHDI-009R $V_{T\ (IDIF)}$ (Logan) [mL/cm^3] quantitative values in 9-month-old mice

	WT (mean±SD)	HET (mean±SD)	*Post hoc p* value	Δ±SE [%]
Striatum	0.46±0.05	0.90±0.11	<0.0001	94.6±6.2
Motor Cortex	0.47±0.06	0.88±0.13	<0.0001	88.8±6.3
Hippocampus	0.45±0.04	0.83±0.11	<0.0001	85.9±6.4
Thalamus	0.46±0.05	0.94±0.13	<0.0001	105.8±6.2
Cerebellum	0.52±0.06	0.81±0.10	<0.0001	55.5±5.4

WT: *n*=20; HET: *n*=19. SD = standard deviation; Δ = phenotypic difference; SE = standard error

612 ***Table 2 [¹¹C]CHDI-009R test-retest $V_{T\,(IDIF)}$ values and relative and absolute test-retest***

613 ***variability in 9-month-old mice***

	Test $V_{T\,(IDIF)}$ (Logan) [mL/cm³]		Retest $V_{T\,(IDIF)}$ (Logan) [mL/cm³]		rTRV [%]		aTRV [%]		ICC
	WT	HET	WT	HET	WT	HET	WT	HET	
Striatum	0.45±0.04	0.84±0.06	0.46±0.04	0.82±0.10	-0.9±6.7	0.9±9.2	5.1±4.0	7.7±4.2	0.62
Motor Cortex	0.49±0.04	0.79±0.07	0.47±0.03	0.84±0.12	4.3±8.3	-4.8±6.2	7.0±5.8	6.3± 4.5	0.79
Hippocampus	0.43±0.04	0.77±0.06	0.44±0.04	0.66±0.04	-1.8± 7.4	1.2 ± 8.3	6.1± 4.2	6.9± 4.1	0.71
Thalamus	0.44±0.03	0.84±0.07	0.46±0.04	0.81±0.08	-3.1±8.5	-0.1±7.8	6.2± 6.2	5.2±5.4	0.72
Cerebellum	0.49±0.04	0.72±0.05	0.50±0.05	0.67±0.04	-2.1± 8.3	-0.1± 8.0	6.2± 5.5	6.3± 4.2	0.63

615 *rTRV = relative test-retest variability; aTRV = absolute test-retest variability; ICC = Intraclass*

616 *correlation coefficient. Data is given as mean±SD. WT: n=8; HET: n=8*

618 ***Table 3 [^{11}C]CHDI-009R $V_{T\ (IDIF)}$ (Logan) [mL/cm^3] quantitative values in 3-month-old mice***

	WT (mean±SD)	HET (mean±SD)	*Post hoc* *p* value	Δ±SE [%]
Striatum	0.46±0.03	0.52±0.04	0.0002	12.3±2.9
Motor Cortex	0.47±0.04	0.50±0.06	0.3989	5.0±2.8
Hippocampus	0.44±0.03	0.48±0.04	0.0194	8.9±3.0
Thalamus	0.46±0.03	0.53±0.05	<0.0001	16.3±2.9
Cerebellum	0.50±0.03	0.54±0.04	0.0104	8.3±2.7

625 *WT: n=18; HET: n=19. SD = standard deviation; Δ = phenotypic difference; SE = standard error*

627 **Figures**

629 *Fig. 1* Overview of the experimental design. Number and age of each experimental group for the

630 different experimental setups. HET = heterozygous; n = number; WT = wildtype

Fig. 2 *[11C]CHDI-009R brain and plasma analysis and IDIF of WT and HET zQ175DN mice. The*

intact radioligand in percentage for brain (a) and plasma (b) over time. c, Plasma-to-whole-blood

ratio over time in both genotypes. d, Averaged IDIF SUV curves over 90 min scan time for both

genotypes. d: Data are represented as mean±SEM. a-c: n=2-5 per genotype and timepoint; d: WT:

n=20; HET: n=19

Fig. 3 *[¹¹C]CHDI-009R brain kinetic modeling and $V_{T\,(IDIF)}$ time stability with Logan. **a**, Exemplary*

*striatal SUV TAC curves of WT and HET animals fitted to 2TCM. **b**, Pearson's correlation of striatal*

$V_{T\,(IDIF)}$ values with 2TCM and Logan for verification with simple linear regression. $V_{T\,(IDIF)}$ time

*stability of Logan in WT (**c**) and HET (**d**) mice. $V_{T\,(IDIF)}$ values were normalized to the 90 min scan*

acquisition time. b-d: WT: n=20; HET: n=19

Fig. 4 *[¹¹C]CHDI-009R mHTT aggregate quantification in 9-month-old mice. a, Group-averaged*

parametric maps for WT and HET zQ175DN mice overlaid onto MRI template. b, [¹¹C]CHDI-009R

$V_{T(IDIF)}$ estimations in WT and HET mice. Two-way ANOVA with post hoc Bonferroni for multiple

*comparisons. ****$p<0.0001$. WT: n=20; HET: n=19. STR = Striatum; MC = Motor Cortex; HC =*

Hippocampus; THAL = Thalamus; CB = Cerebellum

Fig. 5 *[11C]CHDI-009R test-retest variability analysis in 9-month-old mice. a, Group-averaged*

parametric maps for WT and HET zQ175DN mice at test and retest overlaid onto MRI template.

b, Absolute test-retest variability analysis in percentage. c, Bland-Altman plot to display the level

of agreement between test and retest. WT: n=8; HET: n=8. STR = Striatum; MC = Motor Cortex;

HC = Hippocampus; THAL = Thalamus; CB = Cerebellum

Fig. 6 *mHTT quantification in 3-month-old mice. a, Group-averaged parametric maps for WT and*

HET zQ175DN mice overlaid onto MRI template. b, [¹¹C]CHDI-009R $V_{T\,(IDIF)}$ estimations in WT and

HET mice. c, [³H]CHDI-009R autoradiography of the whole brain in the same mice. ns=not

*significant; * p<0.05; *** p<0.001; **** p<0.0001. a and b: WT: n=18; HET: n=19; c: WT: n=20;*

HET: n=22. STR = Striatum; MC = Motor Cortex; HC = Hippocampus; THAL = Thalamus; CB =

Cerebellum

Supplemental material

Fig. S1 [¹¹C]CHDI-009R radiometabolite analysis in WT and HET zQ175DN mice. *a, Decay-*

corrected reconstructed radiochromatogram for the pure tracer. Decay-corrected reconstructed

radiochromatograms for all time points in plasma (b and c) and brain (d and e) for WT (b and d)

and HET mice (c and e). Data is given as the mean. n=2-3 per time point and genotype. CPM =

counts per minute

Fig. S2 Mass dose effect in striatum of HET zQ175DN mice at 9 months. *Spearman's*

correlation of injected masses (µg/kg) and striatal Logan $V_{T\ (IDIF)}$ values. The sigmoidal interpolation

(sigmoidal, 4PL, X is concentration) estimates the influence of higher injected masses on the

$V_{T\ (IDIF)}$ value. The upper mass dose limit was recommended to be 1.7µg/kg. Data is represented

as individual points. n=30

Table S1 Scan parameters during the PET imaging study in WT and HET zQ175DN mice

for the test-retest study at 9 months

	Test	Retest	*p* value	Test	Retest	*p* value
	WT (*n* = 8)		*p* value	HET (*n* = 8)		*p* value
Injected activity (MBq)	6.3±1.6	7.1±2.5	0.1507	5.3±1.7	5.7±2.2	0.5148
A_m (GBq/µmol) at EOS	112.6±28.6	122.8±26.9	0.3682	107.9±30.6	111.2±24.8	0.9375
Weight (g)	34.8±3.4	33.9±2.8	0.0006	30.8±2.6	30.1±2.2	0.0023
Age (d)	257.5±7.5	261.0±10.4	<0.0001	261.1±9.5	267.0±13.9	0.0143
Injected mass (µg/kg)	1.31±0.15	1.27±0.17	0.6242	1.27±0.14	1.22±0.14	0.9275

686 Data is given as mean±SD. Unpaired t-test for comparison of genotypes. A_m = molar activity;

687 EOS = end of synthesis

689 ***Table S2 Animal and dose parameters for the PET imaging study in WT and HET zQ175DN***

690 ***mice for the evaluation at 9 months***

	WT ($n = 20$)	HET ($n = 19$)	*p* value
Injected activity (MBq)	3.8±0.6	4.5 ± 1.2	0.0062
A_m (GBq/*µmol*) at EOS	71.2±16.1	73.2 ± 16.8	0.7122
Weight (g)	29.7±1.8	34.8 ± 2.8	<0.0001
Age (d)	263.3±4.3	263.2 ± 2.7	0.9221
Injected mass (*µg/kg*)	1.38 ± 0.20	1.35 ± 0.28	0.9337

698 Data is given as mean±SD unless stated otherwise. Unpaired t-test for comparison of genotypes.

699 A_m = molar activity; EOS = end of synthesis

701 ***Table S3 Animal and dose parameters for the PET imaging study in WT and HET zQ175DN***

702 ***mice for the evaluation at 3 months***

	WT (n = 18)	HET (n = 19)	p value
Injected activity (MBq)	5.0±0.9	4.8±0.9	0.4608
A_m (GBq/μmol) at EOS	100.8±16.6	99.1±17.9	0.7676
Weight (g)	28.4±2.2	27.7±1.8	0.2975
Age (d)	108.0±3.3	106.6±4.1	0.2523
Injected mass (μg/kg)	1.31 ± 0.14	1.34 ± 0.19	0.6164

709 Data is given as mean±SD. Unpaired t-test for comparison of genotypes. A_m = molar activity;

710 EOS = end of synthesis

712 ***Table S4 [¹¹C]CHDI-009R power analysis of 3- and 9-month-old mice***

age [M]	Brain region	striatal Δ±SE [%]	Cohen's *d*	10%	20%	30%	40%	50%	100%
3	Striatum	12.3±2.9	1.70	551	139	62	36	23	6
	Motor Cortex	5.0±2.8	0.59	4947	1238	540	310	199	37
	Hippocampus	8.9±3.0	1.13	1238	310	136	78	51	11
	Thalamus	16.3±2.9	1.70	632	159	70	41	26	6
	Cerebellum	8.3±2.7	1.13	1238	310	136	78	51	11
9	Striatum	95.8±6.1	5.15	78	21	10	6	4	2
	Motor Cortex	88.8±6.3	4.05	125	32	15	9	6	2
	Hippocampus	85.9±6.4	4.59	105	27	13	8	5	2
	Thalamus	105.8±6.2	4.87	92	24	11	7	5	2
	Cerebellum	55.5±5.4	3.52	148	38	17	10	7	3

714 *One-tailed power analysis with $\alpha=0.05$ and $\beta=80\%$. Animal numbers needed for significance*

715 *between treated and non-treated HET groups in a mHTT-lowering study. 100% equals difference*

716 *between non-treated HET and WT groups. 3 months: WT: n=18; HET: n=19; 9 months: WT:*

717 *n=20; HET: n=19. M = months; Δ = phenotypic difference; SE = standard error*

718