Epimutations driven by RNAi or heterochromatin evoke antifungal drug resistance in human fungal pathogens

Ye-Eun Son[1] and Joseph Heitman[1*]

[1] *Department of Molecular Genetics and Microbiology, Duke University Medical Center, Durham, NC 27710, USA*

[*]**Corresponding Author:**

213 Research Drive, Durham, NC 27710, USA

Tel: 919-684-2824, Fax: 919-684-2790

Email: heitm001@duke.edu

Keywords: *Mucor*, Epigenetic resistance, RNA silencing, Histone modification, *In vivo* stability

Abstract

Antimicrobial resistance is a global health threat emerging through microbe adaptation, driven by genetic variation, genome plasticity or epigenetic regulation. This study investigates how the *Mucor circinelloides* species complex adapts to the antifungal natural product FK506. In *Mucor bainieri*, most resistant strains exhibit unstable phenotypes without genetic changes. Approximately ~50% of FK506-resistant isolates acquire resistance via RNAi-dependent epimutation, where small interfering RNAs (siRNAs) silence *fkbA* transcription. The remaining isolates undergo heterochromatin-mediated silencing via H3K9 methylation and siRNAs spreading, repressing *fkbA* and neighboring genes. One isolate retained only heterochromatin marks without detectable siRNAs. A similar mechanism operates in *Mucor atramentarius*, where FK506 resistance is mediated by ectopic heterochromatin associated with siRNA. Strikingly, heterochromatin-based epimutation inheritance remains stable following *in vivo* infection. These findings reveal that antifungal resistance can arise through distinct, heritable epigenetic pathways involving RNAi, heterochromatin, or both highlighting adaptive strategies employed by ubiquitous eukaryotic microbial pathogens infecting humans.

Introduction

Fungi are ubiquitous eukaryotic organisms that occupy diverse ecological niches. While many fungi contribute to food production, pharmaceutical development, and biotechnology, others are pathogens capable of infecting a wide range of hosts including humans [1]. The global burden of fungal infections has risen dramatically, now accounting for an estimated two million deaths annually [2, 3]. Despite this growing threat, therapeutic options remain limited due to a restricted antifungal arsenal, inadequate diagnostics, and an alarming rise in antifungal resistance. The widespread use of antifungal agents in agriculture and healthcare, combined with environmental pressures such as climate change, has intensified selective pressure on fungal populations, driving the emergence of fungal antimicrobial resistance (fAMR) [4-7].

Fungi employ both genetic and epigenetic strategies to adapt to antifungal stress [8, 9]. Genetic changes include point mutations (missense/nonsense) and insertions and deletions (indels) in drug target genes (e.g., *erg11*, *cyp51*, *fks1*), as well as large-scale chromosomal alterations such as copy number variation and aneuploidy. These changes confer resistance in several human fungal pathogens including *Candida albicans*, *Cryptococcus neoformans*, and *Aspergillus fumigatus* [10, 11]. In contrast, epigenetic modifications enable non-Mendelian inheritance of resistance without altering the underlying DNA sequence. These include processes such as DNA methylation, RNA interference (RNAi), and ectopic heterochromatin [12].

In *Mucor circinelloides*, RNAi-dependent silencing of *fkbA*—which encodes FKBP12, the cellular target of FK506 (tacrolimus)—confers transient FK506 resistance that reverts in the absence of drug pressure [13]. RNAi-mediated silencing of *pyrF* or

pyrG prevents conversion of 5-fluoroorotic acid (5-FOA) into its toxic form, leading to

transient 5-FOA resistance in *Mucor circinelloides* [14]. In addition, heterochromatin-

driven gene silencing has been implicated in stress adaptation. In

Schizosaccharomyces pombe, exposure to caffeine, an environmental stressor, induces

the formation of histone 3 lysine 9 dimethylation (H3K9me2)-enriched heterochromatin

islands that silence genes and promote survival without genetic mutation [15].

Mucormycosis is a life-threatening infection caused by Mucorales species

including *Mucor* and *Rhizopus* [16]. Spores enter the host via inhalation, traumatic injury,

or the rhino-orbital-cerebral route, often resulting in rapid dissemination, especially in

immunocompromised individuals with conditions such as diabetes mellitus, malignancy,

malnutrition, organ transplantation, or trauma [17, 18]. The mortality rate of

mucormycosis ranges from 50 to 90%, and incidence surged globally during the

COVID-19 pandemic, with over 3,000 deaths reported in India alone [19-21].

Recognizing the severity of mucormycosis, the World Health Organization (WHO)

recently designated Mucorales as a high-priority fungal group in the Fungal Priority

Pathogens List to drive research and therapeutic development [9, 22]. Nonetheless,

both intrinsic and acquired antifungal resistance in *Mucor* spp. continue to undermine

treatment efficacy, underscoring the urgent need to define resistance mechanisms.

The *Mucor circinelloides* complex (MCC) comprises at least 16 phylogenetic

species (PS), most of which are predominantly isolated from clinical cases and primarily

involved in infections within Mucorales [23]. Prior studies in *M. lusitanicus* and *M.*

circinelloides have identified both Mendelian and epigenetic mechanisms of FK506

resistance, including RNAi-mediated silencing of *fkbA* in the absence of mutations [13,

24]. However, whether such epigenetic regulation is conserved across other MCC

species—or whether alternative mechanisms exist—remains unclear. In this study, we

demonstrate that antifungal FK506 resistance in MCC arises through diverse genetic

and epigenetic mechanisms, including RNAi-mediated and heterochromatin-mediated

silencing. Notably, RNAi and heterochromatin functionally interact to modulate gene

expression and confer FK506 resistance in some epimutants. Additionally, we

demonstrate that FK506 resistance mediated by H3K9me2 heterochromatin is stably

inherited after *in vivo* infection, providing new insights into the durability and clinical

relevance of epigenetic antifungal resistance.

Results

Screening of *Mucor circinelloides* species complex (MCC) reveals isolates resistant to FK506

Previous studies reported the phylogenetic classification of the MCC and divided the group into 16 phylogenetic species (PS) based on a polyphasic approach that included multi-locus sequence analyses, as well as morphological and physiological characterization [23]. RNAi-mediated epigenetic gene silencing has been shown to play a key role in FK506 resistance in *Mucor lusitanicus* (PS10) and *M. circinelloides* (PS14 and PS15) [13, 24]. In this study, we screened multiple species to isolate strains resistant to FK506, a calcineurin inhibitor known to affect the hyphae-to-yeast transition, a process that has been linked to virulence in *Mucor* [25], and rapamycin which inhibits TOR and blocks nutrient-stimulated growth [26]. Following the strategy outlined in **Fig 1A**, spores were collected and point-inoculated on YPD media with or without FK506, and the emergence of resistant colonies was monitored. Resistant colonies were observed in *Mucor janssenii* (PS1), *Mucor bainieri* (PS3), *Mucor atramentarius* (PS6), and *Mucor pseudocircinelloides* (PS13), as well as in the previously reported *M. lusitanicus* (PS10) and *M. circinelloides* (PS14 and PS15) (**S1 Fig**).

To investigate the underlying mechanisms of resistance, we selected FK506-resistant isolates from *M. janssenii, M. bainieri,* and *M. atramentarius* for further analysis **(Fig 1B)**. The *fkbA* gene, three genes encoding calcineurin catalytic A (*cnaA~C*) subunits, and the calcineurin regulatory B subunit (*cnbR*) gene were amplified and analyzed by Sanger sequencing. As summarized (**Table 1**), four out of ten FK506-resistant isolates from *M. janssenii* exhibited large deletions spanning the *fkbA* locus.

An additional five isolates carried large insertions (~4 to 5 kb) in the *fkbA* promoter,

nonsense mutations within *fkbA* exons, or point mutations at splice sites. In one isolate,

a point mutation in the calcineurin A subunit gene (*cnaA*) was associated with FK506

resistance but retained rapamycin sensitivity. Thus, in *M. janssenii*, all ten isolates

exhibited Mendelian mutations. In contrast, analysis of ten FK506-resistant isolates from

M. bainieri revealed only one isolate with a nonsense mutation in *fkbA*, while the

remaining nine exhibited no detectable sequence changes in *fkbA*, or the genes

encoding calcineurin A or B subunits, suggesting a possible epigenetic regulation

(**Table 1**). Similarly, in *M. atramentarius*, among ten FK506-resistant isolates, three

harbored nonsense mutations in *fkbA*, one contained a synonymous mutation, two had

splice-site point mutations, one had a start-loss mutation, and one had a point mutation

in calcineurin B subunit (*cnbR*) (**Table 1**). Notably, two isolates showed no genetic

alterations in any of the sequenced genes.

These results indicate that all ten FK506-resistant isolates from *M. janssenii* are

Mendelian mutants. In *M. bainieri*, one isolate exhibited a Mendelian mutation, while the

remaining nine were identified as candidate epimutants. In *M. atramentarius*, eight

isolates carried Mendelian mutations, and two were classified as candidate epimutants.

FK506-resistant epimutants display reversible phenotypes characterized by

silenced *fkbA* transcription and FKBP12 protein expression

For FK506-resistant isolates where no DNA sequence alterations were detected

in the FK506 target gene *fkbA* or the calcineurin genes, we hypothesized that resistance

might be regulated by epigenetic mechanisms. Because epimutations are typically

associated with reversible phenotypic plasticity [14], we tested whether FK506

resistance would be lost after successive subculturing on drug-free medium. In fact,

FK506 resistance was progressively lost following serial passages in the absence of

selection pressure (**Fig 2A, 2D** and **Table 2**). Resistant strains exhibited a hyphal

morphology and were capable of sporulation on FK506-containing media, whereas both

wild-type (WT) and revertant strains displayed a yeast-like morphology when exposed

to FK506 treatment (**S2A, S2D Fig**). Additionally, in *M. bainieri*, phenotypic reversion

occurred after an average of 15.5 passages, whereas in *M. atramentarius*, reversion

required approximately 30 passages (**Fig 2A** and **2D**). Next, considering that epigenetic

alterations inhibit target gene expression without changes in the underlying nucleotide

sequence, we assayed *fkbA* transcript and FKBP12 protein levels. As shown in **Fig 2B**

and **2E**, the *fkbA* mRNA levels in all FK506-resistant strains were significantly reduced

compared to WT and were restored in all reverted strains. Consistently, FKBP12 protein

was undetectable in resistant strains and was restored in reverted epimutant isolates

(**Fig 2C** and **2F**). These results suggest that drug-resistant strains lacking genetic

changes represent epimutants, and that resistance reversion varies both between and

within species.

Small RNAs confer FK506 resistance in *M. bainieri*

To determine which epigenetic mechanisms are involved in FK506 resistance in

M. bainieri, we first evaluated the potential role of DNA methylation. Whole-genome

sequencing data revealed the absence of canonical DNA 5-methylcytosine

methyltransferases (DNMT1, DIM-2, RID-1, DNMT4, and DNMT5), consistent with a

previous study [27], suggesting that DNA methylation is not involved in FK506

resistance.

We next examined the involvement of RNAi by performing sRNA sequencing. As

shown in **Fig 3A**, small antisense RNAs mapping to the *fkbA* locus were detected in

most FK506-resistant epimutants but were absent in WT and revertant strains. The

distribution pattern of sRNAs varied across epimutants: E5–E9 exhibited strong and

specific accumulation of sRNAs at the *fkbA* locus. Additional analysis revealed that

fkbA-targeting sRNAs in E5–E9 averaged approximately 1,000 reads per million (RPM)

(**Fig 3B**). The 5′ nucleotide of these sRNAs was predominantly uracil (~82%; **Fig 3C**),

and their length distribution peaked at 21–24 nucleotides (nt) (**Fig 3D**), consistent with

the general features of Argonaute-loaded sRNAs [13, 28]. These results indicate that

E5–E9 epimutants are RNAi-dependent epimutants, in which FK506 resistance is

mediated by *fkbA*-targeting antisense sRNAs that silence *fkbA* gene expression.

In contrast to E5–E9, which exhibited abundant siRNAs mapping to *fkbA*, isolates

E1–E4 lacked detectable *fkbA*-targeting siRNAs (**Fig 3A** and **Fig 3B**). However, E1–E3

displayed siRNAs mapping to neighboring genes (**S3 Fig**), with a strong 5′ uridine bias

and predominant lengths of 21–24 nt. Strikingly, E4 lacked detectable siRNAs at both

fkbA and its neighboring loci. To determine whether sRNAs spreading at *fkbA*-adjacent

loci was associated with transcriptional repression, we performed reverse transcription

quantitative PCR (RT-qPCR). As shown in **S3O Fig**, E1–E4 exhibited significant

downregulation of genes flanking *fkbA*. However, no significant changes were observed

in the expression of *fkbA*-neighboring genes in E7–E9. Meanwhile several genes (e.g.,

PS3_009596, *PS3_009598*) showed significantly altered expression in E5–E6, but these alterations were not associated with sRNA abundance.

Heterochromatin marks, along with RNAi, contribute to epigenetic resistance in *M. bainieri*

Small RNA spreading has been closely linked to heterochromatin formation and epigenetic gene silencing. Small RNAs targeting specific loci can initiate or reinforce histone modifications, particularly H3K9 methylation [15, 29]. To determine whether histone-mediated epigenetic regulation contributes to FK506 resistance in *M. bainieri*, we conducted chromatin immunoprecipitation (ChIP) to assess the abundance of histone H3 lysine 9 dimethylation (H3K9me2) in the following epimutants: E1–E3, which exhibit sRNA spreading; E4, which lacks detectable sRNAs but shows transcriptional repression of *fkbA* and neighboring genes; and E8, a canonical RNAi-dependent *fkbA* epimutant. ChIP-qPCR was conducted across the *fkbA* locus, targeting the 5′ UTR, two exonic regions, and the 3′ UTR. H3K9me2 enrichment at the positive control region, which was enriched with small RNAs, was comparable across all strains (**Fig 4A**). H3K9me2 enrichment at *fkbA* locus was negligible in WT, E8, and its revertant E8P28. In contrast, significant enrichment of H3K9me2 was observed in epimutants E1–E4 and was lost upon reversion, suggesting that H3K9me2 is associated with unstable antifungal drug resistance.

To further investigate heterochromatin-based epimutational gene silencing, we performed H3K9me2 ChIP-sequencing on the epimutant strains E3, E4, and E8. All strains exhibited similar patterns and levels of H3K9me2 and sRNA at positive control loci, supporting the specificity of the assay (**Fig 4B**). However, in E3 and E4, H3K9me2

signals extended beyond the *fkbA* locus into neighboring regions, suggesting

heterochromatin spreading (**Fig 4C**). Notably, while E3 showed both H3K9me2 and

sRNA spreading, E4 displayed H3K9me2 enrichment without detectable sRNAs. In

contrast, the RNAi-dependent epimutant E8 exhibited strong sRNA accumulation

specifically at the *fkbA* locus but lacked H3K9me2 enrichment. Taken together, these

findings reveal three distinct epigenetic strategies underlying FK506 resistance in *M.*

bainieri as summarized in **Table 3**: RNAi-associated heterochromatin-dependent

mechanisms (Group 1), heterochromatin-dependent only (Group 2), and RNAi-

dependent only (Group 3). Interestingly, Group 1 epimutants required fewer passages

(average 6.67) for phenotypic reversion compared to epimutants in Group 2 (16.00) or

Group 3 (18.80), suggesting that dual-layered epigenetic regulation may promote a

more readily reversible resistance state.

Next, we sought to determine whether gene silencing in these epimutants is

regulated at the pre- or post-transcriptional level. Generally, heterochromatin formation

compacts chromatin structure, preventing RNA polymerase II (RNAPII) from accessing

DNA and thereby blocking transcription initiation, which is classified as pre-

transcriptional regulation [30]. In contrast, RNAi pathways typically involve sRNA that

target and degrade transcripts after transcription, representing post-transcriptional

regulation [31]. To evaluate the transcriptional status of the *fkbA* locus, we performed

RNAPII ChIP-qPCR assays. As shown in **Fig 4D**, RNAPII occupancy at a positive

control region, *actA*, was comparable across all strains, while occupancy at the negative

control region, which was enriched for sRNAs and marked by H3K9me2, was negligible

in all strains. In contrast, the 5′-UTR, exon, and adjacent regions of *fkbA,* where

heterochromatin spreading was observed, was significantly reduced in E3 and E4

compared to WT, the RNAi-dependent epimutant E8, and revertant strains. These

findings indicate that gene silencing in Group 1 (RNAi-associated heterochromatin-

dependent mechanisms) and Group 2 (heterochromatin-dependent only) occurs at the

pre-transcriptional level, whereas silencing in Group 3 (RNAi-dependent only) operates

through post-transcriptional regulation.

Heterochromatin mediates FK506 resistance in *M. atramentarius*

We next assessed whether RNAi- and heterochromatin-based resistance

mechanisms identified in *M. bainieri* also operate in *M. atramentarius*, another member

of the *Mucor circinelloides* species complex. To evaluate the potential involvement of

RNAi, we performed sRNA sequencing. As shown in **Fig 5A**, antisense sRNA spreading

was observed across the *fkbA* locus and its flanking regions. Although sRNA

abundance at *fkbA* itself was too low (**Fig 5B**), the sRNAs exhibited a strong 5′ uridine

bias (**Fig 5C**) and a length distribution peaking at 21–24 nt (**Fig 5D**), consistent with

canonical RNAi-derived sRNAs.

Given that the pattern of sRNA spreading resembled that observed in

heterochromatin-associated epimutants in *M. bainieri* (**S4 Fig**), we next examined

histone modification. ChIP-qPCR revealed significant H3K9me2 enrichment at the *fkbA*

locus in the epimutants (**Fig 5E**). While H3K9me2 enrichment at the positive control

locus, which is enriched with small RNAs, was comparable across all strains, expanded

H3K9me2 signals at *fkbA* and its neighboring genes, accompanied by sRNA spreading,

were strikingly observed exclusively in the epimutant (**Fig 5F** and **5G**). Consistently, RT-

qPCR revealed significantly reduced expression of *fkbA*-neighboring genes in the epimutants compared to WT and revertant strains (**S5 Fig**). Furthermore, consistent with observations in *M. bainieri*, we observed a significant decrease in RNAPII enrichment at the 5′-UTR, exon, and neighboring regions of *fkbA* in the *M. atramentarius* epimutant, indicating that heterochromatin-mediated gene silencing in *M. atramentarius* is associated with pre-transcriptional regulation (**Fig 5H**). Collectively, these results support the conservation of heterochromatin-mediated FK506 resistance in *M. atramentarius*, reinforcing its role as a general epigenetic mechanism across the *Mucor circinelloides* complex.

Antifungal resistance mediated by heterochromatin is stably inherited following *in vivo* passage

Previous studies reported that RNAi-mediated antifungal resistance can be stably transmitted following *in vivo* passage [32]. To investigate whether heterochromatin-mediated resistance also persists through cell division and stress during host infection, we first examined the maximum growth temperature of *M. bainieri* and *M. atramentarius*. *M. atramentarius* was able to grow at 37 °C, suggesting potential pathogenicity, whereas *M. bainieri* failed to grow under the same condition (**Fig 6A**). To further assess pathogenicity of *M. atramentarius*, we infected fungal spores (1×10^6) via retro-orbital injection in a murine model. Mice infected with *M. circinelloides*, a known pathogen causing mucormycosis, exhibited a significant reduction in survival (**Fig 6B**). Notably, infection with *M. atramentarius* also led to a marked decrease in survival, demonstrating its pathogenic potential.

Based on these findings, we infected immunosuppressed mice with equal

number of spores from WT, FK506-resistant epimutant (E1), and revertant (E1P32)

strains of *M. atramentarius*. On day 4 post-infection, brain, liver, spleen, and lung

tissues were harvested to assess fungal dissemination. Fungal colonization levels were

comparable across strains, with no significant differences observed (**S6A Fig**).

Subsequently, when fungal colonies recovered from each organ were tested on FK506-

containing media, resistance was retained in colonies derived from epimutant-infected

mice (over 80% across organs) in contrast to those derived from WT and revertant

strains (**Fig 6C**). Interestingly, a few colonies recovered from the lungs of WT- and

revertant-infected mice also exhibited FK506 resistance. Given that heterochromatin

mediates FK506 resistance in epimutants, we next asked whether this epigenetic state

persists following *in vivo* passage. To address this, we performed ChIP-qPCR on fungal

colonies recovered from four organs. FK506-resistant epimutants exhibited robust

H3K9me2 enrichment not only at exonic regions but also across the *fkbA* promoter and

terminator (**Fig 6D** and **S6B**). By contrast, H3K9me2 levels were minimal or

undetectable in WT and revertant strains. These findings indicate that heterochromatin-

mediated antifungal resistance is largely stable during host infection, likely maintained

through mitotic inheritance of H3K9 methylation marks.

Discussion

Fungal antimicrobial resistance (fAMR) represents an urgent and growing global health threat, jeopardizing human and animal health, agriculture, and food security [7]. Although antifungal drugs are widely used to mitigate this burden, the remarkable adaptability of fungi to chemical stressors has rendered them increasingly difficult to control. Paradoxically, dual antifungal therapies intended to improve treatment outcomes may instead accelerate the evolution of multidrug-resistant strains [33-35]. A deeper understanding of the mechanisms driving fAMR is critical for developing effective interventions across clinical and environmental settings.

FK506 targets FKBP12, a conserved protein that when bound to FK506 inhibits the calcineurin pathway—an essential regulator of fungal growth, morphogenesis, stress responses, and virulence [36]. In *M. circinelloides*, FK506 enforces a morphological shift from hyphae to yeast-like cells, thereby attenuating pathogenicity [25]. Here, we show that FK506 regulates morphology across the *Mucor circinelloides* complex (**S1 Fig**), and that resistance arises via both Mendelian and non-Mendelian mechanisms (**Fig 1** and **Table1**). Our findings expand current models of antifungal resistance by revealing that RNAi- or heterochromatin-mediated epimutations represent alternative, and sometimes independent, pathways. These mechanisms may function separately or synergistically to silence *fkbA*, operating through either post-transcriptional or pre-transcriptional regulation to confer FK506 resistance, and resulting in resistance phenotypes with varying degrees of stability, specificity, and reversibility (**Fig 7**).

The interplay between RNAi and histone methylation raises key questions about the sequence and causality of these silencing events. Preexisting heterochromatin can

initiate RNAi, as seen in centromeres and transposable elements in *S. pombe* and

Arabidopsis thaliana, where transcripts from heterochromatic loci are processed into

siRNAs that reinforce silencing [37]. In *S. pombe*, chaperone proteins like Hsp90

stabilize the RNA-induced transcriptional silencing (RITS) complex, which recruits Clr4

to nascent transcripts, facilitating heterochromatin formation [38]. Conversely,

euchromatic loci may sporadically generate double-stranded RNA, initiating RNAi and

leading to heterochromatin deposition via a positive feedback loop[39]. However, in *M.*

circinelloides, RNAi and heterochromatin formation have been shown to function

independently in repressing transposable elements, contributing to genome stability [27].

Interestingly, in our study, we identified FK506-resistant epimutants with varying

combinations of siRNA accumulation and H3K9me2 enrichment at the *fkbA* locus (**Fig**

4). These findings support two non-exclusive models: (1) RNAi-driven initiation of

heterochromatin, or (2) heterochromatin-facilitated RNAi. Given that the *fkbA* gene is

located in a euchromatic region, the RNAi-first model appears more plausible, although

chromatin-initiated RNAi cannot be excluded. Future studies analyzing pathway-specific

mutants will be critical to determine the directionality and mechanistic coupling of these

epigenetic processes.

Maintaining epigenetic silencing is energetically demanding. The RNAi pathway

requires ATP for the cleavage of double-stranded RNA by Dicer and the unwinding of

siRNA duplexes [40]. Histone methylation also relies on ATP-driven production of S-

adenosylmethionine (SAM), a cofactor of histone methyltransferase, and ATP-

dependent chromatin remodeling, which is essential for stable heterochromatin

maintenance [41, 42]. These processes reflect a substantial metabolic cost to the cell.

Notably, epimutants governed by both RNAi and heterochromatin-dependent silencing

exhibited faster phenotypic reversion than those regulated by a single mechanism

(**Table 3**). This reduced stability may stem from the cumulative energetic burden of

maintaining multiple ATP-intensive pathways. The associated metabolic cost could

impose selective pressure favoring reversion, thereby restoring cellular fitness.

Alternatively, redundancy or incomplete reinforcement between the two silencing

systems may inherently destabilize the dual-layered epimutation, rendering it more

prone to reversion than resistance maintained by either pathway alone.

Torres-Garcia et al. demonstrated that external stressors such as caffeine induce

phenotypic plasticity via Clr4/H3K9me "read-write" systems in *S. pombe*, underscoring

the adaptive function of heterochromatin in non-pathogenic fungi [15]. Our findings

extend this concept by providing direct evidence that H3K9 methylation contributes to

FK506 resistance in dimorphic *Mucor* species. Notably, this mechanism operates in

pathogenic potential *M. bainieri* and *M. atramentarius* species, suggesting that

heterochromatin-mediated resistance is conserved across diverse fungal taxa, spanning

both unicellular yeasts and multinuclear hyphal species. While RNAi-dependent

antifungal resistance has been shown to be rapidly induced and reversed following *in*

vivo passage in a murine model [32], the *in vivo* stability of heterochromatin-dependent

epimutants has not been previously characterized. Here, we demonstrate that a

heterochromatin-dependent epimutant remains stable *in vivo* (**Fig 6**). The persistence,

even in the absence of antifungal pressure, raises the possibility that epigenetically

resistant strains could silently disseminate in clinical settings, complicating diagnostics

and treatment.

In nature, diverse microorganisms, including bacteria, viruses, and fungi, interact

continuously, forming complex ecological networks [43-46]. Soil systems are nutrient-

rich environments that support a vast diversity of microorganisms, which in turn

contribute to essential ecological functions such as nutrient cycling and organic matter

decomposition [47, 48]. *Streptomyces* is one of the most widely distributed and

ecologically significant genera within the phylum Actinobacteria, renowned for its

remarkable ability to produce a wide variety of bioactive secondary metabolites,

including antifungal compounds[49]. Here, we used FK506, a natural antifungal

compound that is produced by several *Streptomyces* species, such as *Streptomyces*

hygroscopicus and *Streptomyces tsukubaensis* [50, 51]. Mucorales represent a major

fungal order of soil-dwelling saprophytic fungi [52], and thus the ecological coexistence

of *Streptomyces* and *Mucor* in soil raises the possibility that natural occurrence of the

antifungal natural product FK506 may impose a selective pressure on fungal

populations. This pressure may promote the emergence of epigenetic resistance, even

in the absence of anthropogenic drug use, by enabling fungi to flexibly modulate gene

expression in response to environmental antifungal stress, thereby enhancing their

survival and competitive fitness in chemically dynamic ecosystems.

In summary, our study extends RNAi-dependent gene silencing to additional

Mucor species and uncovers that heterochromatin-mediated silencing also drives

antifungal resistance. These findings underscore the remarkable epigenetic plasticity of

fungal epigenomes and highlight adaptive strategies that enable reversible, yet durable,

resistance. Targeting RNAi and chromatin-modifying pathways may thus represent a

promising therapeutic strategy against fAMR.

Material and methods

Fungal strains, media and growth conditions

Mucor strains studied here are described in **S1 Table**. *Mucor* strains were grown on liquid or solid yeast extract peptone dextrose (YPD; BD, Difco, Franklin Lakes, NJ, USA) at room temperature. To isolate FK506-/rapamycin-resistant strains, 10^5 spores were inoculated onto solid YPD, supplemented with antifungal drug 1 μg/mL of FK506 (Astellas Pharma, Chuo, Tokyo, Japan) and 100 ng/ml of rapamycin (API Chem, Hangzhou, Zhejiang, Chian) and incubated for 3, 5, 7 days [13]. To test the stability of FK506-resistant strains, spores of the resistant strains were collected and inoculated in liquid YPD without any drugs overnight [14]. After several rounds of subculturing, the passaged cells were spotted onto YPD or YPD with FK506 to confirm phenotypic plasticity.

Sanger sequencing analysis

Mucor strains were cultured on YPD medium for three days and gDNA was extracted from with the Tissue & Cell Lysis Buffer (Lucigen, Middleton, WI, USA) and MPC Protein Precipitation Reagent (Lucigen). The promoter, open reading frame, and terminator regions of the *fkbA* gene were amplified by polymerase chain reaction (PCR) using primers listed in **S2 Table**. The purified PCR products were sequenced and analyzed against the reference *fkbA* sequence.

RNA extraction and quantitative PCR (qPCR) analysis

Total RNA was extracted from Mucor cells cultured on YPD, with or without

FK506, at room temperature. After overnight lyophilization, each sample was

homogenized in a FastPrep bead-beater (MP Biomedicals, Santa Ana, CA, USA) with

0.2 mL of zirconia-silica beads (BioSpec Products, Bartlesville, OK, USA). Total RNA

was then purified with the miRNeasy Mini Kit (Qiagen, Hilden, Germany) [53]. The

isolated RNA was treated with DNase and subsequently employed for cDNA synthesis

with the Maxima H Minus cDNA Synthesis Master Mix (ThermoFisher, Waltham, MA,

USA). Real-time PCR was performed with PowerUp SYBR Green Master Mix

(ThermoFisher) on a QuantStudio 3 qPCR system (ThermoFisher) in triplicate. The

primers for qPCR are listed in **S2 Table**. Gene expression was normalized to *actA*,

relative fold expression was calculated using $2^{-\Delta\Delta Ct}$ method.

Small RNA (sRNA) sequencing

sRNA libraries were constructed with the QIAseq miRNA Library Kit (Qiagen) and

sequenced on an NovaSeq X platform (Illumina) by 75-bp single end. Raw reads were

trimmed with adaptor sequence (5′-GTTCAGAGTTCTACAGTCCGACGATC-3′ and 5′-

AACTGTAGGCACCATCAA-3′) utilizing BBDuk. Filtered reads were then mapped to the

reference genomes (*M. bainieri* CBS293.63; PRJNA1277536 or *M. atramentarius*

CBS202.28; PRJNA1277544) using BBMap [54]. The resulting SAM files were

converted to BAM files, and antisense reads were sorted using SAMtools (v1.21) [55].

BAM files were converted to TDF files using IGV tools with a window size of 20 bp and

the mean window function. The data was subsequently visualized in the Integrative

Genome Viewer (IGV v2.16.0) [56]. The Y-axis reflects the average read coverage per

20 bp window. Peak calling was performed with MACS3 using a fold-enrichment range

between 5 and 50. sRNA-seq experiments were performed with one biological sample

per strain.

Western blot

Each *Mucor* strain was cultivated in YPD, with or without FK506, at room

temperature for three days. After overnight lyophilization, each sample was

homogenized using a FastPrep bead-beater (MP Biomedicals) with 0.2 mL of zirconia-

silica beads (BioSpec Products). Pierce RIPA buffer (ThermoFisher) containing a

protease inhibitor tablet (Roche) was then added, and each sample was further

disrupted with a FastPrep bead-beater. The samples were incubated on ice for at least

30 min. Cell extracts were centrifuged at 14,000 RPM, 4°C for 10 min, and the

supernatants were quantified with the Pierce BCA Protein Assay Kit (ThermoFisher).

Next, 20 µg of each sample was mixed with Novex Tris-Glycine SDS Sample Buffer

(Invitrogen, Waltham, MA, USA) and boiled at 85°C for 5 min. After briefly cooling, the

samples were separated by SDS-PAGE, transferred to a PVDF membrane, and

incubated overnight at 4°C with an anti-FKBP12 antibody (Invitrogen) or an anti-GAPDH

antibody (Proteintech, Rosemont, IL, USA). Following incubation with a horseradish

peroxidase-conjugated anti-rabbit antibody, FKBP12 and GAPDH expression were

detected on a ChemiDoc MP imaging system (Bio-Rad, Hercules, CA, USA).

Chromatin immunoprecipitation (ChIP)-qPCR

A total of 10□ spores from each *Mucor* strain were inoculated in liquid YPD

medium with or without FK506 and incubated overnight at 25°C. The samples were then

cross-linked with 1% formaldehyde (ThermoFisher), followed by addition of glycine

(Sigma-Aldrich, St. Louis, MO, USA) at a final concentration of 0.125M. After washing

and freezing using liquid nitrogen, each sample was ground with a pestle and

suspended in ChIP lysis buffer (50 mM HEPES, pH=7.5, 150 mM NaCl, 1 mM EDTA,

pH=8, 1% Triton X-100, and 0.1% sodium deoxycholate) containing a protease inhibitor

tablet (Roche). Disrupted cells were then sonicated with a Bioruptor (Diagenode) for 40

cycles, with 30s on and 30s off and clarified by centrifuge for 5 min. The input samples

were stored at

-80°C, while the IP samples were incubated overnight at 4°C with Dynabeads Protein

A/G (Invitrogen) pre-conjugated with the anti-histone H3K9 dimethyl antibody (Abcam,

Cambridge, UK) or the anti-RNA polymerase II antibody (Active Motif, Carlsbad, CA).

Crosslinking between protein and DNA in both input and IP samples was reversed with

the MAGnify Chromatin Immunoprecipitation System (Invitrogen), and DNA quality was

analyzed using a Qubit (ThermoFisher). qPCR was performed in triplicate with primers

described in **S2 Table**. H3K9 methylation enrichment was calculated by normalizing the

IP DNA to the input DNA with the ΔCt method and further adjusted to the *actA* gene

with the ΔΔCt method, followed by $2^{-\Delta\Delta Ct}$ to determine relative enrichment [57, 58]. RNA

polymerase II enrichment was similarly calculated by normalizing the IP DNA to the

input DNA (ΔCt), followed by normalization to an intronic region enriched for siRNAs

and marked by H3K9me2 (ΔΔCt). Relative enrichment was determined with the $2^{-\Delta\Delta Ct}$

method.

ChIP sequencing

479 Input and IP libraries were constructed with KAPA HyperPrep (Roche) and

480 sequenced on a NovaSeq X platform (Illumina) by 150-bp paired end. Raw FASTA files

481 were trimmed with Trimmomatic (v0.39) and filtered reads were mapped to the

482 reference genomes (*M. bainieri* CBS293.63; PRJNA1277536 or *M. atramentarius*

483 CBS202.28; PRJNA1277544) using Bowtie2 (v2.5.4) [59, 60]. The resulting SAM files

484 were converted to BAM files using SAMtools (v1.21) [55]. Broad ChIP-peak calling was

485 analyzed with MACS3 (v3.0.3) with the parameters --mfold 10 1000 and --extsize 147,

486 and enrichment was analyzed by generating coverage files with BamCoverage (v3.5.6),

487 followed by counts per million (CPM) normalization [61, 62]. Processed BAM files were

488 converted to TDF files IGV tools with a window size of 20 bp and the mean window

489 function and subsequently visualized with IGV (v2.16.0) [63]. ChIP-seq experiments

490 were performed with one biological sample per strain.

492 ***In vivo* infection**

493 Six-week-old male BALB/cAnNCrl mice, 6 to 7 weeks old btained from Charles

494 River laboratory (Wilmington, MA, USA), were analyzed in this study. Five mice were

495 infected with each *Mucor* strain. Two days prior to infection, 200 mg/kg of

496 cyclophosphamide (Sigma-Aldrich), an immunosuppressant, was administered

497 intraperitoneally. Each mouse was infected with 1×10^6 spores (in 50 µL PBS) via retro-

498 orbital injection after anesthetizing the mice with isoflurane for 2.5 min. After infection,

499 the mice were monitored twice daily and administered cyclophosphamide at 200 mg/kg

500 body weight every four days. To assess pathogenicity, weight loss was monitored, and

501 any animal experiencing ≥ 20% loss of body weight was humanely euthanized. All

502 animal experiments in this study were approved by the Duke University Institutional

503   Animal Care and Use Committee (IACUC) (protocol #A224-23-11). Animal care and

504   experiments were conducted according to IACUC ethical guidelines.

Strains recovered after *in vivo* passage

To assess fungal burden and isolate *Mucor* strains from organs, five mice per *Mucor* strain were intravenously infected. At the humane endpoint, the brain, lung, liver, and spleen were dissected. Whole organs were homogenized in 0.8 mL of PBS with a bead beater for 2 min, then diluted and plated on YPD supplemented with 50 µg/mL chloramphenicol. After a two-day incubation at room temperature, fungal colonies were picked and patched onto YPD plates with or without FK506 to assess the stability of FK506 resistance following *in vivo* passage. The stability rate was determined by calculating the proportion of resistant colonies on YPD supplemented with FK506 relative to the total colonies patched on YPD [32].

Quantification and statistical analyses

Statistical differences between the control and target strains were evaluated using ordinary one-way ANOVA using Tukey's Honestly Significant Difference (HSD) test with using GraphPad Prism (v10.1.1). Kaplan-Meier survival curves were generated and analyzed with the log-rank test in R (v4.3.2) to estimate survival probability. Data are reported as mean ± standard deviation. In all graphs, p values or asterisks were listed indicating statistical significance ($^*p < 0.05$; $^{**}p < 0.01$; $^{***}p < 0.001$; $^{****}p < 0.0001$).

Funding Statement

This study was supported by the NIH/National Institute of Allergy and Infectious Diseases (R01 AI170543-03, R01-AI039115-27, and R01-AI050113-20) awarded to J.H.. J.H. is co-director and fellow of the CIFAR program Fungal Kingdom: Threats & Opportunities.

Acknowledgments

We thank Dr. Vikas Yadav for his insightful advice on bioinformatics analyses as well as for his scientific discussions and critical reading. We are grateful to Dr. Carlos Pérez-Arques for his contributions on initial whole genome sequencing, assembly and annotation, as well as for his assistance with *in vivo* training and critical reading, and to Dr. María Isabel Navarro-Mendoza for providing guidance on *in vivo* training. We also thank Anna Floyd Averette for her invaluable support and assistance with *in vivo* assays; Ziyan Xu for preparing DNA for Illumina whole genome sequencing; Dr. Jason E. Stajich (University of California) for providing whole-genome sequence datasets. We appreciate critical reading by Dr. Sheng Sun, Dr. Jun Huang, and Dr. Grit Walther (Hans Knöll Institute).

Author Contributions

Conceptualization: Ye-Eun Son, Joseph Heitman

Methodology: Ye-Eun Son

Software: Ye-Eun Son

Validation: Ye-Eun Son

Investigation: Ye-Eun Son

Visualization: Ye-Eun Son

Resources: Joseph Heitman

Writing – Original Draft: Ye-Eun Son

Writing – Review & Editing: Ye-Eun Son, Joseph Heitman

Supervision: Joseph Heitman

Funding Acquisition: Joseph Heitman

Competing interests

The authors declare no conflicts of interest.

Data Availability

Small RNA sequencing and ChIP sequencing raw data are available in the NCBI

SRA under the project accession numbers PRJNA1242486 (*M. bainieri* CBS293.63)

and PRJNA1242487 (*M. atramentarius* CBS202.28).

References

1.   Case NT, Gurr SJ, Fisher MC, Blehert DS, Boone C, Casadevall A, et al. Fungal impacts on Earth's ecosystems. Nature. 2025;638(8049):49-57. Epub 20250205. doi: 10.1038/s41586-024-08419-4. PubMed PMID: 39910383.

2.   McDermott A. Drug-resistant fungi on the rise. Proc Natl Acad Sci U S A. 2022;119(48):e2217948119. Epub 20221121. doi: 10.1073/pnas.2217948119. PubMed PMID: 36413501; PubMed Central PMCID: PMCPMC9860136.

3.   Fisher MC, Denning DW. The WHO fungal priority pathogens list as a game-changer. Nat Rev Microbiol. 2023;21(4):211-2. doi: 10.1038/s41579-023-00861-x. PubMed PMID: 36747091; PubMed Central PMCID: PMCPMC9901396.

4.   Lockhart SR, Chowdhary A, Gold JAW. The rapid emergence of antifungal-resistant human-pathogenic fungi. Nat Rev Microbiol. 2023;21(12):818-32. Epub 20230830. doi: 10.1038/s41579-023-00960-9. PubMed PMID: 37648790; PubMed Central PMCID: PMCPMC10859884.

5.   Perlin DS, Rautemaa-Richardson R, Alastruey-Izquierdo A. The global problem of antifungal resistance: prevalence, mechanisms, and management. Lancet Infect Dis. 2017;17(12):e383-e92. Epub 20170731. doi: 10.1016/S1473-3099(17)30316-X. PubMed PMID: 28774698.

6.   Nnadi NE, Carter DA. Climate change and the emergence of fungal pathogens. PLoS Pathog. 2021;17(4):e1009503. Epub 20210429. doi: 10.1371/journal.ppat.1009503. PubMed PMID: 33914854; PubMed Central PMCID: PMCPMC8084208.

7.   Fisher MC, Burnett F, Chandler C, Gow NAR, Gurr S, Hart A, et al. A one health roadmap towards understanding and mitigating emerging Fungal Antimicrobial Resistance: fAMR. NPJ Antimicrob Resist. 2024;2(1):36. Epub 20241107. doi: 10.1038/s44259-024-00055-2. PubMed PMID: 39524479; PubMed Central PMCID: PMCPMC11543597.

8.   Lee Y, Robbins N, Cowen LE. Molecular mechanisms governing antifungal drug resistance. NPJ Antimicrob Resist. 2023;1(1):5. Epub 20230717. doi: 10.1038/s44259-023-00007-2. PubMed PMID: 38686214; PubMed Central PMCID: PMCPMC11057204.

9.   Al Aboud NM, Tupper C, Jialal I. Genetics, Epigenetic Mechanism. StatPearls. Treasure Island (FL)2025.

10.   Flowers SA, Colon B, Whaley SG, Schuler MA, Rogers PD. Contribution of clinically derived mutations in *ERG11* to azole resistance in *Candida albicans*. Antimicrob Agents Chemother. 2015;59(1):450-60. Epub 20141110. doi: 10.1128/AAC.03470-14. PubMed PMID: 25385095; PubMed Central PMCID: PMCPMC4291385.

11.   Lewis RE, Liao G, Hou J, Prince RA, Kontoyiannis DP. Comparative in vivo dose-dependent activity of caspofungin and anidulafungin against echinocandin-susceptible and -resistant *Aspergillus fumigatus*. J Antimicrob Chemother. 2011;66(6):1324-31. Epub 20110411. doi: 10.1093/jac/dkr142. PubMed PMID: 21486855.

12.   Chang Z, Yadav V, Lee SC, Heitman J. Epigenetic mechanisms of drug resistance in fungi. Fungal Genet Biol. 2019;132:103253. Epub 20190717. doi:

10.1016/j.fgb.2019.103253. PubMed PMID: 31325489; PubMed Central PMCID: PMCPMC6858951.

13. Calo S, Shertz-Wall C, Lee SC, Bastidas RJ, Nicolas FE, Granek JA, et al. Antifungal drug resistance evoked via RNAi-dependent epimutations. Nature. 2014;513(7519):555-8. Epub 20140727. doi: 10.1038/nature13575. PubMed PMID: 25079329; PubMed Central PMCID: PMCPMC4177005.

14. Chang Z, Billmyre RB, Lee SC, Heitman J. Broad antifungal resistance mediated by RNAi-dependent epimutation in the basal human fungal pathogen *Mucor circinelloides*. PLoS Genet. 2019;15(2):e1007957. Epub 20190211. doi: 10.1371/journal.pgen.1007957. PubMed PMID: 30742617; PubMed Central PMCID: PMCPMC6386414.

15. Torres-Garcia S, Yaseen I, Shukla M, Audergon P, White SA, Pidoux AL, et al. Epigenetic gene silencing by heterochromatin primes fungal resistance. Nature. 2020;585(7825):453-8. Epub 20200909. doi: 10.1038/s41586-020-2706-x. PubMed PMID: 32908306; PubMed Central PMCID: PMCPMC7116710.

16. Hernández JL, Buckley CJ. Mucormycosis. StatPearls. Treasure Island (FL)2025.

17. Binder U, Maurer E, Lass-Florl C. Mucormycosis--from the pathogens to the disease. Clin Microbiol Infect. 2014;20 Suppl 6:60-6. Epub 20140224. doi: 10.1111/1469-0691.12566. PubMed PMID: 24476149.

18. Lax C, Nicolas FE, Navarro E, Garre V. Molecular mechanisms that govern infection and antifungal resistance in Mucorales. Microbiol Mol Biol Rev. 2024;88(1):e0018822. Epub 20240306. doi: 10.1128/mmbr.00188-22. PubMed PMID: 38445820; PubMed Central PMCID: PMCPMC10966947.

19. Cornely OA, Alastruey-Izquierdo A, Arenz D, Chen SCA, Dannaoui E, Hochhegger B, et al. Global guideline for the diagnosis and management of mucormycosis: an initiative of the European Confederation of Medical Mycology in cooperation with the Mycoses Study Group Education and Research Consortium. Lancet Infect Dis. 2019;19(12):e405-e21. Epub 20191105. doi: 10.1016/S1473-3099(19)30312-3. PubMed PMID: 31699664; PubMed Central PMCID: PMCPMC8559573.

20. Roden MM, Zaoutis TE, Buchanan WL, Knudsen TA, Sarkisova TA, Schaufele RL, et al. Epidemiology and outcome of zygomycosis: a review of 929 reported cases. Clin Infect Dis. 2005;41(5):634-53. Epub 20050729. doi: 10.1086/432579. PubMed PMID: 16080086.

21. Devnath P, Dhama K, Tareq AM, Emran TB. Mucormycosis coinfection in the context of global COVID-19 outbreak: A fatal addition to the pandemic spectrum. Int J Surg. 2021;92:106031. Epub 20210724. doi: 10.1016/j.ijsu.2021.106031. PubMed PMID: 34311126; PubMed Central PMCID: PMCPMC8302826.

22. WHO. WHO fungal priority pathogens list to guide research, development and public health action2022.

23. Wagner L, Stielow JB, de Hoog GS, Bensch K, Schwartze VU, Voigt K, et al. A new species concept for the clinically relevant *Mucor circinelloides* complex. Persoonia. 2020;44:67-97. Epub 20190829. doi: 10.3767/persoonia.2020.44.03. PubMed PMID: 33116336; PubMed Central PMCID: PMCPMC7567969.

24.   Perez-Arques C, Navarro-Mendoza MI, Xu Z, Walther G, Heitman J. RNAi
epimutations conferring antifungal drug resistance are inheritable. bioRxiv. 2024. Epub
20241118. doi: 10.1101/2024.10.15.618540. PubMed PMID: 39463932; PubMed
Central PMCID: PMCPMC11507787.
25.   Lee SC, Li A, Calo S, Heitman J. Calcineurin plays key roles in the dimorphic
transition and virulence of the human pathogenic zygomycete *Mucor circinelloides*.
PLoS Pathog. 2013;9(9):e1003625. Epub 20130905. doi: 10.1371/journal.ppat.1003625.
PubMed PMID: 24039585; PubMed Central PMCID: PMCPMC3764228.
26.   Bastidas RJ, Shertz CA, Lee SC, Heitman J, Cardenas ME. Rapamycin exerts
antifungal activity in vitro and in vivo against *Mucor circinelloides via* FKBP12-
dependent inhibition of Tor. Eukaryot Cell. 2012;11(3):270-81. Epub 20111230. doi:
10.1128/EC.05284-11. PubMed PMID: 22210828; PubMed Central PMCID:
PMCPMC3294450.
27.   Navarro-Mendoza MI, Perez-Arques C, Heitman J. Heterochromatin and RNAi
act independently to ensure genome stability in Mucorales human fungal pathogens.
Proc Natl Acad Sci U S A. 2023;120(7):e2220475120. Epub 20230206. doi:
10.1073/pnas.2220475120. PubMed PMID: 36745785; PubMed Central PMCID:
PMCPMC9963178.
28.   Mi S, Cai T, Hu Y, Chen Y, Hodges E, Ni F, et al. Sorting of small RNAs into
Arabidopsis argonaute complexes is directed by the 5' terminal nucleotide. Cell.
2008;133(1):116-27. Epub 20080313. doi: 10.1016/j.cell.2008.02.034. PubMed PMID:
18342361; PubMed Central PMCID: PMCPMC2981139.
29.   Bhattacharjee S, Roche B, Martienssen RA. RNA-induced initiation of
transcriptional silencing (RITS) complex structure and function. RNA Biol.
2019;16(9):1133-46. Epub 20190618. doi: 10.1080/15476286.2019.1621624. PubMed
PMID: 31213126; PubMed Central PMCID: PMCPMC6693537.
30.   Allshire RC, Madhani HD. Ten principles of heterochromatin formation and
function. Nat Rev Mol Cell Biol. 2018;19(4):229-44. Epub 20171213. doi:
10.1038/nrm.2017.119. PubMed PMID: 29235574; PubMed Central PMCID:
PMCPMC6822695.
31.   Filipowicz W, Jaskiewicz L, Kolb FA, Pillai RS. Post-transcriptional gene silencing
by siRNAs and miRNAs. Curr Opin Struct Biol. 2005;15(3):331-41. doi:
10.1016/j.sbi.2005.05.006. PubMed PMID: 15925505.
32.   Chang Z, Heitman J. Drug-resistant epimutants exhibit organ-specific stability
and induction during murine infections caused by the human fungal pathogen *Mucor
circinelloides*. mBio. 2019;10(6):e02579-19. Epub 20191105. doi: 10.1128/mBio.02579-
19. PubMed PMID: 31690679; PubMed Central PMCID: PMCPMC6831780.
33.   Rhodes J, Abdolrasouli A, Farrer RA, Cuomo CA, Aanensen DM, Armstrong-
James D, et al. Genomic epidemiology of the UK outbreak of the emerging human
fungal pathogen *Candida auris*. Emerg Microbes Infect. 2018;7(1):43. Epub 20180329.
doi: 10.1038/s41426-018-0045-x. PubMed PMID: 29593275; PubMed Central PMCID:
PMCPMC5874254.
34.   Fraaije B, Atkins S, Hanley S, Macdonald A, Lucas J. The multi-fungicide
resistance status of *Aspergillus fumigatus* populations in arable soils and the wider
European environment. Front Microbiol. 2020;11:599233. Epub 20201215. doi:

10.3389/fmicb.2020.599233. PubMed PMID: 33384673; PubMed Central PMCID:
PMCPMC7770239.
35. Ikuta KS, Mestrovic T, Naghavi M. Global incidence and mortality of severe
fungal disease. Lancet Infect Dis. 2024;24(5):e268. Epub 20240220. doi:
10.1016/S1473-3099(24)00102-6. PubMed PMID: 38395046.
36. Park HS, Lee SC, Cardenas ME, Heitman J. Calcium-calmodulin-calcineurin
signaling: A globally conserved virulence cascade in eukaryotic microbial pathogens.
Cell Host Microbe. 2019;26(4):453-62. doi: 10.1016/j.chom.2019.08.004. PubMed PMID:
31600499; PubMed Central PMCID: PMCPMC6788756.
37. Djupedal I, Ekwall K. Epigenetics: heterochromatin meets RNAi. Cell Res.
2009;19(3):282-95. doi: 10.1038/cr.2009.13. PubMed PMID: 19188930.
38. Okazaki K, Kato H, Iida T, Shinmyozu K, Nakayama JI, Murakami Y, et al. RNAi-
dependent heterochromatin assembly in fission yeast S*chizosaccharomyces pombe*
requires heat-shock molecular chaperones Hsp90 and Mas5. Epigenetics Chromatin.
2018;11(1):26. Epub 20180604. doi: 10.1186/s13072-018-0199-8. PubMed PMID:
29866182; PubMed Central PMCID: PMCPMC5985592.
39. Yu R, Wang X, Moazed D. Epigenetic inheritance mediated by coupling of RNAi
and histone H3K9 methylation. Nature. 2018;558(7711):615-9. Epub 20180620. doi:
10.1038/s41586-018-0239-3. PubMed PMID: 29925950; PubMed Central PMCID:
PMCPMC6312107.
40. Nykanen A, Haley B, Zamore PD. ATP requirements and small interfering RNA
structure in the RNA interference pathway. Cell. 2001;107(3):309-21. doi:
10.1016/s0092-8674(01)00547-5. PubMed PMID: 11701122.
41. Teperino R, Schoonjans K, Auwerx J. Histone methyl transferases and
demethylases; can they link metabolism and transcription? Cell Metab. 2010;12(4):321-
7. doi: 10.1016/j.cmet.2010.09.004. PubMed PMID: 20889125; PubMed Central PMCID:
PMCPMC3642811.
42. Clapier CR, Iwasa J, Cairns BR, Peterson CL. Mechanisms of action and
regulation of ATP-dependent chromatin-remodelling complexes. Nat Rev Mol Cell Biol.
2017;18(7):407-22. Epub 20170517. doi: 10.1038/nrm.2017.26. PubMed PMID:
28512350; PubMed Central PMCID: PMCPMC8127953.
43. Trivedi P, Leach JE, Tringe SG, Sa T, Singh BK. Plant-microbiome interactions:
from community assembly to plant health. Nat Rev Microbiol. 2020;18(11):607-21. Epub
20200812. doi: 10.1038/s41579-020-0412-1. PubMed PMID: 32788714.
44. Faust K, Raes J. Microbial interactions: from networks to models. Nat Rev
Microbiol. 2012;10(8):538-50. Epub 20120716. doi: 10.1038/nrmicro2832. PubMed
PMID: 22796884.
45. Weiland-Brauer N. Friends or Foes-Microbial Interactions in Nature. Biology
(Basel). 2021;10(6). Epub 20210602. doi: 10.3390/biology10060496. PubMed PMID:
34199553; PubMed Central PMCID: PMCPMC8229319.
46. Getzke F, Thiergart T, Hacquard S. Contribution of bacterial-fungal balance to
plant and animal health. Curr Opin Microbiol. 2019;49:66-72. Epub 20191112. doi:
10.1016/j.mib.2019.10.009. PubMed PMID: 31731228.
47. de Menezes AB, Richardson AE, Thrall PH. Linking fungal-bacterial co-
occurrences to soil ecosystem function. Curr Opin Microbiol. 2017;37:135-41. Epub
20170708. doi: 10.1016/j.mib.2017.06.006. PubMed PMID: 28692866.

48.   Baumgardner DJ. Soil-related bacterial and fungal infections. J Am Board Fam
Med. 2012;25(5):734-44. doi: 10.3122/jabfm.2012.05.110226. PubMed PMID:
22956709.
49.   Procopio RE, Silva IR, Martins MK, Azevedo JL, Araujo JM. Antibiotics produced
by *Streptomyces*. Braz J Infect Dis. 2012;16(5):466-71. Epub 20120911. doi:
10.1016/j.bjid.2012.08.014. PubMed PMID: 22975171.
50.   Kino T, Hatanaka H, Hashimoto M, Nishiyama M, Goto T, Okuhara M, et al. FK-
506, a novel immunosuppressant isolated from a *Streptomyces*. I. Fermentation,
isolation, and physico-chemical and biological characteristics. J Antibiot (Tokyo).
1987;40(9):1249-55. doi: 10.7164/antibiotics.40.1249. PubMed PMID: 2445721.
51.   Arndt C, Cruz MC, Cardenas ME, Heitman J. Secretion of FK506/FK520 and
rapamycin by Streptomyces inhibits the growth of competing *Saccharomyces cerevisiae*
and *Cryptococcus neoformans*. Microbiology (Reading). 1999;145 (Pt 8):1989-2000.
doi: 10.1099/13500872-145-8-1989. PubMed PMID: 10463165.
52.   Ziaee A, Zia M, Bayat M, Hashemi J. Identification of Mucorales isolates from soil
using morphological and molecular methods. Curr Med Mycol. 2016;2(1):13-9. doi:
10.18869/acadpub.cmm.2.1.13. PubMed PMID: 28681007; PubMed Central PMCID:
PMCPMC5490292.
53.   Son YE, Yu JH, Park HS. The novel spore-specific regulator SscA controls
Aspergillus conidiogenesis. mBio. 2023;14(5):e0184023. Epub 20230914. doi:
10.1128/mbio.01840-23. PubMed PMID: 37707170; PubMed Central PMCID:
PMCPMC10653911.
54.   Bushnell B. BBMap: A Fast, Accurate, Splice-Aware Aligner. 2014.
55.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The
Sequence Alignment/Map format and SAMtools. Bioinformatics. 2009;25(16):2078-9.
Epub 20090608. doi: 10.1093/bioinformatics/btp352. PubMed PMID: 19505943;
PubMed Central PMCID: PMCPMC2723002.
56.   Priest SJ, Yadav V, Roth C, Dahlmann TA, Kuck U, Magwene PM, et al.
Uncontrolled transposition following RNAi loss causes hypermutation and antifungal
drug resistance in clinical isolates of *Cryptococcus neoformans*. Nat Microbiol.
2022;7(8):1239-51. Epub 20220802. doi: 10.1038/s41564-022-01183-z. PubMed PMID:
35918426; PubMed Central PMCID: PMCPMC10840647.
57.   Park HS, Man Yu Y, Lee MK, Jae Maeng P, Chang Kim S, Yu JH. Velvet-
mediated repression of beta-glucan synthesis in *Aspergillus nidulans* spores. Sci Rep.
2015;5:10199. Epub 20150511. doi: 10.1038/srep10199. PubMed PMID: 25960370;
PubMed Central PMCID: PMCPMC4426670.
58.   Navarro-Mendoza MI, Perez-Arques C, Panchal S, Nicolas FE, Mondo SJ,
Ganguly P, et al. Early diverging fungus *Mucor circinelloides* lacks centromeric histone
CENP-A and displays a mosaic of point and regional centromeres. Curr Biol.
2019;29(22):3791-802 e6. Epub 20191031. doi: 10.1016/j.cub.2019.09.024. PubMed
PMID: 31679929; PubMed Central PMCID: PMCPMC6925572.
59.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics. 2014;30(15):2114-20. Epub 20140401. doi:
10.1093/bioinformatics/btu170. PubMed PMID: 24695404; PubMed Central PMCID:
PMCPMC4103590.

60. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat
Methods. 2012;9(4):357-9. Epub 20120304. doi: 10.1038/nmeth.1923. PubMed PMID:
22388286; PubMed Central PMCID: PMCPMC3322381.
61. Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, et al.
Model-based analysis of ChIP-Seq (MACS). Genome Biol. 2008;9(9):R137. Epub
20080917. doi: 10.1186/gb-2008-9-9-r137. PubMed PMID: 18798982; PubMed Central
PMCID: PMCPMC2592715.
62. Ramirez F, Dundar F, Diehl S, Gruning BA, Manke T. deepTools: a flexible
platform for exploring deep-sequencing data. Nucleic Acids Res. 2014;42(Web Server
issue):W187-91. Epub 20140505. doi: 10.1093/nar/gku365. PubMed PMID: 24799436;
PubMed Central PMCID: PMCPMC4086134.
63. Robinson JT, Thorvaldsdottir H, Turner D, Mesirov JP. igv.js: an embeddable
JavaScript implementation of the Integrative Genomics Viewer (IGV). Bioinformatics.
2023;39(1). doi: 10.1093/bioinformatics/btac830. PubMed PMID: 36562559; PubMed
Central PMCID: PMCPMC9825295.

Figure legends

Fig 1. FK506-resistant strains isolated from the *Mucor circinelloides* complex.

(A) Schematic representation of the procedure followed to obtain FK506- and/or rapamycin-resistant strains from *Mucor circinelloides* complex species. This Figure was generated with BioRender.com. **(B)** Phenotypic analysis of FK506-resistant isolates in *M. janssenii* CBS185.68 (PS1), *M. bainieri* CBS293.63 (PS3), and *M. atramentarius* CBS202.28 (PS6).

WT, wild-type; M, Mendelian mutant; E, Epimutant.

Fig 2. FK506-resistant epimutants exhibit silencing of *fkbA* mRNA and FKBP12 protein expression and are unstable.

(A, D) Phenotypic reversion of FK506-resistant epimutant strains following serial passage without drug selection in *M. bainieri* (A) or *M. atramentarius* (D). Red arrows indicate the points at which epimutants reverted to the WT phenotype. FK506 was utilized at 1 μg/mL and rapamycin at 100 ng/mL. WT, wild-type; P, number of passages. **(B, E)** Quantification of the *fkbA* mRNA expression in *M. bainieri* (B) *or M. atramentarius* (E). Error bars represent mean ± SEM (n = 3). Statistical significance: ****p ≤ 0.0001. **(C, F)** Western blot analysis of FKBP12 protein expression in *M. bainieri* (C) or *M. atramentarius* (F).

PS3: E1–E9, epimutants; E1P4–E9P24, revertants. PS6: E1 and E2, epimutants; E1P32 and E2P28, revertants.

Fig 3. Small RNAs mediate FK506 resistance in *M. bainieri*.

(A) Antisense small RNA coverage mapped across the *fkbA* gene and its neighboring loci. The peaks shown below represent significant enrichment peaks, validated by MACS3. **(B)** Small RNA abundance (RPM: reads per million) mapped to the *fkbA* gene. **(C)** 5′-end nucleotide preference of small RNAs associated with the *fkbA* gene. **(D)** Size distribution of small RNAs mapped to the *fkbA* gene.

WT, wild-type; E1–E9, epimutants; E1P4–E9P24, revertants.

Fig 4. Heterochromatin marks underlie epigenetic FK506 resistance in *M. bainieri*.

(A) ChIP-qPCR analysis of H3K9me2 enrichment at the positive control region (an siRNA-enriched locus) and *fkbA* locus (targeting the 5′-UTR, exonic regions, and 3′-UTR). Error bars represent mean ± SEM (n = 3). Statistical significance: $*p \le 0.05$, $**p \le 0.01$; $****p \le 0.0001$. **(B, C)** ChIP-seq coverage mapped across the positive control loci (B) or *fkbA* and its neighboring loci (C). The peaks shown below represent significant enrichment peaks, validated by MACS3. **(D)** ChIP-qPCR analysis of RNA polymerase II enrichment at the positive control region (*actA*), *fkbA* locus (targeting the 5′-UTR and exonic region), a *fkbA*-neighboring gene, and negative control region (an siRNA-enriched locus marked by H3K9me2). Statistical significance: $****p \le 0.0001$.

WT, wild-type; E1–E8, epimutants; E1P4–E8P28, revertants

Fig 5. Heterochromatin-associated FK506 resistance is conserved in *M. atramentarius*.

(A) Antisense small RNA coverage mapped across the *fkbA* gene and its neighboring loci. The peaks shown below represent significant enrichment peaks, validated by MACS3. **(B)** Small RNA abundance (RPM: reads per million) mapped to the *fkbA* gene. **(C)** 5′-end nucleotide preference of small RNAs associated with the *fkbA* gene. **(D)** Size distribution of small RNAs mapped to the *fkbA* gene. **(E)** ChIP-qPCR analysis of H3K9me2 enrichment at the positive control region (an siRNA-enriched locus) and *fkbA* locus (targeting the 5′-UTR, exonic regions, and 3′-UTR). Error bars represent mean ± SEM (n = 3). Statistical significance: ****$p \leq 0.0001$. **(F, G)** ChIP-seq coverage mapped across the positive control loci (F) or *fkbA* and its neighboring loci (G). The peaks shown below represent significant enrichment peaks, validated by MACS3. **(H)** ChIP-qPCR analysis of RNA polymerase II enrichment at the positive control region (*actA*), *fkbA* locus (targeting the 5′-UTR and exonic region), a *fkbA*-neighboring gene, and negative control region (an siRNA-enriched locus marked by H3K9me2). Statistical significance: ****$p \leq 0.0001$.

WT, wild-type; E1 and E2, epimutants; E1P32 and E2P28, revertants.

Fig 6. Inherited heterochromatin-mediated antifungal resistance persists after *in vivo* passage.

(A) Fungal growth of *M. bainieri* and *M. atramentarius* under different temperature conditions. Microscopic images show enlarged sporangia (scale bar = 20 µm). **(B)** Survival rates of mice infected with *M. atramentarius*. *Mucor circinelloides* and PBS

served as positive and negative controls, respectively. The log-rank test revealed a

statistically significant difference in survival between groups (*M. atramentarius* vs PBS,

p = 0.014). **(C)** Fungal colonies recovered from four different organs following *in vivo*

infection with wild-type (WT), epimutant (E1), and revertant (E1P32) strains. The bar

plot (right) shows the percentage of FK506-resistant colonies from each organ after

passage. **(D)** ChIP-qPCR analysis of H3K9me2 enrichment at the *fkbA* locus following

in vivo passage, targeting two exonic regions. Error bars represent mean ± SEM (n = 3).

Statistical significance: ****p ≤ 0.0001.

WT, wild-type; E1R1–E1R3, FK506-resistant epimutants; E1S1, FK506-sensitive

epimutant; E1P32, revertant.

Fig 7. Schematic overview of epigenetic genetic silencing contributing to

antifungal resistance in *Mucor* species.

fkbA expression enables FKBP12 production and FK506 sensitivity (yeast-like

morphology). Epigenetic silencing of *fkbA* via RNAi or heterochromatin leads to FK506

resistance and hyphal growth. Generated with BioRender.com.

 Table 1. FK506 resistant strains isolated in this study.

Species	Name	Phenotype	*fkbA* mutation	Impact on FKBP12	Calcineurin mutation
PS1 *Mucor janssenii*	M1	FK506-R, Rapa-S	No	No	G1496C in *cnaA* (G347A missense mutation)
	M2	FK506-R, Rapa-R	4,324 bp insertion (Unknown) at 38 bp upstream *fkbA* ORF	-	-
	M3	FK506-R, Rapa-R	Large deletion	Loss	-
	M4	FK506-R, Rapa-R	Large deletion	Loss	-
	M5	FK506-R, Rapa-R	5,052 bp insertion (RC/Helitron_DNA transposons) at 16 bp upstream *fkbA* ORF	-	-
	M6	FK506-R, Rapa-R	C120 1 bp deletion in exon 2 (Frameshift)	Nonsense mutation	-
	M7	FK506-R, Rapa-R	Large deletion	Loss	-
	M8	FK506-R, Rapa-R	G258C in 5' splice site of intron2	-	-
	M9	FK506-R, Rapa-R	Large deletion	Loss	-
	M10	FK506-R, Rapa-R	G258C in 5' splice site of intron2	-	-
PS3 *Mucor bainieri*	E1	FK506-R, Rapa-R	No	No	No
	E2	FK506-R, Rapa-R	No	No	No
	E3	FK506-R, Rapa-R	No	No	No
	E4	FK506-R, Rapa-R	No	No	No
	E5	FK506-R, Rapa-R	No	No	No
	E6	FK506-R, Rapa-R	No	No	No
	M1	FK506-R, Rapa-R	C357 1 bp deletion in exon 3 (Frameshift)	Nonsense mutation	No
	E7	FK506-R, Rapa-R	No	No	No
	E8	FK506-R, Rapa-R	No	No	No
	E9	FK506-R, Rapa-R	No	No	No
PS6 *Mucor atramentarius*	M1	FK506-R, Rapa-R	C404 1 bp deletion in exon 3 (Frameshift)	Nonsense mutation	No
	M2	FK506-R, Rapa-S	No	No	G477T in *cnbR* (V122F missense mutation)
	M3	FK506-R, Rapa-R	G249A in 5' splice site	No	No
	M4	FK506-R, Rapa-R	G249A in 5' splice site	No	No
	M5	FK506-R, Rapa-S	G189T in exon 2	Synonymous mutation	No
	M6	FK506-R, Rapa-R	C404 1 bp deletion in exon 3 (Frameshift)	Nonsense mutation	No
	M7	FK506-R, Rapa-R	T2C in exon 1 (start-loss mutation)	Loss	No
	E1	FK506-R, Rapa-R	No	No	No
	E2	FK506-R, Rapa-R	No	No	No
	M8	FK506-R, Rapa-R	C404 1 bp deletion in exon 3 (Frameshift)	Nonsense mutation	No

882   **Table 2. Epimutant reverted strains analyzed in this study.**

Species	Name	Background	Phenotype	Description
PS3 *Mucor bainieri*	E1P4	E1	FK506-S, Rapa-S	E1 Revertant after passage 4
	E2P4	E2	FK506-S, Rapa-S	E2 Revertant after passage 4
	E3P12	E3	FK506-S, Rapa-S	E3 Revertant after passage 12
	E3P16	E4	FK506-S, Rapa-S	E4 Revertant after passage 16
	E5P12	E5	FK506-S, Rapa-S	E5 Revertant after passage 12
	E6P12	E6	FK506-S, Rapa-S	E6 Revertant after passage 12
	E7P28	E7	FK506-S, Rapa-S	E7 Revertant after passage 28
	E8P28	E8	FK506-S, Rapa-S	E8 Revertant after passage 28
	E9P24	E9	FK506-S, Rapa-S	E9 Revertant after passage 24
PS6 *Mucor atramentarius*	E1P32	E1	FK506-S, Rapa-S	E1 Revertant after passage 32
	E2P28	E2	FK506-S, Rapa-S	E2 Revertant after passage 28

884   **Table 3. Summary of epigenetic mechanisms in *Mucor bainieri*.**

Group	Mechanisms	Epimutant	Revertant	Average passages for revertant
1	RNAi-associated heterochromatin-dependent silencing	E1 E2 E3	E1P4 E2P4 E3P12	6.67
2	Heterochromatin-dependent silencing	E4	E3P16	16
3	Canonical RNAi-dependent silencing	E5 E6 E7 E8 E9	E5P12 E6P12 E7P28 E8P28 E9P24	18.8

A

Incubate at RT for 2 days   Spore suspension   Point-inoculate 10^5 spore with or without drug   Incubate at RT

B

PS1 *Mucor janssenii* CBS185.68   PS3 *Mucor bainieri* CBS293.63   PS6 *Mucor atramentarius* CBS202.28

A PS3 *Mucor bainieri* CBS293.63: Scaffold 802

B Levels of sRNAs in *fkbA* gene

C 5' end nucleotide

D Analysis of the length of antisense sRNAs

M. bainieri
M. atramentarius

M. bainieri
[Epimutant 5–9]

M. bainieri   *M. atramentarius*
[Epimutant 1–3]   [Epimutant 1–2]

M. bainieri
[Epimutant 4]