1 **Epimutations driven by RNAi or heterochromatin evoke transient antimicrobial**

2 **drug resistance in fungi**

4 Ye-Eun Son[1], Carlos Pérez-Arques[1], and Joseph Heitman[1*]

[1] Department of Molecular Genetics and Microbiology, Duke University Medical Center,

Durham, NC 27710, USA

[*] **Corresponding Author:**

213 Research Drive, Durham, NC 27710, USA

Tel: 919-684-2824, Fax: 919-684-2790

Email: heitm001@duke.edu

Keywords: *Mucor*, Epigenetic resistance, RNA silencing, Histone modification, *In vivo*

stability

Abstract

Antimicrobial resistance (AMR) is a global health threat emerging through microbe adaptation, driven by genetic variation, genome plasticity or epigenetic process. In this study, we investigated how the *Mucor circinelloides* species complex adapts to the antifungal natural product FK506, which binds to FKBP12 and inhibits calcineurin-dependent hyphal growth. In *Mucor bainieri*, most FK506-resistant isolates (90%) were found to be unstable and transient, readily reverting to drug sensitivity when passaged without drug, and with no associated DNA mutations. In half of the isolates (50%), FK506-resistance was conferred by RNAi-dependent epimutation in which small interfering RNAs (siRNAs) silenced the *fkbA* encoding FKBP12 post-transcriptionally. In contrast, most of the remaining FK506-resistant isolates (40%) were found to have undergone heterochromatin-mediated silencing via H3K9 dimethylation, transcriptionally repressing *fkbA* and neighboring genes. In these heterochromatic epimutants, only minimal enrichment of siRNA to the *fkbA* locus was observed, but in three of the four examples, siRNA was significantly enriched at a locus distant from *fkbA*. A similar mechanism operates in *Mucor atramentarius*, where FK506 resistance was mediated by ectopic heterochromatin silencing *fkbA* and associated genes with siRNA spreading across the region. Heterochromatin-mediated *fkbA* epimutants exhibited stability during *in vivo* infection, suggesting epimutation could impact pathogenesis. These findings reveal that antifungal resistance arising through distinct, transient epimutation pathways involving RNAi or heterochromatin, highlighting adaptive AMR strategies employed by ubiquitous eukaryotic microbes.

Introduction

Fungi are ubiquitous eukaryotic organisms that occupy diverse ecological niches. While many fungi contribute to food production, pharmaceutical development, and biotechnology, others are pathogens capable of infecting a wide range of hosts including humans [1]. The global burden of fungal infections has risen dramatically, now accounting for an estimated two million deaths annually [2, 3]. Despite this growing threat, therapeutic options remain limited due to a restricted antifungal arsenal, inadequate diagnostics, and an alarming rise in antifungal resistance. The widespread use of antifungal agents in agriculture and healthcare, combined with environmental pressures such as climate change, has intensified selective pressure on fungal populations, driving the emergence of fungal antimicrobial resistance (fAMR) [4-7].

Fungi employ both genetic and epigenetic strategies to adapt to antifungal stress [8, 9]. Genetic changes include point mutations (missense/nonsense) and insertions and deletions (indels) in drug target genes or drug transporter genes or their regulators, as well as large-scale chromosomal alterations such as copy number variation and aneuploidy. These changes confer resistance in several human fungal pathogens including *Candida albicans*, *Cryptococcus neoformans*, *Aspergillus fumigatus*, and *Mucor circinelloides* [10-13]. In contrast, epigenetic modifications enable non-Mendelian inheritance of resistance without altering the underlying DNA sequence. These include processes such as DNA methylation, RNA interference (RNAi), ectopic heterochromatin, and DNA methylation [14].

In *Mucor circinelloides*, RNAi-dependent silencing of *fkbA*—which encodes FKBP12, the cellular target of FK506 (tacrolimus)—confers transient FK506 resistance

that reverts in the absence of drug pressure [15]. RNAi-mediated silencing of *pyrF* or *pyrG*

prevents conversion of 5-fluoroorotic acid (5-FOA) into its toxic form, leading to transient

5-FOA resistance in *M. circinelloides* [16]. In addition, heterochromatin-driven gene

silencing has been implicated in stress adaptation. In *Schizosaccharomyces pombe*,

exposure to caffeine, an environmental stressor, selects for epimutations in which induces

the formation of histone 3 lysine 9 dimethylation (H3K9me2)-enriched heterochromatin

islands silence genes and promote survival without genetic mutation [17].

Mucormycosis is a life-threatening infection caused by Mucorales species

including *Mucor* and *Rhizopus* [18]. Spores enter the host via inhalation, traumatic injury,

or the rhino-orbital-cerebral route, often resulting in rapid dissemination, especially in

immunocompromised individuals with conditions such as diabetes mellitus, malignancy,

malnutrition, organ transplantation, or trauma [19, 20]. The mortality rate of mucormycosis

ranges from 50 to 90%, and incidence surged globally during the COVID-19 pandemic,

with over 3,000 deaths reported in India alone [21-23]. Recognizing the severity of

mucormycosis, the World Health Organization (WHO) recently designated Mucorales as

a high-priority fungal group in the Fungal Priority Pathogens List to drive research and

therapeutic development [9, 24]. Given both the severity and widespread incidence of

mucormycosis, a deeper understanding of intrinsic resistance and the potential for

acquired antifungal resistance is urgently needed to guide proper and efficient therapeutic

interventions.

The *Mucor circinelloides* complex (MCC) comprises at least 16 phylogenetic

species (PS), 7 of which have been isolated from clinical cases within Mucorales [25].

Prior studies in *M. lusitanicus* and *M. circinelloides* have identified both Mendelian and

epigenetic mechanisms of FK506 resistance, including RNAi-mediated silencing of *fkbA*

in the absence of mutations [15, 26-28]. However, whether such epigenetic regulation is

conserved across other MCC species—or whether alternative mechanisms exist—

remains unclear. In this study, we demonstrate that antifungal FK506 resistance in MCC

arises primarily through diverse genetic and epigenetic mechanisms, involving either

RNAi-mediated or heterochromatin-mediated epimutations that silence the *fkbA* gene

encoding the FK506 target FKBP12. Drug resistance conferred by both classes of

epimutation is unstable, and readily reverts following passage in the absence of drug in

contrast to mutations in the *fkbA* gene which are stably inherited even in the absence of

drug. Additionally, we demonstrate that FK506 resistance mediated by H3K9me2

heterochromatin is stably inherited after *in vivo* infection, providing new insights into the

durability and clinical relevance of epigenetic antifungal resistance.

Results

Screening of *Mucor circinelloides* species complex (MCC) reveals isolates resistant to FK506

Previous studies reported the phylogenetic classification of the MCC and divided the group into 16 phylogenetic species (PS) based on a polyphasic approach that included multi-locus sequence analyses, as well as morphological and physiological characterization [25]. RNAi-mediated epigenetic gene silencing has been shown to play a key role in FK506 resistance in *Mucor lusitanicus* (PS10) and *M. circinelloides* (PS14 and PS15) [15, 26]. In this study, we screened multiple species to isolate strains resistant to FK506, a calcineurin inhibitor known to prevent the dimorphic transition from yeast to hyphae, a process linked to virulence of *Mucor* [28], and rapamycin which also binds FKBP12 and the complex then inhibits TOR and blocks nutrient-stimulated growth [29]. Following the strategy outlined in **Fig 1A**, spores were collected and point-inoculated on YPD media with or without FK506, and the emergence of resistant colonies was monitored. Resistant colonies were observed in *Mucor janssenii* (PS1), *Mucor bainieri* (PS3), and *Mucor atramentarius* (PS6), as well as in the previously reported species *M. lusitanicus* (PS10) and *M. circinelloides* (PS14 and PS15) (**S1 Fig**).

To investigate the underlying mechanisms of resistance, we selected FK506-resistant isolates from *M. janssenii, M. bainieri,* and *M. atramentarius* for further analysis **(Fig 1B)**. The *fkbA* gene, three genes encoding calcineurin catalytic A (*cnaA*, *cnaB*, *cnaC*) subunits, and the calcineurin regulatory B subunit (*cnbR*) gene were amplified and analyzed by sequencing. As summarized (**Table 1**), four out of ten FK506-resistant isolates from *M. janssenii* exhibited large deletions spanning the *fkbA* locus. An additional

five isolates carried large insertions (~4 to 5 kb) in the *fkbA* promoter, nonsense mutations

within *fkbA* exons, or point mutations at splice sites. These *fkbA* mutants were cross-

resistant to rapamycin. In one isolate, a point mutation in the calcineurin A subunit gene

(*cnaA*) was associated with FK506 resistance but retained rapamycin susceptibility. Thus,

in *M. janssenii*, all ten isolates exhibited Mendelian mutations. In contrast, analysis of ten

FK506-resistant isolates from *M. bainieri* revealed only one isolate with a nonsense

mutation in *fkbA*, while the remaining nine exhibited no detectable sequence changes in

fkbA, or the genes encoding calcineurin A or B subunits, suggesting a possible epigenetic

regulation (**Table 1**). Similarly, in *M. atramentarius*, among ten FK506-resistant isolates,

three harbored nonsense mutations in *fkbA*, one contained a synonymous mutation, two

had splice-site point mutations, one had a start-loss mutation, and one had a point

mutation in the calcineurin B subunit (*cnbR*) (**Table 1**). Notably, two isolates showed no

genetic alterations in any of the sequenced genes.

In summary, these results indicate that all ten FK506-resistant isolates from *M.*

janssenii are Mendelian mutants. In *M. bainieri*, one isolate exhibited a Mendelian

mutation, while the remaining nine were identified as candidate epimutants. In *M.*

atramentarius, eight isolates carried Mendelian mutations, and two were classified as

candidate epimutants.

FK506-resistant epimutants display reversible phenotypes characterized by

decreased levels of *fkbA* mRNA and FKBP12 protein expression

For FK506-resistant isolates where no DNA sequence alterations were detected

in the FK506 target gene *fkbA* or the calcineurin genes, we hypothesized that resistance

might be regulated by epigenetic mechanisms. Because epimutations are typically

associated with reversible phenotypic plasticity [16], we tested whether FK506 resistance

would be lost after successive subculturing on drug-free medium. In fact, FK506

resistance was progressively lost following serial passages in the absence of selection

pressure (**Figs 2A, 2D** and **Table 2**). Resistant strains exhibited a hyphal morphology

and were capable of sporulation on FK506-containing media, whereas both wild-type (WT)

and revertant strains displayed a yeast-like morphology when exposed to FK506

treatment (**S2 Fig**). Additionally, in *M. bainieri*, phenotypic reversion occurred after an

average of 15.5 passages, whereas in *M. atramentarius*, reversion required

approximately 30 passages (**Figs 2A** and **2D**). Next, considering that epigenetic

alterations inhibit target gene expression without changes in the underlying nucleotide

sequence, we assayed *fkbA* transcript and FKBP12 protein levels. As shown in **Fig 2B**

and **2E**, the *fkbA* mRNA levels in all FK506-resistant strains were significantly reduced

compared to WT and were restored in all reverted strains. Consistently, FKBP12 protein

was undetectable in resistant strains and was restored in reverted epimutant isolates

(**Figs 2C** and **2F**). These results suggest that drug-resistant strains lacking genetic

changes represent epimutants, and that resistance reversion varies both between and

within species.

Small RNAs confer FK506 resistance in *M. bainieri*

To determine which epigenetic mechanisms are involved in FK506 resistance in

M. bainieri, we first evaluated the potential role of 5mC DNA methylation. Whole-genome

sequencing data revealed the absence of canonical DNA 5-methylcytosine

methyltransferases (DNMT1, DIM-2, RID-1, DNMT4, and DNMT5), consistent with a

previous study (**S3A Fig**) [30]. Furthermore, 5mC analysis via Nanopore sequencing in

the two *Mucor* wild types detected no cytosine methylation, while a clear 5mC signal was

detected in a *Cryptococcus neoformans* clinical isolate with the same analysis pipeline

[31], suggesting that this mechanism is not involved in FK506 resistance (**S3B Fig**).

We next examined the involvement of RNAi by performing small RNA (sRNA)

sequencing. As shown in **Fig 3A** and **S4 Fig**, antisense small RNAs mapping to the *fkbA*

locus were detected in five of the nine FK506-resistant epimutants but were absent in WT

and revertant strains. The distribution pattern of sRNAs varied across epimutants. In

epimutants E5–E9, *fkbA*-specific sRNAs strongly accumulated to ~1,000 RPM, whereas

in WT and most revertants no reads were detected, with only a minimal signal (~1 RPM)

observed in E9 revertants (**Fig 3B**). The 5′ nucleotide of these sRNAs was predominantly

uracil (~82%; **Fig 3C**), and their length distribution peaked at 21–24 nucleotides (nt) (**Fig**

3D), consistent with the general features of Argonaute-loaded sRNAs [15, 32]. These

results indicate that the E5–E9 epimutants are RNAi-dependent epimutants, in which

FK506 resistance is mediated by *fkbA*-targeting antisense sRNAs that silence *fkbA* gene

expression.

In contrast to epimutants E5–E9, which exhibited abundant siRNAs mapping to

fkbA, epimutant isolates E1–E4 lacked detectable *fkbA*-targeting siRNAs (~1 RPM or less)

(**Figs 3A** and **Fig 3B**). However, E1–E3 displayed siRNAs mapping to neighboring genes

distant from *fkbA* (**S5 Fig**), with a strong 5′ uridine bias and predominant lengths of 21–

24 nt. For example, at *PS3_001333*, sRNAs reached ~2,595 RPM in epimutants E1-E3

but was largely absent in WT (0.15 RPM) and in revertants (1–1.6 RPM). Strikingly,

195 epimutant E4 lacked detectable siRNAs at both *fkbA* and its neighboring loci. To

196 determine whether epimutation spreading to *fkbA* from adjacent loci was associated with

197 transcriptional repression, we performed reverse transcription quantitative PCR (RT-

198 qPCR). As shown in **S5K Fig**, E1–E4 exhibited significant downregulation of genes

199 flanking *fkbA*. However, no significant changes were observed in the expression of *fkbA*-

200 neighboring genes in epimutants E7–E9. Meanwhile several genes (e.g., *PS3_001332*,

201 *PS3_001331*) showed significantly altered expression in epimutants E5–E6, but these

202 alterations were not associated with sRNA abundance.

204 **Heterochromatin marks, along with RNAi, contribute to epigenetic resistance in *M.***

205 ***bainieri***

206  Small RNA spreading has been closely linked to heterochromatin formation and

207 epigenetic gene silencing. Small RNAs targeting specific loci can initiate or reinforce

208 histone modifications, particularly H3K9 methylation [17, 33]. To determine whether

209 histone-mediated epigenetic regulation contributes to FK506 resistance in *M. bainieri*, we

210 conducted chromatin immunoprecipitation (ChIP) to assess the abundance of histone H3

211 lysine 9 dimethylation (H3K9me2) in the following epimutants: E1–E3, which exhibit sRNA

212 spreading; E4, which lacks detectable sRNAs but shows transcriptional repression of *fkbA*

213 and neighboring genes; and E8, a canonical RNAi-dependent, heterochromatin-

214 independent *fkbA* epimutant. ChIP-qPCR was conducted across the *fkbA* locus, targeting

215 the 5′ UTR, two exonic regions, and the 3′ UTR. H3K9me2 enrichment at the positive

216 control region (an siRNA-enriched locus validated by H3K9me2) was comparable across

217 all strains (**Fig 4A**). H3K9me2 levels at the *fkbA* locus were markedly increased in

epimutants E1–E4, with fold enrichment ranging from ~5-fold to over 100-fold relative to

WT, although the magnitude varied across loci and strains. By contrast, H3K9me2

enrichment in E8 and its revertant E8P28 was negligible and comparable to WT. These

results indicate that histone modification is associated with unstable antifungal drug

resistance in some epimutants (E1-E4) and not others.

To further investigate heterochromatin-based epimutational gene silencing, we

performed H3K9me2 ChIP-sequencing on the epimutant strains E3, E4, and E8. All

strains exhibited similar patterns and levels of H3K9me2 and sRNA at positive control loci,

supporting the specificity of the assay (**Fig 4B**). However, in E3 and E4, H3K9me2 signals

extended across the *fkbA* locus from neighboring regions, suggesting heterochromatin

spreading (**Fig 4C**). Notably, while E3 showed both H3K9me2 and sRNA (distal to *fkbA*),

E4 displayed H3K9me2 enrichment without detectable sRNAs. Notably, these ectopic

heterochromatic regions spanning *fkbA* were absent in the wild type and revertant isolates.

In contrast, the RNAi-dependent epimutant E8 exhibited strong sRNA accumulation

specifically at the *fkbA* locus but lacked H3K9me2 enrichment. Taken together, these

findings reveal three distinct types of epigenetic strategies underlying FK506 resistance

in *M. bainieri* (**Table 3**): RNAi-associated heterochromatin mechanisms (type 1),

heterochromatin-dependent only (type 2), and RNAi-dependent only (type 3).

Interestingly, type 1 epimutants required fewer passages (average of 6.67) for phenotypic

reversion compared to epimutants in type 2 (16.00) or type 3 (18.80), suggesting that

dual-layered epigenetic regulation may promote a more readily reversible resistance state.

Next, we sought to determine whether gene silencing in these epimutants is

regulated at the pre- or post-transcriptional level. Generally, heterochromatin formation

compacts chromatin structure, preventing RNA polymerase II (RNAPII) from accessing

DNA and thereby blocking transcription initiation, which is classified as pre-transcriptional

regulation [34]. In contrast, RNAi pathways typically involve sRNA that target and degrade

transcripts after transcription, representing post-transcriptional regulation [35]. To

evaluate the transcriptional status of the *fkbA* locus, we performed RNAPII ChIP-qPCR

assays. As shown in **Fig 4D**, RNAPII occupancy at a positive control region, *actA*, was

comparable across all of the strains, while occupancy at the negative control region,

which was enriched for sRNAs and marked by H3K9me2, was negligible in all strains. In

contrast, the 5′-UTR, exon, and adjacent regions of *fkbA,* where heterochromatin

spreading was observed, was significantly reduced in E3 and E4 compared to WT, the

RNAi-dependent epimutant E8, and revertant strains. These findings indicate that gene

silencing in type 1 (RNAi-associated heterochromatin mechanisms) and type 2

(heterochromatin-dependent only) occurs at the pre-transcriptional level, whereas

silencing in type 3 (RNAi-dependent only) operates exclusively through post-

transcriptional regulation.

Heterochromatin mediates FK506 resistance in *M. atramentarius*

In *M. atramentarius*, canonical DNA 5-methylcytosine methyltransferases (DNMT1,

DIM-2, RID-1, DNMT4, and DNMT5) were also absent, consistent with a previous study

(**S3A Fig**) [29]. Moreover, 5mC analysis in the wild type detected no cytosine methylation,

suggesting that DNA methylation is not involved in FK506 resistance (**S3B Fig**). We next

assessed whether RNAi- or heterochromatin-based resistance mechanisms identified in

M. bainieri also operate in *M. atramentarius*, another member of the *Mucor circinelloides*

species complex. To evaluate the potential involvement of RNAi, we performed sRNA

sequencing of WT, two FK506-resistant epimutants, and their corresponding revertant

strains. As shown in **Fig 5A**, antisense sRNA spreading was observed across the *fkbA*

locus and its flanking regions. Although sRNA abundance at *fkbA* itself was limited (**S6**

and **S7B Fig**), overall the sRNAs exhibited a strong 5′ uridine bias (**S7C Fig**) and a length

distribution peaking at 21–24 nt (**S7D Fig**), consistent with canonical RNAi-derived

sRNAs. Interestingly, while the spreading pattern resembled that observed in

heterochromatin-associated epimutants of *M. bainieri* (**Figs 3A** and **5A**), *M. bainieri*

accumulated markedly higher levels of sRNAs at genes flanking the *fkbA* locus compared

to *M. atramentarius* (**S4** and **S6 Figs**).

To further determine whether the resistant isolates of *M. atramentarius* also involve

heterochromatin-associated mechanisms, we next examined histone modification. ChIP-

qPCR revealed significant H3K9me2 enrichment at the *fkbA* locus in the epimutants, with

fold enrichment ranging from ~12-fold to over 60-fold relative to WT (**Fig 5B**). While

H3K9me2 enrichment at the GremLINE1 transposon, used as a positive control, was

comparable across all strains, expanded H3K9me2 signals at *fkbA* and its neighboring

genes, accompanied by sRNA spreading, were strikingly observed exclusively in the

epimutant (**Figs 5C** and **5D**). Consistently, RT-qPCR revealed significantly reduced

expression of *fkbA*-neighboring genes in the epimutants compared to WT and revertant

strains (**S8 Fig**). Furthermore, consistent with observations in *M. bainieri*, we detected a

significant decrease in RNAPII enrichment at the 5′-UTR, exon, and neighboring regions

of *fkbA* in the *M. atramentarius* epimutant, indicating that heterochromatin-mediated gene

silencing in *M. atramentarius* is associated with pre-transcriptional regulation (**Fig 5E**).

Collectively, these results support the conservation of heterochromatin-mediated FK506

resistance in *M. atramentarius*, reinforcing its role as a general epigenetic mechanism

across the *Mucor circinelloides* complex.

Antifungal resistance mediated by heterochromatin is stably inherited following *in*

***vivo* infection**

Previous studies reported that RNAi-mediated antifungal resistance can be stably

transmitted following *in vivo* passage [36]. To investigate whether heterochromatin-

mediated resistance also persists through cell division and stress during host infection,

we first examined the maximum growth temperature of *M. bainieri* and *M. atramentarius*

with *M. circinelloides* serving as a positive control for growth. *M. atramentarius* was able

to grow at 37°C, suggesting potential pathogenicity, whereas *M. bainieri* failed to grow at

37°C (**Fig 6A**). This was further confirmed in a murine model, as mice infected with *M.*

atramentarius exhibited significantly reduced survival, comparable to that observed with

M. circinelloides, a known pathogen that causes mucormycosis (**Fig 6B**). Notably,

infection with *M. atramentarius* also led to a marked decrease in survival, demonstrating

its pathogenic potential.

Based on these findings, we infected immunosuppressed mice with equal number

of spores from WT, FK506-resistant epimutant (E1), and revertant (E1P32) strains of *M.*

atramentarius. On day 4 post-infection, brain, liver, spleen, and lung tissues were

harvested to assess fungal dissemination. Fungal burden was comparable across strains,

with no significant differences observed (**S9A Fig**). Subsequently, when fungal colonies

recovered from each organ were tested on FK506-containing media, resistance was

retained in colonies derived from epimutant-infected mice (over 80% across organs) in

contrast to those derived from WT and revertant strains (**Fig 6C**). Interestingly, a few

colonies recovered from the lungs of WT- and revertant-infected mice also exhibited

FK506 resistance. Sequencing of the *fkbA* locus in FK506-resistant isolates recovered

from each organ of WT-, epimutant-, and revertant-infected mice revealed no mutations,

with sequences identical to the wild type, indicating that these isolates represent

epimutants rather than Mendelian mutants (**S9B Fig** and **S3 Table**). Given that

heterochromatin mediates FK506 resistance in epimutants, we next asked whether this

epigenetic state persists following *in vivo* infection. To address this, we performed ChIP-

qPCR on fungal colonies recovered from four organs. FK506-resistant epimutants

exhibited robust H3K9me2 enrichment not only at exonic regions but also across the *fkbA*

promoter and terminator (**Fig 6D** and **S10B Fig**). By contrast, H3K9me2 levels were

minimal or undetectable in WT and revertant strains. These findings indicate that

heterochromatin-mediated antifungal resistance is largely stable during host infection,

likely maintained through mitotic inheritance of H3K9 methylation marks.

To further examine whether epimutant stability persists over longer periods of

infection, we extended our *in vivo* assays by administering a lower dose of

cyclophosphamide (an immunosuppressant) or without cyclophosphamide treatment and

then assessed FK506 resistance and heterochromatin-associated stability after two or

four weeks. Unlike the results observed at 4 days post-infection, fungal burden at 2 weeks

post-infection with lower or no cyclophosphamide, and at 4 weeks post-infection without

cyclophosphamide, was detected only in the liver and spleen, but not in the brain or lung

(**S11A-S11C Figs**). Next, when fungal colonies recovered from the liver and spleen were

tested on FK506-containing medium, resistance was retained in colonies derived from

epimutant-infected mice at frequencies greater than 50% under all conditions tested (**Figs**

7A, 7C, and 7E). With resistant colonies from liver and spleen, we sequenced the *fkbA*

locus, confirming that there were no mutations, with sequences identical to the wild type

(**S11D-S11F Figs** and **S4-S6 Tables**). Furthermore, ChIP-qPCR analysis revealed that

FK506-resistant epimutants exhibited H3K9me2 enrichment at the *fkbA* locus (**Figs 7B,**

7D, and **7F**). Together, these results demonstrate that heterochromatin-mediated

antifungal resistance in *M. atramentarius* is stably maintained during prolonged host

infection, independent of genetic mutations, and is instead sustained by mitotically

inherited H3K9me2 marks.

Discussion

Fungal antimicrobial resistance (fAMR) represents an urgent and growing global health threat, jeopardizing human and animal health, agriculture, and food security [7]. Although antifungal drugs are widely used to mitigate this burden, the remarkable adaptability of fungi to chemical stressors has rendered them increasingly difficult to control. Paradoxically, dual antifungal therapies intended to improve treatment outcomes may instead accelerate the evolution of multidrug-resistant strains [37-39]. A deeper understanding of the mechanisms driving fAMR is critical for developing effective interventions across clinical and environmental settings.

FK506 targets FKBP12, a conserved protein that when bound to FK506 inhibits the calcineurin pathway—an essential regulator of fungal growth, morphogenesis, stress responses, and virulence [40]. In *M. circinelloides*, FK506 enforces a morphological shift from hyphae to yeast cells, thereby attenuating pathogenicity [28]. Here, we show that FK506 regulates morphology across the *Mucor circinelloides* complex (**S1 Fig**), and that resistance arises via both Mendelian and non-Mendelian mechanisms (**Fig 1** and **Table1**). Our findings expand current models of antifungal resistance by revealing that RNAi- or heterochromatin-mediated epimutations represent alternative, and sometimes independent, pathways. These mechanisms may function separately or synergistically to silence *fkbA*, operating through either post-transcriptional or pre-transcriptional regulation to confer FK506 resistance, and resulting in resistance phenotypes with varying degrees of stability, specificity, and reversibility (**Fig 8**).

In nature, diverse microorganisms, including bacteria, viruses, and fungi, interact continuously, forming complex ecological networks [41-44]. Soil systems are nutrient-rich

environments that support a vast diversity of microorganisms, which in turn contribute to essential ecological functions such as nutrient cycling and organic matter decomposition [45, 46]. *Streptomyces* is one of the most widely distributed and ecologically significant genera within the phylum Actinobacteria, renowned for its remarkable ability to produce a wide variety of bioactive secondary metabolites, including antifungal compounds[47]. Here, we studied FK506, a natural antifungal compound that is produced by several *Streptomyces* species, such as *Streptomyces hygroscopicus* and *Streptomyces tsukubaensis* [48, 49]. Mucorales represent a major fungal order of soil-dwelling saprophytic fungi [50], and thus the ecological coexistence of *Streptomyces* and *Mucor* in soil raises the possibility that natural occurrence of the antifungal natural product FK506 may impose a selective pressure on fungal populations. This pressure may promote the emergence of epigenetic resistance, even in the absence of anthropogenic drug use, by enabling fungi to flexibly modulate gene expression in response to environmental antifungal stress. Interestingly, amphotericin B, the primary first-line therapy for mucormycosis, is also a natural product produced by *Streptomyces nodosus* [51]. This raises the possibility that fungi may encounter antifungal pressures in nature, allowing epigenetic resistance mechanisms to evolve as strategies that enhance survival and competitive fitness in chemically dynamic ecosystems.

Our data revealed that the size distribution of sRNAs at the *fkbA* locus differs from those mapped to neighboring genes (**Fig 3, S5** and **S7 Figs**). The predominant sRNAs derived from *fkbA* were 21 and 24 nt, whereas most neighboring genes displayed stronger accumulation at 23–24 nt. Previous work in *Mucor* proposed that different sRNA size classes may serve distinct roles in silencing, with 21-nt sRNAs acting as the predominant

species involved in target mRNA degradation [52]. In this context, the increased abundance of 21 nt sRNAs derived specifically from *fkbA* suggests that FK506-resistant epimutants may preferentially generate the sRNA class most directly linked to efficient target degradation. A similar phenomenon was also reported at the *pyrG* and *pyrF* loci, where sRNAs fell within the 21–24 nt range but showed locus differences. *pyrG* sRNAs were enriched at 21 and 24 nt, whereas *pyrF* sRNAs were predominantly 23–24 nt [16]. Such variation may result from multiple, non-mutually exclusive mechanisms. First, differential usage of Dicer paralogs may contribute. In *M. circinelloides*, DCL2 produces 23–24 nt siRNAs, whereas DCL1 can result in more heterogeneous size distributions including 21 nt [53, 54]. Similarly, in the model plant *Arabidopsis thaliana*, DCL4 primarily generates 21-nt siRNAs, while DCL3 is responsible for 24-nt siRNAs involved in transcriptional silencing [55, 56]. Second, substrate properties of the transcripts, including secondary structure, GC content, and overall architecture, may influence Dicer processing efficiency and cleavage-site preference, thereby generating distinct size profiles [57, 58]. These observations suggest that even within the canonical RNAi framework, locus-specific features may shape distinct sRNA size distributions.

The interplay between RNAi and histone methylation raises key questions about the sequence and causality of these silencing events. Preexisting heterochromatin can initiate RNAi, as seen in centromeres and transposable elements in *S. pombe* and *Arabidopsis thaliana*, where transcripts from heterochromatic loci are processed into siRNAs that reinforce silencing [59]. In *S. pombe*, small RNAs bound to Argonaute form the RNA-induced transcriptional silencing (RITS) complex with the chromodomain protein Chp1 and the bridging protein Tas3. Chaperone proteins such as Hsp90 stabilize RITS

complex, which recruits Clr4 to nascent transcripts, thereby facilitating heterochromatin

formation [59, 60]. Stc1 associates with Ago1 and recruits the chromatin-modifying CLRC

complex (Clr4-Rik1-Cul4-Dos1-Dos2), underscoring its critical role in connecting RNAi

with heterochromatin formation [61]. Meanwhile, euchromatic loci may sporadically

generate double-stranded RNA, initiating RNAi and leading to heterochromatin deposition

via a positive feedback loop involving the RITS complex and Clr4 [62]. [62]. Interestingly,

in our study, we identified FK506-resistant epimutants with varying combinations of siRNA

accumulation and H3K9me2 enrichment at the *fkbA* locus (**Fig 4**). These findings support

several non-exclusive models: (1) RNAi-driven initiation of heterochromatin, (2)

heterochromatin-facilitated RNAi, or (3) other mechanisms.

To explore the first hypothesis, we conducted a comparative analysis of RNAi,

H3K9 methylation-based heterochromatin formation, and RITS components across *M.*

bainieri, *M. atramentarius*, *M. lusitanicus*, and *M. circinelloides* to assess the potential

coupling between RNAi and heterochromatin pathway. As reported previously in *M.*

lusitanicus [30], the RITS components Chp1 and Tas3 are absent across all four *Mucor*

species, whereas the bridging protein Stc1, the histone methyltransferase Clr4, and

CLRC complex subunits (Rik1, Cul4, and Raf1) are conserved (**S12 Fig**). Thus, although

elements of the coupling machinery such as Stc1 and CLRC are conserved, the apparent

absence of a complete RITS complex suggests that the connection between RNAi and

heterochromatin may operate in a partial or context-dependent manner. Under FK506

selective pressure, RNAi may be initiated at the *fkbA* locus, leading in some cases to the

establishment of heterochromatin, while in others, RNAi may act independently. Even

when heterochromatin is formed, weak or transient interactions between RNAi and

heterochromatin could result in heterochromatin-only epimutants.

Second, preexisting heterochromatin could facilitate RNAi, serving as a template

for small RNA generation. To explore this possibility, we analyzed the distribution of H3K9

dimethylation and sRNA in wild-type strains of four *Mucor* species (*M. bainieri*, *M.*

atramentarius, *M. lusitanicus*, and *M. circinelloides*). As shown in **S13 Fig**, discrete

H3K9me2 islands were detected downstream of *fkbA* in all four species, suggesting that

preexisting heterochromatin on transposon-rich regions may spread toward the *fkbA*

locus under FK506 selective pressure and recruit the RNAi machinery to generate small

RNAs. Similar phenomena have been reported in other organisms. In *S. pombe*, for

example, caffeine stress triggers redistribution of H3K9 methylation and evokes sRNA

generation [17]. Interestingly, these H3K9me2 islands lacked detectable sRNAs in *M.*

lusitanicus and *M. circinelloides*, whereas the co-occurrence of H3K9me2 and sRNAs

was observed in *M. bainieri* and *M. atramentarius*, raising the possibility that the

relationship between RNAi and heterochromatin formation may interact differently across

species.

Third, RNAi and heterochromatin formation may occur simultaneously under

FK506 pressure but act independently without functional coupling. A previous study

reported that RNAi and heterochromatin formation can function independently to repress

GremLINE1 transposable elements and maintain genome stability in *M. lusitanicus* [30].

Thus, it is possible that, upon FK506 exposure, these machineries were independently

activated at the same genomic region, with RNAi generating sRNAs at the *fkbA* locus and

heterochromatin spreading from adjacent transposon-rich regions, resulting in their

apparent coexistence without direct mechanistic interaction. Taken together, future

studies analyzing pathway-specific mutants will be critical to determine the directionality

and mechanistic coupling or uncoupling of these epigenetic processes.

Beyond FK506-induced dynamics at the *fkbA* locus, our comparative genomic

analyses across four *Mucor* species provide additional insight into how RNAi and

heterochromatin interplay may differ among species. GremLINE elements are present in

M. atramentarius, *M. lusitanicus*, and *M. circinelloides*, where they reside within or near

putative centromeric regions, but are absent from the *M. bainieri* genome (**S14 Fig**). The

centromeric GremLINEs in these species consistently show strong H3K9me2 enrichment,

supporting their association with heterochromatic domains. Notably, *M. bainieri* exhibits

a distinct architecture. Although it lacks GremLINEs, its putative centromeric regions are

enriched for a different repeat, TE00000158, which is associated with strong H3K9me2.

This pattern diverges sharply from the GremLINE-associated uncoupling observed in *M.*

circinelloides, raising the possibility that *M. bainieri* may have evolved a species-specific

centromeric repeat landscape and RNAi-heterochromatin relationship. Collectively, these

comparative observations suggest that the interplay between RNAi and heterochromatin

may not be conserved and may vary across species, likely influenced by species-specific

differences in repeat landscapes and centromere architecture.

Torres-Garcia et al. demonstrated that external stressors such as caffeine induce

phenotypic plasticity via Clr4/H3K9me "read-write" systems in *S. pombe*, underscoring

the adaptive function of heterochromatin in non-pathogenic yeast [17]. Our findings

extend this concept by providing direct evidence that H3K9 methylation contributes to

FK506 resistance in dimorphic *Mucor* species. Notably, this mechanism operates in the

non-pathogenic species *M. bainieri* and *M. atramentarius* species, a species with

pathogenic potential, suggesting that heterochromatin-mediated resistance is conserved

across diverse fungal taxa, spanning both unicellular yeasts and multinuclear hyphal

species. While RNAi-dependent antifungal resistance has been shown to be rapidly

induced and reversed following *in vivo* passage in a murine model [36], the *in vivo* stability

of heterochromatin-dependent epimutants has not been previously characterized. Here,

we demonstrate that a heterochromatin-dependent epimutant remains stable *in vivo* (**Fig**

6 and **Fig 7**). The persistence, even in the absence of antifungal pressure, raises the

possibility that epigenetically resistant strains could silently disseminate in clinical settings,

complicating diagnostics and treatment.

In summary, our study extends RNAi-dependent gene silencing to additional

Mucor species and uncovers that heterochromatin-mediated silencing also drives

antifungal resistance. These findings underscore the remarkable epigenetic plasticity of

fungal epigenomes and highlight adaptive strategies that enable reversible, yet durable,

resistance. Targeting RNAi and chromatin-modifying pathways may thus represent a

promising therapeutic strategy against fAMR.

Material and methods

Fungal strains, media and growth conditions

Mucor strains studied here are described in **S1 Table**. Mucor strains were grown on liquid or solid yeast extract peptone dextrose (YPD; BD, Difco, Franklin Lakes, NJ, USA) at room temperature. To isolate FK506-/rapamycin-resistant strains, 10^5 spores were inoculated onto more than ten solid YPD, supplemented with antifungal drug 1 μg/mL of FK506 (Astellas Pharma, Chuo, Tokyo, Japan) and 100 ng/ml of rapamycin (API Chem, Hangzhou, Zhejiang, Chian) and incubated for 3, 5, 7 days [15]. To test the stability of FK506-resistant strains, spores of the resistant strains were collected and inoculated in liquid YPD without any drugs overnight [16]. These were then serially subcultured overnight for several rounds, after which the passaged cells were spotted onto YPD or YPD supplemented with FK506 to confirm phenotypic plasticity.

Illumina sequencing and genome assembly

Mucor mycelium were grown on YPD medium, harvested, frozen in liquid nitrogen, and ground into a find powder. Genomic DNA samples were prepared using Norgen Biotek Yeast/Fungi Genomic DNA Isolation Kit (Ontario, Canada). Genomic DNA libraries were constructed with Roche KAPA HyperPrep Kit. 150-bp paired-end reads were obtained utilizing the NovaSeq X Plus sequencing system (Illumina) at the Duke Sequencing and Genomic Technologies Core (SGT). Raw reads were checked for quality metrics using FastQC v0.11.7 and processed by fastp v0.23.4 to remove adapters and low-quality bases or reads. Genomes were assembled using SPAdes v3.15.5 set to

isolate default mode. Genome assemblies were cleaned, sorted, masked, and annotated

using the Funannotate v1.8.15 [63].

Nanopore sequencing and genome assembly

A total of 10^6 spores from *M. bainieri* and *M. atramentarius* wild-type strains were

inoculated into liquid YPD medium and incubated overnight at 25 °C. Cells were

harvested by centrifugation, washed with distilled water, frozen with liquid nitrogen, and

ground with a mortar and pestle. High-molecular-weight (HMW) DNA was extracted from

each sample using the Monarch HMW DNA Extraction Kit for Tissue (NEB, Ipswich, MA,

USA) according to the manufacturer's protocol. DNA concentration and quality were

assessed using a NanoDrop spectrophotometer (ThermoFisher, Waltham, MA, USA) and

a Qubit fluorometer with the Qubit dsDNA High Sensitivity kit (ThermoFisher). DNA

integrity was confirmed by CHEF electrophoresis performed at 6.0 V in a 1% agarose gel

for 18 hours, which revealed intact HMW DNA. Libraries were prepared using the ligation

sequencing kit V14 (SQK-LSK114; Oxford Nanopore Technologies, Oxford, United

Kingdom) and sequenced on MinION R10.4.1 flow cells (ONT; Oxford Nanopore

Technologies) following to the manufacturer's protocol in-house in our laboratory.

POD5 base calling was performed with the Dorado basecaller (v0.5.2; Oxford

Nanopore Technologies, https://github.com/nanoporetech/dorado). Demultiplexing and

BAM-to-FASTQ conversion were carried out using Dorado demux (v0.5.2). The resulting

FASTQ files were error-corrected with Canu (v2.2) and assembled using Flye (v2.9.5-

b1801) [64, 65]. The expected genome size was set to 50 Mb, and a minimum overlap

length of 5 kb was required between reads. The assembled genome was subsequently

polished in three iterations using Pilon (v1.24; Broad Institute,

https://github.com/broadinstitute/pilon/releases/tag/v1.24) with default parameters

utilizing the raw Illumina sequencing data deposited under NCBI BioProject accessions

PRJNA1277536 (*M. bainieri* CBS293.63) and PRJNA1277544 (*M. atramentarius*

CBS202.28). Repetitive sequences were identified using RepeatMasker (v4.1.5;

https://www.repeatmasker.org) with default parameters and genome annotation was

performed using Funannotate (v1.8.17) with the masked genome assembly and strain-

specific RNA-seq alignments [63]. This pipeline produced high-quality genome

assemblies, with *M. bainieri* assembled into 30 contigs (N50 = 6 Mb; 98.7% BUSCO

completeness; 9,781 predicted genes) and *M. atramentarius* into 33 contigs (N50 = 2 Mb;

98.8% BUSCO completeness; 11,597 predicted genes).

The *fkbA* and calcineurin gene sequencing analysis

Mucor strains were cultured on YPD medium for three days. Approximately 1–5

mg of harvested cells was suspended in 300 µL Tissue & Cell Lysis Buffer (Lucigen,

Middleton, WI, USA) and 1 µL proteinase K (ThermoFisher). The mixture was incubated

at 65°C for 45 min, followed by the addition of 150 µL MPC Protein Precipitation Reagent

(Lucigen). After centrifugation, the pellet was washed with 70% ethanol and resuspended

in distilled water, and genomic DNA (gDNA) quality was analyzed using a Qubit and a

NanoDrop (ThermoFisher). The promoter, open reading frame, and terminator regions of

the *fkbA* gene were amplified by polymerase chain reaction (PCR) using primers listed in

S2 Table. The purified PCR products were sequenced using long read sequencing with

ONT at Plasmidsaurus Inc. (Eugene, CA, USA) and analyzed against the reference *fkbA*

sequence using Clustal Omega (v1.2.3).

RNA extraction and quantitative PCR (qPCR) analysis

Total RNA was extracted from 10 mg of *Mucor* cells cultured on YPD, with or

without FK506, at room temperature. After overnight lyophilization, each sample was

homogenized in a FastPrep bead-beater (MP Biomedicals, Santa Ana, CA, USA) with 0.2

mL of zirconia-silica beads (BioSpec Products, Bartlesville, OK, USA). Total RNA was

then purified with the miRNeasy Mini Kit (Qiagen, Hilden, Germany) and RNA quantity

and quality were analyzed using a Nanodrop (Thermo Fisher) and a Qubit Fluorometer

with Qubit RNA High Sensitivity kit (ThermoFisher) [66]. The isolated RNA was treated

with DNase and subsequently employed for cDNA synthesis with the Maxima H Minus

cDNA Synthesis Master Mix (ThermoFisher, Waltham, MA, USA). Real-time PCR was

performed with PowerUp SYBR Green Master Mix (ThermoFisher) on a QuantStudio 3

qPCR system (ThermoFisher) in technical triplicate. The primers for qPCR are listed in

S2 Table. Gene expression was normalized to *actA*, relative fold expression was

calculated using $2^{-\Delta\Delta Ct}$ method.

Small RNA (sRNA) sequencing

sRNA libraries were constructed with the QIAseq miRNA Library Kit (Qiagen) and

sequenced on an NovaSeq X platform (Illumina) by 75-bp single end. Raw reads were

trimmed with adaptor sequence (5′-GTTCAGAGTTCTACAGTCCGACGATC-3′ and 5′-

AACTGTAGGCACCATCAA-3′) utilizing BBDuk. Filtered reads were then mapped to the

reference genomes (*M. bainieri* CBS293.63; PRJNA1277536 or *M. atramentarius*

CBS202.28; PRJNA1277544) using BBMap with default parameters (last modified

September 15, 2022) [67]. The resulting SAM files were converted to BAM files, and

antisense reads were sorted using SAMtools (v1.21) [68]. BAM files were converted to

TDF files using IGV tools with a window size of 20 bp and the mean window function. The

data was subsequently visualized in the Integrative Genome Viewer (IGV v2.16.0), where

the Y-axis represents the average read coverage per 20-bp window [69]. The Y-axis

reflects the average read coverage per 20 bp window. Peak calling was performed with

MACS3 using the --nomodel option with an extension size of 147 bp. sRNA-seq

experiments were performed with one biological sample per strain.

Western blot

Each *Mucor* strain was cultivated in YPD, with or without FK506, at room

temperature for three days. Ten milligrams of *Mucor* cells were lyophilized overnight and

homogenized using a FastPrep bead-beater (MP Biomedicals) with 0.2 mL of zirconia-

silica beads (BioSpec Products). Pierce RIPA buffer (ThermoFisher) containing a

protease inhibitor tablet (Roche) was then added, and each sample was further disrupted

with a FastPrep bead-beater. The samples were incubated on ice for at least 30 min. Cell

extracts were centrifuged at 14,000 RPM, 4°C for 10 min, and the supernatants were

quantified with the Pierce BCA Protein Assay Kit (ThermoFisher). Next, 20 μg of each

sample was mixed with Novex Tris-Glycine SDS Sample Buffer (Invitrogen, Waltham, MA,

USA) and boiled at 85°C for 5 min. After briefly cooling, the samples were separated by

SDS-PAGE, transferred to a PVDF membrane, and incubated overnight at 4°C with an

anti-FKBP12 antibody (Invitrogen) or an anti-GAPDH antibody (Proteintech, Rosemont,

IL, USA). Following incubation with a horseradish peroxidase-conjugated anti-rabbit

antibody, FKBP12 and GAPDH expression were detected on a ChemiDoc MP imaging

system (Bio-Rad, Hercules, CA, USA).

Chromatin immunoprecipitation (ChIP)-qPCR

A total of 10^6 spores from each *Mucor* strain were inoculated in liquid YPD medium

with or without FK506 and incubated overnight at 25°C. The samples were then cross-

linked with 1% formaldehyde (ThermoFisher), followed by addition of glycine (Sigma-

Aldrich, St. Louis, MO, USA) at a final concentration of 0.125M. After washing and

freezing using liquid nitrogen, each sample was ground with a pestle and suspended in

ChIP lysis buffer (50 mM HEPES, pH=7.5, 150 mM NaCl, 1 mM EDTA, pH=8, 1% Triton

X-100, and 0.1% sodium deoxycholate) containing a protease inhibitor tablet (Roche).

Disrupted cells were then sonicated with a Bioruptor (Diagenode) for 40 cycles, with 30s

on and 30s off and clarified by centrifuge for 5 min. The input samples were stored at

-80°C, while the IP samples were incubated overnight at 4°C with Dynabeads Protein A/G

(Invitrogen) pre-conjugated with the anti-histone H3K9 dimethyl antibody (Abcam,

Cambridge, UK) or the anti-RNA polymerase II antibody (Active Motif, Carlsbad, CA).

Crosslinking between protein and DNA in both input and IP samples was reversed with

the MAGnify Chromatin Immunoprecipitation System (Invitrogen), and DNA quality and

quality were assessed using a Nanodrop (Thermo Fisher) and a Qubit Fluorometer with

Qubit dsDNA High Sensitivity kit (ThermoFisher). qPCR was performed in technical

triplicate with primers described in **S2 Table**. For H3K9me2 ChIP-qPCR, GremLINE1

transposon elements were used as positive controls in *M. atramentarius* [30]. However,

our analysis of the *M. bainieri* genome suggests the GremLINE1 elements are absent in

the Illumina-based genome assembly. Instead, to develop a positive control for the

H3K9me2 ChIP-qPCR analysis in *M. bainieri*, we hypothesized that the loci enriched for

H3K9m2 should at least partially overlap with the genomic regions enriched for small

RNAs. Specifically, we first identified three genomic loci enriched for small RNAs based

on our sRNA analysis and then designed primers targeting these loci and tested them for

the presence of H3K9me2 signals using ChIP-qPCR. One of the three loc, *PS3_010814*

(identified in Illumina-based genome assembly deposited under NCBI SRA

PRJNA1277536), exhibited consistent H3K9me2 enrichment across samples including

WT, epimutants, and revertants, and was selected as the positive control for the

subsequent H3K9me2 ChIP-qPCR analyses. The *actA* gene, which lacks both small RNA

accumulation and heterochromatin marks, was used as a negative control. H3K9

methylation enrichment was calculated by normalizing the IP DNA to the input DNA with

the ΔCt method and further adjusted to the *actA* gene with the ΔΔCt method, followed by

$2^{-\Delta\Delta Ct}$ to determine relative enrichment [70, 71]. For RNA polymerase II ChIP-qPCR, the

actA gene was used as a positive control, as it is transcriptionally active. As a negative

control, we selected an intergenic locus defined as an siRNA-producing region embedded

within transposable element–rich sequences and associated with H3K9me2

heterochromatin. RNA polymerase II enrichment was similarly calculated by using the

$2^{-\Delta\Delta Ct}$ method with a negative control, an intronic region enriched for siRNAs and by

H3K9me2.

ChIP sequencing

Input and IP libraries were constructed with KAPA HyperPrep (Roche) and sequenced on a NovaSeq X platform (Illumina) by 150-bp paired end. Raw FASTA files were trimmed with Trimmomatic (v0.39) and filtered reads were mapped to the reference genomes (*M. bainieri* CBS293.63; PRJNA1277536 or *M. atramentarius* CBS202.28; PRJNA1277544) using Bowtie2 (v2.5.4) with default parameters [72, 73]. The resulting SAM files were converted to BAM files using SAMtools (v1.21) [68]. Broad ChIP-peak calling was analyzed with MACS3 (v3.0.3) with the parameters --mfold 10 1000 and --extsize 147, and enrichment was analyzed by generating coverage files with BamCoverage (v3.5.6), followed by counts per million (CPM) normalization [74, 75]. Processed BAM files were converted to TDF files IGV tools with a window size of 20 bp and the mean window function and subsequently visualized with IGV (v2.16.0) [76]. ChIP-seq experiments were performed with one biological sample per strain.

***In vivo* infection**

Six-week-old male BALB/cAnNCrl mice, 6 to 7 weeks old obtained from Jackson Laboratory (Bar Harbor, Maine, USA), were analyzed in this study. Five mice were infected with each *Mucor* strain. Two days prior to infection, 200 mg/kg of cyclophosphamide (Sigma-Aldrich), an immunosuppressant, was administered intraperitoneally. Each mouse was infected with 1×10^6 spores (in 50 µL PBS) via retro-orbital injection after anesthetizing the mice with isoflurane for 2.5 min. After infection, the mice were monitored twice daily and administered cyclophosphamide at 200 mg/kg body weight every four days. To assess pathogenicity, weight loss was monitored, and any animal experiencing ≥ 20% loss of body weight was humanely euthanized. To examine

epimutant stability, five mice per each strain were subjected under four treatment

conditions: (i) cyclophosphamide at 200 mg/kg every 4 days, (ii) cyclophosphamide at 50

mg/kg every 4 days for 2 weeks, (iii) no cyclophosphamide for 2 weeks, (iv) no

cyclophosphamide treatment for 4 weeks. At the designated time points, mice were

sacrificed, and fungal burden and epimutant stability were evaluated. All animal

experiments in this study were approved by the Duke University Institutional Animal Care

and Use Committee (IACUC) (protocol #A224-23-11). Animal care and experiments were

conducted according to IACUC ethical guidelines.

Stability of epimutants during *in vivo* infection

To assess fungal burden and isolate *Mucor* strains from organs, five mice per

Mucor strain were intravenously infected. At the humane endpoint, the brain, lung, liver,

and spleen were dissected. Whole organs were homogenized in 0.8 mL of PBS with a

bead beater for 2 min, then diluted and plated on YPD supplemented with 50 µg/mL

chloramphenicol. After a two-day incubation at room temperature, fungal colonies were

picked and patched onto YPD plates with or without FK506 to assess the stability of

FK506 resistance following *in vivo* passage. The stability rate was determined by

calculating the proportion of resistant colonies on YPD supplemented with FK506 relative

to the total colonies patched on YPD [36]. To determine whether resistant isolates were

epimutants, genomic DNA was extracted from all resistant isolates, and the *fkbA* region

was PCR-amplified and sequenced using long read sequencing with ONT in

Plasmidsaurus Inc. (Eugene, CA, USA). In addition, to determine whether the resistant

isolates represented heterochromatin-associated epimutants, several resistant isolates

were analyzed by ChIP-qPCR.

702   32

Quantification and statistical analyses

704   Statistical differences between the control and target strains were evaluated using

705 ordinary one-way ANOVA using Tukey's Honestly Significant Difference (HSD) test with

706 using GraphPad Prism (v10.1.1). Kaplan-Meier survival curves were generated and

707 analyzed with the log-rank test in R (v4.3.2) to estimate survival probability. Data are

708 reported as mean ± standard deviation. In all graphs, p values or asterisks were listed

709 indicating statistical significance ($^*p < 0.05$; $^{**}p < 0.01$; $^{***}p < 0.001$; $^{****}p < 0.0001$).

Funding Statement

This study was supported by the NIH/National Institute of Allergy and Infectious Diseases (R01-AI170543-04, R01-AI039115-28, and R01-AI050113-20) awarded to J.H.. J.H. is co-director and fellow of the CIFAR program Fungal Kingdom: Threats & Opportunities.

Acknowledgments

We thank Dr. Vikas Yadav for his insightful advice on bioinformatics analyses as well as for his scientific discussions and critical reading. We are grateful to Dr. María Isabel Navarro-Mendoza for providing guidance on *in vivo* training. We also thank Anna Floyd Averette for her invaluable support and assistance with *in vivo* assays; Ziyan Xu for preparing DNA for Illumina whole genome sequencing; Dr. Jason E. Stajich (University of California) for providing whole-genome sequence datasets and initial preliminary analysis; and Dr. Grit Walther (Hans Knöll Institute) for providing *Mucor* isolates. We appreciate critical reading by Dr. Sheng Sun and Dr. Jun Huang. We thank Devi Swain, Director of the Duke Sequencing and Genomic Technologies Core for technical support.

Author Contributions

Conceptualization: Ye-Eun Son, Joseph Heitman

Methodology: Ye-Eun Son, Carlos Pérez-Arques

Software: Ye-Eun Son, Carlos Pérez-Arques

Validation: Ye-Eun Son

Investigation: Ye-Eun Son, Carlos Pérez-Arques

Visualization: Ye-Eun Son, Carlos Pérez-Arques

Resources: Joseph Heitman

Writing – Original Draft: Ye-Eun Son

Writing – Review & Editing: Ye-Eun Son, Carlos Pérez-Arques, Joseph Heitman

Supervision: Joseph Heitman

Funding Acquisition: Joseph Heitman

Competing interests

The authors declare no conflicts of interest.

Data Availability

Small RNA-seq, ChIP-seq, Nanopore sequencing data, and the corresponding

genome assemblies are available in the NCBI SRA under project accession numbers

PRJNA1242486 (*M. bainieri* CBS293.63) and PRJNA1242487 (*M. atramentarius*

CBS202.28). Illumina whole-genome sequencing data are available under project

accession numbers PRJNA1277536 (*M. bainieri* CBS293.63) and PRJNA1277544 (*M.*

atramentarius CBS202.28).

References

1. Case NT, Gurr SJ, Fisher MC, Blehert DS, Boone C, Casadevall A, et al. Fungal impacts on earth's ecosystems. Nature. 2025;638(8049):49-57. pmid: 39910383.
2. McDermott A. Drug-resistant fungi on the rise. Proc Natl Acad Sci U S A. 2022;119(48):e2217948119. pmid: 36413501.
3. Fisher MC, Denning DW. The WHO fungal priority pathogens list as a game-changer. Nat Rev Microbiol. 2023;21(4):211-2. pmid: 36747091.
4. Lockhart SR, Chowdhary A, Gold JAW. The rapid emergence of antifungal-resistant human-pathogenic fungi. Nat Rev Microbiol. 2023;21(12):818-32. pmid: 37648790.
5. Perlin DS, Rautemaa-Richardson R, Alastruey-Izquierdo A. The global problem of antifungal resistance: prevalence, mechanisms, and management. Lancet Infect Dis. 2017;17(12):e383-e92. pmid: 28774698.
6. Nnadi NE, Carter DA. Climate change and the emergence of fungal pathogens. PLoS Pathog. 2021;17(4):e1009503. pmid: 33914854.
7. Fisher MC, Burnett F, Chandler C, Gow NAR, Gurr S, Hart A, et al. A one health roadmap towards understanding and mitigating emerging fungal antimicrobial resistance: fAMR. NPJ Antimicrob Resist. 2024;2(1):36. pmid: 39524479.
8. Lee Y, Robbins N, Cowen LE. Molecular mechanisms governing antifungal drug resistance. NPJ Antimicrob Resist. 2023;1(1):5. pmid: 38686214.
9. Al Aboud NM, Tupper C, Jialal I. Genetics, epigenetic mechanism. StatPearls. Treasure Island (FL)2025.
10. Flowers SA, Colon B, Whaley SG, Schuler MA, Rogers PD. Contribution of clinically derived mutations in *ERG11* to azole resistance in *Candida albicans*. Antimicrob Agents Chemother. 2015;59(1):450-60. pmid: 25385095.
11. Lewis RE, Liao G, Hou J, Prince RA, Kontoyiannis DP. Comparative *in vivo* dose-dependent activity of caspofungin and anidulafungin against echinocandin-susceptible and -resistant *Aspergillus fumigatus*. J Antimicrob Chemother. 2011;66(6):1324-31. pmid: 21486855.
12. Nagy G, Kiss S, Varghese R, Bauer K, Szebenyi C, Kocsube S, et al. Characterization of three pleiotropic drug resistance transporter genes and their participation in the azole resistance of *Mucor circinelloides*. Front Cell Infect Microbiol. 2021;11:660347. pmid: 33937100.
13. Navarro-Mendoza MI, Perez-Arques C, Parker J, Xu Z, Kelly S, Heitman J. Alternative ergosterol biosynthetic pathways confer antifungal drug resistance in the human pathogens within the *Mucor* species complex. mBio. 2024;15(8):e0166124. pmid: 38980037.
14. Chang Z, Yadav V, Lee SC, Heitman J. Epigenetic mechanisms of drug resistance in fungi. Fungal Genet Biol. 2019;132:103253. pmid: 31325489.
15. Calo S, Shertz-Wall C, Lee SC, Bastidas RJ, Nicolas FE, Granek JA, et al. Antifungal drug resistance evoked via RNAi-dependent epimutations. Nature. 2014;513(7519):555-8. pmid: 25079329.
16. Chang Z, Billmyre RB, Lee SC, Heitman J. Broad antifungal resistance mediated by RNAi-dependent epimutation in the basal human fungal pathogen *Mucor circinelloides*. PLoS Genet. 2019;15(2):e1007957. pmid: 30742617.

17.   Torres-Garcia S, Yaseen I, Shukla M, Audergon P, White SA, Pidoux AL, et al.
Epigenetic gene silencing by heterochromatin primes fungal resistance. Nature.
2020;585(7825):453-8. pmid: 32908306.
18.   Hernández JL, Buckley CJ. Mucormycosis. StatPearls. Treasure Island
(FL)2025.
19.   Binder U, Maurer E, Lass-Florl C. Mucormycosis--from the pathogens to the
disease. Clin Microbiol Infect. 2014;20 Suppl 6:60-6. pmid: 24476149.
20.   Lax C, Nicolas FE, Navarro E, Garre V. Molecular mechanisms that govern
infection and antifungal resistance in Mucorales. Microbiol Mol Biol Rev.
2024;88(1):e0018822. pmid: 38445820.
21.   Cornely OA, Alastruey-Izquierdo A, Arenz D, Chen SCA, Dannaoui E,
Hochhegger B, et al. Global guideline for the diagnosis and management of
mucormycosis: an initiative of the european onfederation of medical mycology in
cooperation with the mycoses study group education and research consortium.
Lancet Infect Dis. 2019;19(12):e405-e21. pmid: 31699664.
22.   Roden MM, Zaoutis TE, Buchanan WL, Knudsen TA, Sarkisova TA, Schaufele
RL, et al. Epidemiology and outcome of zygomycosis: a review of 929 reported
cases. Clin Infect Dis. 2005;41(5):634-53. pmid: 16080086.
23.   Devnath P, Dhama K, Tareq AM, Emran TB. Mucormycosis coinfection in the
context of global COVID-19 outbreak: A fatal addition to the pandemic spectrum.
Int J Surg. 2021;92:106031. pmid: 34311126.
24.   WHO. WHO fungal priority pathogens list to guide research, development and
public health action2022.
25.   Wagner L, Stielow JB, de Hoog GS, Bensch K, Schwartze VU, Voigt K, et al. A
new species concept for the clinically relevant *Mucor circinelloides* complex.
Persoonia. 2020;44:67-97. pmid: 33116336.
26.   Perez-Arques C, Navarro-Mendoza MI, Xu Z, Walther G, Heitman J. RNAi
epimutations conferring antifungal drug resistance are inheritable. Nat Commun.
2025;16(1):7293. pmid: 40775213.
27.   Lee SC, Li A, Calo S, Inoue M, Tonthat NK, Bain JM, et al. Calcineurin
orchestrates dimorphic transitions, antifungal drug responses and host-pathogen
interactions of the pathogenic mucoralean fungus *Mucor circinelloides*. Mol
Microbiol. 2015;97(5):844-65. pmid: 26010100.
28.   Lee SC, Li A, Calo S, Heitman J. Calcineurin plays key roles in the dimorphic
transition and virulence of the human pathogenic zygomycete *Mucor
circinelloides*. PLoS Pathog. 2013;9(9):e1003625. pmid: 24039585.
29.   Bastidas RJ, Shertz CA, Lee SC, Heitman J, Cardenas ME. Rapamycin exerts
antifungal activity in vitro and in vivo against *Mucor circinelloides* via FKBP12-
dependent inhibition of Tor. Eukaryot Cell. 2012;11(3):270-81. pmid: 22210828.
30.   Navarro-Mendoza MI, Perez-Arques C, Heitman J. Heterochromatin and RNAi
act independently to ensure genome stability in Mucorales human fungal
pathogens. Proc Natl Acad Sci U S A. 2023;120(7):e2220475120. pmid:
36745785.
31.   Huang J, Larmore CJ, Priest SJ, Xu Z, Dietrich FS, Yadav V, et al. Distinct
evolutionary trajectories following loss of RNA interference in *Cryptococcus*

neoformans. Proc Natl Acad Sci U S A. 2024;121(47):e2416656121. pmid:
39536081.

32.   Mi S, Cai T, Hu Y, Chen Y, Hodges E, Ni F, et al. Sorting of small RNAs into
Arabidopsis argonaute complexes is directed by the 5' terminal nucleotide. Cell.
2008;133(1):116-27. pmid: 18342361.

33.   Bhattacharjee S, Roche B, Martienssen RA. RNA-induced initiation of
transcriptional silencing (RITS) complex structure and function. RNA Biol.
2019;16(9):1133-46. pmid: 31213126.

34.   Allshire RC, Madhani HD. Ten principles of heterochromatin formation and
function. Nat Rev Mol Cell Biol. 2018;19(4):229-44. pmid: 29235574.

35.   Filipowicz W, Jaskiewicz L, Kolb FA, Pillai RS. Post-transcriptional gene silencing
by siRNAs and miRNAs. Curr Opin Struct Biol. 2005;15(3):331-41. pmid:
15925505.

36.   Chang Z, Heitman J. Drug-resistant epimutants exhibit organ-specific stability
and induction during murine infections caused by the human fungal pathogen
Mucor circinelloides. mBio. 2019;10(6):e02579-19. pmid: 31690679.

37.   Rhodes J, Abdolrasouli A, Farrer RA, Cuomo CA, Aanensen DM, Armstrong-
James D, et al. Genomic epidemiology of the UK outbreak of the emerging
human fungal pathogen *Candida auris*. Emerg Microbes Infect. 2018;7(1):43.
pmid: 29593275.

38.   Fraaije B, Atkins S, Hanley S, Macdonald A, Lucas J. The multi-fungicide
resistance status of *Aspergillus fumigatus* populations in arable soils and the
wider European environment. Front Microbiol. 2020;11:599233. pmid: 33384673.

39.   Ikuta KS, Mestrovic T, Naghavi M. Global incidence and mortality of severe
fungal disease. Lancet Infect Dis. 2024;24(5):e268. pmid: 38395046.

40.   Park HS, Lee SC, Cardenas ME, Heitman J. Calcium-calmodulin-calcineurin
signaling: A globally conserved virulence cascade in eukaryotic microbial
pathogens. Cell Host Microbe. 2019;26(4):453-62. pmid: 31600499.

41.   Trivedi P, Leach JE, Tringe SG, Sa T, Singh BK. Plant-microbiome interactions:
from community assembly to plant health. Nat Rev Microbiol. 2020;18(11):607-
21. pmid: 32788714.

42.   Faust K, Raes J. Microbial interactions: from networks to models. Nat Rev
Microbiol. 2012;10(8):538-50. pmid: 22796884.

43.   Weiland-Brauer N. Friends or Foes-Microbial Interactions in Nature. Biology
(Basel). 2021;10(6). pmid: 34199553.

44.   Getzke F, Thiergart T, Hacquard S. Contribution of bacterial-fungal balance to
plant and animal health. Curr Opin Microbiol. 2019;49:66-72. pmid: 31731228.

45.   de Menezes AB, Richardson AE, Thrall PH. Linking fungal-bacterial co-
occurrences to soil ecosystem function. Curr Opin Microbiol. 2017;37:135-41.
pmid: 28692866.

46.   Baumgardner DJ. Soil-related bacterial and fungal infections. J Am Board Fam
Med. 2012;25(5):734-44. pmid: 22956709.

47.   Procopio RE, Silva IR, Martins MK, Azevedo JL, Araujo JM. Antibiotics produced
by *Streptomyces*. Braz J Infect Dis. 2012;16(5):466-71. pmid: 22975171.

48.   Kino T, Hatanaka H, Hashimoto M, Nishiyama M, Goto T, Okuhara M, et al. FK-
506, a novel immunosuppressant isolated from a *Streptomyces*. I. Fermentation,

isolation, and physico-chemical and biological characteristics. J Antibiot (Tokyo).
1987;40(9):1249-55. pmid: 2445721.

49.   Arndt C, Cruz MC, Cardenas ME, Heitman J. Secretion of FK506/FK520 and
rapamycin by *Streptomyces* inhibits the growth of competing *Saccharomyces*
cerevisiae and *Cryptococcus neoformans*. Microbiology (Reading). 1999;145 (Pt
8):1989-2000. pmid: 10463165.

50.   Ziaee A, Zia M, Bayat M, Hashemi J. Identification of Mucorales isolates from soil
using morphological and molecular methods. Curr Med Mycol. 2016;2(1):13-9.
pmid: 28681007.

51.   Abu-Salah KM. Amphotericin B: an update. Br J Biomed Sci. 1996;53(2):122-33.
pmid: 8757689.

52.   Nicolas FE, Torres-Martinez S, Ruiz-Vazquez RM. Two classes of small
antisense RNAs in fungal RNA silencing triggered by non-integrative transgenes.
EMBO J. 2003;22(15):3983-91. pmid: 12881432.

53.   Calo S, Nicolas FE, Lee SC, Vila A, Cervantes M, Torres-Martinez S, et al. A
non-canonical RNA degradation pathway suppresses RNAi-dependent
epimutations in the human fungal pathogen *Mucor circinelloides*. PLoS Genet.
2017;13(3):e1006686. pmid: 28339467.

54.   Nicolas FE, Moxon S, de Haro JP, Calo S, Grigoriev IV, Torres-Martinez S, et al.
Endogenous short RNAs generated by Dicer 2 and RNA-dependent RNA
polymerase 1 regulate mRNAs in the basal fungus *Mucor circinelloides*. Nucleic
Acids Res. 2010;38(16):5535-41. pmid: 20427422.

55.   Brodersen P, Voinnet O. The diversity of RNA silencing pathways in plants.
Trends Genet. 2006;22(5):268-80. pmid: 16567016.

56.   Hamilton A, Voinnet O, Chappell L, Baulcombe D. Two classes of short
interfering RNA in RNA silencing. EMBO J. 2015;34(20):2590. pmid: 26291654.

57.   Chakravarthy S, Sternberg SH, Kellenberger CA, Doudna JA. Substrate-specific
kinetics of Dicer-catalyzed RNA processing. J Mol Biol. 2010;404(3):392-402.
pmid: 20932845.

58.   Vermeulen A, Behlen L, Reynolds A, Wolfson A, Marshall WS, Karpilow J, et al.
The contributions of dsRNA structure to Dicer specificity and efficiency. RNA.
2005;11(5):674-82. pmid: 15811921.

59.   Djupedal I, Ekwall K. Epigenetics: heterochromatin meets RNAi. Cell Res.
2009;19(3):282-95. pmid: 19188930.

60.   Okazaki K, Kato H, Iida T, Shinmyozu K, Nakayama JI, Murakami Y, et al. RNAi-
dependent heterochromatin assembly in fission yeast S*chizosaccharomyces*
pombe requires heat-shock molecular chaperones Hsp90 and Mas5. Epigenetics
Chromatin. 2018;11(1):26. pmid: 29866182.

61.   Bayne EH, White SA, Kagansky A, Bijos DA, Sanchez-Pulido L, Hoe KL, et al.
Stc1: a critical link between RNAi and chromatin modification required for
heterochromatin integrity. Cell. 2010;140(5):666-77. pmid: 20211136.

62.   Yu R, Wang X, Moazed D. Epigenetic inheritance mediated by coupling of RNAi
and histone H3K9 methylation. Nature. 2018;558(7711):615-9. pmid: 29925950.

63.   Funannotate v1.8.1: Eukaryotic genome annotation [Internet]. Zenodo. 2020.

64. Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. Canu:
scalable and accurate long-read assembly via adaptive k-mer weighting and
repeat separation. Genome Res. 2017;27(5):722-36. pmid: 28298431.
65. Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-prone reads
using repeat graphs. Nat Biotechnol. 2019;37(5):540-6. pmid: 30936562.
66. Son YE, Yu JH, Park HS. The novel spore-specific regulator SscA controls
Aspergillus conidiogenesis. mBio. 2023;14(5):e0184023. pmid: 37707170.
67. Bushnell B. BBMap: A fast, accurate, splice-aware aligner. 2014.
68. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The
Sequence Alignment/Map format and SAMtools. Bioinformatics.
2009;25(16):2078-9. pmid: 19505943.
69. Priest SJ, Yadav V, Roth C, Dahlmann TA, Kuck U, Magwene PM, et al.
Uncontrolled transposition following RNAi loss causes hypermutation and
antifungal drug resistance in clinical isolates of *Cryptococcus neoformans*. Nat
Microbiol. 2022;7(8):1239-51. pmid: 35918426.
70. Park HS, Man Yu Y, Lee MK, Jae Maeng P, Chang Kim S, Yu JH. Velvet-
mediated repression of beta-glucan synthesis in *Aspergillus nidulans* spores. Sci
Rep. 2015;5:10199. pmid: 25960370.
71. Navarro-Mendoza MI, Perez-Arques C, Panchal S, Nicolas FE, Mondo SJ,
Ganguly P, et al. Early diverging fungus *Mucor circinelloides* lacks centromeric
histone CENP-A and displays a mosaic of point and regional centromeres. Curr
Biol. 2019;29(22):3791-802 e6. pmid: 31679929.
72. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics. 2014;30(15):2114-20. pmid: 24695404.
73. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat
Methods. 2012;9(4):357-9. pmid: 22388286.
74. Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, et al.
Model-based analysis of ChIP-Seq (MACS). Genome Biol. 2008;9(9):R137.
pmid: 18798982.
75. Ramirez F, Dundar F, Diehl S, Gruning BA, Manke T. deepTools: a flexible
platform for exploring deep-sequencing data. Nucleic Acids Res. 2014;42:W187-
91. pmid: 24799436.
76. Robinson JT, Thorvaldsdottir H, Turner D, Mesirov JP. igv.js: an embeddable
JavaScript implementation of the Integrative Genomics Viewer (IGV).
Bioinformatics. 2023;39(1). pmid: 36562559.
965

Figure legends

Fig 1. FK506-resistant strains isolated from the *Mucor circinelloides* complex.

(A) Schematic representation of the procedure followed to obtain FK506- and/or rapamycin-resistant strains from *Mucor circinelloides* complex species. This Figure was generated with BioRender.com. **(B)** Phenotypic analysis of FK506-resistant isolates in *M. janssenii* CBS185.68 (PS1), *M. bainieri* CBS293.63 (PS3), and *M. atramentarius* CBS202.28 (PS6). WT, wild-type; M, Mendelian mutant; E, Epimutant.

Fig 2. FK506-resistant epimutants exhibit silencing of *fkbA* mRNA and FKBP12 protein expression and are unstable.

(A, D) Phenotypic reversion of FK506-resistant epimutant strains following serial passage without drug selection in *M. bainieri* (A) or *M. atramentarius* (D). Red arrows indicate the points at which epimutants reverted to the WT phenotype. FK506 was utilized at 1 µg/mL and rapamycin at 100 ng/mL. WT, wild-type; P, number of passages. **(B, E)** Quantification of the *fkbA* mRNA expression in *M. bainieri* (B) *or M. atramentarius* (E). Error bars represent mean ± SEM (n = 3). Statistical significance: ****$p \leq 0.0001$. **(C, F)** Western blot analysis of FKBP12 protein expression in *M. bainieri* (C) or *M. atramentarius* (F). PS3: E1–E9, epimutants; E1P4–E9P24, revertants. PS6: E1 and E2, epimutants; E1P32 and E2P28, revertants.

Fig 3. Small RNAs mediate FK506 resistance in *M. bainieri*.

(A) Small RNA coverage mapped across the *fkbA* gene and its neighboring loci. The right panel presents a detailed view of the *fkbA* gene and its neighboring loci. Peaks shown below the coverage track indicate statistically enriched small RNA intervals identified by the MACS3 peak-calling algorithm, which models background signal to detect significant enrichment. **(B)** Small RNA abundance (RPM: reads per million) mapped to the *fkbA* gene. **(C)** 5′-end nucleotide preference of small RNAs associated with the *fkbA* gene. **(D)** Size distribution of small RNAs mapped to the *fkbA* gene. WT, wild-type; E1–E9, epimutants; E1P4–E9P24, revertants.

Fig 4. Heterochromatin marks underlie epigenetic FK506 resistance in *M. bainieri*.

(A) ChIP-qPCR analysis of H3K9me2 enrichment at the positive control region (an siRNA-enriched locus validated by H3K9me2) and *fkbA* locus (targeting the 5′-UTR, exonic regions, and 3′-UTR). Error bars represent mean ± SEM (n = 3). Statistical significance: $*p \le 0.05$, $**p \le 0.01$; $****p \le 0.0001$. **(B, C)** ChIP-seq coverage mapped across the positive control locus (an siRNA-enriched locus validated by H3K9me2; B) or *fkbA* and its neighboring loci (C). The peaks shown beneath the coverage track denote significant H3K9me2-enriched intervals identified by the MACS3 peak-calling algorithm, which detects statistically enriched regions by modeling background signal. **(D)** ChIP-qPCR analysis of RNA polymerase II enrichment at the positive control region (*actA*), *fkbA* locus (targeting the 5′-UTR and exonic region), a *fkbA*-neighboring gene, and negative control, defined as an siRNA-producing region flanked by transposable elements and associated

1007 with H3K9me2. Statistical significance: ****$p \le 0.0001$. WT, wild-type; E1–E8, epimutants;

1008 E1P4–E8P28, revertants.

1009

Fig 5. Heterochromatin-associated FK506 resistance is conserved in *M.

atramentarius.**

(A) Small RNA coverage mapped across the *fkbA* gene and its neighboring loci. Below

panel presents a detailed view of the *fkbA* gene and its neighboring loci. Peaks shown

below the coverage track indicate statistically enriched small RNA intervals identified by

the MACS3 peak-calling algorithm, which models background signal to detect significant

enrichment. **(B)** ChIP-qPCR analysis of H3K9me2 enrichment at the the GremLINE1

transposon (positive control region) and *fkbA* locus (targeting the 5′-UTR, exonic regions,

and 3′-UTR). Error bars represent mean ± SEM (n = 3). Statistical significance: ****$p \le$

0.0001. **(C, D)** ChIP-seq coverage mapped across the GremLINE1 transposon and its

neighboring loci (C) or *fkbA* and its neighboring loci (D). The peaks shown beneath the

coverage track denote significant H3K9me2-enriched intervals identified by the MACS3

peak-calling algorithm, which detects statistically enriched regions by modeling

background signal. **(E)** ChIP-qPCR analysis of RNA polymerase II enrichment at the

positive control region (*actA*), *fkbA* locus (targeting the 5′-UTR and exonic region), a *fkbA*-

neighboring gene, and negative control locus, defined as an siRNA-producing region

flanked by transposable elements and associated with H3K9me2. Statistical significance:

****$p \le 0.0001$. WT, wild-type; E1 and E2, epimutants; E1P32 and E2P28, revertants.

1028

Fig 6. Inherited heterochromatin-mediated antifungal resistance persists after *in*

***vivo* passage.**

(A) Fungal growth of *M. circinelloides*, *M. bainieri* and *M. atramentarius* under different

temperature conditions. Microscopic images show enlarged sporangia (scale bar =

20 μm). **(B)** Survival rates of mice infected with *M. atramentarius*. *Mucor circinelloides*

and PBS served as positive and negative controls, respectively. The log-rank test

revealed a statistically significant difference in survival between groups (*M. atramentarius*

vs PBS, *p* = 0.014). **(C)** Fungal colonies recovered from four different organs following *in*

vivo infection with wild-type (WT), epimutant (E1), and revertant (E1P32) strains. The bar

plot (right) shows the percentage of FK506-resistant colonies from each organ after

passage. **(D)** ChIP-qPCR analysis of H3K9me2 enrichment at the *fkbA* locus following *in*

vivo passage, targeting two exonic regions. Error bars represent mean ± SEM (n = 3).

Statistical significance: ****p ≤ 0.0001. WT, wild-type; E1R1–E1R3, FK506-resistant

epimutants; E1S1, FK506-sensitive epimutant; E1P32, revertant.

1043

Fig 7. Heterochromatin-mediated antifungal resistance remains stable after *in vivo*

infection in immunosuppressed and non-immunosuppressed mice.

(A−C) Fungal colonies recovered from four different organs following *in vivo* infection with

heterochromatin-mediated epimutant under three different conditions: (A)

cyclophosphamide-treated mice (50 mg/kg, 2 weeks post-infection), (B) untreated mice

(2 weeks post-infection), and (C) untreated mice (4 weeks post-infection). The right bar

plot shows the percentage of FK506-resistant colonies recovered from each organ under

each condition. **(D−F)** ChIP-qPCR analysis of H3K9me2 enrichment at the *fkbA* locus

under the same three conditions: (D) cyclophosphamide-treated mice (50 mg/kg, 2 weeks

post-infection), (E) untreated mice (2 weeks post-infection), and (F) untreated mice (4

weeks post-infection). Error bars represent mean ± SEM (n = 3). Statistical significance:

$^{**}p \leq 0.01$; $^{****}p \leq 0.0001$.

1056

1057 **Fig 8. Schematic overview of epigenetic genetic silencing contributing to antifungal**

1058 **resistance in *Mucor* species.**

1059 *fkbA* expression enables FKBP12 production and FK506 sensitivity (yeast-like

1060 morphology). Epigenetic silencing of *fkbA* via RNAi or heterochromatin leads to FK506

1061 resistance and hyphal growth. Generated with BioRender.com.

 Table 1. FK506 resistant strains isolated in this study.

Species	Name	Phenotype	*fkbA* mutation	Impact on FKBP12	Calcineurin mutation
PS1 *Mucor janssenii*	M1	FK506-R, Rapa-S	No	No	G1496C in *cnaA* (G347A missense mutation)
	M2	FK506-R, Rapa-R	4,324 bp insertion (Unknown) at 38 bp upstream *fkbA* ORF	-	-
	M3	FK506-R, Rapa-R	Large deletion	Loss	-
	M4	FK506-R, Rapa-R	Large deletion	Loss	-
	M5	FK506-R, Rapa-R	5,052 bp insertion (RC/Helitron_DNA transposons) at 16 bp upstream *fkbA* ORF	-	-
	M6	FK506-R, Rapa-R	C120 1 bp deletion in exon 2 (Frameshift)	Nonsense mutation	-
	M7	FK506-R, Rapa-R	Large deletion	Loss	-
	M8	FK506-R, Rapa-R	G258C in 5' splice site of intron2	-	-
	M9	FK506-R, Rapa-R	Large deletion	Loss	-
	M10	FK506-R, Rapa-R	G258C in 5' splice site of intron2	-	-
PS3 *Mucor bainieri*	E1	FK506-R, Rapa-R	No	No	No
	E2	FK506-R, Rapa-R	No	No	No
	E3	FK506-R, Rapa-R	No	No	No
	E4	FK506-R, Rapa-R	No	No	No
	E5	FK506-R, Rapa-R	No	No	No
	E6	FK506-R, Rapa-R	No	No	No
	M1	FK506-R, Rapa-R	C357 1 bp deletion in exon 3 (Frameshift)	Nonsense mutation	No
	E7	FK506-R, Rapa-R	No	No	No
	E8	FK506-R, Rapa-R	No	No	No
	E9	FK506-R, Rapa-R	No	No	No

	M1	FK506-R, Rapa-R	C404 1 bp deletion in exon 3 (Frameshift)	Nonsense mutation	No
	M2	FK506-R, Rapa-S	No	No	G477T in *cnbR* (V122F missense mutation)
	M3	FK506-R, Rapa-R	G249A in 5′ splice site	No	No
	M4	FK506-R, Rapa-R	G249A in 5′ splice site	No	No
PS6 *Mucor atramentarius*	M5	FK506-R, Rapa-S	G189T in exon 2	Synonymous mutation	No
	M6	FK506-R, Rapa-R	C404 1 bp deletion in exon 3 (Frameshift)	Nonsense mutation	No
	M7	FK506-R, Rapa-R	T2C in exon 1 (start-loss mutation)	Loss	No
	E1	FK506-R, Rapa-R	No	No	No
	E2	FK506-R, Rapa-R	No	No	No
	M8	FK506-R, Rapa-R	C404 1 bp deletion in exon 3 (Frameshift)	Nonsense mutation	No

1063

1064 **Table 2. Epimutant reverted strains analyzed in this study.**

Species	Name	Background	Phenotype	Description
PS3 *Mucor bainieri*	E1P4	E1	FK506-S, Rapa-S	E1 Revertant after passage 4
	E2P4	E2	FK506-S, Rapa-S	E2 Revertant after passage 4
	E3P12	E3	FK506-S, Rapa-S	E3 Revertant after passage 12
	E3P16	E4	FK506-S, Rapa-S	E4 Revertant after passage 16
	E5P12	E5	FK506-S, Rapa-S	E5 Revertant after passage 12
	E6P12	E6	FK506-S, Rapa-S	E6 Revertant after passage 12
	E7P28	E7	FK506-S, Rapa-S	E7 Revertant after passage 28
	E8P28	E8	FK506-S, Rapa-S	E8 Revertant after passage 28
	E9P24	E9	FK506-S, Rapa-S	E9 Revertant after passage 24
PS6 *Mucor atramentarius*	E1P32	E1	FK506-S, Rapa-S	E1 Revertant after passage 32
	E2P28	E2	FK506-S, Rapa-S	E2 Revertant after passage 28

1065

1066 **Table 3. Summary of epigenetic mechanisms in *Mucor bainieri*.**

Type	Mechanisms	Epimutant	Revertant	Average passages for revertant
1	RNAi-associated heterochromatin-dependent silencing	E1	E1P4	6.67
		E2	E2P4	
		E3	E3P12	
2	Heterochromatin-dependent silencing	E4	E3P16	16
3	Canonical RNAi-dependent silencing	E5	E5P12	18.8
		E6	E6P12	
		E7	E7P28	
		E8	E8P28	
		E9	E9P24	

1067

A

B

| PS1 *Mucor janssenii* CBS185.68 | PS3 *Mucor bainieri* CBS293.63 | PS6 *Mucor atramentarius* CBS202.28 |

A PS3 *Mucor bainieri* CBS293.63

B *fkbA* mRNA levels

C FKBP12 protein levels

D PS6 *Mucor atramentarius* CBS202.28

E *fkbA* mRNA levels

F FKBP12 protein levels