Title

Evidence for the acquisition of a proteorhodopsin-like rhodopsin by a chrysophyte-infecting

giant virus

Authors

Petra Byl*[1, 2, 3], Christopher R. Schvarcz[1, 2], Julie Thomy[1, 2], Qian Li[1, 2], Cori B. Williams[1], Kurt

LaButti[3], Frederik Schulz[3], Kyle F. Edwards[2], Grieg F. Steward*[1, 2]

Affiliations

[1]Daniel K. Inouye Center for Microbial Oceanography: Research and Education, School of

Ocean and Earth Science and Technology (SOEST), University of Hawaiʻi Mānoa, 1950 East-

West Road, Honolulu, HI 96822.

[2]Department of Oceanography, School of Ocean and Earth Science and Technology (SOEST),

University of Hawaiʻi Mānoa, 1000 Pope Road, Honolulu, HI 96822.

[3]DOE Joint Genome Institute, Lawrence Berkeley National Laboratory, 1 Cyclotron Road,

Berkeley, CA 94720.

*Corresponding author(s): pkbyl@hawaii.edu; grieg@hawaii.edu

Abstract

Abstract

Chrysophytes are widespread nanoflagellate protists in aquatic ecosystems with diverse trophic

strategies ranging from phototrophy to heterotrophy. Here, we report the first isolation of a

chrysophyte-infecting virus, Chrysophyceae clade-H virus SA1, referred to hereafter as

ChrysoHV, on a sympatric mixotrophic chrysophyte. Both host and virus were isolated from

surface waters in the subtropical North Pacific. The ChrysoHV capsid (291 ± 42 nm diameter) is

associated with a loose, sac-like membrane that extends its effective diameter (716 ± 115 nm),

and presents a long, thin tail extending 1,202 (± 243) nm. This unique morphology has not been

previously observed in other isolates or environmental surveys. The sequenced, assembled

genome is 1.19 Mbp, and phylogenetic analysis places ChrysoHV as the third cultivated member

of the *Aliimimivirinae* subfamily in the *Mimiviridae* family of giant viruses. ChrysoHV is the

first cultivated member of *Aliimimivirinae* to encode a rhodopsin, which has high phylogenetic

relatedness and structural similarity to proteorhodopsin sequences retrieved from the same

subtropical research site. The most closely related sequences from cultivated organisms derive

from bacteria in the genus *Pelagibacter*. This suggests that ChrysoHV acquired the

proteorhodopsin-like protein through lateral gene transfer from the chrysophyte's bacterial prey.

Key words

Nucleocytoviricota, horizontal gene transfer (HGT), mixotrophy

Main text

Giant viruses are large double-stranded DNA viruses in the phylum *Nucleocytoviricota* [1]. They display extensive genomic repertoires, including viral homologs of cellular genes, among which rhodopsin homologs are commonly found [2]. These rhodopsin homologs could augment or compensate for host functions compromised during infection [3-5] or interfere with host defenses [6]. Viral rhodopsin homologs have been experimentally confirmed to convert light energy into electrochemical potential across multiple clades: viral rhodopsin groups I and II [3, 7, 8], channelrhodopsin [4], and heliorhodopsin [6]. Furthermore, these rhodopsin homologs are globally distributed in marine habitats, and their abundance decreases over depth with the attenuation of sunlight [3].

Although giant virus rhodopsin homologs are widespread and potentially functional, only a few rhodopsin-encoding giant viruses have been isolated and remain in culture with their native host [5, 6, 9]. Here, we report the isolation of the giant virus ChrysoHV (Fig. 1A-E), which encodes a proteorhodopsin-like protein (Fig. 2A). ChrysoHV was isolated with a co-occurring mixotrophic chrysophyte from an open ocean research site in the North Pacific Subtropical Gyre (Station ALOHA: 22° 45' N, 158° W; Supplementary Information).

We tested the host range of ChrysoHV against eight protist isolates from Stn. ALOHA, resulting in no lysis for six of the eight isolates (Supplementary Fig. S1, S2A,C; Supplementary Information). Both Chrysophyceae clade-H representatives UHM3500 and UHM3501 lysed when challenged with ChrysoHV, but a greater fraction of UHM3501 cells lysed and more virions were produced (Supplementary Fig. S2B), even though virions produced per cell (burst size) were similar for each host strain (22.0-25.5). Lower mortality for UHM3500 may be due to resistant subpopulations, host immune responses, or virophage integrations, which are

widespread in uncultivated members of Chrysophyceae clade-H [10]. The virion particles

produced from the two chrysophytes shared the same notable morphological features, *i.e.*, an

icosahedral capsid (291 ± 42 nm diameter [mean ± SD]) associated with a loose membrane (716

± 115 nm), and a long tail (1,202 ± 243 nm length) ending in tail fibers (206 ± 72 nm length)

(Fig. 1A-C).

The ChrysoHV genome was fully assembled as a putatively linear DNA sequence of 1.19 Mbp

encoding 1,138 predicted genes with a G+C content of 23.42% (Fig. 1B; Supplementary

Information). The ChrysoHV genome encodes genes of diverse putative function, including the

ribosomal proteins eL40 and S27a, making it the second known eukaryote-infecting virus to

encode a ribosomal protein (Supplementary Table S1) [5]. ChrysoHV also encodes cytochrome

b5 and eight other genes involved with Fe use or transport (Supplementary Table S1). Notably,

the genome includes a region of light-sensitive proteins, encompassing a proteorhodopsin-like

protein surrounded by two photolyases (Supplementary Table S1, Fig. 1D, Fig. 2C). The

positioning and shared orientation of the photolyases around the proteorhodopsin-like protein

implies that it may sit on the same operon and function like the heliorhodopsin in *Trichococcus*

flocculiformis, which was proposed to bind to nearby photolyases to enhance DNA repair [11].

ChrysoHV is the third cultivated isolate in the *Aliimimivirinae* subfamily of *Mimiviridae* (Fig.

1E); other known isolates include CroV BV-PW1 [12], infecting a heterotrophic nanoflagellate,

and CV-XW01 [13], infecting a green alga. In this subfamily only two other representatives,

cPacV-1605 and GVMAG-M-3300013131-43, encode rhodopsin, which belong to the viral

rhodopsin group I clade (Fig. 2A). However, 40 other *Aliimimivirinae* representatives encode at

least one photolyase homolog (Fig. 1E) in the (6-4) photolyase or DASH cryptochrome clades.

Only ChrysoHV and cPacV-1605 encode both rhodopsin and photolyase homologs, but the

genome architecture of cPacV-1605 places the rhodopsin (CDS 205) away from the photolyase

cluster (CDS 131, 134) [14]. Given the mismatched distribution of rhodopsin and photolyase

homologs within the *Aliimimivirinae* subfamily, the ChrysoHV proteorhodopsin-like protein may

represent a unique adaptation to high light in the surface waters of the subtropical ocean.

Phylogenetic analysis of ChrysoHV's rhodopsin placed it within the proteorhodopsin clade (Fig.

2A), and ChrysoHV may have acquired the proteorhodopsin-like rhodopsin from the

chrysophyte host's putative bacterial prey, *Pelagibacter* (Fig. 2A). The sequence alignment of

transmembrane domain helix-III (TM3) indicates that the ChrysoHV rhodopsin is spectrally

tuned to blue-light (glutamine, Q; Fig. 2A), which is the dominant spectral tuning amino acid in

proteorhodopsins from Stn. ALOHA at 25-200 m depth [15]. While ChrysoHV lacks the

conserved DTE proton pumping motif found in most proteorhodopsins, it possesses an ETL

motif (Fig. 2A), which has been detected in proteorhodopsins from deep water at Stn. ALOHA

(Fig. 2A) [15]. Compared to rhodopsins with existing functional characterization, the predicted

structure of the ChrysoHV rhodopsin was most similar to the blue-light absorbing

proteorhodopsin from uncultured Gamma-proteobacterium clone HOT 75m4 (Fig. 2B)

confirmed to have ion (H^+) pumping functionality [16]. Although the amino acid identity was not

the highest (35.9%), this HOT 75m4 proteorhodopsin was also retrieved from Stn. ALOHA [16].

ChrysoHV does not encode β-carotene, precursor to retinal, the light-absorbing chromophore in

proteorhodopsin, but the retinal binding site (lysine, K) in transmembrane domain helix VII

(TM7) was conserved (Fig. 2A). Further investigation is needed to confirm whether the

ChrysoHV proteorhodopsin-like protein acts as an ion pump.

Analysis of phylotypes related to ChrysoHV (≥ 90% polB amino acid identity) from the TARA

Oceans survey [17] revealed greatest prevalence in the subtropical surface ocean (Supplementary

Fig. S3). In this environment, diverse microorganisms from proteobacteria to microalgae utilize

rhodopsins to supplement their energetic needs [15, 18]. SAR11 bacteria, of which *Pelagibacter*

is an abundant subclade, account for 25% of all plankton [19], and are a likely prey source for

mixotrophic algae such as Chrysophyceae clade-H sp. UHM3500 and UHM3501. The discovery

of a proteorhodopsin-like protein encoded by a giant virus supports the conceptual model that

horizontal gene transfer is driven by the co-mingling of infectious virions and ingested prey

within the same host cell [20]. By leveraging naturally co-occurring and experimentally

accessible virus-host pairs, this study lays the groundwork for future investigations into how the

virocell environment of predatory protists influences gene acquisition in giant viruses.

Figures

Figure 1. (A) Transmission electron microscopy (TEM) negative stain micrograph of ChrysoHV

propagated on UHM3501. (B, C) TEM negative stain micrograph of ChrysoHV propagated on

UHM3500. (D) Genome plot of ChrysoHV with track placement and color indicating gene

lineage by domain: inner to outer tracks show prokaryote (blue), eukaryote (yellow), and virus

(black). Phylogenetically informative giant virus marker genes are represented as black dots on

the outside track and show gene name. The genes used in the *Nucleocytoviricota* phylogenetic

analysis are bolded. (E) Maximum Likelihood (ML) phylogeny of the *Nucleocytoviricota*

phylum based on a concatenated alignment of translated marker genes (*TopoII*, *RNAPL*, *TFIIB*,

VLTFB, *A32*, *PolB* and *SFII*) using model LG+I+F+G4 with ultrafast bootstrap supports (-bb

1000) shown by node symbol color– white (≥ 80%), gray (≥ 90%), and black (100%) with no

node symbol for supports < 80%. The tree is rooted with *Chitovirales* as the outgroup, and the

scale bar represents one amino acid substitution per site. The frequency of light-sensitive

proteins such as rhodopsin and photolyases (phrB) and (DRPR) present in the *Aliimimivirinae*

subfamily are annotated in the grayscale matrix and the environment the virus was collected

from is shown in jade (freshwater), blue (marine), and brown (terrestrial).

Figure 2. (A) Microbial rhodopsin ML phylogeny with heliorhodopsin as the outgroup, built

with model LG+F+R10. Select representatives of viral rhodopsin homologs and putative

chrysophyte rhodopsins are referred to next to their corresponding collapsed clade. The

proteorhodopsin (PR) clade is annotated with the alignment of transmembrane helix 3 (TM3)

outlining the ion-pumping motif (gray) and the spectral tuning amino acid (green or blue).

Residues that differ from the most common residue at that locus are in bold (B). The predicted

structure of ChrysoHV's proteorhodopsin-like protein conserved domain (yellow) and the HOT

75m4 Gamma-proteobacterium proteorhodopsin (blue). (C) Gene neighborhood of light-

sensitive proteins in ChrysoHV, including the proteorhodopsin (CDS 97) between the two

photolyases (CDS 91 and 100).

Acknowledgments

We thank the personnel of the Hawaiʻi Ocean Time-series program (NSF award 12-60164) and

Research Vessel Kilo Moana for assistance with seawater collection. We thank the Biological

Electron Microscope Facility (BEMF) at the University of Hawaiʻi Mānoa, especially Facility

Supervisors, Tina Weatherby and Dr. Orion Rivers, for their assistance with electron microscopy

procedures. A special thanks to US DOE Joint Genome Institute Scientific Project Manager,

Kerrie Barry, for overseeing DNA library preparation, quality control, and sequencing of

ChrysoHV.

We are grateful for the thoughtful feedback provided by Dr. Karen Selph (Department of

Oceanography, University of Hawaiʻi Mānoa), and Dr. Alison Sherwood (School of Life

Sciences, University of Hawaiʻi Mānoa).

Funding

This work was supported by the US National Science Foundation Graduate Research Fellowship

Program (Grant No. 1842402 to PB), Division of Ocean Sciences (Grant No. 2129697 to GFS

and KFE), and RII Track-2 FEC (Grant No. 1736030 to GFS and KFE). The work conducted by

the US Department of Energy Joint Genome Institute (https://ror.org/04xm1d337), a Department

of Energy (DOE) Office of Science User Facility, was supported by the Office of Science of the

US DOE, operated under Contract No. DE-AC02-05CH11231.

Conflicts of Interest

The authors declare no conflicts of interest.

Data Availability

ChrysoHV sequencing data is available through JGI IMG under Sequencing Project ID 1432190,

and the genome assembly has been deposited to NCBI GenBank under accession ID PV764857.

Additional data and scripts on gene alignments, phylogenies, genome annotation, protein

structural predictions, host range challenges, and biogeography are available in the Dryad

repository:

http://datadryad.org/share/8duyiXJ3riDLgLbseVaWvUBnbOPOM4SkBw9WNBaRYA4. The

virus and protist strains used in this study are available through the Marine Viral Ecology

Laboratories at the University of Hawaiʻi Mānoa upon request to KFE and GFS.

References

1. Aylward, F. O. *et al.* Taxonomic update for giant viruses in the order Imitervirales (phylum

Nucleocytoviricota). *Arch. Virol.* **168**, 283 (2023). 10.1371/journal.pbio.3001430

2. Moniruzzaman, M. *et al.* Virologs, viral mimicry, and virocell metabolism: the expanding

scale of cellular functions encoded in the complex genomes of giant viruses. *FEMS Microbiol.*

Rev. **47**, fuad053 (2023). 10.1093/femsre/fuad053

3. Needham, D. M. *et al.* A distinct lineage of giant viruses brings a rhodopsin photosystem to

unicellular marine predators. *Proc. Natl. Acad. Sci.* **116**, 20574–20583 (2019).

10.1073/pnas.1907517116

4. Rozenberg, A. *et al.* Lateral Gene Transfer of Anion-Conducting Channelrhodopsins between

Green Algae and Giant Viruses. *Curr. Biol.* **30**, 4910-4920.e5 (2020). 10.1016/j.cub.2020.09.056

5. Thomy, J., Schvarcz, C. R., McBeain, K. A., Edwards, K. F. & Steward, G. F. Eukaryotic

viruses encode the ribosomal protein eL40. *Npj Viruses* **2**, 51 (2024).

6. Hososhima, S. *et al.* Proton-transporting heliorhodopsins from marine giant viruses. *eLife* **11**,

e78416 (2022). 10.1038/s44298-024-00060-2

7. Bratanov, D. *et al.* Unique structure and function of viral rhodopsins. *Nat. Commun.* **10**, 4939

(2019). 10.1038/s41467-019-12718-0

8. Zabelskii, D. *et al.* Viral rhodopsins 1 are an unique family of light-gated cation channels. *Nat.*

Commun. **11**, 5707 (2020). 10.1038/s41467-020-19457-7

9. Brussaard, C. P. D., Bratbak, G., Baudoux, A. -C. & Ruardij, P. Phaeocystis and its interaction

with viruses. *Biogeochemistry* **83**, 201–215 (2007). 10.1007/s10533-007-9096-0

10. Castillo, Y. M. *et al.* Assessing the viral content of uncultured picoeukaryotes in the global-

ocean by single cell genomics. *Mol. Ecol.* **28**, 4272–4289 (2019). 10.1111/mec.15210

11. Shim, J. *et al.* Heliorhodopsin Helps Photolyase to Enhance the DNA Repair Capacity.

Microbiol. Spectr. **10**, e02215-22 (2022). 10.1128/spectrum.02215-22

12. Fischer, M. G., Allen, M. J., Wilson, W. H. & Suttle, C. A. Giant virus with a remarkable

complement of genes infects marine zooplankton. *Proc. Natl. Acad. Sci.* **107**, 19508–19513

(2010). 10.1073/pnas.1007615107

13. Sheng, Y., Wu, Z., Xu, S. & Wang, Y. Isolation and Identification of a Large Green Alga

Virus (*Chlorella* Virus XW01) of *Mimiviridae* and Its Virophage (*Chlorella* Virus Virophage

SW01) by Using Unicellular Green Algal Cultures. *J. Virol.* **96**, e02114-21 (2022).

10.1128/jvi.02114-21

14. Needham, D. M. *et al.* Targeted metagenomic recovery of four divergent viruses reveals

shared and distinctive characteristics of giant viruses of marine eukaryotes. *Philos. Trans. R. Soc.*

B Biol. Sci. **374**, 20190086 (2019). 10.1098/rstb.2019.0086

15. Olson, D. K., Yoshizawa, S., Boeuf, D., Iwasaki, W. & DeLong, E. F. Proteorhodopsin

variability and distribution in the North Pacific Subtropical Gyre. *ISME J.* **12**, 1047–1060

(2018). 10.1038/s41396-018-0074-4

16. Ran, T., *et al*. 2013. Cross-protomer interaction with the photoactive site in oligometric

proteorhodopsin complexes. *Acta Crystallogr* **69**, 1965–1980 (2013).

10.1107/S0907444913017575

17. Endo, H. *et al.* Biogeography of marine giant viruses reveals their interplay with eukaryotes

and ecological functions. *Nat. Ecol. Evol.* **4**, 1639–1649 (2020). 10.1038/s41559-020-01288-w

18. Strauss, J. *et al.* Plastid-localized xanthorhodopsin increases diatom biomass and ecosystem

productivity in iron-limited surface oceans. *Nat. Microbiol.* **8**, 2050–2066 (2023).

10.1038/s41564-023-01498-5

19. Giovannoni, S. J. SAR11 Bacteria: The Most Abundant Plankton in the Oceans. *Annu. Rev.*

Mar. Sci. **9**, 231–255 (2017). 10.1146/annurev-marine-010814-015934

20. Filée, J., Siguier, P. & Chandler, M. I am what I eat and I eat what I am: acquisition of

bacterial genes by giant viruses. *Trends Genet.* **23**, 10–15 (2007). 10.1016/j.tig.2006.11.002