Evidence for the acquisition of a proteorhodopsin-like rhodopsin by a chrysophyte-infecting

giant virus

Petra Byl[#, 1, 2, 3, *], Christopher R. Schvarcz[2, 3], Julie Thomy[2, 3], Qian Li[2, 3], Cori B. Williams[2],

Kurt LaButti[4], Frederik Schulz[4], Kyle F. Edwards[3], Grieg F. Steward[#, 2, 3]

[1]Daniel K. Inouye Center for Microbial Oceanography: Research and Education, School of

Ocean and Earth Science and Technology (SOEST), University of Hawai'i at Mānoa, Honolulu,

HI, USA.

[2]Department of Oceanography, School of Ocean and Earth Science and Technology (SOEST),

University of Hawai'i at Mānoa, Honolulu, HI, USA.

[3]DOE Joint Genome Institute, Lawrence Berkeley National Laboratory, Berkeley, CA. USA.

Running Head: Proteorhodopsin encoded by chrysophyte virus

#Address correspondence to Petra Byl pbyl@umces.edu & Grieg Steward grieg@hawaii.edu

*Present address: University of Maryland Center for Environmental Science, Horn Point

Laboratory, Cambridge, MD, USA.

Abstract word count: 239

Importance word count: 144

Main text word count: 3,619

**Abstract**

Chrysophytes are widespread nanoflagellate protists in aquatic ecosystems with diverse trophic

roles as primary producers and bacterivores. While molecular evidence suggests that

chrysophytes are commonly infected by giant viruses, no previous isolates have been reported.

Here, we describe the first isolated chrysophyte-infecting virus, Chrysophyceae Clade H virus

SA1 (ChrysoHV). ChrysoHV and its mixotrophic host were isolated from surface waters in the

tropical North Pacific. The ChrysoHV capsid ($290 \pm 40$ nm diameter) is associated with a loose,

sac-like membrane that extends its effective diameter ($720 \pm 120$ nm), and presents a long, thin

tail extending $1{,}200$ ($\pm 240$) nm in length. This morphology has not been previously observed in

other virus isolates or environmental surveys. The sequenced, assembled genome is 1.19 Mbp,

and phylogenetic analysis places ChrysoHV as the third cultivated member of the

*Aliimimivirinae* subfamily in the *Mimiviridae* family of giant viruses. ChrysoHV is the first

cultivated member of *Aliimimivirinae* to encode a rhodopsin. Moreover, ChrysoHV encodes

three rhodopsins total, two heliorhodopsins and one proteorhodopsin. No other virus-derived

rhodopsin has been found in the proteorhodopsin clade, and the loss of retinal binding hints at a

light-independent function. ChrysoHV also encodes nine genes with closest known homologs

from marine cyanobacteria, mostly related to nutrient uptake. This suggests that lateral gene

transfer occurs between abundant marine bacteria and giant viruses, supporting the conceptual

model that genes are horizontally transferred between ingested prey and infectious virions within

predatory protists.

**Importance**

Chrysophytes are abundant eukaryotic phytoplankton with diverse trophic strategies ranging from photosynthesis to phagotrophy, and have been studied as models of mixotrophy among marine protists. However, no chrysophyte-infecting viruses had been cultivated, leaving a major gap in our understanding of viral diversity. This study reports the functional and genomic characterization of the first isolated chrysophyte-infecting virus, the giant virus ChrysoHV. The virion morphology is unusual, with a loosely associated outer membrane and a long thin tail. The viral genome encodes three rhodopsins, one of which is affiliated with a bacterial proteorhodopsin lineage not previously identified in viruses. ChrysoHV also encodes genes with closest homologs in marine cyanobacteria, suggesting that phagocytic protists may provide a conduit for horizontal gene transfer from bacterial prey to viral pathogens. This isolate offers a useful model for exploring giant virus genomics and the function of a new viral rhodopsin.

**Main text**

**Introduction**

Chrysophytes are single-celled or colonial protists in the class Chrysophyceae, and they are globally abundant in freshwater and marine ecosystems (1, 2). Chrysophytes play multiple ecosystem roles as primary producers and bacterivores, and include phago-mixotrophic forms that perform photosynthesis and ingest prey, strictly heterotrophic phagotrophs, and potentially non-phagotrophic phototrophs (3). In subtropical ocean gyre ecosystems, which span approximately a quarter of the global surface ocean (4), chrysophytes play an important role as predators of marine cyanobacteria (5, 6), influencing the community structure at the base of the food web. Isolates from Chrysophyceae Clade H have been shown to be voracious consumers of

the cyanobacterium *Prochlorococcus* (7), the dominant primary producer in subtropical gyres

and most abundant phototroph in the ocean (8). Additionally, single cell amplified genomes of

Chrysophyceae Clade H have been shown to encode endogenous viral elements from the

*Nucleocytoviricota* phylum (9), referred to as giant viruses (10). Genetic marker surveys of

protist and giant virus distributions from the *Tara* Oceans global dataset also indicate that

chrysophytes commonly co-occur with the giant virus family *Mimiviridae* (11), and further

RNA-seq studies revealed that chrysophytes are in fact native hosts of mimivirids (12).

However, chrysophyte-infecting giant virus isolates have not been previously described. In this

study we detail the isolation and initial characterization of a giant virus that infects a Clade H

chrysophyte, ChrysoHV.

Giant viruses encode vast genomic repertoires that include numerous homologs of

cellular genes (13). These cellular homologs may augment or compensate for host functions

compromised during infection (14–16), or interfere with host defenses (17). Among the most

commonly found viral homologs of cellular genes are microbial rhodopsins (18–20), which are

photoreceptive proteins that may aid in the generation of ion gradients or intracellular signaling

(21, 22). Viral rhodopsin homologs have been experimentally confirmed to convert light energy

into electrochemical potential across multiple clades: viral rhodopsin groups I and II (14, 23, 24),

channelrhodopsin (15), and heliorhodopsin (17). Furthermore, these viral rhodopsin homologs

are globally distributed across aquatic habitats, with their relative abundance decreasing over

depth with the attenuation of sunlight (14, 25).

Although giant viruses are diverse and globally abundant (11, 19, 20), a relatively small

fraction of their known diversity is represented by cultivated isolates (26). Likewise, giant virus

rhodopsin homologs are widespread and potentially functional, but only a few rhodopsin-

encoding giant viruses have been isolated and remain in culture with their native host (16, 17,

27). Cultivated isolates are important for establishing virus-host linkages, and furthermore, can

reveal novel gene lineages that may go undetected in environmental datasets and clarify the

evolutionary history of highly divergent sequences (16). Through the isolation and

characterization of ChrysoHV, we identify a previously undescribed lineage of viral rhodopsin

that appears to derive from bacterial proteorhodopsin.

**Results & Discussion**

*Host range, infection dynamics, and morphology*

We tested the host range of ChrysoHV against seven protist isolates from Station

ALOHA, spanning three classes: three Chrysophyceae, three Dictyochophyceae, and one

Bolidophyceae representative. This resulted in no lysis for six of the seven isolates (Fig. 1B)

(Supplementary Fig. S1). Only the representative of chrysophyte Clade H, lysed when

challenged with ChrysoHV (Fig. 1). The apparent burst size, *i.e.*, estimated number of virions

produced per cell, was $24 \pm 2$ ([mean $\pm$ SD]) (Fig. 1A). The virion particles produced from the

infection of chrysophyte strain UHM3501 displayed the notable morphological features of an

icosahedral capsid ($290 \pm 40$ nm diameter [mean $\pm$ SD]) associated with a loose membrane (720

$\pm 120$ nm), and a long tail ($1,200 \pm 240$ nm length) terminating in tail fibers ($210 \pm 700$ nm

length) (Fig. 2A-C).

*Genomic characterization*

The ChrysoHV genome was assembled as a putatively linear DNA sequence of 1.19 Mbp

encoding 1,138 predicted genes with a G+C content of 23.42% (Fig. 2B). The encoded genes

have diverse putative functions, including the ribosomal proteins eL40 and S27a, making it the

second isolated eukaryote-infecting virus known to encode ribosomal proteins (21), and the first

to encode S27a (Supplementary Table S1). ChrysoHV also encodes electron transport proteins

such as cytochrome b5, widespread in prokaryotes, eukaryotes, and giant viruses (Supplementary

Table S1) (28), along with succinate dehydrogenase flavoprotein and iron-sulfur subunits

(Supplementary Table S1). Additionally, the virus encodes seven ABC transporters including

one specific to phosphate transport (XVM28869.1) and two AmtB ammonium transporters. Of

these ammonium transporters, one (XVM28992.1) has 55% amino acid sequence identity (aaID)

to *Prochlorococcus* and *Synechococcus* homologs (Supplementary Table S1). Likewise,

ChrysoHV encodes three porins with 44-66% aaID to *Prochlorococcus* iron and carbohydrate

porins (XVM28452.1, XVM28569.1, XVM28570.1), two urea ABC transporter substrate-

binding proteins with 81-90% aaID to a *Prochlorococcus* homolog (XVM29005.1,

XVM29006.1), a semiSWEET sugar transporter with 44% aaID to a *Prochlorococcus* homolog

(XVM28124.1), a sulfotransferase family protein with 55% aaID to a *Prochlorococcus* homolog

(XVM28964.1), and a N-acetylmuramoyl-L-alanine amidase for cell wall remodeling with 39%

aaID to a *Prochlorococcus* homolog (XVM29061.1) (Supplementary Table S1). The presence of

genes whose closest known relatives are *Prochlorococcus* homologs suggests extensive

horizontal gene transfer from cyanobacterial prey of the chrysophyte to a giant virus that infects

it.

The ChrysoHV genome also includes a region of putatively light-sensitive proteins,

encompassing a proteorhodopsin-like protein surrounded by two photolyases and followed by a

heliorhodopsin (Fig. 2D, Fig. 3C, Supplementary Table S1). The positioning and shared

orientation of the photolyases and heliorhodopsin around the proteorhodopsin-like protein

implies that they may be part of the same operon.

*Sequence characterization of a distinct lineage of proteorhodopsin*

Phylogenetic analysis of ChrysoHV's rhodopsin placed it within the proteorhodopsin

clade (Fig. 3A). The sequence alignment of transmembrane domain helix-III (TM3) indicates

that the ChrysoHV rhodopsin is spectrally tuned to blue light (glutamine, Q; Fig. 3A), which is

the dominant spectral tuning residue in proteorhodopsins from Station ALOHA, the isolation site

of ChrysoHV (29). The two most closely related rhodopsin sequences were recovered from

metagenomes from the Black Sea at 50 m depth (SAMN27960419) and the North Pacific at 5 m

depth in the 0.22-3µm size fraction (SAMEA2623059). These rhodopsins may also absorb blue

light (Fig. 3A) and share a poly-N (asparagine) tract at the start of the rhodopsin coding

sequence.

ChrysoHV does not encode β-carotene, a precursor to retinal, the light-absorbing

chromophore in proteorhodopsin, and the salt bridge between K102 and E228 most likely

inhibits retinal binding. This loss of retinal binding, if associated with a change in rhodopsin

function, could explain the high rate of sequence divergence of ChrysoHV's proteorhodopsin

compared to the rest of the proteorhodopsin clade (Fig. 3A). Understanding how the rhodopsin

may function without retinal binding could be important to interpreting the function of viral

rhodopsins detected below the photic zone (30, 31). Further investigation is needed to confirm

whether the ChrysoHV proteorhodopsin can confer other physiological benefits in the absence of

light absorption, similar to the *Psychroflexus torquis* proteorhodopsin ptqPR, which acts as a

phospholipid scramblase to preserve membrane potential, fluidity, and integrity (32).

ChrysoHV also lacks the conserved DTE proton pumping motif found in most

proteorhodopsins, and instead, possesses an ETL motif (Fig. 3A), which has been detected in

proteorhodopsins from deep water at Station ALOHA (Fig. 3A) (29). A comparison of the

predicted structure of the conserved domain of the ChrysoHV rhodopsin against existing

functional characterizations revealed that it was most similar (35.9% aaID) to the uncultured

gamma-proteobacterium clone HOT 75m4 rhodopsin (Fig. 3B). This rhodopsin was confirmed to

pump protons ($H^+$) (33), and, like ChrysoHV, was collected from Station ALOHA (33). These

results suggest that ChrysoHV may have acquired the proteorhodopsin from a marine bacterium,

which, like *Prochlorococcus*, may be prey for chrysophyte Clade H.

*Comparison to photoreceptive proteins across Nucleocytoviricota*

ChrysoHV is the third cultivated isolate in the *Aliimimivirinae* subfamily of *Mimiviridae*

(Fig. 2E); other known isolates include CroV BV-PW1 (34), infecting a heterotrophic

nanoflagellate, and CV-XW01 (35), infecting a green alga. In this subfamily, only two other

representatives, cPacV-1605 and GVMAG-M-3300023174-43, encode rhodopsins, which belong

to viral rhodopsin group II (Fig. 2A). However, 40 other *Aliimimivirinae* representatives encode

at least one photolyase homolog (Fig. 2E), either the (6-4) photolyase or DASH-type

cryptochrome. Only ChrysoHV and cPacV-1605 encode multiple rhodopsin and photolyase

types, including two heliorhodopsins (Fig. 3A). However, cPacV-1605 encodes a viral rhodopsin

(group II) and ChrysoHV encodes a proteorhodopsin (Fig. 3A). Furthermore, instead of being

nested within a photoreceptive gene neighborhood, the position of the viral rhodopsin in cPacV-

1605 (locus tag: CPAV1605_207, GenBank ID: VVU94485.1) is further away from the

photolyases (locus tag: CPAV1605_133, CPAV1605_136; GenBank ID: VVU94411.1,

VVU94414.1) and heliorhodopsins (locus tag: CPAV1605_533, CPAV1605_1309; GenBank ID:

VVU95557.1, VVU94808.1) (36). ChrysoHV is the only cultivated giant virus to encode the

proteorhodopsin lineage of rhodopsin along with two heliorhodopsins (Fig 2E, Fig. 3A).

*Global ocean distribution of ChrysoHV phylotypes and proteorhodopsins*

Analysis of the distribution of ChrysoHV phylotypes ($\geq$ 90% PolB aaID) from the *Tara*

Oceans dataset (11) revealed their prevalence in subtropical surface ocean sites in the Pacific and

Indian Ocean as well as the Mediterranean Sea (Fig. 4A). Notably, at station 132, north of the

Hawaiian archipelago, ChrysoHV phylotypes were detected in both the mixed layer and

mesopelagic zone, suggesting connectivity between surface and deep ocean virus populations,

which has previously been observed in the tropical North Pacific (25). The abundance of the

ChrysoHV proteorhodopsin-like protein ($\geq$ 80% PolB aaID) was also highest in the subtropical

surface ocean (Fig. 4B), but detected at fewer overall sites than the ChrysoHV PolB.

Additionally, the ChrysoHV proteorhodopsin relative abundance was highest at station 132,

indicating that these proteorhodopsin sequences may be derived from a ChrysoHV phylotype.

**Conclusion**

In the marine environment, diverse microorganisms from proteobacteria to microalgae

utilize rhodopsins to supplement their energetic needs (29, 37). SAR11 bacteria, of which

*Pelagibacter* is an abundant subclade, account for 25% of all plankton (38), and are a likely prey

source for mixotrophic algae such as Clade H chrysophytes. The discovery of a proteorhodopsin-

like protein encoded by a giant virus supports the conceptual model that lateral gene transfer is

driven by the co-mingling of infectious virions and ingested prey within the same host cell (39).

The presence of nine genes whose closest known relatives are *Prochlorococcus* homologs further

suggests that prey-virus HGT has played an important role in the evolution of ChrysoHV. By

leveraging naturally co-occurring and experimentally accessible virus-host pairs, this study lays

the groundwork for future investigations into how the virocell environment of predatory protists

influences gene acquisition in giant viruses. As the only cultivated isolate known to encode one

microbial rhodopsin and two heliorhodopsins, this system provides the opportunity for novel

insights into the role of rhodopsins in giant virus infection strategies.

**Materials and Methods**

*Isolation and cultivation*

The five mixotrophic protists used in this study include one chrysophyte (UHM3501),

one bolidophyte (UHM3600), and three dictyochophytes (UHM3020, UHM3053, UHM3071),

all of which were isolated from the epipelagic zone (5–100m) of Station ALOHA through serial

dilutions of whole seawater samples enriched with *Prochlorococcus* (7). The mixotrophs were

cultivated at 24°C at approximately 200–220 µmol photons $m^{-2}$ $s^{-1}$ irradiance to mimic the high

light conditions at station ALOHA (40). The unialgal cultures were grown on K medium without

dissolved, inorganic nitrogen sources ($NaNO_3$ and $NH_4Cl$). Cultures were instead amended with

cyanobacteria prey (*Prochlorococcus* MIT9301) for their nitrogen source (7, 41). In addition,

two heterotrophic chrysophytes (UHM3530 and UHM3560) were used in this study. These

strains were isolated from seawater collected from the mixed layer (5–50 m) at Station ALOHA

in 2022 and 2024 through serial dilutions of seawater samples. The heterotrophs were cultivated

on filtered seawater amended with sterile quinoa grains to stimulate growth of ambient bacteria

as their food source.

The Chrysophyceae Clade H virus SA1 (ChrysoHV) was isolated by challenging the

chrysophyte isolate UHM3500 with unfiltered seawater collected from 25 m depth at Station

ALOHA on 04 May 2019. The lytic agent detected in the challenge was rendered clonal by serial

dilutions to extinction (42). This is the first chrysophyte-infecting virus isolated from Station

ALOHA, thus the strain designation SA1. In addition to the host used for isolation, the virus was

found to infect a related chrysophyte strain UHM3501 (Supplementary Fig. S1). Both strains are

members of Chrysophyceae Clade H, but differ in their grazing kinetics and 18S rRNA gene

sequences (7). The virus was maintained on both chrysophytes through serial, monthly

challenges of exponential phase culture with lysate from the preceding challenge until

chrysophyte strain UHM3500 was lost. Approximately three to four days post-infection (DPI),

infected cultures were stored at 4°C to serve as lysate for the next round of infection.

*Host range tests and flow cytometry*

We challenged isolates of Ochrophyta protists, both mixotrophs and heterotrophs, with

raw ChrysoHV lysate, then tracked the population dynamics of the protists and viruses over time

using flow cytometry. The mixotrophic algal protists were tracked every two days up to ten DPI,

and the heterotrophic protists were tracked every day up to five DPI. The experimental design

included triplicate, 25 mL protist cultures for both negative controls and infection treatments. To

prepare the infection treatment, 500 μL of unfiltered lysate (3-6 × $10^5$ virus-like particles mL$^{-1}$

final concentration) was added to exponential or early stationary phase protist cultures. The

samples were fixed with ice-cold glutaraldehyde at 0.5% final concentration, incubated for 15

min at 4°C, and then flash frozen in liquid nitrogen (43).

The samples were stored at -80°C up to four days, then thawed at room temperature and

processed within the same day on a CytoFlex-S (Beckman Coulter Life Sciences, Indianapolis,

IN, USA) in the Biological Electron Microscope Facility (BEMF) at the University of Hawaiʻi at

Mānoa. The samples were split into two fractions, with 200 μL allocated towards quantifying

cell abundances and loaded directly onto a 96-well plate, and 5 μL allocated toward quantifying

virus abundances. The latter fractions were diluted into 195 μL solutions of TE (pH 8.0) with

2.5X SYBR Green I nucleic acid stain (Invitrogen, Thermo Fisher Scientific, Waltham, MA,

USA) and incubated at 80°C for 20 min in darkness (44), then loaded onto a 96-well plate. The

cell counts were run on auto-record for one minute at a flow rate of 20 μL min$^{-1}$ for the algae and

30 μL min$^{-1}$ for the heterotrophs. The mixotroph populations were distinguished through

chlorophyll fluorescence and 488 nm, 90° side scatter (SSC), whereas the heterotrophs were

detected using SYBR Green fluorescence and 488 nm, 90° SSC (45). The viral-like particle

(VLP) population was identified through SYBR Green fluorescence and 405 nm, 90° violet side

scatter (VSSC), enhancing marine virus detection (46). The samples were run on auto-record for

one minute at a flow rate of 10 μL min$^{-1}$. Both algae and virus populations were delineated

through a custom Python script using the following packages: fcsparser (v0.2.8), pandas (v2.2.2),

matplotlib (v3.7.1), while the FlowJo (v.10.10.0) software was used to delineate the

heterotrophic protist population. Apparent burst size was determined by dividing the maximum

concentration of VLPs post-infection by the inferred number of lysed cells, calculated as the

difference between the cell concentration on the day of infection minus the minimum

concentration post-infection.

*Electron microscopy*

Transmission electron microscope (TEM) micrographs of viral particles were prepared

from raw, unfixed lysate and negatively stained with 2% uranyl acetate for 45 sec on 200-mesh

copper grids with carbon-stabilized formvar support that had been rendered hydrophilic by glow

discharge (47). Images were captured with a Hitachi HT7700 transmission electron microscope

(Tokyo, Japan) at the Biological Electron Microscope Facility (BEMF) at the University of

Hawaiʻi at Mānoa. Particle dimensions were measured with ImageJ from Fiji (v2.16.0/1.54p)

(48), including: capsid diameter (n = 38), envelope (n = 34), tail (n = 17), and tail fibers (n = 6).

*DNA extraction, sequencing, assembly, and annotation*

To prepare sufficient viral biomass for DNA sequencing, DNA was extracted from 610

mL of lysate ($6.4 \times 10^6$ VLP/mL), after allowing the infection of chrysophyte Clade H sp.

UHM3501 to clear four DPI. The unfiltered lysate was concentrated and pooled through serial

centrifugations (6X) in a 100 kDA Centricon Plus centrifugal filter (MilliporeSigma, Burlington,

MA, USA) at 3,500 × $g$ at 20°C for 15 minutes. Approximately 200 ng of DNA was extracted

from the concentrate using the ZymoBiomics Quick-DNA High Molecular Weight MagBead Kit

(Zymo Research, Irvine, CA, USA), flash frozen in liquid nitrogen, stored at -20°C, then shipped

on dry ice to the US Department of Energy Joint Genome Institute (JGI). There, the DNA was

sequenced using PacBio Multiplexed 6–10 kb Low Input parameters.

A PacBio Low Input (DNA) library was constructed and sequenced using the PacBio

SEQUELIIe platform (Pacific Biosciences, Menlo Park, CA USA), which generated 3,267,906

reads totaling 31,033,080,767 bp. Reads were deduplicated using pbmarkdup (v1.0.3) (49).

BBDuk (v38.99) (50) was used to remove reads that contained PacBio control sequences and

PCR adapters, and IceCreamFinder (v38.99) (51) was used to remove reads without SMRTbell

sequences. The final filtered fastq contained 3,257,268 reads totaling 30,921,351,264 bp with

mean read length of 9,493 bp.

The filtered reads (31 Gb) were assembled using Flye (v2.9) (52) with the "--meta --

pacbio-hifi" argument. From this initial assembly, viral contigs were identified and grouped

through a low-complexity binning workflow, utilizing CONCOCT (v1.1.0) (53), MaxBin

(v2.2.7) (54), MetaBAT2 (v2.15-5) (55), and SemiBin2 (v1.5.1) (56). For each binning strategy,

the viral bin was identified by assessing the presence of giant virus marker genes (10) with

GVClass (v1.0) (57), and the MetaBAT2-generated viral bin was used in downstream analysis.

Subsequently, circular consensus sequencing reads were recruited against these identified viral

contigs, and a 100 Mb subsample of these reads was assembled using Flye without the "--meta"

argument (v2.9) (52). This iterative assembly resulted in one high-quality viral contig, which was

polished using two rounds of Racon (v1.6.0) (58) to ensure accuracy.

Gene calling was performed with GVClass (v1.0) (57), and the 1.196 Mb viral genome

had an estimated 93.77% coding density. Gene annotation was performed with KofamScan

(v1.3.0) (59) and eggnog mapper (v2.1.12) (60) with annotations of genes of interests confirmed

with BLASTp (61). tRNAs were annotated with tRNAscan-SE (v2.0.12) (62).

*Phylogenetic analysis*

Phylogenetic analysis of ChrysoHV with IQ-TREE (v2.2.6) (63) was based on a

concatenated alignment of seven core proteins: SFII, RNAPL, PolB, TFIIB, TopoII, A32, and

VLTF3 (10). Query sequences were retrieved from the GVMAGs (20) and GenBank Assembly

(64) databases, and quality filtered to contain at least four out of seven core proteins determined

through hmmsearch (v3.3.2, E-value $\leq 1 \times 10^{-10}$) (65). Additionally, the query sequences were

filtered to remove genomes with more than 10 copies of any core protein and more than four

core proteins with multiple copies. The remaining 990 query sequences were aligned with

MAFFT (v7.520) (66) and trimmed with trimAl (-gt 0.1; v1.4.1) (67), then concatenated with a

custom python script (68). The trimmed, concatenated alignment was used for phylogenetic

analysis using the mixture model LG+I+F+G4 and ultrafast bootstrap supports (option: -bb

1000) (v2.2.6) (63). The tree was annotated with iTOL (v6) (69).

Gene phylogeny of microbial rhodopsins with a heliorhodopsin outgroup were built with

IQ-TREE (v2.2.6) (63) using model LG+F+R10 and ultrafast bootstrap supports (option: -bb

1000) and annotated with iTOL (v6) (69). Representatives of rhodopsin homologs from

cultivated virus isolates include: EhV-202 (heliorhodopsin) (17), PgV-12T and -16T (viral

rhodopsin I) (27), and FloV-SA2 (viral rhodopsin II) (16). Additionally, the gene phylogeny

includes synthetic construct: vPyACR_21921 (channelrhodopsin) (15), and rhodopsin homologs

from single amplified genomes (SAGs) and metagenome assembled genomes (MAGS):

ChoanoV1 (viral group II) (14), cPacV-1605 (viral group II) (70), and OLPV 1 and 2 (viral

groups I & II) (36). Query rhodopsin sequences were retrieved from NCBI and the

supplementary materials of Olson *et al*. (2018), which included proteorhodopsin sequences

recovered from metagenomic sequences at Station ALOHA (29). The alignment was built with

MAFFT (L-INS-i; v7.520) (66) and trimmed with trimAl (-gt 0.1; v1.4.1) (67).

The phylogeny of near-complete 18S rRNA genes from select algal isolates were built

with IQ-TREE (v2.2.6) (63) using GTR+I+G model with ultrafast bootstrap support (option: -bb

1000) and annotated with iTOL (v6) (69). The nucleotide sequences were retrieved from NCBI,

aligned with MAFFT (L-INS-i; v7.520) (66) and trimmed with trimAl (-gt 0.1; v1.4.1) (67).

*Gene distribution in the global ocean*

Comparisons to *Nucleocytoviricota* PolB amino acid sequences from the *Tara* Oceans dataset (11) were performed using BLASTp (v2.13.0, E-value $\leq 1 \times 10^{-120}$) (61), and sequences with 90% or greater aaID were plotted using a custom R script with the following packages: maps (v3.4.2), mapdata (v2.3.1), and ggplot2 (v.3.5.1). The relative abundance of the PolB genes were normalized to the total number of *Nucleocytoviricota* PolBs (11). Likewise, ChysoHV proteorhodopsin-like protein sequences from the *Tara* Oceans dataset with a minimum of 80% aaID were identified through Ocean Gene Atlas (71, 72). The relative abundance of ChysoHV proteorhodopsin-like protein sequences were normalized to total metagenomic reads in the given sample (71, 72).

*Protein structure predictions*

The conserved domain of the ChrysoHV rhodopsin structure was predicted using ColabFold (v1.5.5) (73) and compared against the AlphaFold Protein Structure Databases– SwissProt (74) and CATH (v4.4) (75) using Foldseek (76). The conserved domain of the ChrysoHV rhodopsin and HOT 75m4 Gamma-proteobacterium proteorhodopsin structures were rendered in ChimeraX (v1.8.dev202403010247) (77).

**Acknowledgments**

We thank the personnel of the Hawai'i Ocean Time-series program (NSF award 12-60164) and Research Vessel Kilo Moana for assistance with seawater collection. We thank the Biological Electron Microscope Facility (BEMF) at the University of Hawai'i at Mānoa, especially facility supervisors, Tina Weatherby and Dr. Orion Rivers, for their assistance with electron microscopy

procedures. A special thanks to laboratory technician Kelsey McBeain for maintenance of the

UHM library culture collection, and US DOE Joint Genome Institute Scientific Project Manager,

Kerrie Barry, for overseeing DNA library preparation, quality control, and sequencing of

ChrysoHV.

We are grateful for thoughtful feedback provided by Dr. Karen Selph (Department of

Oceanography, University of Hawaiʻi at Mānoa) and Dr. Alison Sherwood (School of Life

Sciences, University of Hawaiʻi at Mānoa) on manuscript preparation, and insights into

rhodopsin structure and rhodopsin sequences (SAMN27960419 and SAMEA2623059), provided

by Dr. Oded Béjà (Faculty of Biology, Technion-Israel Institute of Technology) and Dr. Andrey

Rozenberg (Faculty of Biology, Technion-Israel Institute of Technology).

**Funding**

This work was supported by the US National Science Foundation Graduate Research Fellowship

Program (Grant No. 1842402 to PB), Division of Ocean Sciences (Grant No. 2129697 to GFS

and KFE), and RII Track-2 FEC (Grant No. 1736030 to GFS and KFE). The work conducted by

the US Department of Energy Joint Genome Institute (https://ror.org/04xm1d337), a Department

of Energy (DOE) Office of Science User Facility, was supported by the Office of Science of the

US DOE, operated under Contract No. DE-AC02-05CH11231.

**Author Contributions**

GFS, KFE, and PB developed, designed, and acquired funding support for the project. CRS

isolated the virus and algae strains. CW and QL assisted with the collection of virus host-range

data. FS provided sequencing support and supervised the bioinformatics analyses. KL assisted

with the virus genome assembly. JT assisted with gene phylogeny workflows. CRS and PB

imaged the virus. PB wrote the manuscript with input from all authors.

**Conflicts of Interest**

The authors declare no conflicts of interest.

**Data Availability**

ChrysoHV sequencing data is available through the NCBI sequence read archive SRR34002084,

and the genome assembly has been deposited to NCBI GenBank under accession ID PV764857.

The mixotrophic protists used in the study, GenBank IDs: MZ611734.1 (Chrysophyceae Clade H

sp. UHM3500), MZ611720.1 (Chrysophyceae Clade H sp. UHM3501), MN615710.1

(*Florenciella* sp. UHM3020), MZ611733.1 (Dictyochophyceae clade X sp. UHM3020),

MZ611705.1 (*Rhizochromulina* sp. UHM3071), MZ611730.1 (*Triparma* sp. UHM3600), are

available through NCMA. The virus and heterotrophic protists used in this study, GenBank IDs:

PV717318.1 (*Paraphysomonas* sp. UHM3560) and PV717319.1 (*Spumella* sp. UHM3530), are

available through the Marine Viral Ecology Laboratories at the University of Hawaiʻi at Mānoa

upon request to KFE and GFS. Additional data and scripts on gene alignments, phylogenies,

genome annotation, protein structural predictions, host range challenges, and biogeography are

available on the Dryad repository:

http://datadryad.org/share/8duyiXJ3riDLgLbseVaWvUBnbOPOM4SkBw9WNBaRYA4.

**Figures**

**Figure 1**. **ChrysoHV infection dynamics and host range tests.** (A) The left panel shows a time

series of Chrysophyceae Clade H sp. UHM3501 abundance (cells/mL) in response to the

addition of raw ChrysoHV lysate. The control treatment is shown in the black dashed line with

white-fill points, and the infection treatment shown in the solid black line with black-fill points.

The right panel shows the abundance of ChrysoHV (VLP/mL) during the same time series. Days

pre- and post-infection are displayed on the x-axis. Error bars represent the standard deviation of

the triplicate counts. (B) Maximum Likelihood (ML) 18S rRNA phylogeny (GTR+I+R model)

420    of the Ochrophyta phylum with isolates challenged in bold. The white or black boxes by the

421    bolded isolate names indicate whether the protist culture did not lyse (white with "X") or lysed

422    and produced virions (black) in response to the ChrysoHV challenge. The node symbols indicate

423    the bootstrap values equal to or greater than 80% (white), 90% (gray) or 100% (black). There is

424    no node symbol for branches with less than 80% support.

**Figure 2**. **ChrysoHV ultrastructure, genome, and phylogenetic analysis.** (A) Transmission

electron microscopy (TEM) micrograph of ChrysoHV propagated on UHM3501. (B, C) TEM

micrograph of negatively stained ChrysoHV propagated on UHM3500. (D) Genome plot of

ChrysoHV with track placement and color showing gene lineage annotated according to domain,

inner to outer tracks showing: archaeal and bacterial (blue), eukaryotic (yellow), and viral

(black) lineages. Phylogenetically informative giant virus marker genes are represented as black

dots on the outside track and show gene name. The genes used in the *Nucleocytoviricota*

phylogenetic analysis are bolded. (E) Maximum Likelihood (ML) phylogeny of the

*Nucleocytoviricota* phylum inferred from a concatenated alignment of amino acid sequences of

marker genes (SFII, RNAPL, PolB, TFIIB, TopoII, A32, VLTF3) using model LG+I+F+G4 with

ultrafast bootstrap supports (-bb 1000) shown by node symbol color– white ($\geq$ 80%), gray ($\geq$

90%), and black (100%) with no node symbol for supports < 80%. The tree is rooted with

*Chitovirales* as the outgroup, and the scale bar represents one amino acid substitution per site.

The frequency of microbial rhodopsins and photolyases present in *Aliimimivirinae*

representatives are annotated in the grayscale matrix, and the environment the representative was

collected from is shown in jade (freshwater), blue (marine), and brown (terrestrial).

**Figure 3**. **Rhodopsin gene phylogeny, predicted structure, and gene neighborhood.** (A)

Microbial rhodopsin ML phylogeny with heliorhodopsin as the outgroup, built with model

LG+F+R10. Select representatives of viral rhodopsin homologs and putative chrysophyte

rhodopsins are referred to in the parenthesis of their corresponding collapsed clade. The

proteorhodopsin (PR) clade is annotated with the alignment of transmembrane helix 3 (TM3)

outlining the ion-pumping motif (gray) and the spectral tuning amino acid (green or blue).

Residues that differ from the most common residue at that locus are in bold (B). The predicted

structure of ChrysoHV's proteorhodopsin-like protein's conserved domain (yellow) and the HOT

75m4 Gamma-proteobacterium proteorhodopsin (blue). (C) Gene neighborhood of

photoreceptive proteins in ChrysoHV, showing the proteorhodopsin (locus tag: ChrysoHV_0101;

GenBank ID: XVM28022.1) next to the enzyme lysine decarboxylase (locus tag:

ChrysoHV_0099, GenBank ID: XVM28020.1), between two photolyases (locus tag:

ChyrsoHV_0095, ChyrsoHV_0104; GenBank ID: XVM28016.1, XVM28025.1), and one

heliorhodopsin (locus tag: ChrysoHV_0106, GenBank ID: XVM28027.1).

**Figure 4**. **Distribution and relative abundance of ChrysoHV PolB and PR**

**(proteorhodopsin)**. (A) Global distribution of phylotypes with ≥ 90% PolB aaID to ChrysoHV

from the *Tara* Oceans dataset. Size of the circle indicates the relative abundance of the summed

phylotypes at each given sample site to total *Nucleocytoviricota* PolB sequences per site. (B)

Global distribution of homologous PR with ≥ 80% aaID to the ChrysoHV proteorhodopsins with

size of the circle indicating relative abundance normalized to total metagenomic reads per site.

Sample depth is shown through color: surface (SRF: 3-7 m, orange), deep chlorophyll maximum

(DCM: 7-200 m, green), and mesopelagic (MES: 200-1,000 m, purple).

**References**

1.     Del Campo J, Massana R. 2011. Emerging diversity within chrysophytes, choanoflagellates

and bicosoecids based on molecular surveys. Protist 162:435–448.

2.     Seeleuthner Y, Mondy S, Lombard V, Carradec Q, Pelletier E, Wessner M, Leconte J,

Mangot J-F, Poulain J, Labadie K, Logares R, Sunagawa S, De Berardinis V, Salanoubat

M, Dimier C, Kandels-Lewis S, Picheral M, Searson S, Tara Oceans Coordinators, Acinas

SG, Boss E, Follows M, Gorsky G, Grimsley N, Karp-Boss L, Krzic U, Not F, Ogata H,

Raes J, Reynaud EG, Sardet C, Speich S, Stemmann L, Velayoudon D, Weissenbach J,

Pesant S, Poulton N, Stepanauskas R, Bork P, Bowler C, Hingamp P, Sullivan MB,

Iudicone D, Massana R, Aury J-M, Henrissat B, Karsenti E, Jaillon O, Sieracki M, De

Vargas C, Wincker P. 2018. Single-cell genomics of multiple uncultured stramenopiles

reveals underestimated functional diversity across oceans. Nat Commun 9:310.

3.     Olefeld JL, Majda S, Albach DC, Marks S, Boenigk J. 2018. Genome size of chrysophytes

varies with cell size and nutritional mode. Org Divers Evol 18:163–173.

4.    Dai M, Luo Y, Achterberg EP, Browning TJ, Cai Y, Cao Z, Chai F, Chen B, Church MJ, Ci

485           D, Du C, Gao K, Guo X, Hu Z, Kao S, Laws EA, Lee Z, Lin H, Liu Q, Liu X, Luo W,

486           Meng F, Shang S, Shi D, Saito H, Song L, Wan XS, Wang Y, Wang W, Wen Z, Xiu P,

487           Zhang J, Zhang R, Zhou K. 2023. Upper ocean biogeochemistry of the oligotrophic North

488           Pacific Subtropical Gyre: from nutrient sources to carbon export. Reviews of Geophysics

489           61:e2022RG000800.

490    5.    Frias-Lopez J, Thompson A, Waldbauer J, Chisholm SW. 2009. Use of stable isotope-

491           labelled cells to identify active grazers of picocyanobacteria in ocean surface waters.

492           Environmental Microbiology 11:512–525.

493    6.    Wilken S, Yung CCM, Poirier C, Massana R, Jimenez V, Worden AZ. 2023.

494           Choanoflagellates alongside diverse uncultured predatory protists consume the abundant

495           open-ocean cyanobacterium *Prochlorococcus*. Proc Natl Acad Sci USA 120:e2302388120.

496    7.    Li Q, Edwards KF, Schvarcz CR, Steward GF. 2022. Broad phylogenetic and functional

497           diversity among mixotrophic consumers of *Prochlorococcus*. The ISME Journal 16:1557–

498           1569.

499    8.    Flombaum P, Gallegos JL, Gordillo RA, Rincón J, Zabala LL, Jiao N, Karl DM, Li WKW,

500           Lomas MW, Veneziano D, Vera CS, Vrugt JA, Martiny AC. 2013. Present and future

501           global distributions of the marine Cyanobacteria *Prochlorococcus* and *Synechococcus*. Proc

502           Natl Acad Sci USA 110:9824–9829.

9.    Castillo YM, Mangot J, Benites LF, Logares R, Kuronishi M, Ogata H, Jaillon O, Massana

R, Sebastián M, Vaqué D. 2019. Assessing the viral content of uncultured picoeukaryotes

in the global-ocean by single cell genomics. Molecular Ecology 28:4272–4289.

10.   Aylward FO, Moniruzzaman M, Ha AD, Koonin EV. 2021. A phylogenomic framework for

charting the diversity and evolution of giant viruses. PLoS Biol 19:e3001430.

11.   Endo H, Blanc-Mathieu R, Li Y, Salazar G, Henry N, Labadie K, De Vargas C, Sullivan

MB, Bowler C, Wincker P, Karp-Boss L, Sunagawa S, Ogata H. 2020. Biogeography of

marine giant viruses reveals their interplay with eukaryotes and ecological functions. Nat

Ecol Evol 4:1639–1649.

12.   Fromm A, Hevroni G, Vincent F, Schatz D, Martinez-Gutierrez CA, Aylward FO, Vardi A.

2024. Single-cell RNA-seq of the rare virosphere reveals the native hosts of giant viruses in

the marine environment. Nat Microbiol 9:1619–1629.

13.   Moniruzzaman M, Erazo Garcia MP, Farzad R, Ha AD, Jivaji A, Karki S, Sheyn U, Stanton

J, Minch B, Stephens D, Hancks DC, Rodrigues RAL, Abrahao JS, Vardi A, Aylward FO.

2023. Virologs, viral mimicry, and virocell metabolism: the expanding scale of cellular

functions encoded in the complex genomes of giant viruses. FEMS Microbiology Reviews

47:fuad053.

14.   Needham DM, Yoshizawa S, Hosaka T, Poirier C, Choi CJ, Hehenberger E, Irwin NAT,

Wilken S, Yung C-M, Bachy C, Kurihara R, Nakajima Y, Kojima K, Kimura-Someya T,

Leonard G, Malmstrom RR, Mende DR, Olson DK, Sudo Y, Sudek S, Richards TA,

DeLong EF, Keeling PJ, Santoro AE, Shirouzu M, Iwasaki W, Worden AZ. 2019. A

distinct lineage of giant viruses brings a rhodopsin photosystem to unicellular marine

predators. Proc Natl Acad Sci USA 116:20574–20583.

15.   Rozenberg A, Oppermann J, Wietek J, Fernandez Lahore RG, Sandaa R-A, Bratbak G,

Hegemann P, Béjà O. 2020. Lateral gene transfer of anion-conducting channelrhodopsins

between green algae and giant viruses. Current Biology 30:4910-4920.e5.

16.   Thomy J, Schvarcz CR, McBeain KA, Edwards KF, Steward GF. 2024. Eukaryotic viruses

encode the ribosomal protein eL40. npj Viruses 2:51.

17.   Hososhima S, Mizutori R, Abe-Yoshizumi R, Rozenberg A, Shigemura S, Pushkarev A,

Konno M, Katayama K, Inoue K, Tsunoda SP, Béjà O, Kandori H. 2022. Proton-

transporting heliorhodopsins from marine giant viruses. eLife 11:e78416.

18.   Yutin N, Koonin EV. 2012. Proteorhodopsin genes in giant viruses. Biol Direct 7:34.

19.   Moniruzzaman M, Martinez-Gutierrez CA, Weinheimer AR, Aylward FO. 2020. Dynamic

genome evolution and complex virocell metabolism of globally-distributed giant viruses.

Nat Commun 11:1710.

20.   Schulz F, Roux S, Paez-Espino D, Jungbluth S, Walsh DA, Denef VJ, McMahon KD,

Konstantinidis KT, Eloe-Fadrosh EA, Kyrpides NC, Woyke T. 2020. Giant virus diversity

and host interactions through global metagenomics. Nature 578:432–436.

21.   Pinhassi J, DeLong EF, Béjà O, González JM, Pedrós-Alió C. 2016. Marine bacterial and

archaeal ion-pumping rhodopsins: genetic diversity, physiology, and ecology. Microbiol

Mol Biol Rev 80:929–954.

22.  Rozenberg, A Inoue, K, Kandori, H, Béjà O. Microbial rhodopsins: the last two decades.

Annu Rev Microbiol 75:427-447.

23.  Bratanov D, Kovalev K, Machtens J-P, Astashkin R, Chizhov I, Soloviov D, Volkov D,

Polovinkin V, Zabelskii D, Mager T, Gushchin I, Rokitskaya T, Antonenko Y, Alekseev A,

Shevchenko V, Yutin N, Rosselli R, Baeken C, Borshchevskiy V, Bourenkov G, Popov A,

Balandin T, Büldt G, Manstein DJ, Rodriguez-Valera F, Fahlke C, Bamberg E, Koonin E,

Gordeliy V. 2019. Unique structure and function of viral rhodopsins. Nat Commun

10:4939.

24.  Zabelskii D, Alekseev A, Kovalev K, Rankovic V, Balandin T, Soloviov D, Bratanov D,

Savelyeva E, Podolyak E, Volkov D, Vaganova S, Astashkin R, Chizhov I, Yutin N, Rulev

M, Popov A, Eria-Oliveira A-S, Rokitskaya T, Mager T, Antonenko Y, Rosselli R, Armeev

G, Shaitan K, Vivaudou M, Büldt G, Rogachev A, Rodriguez-Valera F, Kirpichnikov M,

Moser T, Offenhäusser A, Willbold D, Koonin E, Bamberg E, Gordeliy V. 2020. Viral

rhodopsins 1 are an unique family of light-gated cation channels. Nat Commun 11:5707.

25.  Sheam MM, Luo E. 2025. Vertical transport and spatiotemporal dynamics of giant viruses

in the North Pacific subtropical gyre. The ISME Journal 19:wraf094.

26.  Bosmon T, Abergel C, Claverie J-M. 2025. 20 years of research on giant viruses. npj

Viruses 3:9.

27.  Brussaard CPD, Bratbak G, Baudoux A-C, Ruardij P. 2007. Phaeocystis and its interaction

with viruses. Biogeochemistry 83:201–215.

28.    Lamb DC, Goldstone JV, Belhaouari DB, Andréani J, Farooqi A, Allen MJ, Kelly SL, La

Scola B, Stegeman JJ. 2025. Cytochrome b5 occurrence in giant and other viruses

belonging to the phylum Nucleocytoviricota. npj Viruses 3:8.

29.    Olson DK, Yoshizawa S, Boeuf D, Iwasaki W, DeLong EF. 2018. Proteorhodopsin

variability and distribution in the North Pacific Subtropical Gyre. The ISME Journal

12:1047–1060.

30.    Zhang L, Meng L, Fang Y, Ogata H, Okazaki Y. 2024. Spatiotemporal dynamics of giant

viruses within a deep freshwater lake reveal a distinct dark-water community. The ISME

Journal 18:wrae182.

31.    Pitot TM, Girard C, Rapp JZ, Somerville V, Culley AI, Vincent WF, Moineau S, Roux S.

2025. Viral niche-partitioning: comparative genomics of giant viruses across environmental

gradients in a high Arctic freshwater-saltwater lake. ISME Communications 5:ycae155.

32.    Fang J, Zhang Y, Zhu T, Li Y. 2023. Scramblase activity of proteorhodopsin confers

physiological advantages to Escherichia coli in the absence of light. iScience 26:108551.

33.    Ran T, Ozorowski G, Gao Y, Sineshchekov OA, Wang W, Spudich JL, Luecke H. 2013.

Cross-protomer interaction with the photoactive site in oligomeric proteorhodopsin

complexes. Acta Crystallogr D Biol Crystallogr 69:1965–1980.

34.    Fischer MG, Allen MJ, Wilson WH, Suttle CA. 2010. Giant virus with a remarkable

complement of genes infects marine zooplankton. Proc Natl Acad Sci USA 107:19508–

19513.

35.   Sheng Y, Wu Z, Xu S, Wang Y. 2022. Isolation and identification of a large green alga

585            virus (*Chlorella* Virus XW01) of *mimiviridae* and its virophage (*Chlorella* Virus Virophage

586            SW01) by using unicellular green algal cultures. J Virol 96:e02114-21.

587    36.   Yau S, Lauro FM, DeMaere MZ, Brown MV, Thomas T, Raftery MJ, Andrews-Pfannkoch

588            C, Lewis M, Hoffman JM, Gibson JA, Cavicchioli R. 2011. Virophage control of antarctic

589            algal host–virus dynamics. Proc Natl Acad Sci USA 108:6163–6168.

590    37.   Strauss J, Deng L, Gao S, Toseland A, Bachy C, Zhang C, Kirkham A, Hopes A, Utting R,

591            Joest EF, Tagliabue A, Löw C, Worden AZ, Nagel G, Mock T. 2023. Plastid-localized

592            xanthorhodopsin increases diatom biomass and ecosystem productivity in iron-limited

593            surface oceans. Nat Microbiol 8:2050–2066.

594    38.   Giovannoni SJ. 2017. SAR11 Bacteria: The most abundant plankton in the oceans. Annu

595            Rev Mar Sci 9:231–255.

596    39.   Filée J. 2018. Giant viruses and their mobile genetic elements: the molecular symbiosis

597            hypothesis. Current Opinion in Virology 33:81–88.

598    40.   Letelier RM, White AE, Bidigare RR, Barone B, Church MJ, Karl DM. 2017. Light

599            absorption by phytoplankton in the North Pacific Subtropical Gyre. Limnology &

600            Oceanography 62:1526–1540.

601    41.   Keller MD, Selvin RC, Claus W, Guillard RRL. 1987. Media for the culture of oceanic

602            ultraphytoplankton. Journal of Phycology 23:633–638.

42.    Bratbak G, Jacobsen A, Heldal M, Nagasaki K, Thingstad F. 1998. Virus production in

Phaeocystis pouchetii and its relation to host cell growth and nutrition. Aquat Microb Ecol

16:1–9.

43.    Li Q, Edwards KF, Schvarcz CR, Selph KE, Steward GF. 2021. Plasticity in the grazing

ecophysiology of *Florenciella* (Dichtyochophyceae), a mixotrophic nanoflagellate that

consumes *Prochlorococcus* and other bacteria. Limnology & Oceanography 66:47–60.

44.    Brussaard CPD. 2004. Optimization of Procedures for Counting Viruses by Flow

Cytometry. Appl Environ Microbiol 70:1506–1513.

45.    Christaki U, Courties C, Massana R, Catala P, Lebaron P, Gasol JM, Zubkov MV. 2011.

Optimized routine flow cytometric enumeration of heterotrophic flagellates using SYBR

Green I. Limnology & Ocean Methods 9:329–339.

46.    Zhao Y, Zhao Y, Zheng S, Zhao L, Zhang W, Xiao T, Grégori G. 2023. Enhanced

resolution of marine viruses with violet side scatter. Cytometry Pt A 103:260–268.

47.    Doane FW, Anderson N. 1987. Electron microscopy in diagnostic virology: a practical

guide and atlas. Cambridge University Press, New York.

48.    Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,

Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K,

Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-image analysis.

Nat Methods 9:676–682.

49.    PacificBiosciences. pbdmarkdup. https://github.com/PacificBiosciences/pbmarkdup (25

623           February 2025, date last accessed)

624    50.    BioInfoTools. BBMap. https://github.com/BioInfoTools/BBMap/blob/master/sh/bbduk.sh

625           (21 February 2021, date last accessed)

626    51.    Kbaseapps. BBTools. https://github.com/kbaseapps/BBTools?tab=readme-ov-file (21

627           February 2021, date last accessed)

628    52.    Kolmogorov M, Yuan J, Lin Y, Pevzner PA. 2019. Assembly of long, error-prone reads

629           using repeat graphs. Nat Biotechnol 37:540–546.

630    53.    Alneberg J, Bjarnason BS, De Bruijn I, Schirmer M, Quick J, Ijaz UZ, Lahti L, Loman NJ,

631           Andersson AF, Quince C. 2014. Binning metagenomic contigs by coverage and

632           composition. Nat Methods 11:1144–1146.

633    54.    Wu Y-W, Tang Y-H, Tringe SG, Simmons BA, Singer SW. 2014. MaxBin: an automated

634           binning method to recover individual genomes from metagenomes using an expectation-

635           maximization algorithm. Microbiome 2:26.

636    55.    Kang DD, Li F, Kirton E, Thomas A, Egan R, An H, Wang Z. 2019. MetaBAT 2: an

637           adaptive binning algorithm for robust and efficient genome reconstruction from

638           metagenome assemblies. PeerJ 7:e7359.

639    56.    Pan S, Zhao X-M, Coelho LP. 2023. SemiBin2: self-supervised contrastive learning leads

640           to better MAGs for short- and long-read sequencing. Bioinformatics 39:i21–i29.

57.  Pitot TM, Brůna T, Schulz F. 2024. Conservative taxonomy and quality assessment of giant

virus genomes with GVClass. npj Viruses 2:60.

58.  Vaser R, Sović I, Nagarajan N, Šikić M. 2017. Fast and accurate de novo genome assembly

from long uncorrected reads. Genome Res 27:737–746.

59.  Aramaki T, Blanc-Mathieu R, Endo H, Ohkubo K, Kanehisa M, Goto S, Ogata H. 2020.

KofamKOALA: KEGG Ortholog assignment based on profile HMM and adaptive score

threshold. Bioinformatics 36:2251–2252.

60.  Cantalapiedra CP, Hernández-Plaza A, Letunic I, Bork P, Huerta-Cepas J. 2021. eggNOG-

mapper v2: functional annotation, orthology assignments, and domain prediction at the

metagenomic scale. Molecular Biology and Evolution 38:5825–5829.

61.  Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009.

BLAST+: architecture and applications. BMC Bioinformatics 10:421.

62.  Chan PP, Lin BY, Mak AJ, Lowe TM. 2021. tRNAscan-SE 2.0: improved detection and

functional classification of transfer RNA genes. Nucleic Acids Research 49:9077–9096.

63.  Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, Von Haeseler A,

Lanfear R. 2020. IQ-TREE 2: New models and efficient methods for phylogenetic

inference in the genomic era. Molecular Biology and Evolution 37:1530–1534.

64.  Benson DA, Cavanaugh M, Clark K, Karsch-Mizrachi I, Lipman DJ, Ostell J, Sayers EW.

2012. GenBank. Nucleic Acids Research 41:D36–D42.

65.  Eddy SR. 2011. Accelerated profile HMM searches. PLoS Comput Biol 7:e1002195.

66.   Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. Molecular Biology and Evolution 30:772–780.

67.   Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: a tool for automated

alignment trimming in large-scale phylogenetic analyses. Bioinformatics 25:1972–1973.

68.   NeLli-team. nsgtree. https://github.com/NeLLi-team/nsgtree (03 March 2024, date last

accessed)

69.   Letunic I, Bork P. 2024. Interactive Tree of Life (iTOL) v6: recent updates to the

phylogenetic tree display and annotation tool. Nucleic Acids Research 52:W78–W82.

70.   Needham DM, Poirier C, Hehenberger E, Jiménez V, Swalwell JE, Santoro AE, Worden

AZ. 2019. Targeted metagenomic recovery of four divergent viruses reveals shared and

distinctive characteristics of giant viruses of marine eukaryotes. Phil Trans R Soc B

374:20190086.

71.   Villar E, Vannier T, Vernette C, Lescot M, Cuenca M, Alexandre A, Bachelerie P, Rosnet

T, Pelletier E, Sunagawa S, Hingamp P. 2018. The Ocean Gene Atlas: exploring the

biogeography of plankton genes online. Nucleic Acids Research 46:W289–W295.

72.   Vernette C, Lecubin J, Sánchez P, Tara Oceans Coordinators, Acinas SG, Babin M, Bork P,

Boss E, Bowler C, Cochrane G, De Vargas C, Gorsky G, Guidi L, Grimsley N, Hingamp1

P, Iudicone D, Jaillon O, Kandels-Lewis S, Karp-Boss L, Karsenti E, Not F, Ogata H,

Poulton N, Pesant S, Sardet C, Speich S, Stemmann L, Sullivan MB, Sunagawa S, Wincker

P, Sunagawa S, Delmont TO, Acinas SG, Pelletier E, Hingamp P, Lescot M. 2022. The

Ocean Gene Atlas v2.0: online exploration of the biogeography and phylogeny of plankton

genes. Nucleic Acids Research 50:W516–W526.

73.  Mirdita M, Schütze K, Moriwaki Y, Heo L, Ovchinnikov S, Steinegger M. 2022.

ColabFold: making protein folding accessible to all. Nat Methods 19:679–682.

74.  The UniProt Consortium. 2017. UniProt: the universal protein knowledgebase. Nucleic

Acids Res 45:D158–D169.

75.  Waman VP, Bordin N, Lau A, Kandathil S, Wells J, Miller D, Velankar S, Jones DT,

Sillitoe I, Orengo C. 2025. CATH v4.4: major expansion of CATH by experimental and

predicted structural data. Nucleic Acids Research 53:D348–D355.

76.  Van Kempen M, Kim SS, Tumescheit C, Mirdita M, Lee J, Gilchrist CLM, Söding J,

Steinegger M. 2024. Fast and accurate protein structure search with Foldseek. Nat

Biotechnol 42:243–246.

77.  Meng EC, Goddard TD, Pettersen EF, Couch GS, Pearson ZJ, Morris JH, Ferrin TE. 2023.

UCSF CHIMERAX: Tools for structure building and analysis. Protein Science 32.

695