Multi-omics reveal critical differentiation target for Parkinson's Disease-vulnerable midbrain dopaminergic neurons

Author Information

Roberto Garcia-Swinburn[1]*#, Guochang Lyu[1,3]#, Judith C. Kreutzmann[1,2] , Anqi Xiong[1], Rika Kojima[2], Carmen Abaurre[1], Chiara Tremolanti[1], Sandra Gellhaar[1], Per Uhlén[1], Per Svenningsonn[2,4], Gonçalo Castelo-Branco[1], Carmen Salto[1], Onur Dagliyan[1]*, Ernest Arenas[1,3]†

Affiliations

[1]Laboratory of Molecular Neurobiology, Department of Medical Biochemistry and Biophysics, Karolinska Institutet, 171 77 Stockholm, Sweden

[2]Department of Clinical Neuroscience, Karolinska Institutet, 171 76 Stockholm, Sweden

[3]Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD, 20815.

[4]Department of Basic and Clinical Neuroscience, King's College London, WC2R 2LS London, UK

*Corresponding author: roberto.garcia.swinburn@ki.se or onur.dagliyan@ki.se

#These authors contributed equally

† Deceased 15/09/2024

Abstract

Parkinson's disease (PD) is characterized by the progressive loss of midbrain dopaminergic (mDA) neurons, leading to severe debilitating motor impairment. Recent single-cell analyses have revealed that specific mDA subpopulations, such as SOX6[+] AGTR1[+] neurons, are most vulnerable to degeneration. Current human pluripotent stem cell differentiation protocols fail to selectively generate these subtypes. Here we describe a multiomic-guided strategy that enriches SOX6[+] mDA neurons by combining enhancing Sonic Hedgehog agonism with prolonged Wnt activation. This approach accelerates floor plate specification, increases expression of mDA developmental

markers, and increases substantially the yield of SOX6$^+$ mature mDA neurons compared to prior

attempts. Following intrastriatal transplantation into hemiparkinsonian mice, these cells restore

motor function in approximately 4 months and generated grafts containing SOX6$^+$ A9-like

neurons. Our work thus establishes a reproducible differentiation platform for generating the PD-

susceptible mDA subtype, providing a foundation for precision disease modelling and subtype-

targeted cell replacement therapies.

Introduction

Parkinson's disease (PD), the second most prevalent neurodegenerative disorder, lacks curative

treatments, severely impacting patients' quality of life. This condition is characterized by the

progressive degeneration of midbrain dopaminergic (mDA) neurons in the substantia nigra pars

compacta (SNpc). The loss of these neurons, which project primarily to the dorsal striatum,

depletes dopaminergic input and leads to the characteristic motor symptoms of PD[1]. Although

advances in pharmacological therapies have been made, these treatments are often insufficient in

halting disease progression or restoring lost neuronal function in later stages[2]. Cell therapy offers

a promising alternative, with the potential to replace the damaged or lost DA neurons critical to

PD pathology[3].

Advances in stem cell and midbrain development research have enabled the generation of

transplantable mDA progenitors from human pluripotent stem cells (hPSCs)[4–7]. Success in

preclinical models has led to the development of clinical-grade mDA progenitors for PD clinical

trials[8–11] with recent phase I/II clinical trials reporting safe outcomes, thereby paving the way for

future medical use of hPSCs[12,13].

However, the translational promise of hPSC-derived cell therapies hinges on the precise

composition and quality of the grafted cells. Single-cell transcriptomic studies have begun to

molecularly define the diversity of human mDA neurons[14,15], both in development and in hPSC-

derived midbrain cultures[7,15–17]. These studies have revealed a critical challenge: not all mDA

neurons are equally susceptible to PD. Recent work has reported complications like dyskinesias,

which is linked to serotonergic neuron contamination in fetal transplants[18]. Thus, the inability of

existing differentiation protocols, which generate heterogeneous mixes of mDA subtypes, to

specifically target this critical population represents a significant limitation for both disease

modelling and the development of precision cell therapies.

Single-cell transcriptomics has revealed significant diversity within mDA neurons in the adult

human SNpc, categorizing them into three major subtypes based on the expression of *SOX6*,

CALB1, and *GAD2*, establishing that the SOX6 subpopulation is the most susceptible to

degeneration[19,20]. These subtypes emerge during human mDA neuron development[14], and recent

single-cell epigenomic profiling (e.g., ATAC-seq) further elucidates midbrain differentiation[21]. To

generate PD-vulnerable SOX6$^+$ mDA neurons, we first developed a simplified differentiation

protocol based on recently identified key developmental regulators[7], but molecular phenotyping

revealed unstable phenotypes at later stages, indicating insufficient patterning. This prompted

reviewing multiomic data leading to implement two key protocol modifications: enhanced

ventralization via stronger Sonic Hedgehog (SHH) pathway agonization and prolonged Wnt

activation improved midbrain patterning, with elevated *OTX2*, *EN1* and *SOX6* expression in

engineered cells. Transplanting these engineered cells into hemiparkinsonian (hemiPD) rodent

models demonstrated their therapeutic efficacy, yielding grafts enriched with SOX6$^+$ mDA

neurons and restoring motor function. Thus, we here report a multiomic-guided differentiation

strategy that specifically enriches for the critical PD-vulnerable mDA subtype and validates its

therapeutic potential *in vivo*.

Results

Longitudinal multiomic single-nuclei sequencing delineates the cell-type

composition throughout the process of midbrain dopaminergic neuron

differentiation *in vitro*

Building upon our recently published method[22], we designed a new protocol focused on precise

midbrain floor plate (mFP) patterning and improving cell viability to generate sufficient progenitor

cells, while expanding compatibility to a broader range of human induced pluripotent stem cell

lines. Our protocol enabled sequential differentiation of human induced pluripotent stem cells

(hiPSCs) into midbrain progenitors, positive in mFP progenitor markers such as FOXA2, OTX2

and LMX1A, and functional neurons expressing canonical mDA markers like CORIN, TH or

Nurr1 (Fig. 1a, Extended Fig. 1).

To delineate developmental trajectories, we performed single-cell multiomic sequencing

(scATAC-seq + scRNAseq) in benchmarked wild-type iPSCs (KOLF2.1J) at five developmental

stages (Days *in vitro* (DIV): DIV11, 16, 28, 42 and 56; Fig. 1b). Following quality control and

filtering, we annotated 9,233 high-quality nuclei using integrated manual-automatic classification

based on cluster-specific markers and developing human brain references[14] categorized into major

types, with sequential differentiation to a mature DA identity based on their enriched GO terms

(Fig. 1c,d). Chronologically, early stages (DIV11-16) showed high expression of regional

specification markers (LMX1A, OTX2). By DIV28, dopaminergic commitment marker NR4A2

upregulated, while late stages (DIV42-56) exhibited elevated expression of maturation markers

(such as TH, NR4A2, KCNJ6), indicating functional specification (Fig. 1e, Extended Fig. 2).

Further chromatin accessibility analysis showed precursor cell types such as floor plate progenitors

(ProgFP), medial neuroblasts (NbM) and DA clusters to recapitulate expected chromatin access

(*SOX6* and *LMX1A* for mDA progenitors, and *DDC* for mature functional mDA neurons) with

ProgFP having a higher potential to differentiate to SOX6 populations (Fig. 1g). We then validated

our annotations by comparing them to publicly available scRNA-seq datasets from human fetal

midbrain samples[14], by quantifying cell type identity across individual cells. As a result, our

annotated cell types closely resembled their endogenous counterparts, as demonstrated by

hierarchical clustering (Fig. 1f). High similarity scores (>0.8) were predominantly observed during

early stages of the midbrain differentiation and were primarily enriched in ProgFP and NbM,

underscoring the effectiveness of our approach in recapitulating endogenous early mDA neuron

development. Radial glia cell subtypes, Rgl2 and Rgl3, also displayed transcriptomic profiles akin

to their counterparts *in vivo*, hinting at the establishment of an mDA signaling niche to deliver cell

type-specific developmental factors for neuronal maturation *in vitro*[23]. At advanced stages, mDA

neurons, closely matched their endogenous cells from reference datasets (reference DA and DA0

vs experimental DA similarity score >0.8). These findings show that while mDA neurons matched

reference profiles, later stages may benefit from extended maturation cues to enhance postmitotic

neuronal survival.

Reconstructed differentiation trajectory resolves transcriptional dynamics during

mDA lineage specification and maturation

To move beyond chronological staging and define the molecular logic of *in vitro* differentiation,

we reconstructed the developmental trajectory of the dopaminergic lineage. We isolated

progenitor and neuronal populations (ProgFP, NbM and mDA neurons) from our multiomic dataset and inferred their relationships using pseudotime ordering and RNA velocity. Both analyses confirmed a direct progression from ProgFP and NbM progenitors to mature mDA neurons, validating the protocol's ability to recapitulate this fundamental lineage path (Fig. 1h). Uniform manifold approximation and projection (UMAP) analysis revealed a continuous maturation trajectory for mDA neurons. Gene Ontology (GO) analysis of genes ordered along this path showed a developmentally-relevant progression of biological processes: initial neurite outgrowth was followed by neurotransmitter release, synaptic assembly, and finally synaptic modulation (Extended Fig. 2b), matching the pattern observed in pseudotime analysis. Intriguingly, we observed heterogeneity in the maturation routes. Intriguingly, we observed heterogeneity in the maturation routes. While most cells followed the canonical path, a subset of late-stage cells (DIV42–56) appeared to bypass the axonal development phase characteristic of DIV28, suggesting the existence of alternative maturation mechanisms, potentially influenced by laminar identity cues established earlier in development[24].

We next applied single-cell regulatory network inference and clustering (SCENIC) to identify stage-specific regulons. Clustering of AUCell scores for activities of each regulon revealed distinct segregation between early neural progenitors, ProgFP, and NbM, and their descendant mDA neurons along the development trajectory (Fig. 1i). We identified key transcription factors (TFs) orchestrating each developmental transition. In ProgFP cells, the NR3C1 (glucocorticoid receptor) regulon was upregulated, promoting progenitor pool expansion by skewing cells toward proliferation over differentiation[25]. Concurrently, SREBF1, a pro-neural transcription factor controlling mDA neurogenesis[26], activity drove the subsequent transition to neurogenic progenitors (NProg). In NbM cells, pro-neural TFs NEUROD1 and NHLH1 emerged as top

regulons, initiating neurogenesis and differentiation in postmitotic cells[27,28]. Subtypes of mDA

neurons shared a core set of transcription factors, including PBX1, a canonical regulator of mDA

specification and survival[29]. In contrast, transcription factors as POU2F2 and BCL11A displayed

more specialized roles at distinct maturation stages[17,30]. In conclusion, trajectory reconstruction

confirms that our differentiation protocol recapitulates key embryological processes of mDA

development. More importantly, this approach provides a powerful *de novo* framework for

identifying the developmental regulatory modules, offering deeper insights into dopaminergic

neuron differentiation and maturation.

147 An iterative enhancement in ventralization improves floorplate patterning and

148 accelerates neuronal conversion

Given that robust regional patterning underpins high-quality mDA neuron generation, and our

single-cell data suggested suboptimal ventralization, we hypothesized that enhanced ventralization

would improve patterning outcomes. We induced Engrailed 1 (EN1), a well-established marker

for mDA neuron specification[36], while streamlining the protocol for scalability. EN1 plays a

pivotal role from the earliest stages of regional specification and maintenance of mDA neurons in

adulthood[31]. It supports the midbrain-hindbrain boundary formation during development and

sustains mDA function by regulating key genes (*TH, VMAT2, DAT,* and *ALDH1A1*) in postmitotic

neurons[31,32]. Thus, EN1 serves as both a surrogate for enhanced mDA neuron specification quality

and a predictor of transplantation outcome[9].

To enhance EN1 expression, we applied a higher concentration of the SHH agonist

purmorphamine[6], and optimized culture conditions to maintain cell viability (Fig. 2a).

Subsequently, elevated purmorphamine significantly boosted EN1 expression at DIV11

(Optimized: 88.53% ± 6.8%; Original: 20.00% ± 3.65%) without altering core floor plate markers

FOXA2 or LMX1A (Fig.2b,e; Extended Fig.3a). The effects of this enhanced ventralization were

more pronounced at later stages. By DIV16, during the fate transition from NbM towards early

mDA neurons, new protocol cultures exhibited an accelerated TH$^+$ neuron generation (Optimized:

26.12% ± 3.93%; Original: 3.57% ± 1.68%) and increased co-expression of TH$^+$, EN1$^+$ and

SOX6$^+$ (Optimized: 75.92% ± 1.26%; Original: 7.56% ± 5.11%; Fig. 2c,f). This improved

molecular signature persisted through maturation (DIV28), with sustained co-expression of EN1

and SOX6 (Optimized: 65.44% ± 7.06%; Original: 5.35% ± 3.75%; Fig. 2d,g, Extended Fig. 3b).

These neurons exhibited functional maturity, displaying calcium dynamics characteristic of mDA

neurons by DIV42 (Fig. 2h). Furthermore, we applied this differentiation protocol to GBA^{E236K}

hiPSC NH50143 and isogenic corrected cells NH50142, which showed similar enhancements in

EN1 expression and DA marker progression (Extended Fig. 4).

To molecularly characterize the improvements driven by enhanced ventralization, we performed

single-nuclei RNA sequencing (snRNA-seq) on cells from the new protocol at DIV14,16 and 28.

The new protocol accelerated differentiation, with a higher proportion of NProg and NbM

emerging at earlier stages, accompanied by a reduction in ProgFP cells (Fig. 3b). Using the same

annotation methods as in the previous dataset, the resulting mDA lineage cell types were enriched

with classic mDA neuron marker genes (Fig. 3c), including *FOXA2, LMX1A, OTX2, TH,* and

NR4A2, and showed an expected developmental trajectory across different cell populations (Fig.

3d). Critically, postmitotic cells exhibited higher EN1, sustained OTX2 and SOX6, and elevated

maturation markers TH, SLC18A2 (VMAT2), and CALB1[33] (Fig. 3f).

SCENIC analysis identified the key transcription factors activated by the new protocol (Fig. 3e).

The paired-box transcription factor PAX7 transiently became the dominant regulon in NProg,

suggesting precise rostro-caudal patterning as it contributes to determine the midbrain-hindbrain

boundary during regionalization[34,35]. Other leading regulons in NProg and NbM cells are also

actively involved in guiding the neuronal identity acquisition. For instance, POU3F1 and ZEB1

are essential regulators of neural lineage commitment, often working synergistically to coordinate

correct orders of neuronal development, as POU3F1 promotes early neural fate decisions, while

ZEB1 modulates the developmental timing and prevents premature differentiation[36,37]. Notably,

PITX3 emerged as a leading regulon in mDA neurons, which reflects the progressive acquisition

of mature mDA phenotypes[38] and highlights their molecular similarity to endogenous SNpc mDA

neurons (Fig. 3e,f). Regulon activity profiles from the integrated dataset revealed a distinct and

significant regulatory impacts of TCF4 and TCF12 under the new protocol (Extended Fig. 5c).

Interestingly, the new protocol also appeared to enhance SREBF1 expression throughout

differentiation, linking it to its known role in regulating midbrain dopaminergic neurogenesis[26],

and its involvement in controlling neuronal dendrite expansion through lipid homeostasis[39].

Overall, a targeted intervention to enhance ventralization, guided by multiomic insight, profoundly

accelerated and improved the specification of mDA neurons, driving them toward a transcriptomic

profile that closely resembles that of endogenous, vulnerable SNpc neurons.

Transplanted hiPSC-derived neurons mature into striatum-innervating dopaminergic

neurons and restore motor function in hemiPD mice

Having established that our optimized protocol enriches ventralized mDA subpopulation *in vitro*,

we next evaluated the therapeutic potential of our engineered cells *in vivo* using hemiparkinsonian

rodent models.. We unilaterally lesioned immunocompromised B6RGS mice by injecting 6-

OHDA into the medial forebrain bundle (MFB). Upon confirming hemiparkinsonism, we

intrastriatally transplanted 200,000 cells into randomly selected mice. Motor recovery was

assessed longitudinally by amphetamine-induced ipsilateral rotations, showing that grafted

hemiPD animals returned to pre-lesion state 4 months post-implant (m.p.i.), which was maintained

1 month later, ending our grafting experiment (Sham: -0.24 ± 0.73; 6-OHDA: 13.41 ± 1.43;

Grafted: -0.70 ±1.18 net ipsilateral rotations/min). To ensure overall motor recovery, we also

analysed complex motor coordination by tape removal test (Sham: -0.40 ± 0.34; 6-OHDA: 19.51

± 2.50; Grafted: -2.82 ±0.96 net ipsilateral removal time) and pole test (Sham: 5.00 ± 0.59; 6-

OHDA: 21.25 ± 3.53; Grafted: 8.56 ± 1.13 seconds), and spontaneous motor activity by cylinder

wall test (Sham: 51.82 ± 3.73%; 6-OHDA: 8.85 ± 4.75%; Grafted: 40.89 ± 3.73% ipsilateral

contacts). Human cells were detected in these 5-month grafts by STEM121[+] staining, confirming

that grafted cells did not migrate to other regions (Fig. 4c). Furthermore, human DANs were

detected by double positive staining of TH and HuNu, with these cells showing extended

processes, characteristic of mDA neurons (Fig. 4d,e). Using stereological quantification of

TH[+]/HuNu[+] cells (Fig. 4F), grafts were estimated to have 160000 human nuclei (155700 ± 35893),

of which about 10% were also stained positively for TH (10.36 ± 0.73%). Additionally, no actively

proliferative human cells were found within the grafts (Extended Fig. 6)

After establishing the presence of DANs within our grafts, we aimed to analyse if these DANs

recapitulated the phenotypical aspects of the most PD-vulnerable portion of nigral DANs which

have been reported to be A9-subtype (DANs from the SNpc), more specifically the

SOX6[+]/AGTR1a[+] subpopulation located in the ventral SNpc[19,40]. To this aim, we performed a

series of IHC analysis of the grafted DANs. A majority of grafted TH[+] cells were found to be

positive for several markers distinctive to the PD-vulnerable A9-subtype (Fig 4g,h), such as SOX6

(84.67 ± 3.03%), ALDH1A1 (50.67 ± 2.37%) and AGTR1 (64.16 ± 4.97%). In contrast, a minority

of the DANS stained positively for the A10-subtype marker Calbindin (29.92 ± 3.45%). Another

aspect that we wanted to evaluate is the expression of canonical markers for mDA neurons in the

grafted TH$^+$ cells (Fig. 4i,j). We performed IHC and we identified approximately all of the grafted

DANs as LMX1A$^+$ (92.09 ± 3.37%), FOXA2$^+$ (96.66 ± 1.88%), NR4A2$^+$ (96.19 ± 2.16%) and

PITX3$^+$ (85.14 ± 1.80%). Finally, we performed another series of IHC to evaluate expression of

proteins involved in DA regulation and secretion (Fig. 4k,l). We were able to determine that the

grafted DANs expressed DAT (90.03 ± 1.23%), GIRK2 (96.78 ± 2.12%), VMAT (94.24 ± 1.47%)

and DDC (100.00 ± 0.00%). Overall, these results show that grafted cells mature into a large

percentage of DANs expressing canonical markers for A9-subtype mDA neurons that are able to

ameliorate motor degeneration in rodent PD models.

snRNA-seq reveals A9-subtype mDA neuron enrichment in grafts

To unbiasedly characterize graft compositionand transcriptional fidelity to the transplanted

neurons to native human neuron subtypes, we performed snRNA-seq on graft tissue from three

transplanted mice. After quality control, we retained 9196 high-quality human nuclei.

Unsupervised clustering revealed that the graft composition consisted of four main cell lineages:

neurons, radial glia, gliobasts and oligodendroglial lineage cells, with no presence of vascular

leptomeningeal meningeal cells (VLMC) contamination, an off-target population in mDA neuron

differentiation[41] (Fig. 5a, Extended Fig. 7). Further analysis of the neuronal clusters (defined by

RBFOX3) resolved into four distinct populations: dopaminergic (DAergic), glutamatergic,

GABAergic ,and a small cluster of neural progenitors (defined by high expression of INA)

(Extended Fig. 8a, Extended Fig. 9a).

Critically, the DAergic cluster was comprised of 3 distinct subclusters: DA_SOX6, DA_CALB1

and DA_ONECUT2 (Fig. 5b). DA clusters exhibited high expression of canonical dopaminergic

markers (*TH, DDC, NR4A2, LMX1A, PBX1, SLC18A2* [VMAT2], *SLC6A3* [DAT], *PITX3, KCNJ6*

[GIRK2]), whereas GABA and glutamatergic clusters lacked these but expressed *GAD1* and *GAD2*

and *SLC17A6* [VGLUT2], respectively. INA clusters were distinguished from glutamatergic

clusters by elevated *SLC17A6*, *STMN2* and *INA* expression (Extended Fig. 8b,c, Extended Fig.

9b,c) . Within the DA lineage, subclusters were resolved by differential expression of SOX6,

CALB1 and ONECUT2 (Fig. 5c,d). Roughly half of DA cells were classified as DA_SOX6 (44.73

± 4.79%), with the remainder predominantly DA_CALB1 and a minority DA_ONECUT2 (Fig.

5e). These results confirm our protocol's capacity to produce PD-susceptible SOX6$^+$ mDA subtype

in vivo.

Transcriptomic comparison validates fidelity of grafted neurons to native human midbrain subtypes

To rigorously benchmark the identity of our grafted dopaminergic neurons, we compared their

transcriptional profiles directly to those of native adult human mDA cells. We performed a

MetaNeighbor analysis, a quantitative method for assessing cell type similarity, comparing our

graft-derived DA subclusters to the well-annotated dopaminergic subtypes from a reference single

cell atlas of the human substantia nigra[19] (Fig. 5f). This analysis showed high distinctive

resemblance with a score >0.8 between our DA_SOX6 subcluster and the SOX6_AGTR1

subcluster from Kamath dataset, which have been reported to be the most susceptible

subpopulation to PD[19,40]. In contrast, the DA_ONECUT2 subcluster showed resemblance to the

human CALB1_TRHR reported to be the most resilient. However, the DA_CALB1 subcluster

lacked exclusive alignment with any Kamath cluster, instead showing broad similarity to multiple

CALB1$^+$ subtypes, indicating incomplete subtype commitment or ongoing differentiation in this

population. These results confirm successful *in vivo* differentiation into SOX6$^+$ mDA neurons

mirroring PD-vulnerable human counterparts reinforces that we were successfully able to generate

cells that differentiate *in vivo* to cells resembling human SOX6 mDA neurons.

Gene regulatory network analysis confirms stable commitment of SOX6$^+$

dopaminergic neurons

Given the transcriptional similarity of grafted DA neurons to native human subtypes, we assessed

whether SOX6$^+$ subtype represented a stably committed fate or a transitional state. GO analysis

revealed distinct functional specializations among the dopaminergic subclusters, consistent with

advanced maturation (Fig. 5g). The DA_SOX6 population was enriched for terms related to

synaptic transmission and organization, hallmarks of functional, integrated neurons. In contrast,

the DA_CALB1 and DA_ONECUT2 clusters were associated with terms like "learning" and

"regulation of cell projection," suggesting nuanced, subtype-specific functional roles. (Fig. 5g).

To directly infer developmental potential, we performed RNA velocity analysis on our DA and

immature neurons subclusters. We found that immature neurons have the potential to become

DA_CALB1 cells. Interestingly, the DA_CALB1 subclusters have the potential to become either

DA_ONECUT2 or DA_SOX6, which in contrast their velocity show that these identities were a

stable endpoint, not a transient phase (Fig. 5h, Extended Fig. 9d,e, Extended Fig. 10).

Further evaluation of the gene regulatory network by SCENIC established that the DA clusters

were enriched in exclusive TFs expression profile, not found in the immature neuron, GABA or

Glut clusters (Fig. 5i, Extended Fig. 9f). For example, PBX1 regulon networks have been

associated with ventral midbrain DA organization and is highly enriched (score >0.8) in our DA

cluster. Moreover, the DA_SOX6 is differentially highly enriched in SOX6 (score >0.8) and

CUX2 (score >0.6) within the DA clusters. Interestingly, the CUX2 regulon network has been

reported to be highly associated with the SOX6$^+$AGTR1$^+$ clusters of neurons in human datasets,

further validated in PD macaque models[19,42]. In conclusion, this multi-faceted computational

analysis demonstrates that our grafts contain mature, synaptically active neurons whose subtype

identities are locked in by stable gene regulatory networks. The DA_SOX6 population, in

particular, represents a terminal, committed fate, confirming that our protocol generates the

precise, stable, and vulnerable neuronal subtype required for accurate disease modeling and

effective cell therapy for PD.

Fig. 1: Single nuclei multi-omic sequencing revealed the diversity of cellular composition

 during mDA neuron differentiation *in vitro*.

a, Schematic diagram of the initial differentiation protocol used on the iPSC. **b,** UMAP of time-course

samples profiled with simultaneous snRNA-seq and snATAC-seq showing clustering of cells based on both

transcriptomic and chromatin accessibility profiles. **c,** Annotated bimodal UMAP clusterings based on

major cell types. **d,** Subsets of cell types were manually annotated with top-ranked marker genes and their

enriched GO terms. **e,** Dot plot visualizing the expression levels and the percentage of cells expressing

curated key genes across cell types during mDA neuron differentiation. **f,** Heatmap of Area Under the

Receiver Operating Characteristic Curve (AUROC) scores comparing annotated cell types derived from in

vitro differentiation in this study with those from a publicly available first-trimester human midbrain dataset.

g, Normalized chromatin accessibility tracks at genomic loci of genes of interest (SOX6, LMX1A and

DDC), along with enriched accession peaks and links. **h,** UMAP clustering showing RNA velocity

projection in mDA-related cell types (left), and a heatmap displaying gene expression dynamics of cluster

driver genes ordered by their inferred latent time (right). **i,** Regulon specificity score (RSS) plots

highlighting the top five cell type-specific transcription factors involved in mDA differentiation.

Fig. 2: Prolonged ventralization promotes accelerated differentiation of mDA neurons.

a, Schematic of the optimized differentiation protocol, illustrating the application of pharmacological compounds and growth factors. Representative immunocytochemistry images and quantification from both original protocol and new protocol showing **b,e**, efficacies of midbrain floorplate patterning at DIV11, showing co-localizations of LMX1$^+$/EN1$^+$ and SOX6$^+$/OTX2$^+$; and EN1$^+$/SOX6$^+$/TH$^+$ neurons at **c,f,** DIV 16 and **d,g,** 28 . White scale bars = 1 mm in B, 50 µm in **c,d** (n = 3 biological replicates). **h**, Calcium

330   imaging recordings of spontaneous and KCl-induced activities on DIV 28 and 42, presenting the total

331   number of calcium events per neuron recorded, and the percentages of cells responding to KCl-induced

332   depolarization. Metrics of activities, including amplitude, rise time, and fall time of calcium signals, are

333   depicted in the subsequent bar plots. Each dot represents one cell. Two-tailed T-tests in **e-h** were performed.

334   * = $p<0.05$; ** = $p<0.01$; *** = $p<0.001$. Error bars = SEM.

Fig. 3: snRNA-seq confirms enhanced molecular patterning under the optimized protocol.

a, UMAP embeddings of mDA differentiation in vitro using KOLF2.1J iPSCs under the new protocol, color

coded by different time points at DIV 14, 16 and 28. **b,** UMAP embeddings of detailed cell types were

manually annotated using top-ranked marker genes and their enriched GO terms, and proportion of cells

classified into each cell type per differentiation time point. **c,** Dot plot visualizing the expression levels and

the percentage of cells expressing curated key genes across distinct cell types generated under the new

protocol. **d**, Pseudotime trajectory analysis showing progression from progenitor-like populations (ProgFP,

NProg, and NbM) to differentiated DA neurons. Cells are ordered by inferred pseudotime along the x-axis.

e, Regulon specificity score (RSS) plots highlighting the top 5 cell type-specific transcription factors driving

mDA differentiation under the new protocol. (f) Dot plots showing the expression levels and the percentage

of cells expressing midbrain dopaminergic lineage marker genes after integration in (left) midbrain cells

categorized as dopaminergic neurons (DA) (right), dopaminergic lineage cell types across all timepoints

and both conditions.

351 Fig. 4: Transplanted neurons restore motor function and mature into A9-subtype mDA

352 neurons *in vivo*.

a, Diagram of the experimental design for the implantation of day 14-differentiated hiPSC in B6RGS mice.

b, Motor recovery is observed by continuous analysis of amphetamine-induced ipsilateral rotation, endpoint

tape removal test, endpoint pole test and endpoint cylinder wall analysis (right). Of note, grafted animals

recover pre-lesion state while the 6-OHDA control animals maintain the lesioned state. **c**, Example of

grafted tissue, stained against for STEM121, showing the human cells within the graft. **d,** Micrograph of

graft stained against for TH and HuNu, showing hDANs all over the graft. **e,** Closeup of grafted hDANs

showing elongated and ramified extensions. **f,** Stereological estimate of HuNu+ nuclei (left) and percentage

of hDANs (TH+-HuNu+) in grafts (right). **g,** Representative examples of hDANs stained positive for A9-

subtype marker, SOX6, ALDH1A1, AGTR1A and A10 marker, Calbindin. **h,** Quantification of percentage

of hDANs (TH+-HuNu+) expressing SOX6, ALDH1A1, AGTR1A and Calbindin. **i,** Representative

examples of hDANs stained positive for mDA neuron makers, LMX1A, FOXA2, NR4A2 and PITX3. **j,**

Quantification of percentage of hDANs (TH+-HuNu+) expressing LMX1A, FOXA2, NR4A2 and PITX3.

k, Representative examples of hDANs stained positive for proteins involved in DA excretion, DAT, GIRK2,

VMAT2 and DDC. **l,** Quantification of percentage of hDANs (TH+-HuNu+) expressing DAT, GIRK2,

VMAT2 and DDC. m.p.i = months post-implant. n for animal behavioral experiment **b** = sham: 6; 6-OHDA:

8; hIPSC-grafted: 8. N for IHC **f,h,j,l**, = 5. Scale bar = **c**: 1mm; d: 500 μm; **e**: 100 μm; **g,i,k**: 10 μm. Means

are represented as bars (**b**) or lines (**f,h,j,l**). One-way ANOVA with Tukey correction between treatments

was performed separately on each timepoint in the continuous amphetamine test ($F_{12,114}$ = 7.876), as well

as for each separate timepoint in the endpoint experiments (Pole test $F_{4,38,}$ = 27.32; Tape Removal $F_{4,38}$ =

8.40; Cylinder Wall $F_{4,38}$ = 3.97). * = p<0.05; ** = p<0.01; *** = p<0.001. Error bars = SEM.

Fig. 5: snRNA-seq of grafts reveals presence of transcriptionally-defined A9-subtype

neurons

a, Annotated UMAP embeddings of 9,196 high-quality human nuclei recovered from three grafted mice at

five months post-transplantation, with cells classified into neurons, radial glia, gliobasts and

oligodendrocyte precursors; and **b**, neuronal population was consisted of immature neurons, GABAergic

neurons, Glutamatergic neurons and 3 subtypes of mDA neurons, each characterized by specific

transcription factors, as DA_CALB1, DA_ONECUT2, and DA_SOX6. **c,** Dot plot showing average gene

expression and percentage of expressing cells for selected marker genes across annotated neuronal cell

types. **d**, Violin plots of representative marker genes for dopaminergic subtypes (SOX6, ONECUT2,

CALB1), pan-DA markers (TH, NR4A2, KCNJ6), illustrating subtype-specific and shared expression

patterns. **e**, Average proportion of yielded human nuclei in each cellular hierarchy across three individual

datasets, demonstrating consistent representation of DA subtypes (N=3). **f**, MetaNeighbor analysis between

transplanted mDA neurons in this study and a reference dataset of DA neurons from the SNpc region of PD

patients and matched controls, suggesting transcriptional similarities between DA_SOX6 subtype and

SOX6_AGTR1 subtype from the reference dataset. **g,** GO enrichment analysis of DA subtype-specific

markers, showing distinct biological processes enriched in DA_SOX6, DA_CALB1, and DA_ONECUT2

populations. **h**, RNA velocity projection of mDA neuron trajectories in UMAP embeddings, suggesting

differentiation pathways among different DA subtypes and immature neurons. **i**, Heatmap of average

regulon activities of the top 30 transcription factors across mDA and other neuronal subtypes.

Discussion

This study establishes a multiomic-guided framework for differentiating human iPSCs into mDA

neurons with unprecedented molecular and functional fidelity to the PD-vulnerable

$SOX6^+AGTR1^+$ subtype of the substantia nigra. By leveraging single-cell transcriptomic and

epigenomic data to identify and overcome a key bottleneck (insufficient ventral patterning), we

developed an optimized protocol that achieves accelerated, reproducible production of these

therapeutically critical neurons across multiple cell lines.

Our longitudinal multi-omic atlas of *in vitro* mDA differentiation revealed stage-specific

chromatin accessibility and transcriptional shifts mirroring human midbrain development. Late

maturation (DIV42-56) coincided with elevated TH, DDC and KCNJ6 expression, and PBX1

regulon activity, a known regulator of mDA survival and functional maturation. Crucially,

transcriptome-informed ventralization enhancement (via strategic agonism of SHH signaling)

boosted EN1 expression, a key regulator of midbrain identity and a known predictor of transplant

success, yielding ventral SNpc neurons with enhanced molecular fidelity to human native

counterparts. Furthermore, we were able to identify regulatory networks involving mDA neuron

maturation enriched in the new protocol, such as TCF12 and TCF4. Specifically, TCF12 plays a

dual role in the early commitment of neural progenitors and the late specification of neuronal

subsets, with its loss leading to altered expression of TH and AHD2 (ALDH1A1), the latter being

crucial for the specification and maintenance of dopaminergic neurons, particularly in the A9

dopaminergic neuronal group[43]. TCF4, on the other hand, is likely involved in the fine-tuning of

neuron identity during the onset of mDA neuron subtype specification. In a mouse study, deletion

of TCF4 led to delayed maturation of dopaminergic neurons, primarily through transient

suppression of *Ahd2* and *Nxph3*[44]. A key advancement in the optimized protocol was the precise

modulation of midbrain floor plate (mFP) patterning by the biphasic WNT activation (CHIR boost)

and SSH agonism, combined with optimized coating conditions using LN511, which generated a

substantial population of mFP progenitors by day 11 and enhanced the expression of critical mDA

markers such as FOXA2, LMX1A, and OTX2. This improved patterning was accompanied by a

reduced contamination by non-neural cells (a common issue in region-specific neuron generation).

Furthermore, the specification to committed progenitors allowed us to omit the MEK/ERK

inhibitor PD0325901 and the FGF receptor inhibitor SU5402, previously used to suppress

expansion of SOX2[+] progenitors[45], therefore harnessing the intrinsic neuronal differentiation

mechanisms without any further external influence for producing mature mDA neurons

downstream. The addition of synthetic LXR ligands (GW3965) further accelerated neurogenesis,

leading to the earlier appearance of TH[+] neurons at day 16 compared to day 21 in prior protocols.

This earlier establishment enables functional assays (such as calcium imaging and neurotransmitter

release analysis) to be performed sooner, thereby expediting detection of phenotypic differences.

As an example, live-cell calcium recordings confirmed that the derived mDA neurons displayed

spontaneous electrical activity and robust depolarization responses, consistent with functionally

active neurons[46]. This protocol also proved reproducible across multiple iPSC lines, including

those from Parkinsonian patients, highlighting its potential as a standardized approach for

generating patient-targeted high-quality dopaminergic neurons, a crucial step for modeling

neurodegenerative diseases and testing therapeutic strategies. Finally, as a proof-of concept

validation for therapeutic approaches, we determined that the new protocol could produce mFP

progenitors and NbM able to differentiate *in vivo* to mDA neurons resembling the A9-subtype

nigral neurons in majority, based on expression of proteins such as SOX6, AGTR1, GIRK2 or

ALDH1A, reported to be more susceptible to PD in human and animal models[19,40,47,48].

Furthermore, evaluation of the transcriptomic profile of the DA neurons in the graft showed high

resemblance to the human counterpart, specially the SOX6-AGTR1A mDA neurons. Moreover,

the lack of proliferative progenitor cells at our endpoint is promising for putative long-term

transplants. This proposes that these cells could have great potential for directed replacement

therapies. While these advancements could be of great interest to the PD research community, we

have to address several caveats. First, we only have established the replicability of this protocol

on hIPSC, both WT and with GBA mutations. But further in-depth characterization with more well

known mutations found in familial PD, such as GBA^{E326K}, LRRK2^{G2019S} or the triplicate SNCA,

could elucidate the validity of this enhanced protocol. Furthermore, the use of human embryonic

stem cells has not been proposed for this study. More *in vitro* studies would be of interest to

validate further this new protocol. Another caveat to address is the pre-clinical aspect, as the *in*

vivo experiments were only used as proof-of-concept for therapeutical approaches. Further grafting

experiments with PD-mutated hiPSC cell lines and a more in-depth preclinical analysis could detail

if this protocol would be a much better advantage that the current available protocols, which so far

have shown to be safe in clinical trials[12,13]. Moreover, the longevity of the grafting experiment was

decided upon stable motor recovery, so we do not have an overview of the long-term effects of

these grafts, whether other contaminants can appear or tumorigenesis in aged individuals. In

conclusion, we have developed a multiomic-guided differentiation protocol which addresses the

major challenge of subtype specificity in generating human mDA neurons relevant to PD. By

leveraging single-cell genomics and epigenomics to identify and correct insufficient ventral

patterning, we achieved robust enrichment of the PD-vulnerable SOX6$^+$ A9 subtype, which yielded

neurons with enhanced molecular fidelity, functional maturity, and the capacity to restore motor

function upon transplantation. Our work provides a robust platform for generating more accurate

human cellular models of PD and establishes a foundational strategy for developing next-

generation, subtype-enriched cell replacement therapies.

Methods

Cell lines

KOLF 2.1J cell line was obtained through Jackson Laboratories[49]. GBA1[E326K] heterozygous PD

patient-derived iPSCs (NH50143) along with its edited isogenic control iPSCs (NH50142) were

obtained from the Rutgers University Cell and DNA Repository (RUCDR). All experiments

performed with these cells followed the Ethical Permit 2021-06727-01 issued by the Swedish

Ethical Review Authority. Karyotipe by G-banding, genotypes by polymerase chain reaction

(PCR), pluripotency by pluripotency marker microscopy analysis, and mycoplasma infection by

PCR were routinely assessed before starting new experiments. Applied iPSC lines were adapted

to grow on Laminin 521 (Biolamina, LN521)-coated plates and maintained in Essential 8 Flex

media (E8 Flex, Thermo Fisher Scientific) at 37ºC, 5% CO_2, until starting differentiation. The

medium was changed daily, and cells were passaged at 70% confluency using TrypLE™ Select

Enzyme (Thermo Fisher Scientific) at a 1:20 ratio. When thawing and replating, 10 µM Rho-

associated protein kinase (ROCK) inhibitor (Y-27632) (Tocris) was added into the culture media

overnight.

Midbrain dopaminergic neuron differentiation

Two days prior to the mDA differentiation, adapted iPSC cells were seeded at a density of 200,000

cells/cm2 onto plates coated with Geltrex (Life Technologies) and Laminin-511 (Biolamina,

LN511), and maintained in E8 Flex media (Thermo Fisher Scientific) for expansion. The

differentiation protocol was initiated when the cells reach approximately 80% confluency. For the

first iteration, cells were cultured at day 0-4 on plates coated with Geltrex and LN511, replated at

day 11 on LN511, and replated again at day 16 on LN511 + poly-L-ornithine. Cultured cells were

kept in Neurobasal/N2/B27 (Life Technologies) medium and 2 mM L-glutamine (Invitrogen),

along with various supplements for the first stage of the protocol. From day 0 to day 4, the culture

medium was conditioned with 250 nM LDN193189 (Stemgent), 10 µM SB431542, 1 µM

Purmorphamine, and 0.7 μM CHIR99021 from day 0 to day 3, with the inclusion of 10 μM Y27632

(Tocris Bioscience) during the first 24 hours. Boosted CHIR99021 (7.5μM) was administered from

day 4 to day 11, while LDN193189, SB431542 and Purmorphamine were withdrawn after day 7

and N2 supplement after day 10. From day 10 to day 28, cells were switched to the differentiation

media: Neurobasal/B27/L-Glu media supplemented with 20 ng/mL brain-derived neurotrophic

factor (BDNF, R&D Systems), 20 ng/ml glial cell line-derived 455 neurotrophic factor (GDNF,

R&D Systems), 1 ng/ml transforming growth factor type b3 (TGFb3, R&D Systems), 0.2 mM

ascorbic acid (Sigma), 0.2 mM Dibutyryl cAMP (Sigma). On day 11 and 16, the resulting cells

were dissociated into single cells, replated at a density of 800,000 cells/cm2 onto plates coated

with LN511 and polyornithine/LN511 respectively, and treated with 10 μM Y27632 for 48 hours.

To promote neurogenesis, the mDA differentiation media was supplemented with 10 μM DAPT

(Sigma) from day 16. The small molecule GW3965 (10 μM) was administered from day 12 to day

15, followed by a cocktail of small molecules consisting of 1 μM PD0325901 and 5 μM SU5402

(Sigma), from day 16 to day 21. For the second iteration, conditions were similar except for the

following: Culture plates were coated with Geltrex + LN511 until day 14, and poly-L-ornithine +

LN511 from day 14 onwards. Cells were only replated once at day 14 and boosted CHIR99021

(7.5μM) was administered from day 4 to day 14. Day 0-7 Purmorphamine concentration was

increased to 2 μM, and both PD0325901 and SU5402 were removed from the protocol.

Immunocytochemistry

Progenitor cells at different stages were seeded in a 96-well plate at a density of about 300,000

cells per well for the intended cell culture purposes. Harvested cells were fixed by 4%

paraformaldehyde (PFA) for 20 min at 4ºC, and then pre-incubated by 5% donkey serum in

phosphate buffered saline containing 0.3% Triton X-100 (PBST) for 1 hour. The samples were

incubated with primary antibodies at 4ºC overnight. The samples were incubated with either

Alexa488 or Alexa555 or Alexa647-conjugated secondary antibodies (Thermo Fisher Scientific)

for 30 min and then were incubated with 4',6-diamidino-2-phenylindole (DAPI, Sigma, D9542)

for 15 min. The primary antibodies were used as follows: CORIN (rat, 1:1,000, R&D Systems,

MAB2209), DCX (goat, 1:500, SantaCruz, sc-8066), EN1 (mouse 1:50, DSHB, 4G11), FOXA2

(goat, 1:500, R&D Systems, AF2400), GIRK2 (rabbit, 1:400, Alomone, APC006), LMX1A

(rabbit, 1:4,000, Millipore, AB10533), LMO3 (goat, 1:200, SantaCruz, sc-82647), MAP2 (mouse

1:1,000, Sigma, M4403), NURR1 (rabbit, 1:500, SantaCruz, sc-990), OTX2 (goat, 1:1,000, R&D

Systems, AF1979), TH (rabbit, 1:1,000, Millipore, AB152), TH (mouse, 1:500, ImmunoStar,

22941), and TH (sheep, 1:500, Novus Biological, NB300). Mounted cells were further visualized

using confocal microscopy (Zeiss LSM880 or 980) with a 10X or 20X lens. Quantification of

images was performed using QuPath[58] by generating a "single measurement classifier" or

"composite classifier" for positive or double-positive cells respectively.

Calcium imaging

The 35mm culture dishes were loaded with the $Ca2^+$-sensitive fluorescence indicator Fluo-4/AM

(5 µM; Invitrogen, USA) and Pluronic F-127 (0.625%, ThermoFisher Scientific) and incubated

for 30 min at 37ºC in Krebs-Ringer buffer, containing: NaCl (119 mM), KCl (2.5 mM), NaH2PO4

(1 mM), CaCl2 (2.5 mM), MgCl2 (1.3 mM), HEPES (20 mM) and D-Glucose (11 mM), with pH

adjusted to 7.4. Dishes were washed twice and $Ca2^+$-measurements were performed in Krebs-

Ringer buffer at 37ºC on a confocal spinning disk microscope (Nikon CrEST X-Light V3)

equipped with a 20x Air lens. Excitation was assessed at 480 nm at a sampling frequency of 1 Hz.

The equipment was controlled, and data was acquired using the Nikon NIS-Elements software.

FIJI, MATLAB (R2020b, MathWorks, USA) and FluoroSNNAP (https://github.com/tapan-

patel/FluoroSNNAP) were used to process and analyze the data.

Single cell sequencing libraries preparation and data processing

A total of 5 samples (DIV 11, 16, 28, 42, and 56) from KOLF2.1J mDA neuron cultures generated

with the optimized protocol were collected for time-course multiomic analysis (ATAC + Gene

Expression); in addition, 3 samples (DIV 14, 16, and 28) from the new protocol trial of KOLF2.1J

mDA differentiation were collected for single-nuclei transcriptomic profiling. All 8 samples were

prepared with commercially available protocols (10x Genomics) and the resulting Fastqc files

were processed through the 10X Genomics Cellranger pipeline[50], while 5 multiomic libraries were

aligned to the hg38 reference genome using Cellranger-arc default settings[51]. After quality-control

filtering, we obtained high-quality 23,534 cells (Fig. 3a, Extended Fig. 5a) and then we integrated

transcriptomic profiles from both conditions using Harmony[52], enabling a comparative analysis by

isolating the mDA neuronal lineage from different datasets (Extended Fig. 5b). Resulting

transcriptomic and epigenomic data from individual samples were combined by projects using the

'aggr' function in Cellranger or Cellranger-arc, without normalization applied. Objects produced

with Seurat V4.4.0 were employed to accommodate aggregated count matrix[53], with the fragment

files specifically integrated into Signac v1.9.0 chromatin assays[54]. Quality metrics for gene

expression and peak accessibility, such as mitochondrial ratio, cell cycle score, nucleosome signal,

and Transcription Start Site (TSS) enrichment, were thereafter calculated to determine quality

control cutoffs. Depending on sample quality, following thresholds were applied for downstream

analysis: a maximum of 300,000 and a minimum of 1,000 ATAC counts per cell, a maximum of

25,000 and a minimum of 600 RNA counts, at least 250 genes detected, a maximum nucleosomal

signal of 1.5, TSS enrichment of at least 2, and mitochondrial content no higher than 15%. These

cutoffs reduced the cell count by less than 20% without significantly affecting clustering

downstream. Peak calling was performed with MACS2 v2.2.9 using the aggregated fragments

file[55], and those mapped to non-standard chromosomes or overlapping hg38 genome blacklist

regions were excluded. DoubletFinder v2.0 was applied to identify doublets in gene expression

data by flagging instances with a high resemblance to artificial neighbors in gene expression

space[56]. AMULET (ATAC-seq MULtiplet Estimation Tool) was additionally utilized to detect

potential multiplets in snATAC-seq datasets by enumerating regions with greater than two

uniquely aligned reads across the genome[57]. All predicted doublets were removed from the

downstream analysis. After merging high-quality cells from the relevant projects, count data was

normalized using SCTransform V2 function with regression out of mitochondrial or cell-cycle

genes. The top 2,000 most variable genes were initially identified through variance-stabilizing

transformation, along with 95% of the most frequent peaks in multiomic datasets, as a part of the

significant feature selection process. Subsequent dimensionality reduction was performed using

the principal components analysis (PCA), in which the top 30/40 principal components were

chosen from the elbow plot to represent most of the diversity of datasets. For single-nuclei ATAC-

seq datasets, singular value decomposition (SVD) was performed with the top 25 of the Latent

Semantic Indexing (LSI) layers to reduce the dimensions of the most common peaks. Notably, the

first LSI component was excluded from downstream analysis as it often captures sequencing depth

(technical variation) rather than biological variation. Putative clusters were identified by a spiking

neural network (SNN) modularity optimization-based clustering algorithm, and the optimal

resolution of clustering was determined by iterating over different resolution for the most stable

segregation. Finally, the calculated nearest-neighbor graph was projected to two dimensions to

visualize the distribution of instances by representing with uniform manifold approximation and

projection (UMAP). To accurately identify and interpret cell states *in vitro*, we leveraged

combined manual cell type annotation using cluster-specific marker genes[58,59], with automatic

annotation methods using reference datasets that characterize first-trimester developing human

brain[14,60]. Initially, a label transfer was performed with Celltypist[61] and 'TransferData' function in

Seurat to automatically classify selected dataset with cell type labels from the reference dataset,

based on the similarity of query gene expression profiles to selected anchors in the reference

dataset. The identity of each cell will be determined by the highest prediction score, and for

stringent comparison, cells with a prediction score below 0.51 will be excluded from downstream

analysis. After data trimming and reclustering, a manual cluster-based annotation was performed,

using both expression levels of canonical midbrain development marker genes and top 50

upregulated genes identified by the 'FindAllMarkers' function. These upregulated genes were used

for pathway and biological process enrichment analysis via Metascape[59], and the top-ranked terms

were used to refine annotations and determine cell subtypes or states. To reconstruct the putative

trajectory of mDA differentiation *in vitro*, we employed Monocle3[62] and scVelo[63,64] to infer

pseudo- and latent-temporal ordering of cells using default parameters[65]. The root of trajectory for

pseudotime analysis was set as ProgFP or NbM at the earliest timepoint. at the earliest timepoint.

To represent quantitatively the cell identity similarities with other published human datasets, we

performed Metaneighbor[66]. SCENIC[67] was used to visualize regulatory networks and regulon in

our developing cells using AUCell scores.

Animals

Healthy female adult (8-12 weeks old) C57BL/6N-Rag2Tm1-IL2rgTm1-SirpaNOD/Rj (B6RGS)

mice were purchased from Janvier Labs. Groups of 5 animals were housed in ventilated cages

under a 12 h light-dark cycle with food (standard chow) and water ad libitum at 20-22ºC and 50-

55% humidity. Tunnels, cardboard nesting material and chewing wooden bars were used for

environmental enrichment. All surgical procedures followed aseptic techniques, and for post-

surgical recovery, additional Dietgel Recovery (Scanbur) and food pellets were supplied on the

cage floor. Animals were randomly allocated to sham control group (n=6), hemiPD group (n=8)

and a grafted group (n=8), ensuring all cages had individuals of each group. Briefly, the exclusion

criteria were animals which lost over 10% of their pre-surgical weight after 4 weeks were discarded

from the experiment. No animals were discarded based on this criterion. The experimental unit

were the individual mouse. All procedures were approved by the local ethical committee

(Stockholms djurforsoksetiska namnd) and followed the EU directive 2010/63/EU under the

ethical permit of Ernesto Arenas Cases and Sandra Gellhaar (12265-2021; 2-3143/23; 2-3595/24;

2-3147/24).

6-OHDA mouse model

B6RGS mice were acclimatized 2-3 weeks before surgery to the animal facility and the researcher

and to a recovery diet 2-3 days before surgery. For 6-hydroxidopamine (6-OHDA, Sigma) lesions

(or sham), animals were injected analgesia (Carprofen 5mg/kg + Buprenorphine 0.1 mg/kg) 15

minutes before isoflurane induction (4%). Animals were anesthetized at 1-2% isoflurane in air and

injected intracranially (ICR) using a Neurostar Robotic Stereotaxic Frame equipped with a 10 µl

syringe with 33G blunt needle (Hamilton) 1 µl of 6-OHDA 3.6 µg/ µl in sterile ascorbic acid/saline

0.02% (or sham vehicle) in the right medial forebrain bundle [respect to Bregma: AP -1.2 mm,

ML +1.2 mm, DV -5.0 mm] at a flow of 0.2 µl/min, raising the needle 500 µm for 3 minutes. The

needle was left inside for 2 additional minutes. Animals were placed in warm cages (28 °C)

between 3 days and 1 week for better recovery and rehydrated with Ringer Acetate solution if

weight decreased by 5%. 6-OHDA hemiPD was confirmed by amphetamine-induced rotation.

Briefly, 4 weeks after the lesion, mice were injected intraperitoneally with d-amphetamine in saline

(5 mg/kg). 10 minutes after injection, the mice were recorded for 30 minutes inside a 15 cm wide

cylinder in dark conditions, and the number of net ipsilateral rotations (ipsilateral-contralateral)

per minute in active movement was calculated using Ethovision XT 17 (Noldus). Mice are

considered hemiPD if the values are >6 rotations/minutes.

hiPSC transplant

HemiPD-confirmed mice of healthy weight (22-30 g) were randomly selected for ICR hPSC

injection in the right striatum [respect to Bregma: AP +0.5 mm, ML +2 mm, DV -3.5 mm].

Analgesia and anesthesia were followed as above. For ICR transplants, 2 µl of 100.000 ± 20.000

hPSCs/µl in Neurobasal medium was injected to achieve 200.000 hPSC transplants. For cell

injection, the Neurostar robot was coupled with a 10 µl syringe with a 27G sharp syringe. A flow

of 0.5 µl/min was used, and the needle remained for 5 additional minutes. Cells were injected

between 1-6 hours after thawing.

Amphetamines-induced rotations

Mice were analysed a week before surgery, 4 weeks after surgery and hereon once a month until

endpoint was reached. Mice were injected intraperitoneally with 5 mg/kg d-amphetamine in saline

(0.5 mg/ml). 10 minutes after injection, the mice were recorded for 30 minutes inside a 15 cm wide

cylinder in dark conditions, and the number of net ipsilateral rotations (ipsilateral-contralateral)

per minute in active movement (body center traveling at >0.5 cm/s) was calculated using

Ethovision XT 17.

Pole test

Mice were analysed a week before surgery, 4 weeks after surgery and at the endpoint. Animals

were placed facing upwards on the top of a 20 cm vertical smooth wooden pole with 2 mm grooves

notched every 0.5 cm down. The time to climb down was counted starting when the mouse rotates

facing downwards. Animals were trained for a minimum of 2 days before testing day, by

performing 3 mock tests each day. If the animals took more than 1 minute to descend, the trial was

interrupted and repeated at least 10 minutes later. A total of 3 trials were used to calculate a mean

value per animal.

Cylinder wall

Mice were analysed a week before surgery, 4 weeks after surgery and at the endpoint. Animals

were placed inside a 12 cm diameter glass cylinder in red light and quiet conditions. For 15

minutes, animals were recorded and videos were analysed at x8 speed. Unilateral full paw contacts

were counted. Paw dragging or paws not fully placed on glass were disregarded.

Tape removal test

Mice were analysed a week before surgery, 4 weeks after surgery and at the endpoint. A green

adhesive paper tape 2x3 mm was placed on each paw individually. The time it took to remove the

tape was counted, beginning when the mouse recognized it and ending when it was removed from

the paw. Net contralateral time (ipsilateral-contralateral) was calculated to better visualize and

eliminate external factors. Before testing day, animals were trained for a minimum of 2 days,

performing 3 mock trials for each paw. If the animal took more than a minute, the trial was

interrupted and repeated at least 15 minutes later. A total of 3 trials were used to calculate a mean

value.

Immunohistochemistry

5 animals were terminally anesthethized with IP pentobarbital 100 mg/kg dose and then perfused

intracardially with 50 ml of pH-corrected PBS1X followed by 50 ml of formaldehyde 4 % (FA)

both at a flow of 9 ml/min. Brains were extracted and placed in FA overnight, after which they

were placed in sucrose 30%/PBS solution at 4ºC for 2-3 days. After, brains were kept at -80ºC.

Brains were equilibrated at -20ºC for 4 hours before slicing. Samples were sliced in series at -20ºC

at 20 µm using a Cryostar NX70 (Epredia) and stored for further use at -20ºC in cryoprotective

solution (Ethylenglycol 30%, Glycerol 30%, PBS1X). A total of 5 brains were processed.

Samples were stained using free-floating protocols. Briefly, 2 series of 5 minutes of PBS1X,

followed by 15 minutes of PBS-TritonX 0.3% (PBTx0.3%), then 2 series of 5 minutes of

PBTx0.1%. Samples were blocked for 1 hour with blocking solution (BSA 1 mg/ml, Normal

Donkey Serum 5%, PBTx0.1%) and incubated o/n at 4ºC with primary antibody diluted in

blocking solution. Primary antibody was washed with 2 series of 5 minutes PBTx0.1% and

samples were incubated for 4 hours with secondary antibody diluted in blocking solution and

washed twice for 5 minutes with PBS1X. Tissue was counterstained with DAPI 1:1000 in PBS1X

for 10 minutes and then washed twice for 5 minutes with PBS1X. Stained samples were mounted

on Superfrost Plus slides (Epredia) and Fluoromount-G mounting medium (00-4958-02, Thermo

Scientific). Primary antibodies used were STEM121 (mouse, 1:1000, Tanaka Bio, Y40410), TH

(rabbit, 1:1000, Pelfreeze, P4010; sheep, 1:250 Pelfreeze, P60101), HuNu (mouse, 1:1000,

Millipore, MAB1281), SOX6 (rabbit, 1:200, Abcam, ab30455), ALDH1A (rabbit, 1:500, Atlas

Antibodies, HPA002123), AGTR1 (rabbit, 1:100, Invitrogen, MA5-56587), Calbindin-1 (rabbit,

1:200, Invitrogen, PA5-85669), LMX1 (rabbit, 1:100, Millipore, AB10533), FOXA2 (rabbit,

1:200, Cell Signaling, 3143S), NR4A2 (rabbit, 1:500, Invitrogen, PA5-78097), PITX3 (rabbit,

1:100, Invitrogen, 35-2850), DAT (rat, 1:250, Abcam, ab5990), GIRK2 (rabbit, 1:500, Alomone

Labs, APC006), VMAT2 (rabbit, 1:500, Sigma-Aldrich, AB1598P), DDC (rabbit, 1:500, Novus

Biological, NB300-252), Ki67 (rabbit, 1:200, Abcam, ab15580), SOX2 (goat, 1:200, R&D

Systems, AF2018). All fluorescent secondary antibodies were from Invitrogen, donkey as host and

conjugated with Alexa Fluor 488, 555 or 647. The HRP-conjugated antibody was a goat anti-

mouse from Invitrogen. Images were captured either with laser confocal microscope Zeiss

LSM800 equipped with x20 lens or high-throughput epifluorescence microscope Zeiss Axioscan

equipped with x10 lens. Quantification of images was performed using QuPath[58] by generating a

'single measurement classifier' or 'composite classifier' for positive or double-positive cells

respectively.

Single-nuclei RNA sequencing of grafts and data analysis

3 animals were euthanized by cervical dislocation and brains were promptly extracted and flash-

frozen in dry-ice chilled isopentane for 10-15 seconds. Flash-frozen brains were stored at -80 °C

until graft extraction. A total of 3 brains with grafts were processed independently.

On the day of nuclei extraction, brains were equilibrated to -20ºC and sliced in -20ºC cooled

chambers. Brain matrices were used to stabilize the brain, and 1 mm-thick slices for a total of 4 (2

before the center of graft implant location and 2 after) were cut using chilled razors. 1 mm punches

of each slice of the dorsal striatum were extracted using 1 mm biopsy-punchers (49101, PFM

Medical).

Nuclei were dissociated using the 10X Genomics Chromium nuclei extraction kit (1000494) using

manufacturer's instructions. For better extraction, eppendorfs used for tissue dissociation were

coated with BSA 1% o/n at 4ºC, briefly washed with filtered sterile PBS1X before procedure. 10X

Genomics Chromium GEM-X Single Cell 3' Kit v4 (1000691), Chromium GEM-X Single Cell 3'

Chip Kit v4 (1000690) and Dual Index Kit TT Set A (1000215) were used for GEM generation

using manufacturer's instructions with the 10X Controller iX. GEMs were sent for RNA-

sequencing to National Genomics Infrastructure of SciLifeLab (Karolinska Institute, Sweden)

using the NovaSeq XPlus 25B. Data analysis was performed as the *in vitro* RNAseq dataset,

described above. To ensure analysis of human cells, we aligned the libraries to the h38 human

genome dataset and the m39 mouse genome dataset, and cells with a ratio of human:mouse gene

count >1 were processed.

Statistics

The N for the animals was calculated using G-Power 3.1[68] based on prior data from several

publications of hemiPD rodent model behavior[69]. Briefly, using the "ANOVA: repeated measures,

between factors" a priori statistical test, the effect size was calculated to be 0.82, α =0.05, β=0.95,

groups =3, measurements=7 and correlation among repeated measurements=0.8.

GraphPad Prism v10.00 was used for generating all graphs and performing statistical analyses. For

pairwise comparisons, Two-tailed Student's t-test was utilized. One-way ANOVA or two-way

ANOVA with Tukey's or Bonferroni's multiple comparison tests were used to compare multiple

variables. Error bars indicate the standard error of the mean (SEM). F are reported with degrees of

freedom for numerator and denominator ($F_{DFn,DFd}$). Significance levels were set as follows: ns =

not significant; * = p<0.05; ** = p<0.01; *** = p<0.001; **** = p<0.0001.

Data Availability

Single-cell RNA-seq and ATAC-seq raw data have been deposited at EGA (EGA accession

number: EGAD50000002054). Key Resource Table (including Material RRID, protocol.io links

for protocols used, accession numbers for sequencing datasets), microscopy data and images have

been deposited at Zenodo at https://doi.org/10.5281/zenodo.17099441 and are publicly available

as of the date of publication. All reused code used for this paper has been deposited at Github and

linked to Zenodo at https://doi.org/10.5281/zenodo.17099441 and is publicly available as of the

date of publication. This paper analyzes existing, publicly available data, accessible at

https://doi.org/10.1038/s41593-022-01061-1; https://doi.org/10.1126/science.adf1226;

https://doi.org/10.1126/science.add7046. Any additional information required to reanalyze the

data reported in this paper is available from the lead contact upon request.

Code Availability

No new or original code was generated for this paper.

References

1. Poewe, W. *et al.* Parkinson disease. *Nat Rev Dis Primers* **3**, 17013 (2017).

2. Brichta, L., Greengard, P. & Flajolet, M. Advances in the pharmacological treatment of Parkinson's disease: targeting neurotransmitter systems. *Trends Neurosci* **36**, 543–54 (2013).

3. Björklund, A. & Parmar, M. Dopamine Cell Therapy: From Cell Replacement to Circuitry Repair. *J Parkinsons Dis* **11**, S159 (2021).

4. Doi, D. *et al.* Isolation of human induced pluripotent stem cell-derived dopaminergic progenitors by cell sorting for successful transplantation. *Stem Cell Reports* **2**, 337–350 (2014).

5. Kim, T. W. *et al.* Biphasic Activation of WNT Signaling Facilitates the Derivation of Midbrain Dopamine Neurons from hESCs for Translational Use. *Cell Stem Cell* **28**, 343-355.e5 (2021).

6. Kriks, S. *et al.* Dopamine neurons derived from human ES cells efficiently engraft in animal models of Parkinson's disease. *Nature* **480**, 547–551 (2011).

7. Nishimura, K. *et al.* Single-cell transcriptomics reveals correct developmental dynamics and high-quality midbrain cell types by improved hESC differentiation. *Stem Cell Reports* **18**, 337–353 (2023).

8.   Kirkeby, A. *et al.* Preclinical quality, safety, and efficacy of a human embryonic stem cell-
derived product for the treatment of Parkinson's disease, STEM-PD. *Cell Stem Cell* **30**,
1299-1314.e9 (2023).

9.   Piao, J. *et al.* Preclinical Efficacy and Safety of a Human Embryonic Stem Cell-Derived
Midbrain Dopamine Progenitor Product, MSK-DA01. *Cell Stem Cell* **28**, 217-229.e7
(2021).

10.   Schweitzer, J. S. *et al.* Personalized iPSC-Derived Dopamine Progenitor Cells for
Parkinson's Disease. *N Engl J Med* **382**, 1926–1932 (2020).

11.   Takahashi, J. iPS cell-based therapy for Parkinson's disease: A Kyoto trial. *Regen Ther*
13, 18–22 (2020).

12.   Tabar, V. *et al.* Phase I trial of hES cell-derived dopaminergic neurons for Parkinson's
disease. *Nature 2025 641:8064* **641**, 978–983 (2025).

13.   Sawamoto, N. *et al.* Phase I/II trial of iPS-cell-derived dopaminergic cells for Parkinson's
disease. *Nature 2025 641:8064* **641**, 971–977 (2025).

14.   Braun, E. *et al.* Comprehensive cell atlas of the first-trimester developing human brain.
Science **382**, eadf1226 (2023).

15.   La Manno, G. *et al.* Molecular Diversity of Midbrain Development in Mouse, Human, and
Stem Cells. *Cell* **167**, 566-580.e19 (2016).

16.   Xu, P. *et al.* Human midbrain dopaminergic neuronal differentiation markers predict cell
therapy outcomes in a Parkinson's disease model. *J Clin Invest* **132**, (2022).

17.   Tiklová, K. *et al.* Single-cell RNA sequencing reveals midbrain dopamine neuron
diversity emerging during mouse brain development. *Nature Communications 2019 10:1*
10, 1–12 (2019).

18.   Greene, P. E. *et al.* Persistent dyskinesias in patients with fetal tissue transplantation for
Parkinson disease. *NPJ Parkinsons Dis* **7**, (2021).

19.   Kamath, T. *et al.* Single-cell genomic profiling of human dopamine neurons identifies a
population that selectively degenerates in Parkinson's disease. *Nature Neuroscience 2022*
25:5 **25**, 588–595 (2022).

20.   Siletti, K. *et al.* Transcriptomic diversity of cell types across the adult human brain.
Science **382**, eadd7046 (2023).

21.   Mannens, C. C. A. *et al.* Chromatin accessibility during human first-trimester
neurodevelopment. *Nature* 1–8 (2024) doi:10.1038/s41586-024-07234-1.

22.   Nishimura, K., Ásgrímsdóttir, E. S., Yang, S. & Arenas, E. A protocol for the
differentiation of human embryonic stem cells into midbrain dopaminergic neurons. *STAR*
Protoc **4**, 102355 (2023).

23.   Sif Ásgrímsdóttir, E. *et al.* Midbrain Dopaminergic Neuron Development is Regulated by
Two Molecularly Distinct Subtypes of Radial Glia Cells. *bioRxiv* 2024.03.08.584031
(2024) doi:10.1101/2024.03.08.584031.

24.   Desai, A. R. & McConnell, S. K. Progressive restriction in fate potential by neural
progenitors during cerebral cortical development. *Development* **127**, 2863–2872 (2000).

25.   Nürnberg, E., Horschitz, S., Schloss, P. & Meyer-Lindenberg, A. Basal glucocorticoid
receptor activation induces proliferation and inhibits neuronal differentiation of human
induced pluripotent stem cell-derived neuronal precursor cells. *J Steroid Biochem Mol*
Biol **182**, 119–126 (2018).

26.   Toledo, E. M. *et al.* Srebf1 Controls Midbrain Dopaminergic Neurogenesis. *Cell Rep* **31**,
107601 (2020).

27.   Kim, W. Y. NeuroD1 is an upstream regulator of NSCL1. *Biochem Biophys Res Commun*
419, 27–31 (2012).

28.   Murdoch, J. N., Eddleston, J., Leblond-Bourget, N., Stanier, P. & Copp, A. J. Sequence
and expression analysis of Nhlh1: a basic helix-loop-helix gene implicated in
neurogenesis. *Dev Genet* **24**, 165–177 (1999).

29.   Dennis, D. J., Han, S. & Schuurmans, C. bHLH transcription factors in neural
development, disease, and reprogramming. *Brain Res* **1705**, 48–65 (2019).

30.   Tolve, M. *et al.* The transcription factor BCL11A defines distinct subsets of midbrain
dopaminergic neurons. *Cell Rep* **36**, 109697 (2021).

31.   Simon, H. H., Saueressig, H., Wurst, W., Goulding, M. D. & O'Leary, D. D. M. Fate of
Midbrain Dopaminergic Neurons Controlled by the Engrailed Genes. *Journal of*
Neuroscience **21**, 3126–3134 (2001).

32.   Albéri, L., Sgadò, P. & Simon, H. H. Engrailed genes are cell-autonomously required to
prevent apoptosis in mesencephalic dopaminergic neurons. *Development* **131**, 3229–3236
(2004).

33.   Poulin, J. F. *et al.* Mapping projections of molecularly defined dopamine neuron subtypes
using intersectional genetic approaches. *Nature Neuroscience 2018 21:9* **21**, 1260–1271
(2018).

34.   Agoston, Z., Li, N., Haslinger, A., Wizenmann, A. & Schulte, D. Genetic and physical
interaction of Meis2, Pax3 and Pax7 during dorsal midbrain development. *BMC Dev Biol*
12, 10 (2012).

35.   Nakamura, H., Katahira, T., Matsunaga, E. & Sato, T. Isthmus organizer for midbrain and
hindbrain development. *Brain Res Rev* **49**, 120–126 (2005).

36.   Di Bella, D. J. *et al.* Molecular logic of cellular diversification in the mouse cerebral
cortex. *Nature 2021 595:7868* **595**, 554–559 (2021).

37. Gheldof, A., Hulpiau, P., van Roy, F., De Craene, B. & Berx, G. Evolutionary functional
analysis and molecular regulation of the ZEB transcription factors. *Cell Mol Life Sci* **69**,
2527–41 (2012).

38. Smidt, M. P. *et al.* Early developmental failure of substantia nigra dopamine neurons in
mice lacking the homeodomain gene Pitx3. *Development* **131**, 1145–1155 (2004).

39. Ziegler, A. B. *et al.* Cell-Autonomous Control of Neuronal Dendrite Expansion via the
Fatty Acid Synthesis Regulator SREBP. *Cell Rep* **21**, 3346–3353 (2017).

40. Yaghmaeian Salmani, B. *et al.* Transcriptomic atlas of midbrain dopamine neurons
uncovers differential vulnerability in a Parkinsonism lesion model. *Elife* **12**, (2024).

41. Storm, P. *et al.* Lineage tracing of stem cell-derived dopamine grafts in a Parkinson's
model reveals shared origin of all graft-derived cells. *Science Advances* **10**, 3057 (2024).

42. Tang, L. *et al.* A primate nigrostriatal atlas of neuronal vulnerability and resilience in a
model of Parkinson's disease. *Nature Communications 2023 14:1* **14**, 1–13 (2023).

43. Chung, S. *et al.* The homeodomain transcription factor Pitx3 facilitates differentiation of
mouse embryonic stem cells into AHD2-expressing dopaminergic neurons. *Molecular and*
Cellular Neuroscience **28**, 241–252 (2005).

44. Mesman, S., Wever, I. & Smidt, M. P. Tcf4 Is Involved in Subset Specification of
Mesodiencephalic Dopaminergic Neurons. *Biomedicines 2021, Vol. 9, Page 317* **9**, 317
(2021).

45. Koch, P., Opitz, T., Steinbeck, J. A., Ladewig, J. & Brüstle, O. A rosette-type, self-
renewing human ES cell-derived neural stem cell with potential for in vitro instruction and
synaptic integration. *Proc Natl Acad Sci U S A* **106**, 3225–30 (2009).

46. Pissadaki, E. K. & Bolam, J. P. The energy cost of action potential propagation in
dopamine neurons: clues to susceptibility in Parkinson's disease. *Front Comput Neurosci*
7, (2013).

47. del Rey, N. L. G. *et al.* Calbindin and Girk2/Aldh1a1 define resilient vs vulnerable
dopaminergic neurons in a primate Parkinson's disease model. *npj Parkinson's Disease*
2024 10:1 **10**, 1–15 (2024).

48. Nie, S. *et al.* Sox6 and ALDH1A1 Truncation by Asparagine Endopeptidase Defines
Selective Neuronal Vulnerability in Parkinson's Disease. *Advanced Science* **12**, 2409477
(2025).

49. Pantazis, C. B. *et al.* A reference human induced pluripotent stem cell line for large-scale
collaborative studies. *Cell Stem Cell* **29**, 1685-1702.e22 (2022).

50. Zheng, G. X. Y. *et al.* Massively parallel digital transcriptional profiling of single cells.
Nature Communications 2017 8:1 **8**, 1–12 (2017).

51.   Satpathy, A. T. *et al.* Massively parallel single-cell chromatin landscapes of human
immune cell development and intratumoral T cell exhaustion. *Nature Biotechnology 2019*
37:8 **37**, 925–936 (2019).

52.   Korsunsky, I. *et al.* Fast, sensitive, and accurate integration of single cell data with
Harmony. *Nat Methods* **16**, 1289 (2019).

53.   Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902.e21
(2019).

54.   Stuart, T., Srivastava, A., Madad, S., Lareau, C. A. & Satija, R. Single-cell chromatin state
analysis with Signac. *Nature Methods 2021 18:11* **18**, 1333–1341 (2021).

55.   Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**, R137
(2008).

56.   McGinnis, C. S., Murrow, L. M. & Gartner, Z. J. DoubletFinder: Doublet Detection in
Single-Cell RNA Sequencing Data Using Artificial Nearest Neighbors. *Cell Syst* **8**, 329-
337.e4 (2019).

57.   Thibodeau, A. *et al.* AMULET: a novel read count-based method for effective multiplet
detection from single nucleus ATAC-seq data. *Genome Biol* **22**, 252 (2021).

58.   Clarke, Z. A. *et al.* Tutorial: guidelines for annotating single-cell transcriptomic maps
using automated and manual methods. *Nature Protocols 2021 16:6* **16**, 2749–2764 (2021).

59.   Zhou, Y. *et al.* Metascape provides a biologist-oriented resource for the analysis of
systems-level datasets. *Nat Commun* **10**, 1523 (2019).

60.   Xu, C. *et al.* Automatic cell-type harmonization and integration across Human Cell Atlas
datasets. *Cell* **186**, 5876-5891.e20 (2023).

61.   Domínguez Conde, C. *et al.* Cross-tissue immune cell analysis reveals tissue-specific
features in humans. *Science* **376**, eabl5197 (2022).

62.   Trapnell, C. *et al.* The dynamics and regulators of cell fate decisions are revealed by
pseudotemporal ordering of single cells. *Nature Biotechnology 2014 32:4* **32**, 381–386
(2014).

63.   Bergen, V., Lange, M., Peidli, S., Wolf, F. A. & Theis, F. J. Generalizing RNA velocity to
transient cell states through dynamical modeling. *Nature Biotechnology 2020 38:12* **38**,
1408–1414 (2020).

64.   Gayoso, A. *et al.* Deep generative modeling of transcriptional dynamics for RNA velocity
analysis in single cells. *Nature Methods 2023 21:1* **21**, 50–59 (2023).

65.   Cao, J. *et al.* The single-cell transcriptional landscape of mammalian organogenesis.
Nature 2019 566:7745 **566**, 496–502 (2019).

66. Crow, M., Paul, A., Ballouz, S., Huang, Z. J. & Gillis, J. Characterizing the replicability
of cell types defined by single cell RNA-sequencing data using MetaNeighbor. *Nature*
Communications 2018 9:1 **9**, 1–12 (2018).

67. Aibar, S. *et al.* SCENIC: Single-cell regulatory network inference and clustering. *Nat*
Methods **14**, 1083–1086 (2017).

68. Faul, F., Erdfelder, E., Lang, A. G. & Buchner, A. G*Power 3: a flexible statistical power
analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods*
39, 175–191 (2007).

69. Björklund, A. & Dunnett, S. B. The Amphetamine Induced Rotation Test: A Re-
Assessment of Its Use as a Tool to Monitor Motor Impairment and Functional Recovery in
Rodent Models of Parkinson's Disease. *J Parkinsons Dis* **9**, 17–29 (2019).

Acknowledgements

This work was mainly supported by the European Union's Horizon 2020 (H2020) Research and

Innovation Programme (Neurostemcell-Reconstruct 874758/EA) and the European Research

Council (ERC) PreciseCellPD grant (Grant No. 884608/EA). Other contributors: Knut and Alice

Wallenberg Foundation Wallenberg Scholar 2018.0232/EA, Vetenskaprådet VR2020-01426/EA,

Chan Zuckerberg Initiative and the Silicon Valley Community Foundation (2018-191929)/EA,

H2020 MSCA-ITN ASCTN/EA, H2020 Grant No. 899687 - HS-SEQ, Aligning Science Across

Parkinson's (ASAP-20505)/EA, Hjärnfonden project FO2024-0195/EA. JCK and PU were

supported by the German Science Foundation (516641042) and the Wenner-Gren foundation

(UPD2022-0159) respectively.

The authors acknowledge support from the National Genomics Infrastructure in Stockholm funded

by Science for Life Laboratory, the Knut and Alice Wallenberg Foundation and the Swedish

Research Council, and NAISS/Uppsala Multidisciplinary Center for Advanced Computational

Science for assistance with massively parallel sequencing and access to the UPPMAX

computational infrastructure. We are grateful for the assistance of the Molecular Neurobiology

unit within the Department of Medical Biochemistry and Biophysics at the Karolinska Institute.

We would like to posthumously thank Prof. Ernest Arenas, without whom this project would not

have been possible.

Ethics declarations

Competing interests

940 None of the authors declare competing interests.

Contributions

R.G.S., G.L. and E.A. conceptualized the experiments. R.G.S. and G.L. redacted the manuscript.

R.G.S. performed *in vivo* experiments; G.L., R.K., C.S., C.T. and A.X. contributed to the *in vitro*

protocol development. J.C.K. contributed to the calcium imaging experiment. G.L. and A.X.

contributed to the *in vitro* transcriptional analysis; R.G.S. and C.A. performed the graft nuclei

isolation, and R.G.S. and G.L. the transcriptional analysis. C.S., G.C.B. and O.D. contributed to

supervision and reviewed the manuscript.