1   **Development of a protocol utilizing single-cell analysis for the differentiation of human**
2   **iPSCs into SOX6+ midbrain dopaminergic neurons.**

3   Guochang Lyu[1]*, Roberto Garcia-Swinburn[1]*#, Judith C. Kreutzmann[1,2] , Anqi Xiong[1], Rika
4   Kojima[2], Carmen Abaurre[1], Chiara Tremolanti[1], Sandra Gellhaar[1], Per Uhlén[1], Per
5   Svenningsonn[2], Gonçalo Castelo-Branco[1], Onur Dagliyan[1]#, Carmen Salto[1], Ernest Arenas[1]†

6   [1]Department of Medical Biochemistry and Biophysics, Karolinska Institutet, 171 77 Stockholm,
7   Sweden

8   [2]Department of Clinical Neuroscience, Karolinska Institutet, 171 76 Stockholm, Sweden

9   *These authors contributed equally

10 # Correspondence to roberto.garcia.swinburn@ki.se or onur.dagliyan@ki.se

11 † Deceased 15/09/2024

12

13 **ABSTRACT**

14 Parkinson's disease (PD) is a neurodegenerative disorder characterized by the loss of midbrain
15 dopaminergic (mDA) neurons in the substantia nigra, leading to motor symptoms. Current
16 pharmacological treatments are insufficient in halting disease progression, making cell therapy
17 a promising alternative for replacing lost neurons. Advances in stem cell research have enabled
18 the generation of mDA progenitors from human pluripotent stem cells (hPSCs), demonstrating
19 success in preclinical models and clinical trials. However, challenges such as dyskinesias from
20 non-mDA contaminations, graft rejection, and the need for more specific cell types persist. This
21 study focuses on utilizing single-cell multiomic and transcriptomic data to design mDA
22 progenitor differentiation protocols, aiming to generate more specific subtypes of mDA
23 neurons, particularly those vulnerable to degeneration in PD. By refining the protocol with
24 enhanced ventralization and prolonged Wnt activation, we achieved improved midbrain
25 patterning, cell viability, and the accelerated generation of SOX6-expressing mDA neurons *in*
26 *vitro* with proven therapeutical properties in grafted hemiparkinsonian mice models. This
27 approach aims to present a protocol designed using single-cell analysis to enrich the most PD-
28 affected mDA neuron populations, offering potential for better cell modelization, and more
29 targeted cell replacement therapies.

30

31 **INTRODUCTION**

32 Parkinson's disease (PD), the second most prevalent neurodegenerative disorder, presents
33 significant challenges in treatment and patient quality of life. The progressive degeneration of
34 midbrain dopaminergic (mDA) neurons in the substantia nigra pars compacta (SNpc), which
35 primarily project to the dorsal striatum, leads to the loss of dopaminergic input to the striatum
36 and results in the characteristic motor symptoms of PD (Poewe et al., 2017). Although advances
37 in pharmacological therapies have been made, these treatments are often insufficient in halting
38 disease progression or restoring lost neuronal function in later stages (Brichta et al., 2013). Cell

therapy offers a promising alternative, with the potential to replace the damaged or lost DA
neurons critical to Parkinson's pathology (Björklund & Parmar, 2021).

Advances in stem cell research and a deeper understanding of midbrain development have
enabled the production of mDA progenitors from human pluripotent stem cells (hPSCs) for
transplantation (Doi et al., 2014; Kim et al., 2021; Kriks et al., 2011; Nishimura et al., 2022).
Success in preclinical models has led to the development of clinical-grade mDA progenitors for
PD clinical trials (Kirkeby et al., 2023; Piao et al., 2021; Schweitzer et al., 2020; Takahashi, 2020).
Furthermore, phase I/II clinical trials have recently been completed using hPSCs demonstrating
safe outcomes for these therapies, paving the way for future medical use of hPSCs (Tabar et al.,
2025; Sawamoto et al., 2025).

As a result, there has been a growing focus on assessing the quality and composition of cells
derived from hPSCs. Research is underway to better define and molecularly characterize
human mDA neuron development at the single-cell level (Braun et al., 2023; Manno et al., 2016).
These studies have provided detailed analyses of the cellular composition and molecular
features of individual cell types in hPSC-derived midbrain preparations (Manno et al., 2016;
Nishimura et al., 2022; Tiklová et al., 2020; P. Xu et al., 2022).

However, challenges remain in the effectiveness of cell-based therapies, including
complications like dyskinesias reported in fetal tissue transplants. These have been reported
years after transplantation and are believed to result from serotonergic neuron contamination
within the graft (Greene et al., 2021). Another major challenge is graft rejection by the host,
typically addressed through pharmacological immunosuppression. The use of human-induced
pluripotent stem cells (hiPSCs) presents a potential solution by enabling the generation of
patient-specific neurons, thereby overcoming immune rejection (Schweitzer et al., 2020).

Single-cell transcriptomics has revealed significant diversity within mDA neurons in the adult
human SNpc, categorizing them into three major classes based on the expression of SOX6,
CALB1, and GAD2 (Kamath et al., 2022; Siletti et al., 2023). These classes are also detectable
during human mDA neuron development (Braun et al., 2023). Additionally, advances in single-
cell technologies have now made it possible to assess developmental epigenetics by the use of
ATAC-seq in developing human midbrain (Mannens et al., 2024). To better understand how to
generate specific mDA neuron subtypes, we analyzed multiomic data from established
protocols to identify approaches that enrich for the SOX6 class, as a SOX6$^+$AGTR1$^+$
subpopulation has been linked to increased vulnerability to neurodegeneration in PD in humans
and parkinsonian models (Kamath et al., 2022; Salmani et al., 2024).

In summary, we initially used a simplified mDA neuron differentiation protocol based on key
developmental components identified through single cell multiomics. This protocol was
reproducible across distinct iPSC lines. However, insights from molecular phenotyping
indicated that the resulting phenotypes were not maintained at later stages, suggesting
insufficient patterning, leading us to modify the protocol. A new protocol involving enhanced
ventralization and prolonged Wnt activation improved midbrain patterning, increased cell
viability, and generated more mature mDA neurons with higher expression levels of *OTX2*, *EN1*
and *SOX6* compared to the original protocol. Furthermore, we tested *in vivo* differentiation of

our cells in hemiPD rodent models, achieving a high number of SOX6$^+$ mDA neurons with
effective motor recovery.

While various differentiation protocols exist, many lack specificity to target the DA neuron
subtypes most vulnerable to degeneration in PD. Our work aims to address this gap by
identifying key developmental factors using multiomic data and evaluating a novel protocol to
enrich *in vitro* the mDA neuronal population most affected by Parkinson's disease and further
validate as proof-of-concept in *in vivo* PD models.

MATERIAL AND METHODS

Cell lines

KOLF 2.1J cell line was obtained through collaboration with the Aligning Science Across
Parkinson's (ASAP) consortium. Genotypes, pluripotency, and mycoplasma infection were
routinely assessed before starting new experiments. Applied iPSC lines were adapted to grow
on Laminin 521 (Biolamina, LN521)-coated plates and maintained in Essential 8 Flex media
(Thermo Fisher Scientific) at 37ºC, 5% CO_2, until starting differentiation. The medium was
changed daily, and cells were passaged at 70% confluency using TrypLE™ Select Enzyme
(Thermo Fisher Scientific) at a 1:20 ratio. When thawing and replating, 10 μM Rho-associated
protein kinase (ROCK) inhibitor (Y-27632) (Tocris) was added into the culture media overnight.

Midbrain dopaminergic neuron differentiation

Prior to the mDA differentiation, adapted iPSC cells were seeded at a density of 200,000
cells/cm^2 onto vessels coated with Geltrex (Life Technologies) and Laminin-511 (Biolamina,
LN511), and maintained in Essential 8 Flex media (Thermo Fisher Scientific) for expansion. The
differentiation protocol should be initiated when the cells reach approximately 80% confluency.
Cultured cells will be kept in Neurobasal/N2/B27 (Life Technologies) medium and 2 mM L -
glutamine (Invitrogen), along with various supplements for the first stage of the protocol. From
day 0 to day 3, the culture medium was conditioned with 250 nM LDN193189 (Stemgent), 10 μM
SB431542, 2 μM Purmorphamine, and 0.7 μM CHIR99021 from day 0 to day 3, with the inclusion
of 10 μM Y27632 (Tocris Bioscience) during the first 24 hours. Boosted CHIR9901 (7.5 μM) was
administered from day 4 to day 10, while LDN193189, SB431542 and Purmorphamine were
withdrawn after day 7. From day 10 to day 28, cells were switched to the differentiation media:
Neurobasal/B27/L-Glu media supplemented with 20 ng/mL brain-derived neurotrophic factor
(BDNF, R&D Systems), 20 ng/ml glial cell line-derived 455 neurotrophic factor (GDNF, R&D
Systems), 1 ng/ml transforming growth factor type b3 (TGFb3, R&D Systems), 0.2 mM ascorbic
acid (Sigma), 0.2 mM Dibutyryl cAMP (Sigma). On day 11 and 16, the resulting cells were
dissociated into single cells, replated at a density of 800,000 cells/cm^2 onto plates coated with
LN511 and polyornithine/LN511 respectively, and treated with 10 μM Y27632 for 48 hours. To
promote neurogenesis, the mDA differentiation media was supplemented with 10 mM DAPT
(Sigma) from day 12-28. The small molecule GW3965 (10 μM) was administered from day 12 to
day 15, followed by a cocktail of small molecules consisting of 1 μM PD0325901 and 5 μM
SU5402 (Sigma), from day 16 to day 21.

Immunocytochemistry

Progenitor cells at different stages were seeded in a 96-well plate at a density of about 300,000
cells per well for the intended cell culture purposes. Harvested cells were fixed by 4%
paraformaldehyde (PFA) for 20 min at 4ºC, and then pre-incubated by 5% donkey serum in
phosphate buffered saline containing 0.3% Triton X-100 (PBST) for 1 hr. The samples were
incubated with primary antibodies at 4ºC overnight. The samples were incubated with either
Alexa488 or Alexa555 or Alexa647-conjugated secondary antibodies (Thermo Fisher Scientific)
for 30 min and then were incubated with 4',6-diamidino-2-phenylindole (DAPI) for 15 min. The
primary antibodies were used as follows: CORIN (rat, 1:1,000, R&D Systems, MAB2209), DCX
(goat, 1:500, SantaCruz, sc-8066), EN1 (mouse 1:50, DSHB, 4G11), FOXA2 (goat, 1:500, R&D
Systems, AF2400), GIRK2 (rabbit, 1:400, Alomone, APC006), LMX1A (rabbit, 1:4,000, Millipore,
AB10533), LMO3 (goat, 1:200, SantaCruz, sc-82647), MAP2 (mouse 1:1,000, Sigma, M4403),
NURR1 (rabbit, 1:500, SantaCruz, sc-990), OTX2 (goat, 1:1,000, R&D Systems, AF1979), TH
(rabbit, 1:1,000, Millipore, AB152), TH (mouse, 1:500, ImmunoStar, 22941), and TH (sheep,
1:500, Novus, NB300). Mounted cells were further visualized using Zeiss LSM880 or 980.
Quantification of images was performed using QuPath (Bankhead et al., 2017) by generating a
"single measurement classifier" or "composite classifier" for positive or double-positive cells
respectively.

Calcium imaging

The 35mm culture dishes were loaded with the Ca2+-sensitive fluorescence indicator Fluo-
4/AM (5 µM; Invitrogen, USA) and Pluronic F-127 (0.625%, ThermoFisher Scientific) and
incubated for 30 min at 37ºC in Krebs-Ringer buffer, containing: NaCl (119 mM), KCl (2.5 mM),
NaH2PO4 (1 mM), CaCl2 (2.5 mM), MgCl2 (1.3 mM), HEPES (20 mM) and D-Glucose (11 mM),
with pH adjusted to 7.4. Dishes were washed twice and Ca2+-measurements were performed in
Krebs-Ringer buffer at 37ºC on a confocal spinning disk microscope (Nikon CrEST X-Light V3)
equipped with a 20x Air lens. Excitation was assessed at 480 nm at a sampling frequency of 1
Hz. The equipment was controlled with, and data was acquired using the Nikon NIS-Elements
software. FIJI, MATLAB (R2020b, MathWorks, USA) and FluoroSNNAP
(https://github.com/tapan-patel/FluoroSNNAP) were used to process and analyze the data.

Single cell sequencing libraries preparation and data processing

A total of 5 samples (DIV 11, 16, 28, 42, and 56) from KOLF2.1J mDA neuron cultures generated
with the optimized protocol were collected for time-course multiomic analysis (ATAC + Gene
Expression); while 3 samples (DIV 14, 16, and 28) from the new protocol trial of KOLF2.1J mDA
differentiation were collected for single-nuclei transcriptomic profiling. All 8 samples were
prepared with demonstrated protocols (10x Genomics) and the resulting Fastqc files were
processed through the 10X Genomics Cellranger pipeline (Zheng et al., 2017), while 5 multiomic
libraries were aligned to the hg38 reference genome using Cellranger-arc default settings
(Satpathy et al., 2019). Resulting transcriptomic and epigenomic data from individual samples
were combined by projects using the 'aggr' function in Cellranger or Cellranger-arc, without
normalization applied. Objects produced with Seurat V4.4.0 was employed to accommodate
aggregated count matrix (Stuart et al., 2019), with the fragment files specifically integrated into
Signac v1.9.0 chromatin assays (Stuart et al., 2021). Quality metrics for gene expression and
peak accessibility, such as mitochondrial ratio, cell cycle score, nucleosome signal, and
Transcription Start Site (TSS) enrichment, were thereafter calculated to determine quality

control cutoffs. Depending on sample quality, following thresholds were applied for
downstream analysis: a maximum of 300,000 and a minimum of 1,000 ATAC counts per cell, a
maximum of 25,000 and a minimum of 600 RNA counts, at least 250 genes detected, a
maximum nucleosomal signal of 1.5, TSS enrichment of at least 2, and mitochondrial content
no higher than 15%. These cutoffs reduced the cell count by less than 20% without significantly
affecting clustering downstream. Peak calling was performed with MACS2 v2.2.9 using the
aggregated fragments file (Y. Zhang et al., 2008), and those mapped to non-standard
chromosomes or overlapping hg38 genome blacklist regions were excluded. DoubletFinder v2.0
was applied to identify doublets in gene expression data by flagging instances with a high
resemblance to artificial neighbors in gene expression space (McGinnis et al., 2019). AMULET
(ATAC-seq MULtiplet Estimation Tool) was additionally utilized to detect potential multiplets in
snATAC-seq datasets by enumerates regions with greater than two uniquely aligned reads
across the genome (Thibodeau et al., 2021). All predicted doublets were removed from the
downstream analysis. After merging high-quality cells from the relevant projects, count data
was normalized using SCTransform V2 function with regression out of mitochondrial or cell-
cycle genes. The top 2,000 most variable genes were initially identified through variance-
stabilizing transformation, along with 95% of the most frequent peaks in multiomic datasets, as
a part of the significant feature selection process. Subsequent dimensionality reduction was
performed using the principal components analysis (PCA), in which the top 30/40 principal
components were chosen from the elbow plot to represent most of the diversity of datasets. For
single-nuclei ATAC-seq datasets, singular value decomposition (SVD) was performed with the
top 25 of the Latent Semantic Indexing (LSI) layers to reduce the dimensions of the most
common peaks. Notably, the first LSI component was excluded from downstream analysis as it
often captures sequencing depth (technical variation) rather than biological variation. Putative
clusters were identified by a spiking neural network (SNN) modularity optimization-based
clustering algorithm, and the optimal resolution of clustering was determined by iterating over
different resolution for the most stable segregation. Finally, the calculated nearest-neighbor
graph was projected to two dimensions to visualize the distribution of instances by representing
with uniform manifold approximation and projection (UMAP). To accurately identify and
interpret cell states in vitro, we leveraged combined manual cell type annotation using cluster-
specific marker genes (Clarke et al., 2021; Zhou et al., 2019), with automatic annotation
methods using reference datasets that characterize first-trimester developing human brain
(Braun et al., 2023; C. Xu et al., 2023). Initially, a label transfer was performed with Celltypist
(Conde et al., 2022) and 'TransferData' function in Seurat to automatically classify selected
dataset with cell type labels from the reference dataset, based on the similarity of query gene
expression profiles to selected anchors in the reference dataset. The identity of each cell will be
determined by the highest prediction score, and for stringent comparison, cells with a
prediction score below 0.51 will be excluded from downstream analysis. After data trimming
and reclustering, a manual cluster-based annotation was performed, using both expression
levels of canonical midbrain development marker genes and top 50 upregulated genes
identified by the 'FindAllMarkers' function. These upregulated genes were used for pathway and
biological process enrichment analysis via Metascape (Zhou et al., 2019), and the top-ranked
terms were used to refine annotations and determine cell subtypes or states. To reconstruct the
putative trajectory of mDA differentiation in vitro, we employed Monocle3 (Trapnell et al., 2014)
and scVelo (Bergen et al., 2020; Gayoso et al., 2024) to infer pseudo- and latent-temporal

ordering of cells using default parameters. The root of trajectory for pseudotime analysis was
set as ProgFP or NbM at the earliest timepoint.

Animals

Healthy female adult (8-12 weeks old) C57BL/6N-Rag2Tm1-IL2rgTm1-SirpaNOD/Rj (B6RGS)
mice were purchased from Janvier Labs. Groups of 5 animals were housed in ventilated cages
under a 12 h light-dark cycle with food (standard chow) and water ad libitum at 20-22 ºC and 50-
55% humidity. All surgical procedures followed aseptic techniques, and for post-surgical
recovery, additional Dietgel Recovery (Scanbur) and food pellets were supplied on the cage
floor. All procedures were approved by the local ethical committee (Stockholms
djurforsoksetiska namnd) and followed the EU directive 2010/63/EU under the ethical permit of
Ernesto Arenas Cases and Sandra Gellhaar (12265-2021; 2-3143/23; 2-3595/24; 2-3147/24).

6-OHDA mouse model

B6RGS mice were acclimatized 2-3 weeks before surgery to the animal facility and the
researcher and to a recovery diet 2-3 days before surgery. For 6-hydroxidopamine (6-OHDA)
lesions (or sham), animals were injected analgesia (Carprofen 5mg/kg + Buprenorphine 0.1
mg/kg) 15 minutes before isoflurane induction (4%). Animals were anesthetized at 1-2%
isoflurane in air and injected intracranially (ICR) using a Neurostar Robotic Stereotaxic Frame
equipped with a 10 µl syringe with 33G blunt needle (Hamilton) 1 µl of 6-OHDA 3.6 µg/ µl in
sterile ascorbic acid/saline 0.02% (or sham vehicle) in the right medial forebrain bundle [respect
to Bregma: AP -1.2 mm, ML +1.2 mm, DV -5.0 mm] at a flow of 0.2 µl/min, raising the needle 500
µm for 3 minutes. The needle was left inside for 2 additional minutes. Animals were placed in
warm cages (28 ºC) between 3 days and 1 week for better recovery and rehydrated with Ringer
Acetate solution if weight decreased by 5%. 6-OHDA hemiparkinsonism was confirmed by
amphetamine-induced rotation. Briefly, 4 weeks after the lesion, mice were injected
intraperitoneally with d-amphetamine in saline (5 mg/kg). 10 minutes after injection, the mice
were recorded for 30 minutes inside a 15 cm wide cylinder in dark conditions, and the number of
net ipsilateral rotations (ipsilateral-contralateral) per minute in active movement was
calculated using Ethovision XT 17. Mice are considered hemiparkinsonian if the values are >6
rotations/minutes.

hiPSC transplant

HemiPD-confirmed mice of healthy weight (22-30 g) were randomly selected for ICR hPSC
injection in the right striatum [respect to Bregma: AP +0.5 mm, ML +2 mm, DV -3.5 mm].
Analgesia and anesthesia were followed as above. For ICR transplants, 2 µl of 100.000 ± 20.000
hPSCs/µl in Neurobasal medium was injected to achieve 200.000 hPSC transplants. For cell
injection, the Neurostar robot was coupled with a 10 µl syringe with a 27G sharp syringe. A flow
of 0.5 µl/min was used, and the needle remained for 5 additional minutes. Cells were injected
between 1-6 hours after thawing.

Amphetamines-induced rotations

247   Mice were analysed once a month until endpoint was reached. Mice were injected
248   intraperitoneally with d-amphetamine in saline (5 mg/kg). 10 minutes after injection, the mice
249   were recorded for 30 minutes inside a 15 cm wide cylinder in dark conditions, and the number of

250   net ipsilateral rotations (ipsilateral-contralateral) per minute in active movement was
251   calculated using Ethovision XT 17.

Pole test

253   Animals were placed facing upwards on the top of a 20 cm vertical pole. The time to climb down
254   was counted starting when the mouse rotates facing downwards. Animals were trained for a
255   minimum of 2 days before testing day.

Cylinder wall

257   Animals were placed inside a 12 cm diameter glass cylinder in red light and quiet conditions. For
258   15 minutes, animals were recorded and videos were analysed at x8 speed. Unilateral contacts
259   were counted.

Tape removal test

261   A green adhesive tape 2x3 mm was placed on each paw individually. The time it took to remove
262   the tape was counted, beginning when the mouse recognized it and ending when it was
263   removed from the paw. Net contralateral time (ipsilateral-contralateral) was calculated to
264   better visualize and eliminate external factors. Before testing day, animals were trained for a
265   minimum of 2 days.

Immunohistochemistry

Animals were terminally anesthethised with pentobarbital and then perfused intracardially with
50 ml of pH-corrected PBS1X followed by 50 ml of formaldehyde 4 % (FA) both at a flow of 9
ml/min. Brains were extracted and placed in FA overnight, after which they were placed in
sucrose 30%/PBS solution at 4ºC for 2-3 days. After, brains were kept at -80ºC.

Brains were equilibrated at -20ºC for 4 hours before slicing. Samples were sliced in series at -
20ºC at 20 µm using a Cryostar NX70 (Epredia) and stored for further use at -20ºC in
cryoprotective solution (Ethylenglycol 30%,Glycerol 30%,PBS1X). A total of 5 brains were
processed.

Samples were stained using free-floating protocols. Briefly, 2 series of 5 minutes of PBS1X,
followed by 15 minutes of PBS-TritonX 0.3% (PBTx0.3%), then 2 series of 5 minutes of PBTx0.1%.
Samples were blocked for 1 hour with blocking solution (BSA 1 mg/ml, Normal Donkey Serum
5%, PBTx0.1%) and incubated o/n at 4ºC with primary antibody diluted in blocking solution.
Primary antibody was washed with 2 series of 5 minutes PBTx0.1% and samples were incubated
for 4 hours with secondary antibody diluted in blocking solution and washed twice for 5 minutes
with PBS1X. Tissue was counterstained with 4',6-diamidino-2-phenylindole (DAPI, Sigma,
D9542) 1:1000 in PBS1X for 10 minutes and then washed twice for 5 minutes with PBS1X.
Stained samples were mounted on Superfrost Plus slides (Epredia) and Fluoromount-G
mounting medium (00-4958-02, Thermo Scientific). Primary antibodies used were STEM121
(mouse, 1:1000, Tanaka Bio, Y40410), TH (rabbit, 1:1000, Pelfreeze, P4010; sheep, 1:250
Pelfreeze, P60101), HuNu (mouse, 1:1000, Millipore, MAB1281), SOX6 (rabbit, 1:200, Abcam,
ab30455), ALDH1A (rabbit, 1:500, Atlas Antibodies, HPA002123), AGTR1 (rabbit, 1:100,
Invitrogen, MA5-56587), Calbindin-1 (rabbit, 1:200, Invitrogen, PA5-85669), LMX1 (rabbit, 1:100,
Millipore, AB10533), FOXA2 (rabbit, 1:200, Cell Signaling, 3143S), NR4A2 (rabbit, 1:500,

Invitrogen, PA5-78097), PITX3 (rabbit, 1:100, Invitrogen, 35-2850), DAT (rat, 1:250, Abcam,
ab5990), GIRK2 (rabbit, 1:500, Alomone Labs, APC006), VMAT2 (rabbit, 1:500, Sigma-Aldrich,
AB1598P), DDC (rabbit, 1:500, Novus Biological, NB300-252), Ki67 (rabbit, 1:200, Abcam,
ab15880), SOX2 (goat, 1:200, R&D Systems, AF2018). All secondary antibodies were from
Invitrogen, donkey as host and conjugated with Alexa Fluor 488, 555 or 647. Images were
captured either with laser confocal microscope Zeiss LSM800 or high-throughput
epifluorescence microscope Zeiss Axioscan. Quantification of images was performed using
QuPath (Bankhead et al., 2017) by generating a 'single measurement classifier' or 'composite
classifier' for positive or double-positive cells respectively.

Single-nuclei RNA sequencing of grafts and data analysis

Animals were euthanized by cervical dislocation and brains were promptly extracted and flash-
frozen in dry-ice chilled isopentane for 10-15 seconds. Flash-frozen brains were stored at -80 °C
until graft extraction. A total of 3 brains with grafts were processed.

On the day of nuclei extraction, brains were equilibrated to -20ºC and sliced in -20ºC cooled
chambers. Brain matrices were used to stabilize the brain, and 1 mm-thick slices for a total of 4
(2 before the graft implant location and 2 after) were cut using chilled razors. 1 mm punches of
each slice of the dorsal striatum were extracted using 1 mm biopsy-punchers (49101, PFM
Medical).

Nuclei were dissociated using the 10X Genomics Chromium nuclei extraction kit (1000494)
using manufacturer's instructions. For better extraction, eppendorfs used for tissue
dissociation were coated with BSA 1% o/n at 4ºC, briefly washed with filtered sterile PBS1X
before procedure. 10X Genomics Chromium GEM-X Single Cell 3' Kit v4 (1000691), Chromium
GEM-X Single Cell 3' Chip Kit v4 (1000690) and Dual Index Kit TT Set A (1000215) were used for
GEM generation using manufacturer's instructions with the 10X Controller iX. GEMs were sent
for RNA-sequencing to National Genomics Infrastructure of SciLifeLab (Karolinska Institute,
Sweden) using the NovaSeq XPlus 25B. Data analysis was performed as the *in vitro* RNAseq
dataset, described above. To ensure analysis of human cells, we aligned the libraries to the h38
human genome dataset and the m39 mouse genome dataset, and cells with a ratio of
human:mouse gene count >1 were processed.

Statistics

The N for the animals was calculated using G-Power 3.1 (Faul et al., 2007) based on prior data
from several publications of hemiparkinsonian rodent model behavior (Bjorklund and Dunnett,
2019).

GraphPad Prism v10.00 was used for generating all graphs and performing statistical analyses.
For pairwise comparisons, Student's t-test was utilized. One-way ANOVA or two-way ANOVA
with Tukey's or Bonferroni's multiple comparison tests were used to compare multiple
variables. Error bars indicate the standard error of the mean (SEM). Significance levels were set
as follows: ns = not significant; * = $p<0.05$; ** = $p<0.01$; *** = $p<0.001$; **** = $p<0.0001$.

RESULTS

Longitudinal multiomic single-nuclei sequencing delineates the cell-type composition throughout the process of midbrain dopaminergic neuron differentiation in vitro

Building upon our recently published method (Nishimura et al., 2022), we initially designed a new protocol focused on precise midbrain floor plate (mFP) patterning and improving cell viability to generate sufficient progenitor cells, while expanding compatibility to a broader range of human induced pluripotent stem cell lines. We were able to successfully sequentially differentiate iPSC into midbrain progenitor cells, to functional neurons replicating the canonical markers of mDA neurons (Sup. Figure 1).

Our primary goal was to investigate the developmental trajectories leading to the generation of mDA neurons *in vitro* from both transcriptomic and epigenomic perspectives. We thus conducted single cell multiomic (ATAC + Gene Expression) sequencing in a wild-type iPSC line, KOLF2.1J, at different developmental stages (Days in vitro (DIV): DIV11, 16, 28, 42 and 56; Figure 1A). Following quality control and filtering, we combined manual cell type annotation using cluster-specific marker genes (Clarke et al., 2021; Zhou et al., 2019), with automatic annotation methods (Conde et al., 2022), utilizing reference datasets representing the developing human brain (La Manno et al., 2016). We identified 9,233 high-quality cells, categorized into 8 major types, with sequential differentiation to a mature DA identity (Figure 1B,C). Chronologically, cells from early differentiation phases (DIV11 and 16) exhibited high expression densities of classic markers associated with regional brain identities, such as *LMX1A*, and *OTX2*, reflecting their roles in early neural progenitor stages and initial dopaminergic specification. At DIV28, the expression of *NR4A2* is upregulated; and at advanced phases as DIV42 and 56, markers such as *TH*, *DDC*, and *KCNJ6*, which are associated with mature mDA neuronal identities, displayed high expression densities and chromatin accessibilities, signaling the maturation of dopaminergic neurons (Figure 1F, Sup. Figure 2).

We then validated our annotations by comparing them to publicly available scRNA-seq datasets from human fetal midbrain samples (Braun et al., 2023; C. Xu et al., 2023), by quantifying cell type identity across individual cells (Crow et al., 2018). As a result, our annotated cell types closely resembled their endogenous counterparts, as demonstrated by reasonable hierarchical clustering (Figure 1D). High similarity scores (>0.8) were predominantly observed during early stages of the midbrain differentiation and were primarily enriched in precursor cell types such as ProgFP and NbM, underscoring the effectiveness of our approach in recapitulating endogenous early mDA neuron development. Radial glia cell subtypes, Rgl2 and Rgl3, also displayed transcriptomic profiles akin to their counterparts in vivo, hinting at the establishment of an mDA signaling niche to deliver cell type-specific developmental factors for neuronal maturation *in vitro* (Ásgrímsdóttir et al., 2024). At advanced stages, mDA neurons, closely matched their endogenous cells from reference datasets. These findings support the effectiveness and reliability of this protocol in generating desired cell populations, while also implying that additional or prolonged developmental cues may be needed to promote the maturation and survival of postmitotic neurons at later stages.

Reconstructed midbrain dopaminergic neuron differentiation trajectory resolves evolving transcriptional profiles throughout lineage specification and maturation

Beyond chronological time, which may not sufficiently capture variations in transcriptomic and/or epigenetic regulation and can be significantly confounded by cellular heterogeneity during differentiation, we sought to delineate the lineage trajectory from neural progenitors to functional neurons using single-cell transcriptional profiles as the roadmap. We isolated and further subcategorized dopaminergic lineage cells (ProgFP, NbM and mDA neurons) from our multimodal dataset and inferred putative dopaminergic trajectories through both transcriptome pseudotime (Cao et al., 2019) and RNA velocity (Bergen et al., 2020). Through calculating the RNA velocity stream and predicted latent time from selected cell types, we discovered that ProgFP and NbM were on a neuronal differentiation path toward mature mDA neurons, aligned with their chronological time (Figure 1G). UMAP-based embeddings of selected cell types revealed that mDA neurons can be arranged in a maturation hierarchy, progressing from the early *Neuron projection* to acquiring the ability for *Neurotransmitter release*, followed by structural maturation through *Synaptic assembly*, and ultimately culminating in functional refinement via *Synaptic modulation*, which aligns with the positioning from the independent pseudotime analysis. Interestingly, a shortcut bypassing the cluster for axonal development on DIV 28 and directly leading to mature mDA neurons, was observed at the advanced differentiation time points (DIV 42 and 56), suggesting that the laminar fate, as an alternative mechanism of neurogenesis and maturation, could be involved in different stages of midbrain cell differentiation (Desai & McConnell, 2000).

Furthermore, by employing single-cell regulatory network inference and clustering (SCENIC) methods to analyze active modules of co-regulated genes, we aimed to reinforce our reconstructed developmental trajectories with an additional layer of regulon activity, thereby capturing critical shifts in gene regulatory networks that coincide with transitions in cell identities and functional states (Aibar et al., 2017). Clustering of AUCell scores for activities of each regulon revealed distinct segregation between early neural progenitors, ProgFP, and NbM, and their descendant mDA neurons along the development trajectory (Figure 1H). Regulon similarities were obviously higher among various mDA neuron subcategories, suggesting that mDA lineage specification and maturation are governed by gene networks regulated by a core set of transcription factors, which may undergo slight modifications at different stages of differentiation. Particularly, we observed increased activity of several recognized transcription factors that drive neurogenesis along the mDA lineage. For instance, upregulated NR3C1 regulon in ProgFP cells skews neural precursors towards proliferation over differentiation, thereby facilitating the progenitor pool expansion (Nürnberg et al., 2018). SREBF1, a pro-neural transcription factor controlling mDA neurogenesis (Toledo et al., 2020), was identified as one of the most active regulons in ProgFP development, driving its transition to NProg. Similarly, NEUROD1 and NHLH1, neurogenic transcription factors essential for mDA neuron development, were subsequently ranked as the top regulons in NbM, triggering neurogenesis and differentiation in postmitotic cells throughout neural development (W.-Y. Kim, 2012; Murdoch et al., 1999). Subtypes of mDA neurons appeared to share a core set of transcription factors, including PBX1, a classic regulator of mDA specification and survival (Dennis et al., 2019). In contrast, transcription factors as POU2F2 and BCL11A displayed more specialized roles at distinct maturation stages (Tiklová et al., 2019; Tolve et al., 2021). By reconstructing the *in vitro* developmental trajectory, we confirmed that the differentiation protocol successfully replicated mDA neuron embryological development, reflecting similar gene expression

dynamics and co-expression regulatory networks. This approach holds potential for the *de novo*
identification of developmental regulatory modules, offering deeper insights into dopaminergic
neuron differentiation and maturation.

An iterative enhancement in ventralization yields improved floorplate patterning and
accelerated neuronal conversion in a new protocol

Given that robust regional patterning forms the cornerstone to generate mDA neurons of high
quality, and our single-cell datasets suggested that the current cultures could benefit from
further ventralization, we aimed to investigate whether enhanced ventralization could improve
patterning outcomes, using EN1 as a marker for mDA neuron specification (Simon et al., 2001),
and to streamline the protocol to increase scalability and reduce labor intensity.

EN1 plays a pivotal role from the earliest stages of regional specification and maintenance of
mDA neurons in adulthood (Rekaik et al., 2015). It supports midbrain-hindbrain boundary
formation during midbrain development and regulates a critical set of genes (including *TH*,
VMAT2, *DAT*, and *ALDH1A1*) in postmitotic neurons to sustain mDA function (Albéri et al., 2004;
Simon et al., 2001). Thus, EN1 expression serves as a reliable surrogate for enhanced mDA
neuron specification and a promising indicator of successful transplantation in vivo (Piao et al.,
2021). To replicate an EN1 overexpression, we applied a higher concentration of the SHH
agonist purmorphamine (Kriks et al., 2011), and optimized culture conditions to minimize the
cytotoxicity, thereby preserving the desired cell viabilities (Figure 2A). Subsequently, increasing
concentrations of purmorphamine dramatically boosted EN1 expression, while traditional
midbrain floor plate markers, FOXA2 and LMX1A, remained unaffected compared to the original
protocol at DIV11 (Figure 2B,E; Sup. Figure 3A,B). By DIV16, during the fate transition from NbM
towards early mDA neurons, cell cultures from the new protocol showed a rapid generation of
TH$^+$ neurons, and higher proportion of TH+ cells that co-expressed EN1 and SOX6 (Figure 2C,F),
another key transcription factor that is typical of neurons derived from the human SNpc region
(Oosterveen et al., 2021). Moreover, in prolonged cell cultures, mDA neurons with enhanced
ventralization retained substantial co-expression of EN1 and SOX6 by DIV28 (Figure 2D,G) and
exhibited active electrophysiological properties by DIV42 (Figure 2H).

We then applied single nuclei sequencing (snRNA-seq) to DIV 14, 16, and 28 cell preparations to
assess the quality of iPSC-derived midbrain cells under the new protocol. After filtering, we
obtained high-quality 23,534 cells (Figure 3A) and then we integrated transcriptomic profiles
from both conditions with Harmony approach (Korsunsky et al., 2019), enabling a comparative
analysis by isolating the mDA neuronal lineage from different datasets (Sup. Figure 4B). Under
the new protocol, we observed an accelerated differentiation process, with a higher proportion
of NProg and NbM emerging at earlier stages, accompanied by a reduction in ProgFP cells
(Figure 3B). Using the same annotation methods as in the previous dataset, the resulting mDA
lineage cell types were enriched with classic mDA neuron marker genes (Figure 3C), including
FOXA2, *LMX1A, OTX2*, *TH*, *and NR4A2*, and showed an expected developmental trajectory
across different cell populations (Figure 3D). In contrast, the new protocol was marked by
higher EN1 levels and the sustained expression of OTX2 and SOX6 in postmitotic cells. A
transverse comparison of mDA neurons from the new protocol also revealed higher levels of TH,
SLC18A2 (VMAT2), and CALB1 (Calbindin 1), suggesting that these cells had preceded further in
their maturation (Poulin et al., 2018) (Figure 3F). We then conducted SCENIC analysis to

pinpoint key transcription factors activated by the new protocol (Figure 3E). The paired-box
transcription factor PAX7 transiently became the dominant regulon in NProg, suggesting precise
rostral-caudal patterning as it contributes to determine the midbrain-hindbrain boundary during
regionalization (Agoston et al., 2012; Nakamura et al., 2005). Other leading regulons in NProg
and NbM cells are also actively involved in guiding the neuronal identity acquisition. For
instance, POU3F1 and ZEB1 are essential regulators of neural lineage commitment, often
working synergistically to coordinate correct orders of neuronal development, as POU3F1
promotes early neural fate decisions, while ZEB1 modulates the developmental timing and
prevents premature differentiation (Bella et al., 2021; Gheldof et al., 2011). Notably, PITX3
emerged as a leading regulon in mDA neurons, which reflects the progressive acquisition of
mature mDA phenotypes (Smidt et al., 2004) and highlights their molecular similarity to
endogenous SNpc mDA neurons (Figure 3F,E). Regulon activity profiles from the integrated
dataset revealed a distinct and significant regulatory impacts of TCF4 and TCF12 under the new
protocol (Sup. Figure 4C). Interestingly, the new protocol also appeared to enhance SREBF1
expression throughout differentiation, linking it to its known role in regulating midbrain
dopaminergic neurogenesis (Toledo et al., 2020), and its involvement in controlling neuronal
dendrite expansion through lipid homeostasis (Ziegler et al., 2017).

Overall, the optimization of mDA neuron differentiation protocol resulted in significant
advancements in mDA neuron specification and maturation, with higher yield of mDA neurons
resembling transcriptomically the ventral subset of human native DA nigral neurons.

478　Differentiated hiPSC instrastriatal transplant matures into innervated DANs and
479　establishes motor recovery in hemiPD mice

After establishing that this protocol produces an enriched ventralized subpopulation *in vitro*, we
proceeded to further evaluate these cells *in vivo* using rodent hemiparkinsonian models as
proof-of-concept for potential therapeutic approaches and cell differentiation in a more
physiological and pathological context. We lesioned unilaterally B6RGS mice by injecting 6-
OHDA in the MFB, and once confirmed hemiPD, we randomly selected mice to be injected with
200K cells intrastriatally. Motor recovery was assessed continuously by amphetamine-induced
ipsilateral rotations, showing that grafted hemiPD animals returned to pre-lesion state 4 months
post-implant (m.p.i.), which was maintained 1 month later, ending our grafting experiment
(Sham: -0.24 ± 0.73; 6-OHDA: 13.41 ± 1.43; Grafted: -0.70 ±1.18). To ensure overall motor
recovery, we also analysed complex motor coordination by tape removal test (Sham: -0.40 ±
0.34; 6-OHDA: 19.51 ± 2.50; Grafted: -2.82 ±0.96) and spontaneous motor activity by cylinder
wall test (Sham: 51.82 ± 3.73%; 6-OHDA: 8.85 ± 4.75%; Grafted: 40.89 ± 3.73%). Human cells
were detected in these 5-month grafts by STEM121$^+$ staining, showing that grafted cells did not
migrate to other regions (Fig. 4C). Furthermore, human DANs were detected by double positive
staining of TH and HuNu, with these cells showing extended prolongations, characteristic of
mDA neurons (Fig. 4D, E). Using stereological quantification of TH$^+$/HuNu$^+$ cells (Fig. 4F), grafts
were estimated to have 160000 human nuclei (155700 ± 35893), of which about 10% were also
stained positively for TH (10.36 ± 0.73%). After establishing the presence of DANs within our
grafts, we aimed to analyse if these DANs recapitulated the phenotypical aspects of the most
PD-vulnerable portion of nigral DANs which have been reported to be A9-subtype (DANs from
the SNpc), more specifically the SOX6$^+$/AGTR1a$^+$ subpopulation located in the ventral SNpc

(Kamath et al., 2022; Salmani et al., 2024). To this aim, we performed a series of IHC analysis of
the grafted DANs. A grand majority of grafted TH$^+$ cells were found to be positive for several
markers distinctive to the PD-vulnerable A9-subtype (Fig 4G,H), such as SOX6 (84.67 ± 3.03%),
ALDH1A1 (50.67 ± 2.37%) and AGTR1 (64.16 ± 4.97%). In contrast, a minority of the DANS
stained positively for the A10-subtype marker Calbindin (29.92 ± 3.45%). Another aspect that we
wanted to evaluate is the expression of canonical markers for mDA neurons in the grafted TH$^+$
cells (Fig. 4I,J). We performed IHC and we were able to identify approximately all of the grafted
DANs as LMX1A$^+$ (92.09 ± 3.37%), FOXA2$^+$ (96.66 ± 1.88%), NR4A2$^+$ (96.19 ± 2.16%) and PITX3$^+$
(85.14 ± 1.80%). Finally, we performed another series of IHC to evaluate expression of proteins
involved in DA regulation and secretion (Fig. 4K,L). We were able to determine that the grafted
DANs expressed DAT (90.03 ± 1.23%), GIRK2 (96.78 ± 2.12%), VMAT (94.24 ± 1.47%) and DDC
(100.00 ± 0.00%). Overall, we were able to demonstrate by IHC that grafted cells mature into a
large percentage of DANs expressing canonical markers for A9-subtype mDA neurons that are
able to ameliorate motor degeneration in rodent PD models.

snRNA sequencing of grafts identifies a population of A9-subtype mDA neurons *in vivo*

One of the several caveats to approach regarding grafted cells is the overall composition of the
transplant, as they may contain non-mDA cells that could disturb the putative therapeutic
population, and how much do the grafted cells resemble the native human cells. To do so, we
proceeded to analyse the transcriptomic profile of snap-frozen endpoint graft tissue punches of
3 grafted mice by snRNA-sequencing, and after data quality filtering we obtained 9196 high-
quality human nuclei. Grafted differentiated human cells separated into 4 different cluster
comprised of neurons, radial glia, gliobasts and oligodendrocyte precursors (Fig. 5A). An in-
depth analysis of the neuronal clusters classified into DAergic, Glutamatergic, GABAergic
(defined by RBFOX3) and neural progenitors (defined by INA). Interestingly, the DAergic cluster
was comprised of 3 subclusters: DA_SOX6, DA_CALB1 and DA_ONECUT2 (Fig. 5B).
Furthermore, evaluating the expression levels of genes of interest to each cluster, we were able
to identify that each DA subcluster has individually identifiable transcriptomics profile. DA
clusters presented high expression levels of genes related to DA cells (*TH, DDC, NR4A2, LMX1A,*
PBX1, SLC18A2 [VMAT2], *SLC6A3* [DAT], *PITX3, KCNJ6* [GIRK2]), while in contrast, GABA cluster
and Glut clusters did not, expressing however high levels of *GAD1* and *GAD2*, and *SLC17A6*
[VGLUT2] respectively. INA clusters were able to be distinctly separated from Glut clusters by
expression levels of *SLC17A6, STMN2 and INA*. Furthermore, DA subclusters can be distinctly
separated by differential expression levels of *SOX6*, *CALB1* and *ONECUT2*. (Fig. 5C,D).
Approximately half of the cells from the DA subclusters were classified as DA_SOX6 (44.73 ±
4.79%), with the other half comprised of a majority of DA_CALB1 and a small proportion of
DA_ONECUT2 (Fig. 5E). Our findings show that grafted cells differentiated with this optimized
protocol show promising potential to replace SOX6$^+$ specific subtypes of PD-susceptible mDA
neurons.

Comparison with pre-established transcriptomic dataset finds similarities between adult
human midbrain and grafted cells

As we have been able to identify DANs within our human grafted cells, , we wanted to reassure
that our hPSCs resembled the expected transcriptomic profile of the human native mDA cells.
For such purpose, we performed a MetaNeighbor analysis (Crow et al., 2018) with our DA

clusters and DA clusters of the human substantia nigra dataset from Kamath et al. (Kamath et
al., 2022) after random downsampling the human dataset to avoid false correlations (Fig 5F).
Our analysis showed high distinctive resemblance with a score >0.8 between our DA_SOX6
subcluster and the SOX6_AGTR1 subcluster from Kamath dataset, which have been reported to
be the most susceptible subpopulation to PD (Kamath et al., 2022; Salmani et al., 2024) . In
contrast, the DA_ONECUT2 subcluster showed resemblance to the human CALB1_TRHR
reported to be the most resilient. However, the DA_CALB1 subcluster was not able to achieve a
distinct resemblance to a specific subpopulation but rather most of the CALB1 human clusters,
indicating that these cells are not fully dedicated to a specific subtype and may be prone to
further differentiation. This data reinforces that we were successfully able to generate cells that
differentiate *in vivo* to cells resembling human SOX6 mDA neurons.

555   Analysis of gene regulatory networks of interest show that the distinct subpopulations of
556   SOX6[+] DANs do not shift back to other subtypes

Based on our previous finding about the resemblance of human transcriptomic profile of the DA
subcluster, we wanted to evaluate if these cells could potentially lose their distinct identity. We
first evaluated the functional transcriptome profile of each cluster based on the GO terms.
Based on this analysis, we observed that the DA_SOX6 subcluster is associated with GO terms
related to late-stage mature neurons, for example, modulation of synaptic transmission or
synapse organization. On contrast, DA_CALB1 is related to GO terms associated with learning,
and DA_ONECUT2 with regulation of neuron and cell projection. This indicates that while all
subclusters share similar functions as late-stage neurons, each one has their own defining GO
clustering. We continued by performing RNA velocity analysis with scVelo (Bergen et al., 2020;
Gayoso et al., 2024) on our DA and Immature neurons subclusters and we found that Immature
neurons have the potential to become DA_CALB1 cells. Interestingly, the DA_CALB1
subclusters have the potential to become either DA_ONECUT2 or DA_SOX6, which in contrast
their velocity show that these identities are an endpoint (Fig 5H). Further evaluation of the gene
regulatory network by SCENIC establishes that the DA clusters are enriched in exclusive TFs
expression profile, not found in the immature neuron, GABA or Glut clusters. For example, PBX1
regulon networks have been associated with ventral midbrain DA organization and is highly
enriched (score >0.8) in our DA cluster. Moreover, the DA_SOX6 is differentially highly enriched
in SOX6 (score >0.8) and CUX2 (score >0.6) within the DA clusters. Interestingly, the CUX2
regulon network has been reported to be highly associated with the SOX6[+]AGTR1[+] clusters of
neurons in human datasets, further validated in PD macaque models (Kamath et al, 2022; Tang
et al., 2023).

Overall, as proof-of-concept the combination of the immunohistochemical and transcriptomic
data reflects that this new accelerated *in vitro* protocol is able to produce mDA precursors by
day 14, which, upon transplantation into a PD model, exhibiting successful motor recovery,
selectively differentiate in a majority of PD-susceptible neurons.

583   DISCUSSION

The study presents an enhanced protocol for differentiating midbrain dopaminergic (mDA)
neurons from human induced pluripotent stem cells (hiPSCs) based on multiomic

transcriptomic analysis of different itterations, highlighting improvements in patterning
efficiency and biological reproducibility. This development holds significant promise for PD
research, as it addresses key limitations in the previously published differentiation protocol,
such as prolonged differentiation timelines, suboptimal ventral patterning, and inconsistent
generation of mature mDA neurons across different cell lines.

The single nuclei multiomic profiling of differentiated cells at sequential timepoints revealed
distinct regulatory networks governing the transition from neural progenitors to mature
dopaminergic neurons. By integrating scRNA-seq and scATAC-seq, the study identified
temporal shifts in chromatin accessibility and transcriptional activity that mirrored the
development of the human midbrain. For instance, the emergence of neurons expressing TH,
DDC and KCNJ6 at DIV42 correlated with elevated activity of transcription factors such as PBX1,
which is known to play a crucial role in mDA neuron survival and functional maturation.

Based on the transcriptomic profiling, it was determined that the current protocol could benefit
from the iterative enhancement of the ventralization process, which improved mDA patterning
evidenced by elevated EN1 expression (a key marker for mDA neuron specification). This
adjustment was particularly beneficial in producing neurons that more closely resemble the
human SNpc dopaminergic neurons, the primary neuronal population affected in Parkinson's
disease. Furthermore, we were able to identify regulatory networks involving mDA neuron
maturation enriched in the new protocol, such as TCF12 and TCF4. Specifically, TCF12 plays a
dual role in the early commitment of neural progenitors and the late specification of neuronal
subsets, with its loss leading to altered expression of TH and AHD2 (ALDH1A1), the latter being
crucial for the specification and maintenance of dopaminergic neurons, particularly in the A9
dopaminergic neuronal group (Chung et al., 2005). TCF4, on the other hand, is likely involved in
the fine-tuning of neuron identity during the onset of mDA neuron subtype specification. In a
mouse study, deletion of TCF4 led to delayed maturation of dopaminergic neurons, primarily
through transient suppression of *Ahd2* and *Nxph3* (Mesman et al., 2021).

One of the critical advancements was the precise modulation of the midbrain floor plate (mFP)
patterning using optimized coating conditions (LN511) and biphasic activation of WNT signaling
(CHIR boost), which resulted in a substantial population of mFP progenitors by day 11 of
differentiation and improved the expression of essential mDA markers such as FOXA2, LMX1A,
and OTX2. This suggests that the updated protocol not only enhances patterning but also
reduces contamination by non-neural cell types, a persistent issue in generating region-specific
neurons for disease modelling. Notably, the cellular composition differed from that generated
using the original protocol due to enhanced patterning efficacies of floorplate progenitors, and
the removal of the MEK/ERK pathway inhibitor PD0325901 and the FGF receptor inhibitor
SU5402, that mainly used to repress expansion of SOX2+ radial glia and progenitor cells (Koch
et al., 2009).

Despite the absence of pro-neural effects from PD0325901 and SU5402 in the new protocol, our
cell cultures still produced a greater number of mDA neurons with more mature morphological
and molecular phenotypes, particularly in terms of synapse assembly, relying solely on intrinsic
mechanisms. Moreover, the integration of synthetic LXR ligands (GW3965) further accelerated
neurogenesis process, leading to the earlier emergence of mature dopaminergic neurons.
Specifically, TH$^+$ neurons first appeared at day 16, notably earlier than in previously published

protocols, where they typically emerged around day 21. The early establishment of mDA
neurons facilitates downstream functional assays, such as live-cell calcium recordings and
neurotransmitter release analysis, thus shortening the experimental timeline and allowing for
earlier detection of phenotypic differences.

Subsequent electrophysiological analysis via live-cell calcium recordings demonstrated that
mDA neurons derived using this protocol manifested hallmarks of functionally active neurons,
such as spontaneous electrical activity and robust responses to depolarization stimuli (Chan et
al., 2007; Pissadaki and Bolam, 2013) . These findings are significant, as they confirm that the
cells generated are not merely phenotypically similar to mDA neurons but also functionally
active, which is critical for accurate disease modeling and therapeutic screening.

The study successfully achieved scalable and reproducible mDA neuron differentiation across
multiple iPSC lines, including those derived from Parkinsonian patients, suggesting that this
protocol can serve as a standardized approach for large-scale production of high-quality
dopaminergic neurons. Such advancements are pivotal for the development of robust in vitro
models to study neurodegenerative diseases and for testing potential therapeutic interventions.

Finally, as proof-of concept validation for therapeutic approaches, we were able to determine
that the new protocol could produce mFP progenitors and NbM able to differentiate *in vivo* to
mDA neurons resembling the A9-subtype nigral neurons in majority, based on expression of
proteins such as SOX6, AGTR1, GIRK2 or ALDH1A, reported to be more susceptible to PD in
human and animal models (Kamath et al., 2022, del Rey et al.,2024, Nie et al., 2024, Salmani et
al., 2024). Furthermore, evaluation of the transcriptomic profile of the DA neurons in the graft
showed high resemblance to the human counterpart, specially the SOX6-AGTR1A mDA
neurons. This proposes that these cells could have great potential for directed replacement
therapies.

While these advancements could be of great interest to the PD research community, we have to
address several caveats. First, we only have established the replicability of this protocol on
hIPSC, both WT and with GBA mutations. But further characterization with more well known
mutations found in familial PD, such as GBA E326K, LRRK2 G2019S or the triplicate SNCA,
could elucidate the validity of this enhanced protocol. Furthermore, the use of human
embryonic stem cells has not been proposed for this study. More *in vitro* studies would be of
interest to validate further this new protocol. Another caveat to address is the pre-clinical
aspect, as the *in vivo* experiments were only used as proof-of-concept for therapeutical
approaches. Further grafting experiments with PD-mutated hiPSC cell lines and a more in-depth
preclinical analysis could detail if this protocol would be a much better advantage that the
current available protocols, which so far have shown to be safe in clinical trials (Sawamoto et
al., 2025, Tabar et al;, 2025). Moreover, the longevity of the grafting experiment was decided
upon stable motor recovery, so we do not have an overview of the long-term effects of these
grafts, whether other contaminants can appear or tumorigenesis in aged individuals.

CONCLUSIONS

In conclusion, the optimized differentiation protocol represents a significant step forward for stem cell-based approaches in modeling PD and for future applications in regenerative medicine. By boosting the timed expression of EN1, we have achieved a more efficient and consistent generation of mDA neurons, with mature phenotypes targeting the A9 subtype defined mainly by the expression of SOX6, ALDH1A, GIRK2 and AGTR1. The study sets the stage for more accurate disease modeling and lays the groundwork for potential directed cell replacement strategies.

ACKNOWLEDGMENTS

RGS., GL and EA conceptualized the experiments. RGS and GL redacted the manuscript. RGS performed *in vivo* experiments; GL, RK, CS, CT and AX contributed to the *in vitro* protocol development. JCK contributed to the calcium imaging analysis. GL and AX contributed to the *in vitro* transcriptional analysis; RGS and CA performed the graft nuclei isolation, and GL the transcriptional analysis. CS, GCB and OD reviewed the manuscript.

This work was mainly supported by the European Union's Horizon 2020 (H2020) Research and Innovation Programme (Neurostemcell-Reconstruct 874758/EA) and the European Research Council (ERC) PreciseCellPD grant (Grant No. 884608/EA). Other contributors: Knut and Alice Wallenberg Foundation Wallenberg Scholar 2018.0232/EA, Vetenskaprådet VR2020-01426/EA, Chan Zuckerberg Initiative and the Silicon Valley Community Foundation (2018-191929)/EA, H2020 MSCA-ITN ASCTN/EA, H2020 Grant No. 899687 - HS-SEQ, Aligning Science Across Parkinson's (ASAP-20505)/EA, Hjärnfonden project FO2024-0195/EA. JCK and PU were supported by the German Science Foundation (516641042) and the Wenner-Gren foundation (UPD2022-0159) respectively.

The authors acknowledge support from the National Genomics Infrastructure in Stockholm funded by Science for Life Laboratory, the Knut and Alice Wallenberg Foundation and the Swedish Research Council, and NAISS/Uppsala Multidisciplinary Center for Advanced Computational Science for assistance with massively parallel sequencing and access to the UPPMAX computational infrastructure. We are really thankful for the help of the Molecular Neurobiology unit of the Medical Biochemistry and Biophysics of the Karolinska Institute, especially Gonçalo Castelo-Branco and Onur Dagliyan for their input. We posthumously want to thank Prof. Ernest Arenas, whom without this project could not have been possible.

704 **Figure 1. Multiomic single-nuclei RNA sequencing revealed the diversity of cellular**
705 **composition throughout of mDA neuron differentiation *in vitro*.** (A) Uniform manifold

approximation and projection (UMAP) of time-course cell preparations profiled with simultaneous snRNA-seq and snATAC-seq, along with their derivative joint UMAP embedding showing clustering of cells based on both transcriptomic and chromatin accessibility profiles, colored by different time points from DIV 11 to 56. (B) Annotated bimodal UMAP embedding of mDA differentiation in vitro using KOLF2.1J iPSCs. Cell types were manually annotated with top-ranked marker genes and their enriched GO terms. (C) Proportion of cells belonging to each cell type at each differentiation time point. (D) A heatmap of AUROC scores comparing annotated cell types derived from *in vitro* differentiation in this study with those from a publicly available first-trimester human midbrain dataset. (E) A dot plot visualizing the expression levels and the percentage of cells expressing curated key genes across cell types during mDA neuron differentiation. (F) Normalized chromatin accessibility tracks of representative marker genes (LMX1A, TH and NR4A2) of dopaminergic lineage cell types. (G) UMAP embeddings showing RNA velocity projection in mDA-related cell types, and a heatmap displaying gene expression dynamics of cluster driver genes ordered by their inferred latent time. (H) Regulon specificity score (RSS) plots highlighting the top 5 cell type-specific transcription factors involved in mDA differentiation.

Figure 2. Prolonged ventralization promotes accelerated differentiation of mDA neurons.
(A) Schematic depiction of the protocol for differentiating mDA neurons, illustrating the
application of pharmacological compounds and growth factors. Representative
immunocytochemical images and quantification from both original protocol and new protocol
showing (B, E) efficacies of midbrain floorplate patterning at DIV11, showing co-localizations of
LMX1+/EN1+ and SOX6+/OTX2+; and EN1+/SOX6+/TH+ neurons at DIV 16 (C, F) and 28 (D, G).
White scale bars = 50, 100 μm as indicated (n = 3 independent experiments, error bars are SEM,
ns = not significant; * = p<0.05; ** = p<0.01; *** = p<0.001; **** = p<0.0001). (H) Calcium
imaging recordings of spontaneous and KCl-induced activities on DIV 28 and 42, presenting the
total number of calcium events per neuron recorded, and the percentages of cells responding to
KCl-induced depolarization. Metrics of activities, including amplitude, rise time, and fall time of
calcium signals, are depicted in the subsequent bar plots.

738
Figure 3. **Cell preparations under the new protocol suggested improved molecular**
phenotypes during *in vitro* midbrain differentiation. (A) UMAP embeddings of mDA
differentiation in vitro using KOLF2.1J iPSCs under the new protocol, color coded by different
time points at DIV 14, 16 and 28. (B) UMAP embeddings of detailed cell types were manually
annotated using top-ranked marker genes and their enriched GO terms, and proportion of cells
classified into each cell type per differentiation time point. (C) Dot plot visualizing the
expression levels and the percentage of cells expressing curated key genes across distinct cell
types generated under the new protocol. (D) Pseudotime trajectory analysis showing
progression from progenitor-like populations (ProgFP, NProg, and NbM) to differentiated DA
neurons. Cells are ordered by inferred pseudotime along the x-axis. (E) Regulon specificity score
(RSS) plots highlighting the top 5 cell type-specific transcription factors driving mDA
differentiation under the new protocol. (F) Dot plots showing the expression levels and the
percentage of cells expressing midbrain dopaminergic lineage marker genes after integration in
(left) midbrain cells categorized as dopaminergic neurons (DA) (right), dopaminergic lineage cell
types across all timepoints and both conditions.

Figure 4. Transplantation of hiPSC differentiated with new protocol. (A) Diagram of the experimental design for the implantation of day 14-differentiated hiPSC in B6RGS mice. (B) Motor recovery is observed by continuous analysis of amphetamine-induced ipsilateral rotation (left), endpoint tape removal test (middle) and endpoint cylinder wall analysis (right). Of note, grafted animals recover pre-lesion state while the 6-OHDA control animals maintain the lesioned state. (C) Example of grafted tissue, stained against STEM121, showing the human cells within the graft. (D) Micrograph of graft stained against TH and HuNu, showing hDANs all over the graft. (E) Closeup of grafted hDANs showing elongated and ramified extensions. (F) Stereological estimate of HuNu$^+$ nuclei (left) and percentage of hDANs (TH$^+$-HuNu$^+$) in grafts (right). (G) Representative examples of hDANs stained positive for A9-subtype marker, SOX6, ALDH1A1, AGTR1A and A10 marker, Calbindin. (H) Quantification of percentage of hDANs (TH$^+$-HuNu$^+$) expressing SOX6, ALDH1A1, AGTR1A and Calbindin. (I) Representative examples of hDANs stained positive for mDA neuron makers, LMX1A, FOXA2, NR4A2 and PITX3. (J) Quantification of percentage of hDANs (TH$^+$-HuNu$^+$) expressing LMX1A, FOXA2, NR4A2 and PITX3. (K) Representative examples of hDANs stained positive for proteins involved in DA excretion, DAT, GIRK2, VMAT2 and DDC. (L) Quantification of percentage of hDANs (TH$^+$-HuNu$^+$) expressing DAT, GIRK2, VMAT2 and DDC. m.p.i = months post-implant. N for animal behavioral experiment = sham: 6; 6-OHDA: 8; hIPSC-grafted: 8. N for IHC = 5. Scale bar = C: 1mm; D: 500 µm; E: 100 µm; G,I,K: 10 µm.

Figure 5. An in-depth single nucleus RNA sequencing characterization of *in vivo* xenograft cellular composition. Annotated UMAP embeddings of (A) 9,196 high-quality human nuclei recovered from three grafted mice at five months post-transplantation, with cells classified into neurons, radial glia, gliobasts and oligodendrocyte precursors; and (B) neuronal population was consisted of immature neurons, GABAergic neurons, Glutamatergic neurons and 3 subtypes of mDA neurons, each characterized by specific transcription factors, as DA_CALB1, DA_ONECUT2, and DA_SOX6. (C) Dot plot showing average gene expression and percentage of expressing cells for selected marker genes across annotated neuronal cell types. (D) Violin plots

of representative marker genes for dopaminergic subtypes (SOX6, ONECUT2, CALB1), pan-DA
markers (TH, NR4A2, KCNJ6), illustrating subtype-specific and shared expression patterns. (E)
Average proportion of yielded human nuclei in each cellular hierarchy across three individual
datasets, demonstrating consistent representation of DA subtypes (N=3). (F) MetaNeighbor
analysis between transplanted mDA neurons in this study and a reference dataset of DA
neurons from the SNpc region of PD patients and matched controls, suggesting transcriptional
similarities between DA_SOX6 subtype and SOX6_AGTR1 subtype from the reference dataset.
(G) Gene Ontology (GO) enrichment analysis of DA subtype-specific markers, showing distinct
biological processes enriched in DA_SOX6, DA_CALB1, and DA_ONECUT2 populations. (H) RNA
velocity projection of mDA neuron trajectories in UMAP embeddings, suggesting differentiation
pathways among different DA subtypes and immature neurons. (I) Heatmap of average regulon
activities of the top 30 transcription factors across mDA and other neuronal subtypes.

BIBLIOGRAPHY

Agoston, Z., Li, N., Haslinger, A., Wizenmann, A., & Schulte, D. (2012). Genetic and physical
interaction of Meis2, Pax3 and Pax7 during dorsal midbrain development. *BMC*
Developmental Biology, *12*(1), 10. https://doi.org/10.1186/1471-213X-12-10

Aibar, S., González-Blas, C. B., Moerman, T., Huynh-Thu, V. A., Imrichova, H., Hulselmans, G.,
Rambow, F., Marine, J. C., Geurts, P., Aerts, J., Van Den Oord, J., Atak, Z. K., Wouters, J., &
Aerts, S. (2017). SCENIC: Single-cell regulatory network inference and clustering. *Nature*
Methods, *14*(11), 1083–1086. https://doi.org/10.1038/NMETH.4463

Albéri, L., Sgadò, P., & Simon, H. H. (2004). Engrailed genes are cell-autonomously required to
prevent apoptosis in mesencephalic dopaminergic neurons. *Development*, *131*(13), 3229–
3236. https://doi.org/10.1242/DEV.01128

Alquicira-Hernandez, J., & Powell, J. E. (2021). Nebulosa recovers single-cell gene expression
signals by kernel density estimation. *Bioinformatics*, *37*(16), 2485–2487.
https://doi.org/10.1093/bioinformatics/btab003

Andersson, E., Jensen, J. B., Parmar, M., Guillemot, F., & Björklund, A. (2006). Development of
the mesencephalic dopaminergic neuron system is compromised in the absence of
neurogenin 2. *Development*, *133*(3), 507–516. https://doi.org/10.1242/dev.02224

Bankhead, P., Loughrey, M. B., Fernández, J. A., Dombrowski, Y., McArt, D. G., Dunne, P. D.,
McQuaid, S., Gray, R. T., Murray, L. J., Coleman, H. G., James, J. A., Salto-Tellez, M., &
Hamilton, P. W. (2017). QuPath: Open source software for digital pathology image analysis.
Scientific Reports 2017 7:1, *7*(1), 1–7. https://doi.org/10.1038/s41598-017-17204-5

Bergen, V., Lange, M., Peidli, S., Wolf, F. A., & Theis, F. J. (2020). Generalizing RNA velocity to
transient cell states through dynamical modeling. *Nature Biotechnology 2020 38:12*,
38(12), 1408–1414. https://doi.org/10.1038/s41587-020-0591-3

Björklund, A., & Dunnett, S. B. (2019). The Amphetamine Induced Rotation Test: A Re-
Assessment of Its Use as a Tool to Monitor Motor Impairment and Functional Recovery in
Rodent Models of Parkinson's Disease. *Journal of Parkinson's Disease*, *9*(1), 17–29.
https://doi.org/10.3233/JPD-181525

Björklund, A., & Parmar, M. (2021). Dopamine Cell Therapy: From Cell Replacement to Circuitry
Repair. *Journal of Parkinson's Disease*, *11*(Suppl 2), S159. https://doi.org/10.3233/JPD-
212609

Bodea, G. O., Spille, J.-H., Abe, P., Andersson, A. S., Acker-Palmer, A., Stumm, R., Kubitscheck,
U., & Blaess, S. (2014). Reelin and CXCL12 regulate distinct migratory behaviors during the
development of the dopaminergic system. *Development*, *141*(3), 661–673.
https://doi.org/10.1242/dev.099937

Braun, E., Danan-Gotthold, M., Borm, L. E., Lee, K. W., Vinsland, E., Lönnerberg, P., Hu, L., Li, X.,
He, X., Andrusivová, Ž., Lundeberg, J., Barker, R. A., Arenas, E., Sundström, E., &
Linnarsson, S. (2023). Comprehensive cell atlas of the first-trimester developing human
brain. *Science (New York, N.Y.)*, *382*(6667), eadf1226.
https://doi.org/10.1126/science.adf1226

Brichta, L., Greengard, P., & Flajolet, M. (2013). Advances in the pharmacological treatment of
Parkinson's disease: targeting neurotransmitter systems. *Trends in Neurosciences*, *36*(9),
543–554. https://doi.org/10.1016/j.tins.2013.06.003

Cao, J., Spielmann, M., Qiu, X., Huang, X., Ibrahim, D. M., Hill, A. J., Zhang, F., Mundlos, S.,
Christiansen, L., Steemers, F. J., Trapnell, C., & Shendure, J. (2019). The single-cell
transcriptional landscape of mammalian organogenesis. *Nature 2019 566:7745*, *566*(7745),
496–502. https://doi.org/10.1038/s41586-019-0969-xr

Chan, C. S., Guzman, J. N., Ilijic, E., Mercer, J. N., Rick, C., Tkatch, T., Meredith, G. E., &
Surmeier, D. J. (2007). "Rejuvenation" protects neurons in mouse models of Parkinson's
disease. *Nature*, *447*(7148), 1081–1086. https://doi.org/10.1038/NATURE05865

Chung, S., Hedlund, E., Hwang, M., Kim, D. W., Shin, B. S., Hwang, D. Y., Kang, U. J., Isacson, O.,
& Kim, K. S. (2005). The homeodomain transcription factor Pitx3 facilitates differentiation of
mouse embryonic stem cells into AHD2-expressing dopaminergic neurons. *Molecular and*
Cellular Neuroscience, *28*(2), 241–252. https://doi.org/10.1016/J.MCN.2004.09.008

Clarke, Z. A., Andrews, T. S., Atif, J., Pouyabahar, D., Innes, B. T., MacParland, S. A., & Bader, G.
D. (2021). Tutorial: guidelines for annotating single-cell transcriptomic maps using
automated and manual methods. *Nature Protocols 2021 16:6*, *16*(6), 2749–2764.
https://doi.org/10.1038/s41596-021-00534-0

Crow, M., Paul, A., Ballouz, S., Huang, Z. J., & Gillis, J. (2018). Characterizing the replicability of
cell types defined by single cell RNA-sequencing data using MetaNeighbor. *Nature*
Communications 2018 9:1, *9*(1), 1–12. https://doi.org/10.1038/s41467-018-03282-0

Damier, P., Hirsch, E. C., Agid, Y., & Graybiel, A. M. (1999). The substantia nigra of the human
brain: II. Patterns of loss of dopamine-containing neurons in Parkinson's disease. *Brain*,
122(8), 1437–1448. https://doi.org/10.1093/BRAIN/122.8.1437

del Rey, N. L. G., Hernández-Pinedo, N., Carrillo, M., del Cerro, M., Esteban-García, N., Trigo-
Damas, I., Monje, M. H. G., Lanciego, J. L., Cavada, C., Obeso, J. A., & Blesa, J. (2024).
Calbindin and Girk2/Aldh1a1 define resilient vs vulnerable dopaminergic neurons in a
primate Parkinson's disease model. *Npj Parkinson's Disease 2024 10:1*, *10*(1), 1–15.
https://doi.org/10.1038/s41531-024-00777-0

Dennis, D. J., Han, S., & Schuurmans, C. (2019). bHLH transcription factors in neural
development, disease, and reprogramming. *Brain Research*, *1705*, 48–65.
https://doi.org/10.1016/J.BRAINRES.2018.03.013

Desai, A. R., & McConnell, S. K. (2000). Progressive restriction in fate potential by neural
progenitors during cerebral cortical development. *Development*, *127*(13), 2863–2872.
https://doi.org/10.1242/DEV.127.13.2863

Di Bella, D. J., Habibi, E., Stickels, R. R., Scalia, G., Brown, J., Yadollahpour, P., Yang, S. M.,
Abbate, C., Biancalani, T., Macosko, E. Z., Chen, F., Regev, A., & Arlotta, P. (2021).
Molecular logic of cellular diversification in the mouse cerebral cortex. *Nature 2021*
595:7868, *595*(7868), 554–559. https://doi.org/10.1038/s41586-021-03670-5

Doi, D., Samata, B., Katsukawa, M., Kikuchi, T., Morizane, A., Ono, Y., Sekiguchi, K., Nakagawa,
M., Parmar, M., & Takahashi, J. (2014). Isolation of human induced pluripotent stem cell-
derived dopaminergic progenitors by cell sorting for successful transplantation. *Stem Cell*
Reports, *2*(3), 337–350. https://doi.org/10.1016/j.stemcr.2014.01.013

Domínguez Conde, C., Xu, C., Jarvis, L. B., Rainbow, D. B., Wells, S. B., Gomes, T., Howlett, S. K.,
Suchanek, O., Polanski, K., King, H. W., Mamanova, L., Huang, N., Szabo, P. A., Richardson,
L., Bolt, L., Fasouli, E. S., Mahbubani, K. T., Prete, M., Tuck, L., … Teichmann, S. A. (2022).
Cross-tissue immune cell analysis reveals tissue-specific features in humans. *Science*
(New York, N.Y.), *376*(6594), eabl5197. https://doi.org/10.1126/science.abl5197

Faul, F., Erdfelder, E., Lang, A. G., & Buchner, A. (2007). G*Power 3: a flexible statistical power
analysis program for the social, behavioral, and biomedical sciences. *Behavior Research*
Methods, *39*(2), 175–191. https://doi.org/10.3758/BF03193146

Ferri, A. L. M., Lin, W., Mavromatakis, Y. E., Wang, J. C., Sasaki, H., Whitsett, J. A., & Ang, S. L.
(2007). Foxa1 and Foxa2 regulate multiple phases of midbrain dopaminergic neuron
development in a dosage-dependent manner. *Development*, *134*(15), 2761–2769.
https://doi.org/10.1242/DEV.000141

Garritsen, O., van Battum, E. Y., Grossouw, L. M., & Pasterkamp, R. J. (2023). Development,
wiring and function of dopamine neuron subtypes. *Nature Reviews Neuroscience*, *24*(3),
134–152. https://doi.org/10.1038/s41583-022-00669-3

Gayoso, A., Weiler, P., Lotfollahi, M., Klein, D., Hong, J., Streets, A., Theis, F. J., & Yosef, N.
(2023). Deep generative modeling of transcriptional dynamics for RNA velocity analysis in
single cells. *Nature Methods 2023 21:1*, *21*(1), 50–59. https://doi.org/10.1038/s41592-023-
01994-w

Gheldof, A., Hulpiau, P., van Roy, F., De Craene, B., & Berx, G. (2012). Evolutionary functional
analysis and molecular regulation of the ZEB transcription factors. *Cellular and Molecular*
Life Sciences : CMLS, *69*(15), 2527–2541. https://doi.org/10.1007/s00018-012-0935-3

Gleeson, J. G., Lin, P. T., Flanagan, L. A., & Walsh, C. A. (1999). Doublecortin is a microtubule-
associated protein and is expressed widely by migrating neurons. *Neuron*, *23*(2), 257–271.
https://doi.org/10.1016/s0896-6273(00)80778-3

Greene, P. E., Fahn, S., Eidelberg, D., Bjugstad, K. B., Breeze, R. E., & Freed, C. R. (2021).
Persistent dyskinesias in patients with fetal tissue transplantation for Parkinson disease.
NPJ Parkinson's Disease, 7(1). https://doi.org/10.1038/S41531-021-00183-W

Jiang, Y. Q., & Oblinger, M. M. (1992). Differential regulation of βiii and other tubulin genes during
peripheral and central neuron development. *Journal of Cell Science*, *103*(3), 643–651.
https://doi.org/10.1242/JCS.103.3.643

Kamath, T., Abdulraouf, A., Burris, S. J., Langlieb, J., Gazestani, V., Nadaf, N. M., Balderrama, K.,
Vanderburg, C., & Macosko, E. Z. (2022). Single-cell genomic profiling of human dopamine
neurons identifies a population that selectively degenerates in Parkinson's disease. *Nature*
Neuroscience 2022 25:5, *25*(5), 588–595. https://doi.org/10.1038/s41593-022-01061-1

Kawamoto, E. M., Vivar, C., & Camandola, S. (2012). Physiology and Pathology of Calcium
Signaling in the Brain. *Frontiers in Pharmacology*, *3*, 22359.
https://doi.org/10.3389/fphar.2012.00061

Kele, J., Simplicio, N., Ferri, A. L. M., Mira, H., Guillemot, F., Arenas, E., & Ang, S. L. (2006).
Neurogenin 2 is required for the development of ventral midbrain dopaminergic neurons.
Development, *133*(3), 495–505. https://doi.org/10.1242/DEV.02223

Kim, T. W., Piao, J., Koo, S. Y., Kriks, S., Chung, S. Y., Betel, D., Socci, N. D., Choi, S. J.,
Zabierowski, S., Dubose, B. N., Hill, E. J., Mosharov, E. V., Irion, S., Tomishima, M. J., Tabar,
V., & Studer, L. (2021). Biphasic Activation of WNT Signaling Facilitates the Derivation of
Midbrain Dopamine Neurons from hESCs for Translational Use. *Cell Stem Cell*, *28*(2), 343-
355.e5. https://doi.org/10.1016/j.stem.2021.01.005

Kim, W. Y. (2012). NeuroD1 is an upstream regulator of NSCL1. *Biochemical and Biophysical*
Research Communications, *419*(1), 27–31. https://doi.org/10.1016/J.BBRC.2012.01.100

Kirkeby, A., Nelander, J., Hoban, D. B., Paul, G., Barker, R. A., Parmar, M., Rogelius, N., Lmar
Bjartmarz, H., Nordisk, N., Storm, P., Fiorenzano, A., Adler, A. F., Vale, S., Mudannayake, J.,
Zhang, Y., Cardoso, T., Mattsson, B., Landau, A. M., Glud, A. N., … Kan Widner, H. (2023).
Clinical and Translational Report ll Clinical and Translational Report Preclinical quality,
safety, and efficacy of a human embryonic stem cell-derived product for the treatment of
Parkinson's disease, STEM-PD. *Stem Cell*, *30*, 1299-1314.e9.
https://doi.org/10.1016/j.stem.2023.08.014

Kirkeby, A., Nelander, J., Hoban, D. B., Rogelius, N., Bjartmarz, H., Storm, P., Fiorenzano, A.,
Adler, A. F., Vale, S., Mudannayake, J., Zhang, Y., Cardoso, T., Mattsson, B., Landau, A. M.,
Glud, A. N., Sørensen, J. C., Lillethorup, T. P., Lowdell, M., Carvalho, C., … Parmar, M.
(2023). Preclinical quality, safety, and efficacy of a human embryonic stem cell-derived
product for the treatment of Parkinson's disease, STEM-PD. *Cell Stem Cell*, *30*(10), 1299-
1314.e9. https://doi.org/10.1016/j.stem.2023.08.014

Koch, P., Opitz, T., Steinbeck, J. A., Ladewig, J., & Brüstle, O. (2009). A rosette-type, self-renewing
human ES cell-derived neural stem cell with potential for in vitro instruction and synaptic
integration. *Proceedings of the National Academy of Sciences of the United States of*
America, *106*(9), 3225–3230. https://doi.org/10.1073/pnas.0808387106

Korsunsky, I., Millard, N., Fan, J., Slowikowski, K., Zhang, F., Wei, K., Baglaenko, Y., Brenner, M.,
Loh, P. ru, & Raychaudhuri, S. (2019). Fast, sensitive, and accurate integration of single cell
data with Harmony. *Nature Methods*, *16*(12), 1289. https://doi.org/10.1038/S41592-019-
0619-0

Kriks, S., Shim, J.-W., Piao, J., Ganat, Y. M., Wakeman, D. R., Xie, Z., Carrillo-Reid, L., Auyeung, G., Antonacci, C., Buch, A., Yang, L., Beal, M. F., Surmeier, D. J., Kordower, J. H., Tabar, V., & Studer, L. (2011). Dopamine neurons derived from human ES cells efficiently engraft in animal models of Parkinson's disease. *Nature*, *480*(7378), 547–551. https://doi.org/10.1038/nature10648

La Manno, G., Gyllborg, D., Codeluppi, S., Nishimura, K., Salto, C., Zeisel, A., Borm, L. E., Stott, S. R. W., Toledo, E. M., Villaescusa, J. C., Lönnerberg, P., Ryge, J., Barker, R. A., Arenas, E., & Linnarsson, S. (2016). Molecular Diversity of Midbrain Development in Mouse, Human, and Stem Cells. *Cell*, *167*(2), 566-580.e19. https://doi.org/10.1016/J.CELL.2016.09.027

Mannens, C. C. A., Hu, L., Lönnerberg, P., Schipper, M., Reagor, C. C., Li, X., He, X., Barker, R. A., Sundström, E., Posthuma, D., & Linnarsson, S. (2024). Chromatin accessibility during human first-trimester neurodevelopment. *Nature 2024*, 1–8. https://doi.org/10.1038/s41586-024-07234-1

McGinnis, C. S., Murrow, L. M., & Gartner, Z. J. (2019). DoubletFinder: Doublet Detection in Single-Cell RNA Sequencing Data Using Artificial Nearest Neighbors. *Cell Systems*, *8*(4), 329-337.e4. https://doi.org/10.1016/j.cels.2019.03.003

Mesman, S., Wever, I., & Smidt, M. P. (2021). Tcf4 Is Involved in Subset Specification of Mesodiencephalic Dopaminergic Neurons. *Biomedicines 2021, Vol. 9, Page 317*, *9*(3), 317. https://doi.org/10.3390/BIOMEDICINES9030317

Murdoch, J. N., Eddleston, J., Leblond-Bourget, N., Stanier, P., & Copp, A. J. (1999). Sequence and expression analysis of Nhlh1: a basic helix-loop-helix gene implicated in neurogenesis. *Developmental Genetics*, *24*(1–2), 165–177. https://doi.org/10.1002/(SICI)1520-6408(1999)24:1/2<165::AID-DVG15>3.0.CO;2-V

Nakamura, H., Katahira, T., Matsunaga, E., & Sato, T. (2005). Isthmus organizer for midbrain and hindbrain development. *Brain Research Reviews*, *49*(2), 120–126. https://doi.org/10.1016/J.BRAINRESREV.2004.10.005

Nie, S., Li, B., Wang, M., Chen, Z., Ren, J., Li, Z., Xu, X., Qian, Z., Xie, Z., Han, J., Zhang, Z., Zhang, Z., Zhu, Y., Chen, Z., Yang, X., & Ye, K. (2025). Sox6 and ALDH1A1 Truncation by Asparagine Endopeptidase Defines Selective Neuronal Vulnerability in Parkinson's Disease. *Advanced Science*, *12*(2), 2409477. https://doi.org/10.1002/ADVS.202409477

Nishimura, K., Ásgrímsdóttir, E. S., Yang, S., & Arenas, E. (2023). A protocol for the differentiation of human embryonic stem cells into midbrain dopaminergic neurons. *STAR Protocols*, *4*(3), 102355. https://doi.org/10.1016/J.XPRO.2023.102355

Nishimura, K., Yang, S., Lee, K. W., Ásgrímsdóttir, E. S., Nikouei, K., Paslawski, W., Gnodde, S., Lyu, G., Hu, L., Saltó, C., Svenningsson, P., Hjerling-Leffler, J., Linnarsson, S., & Arenas, E. (2023). Single-cell transcriptomics reveals correct developmental dynamics and high-quality midbrain cell types by improved hESC differentiation. *Stem Cell Reports*, *18*(1), 337–353. https://doi.org/10.1016/J.STEMCR.2022.10.016

Nürnberg, E., Horschitz, S., Schloss, P., & Meyer-Lindenberg, A. (2018). Basal glucocorticoid receptor activation induces proliferation and inhibits neuronal differentiation of human induced pluripotent stem cell-derived neuronal precursor cells. *The Journal of Steroid*

Biochemistry and Molecular Biology, *182*, 119–126.
https://doi.org/10.1016/J.JSBMB.2018.04.017

Oosterveen, T., Garção, P., Moles-Garcia, E., Soleilhavoup, C., Travaglio, M., Sheraz, S., Peltrini,
R., Patrick, K., Labas, V., Combes-Soia, L., Marklund, U., Hohenstein, P., & Panman, L.
(2021). Pluripotent stem cell derived dopaminergic subpopulations model the selective
neuron degeneration in Parkinson's disease. *Stem Cell Reports*, *16*(11), 2718–2735.
https://doi.org/10.1016/j.stemcr.2021.09.014

Pantazis, C. B., Yang, A., Lara, E., McDonough, J. A., Blauwendraat, C., Peng, L., Oguro, H.,
Kanaujiya, J., Zou, J., Sebesta, D., Pratt, G., Cross, E., Blockwick, J., Buxton, P., Kinner-
Bibeau, L., Medura, C., Tompkins, C., Hughes, S., Santiana, M., … Merkle, F. T. (2022). A
reference human induced pluripotent stem cell line for large-scale collaborative studies.
Cell Stem Cell, *29*(12), 1685-1702.e22. https://doi.org/10.1016/J.STEM.2022.11.004

Piao, J., Zabierowski, S., Dubose, B. N., Hill, E. J., Navare, M., Claros, N., Rosen, S., Ramnarine,
K., Horn, C., Fredrickson, C., Wong, K., Safford, B., Kriks, S., El Maarouf, A., Rutishauser, U.,
Henchcliffe, C., Wang, Y., Riviere, I., Mann, S., … Tabar, V. (2021). Preclinical Efficacy and
Safety of a Human Embryonic Stem Cell-Derived Midbrain Dopamine Progenitor Product,
MSK-DA01. *Cell Stem Cell*, *28*(2), 217-229.e7. https://doi.org/10.1016/j.stem.2021.01.004

Pissadaki, E. K., & Bolam, J. P. (2013). The energy cost of action potential propagation in
dopamine neurons: clues to susceptibility in Parkinson's disease. *Frontiers in
Computational Neuroscience*, *7*(FEB). https://doi.org/10.3389/FNCOM.2013.00013

Poewe, W., Seppi, K., Tanner, C. M., Halliday, G. M., Brundin, P., Volkmann, J., Schrag, A. E., &
Lang, A. E. (2017). Parkinson disease. *Nature Reviews Disease Primers 2017 3:1*, *3*(1), 1–21.
https://doi.org/10.1038/nrdp.2017.13

Poulin, J. F., Caronia, G., Hofer, C., Cui, Q., Helm, B., Ramakrishnan, C., Chan, C. S., Dombeck,
D. A., Deisseroth, K., & Awatramani, R. (2018). Mapping projections of molecularly defined
dopamine neuron subtypes using intersectional genetic approaches. *Nature Neuroscience
2018 21:9*, *21*(9), 1260–1271. https://doi.org/10.1038/s41593-018-0203-4

Prakash, N., Brodski, C., Naserke, T., Puelles, E., Gogoi, R., Hall, A., Panhuysen, M., Echevarria,
D., Sussel, L., Vogt Weisenhorn, D. M., Martinez, S., Arenas, E., Simeone, A., & Wurst, W.
(2006). A Wnt1-regulated genetic network controls the identity and fate of midbrain-
dopaminergic progenitors in vivo. *Development*, *133*(1), 89–98.
https://doi.org/10.1242/DEV.02181

Puelles, E., Annino, A., Tuorto, F., Usiello, A., Acampora, D., Czerny, T., Brodski, C., Ang, S. L.,
Wurst, W., & Simeone, A. (2004). Otx2 regulates the extent, identity and fate of neuronal
progenitor domains in the ventral midbrain. *Development*, *131*(9), 2037–2048.
https://doi.org/10.1242/DEV.01107

Rekaik, H., Blaudin De Thé, F. X., Prochiantz, A., Fuchs, J., & Joshi, R. L. (2015). Dissecting the
role of Engrailed in adult dopaminergic neurons – Insights into Parkinson disease
pathogenesis. *FEBS Letters*, *589*(24PartA), 3786–3794.
https://doi.org/10.1016/J.FEBSLET.2015.10.002

Satpathy, A. T., Granja, J. M., Yost, K. E., Qi, Y., Meschi, F., McDermott, G. P., Olsen, B. N.,
Mumbach, M. R., Pierce, S. E., Corces, M. R., Shah, P., Bell, J. C., Jhutty, D., Nemec, C. M.,

Wang, J., Wang, L., Yin, Y., Giresi, P. G., Chang, A. L. S., ... Chang, H. Y. (2019). Massively
parallel single-cell chromatin landscapes of human immune cell development and
intratumoral T cell exhaustion. *Nature Biotechnology 2019 37:8*, *37*(8), 925–936.
https://doi.org/10.1038/s41587-019-0206-z

Sawamoto, N., Doi, D., Nakanishi, E., Sawamura, M., Kikuchi, T., Yamakado, H., Taruno, Y.,
Shima, A., Fushimi, Y., Okada, T., Kikuchi, T., Morizane, A., Hiramatsu, S., Anazawa, T.,
Shindo, T., Ueno, K., Morita, S., Arakawa, Y., Nakamoto, Y., ... Takahashi, J. (2025). Phase I/II
trial of iPS-cell-derived dopaminergic cells for Parkinson's disease. *Nature 2025 641:8064*,
641(8064), 971–977. https://doi.org/10.1038/s41586-025-08700-0

Schweitzer, J. S., Song, B., Herrington, T. M., Park, T.-Y., Lee, N., Ko, S., Jeon, J., Cha, Y., Kim, K.,
Li, Q., Henchcliffe, C., Kaplitt, M., Neff, C., Rapalino, O., Seo, H., Lee, I.-H., Kim, J., Kim, T.,
Petsko, G. A., ... Kim, K.-S. (2020). Personalized iPSC-Derived Dopamine Progenitor Cells
for Parkinson's Disease. *The New England Journal of Medicine*, *382*(20), 1926–1932.
https://doi.org/10.1056/NEJMoa1915872

Sharma, Y., Saha, S., Joseph, A., Krishnan, H., & Raghu, P. (2020). In vitro human stem cell
derived cultures to monitor calcium signaling in neuronal development and function.
Wellcome Open Research, *5*. https://doi.org/10.12688/WELLCOMEOPENRES.15626.1

Sif Ásgrímsdóttir, E., Fusar Bassini, L., Rivetti, P., Cervo, V., Gyllborg, D., Puigsasllosas Pastor,
C., Lee, K., Grigsby, C., Islam, S., Lönneberg, P., Villaescusa, C., Saltó, C., Linnarsson, S.,
Manno, G. La, Toledo, E. M., & Arenas, E. (2024). Midbrain Dopaminergic Neuron
Development is Regulated by Two Molecularly Distinct Subtypes of Radial Glia Cells.
BioRxiv, 2024.03.08.584031. https://doi.org/10.1101/2024.03.08.584031

Siletti, K., Hodge, R., Mossi Albiach, A., Lee, K. W., Ding, S.-L., Hu, L., Lönnerberg, P., Bakken, T.,
Casper, T., Clark, M., Dee, N., Gloe, J., Hirschstein, D., Shapovalova, N. V, Keene, C. D.,
Nyhus, J., Tung, H., Yanny, A. M., Arenas, E., ... Linnarsson, S. (2023). Transcriptomic
diversity of cell types across the adult human brain. *Science (New York, N.Y.)*, *382*(6667),
eadd7046. https://doi.org/10.1126/science.add7046

Simon, H. H., Saueressig, H., Wurst, W., Goulding, M. D., & O'Leary, D. D. M. (2001). Fate of
Midbrain Dopaminergic Neurons Controlled by the Engrailed Genes. *Journal of
Neuroscience*, *21*(9), 3126–3134. https://doi.org/10.1523/JNEUROSCI.21-09-03126.2001

Smidt, M. P., Smits, S. M., Bouwmeester, H., Hamers, F. P. T., van der Linden, A. J. A.,
Hellemons, A. J. C. G. M., Graw, J., & Burbach, J. P. H. (2004). Early developmental failure of
substantia nigra dopamine neurons in mice lacking the homeodomain gene Pitx3.
Development, *131*(5), 1145–1155. https://doi.org/10.1242/DEV.01022

Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W. M., Hao, Y.,
Stoeckius, M., Smibert, P., & Satija, R. (2019). Comprehensive Integration of Single-Cell
Data. *Cell*, *177*(7), 1888-1902.e21. https://doi.org/10.1016/j.cell.2019.05.031

Stuart, T., Srivastava, A., Madad, S., Lareau, C. A., & Satija, R. (2021). Single-cell chromatin state
analysis with Signac. *Nature Methods 2021 18:11*, *18*(11), 1333–1341.
https://doi.org/10.1038/s41592-021-01282-5

Tabar, V., Sarva, H., Lozano, A. M., Fasano, A., Kalia, S. K., Yu, K. K. H., Brennan, C., Ma, Y., Peng,
S., Eidelberg, D., Tomishima, M., Irion, S., Stemple, W., Abid, N., Lampron, A., Studer, L., &

Henchcliffe, C. (2025). Phase I trial of hES cell-derived dopaminergic neurons for Parkinson's disease. *Nature 2025 641:8064*, *641*(8064), 978–983. https://doi.org/10.1038/s41586-025-08845-y

Takahashi, J. (2020). iPS cell-based therapy for Parkinson's disease: A Kyoto trial. *Regenerative Therapy*, *13*, 18–22. https://doi.org/10.1016/J.RETH.2020.06.002

Tang, L., Xu, N., Huang, M., Yi, W., Sang, X., Shao, M., Li, Y., Hao, Z. zhe, Liu, R., Shen, Y., Yue, F., Liu, X., Xu, C., & Liu, S. (2023). A primate nigrostriatal atlas of neuronal vulnerability and resilience in a model of Parkinson's disease. *Nature Communications 2023 14:1*, *14*(1), 1–13. https://doi.org/10.1038/s41467-023-43213-2

Theofilopoulos, S., Wang, Y., Kitambi, S. S., Sacchetti, P., Sousa, K. M., Bodin, K., Kirk, J., Saltó, C., Gustafsson, M., Toledo, E. M., Karu, K., Gustafsson, J. Å., Steffensen, K. R., Ernfors, P., Sjövall, J., Griffiths, W. J., & Arenas, E. (2012). Brain endogenous liver X receptor ligands selectively promote midbrain neurogenesis. *Nature Chemical Biology 2012 9:2*, *9*(2), 126–133. https://doi.org/10.1038/nchembio.1156

Thibodeau, A., Eroglu, A., McGinnis, C. S., Lawlor, N., Nehar-Belaid, D., Kursawe, R., Marches, R., Conrad, D. N., Kuchel, G. A., Gartner, Z. J., Banchereau, J., Stitzel, M. L., Cicek, A. E., & Ucar, D. (2021). AMULET: a novel read count-based method for effective multiplet detection from single nucleus ATAC-seq data. *Genome Biology*, *22*(1), 252. https://doi.org/10.1186/s13059-021-02469-x

iklová, K., Björklund, Å. K., Lahti, L., Fiorenzano, A., Nolbrant, S., Gillberg, L., Volakakis, N., Yokota, C., Hilscher, M. M., Hauling, T., Holmström, F., Joodmardi, E., Nilsson, M., Parmar, M., & Perlmann, T. (2019). Single-cell RNA sequencing reveals midbrain dopamine neuron diversity emerging during mouse brain development. *Nature Communications 2019 10:1*, *10*(1), 1–12. https://doi.org/10.1038/s41467-019-08453-1

Tiklová, K., Nolbrant, S., Fiorenzano, A., Björklund, Å. K., Sharma, Y., Heuer, A., Gillberg, L., Hoban, D. B., Cardoso, T., Adler, A. F., Birtele, M., Lundén-Miguel, H., Volakakis, N., Kirkeby, A., Perlmann, T., & Parmar, M. (2020). Single cell transcriptomics identifies stem cell-derived graft composition in a model of Parkinson's disease. *Nature Communications*, *11*(1). https://doi.org/10.1038/S41467-020-16225-5

Toledo, E. M., Yang, S., Gyllborg, D., van Wijk, K. E., Sinha, I., Varas-Godoy, M., Grigsby, C. L., Lönnerberg, P., Islam, S., Steffensen, K. R., Linnarsson, S., & Arenas, E. (2020). Srebf1 Controls Midbrain Dopaminergic Neurogenesis. *Cell Reports*, *31*(5), 107601. https://doi.org/10.1016/j.celrep.2020.107601

Tolve, M., Ulusoy, A., Patikas, N., Islam, K. U. S., Bodea, G. O., Öztürk, E., Broske, B., Mentani, A., Wagener, A., van Loo, K. M. J., Britsch, S., Liu, P., Khaled, W. T., Metzakopian, E., Baader, S. L., Di Monte, D. A., & Blaess, S. (2021). The transcription factor BCL11A defines distinct subsets of midbrain dopaminergic neurons. *Cell Reports*, *36*(11), 109697. https://doi.org/10.1016/j.celrep.2021.109697

Trapnell, C., Cacchiarelli, D., Grimsby, J., Pokharel, P., Li, S., Morse, M., Lennon, N. J., Livak, K. J., Mikkelsen, T. S., & Rinn, J. L. (2014). The dynamics and regulators of cell fate decisions are revealed by pseudotemporal ordering of single cells. *Nature Biotechnology 2014 32:4*, *32*(4), 381–386. https://doi.org/10.1038/nbt.2859

Xi, J., Liu, Y., Liu, H., Chen, H., Emborg, M. E., & Zhang, S. C. (2012). Specification of Midbrain Dopamine Neurons from Primate Pluripotent Stem Cells. *Stem Cells*, *30*(8), 1655–1663. https://doi.org/10.1002/STEM.1152

Xu, C., Prete, M., Webb, S., Jardine, L., Stewart, B. J., Hoo, R., He, P., Meyer, K. B., & Teichmann, S. A. (2023). Automatic cell-type harmonization and integration across Human Cell Atlas datasets. *Cell*, *186*(26), 5876-5891.e20. https://doi.org/10.1016/j.cell.2023.11.026

Xu, P., He, H., Gao, Q., Zhou, Y., Wu, Z., Zhang, X., Sun, L., Hu, G., Guan, Q., You, Z., Zhang, X., Zheng, W., Xiong, M., & Chen, Y. (2022). Human midbrain dopaminergic neuronal differentiation markers predict cell therapy outcomes in a Parkinson's disease model. *The Journal of Clinical Investigation*, *132*(14). https://doi.org/10.1172/JCI156768

Yaghmaeian Salmani, B., Lahti, L., Gillberg, L., Jacobsen, J. K., Mantas, I., Svenningsson, P., & Perlmann, T. (2024). Transcriptomic atlas of midbrain dopamine neurons uncovers differential vulnerability in a Parkinsonism lesion model. *ELife*, *12*. https://doi.org/10.7554/ELIFE.89482

Zetterström, R. H., Solomin, L., Jansson, L., Hoffer, B. J., Olson, L., & Perlmann, T. (1997). Dopamine Neuron Agenesis in Nurr1 -Deficient Mice. *Science*, *276*(5310), 248–250. https://doi.org/10.1126/SCIENCE.276.5310.248

Zhang, D., Yang, S., Toledo, E. M., Gyllborg, D., Saltó, C., Carlos Villaescusa, J., & Arenas, E. (2017). Niche-derived laminin-511 promotes midbrain dopaminergic neuron survival and differentiation through YAP. *Science Signaling*, *10*(493). https://doi.org/10.1126/scisignal.aal4165

Zhang, Y., Liu, T., Meyer, C. A., Eeckhoute, J., Johnson, D. S., Bernstein, B. E., Nusbaum, C., Myers, R. M., Brown, M., Li, W., & Liu, X. S. (2008). Model-based analysis of ChIP-Seq (MACS). *Genome Biology*, *9*(9), R137. https://doi.org/10.1186/gb-2008-9-9-r137

Zheng, G. X. Y., Terry, J. M., Belgrader, P., Ryvkin, P., Bent, Z. W., Wilson, R., Ziraldo, S. B., Wheeler, T. D., McDermott, G. P., Zhu, J., Gregory, M. T., Shuga, J., Montesclaros, L., Underwood, J. G., Masquelier, D. A., Nishimura, S. Y., Schnall-Levin, M., Wyatt, P. W., Hindson, C. M., … Bielas, J. H. (2017). Massively parallel digital transcriptional profiling of single cells. *Nature Communications 2017 8:1*, *8*(1), 1–12. https://doi.org/10.1038/ncomms14049

Zhou, Y., Zhou, B., Pache, L., Chang, M., Khodabakhshi, A. H., Tanaseichuk, O., Benner, C., & Chanda, S. K. (2019). Metascape provides a biologist-oriented resource for the analysis of systems-level datasets. *Nature Communications*, *10*(1), 1523. https://doi.org/10.1038/S41467-019-09234-6

Ziegler, A. B., Thiele, C., Tenedini, F., Richard, M., Leyendecker, P., Hoermann, A., Soba, P., & Tavosanis, G. (2017). Cell-Autonomous Control of Neuronal Dendrite Expansion via the Fatty Acid Synthesis Regulator SREBP. *Cell Reports*, *21*(12), 3346–3353. https://doi.org/10.1016/j.celrep.2017.11.069