Identification of High Blanchability Donors, Candidate genes and Markers in Groundnut

Priya Shah[1,2], Sunil S. Gangurde[1], Ramachandran Senthil[1], Kuldeep Singh[1], Prashant Singham[2], Pasupuleti Janila[1], Sean Mayes[1], Manish K. Pandey[1]*

[1]Center of Excellence in Genomics & Systems Biology (CEGSB) and Center for Pre-Breeding Research (CPBR), International Crops Research Institute for the Semi-Arid Tropics (ICRISAT), Hyderabad-502324, Telangana, India

[2]Department of Genetics, Osmania University, Hyderabad-500007, Telangana, India

*Correspondence: manish.pandey@icrisat.org

Abstract

Blanchability in groundnut, the ability of seeds to shed their seed coat (testa), is a trait of economic importance in the food processing industry, yet remains underexplored in breeding programs. This study aimed to assess blanchability in 184 accessions from the ICRISAT minicore collection and identify associated genomic regions, candidate genes, and molecular markers. Significant variability was observed over two seasons, with blanchability ranging from 3.98% to 70.08%. Ten genotypes, including ICG10890, ICG9507, ICG13982, and ICG297, exhibited good blanchability, with ICG297 emerging as a promising donor based on cluster analysis of blanchability and agronomic traits. Genome-wide association studies (GWAS) using the 58K 'Axiom_*Arachis*' SNP array revealed 58 significant SNP-trait associations and important candidate genes such as *isocitrate dehydrogenase* and *ubiquitin ligase*, which influence seed coat structure and cell wall integrity. Nine SNPs were validated via allele mining, with four markers—on chromosomes A01 (snpAH00551), A06 (snpAH00554), B04 (snpAH00558), and B07 (snpAH00559), effectively distinguishing between high and low blanchability genotypes. These validated SNPs present valuable tools for genomics-assisted breeding. Overall, the findings is the first study contributing to a better understanding of the genomics and genetic basis of blanchability and offer resources for improving processing traits in groundnut.

Keywords: Blanchability, Candidate genes, Markers

Abbreviations: GWAS: Genome-wide association studies**,** STAs: SNP-trait associations**,** MUFA: Monounsaturated fatty acid**,** PR: post-rainy, R: rainy, PVE: Phenotypic variance explained, SNP: Single nucleotide polymorphism, PPP: Pods per plant **,** SDL: Seed length, SWT: Seed weight**,**

PDWT: Pod weight, PDWD: Pod width, PLN: Pod length, DM: Days to maturity, BLCB: Blanchability, AFTX: Aflatoxin, AFLV: *A. flavus* infection, ANOVA: Analysis of variance, QC: Quality Control, LD: Linkage disequilibrium , KASP: Kompetitive Allele-Specific PCR, MAF: Minor allele frequency, XETs: Xyloglucan endotransglucosylase, OST: Oligosaccharyltransferase, FADs: Fatty acid desaturases, GAB: Genomic-assisted breeding, Q-Q: Quantile-Quantile.

1. Introduction

Groundnut (*Arachis hypogaea* L.) also known as peanut, is a major leguminous crop, grown mostly as a food and feed crop globally [1] . The cultivated species of groundnut is a self-pollinated allotetraploid crop ($2n = 4x = 40$) with 2.54 Gb genome size [2–4] and belongs to the *Fabaceae* family [5, 6]. It is a key seasonal herbaceous legume crop grown in arid and semi-arid regions and is recognized as one of the important sources of edible oil and protein. Groundnut kernels contain 16% to 36% of protein, 36% to 54% oil and between 10% and 20% of total carbohydrates [7]. Groundnut has significant market value and enormous nutritional value by virtue of high mono-unsaturated fatty acid (MUFA) in oil content and possess significant amount of vitamins and minerals [8]. It also possesses several nutritional qualities such as soluble sugars and possesses rich amounts of calcium (Ca), iron (Fe), Vitamin B and E (tocopherol). Additionally, anti-nutrients like trypsin inhibitor and phytic acid are present in groundnut that may be deactivated by the process of boiling and roasting [9, 10]. It has secured 13[th] position among the most vital food crops and secured rank 4[th] in the list of the most vital oilseed crops[11].

According to industrial importance, edible food products such as raw or roasted nuts, refined oil, groundnut butter, salted groundnuts, groundnut flour and other confectionery products are processed groundnut products. During the processing of groundnut to prepare the edible products, an important operation is the separation of the testa or seed coat (skin) and the process is called blanching. The blanchability is the capacity of a genotype to release its testa very easily. This characteristic is of great economic value in the processing of groundnut raw and processed food products since it reduces the cost of food processing. Additionally, this trait also holds importance in terms of quality control as blanchability leads to removal of damaged kernels, including aflatoxin contamination. Moreover, in terms of processing applications,

globally, about 48% of groundnuts are used for the preparation of raw and processed food products

such as groundnut milk, groundnut butter, groundnut chikki, groundnut cake, salted and roasted

groundnut, groundnut bar chocolate, chutney, while oil extraction accounts for 52% usage,

however in India, the groundnut utility accounts for food, seed, and oil extraction is 44%, 24%

and 30% respectively [12, 13].

Blanchability has enormous economic significance in the production of food products made from

groundnut. If the groundnut variety has low blanchability, the processing of

the product becomes laborious, as it results in high re-processing and thus higher costs and time

to get marketable products. This trait is associated with the seed coat and its structural properties

as well as its chemical composition, such as lignin, pectin, polyphenols. Blanchability is reported

to be a trait with high heritability and genetically regulated, hence the breeding and selection using

conventional breeding could help in improving the groundnut cultivars with optimum blanchability

(~75%). In this context, GAB holds higher ability for accelerated improvement of crop varieties

as the conventional plant breeding is prolonged and laborious [14]. Though there are several

studies reported on germplasm screening for blanchability [14, 15], despite that only one study has

reported the genomic regions for blanchability using QTL-seq approach on a RIL population [16]

SNP-trait association mapping of groundnut diverse panels has speed up the genomic region

identification that are linked to agronomic traits by identifying ancestral or natural recombination

events that led to non-random allele association at various loci throughout the genome [17,

18]. Compared to   biparental   linkage   mapping,   association

mapping, respectively, allows greater mapping resolution [19]. With the availability of high-

density SNP (Single Nucleotide Polymorphism) array and WGRS genotyping data, there is much

scope to study the economically important and other traits in groundnut, identify STAs using the

sequencing based molecular markers, and deploying them in crop breeding programs

Association   mapping   has acquired considerable pace in   legumes, and there are

several studies indicating markers associated with nutritional   and   agronomic   traits   [20].

Considering the above knowledge gap and available information, the present study aimed to

identify STAs for economically important trait, blanchability, in the groundnut minicore set [21].

In this association mapping study, we have used three different algorithms, viz., Bayesian-

information and linkage-disequilibrium iteratively nested keyway (BLINK), Settlement of

Unmappable Positions via Extensive Rearrangement (SUPER), and Fixed and random model Circulating Probability Unification (FarmCPU). All these algorithms provide better statistical accuracy and computational efficiency and improve the accuracy of significant marker selection to be used in breeding programs.

Genome-wide association study (GWAS) has a potential to elucidate the genetic structure and the causative loci between different genotypes [22, 23]. This method has been widely utilized for the genetic dissection of complex traits in plants. As next-generation sequencing (NGS) technology has developed rapidly, high-density SNP can be invoked and additional GWAS analysis can be conducted. In the present study, a minicore collection comprising 184 diverse groundnut accessions were phenotyped for blanchability in two subsequent seasons (rainy (R) 2022 and post-rainy (PR) 2022-2023). We conducted GWAS for the identification of significant SNP markers related to blanchability in groundnut. Further, genes associated with blanchability were identified from the LD region of identified STAs. Also, the correlation and cluster analysis of blanchability with agronomic traits have revealed genotypes with superior traits. These results would provide a better and precise understanding of the blanchability trait with potential governing regions and its underlying molecular mechanism. Moreover, this study will further contribute to the improvisation of the blanchability of groundnut varieties with the help of marker-assisted breeding.

2. Material and Methods

2.1. Plant material and Phenotyping for blanchability

The groundnut minicore collection of 184 accessions developed at ICRISAT, Patancheru, was used in the experimental study [21]. Groundnut minicore collection encompasses six botanical types *viz.*, *hypogaea*, *hirsuta*, *fastigiata*, *peruviana*, *aequatoriana* and *vulgaris,* that collectively constitutes the genetic diversity present in the complete groundnut germplasm at ICRISAT **[Supplementary Figure S4 (A)].** The experimental design used here was the alpha-lattice design for a total of two seasons: rainy (R) season of 2022 and post-rainy (PR) 2022-2023. Phenotyping data generated in two replications during each season at ICRISAT, Patancheru, (17° 31' 48.00" N latitude and 78° 16' 12.00" E longitude).

Each accession was planted in a 4 m row plot with inter-plant spacing of 10 cm and between-row

spacing of 60 cm. The experimental site consists of red soils with

pH ranging from 6.0 to 6.3. In the rainy season of 2022, the highest temperature was upto

31.26°C whereas the range of minimum temperature was recorded to be 20.38°C, with 198.2 mm

rainfall, 154.24 mm of evaporation, and 87.60% accounts for relative humidity. But in the post-

rainy season of 2022–2023, the highest temperature was 33.25°C, whereas the

lowest temperature was recorded to be 18.61°C, with 65.5 mm rainfall, 98.75 mm rate of

evaporation, and 82.70% of relative humidity. Throughout both the growth seasons, standard

recommended agronomical practices essential for groundnut cultivation were followed for healthy

and good crop growth. Further, post-harvesting, the phenotypic observations for blanchability

were recorded for each replication at ICRISAT, Hyderabad, India **(Supplementary Table S1).**

Additionally, the previously documented phenotyping data for various agronomical traits such as

pods per plant (PPP), seed length (SDL) seed weight (SWT), days to maturity (DM), pod weight

(PDWT), pod width (PDWD), and pod length (PLN), was also used for correlation and cluster

analysis **(Supplementary Table S6 and S7).**

2.2. Phenotypic evaluation for blanchability

For phenotypic assessment of groundnut blanchability, first, the regular groundnut grading

was done where the variability in seed size observed from 7-8 mm diameter depending on the

genotype. The primary objective of grading was to reduce the seed size and

maturity variability. Subsequently, the pre-blanching weight of every genotype

was measured. Then the phenotyping of groundnut germplasm was done, for obtaining the

blanchability phenotypic data, seeds were subjected to hot air oven. Then the heated seeds were

cooled to room temperature, for more than 8 hrs. The samples were treated on a blancher for sixty

seconds per sample. According to the guidelines, seed sample weight of approximately 200 gm

and 50 gm was pre-heated at 110°C for 30-35 min, the pre-heating results in reduction of moisture

content to the range of 3.75-4.0 %. The samples were then allowed to cool down at room

temperature. For medium size and extra-large seeds, the blanching time was determined to be

180±25 sec and 240±25 sec, respectively, under air pressure at 121±0.5 kPa (17.6 ± 0.1 psi)

(Figure 1) [11]**.** In early generations, seed-set was low, thus a

phenotyping technique that is based on small sample size is more desirable since it

also offers a chance for high blanchability phenotypes in pedigree or in single-seed-descent

programs [14]. Hence, in this study, we have considered two different sample sizes, 50 gm and

200 gm, with a blanching duration of 60 sec and 120 sec, respectively, at 30.00 rpm. The weight

of the blanched seeds as well as blanched splits should be then observed and recorded **(Figure**

2A), followed by calculating the blanching percentage, using the formula below:

Blanching % = 100 × $\frac{\text{Blanched weight}}{\text{Pre}-\text{blanching weight}}$

2.3. Phenotypic data interpretation using statistical analysis

The analysis of variance (ANOVA) and descriptive statistics was performed over the rainy (R)

season of 2022 and post-rainy (PR) 2022-2023 using GenStat software version 15.0 (VSN

International Ltd., Hemel Hempstead, UK) **(Supplementary Table S2, S3 and S4).** The

phenotypic variation was observed in the minicore collection of groundnuts for blanchability trait

was analyzed. In descriptive statistics, the Shapiro-wilk test was performed to test if the data

follows normal distribution [24]. Analysis of differences between genotype replication through

one-way ANOVA across different seasons was evaluated using the F-test test (0.05 significance

level) **(Supplementary Table S2)**. Similarly, two-way ANOVA was used to study the genotype

and environment interaction **(Supplementary Table S4).** The 'GGplot' package of RStudio (R

Project for Statistical Computing) was used to analyze and plot the frequency curve on normal

histograms and to plot the boxplot for showing distribution of phenotypic data across the

agronomic types of groundnuts.

Pearson correlation analysis was conducted with customized RStudio script with 'Hmisc' package,

and the correlation heatmap was generated with 'pheatmap' package in RStudio. Principal

component analysis factor graph was employed to visualize the correlation among several traits,

was calculated with 'FactoMineR' package in RStudio **(Supplementary Table S5 and S6, Figure**

4). Minicore collection of 184 accessions was utilized for subsequent analyses. The frequency

distribution plots were generated using the "ggplot2" package in the RStudio. The objective was

to find a group of accessions to be utilized in breeding programs for good blanchability

without compromising other agronomic qualities screened under various environments.

2.4. Principal Component analysis of phenotypic data

Principal component analysis was performed to examine phenotypic variability in terms of agronomic types and blanchability, and the contribution of these traits to the observed variation, for PCA analysis for each of these traits the phenotypic means were used. Further, based on this result of PCA, the clustering was done for phenotypic characters. The PCA was performed using FactomineR package of RStudio **(Figure 3 and Supplementary Figure S2).**

2.5. Identification of accessions as potential donors for blanchability

The accessions promising for blanchability and other agronomic qualities along with yield advantage, noticed in phenotypic screening at various environments, were chosen, based on favorable alleles. We analyzed the blanchability data produced in the current study and agronomic trait datasets to identify accessions that could serve as donors in the breeding programmes targeting high-yielding varieties with satisfactory blanchability. Two strategies were employed to select desirable accessions from the minicore collection, for good blanchability and agronomic performance across varying environments.

In this approach, the phenotypic data for blanchability and other agronomic traits, days to maturity (DM), seed length (SDL) seed weight (SWT), pod weight (PDWT), pods per plant (PPP), pod width (PDWD), pod length (PLN), were utilized in the identification of superior accessions. The accessions utilized here were based on trait correlations and hierarchical clustering **(Supplementary Figure S5)**. Three distinct clusters were formed and further then from each cluster to enhance the groundnut genetic potential superior accessions were identified to be used as potential donors in breeding programs. The basis of this selection was the presence of a favorable combination of at least two or more traits **[Supplementary Table S7 (a), (b), and (c)].**

2.6. 58 K 'Axiom_Arachis' SNP Array: A tool for high density genotyping in groundnut

A high-density assay of 58 K 'Axiom_Arachis' SNP Array [25] that encompasses genotypic information of 184 groundnut minicore accessions was utilized to find the SNPs-traits association. Genomic DNA was extracted from tender leaves of 25-30 days old seedlings of groundnut minicore collection accessions using Nucleospin Plant II kit (Macherey-Nagel, Düren, Germany). For genotyping, 20 ng/µl of each sample's DNA was utilized with Affymetrix SNP array by the Affymetrix GeneTitan® system. Affymetrix GeneTitan® System was utilized as per protocol described by

Affymetrix (Axiom® 2.0 Assay) and derived data for each accession was created and stored in CEL file format [25]. Genotyping was performed using 'Axiom_*Arachis*' SNP array that consists of 58,233 SNP markers that were obtained from DNA re-sequencing of 41 wild diploid ancestors and tetraploid accessions of groundnut [25, 26]. The 58 K SNPs correspond to an average of ~2900 SNPs for each of 20 chromosomes.

Following SNP calling and data analysis conducted using Axiom™ Analysis Suite version 1.0 (Thermo Fisher Scientific, USA) to conduct quality control (QC) measures and choose samples that successfully cleared the QC test. SNPs were sorted into different classes followed by the "Axiom Best Practices Genotyping Workflow". Polymorphic markers "Poly High Resolution", and quality control passed samples were used in the study to assess the introgression lines with maximum recovery of the polymorphic SNPs. Monomorphic SNPs, low-intensity clusters, and SNPs having lower genotype call rates were removed while analysing the data.

TASSEL software was used to extract the polymorphic markers to further analyse the data. Of the 58,233 SNPs for 300 groundnut reference set were retrieved from Axiom™ analysis suit, the derived SNPs were further filtered for removing low quality and rare SNPs with ≤ 5% minor allele frequency (MAF) and >20% missing data using TASSEL (Trait Analysis by Association Evolution and Linkage) v5.2.93 software. After stringent filtration, high-quality SNPs across 167 accessions were subsequently employed for association mapping studies using GAPIT package in RStudio [27].

2.7. Population Structure analysis and genome-wide linkage disequilibrium

The population structure was estimated by PCA (principal component analysis), conducted using a genetic distance Roger's matrix, to identify clusters of individual belonging to similar genetic background, which was calculated using selection tool package in RStudio **(Figure 5)**. The squared Pearson's correlation coefficients (r^2) for the pairwise analysis of SNPs was calculated to evaluate genome wise linkage disequilibrium (LD) decay [28]. LD decay in the groundnut minicore collection panel was calculated using PopLDdecay v.3.29. and selection tool package in RStudio **(Supplementary Figure S4 (B))**.

2.8. Genome-wide association analysis for identifying SNPs-associated with blanchability

SNP-trait association was carried out utilizing advanced GAPIT models, namely, BLINK,
FarmCPU, and SUPER that employed both population structure (Q) and kinship (K)
matrices, utilizing GAPIT package of RStudio. Customize Quantile-Quantile (Q-Q) plots
were created by graphing observed negative Log10 'p' values against a computed expected
negative Log10 'p' value for all the SNPs available in CMplot package of RStudio [29].
A variation from 'p' values at the starting point reflects the population stratification present.
Manhattan plots were employed to represent chromosome-wise SNPs acquired through the
marker-trait association conducted throughout the genome. Log10 of 'p' values for every SNP
were graphed against twenty chromosomes for the blanchability trait with
the corresponding season and sample size. The false associations in GWAS
were adjusted by applying "Bonferroni Correction" (95% confidence interval).

The threshold of association to establish significant SNP-
trait association was calculated by applying Bonferroni correction for p-value of $4.9682 \times 10\text{-}6$,
through negative log transformation of α/n (where, α refers to the overall significance
level, n is the total number of SNPs employed for GWAS
analysis). Phenotypic observations were taken on four sets of data: 200 gm (post-rainy), 50 gm
(post-rainy), 50 gm (rainy), and 50 gm (average data of rainy and post-rainy).
The percentage of phenotypic variance explained by all significant
SNPs identified was reported from each model employed. The PVE% explained for each
significant SNP was determined by the squared correlation between the SNP's phenotype and
genotype.

According to the distribution of SNPs, the threshold level of significance of associations
between traits and SNPs was set at [–log 10 (p)< 10^{-05}] for GWAS analysis by using SNP
array that provided the best number of valid SNPs. SNP density plots, Q-Q plots, and Manhattan
plots were created by the CMplot package of RStudio **(Figure 6, Supplementary Figure S1, S6**
and Supplementary Figure S7) (Table 1).
In addition, as per the previously described methods by [30, 31], the impacts of the
alleles behind the significant stable SNP markers were investigated. Genotypes
were classified into independent groups based on their respective SNP alleles, and means were
compared by Turkey's HSD test **(Supplementary Figure S3).** In addition, the genes related to the

STAs regions were found using the physical locations of significant SNPs in the reference genome

sequence of Peanutbase (www.peanutbase.org). Gene function and effect of

the corresponding SNPs were identified from the annotation file.

2.9. Candidate gene discovery associated with superior blanchability

267 Peanut base (https://Peanutbase.org/) was employed to detect the candidate genes using GBrowse

268 (cultivated peanut) version 1. The SNP subsiding beginning and end position of the gene

269 were investigated for the discovery of the candidate gene based on their biological function

270 annotation towards the targeted trait **(Table 1).**

2.10. Development of Kompetitive allele specific polymerase chain reaction (KASP) -based markers

The present association mapping study identified 58 SNPs with a significant association, of which

10 highly significant markers were used for Kompetitive Allele-Specific PCR (KASP) assay

development **(Supplementary Figure S8).** These SNPs were selected from

genomic locations near candidate genes on three distinct chromosomes. The chosen SNPs

were redesigned into KASP markers using 50-bp upstream and

downstream regions to create user-friendly and economical markers [32]. Two allele-specific

forward primers and a generic reverse primer were designed for every SNP marker (Intertek Pvt.

Ltd.). The new KASP markers were validated across a panel of 25

genotypes, ranging from high blanchability and low blanchability genotypes as well

as ICRISAT breeding lines **(Figure 7).**

3. Results

3.1. Phenotypic analysis of blanchability

Blanchability is one of the important economic traits and phenotypic variation is an essential

component of GWAS analysis. The range and mean phenotypic distribution in 184 accessions

from the groundnut minicore collection across different seasons and sample size are presented in

Figure 2C. For season 2, traits exhibit a symmetric distribution, on the contrary, for season 1, with

sample size 50gm, the distribution is slightly skewed to the right, indicating more values

concentrated around 20 to 40 percent of blanchability, with fewer values beyond 60.

Supplementary Table S1 shows the wide variability range in the blanchability in the minicore collection. This includes blanchability_50R: 92.63 (mean 32.79), blanchability_50PR: 96.76 (mean 33.11) and blanchability_200R: 94.52 (mean 42.43). The trait and agronomic type wise phenotypic variation was depicted by box plot **(Figure 2B).** The agronomic type, spanish bunch and virginia runner, depicts the highest and lowest blanchability percentage; respectively. The blanchability trait was depicted across different seasons and sample sizes and it was observed that the variation has been profound in sample size 50 gm and rainy season.

Further, when the blanchability was correlated with the different agronomic types of groundnuts, Valencia Bunch, Virginia Bunch, Virginia Runner, and Spanish Bunch) across two seasons (Season 1: rainy and Season 2: post-rainy) and varying sample size (50 gm and 200 gm) **(Figure 2B).** The analysis determines that, in season 1 (rainy, 50 gm), spanish bunch has the highest blanchability, followed by valencia Bunch, while virginia runner shows the lowest. In season 2 (post-rainy, 50 gm), the general pattern remains, with spanish bunch still having the highest blanchability. However, with increased seed weight in season 2 (200 gm), blanchability increases overall, with the Spanish bunch maintaining its high blanchability and the other varieties showing a wider range of variation. The whiskers and outliers reflect variability within each variety **(Figure 2B).**

In the descriptive statistics analysis, it was observed that for rainy season (50 gm), values are generally lower and more around the mean, with a higher spread in the lower ranges, while post-rainy season (50 gm), the distribution is like the rainy season, though the mean is slightly higher. For post-rainy season (200 gm), values are significantly higher, with the mean around 42.57. Additionally, one-way and two-way ANOVA was performed using phenotypic data. In one-way ANOVA, it was observed that for the rainy season (50 gm), post-rainy season (50 gm), and post-rainy season (200 gm), the variation across genotypes was significant while the non-significant variation among replication suggests the accuracy of phenotypic data. In two-way ANOVA, the genotype and environment possess significant variation suggesting the effect of environment, while the genotype and sample size interaction is non-significant, that implies all genotypes respond similarly to sample size variations **(Supplementary Table S4).**

The PCA of the blanchability indicates that the PCs explained 93.3% of the total phenotypic variance **(Figure 3)**. The basis of clustering for accessions was not made on their agronomic type

but had found to be varying degrees of relatedness, with Spanish bunch and Valencia bunch showing distinct separation, while Virginia bunch and Virginia runner overlap, indicating similarities. Blanchability_50PR and 200PR, were inversely related to the blanchability_50R. In addition, the PCA analysis of the blanchability and the agronomic traits showed that the first two PCs explained 62.3% of the total phenotypic variance. The analysis has found to highlight an interesting trend for the blanchability and the agronomic traits. AFTX_PR negatively related with PPP_PR, as did BLCB (200PR, 50PR, 50R) with DM (R and PR), and SWT have been found to highly significantly negatively correlated with BLCB (200PR, 50PR). In contrast, AFTX and AFLV are positively correlated with BLCB (200PR, 50PR, 50R) as did all other agronomic trait with BLCB (200PR, 50PR, 50R), however with AFTX and AFLV the relatedness is not significant **(Supplementary Figure S2)**.

These insights would find to be useful for targeting superior accessions in breeding programs. Pearson correlation analysis of blanchability, aflatoxin and agronomic traits were evaluated across diverse environments post-rainy (PR) and rainy (R). The four traits that include pods per plant (PPP), blanchability (BLCB), aflatoxin (AFTX), *A. flavus* infection (AFLV), days to maturity (DM), pod length (PLN), seed length (SDL), seed weight (SWT), seed width (SWD), and pod width (PDWD) were evaluated in rainy and post-rainy environment. The color reflect the strength of correlation and further cluster has been made. PPP (R) positively correlated with blanchability and hence clustered together and other agronomic traits and aflatoxin make separate clusters **(Supplementary Table S5 and S6) (Figure 4)**.

3.2. Potential donors for blanchability

Promising groundnut accessions with high productivity and blanchability can serve as donors for the breeding programs to be carried out to develop better varieties for fulfilling global demand. Employing information on blanchability and agronomic traits, for selecting better donors with favorable values of desirable traits. **Figure 4** indicates the correlation between blanchability and agronomic traits, where negative correlation exists for certain traits (days to maturity, seed weight). On the contrary, PPP (R) positively correlated with blanchability and hence clustered together and other agronomic traits and aflatoxin make separate clusters.

In identifying the possible donors in this strategy, we considered the correlations among the blanchability and agronomic characteristics, from various environments.

All the characteristics describe nearly 62.3% of total variation. The PCA factor plot revealed that PC1 and PC2 explained 46.3% and 16% of the variation in measured characteristics, respectively **(Supplementary Figure S2).** Here, seed width (SWD_PR, SWD_R) and pod width (PDWD_PR, PDWD_R) were closely related to pods per plant (PPP_R) and pod length (PLN_PR, PLN_R) and distantly related to days to maturity (DM_R, DM_PR), similarly, blanchability (BLCB_50R, BLCB_50PR, BLCB_200PR) and seed weight (SWT_PR, SWT_R) are negatively correlated.

Hierarchical cluster analysis has made the classification of the traits into three distinct clusters: (1) days to maturity (DM_R, DM_PR); (2) seed width (SWD_PR, SWD_R), pod width (PDWD_PR, PDWD_R), pods per plant (PPP_R), seed length (SDL_PR, SDL_R); and (3) blanchability (BLCB_50R, BLCB_50PR, BLCB_200PR), seed weight (SWT_PR, SWT_R), pod length (PLN_PR, PLN_R), pods per plant (PPP_PR) **(Supplementary Figure S5).** The top accessions were selected based on phenotypic performance for the blanchability and agronomic traits. Accessions grouped in the same cluster were allowed to be compared among themselves, to identifying accessions for favorable combinations of two or more traits. For instance, cluster 1 had 21 accessions with higher days to maturity (**Supplementary Table 7 (a)**), twenty-seven accessions with cluster 2 had eight accessions that had higher SWD_PR, SWD_R, PDWD_PR, PDWD_R **(Supplementary Table 7 (b))**. In cluster 3, eight accessions had higher BLCB_50R, BLCB_50PR, BLCB_200PR, SWT_PR, SWT_R, PLN_PR, PLN_R, PPP_PR, and **[Supplementary Table 7 (c)]**.

Selected intercrossing allows the development of improved accessions that have found to harbor beneficial alleles for blanchability and agronomical traits in groundnut. For example, ICG297 (cluster 1) could be crossed with ICG15419, ICG11219, ICG332, ICG6022, ICG9809, ICG 4729 and ICG11687 (cluster 3) to breed long-seeded groundnut varieties with good blanchability and seed weight. ICG6766 (cluster 2) could be crossed with ICG332 (cluster 1) to breed more days to maturity, seed width, pod width sand pod length varieties. We also made the comparison between the accessions identified in different cluster to identify common accessions between cluster 2 and 3, eleven accessions (ICG15419, ICG6022, ICG 11219, ICG4538, ICG11862, ICG6766, ICG9777, ICG4746, ICG2381, ICG8760 and ICG297), and accessions (ICG332) and (ICG 14630) common between cluster 1 and 3, and cluster 1 and 2, respectively. One accession (ICG297) was identified to be common between all the three clusters, cluster 1, 2 and 3, that could be used in

381 breeding programs as potential donors to enhance both, groundnut blanchability and agronomic
382 traits.

3.3. Genotyping and SNP density

The 58 K 'Axiom_*Arachis*' SNP array was used for the genotyping of minicore collection. After
stringent filtration a total of 5044 high-quality SNPs across 167 accessions out of 184 as for some
genotypes data was missing, this data was subsequently utilized for GWAS using GAPIT package
of RStudio. For SNP array, SNP density across chromosomes, segmented into 1Mb windows,
certain chromosomes, such as Chr A02, Chr A03, Chr A04, and Chr B03, show the highest overall
SNP counts, with totals of 323, 293, 296 and 291 SNPs, respectively. In contrast, chromosomes
Chr A10 and Chr B08 have relatively lower SNP counts; 198 and 206; respectively.
(**Supplementary Figure S1**)

3.4. Population structure analysis and genome-wide linkage disequilibrium

Heatmaps and dendrograms of the kinship matrix indicated that there was clear clustering among
the genotypes. The basis of this clustering was observed to be based on polymorphic SNPs for the
studied genotypes, in the SNP array (**Figure 5) [Supplementary Figure S4 (A) and (B)]**
(Supplementary Table S8). For SNP array, the population structure based on groundnut sub-
species, botanical variety and agronomic type also revealed four distinct clusters: cluster 1 (46
genotypes): *fastigata*, vulgaris, spanish bunch; cluster 2 (75 genotypes): hypogaea, hypogaea,
virginia bunch; cluster 3 (27 genotypes): fastigata, fastigata, valencia bunch; cluster 4 (19
genotypes): fastigata, fastigata, vulgaris, spanish bunch, respectively **(Supplementary Table S8).**
The graphical representation of the LD characteristics of the minicore collection is presented for
SNP array data in **Supplementary Figure S4 (B).** In the case of SNP array data, the mean r^2
value for the genome was 0.12 and the LD decay was observed to start at an r^2 value of 0.95
and half-decay at 0.15. The curve of LD decay cut off at 1Mbp which is the genome-wide critical
distance to observe linkage. Therefore, markers linked to the same trait within this distance
were classified as linked or associated.

3.5. Identifying SNPs and its associated candidate genes for blanchability

GWAS analysis was conducted based on 58,233 high-density SNPs having less than 20% missing data. They were scattered among the twenty groundnut chromosomes. SNP genotyping data of 58,233 SNPs together with population structure information and kinship matrix were utilized for conducting genome-wide association analysis for blanchability in the 2022 rainy season and 2022-2023 post-rainy seasons. The threshold level of -log10 p value was fixed at 5.0 for SNP array analysis above which the SNPs are reported to be significantly associated **(Figure 6)**.

A total of 58 highly significant STAs for blanchability were identified. Of the 58 STAs, 21 were identified for 200PR with -log10 p values ranging from 1.61×10^{-8} to 6.28×10^{-6} which explained 0 to 35.28% of phenotypic variation (PVE), and 28 were identified from 50PR_R mean data set, with -log10 p values ranging from 5.45×10^{-13} to 9.36×10^{-6} which explained 0 to 39.03% of PVE, further from 50R, 4 STAs were identified with -log10 'p' values ranging from 1.18×10^{-8} to 1.18×10^{-6}, and PVE range from 0 to 45.93% while for 50PR, 7 STAs were identified with -log10 'p' values ranging from 3.79×10^{-7} to 8.80×10^{-6} and PVE from 0 to 32.24%. Finally, the 9 markers that were identified for blanchability from the analysis with the 4 different data set, and these were distributed across chromosome A01 (1), A05 (2), A06 (1), A09 (2), B04 (2), and B07 (1). Among these markers, B07_2964357 (AX_147254562) was observed to possess the highest value for phenotypic variation of 35.28% with a -log10 p value of 3.69×10^{-6} **(Table 1) (Figure 6).**

Groundnut genome sequencing accounted for genome size of 1.2 Gb in *A. duranensis* (36,734 genes) and for *A. ipaensis* it was 1.5 Gb (41,840 genes) and demystified the role of several genes for blanchability. Physical locations of each SNP marker of the current study were compared with the groundnut genome sequence to identify the gene function that underlies the corresponding SNP using Peanut base (https://Peanutbase.org/). Based on the previously reported genes documented in the literature, candidate genes which were previously reported to control seed coat integrity and adhesion were subsequently found to be candidate genes controlling blanchability. A total number of 13 SNPs were identified to possess candidate genes that were associated with blanchability **(Table 1)** viz., *Oligosaccharyl transferase STT3 subunit homolog (Aradu.B0E28), NRAMP metal ion transporter 6 (Araip.V0BSS); Transcription factor jumonji (jmjC) domain-containing protein (Aradu.N9ZS9); 5-methylthioadenosine / S-adenosylhomocysteine deaminase (Aradu.YDH5Z),*

fatty acid desaturase 2 (Aradu.G1YNF), ethylene-responsive transcription factor 7 (Araip.LL89K), isocitrate dehydrogenase (Araip.BZ4NA); 4-hydroxy-3-methylbut-2-enyl diphosphate synthase (Araip.2B9XL); xyloglucan endotransglucosylase / hydrolase 5 (Araip.2H071); putative membrane-bound O-acyltransferase (Araip.Q36GV), ubiquitin ligase SINAT3 (Aradu.G23Y9); N-terminal nucleophile aminohydrolases (Ntn hydrolases) superfamily protein (Araip.PG50R); charged multivesicular body protein (Araip.1D5HT) were found associated with SNPs identified.

3.6. Allelic effects of stable STAs and validation of blanchability SNPs

A total of 9 significant STAs were observed to be polymorphic for the blanchability identified with SNP array analysis (**Supplementary Figure S8**). All the nine SNP markers A01_230616 (50PR_R), A05_71291610 (50PR_R,200PR), A05_85198860 (50PR_R), A06_57729112 (50PR_R,200PR), A09_114692575 (50PR_R,50PR), A09_115432582 (50PR_R,200PR), B04_105437967 (50PR_R,200PR,50PR), B04_108209828 (50PR_R,50PR), and B07_2964357 (50PR_R,200PR), showed stable STAs with blanchability traits were further allowed to be used for determining the individual allelic effects on the studied traits**.** The identified alleles of these nine SNP markers were observed to have substantial effects on blanchability (**Supplementary Figure S3**).

3.7. KASP markers development and validation for blanchability

KASP markers epitomized low-cost and less tedious genotyping assays to conduct early generation selection among breeding lines to indirectly select a specific phenotype [33]. For the development and validation of the diagnostic markers for blanchability, out of nine SNPs, six SNPs were located on A05, A09 and B04 with two SNPs on each chromosome, and three SNPs with one each on A01, A06, B07. These SNPS were substantially targeted for KASP markers development, Primers were then successfully designed and developed for 9 SNPs; A01_230616 (TT/CC), A05_71291610 (CC/TT), A05_85198860 (TT/GG), A06_57729112 (TT/CC), A09_115432582 (GG/AA), B04_105437967 (CC/TT), A09_114692575 (GG/AA), B04_108209828 (GG/AA) and B07_2964357 (CC/TT) and validated on a validation panel with contrasting blanchability. The validation panel comprised of good and poor blanchability genotypes and breeding lines ranging from 6% to 80%. Of the nine KASP markers selected for validation, four KASPs (snpAH00551, snpAH00554, snpAH00558 and snpAH00559) showed polymorphism in the designed validation

panel. Interestingly, these four KASP markers were polymorphic and were clearly differentiating
between genotypes that possess good and poor blanchability. These KASP markers were localized
on genomic region detected on chromosome A01, A06, B04 and B07 **(Figure 7)**. The validated
highly polymorphic KASP markers for good and poor blanchability genotypes and can be used as
potential diagnostic markers for very early-stage selection of blanchability breeding material at the
varietal development process.

4. Discussion

Groundnut is a globally cultivated food legume, celebrated for its high protein and unsaturated oil
content. Blanchability quality becomes a significant trait regarding the development and utility of
groundnut breeding lines for edible consumption [34]. In the research conducted so far for the
blanchability trait, runner type groundnut has been utilized much more than other types and several
laboratory-scale blanchers were customized and designed to assist breeding programs for the
identification the high blanchability genotypes [35–38] . In the present study, a minicore collection
panel was analysed using 58 K 'Axiom_*Arachis*' array and multi-environment phenotyping for
the identification of the genomic regions and candidate genes that were associated with
blanchability. This approach aims to enhance the understanding of the genetic factors influencing
this economically important trait in groundnut cultivation.

In addition, some accessions identified through cluster analysis that can serve as potential donors
and would further facilitate for the development of accessions with improved traits. The accessions
would be harboring beneficial alleles governing blanchability along with agronomical traits in
groundnut. Accession ICG297 (cluster 1) could be crossed with ICG15419, ICG11219, ICG332,
ICG6022, ICG9809, ICG 4729 and ICG11687 (cluster 3) to breed long-seeded groundnut varieties
with high blanchability and seed weight. ICG6766 (cluster 2) could be crossed with ICG332
(cluster 1) to breed high days to maturity, seed width, pod width and pod length varieties.
Accessions (ICG297) between all three clusters, cluster 1, 2 and 3, combine blanchability and
agronomic traits, that could be effectively used in breeding programs as potential donors.
Therefore, these assays could significantly enhance future molecular breeding initiatives,
providing a targeted and efficient approach to selecting lines with desired traits.

The cultivated agronomic type of groundnut includes, 'Spanish Bunch', 'Valencia Bunch',
'Virginia Runner', and 'Virginia Bunch'. These botanical types have been found to possess vary

distinct phenotypic characters, including branching habits, seed, and pod size which is according
to the suitability to the specific cultivable environments. Spanish Bunch has been observed to
possess good blanchability, while the maximum of virginia runner were with poor blanchability.
So as per the industrial demand to produce groundnut processed food such as peanut butter,
cookies, chocolate industries can prefer spanish bunch agronomic type (good blanchability), while
for beer nuts and other seed based confectionary products virginia runner can be considered (Poor
blanchability).

Genomics assistedbreeding has been an emerging technology for improvisation of several
significantly important traits such as disease resistant, improving nutritional content, abiotic stress
resistance in groundnut [39–41]. GWAS have become the primary method for identifying STAs
related to complex traits of interest. Utilizing advanced models like FarmCPU, BLINK and
SUPER in GWAS significantly enhances the reliability and precision of identifying genetic
associations in plant genomics [42–44]. Resequencing-based genotyping reveals extensive natural
variations, allowing for the exploration of functional genes and elite alleles within natural
populations through GWAS. This approach enables us to identify valuable genetic traits that can
enhance breeding programs and improve crop performance [45, 46]. In this study, we identified a
total of 58 significant STAs for the blanchability trait using 58 K 'Axiom_*Arachis*' array; through
GWAS using a minicore collection panel. Among which four SNP markers were validated, that
were distinguishing the good (50-80%) and poor blanchability lines (1-30%), snpAH00551 (A01),
snpAH00554 (A06), snpAH00558 (B04) and snpAH00559 (B07).

These results are supported by the previous study reporting the same chromosome to be associated
with blanchability, however the genomic region varies [16]. The marker found in previous studies
on A06 were located at *Arahy.06_108,665,514, Arahy.06 _108,812,907* which is located at a
distant from the polymorphic marker, *Aradu.06_57,729,112*; we have found and validated in our
study. These findings suggest that chromosome 6 (A06) comprises of an important region to be
considered for blanchability. Additionally, the markers from chromosome A06 have also been
validated by the Peanut Company of Australia Pty Ltd, A Bega Company, Australia.

Groundnut blanchability, is a trait influenced by genetic and environmental factors. Studies have
associated blanchability with specific genomic regions or quantitative trait loci (QTLs). Previous
research suggests that QTLs linked to groundnut blanchability are often identified on chromosome

A06 and B01, these associations are determined through performing QTL-seq with two bulks of

breeding lines from three populations. In addition, the KASP markers were developed from the

most important SNPs from the QTLs on A06 *(Arahy.06_108,665,514, Arahy.06 _108,812,907)*

and B01 (Arahy.11_15,264,657 Arahy.11_16,329,544, Arahy.11_18,994,278) [16]. The linkage-

drag that's reported to be associated with two prominent *A. cardenasii* introgressions (A02 and

A03) have been conferred for disease resistance and negatively affects blanchability. As a result,

when breeding for disease resistance using wild germplasm, there is a risk of co-introduction of

undesirable traits such as reduced blanchability due to their close genetic linkage [16]. However,

if there is a requirement of skin for processing application, then this linkage-drag could be proven

to be beneficial such as in case of beer nut and other seed based confectionary products. As it

would enable to select disease resistant and blanching resistant varieties.

Blanchability was neglected until now, however   efforts   have been in progress to identify the

SNP-trait   associations   (STAs) associated with the high   blanchability in the panel

for diversity of characteristics such as blanchability of groundnut with   an

aim to develop parental varieties of groundnuts with enhanced blanchability in order to serve as

donors of breeding programs groundnut improvement. The SNP allele analysis of the diverse

minicore collection accessions for the identified significant STAs indicated that accessions

carrying all favorable blanchability alleles tend to exhibit good blanchability. The validation panel

for blanchability includes extreme genotypes representing good blanchability and poor

blanchability genotypes, along with breeding lines. Those STAs which distinguish clearly good

and poor blanchability lines hold promise for developing molecular markers, and accessions with

all favorable good blanchability alleles can serve as donors in MABC programs aimed at achieving

good blanchability lines. Among these, four markers snpAH00551 (A01), snpAH00554 (A06),

snpAH00558 (B04) and snpAH00559 (B07) were successfully validated across minicore

collection and groundnut breeding lines.

Blanchability in groundnuts is more commonly linked to factors such as, water absorption and

seed coat integrity and adhesion, moisture content, and cell wall structure and biochemical

components like pectin and lignin. Moreover, there are studies identifying other genes related to

seed coat properties and adhesion, which can influence blanchability. The cell wall composition,

pectin, cellulose, lignin, and hemicellulose are key components, while adhesion molecules include

proteins involved in cell wall modification or strengthening. Seed coat adhesion and seed coat integrity are key factors influencing traits like blanchability, how tightly the seed coat adheres to the cotyledons, which are influenced by both mechanical properties and biochemical interactions.

Most of the genes found have expressions that are distant from the pathways typically involved in the synthesis and remodeling of cell wall components, which are more directly responsible for seed coat properties. The SNPs, snpAH00551 (A01), snpAH00554 (A06), snpAH00558 (B04) and snpAH00559 (B07) that were found to be polymorphic in the validation analysis are found to be associated with genes *ubiquitin protein- ligase (Aradu.G23Y9)* (A06) and *isocitrate dehydrogenase (Araip.BZ4NA)* (B04). These genes have found to have effect on cell wall dynamics influences the ease of seed coat removal, a key factor in groundnut blanchability. *Isocitrate dehydrogenase* produces α-ketoglutarate (α-KG), which regulates enzymes involved in cell wall loosening (e.g., *pectin methylesterases*, *expansins*). Changes in seedcoat integrity affect adhesion property, which is a determinant of blanchability [47].

Ubiquitin protein-ligase and ubiquitin ligase SINAT3 (a RING-type ubiquitin ligase) regulates protein degradation via the ubiquitin-proteasome system (UPS), affecting stress response, hormonal signalling, and cell wall modifications. SINAT3 is involved in abiotic stress response pathways, including oxidative stress and hormone signalling (e.g., auxin, ethylene), which regulate seed coat integrity and blanchability by influencing pectin breakdown and adhesion strength [48]. It likely interacts with transcription factors (e.g., NAC, MYB) involved in cell wall degradation, modulating lignin biosynthesis and pectin remodelling [49].

The interplay between oxidative metabolism (*Isocitrate dehydrogenase*) and protein turnover (*Ubiquitin protein-ligase* and *Ubiquitin ligase SINAT3*) determines seed coat adhesion strength, thereby influencing blanchability. *Isocitrate dehydrogenase* contributes to blanchability by regulating α-KG-dependent enzymes involved in seed coat loosening. *Ubiquitin protein-ligase* and *Ubiquitin ligase SINAT3* impacts blanchability by targeting proteins that control seed coat adhesion and degradation. This pathway can be further validated by transcriptomic and proteomic analysis of high- and low-blanchability groundnut genotypes to assess differential expression of *ubiquitin protein-ligase/ubiquitin ligase SINAT3*, *isocitrate dehydrogenase* and cell-wall-modifying enzymes.

If *isocitrate dehydrogenase* activity is high, lignin/pectin biosynthesis is enhanced, leading to a
stronger seed coat and lower blanchability. If SINAT3 degrades key regulators of cell wall
biosynthesis, this can lead to a weaker seed coat and higher blanchability. *Isocitrate*
dehydrogenase affects blanchability via metabolic flux regulation, providing intermediates for
seed coat biosynthesis. SINAT3 influences blanchability through the ubiquitin-proteasome
system, modulating stress responses and cell wall composition. Their interaction is likely mediated
by oxidative stress pathways, hormonal regulation, and protein degradation, ultimately impacting
seed coat adhesion and blanchability in groundnuts.

Furthermore, we have determined genes that are associated to cell wall biosynthesis,
modifications, remodeling and adhesion proteins such as *xyloglucan endotransglucosylase*, and
oligosaccharyl transferase STT3, putative membrane-bound O-acyltransferase that influence seed
coat's biochemical properties. For instance, x*yloglucan endotransglucosylase (XETs),* participated
in cell wall modification and maintained cell wall plasticity, helping plant in tolerating stress such
as salinity stress [50]**.**

XETs are enzymes that modify xyloglucans, major hemicellulosic components of the plant cell
wall. They play a pivotal role in restructuring the cell wall by cleaving and rejoining xyloglucan
chains, facilitating cell wall loosening and expansion. This activity is crucial for maintaining cell
wall integrity and modulating adhesion between cells, directly impacting plant growth and
development. Their function is vital during seed coat formation, where precise cell wall
modifications are necessary for proper development [51].

The roles of *STT3* are a component of the *oligosaccharyltransferase (OST)* complex, which is
involved in N-linked glycosylation, and glycoproteins in the cell wall or membrane could influence
cell wall remodeling or stability and *stt3a* mutation disturbs the essential adaptation mechanisms
[52]. *STT3* is a catalytic subunit of the *OST* complex, responsible for transferring oligosaccharides
to nascent proteins during N-linked glycosylation. Proper glycosylation is essential for the stability
and function of many cell wall proteins, thereby influencing cell wall integrity and cell-cell
adhesion.

The structure of the seed coat is tightly regulated during seed maturation, and fatty acids,
synthesized and modified by *Fatty acid desaturases (FADs) FADs*, are part of the cuticle (a waxy
protective layer) that forms on the outer surface. In seed coats, the fatty acid profile affects the

permeability and protective functions of the seed coat, thereby influencing seed development and

viability [53]. *FADs* play a vital role for determining the lipid composition of the seed coat,

affecting its physical properties and functions, including protection, permeability, and regulation

of dormancy and germination.

All these enzymes are integral to the dynamic processes of cell wall construction, modification,

and maintenance, directly affecting seed coat development, cell wall integrity, and adhesion in

plants. By modifying interactions within the cell wall, these genes may influence how tightly the

seed coat adheres to the cotyledons. Strong adhesion can make blanching more difficult, while

effective modification might facilitate easier removal of the seed coat.

626 **5. Conclusion**

Blanchability is a trait of tremendous economic importance as there is a significant energy, time

and cost requirement to remove the groundnut seedcoats from poor blanching lines varieties.

Product processing ultimately becomes cumbersome leading to re-processing. This ultimately

leads to considerable loss of kernels during the processing and sorting of the groundnut. The

present study utilized 58 K 'Axiom_*Arachis*' array genotypic data from a diverse minicore

collection panel phenotypic data collected over two seasons, to conduct a genome-wide association

study. This phenotypic analysis reveals that the spanish bunch agronomic types are possessed to

have good blanchability, additionally, the statistical analysis has revealed the effect of sample size

and environment on blanchability trait. ICG297 genotype has been identified to be the superior

potential donor for blanchability along with other agronomic traits. Further, the GWAS analysis

has identified significant STAs in 58 K 'Axiom_*Arachis*' array genotypic data for blanchability.

The analysis of these STAs revealed potential candidate genes such as *isocitrate dehydrogenase*

and *ubiquitin ligase protein* associated with blanchability, highlighting the critical role of these

genes in the dynamic processes of cell wall construction, modification, and maintenance and have

a direct impact on the development of seed coats, the integrity of cell walls, and plant adhesion.

Ultimately, four KASP markers snpAH00551 (A01), snpAH00554 (A06), snpAH00558 (B04) and

snpAH00559 (B07) were validated from 58 K 'Axiom_*Arachis*' array exhibiting clear

polymorphism between good and poor blanchability genotypes were identified, to be incorporated

in breeding programs. Among these A06 chromosome, has been found common with previous

study which suggests it could prove to be the future potential marker for blanchability.

Furthermore, conducting haplotype analysis on these candidate genes could facilitate the
identification of superior haplotypes for blanchability, which can be leveraged in haplotype-based
breeding strategies.

Data availability statement:

The phenotypic data used in this work is provided in Supplementary Table S1. The sequencing
data generated in this study is deposited in NCBI with bio-project ID PRJNA1002116.

Institutional Review Board Statement:

Not Applicable

Informed Consent Statement:

Not Applicable

CRediT authorship Contribution Statement:

Manish K. Pandey: Conceived the idea and supervised and finalized the manuscript. **Kuldeep**
Singh and Ramachandran Senthil: Contributed seed material and in seed multiplication of the
minicore collection. **Pasupuleti Janila:** Contributed the breeding lines for marker validation.
Priya Shah: Phenotyped minicore collection, performed the analysis and wrote the manuscript.
Sunil S. Gangurde, Prashant Singham and Sean Mayes: Reviewed, edited and improvised the
manuscript. All authors have read and agreed to the published version of the manuscript.

Funding:

The authors are thankful to the Indian Council of Agricultural Research (ICAR) through ICAR-
ICRISAT collaborative project, MARS Inc. USA, and Bill & Melinda Gates Foundation (BMGF),
USA through Tropical Legumes III project.

Acknowledgments:

Priya Shah acknowledges Joint Council of Scientific and Industrial Research-University Grant
Commission (CSIR-UGC), Government of India for the award of fellowship for a Ph.D. and
ICRISAT for providing research facilities. The authors are thankful to Khaja D. Mohinuddin, for
his assistance in improving manuscript figures.

Conflicts of Interest:

The authors declare there is no conflict of interest.

References:

[1] K.A. Bediako, K. Ofori, S.K. Offei, D. Dzidzienyo, J.Y. Asibuo, R.A. Amoah, Aflatoxin
contamination of groundnut (*Arachis hypogaea* L.): Predisposing factors and management
interventions, Food Control 98 (2019) 61–67.

[2] D.J. Bertioli, J. Jenkins, J. Clevenger, O. Dudchenko, D. Gao, G. Seijo, S. Leal-Bertioli, L.
Ren, A.D. Farmer, M.K. Pandey, S.S. Samoluk, B. Abernathy, G. Agarwal, C. Ballén-
Taborda, C. Cameron, J. Campbell, C. Chavarro, A. Chitikineni, Y. Chu, S. Dash, M. El
Baidouri, B. Guo, W. Huang, K.D. Kim, W. Korani, S. Lanciano, C.G. Lui, M. Mirouze,
M.C. Moretzsohn, M. Pham, J.H. Shin, K. Shirasawa, S. Sinharoy, A. Sreedasyam, N.T.
Weeks, X. Zhang, Z. Zheng, Z. Sun, L. Froenicke, E.L. Aiden, R. Michelmore, R.K.
Varshney, C.C. Holbrook, E.K.S. Cannon, B.E. Scheffler, J. Grimwood, P. Ozias-Akins,
S.B. Cannon, S.A. Jackson, & J. Schmutz. The genome sequence of segmental
allotetraploid peanut *Arachis hypogaea*, Nat Genet 51 (2019) 877–884.

[3] X. Chen, Q. Lu, H. Liu, J. Zhang, Y. Hong, H. Lan, H. Li, J. Wang, H. Liu, S. Li, M.K.
Pandey, Z. Zhang, G. Zhou, J. Yu, G. Zhang, J. Yuan, X. Li, S. Wen, F. Meng, S. Yu, X.
Wang, K.H.M. Siddique, Z.-J. Liu*, A.H. Paterson, R.K. Varshney, X. Liang, Sequencing
of cultivated peanut, *Arachis hypogaea*, yields insights into genome evolution and oil
improvement, Mol Plant 12 (2019) 920–934.

[4] W. Zhuang, H. Chen, M. Yang, J. Wang, M.K. Pandey, C. Zhang, W.-C. Chang, L. Zhang,
X. Zhang, R. Tang, V. Garg, X. Wang, H. Tang, C.-N. Chow, J. Wang, Y. Deng, D. Wang,
A.W. Khan, Q. Yang, T. Cai, P. Bajaj, K. Wu, B. Guo, X. Zhang, J. Li, F. Liang, J. Hu, B.
Liao, S. Liu, A. Chitikineni, H. Yan, Y. Zheng, S. Shan, Q. Liu, D. Xie, Z. Wang, S.A.
Khan, N. Ali, C. Zhao, X. Li, Z. Luo, S. Zhang, R. Zhuang, Z. Peng, S. Wang, G.
Mamadou, Y. Zhuang, Z. Zhao, W. Yu, F. Xiong, W. Quan, M. Yuan, Y. Li, H. Zou, H.
Xia, L. Zha, J. Fan, J. Yu, W. Xie, J. Yuan, K. Chen, S. Zhao, W. Chu, Y. Chen, P. Sun, F.
Meng, T. Zhuo, Y. Zhao, C. Li, G. He, Y. Zhao, C. Wang, P.B. Kavikishor, R.-L. Pan,
A.H. Paterson, X. Wang, R. Ming & R.K. Varshney, The genome of cultivated peanut

provides insight into legume karyotypes, polyploid evolution and crop domestication, Nat
Genet 51 (2019) 865–876.

[5]   J.F.M. Valls, C.E. Simpson, New species of *Arachis* (*leguminosae*) from Brazil, Paraguay
and Bolivia, Bonplandia (2005) 35–63.

[6]   H.T. Stalker, Peanut (*Arachis hypogaea* L.), Field Crops Res 53 (1997) 205–217.

[7]   J.C. Jimenez-Lopez, Grain and seed proteins functionality, BoD–Books on Demand, 2021.

[8]   S.S. Arya, A.R. Salve, S. Chauhan, Peanuts as functional food: a review, J Food Sci Technol
53 (2016) 31–41.

[9]   E.M. Ahmed, J.A. Applewhite, Characterization of trypsin inhibitor in Florunner Peanut
Seeds (*Arachis hypogaea* L.), Peanut Sci. 15 (1988) 81–84.

[10] S.B. Mada, A. Garba, A. Mohammed, A. Muhammad, A. Olagunju, H.A. Mohammed,
Effects of boiling and roasting on antinutrients and proximate composition of local and
some selected improved varieties of *Arachis hypogaea* L (Groundnut),).Int. J. Food Nutr.
Saf. (2012) 45–53.

[11] E.C. Reddy, C. Sudhakar, N.P. Reddy, Aflatoxin contamination in groundnut induced by
aspergillus flavus type fungi: A critical review., (2011).

[12] B. Singh, U. Singh, Peanut as a source of protein for human foods, Plant Foods for Hum.
Nutr. 41 (1991) 165–177.

[13] S. Parmar, V. Sharma, D. Bomireddy, P. Soni, P. Joshi, S.S. Gangurde, J. Wang, S.K. Bera,
R.S. Bhat, H. Desmae, K. Shirasawa, B. Guo, R.K. Varshney, M.K. Pandey, Recent
advances in genetics, genomics, and breeding for nutritional quality in groundnut,
Accelerated Plant Breeding, Volume 4: Oil Crops (2022) 111–137.

[14] G.C. Wright, M.G. Borgognone, D.J.O. Connor, R.C.N. Rachaputi, R.J. Henry, A. Furtado,
N.L. Anglin, D.B. Freischfresser, Breeding for improved blanchability in peanut:
phenotyping, genotype× environment interaction and selection, Crop Pasture Sci 69 (2018)
1237–1250.

[15]   P. Janila, R. Aruna, E. Jagadish Kumar, S.N. Nigam, Variation in blanchability in Virginia
groundnut (*Arachis hypogaea* L.), J Oilseeds Res 29 (2012) 116–120.

[16]   W. Korani, D. O'Connor, Y. Chu, C. Chavarro, C. Ballen, B. Guo, P. Ozias-Akins, G.
Wright, J. Clevenger, De novo QTL-seq Identifies Loci Linked to Blanchability in Peanut
(*Arachis hypogaea*) and Refines Previously Identified QTL with Low Coverage Sequence,
Agronomy 11 (2021) 2201.

[17]   Q. Lu, L. Huang, H. Liu, V. Garg, S.S. Gangurde, H. Li, A. Chitikineni, D. Guo, M.K.
Pandey, S. Li, A genomic variation map provides insights into peanut diversity in China
and associations with 28 agronomic traits, Nat Genet 56 (2024) 530–540.

[18]   Z. Zheng, Z. Sun, F. Qi, Y. Fang, K. Lin, S. Pavan, B. Huang, W. Dong, P. Du, M. Tian,
Chloroplast and whole-genome sequencing shed light on the evolutionary history and
phenotypic diversification of peanuts, Nat Genet 56 (2024) 1975–1984.

[19]   C. Zhu, M. Gore, E.S. Buckler, J. Yu, Status and prospects of association mapping in plants,
Plant Genome 1 (2008).

[20]   H. Khazaei, R. Podder, C.T. Caron, S.S. Kundu, M. Diapari, A. Vandenberg, K.E. Bett,
Marker–trait association analysis of iron and zinc concentration in lentil (*Lens culinaris*
Medik.) seeds, Plant Genome 10 (2017) plantgenome2017-02.

[21]   H.D. Upadhyaya, P.J. Bramel, R. Ortiz, S. Singh, Developing a mini core of peanut for
utilization of genetic resources, Crop Sci 42 (2002) 2150–2156.

[22]   H.F. Jiang, R.E.N. Xiao-Ping, X.J. Zhang, J.Q. Huang, L.E.I. Yong, Y.A.N. Li-Ying, L.
Bo-Shou, H.D. Upadhyaya, C.C. Holbrook, Comparison of genetic diversity based on SSR
markers between peanut mini core collections from China and ICRISAT, Acta Agronomica
Sinica 36 (2010) 1084–1091.

[23]   S.S. Gangurde, H. Wang, S. Yaduru, M.K. Pandey, J.C. Fountain, Y. Chu, T. Isleib, C.C.
Holbrook, A. Xavier, A.K. Culbreath, Nested-association mapping (NAM)-based genetic
dissection uncovers candidate genes for seed and pod weights in peanut (*Arachis hypogaea*),
Plant Biotechnol J 18 (2020) 1457–1471.

[24] S.S. Shapiro, M.B. Wilk, An analysis of variance test for normality (complete samples),
Biometrika 52 (1965) 591–611.

[25] M.K. Pandey, G. Agarwal, S.M. Kale, J. Clevenger, S.N. Nayak, M. Sriswathi, A.
Chitikineni, C. Chavarro, X. Chen, H.D. Upadhyaya, M.K. Vishwakarma, S. Leal-Bertioli,
X. Liang, D.J. Bertioli, B. Guo, S.A. Jackson, P. Ozias-Akins, R.K. Varshney, Development
and evaluation of a high-density genotyping 'Axiom_*Arachis*'' array with 58 K SNPs for
accelerating genetics and breeding in groundnut,' Sci Rep 7 (2017) 1–10.

[26] J. Clevenger, Y. Chu, C. Chavarro, G. Agarwal, D.J. Bertioli, S.C.M. Leal-Bertioli, M.K.
Pandey, J. Vaughn, B. Abernathy, N.A. Barkley, R. Hovav, M. Burow, S.N. Nayak, A.
Chitikineni, T.G. Isleib, C.C. Holbrook, S.A. Jackson, R.K. Varshney & P. Ozias-Akins,
Genome-wide SNP genotyping resolves signatures of selection and tetrasomic
recombination in peanut, Mol Plant 10 (2017) 309–322.

[27] Y. Tang, X. Liu, J. Wang, M. Li, Q. Wang, F. Tian, Z. Su, Y. Pan, D. Liu, A.E. Lipka, E.S.
Buckler, Z. Zhang, GAPIT version 2: an enhanced integrated tool for genomic association
and prediction, Plant Genome 9 (2016) plantgenome2015-11.

[28] W.G. Hill, A. Robertson, Linkage disequilibrium in finite populations Theor. Appl. Genet
38 (1968) 226–231.

[29] L. Yin, Package" CMplot". 2019, URL Https://Github. Com/YinLiLin/R-
CMplot/Blob/Master/CMplot. r (2019).

[30] J. Su, F. Zhang, X. Chong, A. Song, Z. Guan, W. Fang, F. Chen, Genome-wide association
study identifies favorable SNP alleles and candidate genes for waterlogging tolerance in
chrysanthemums, Hortic Res 6 (2019).

[31] A. Alemu, T. Feyissa, M. Maccaferri, G. Sciara, R. Tuberosa, K. Ammar, A. Badebo, M.
Acevedo, T. Letta, B. Abeyo, Genome-wide association analysis unveils novel QTLs for
seminal root system architecture traits in Ethiopian durum wheat, BMC Genomics 22 (2021)
1–16.

[32] C. He, J. Holme, J. Anthony, SNP genotyping: the KASP assay, Crop Breeding: Methods
and Protocols (2014) 75–86.

[33]   R. Bohar, A. Chitkineni, R.K. Varshney, Genetic molecular markers to accelerate genetic
784   gains in crops, Biotechniques 69 (2020) 158–160.

785   [34]   P. Shah, G. Wright, C. V Nwosu, D. O'Connor, P. Tsatsos, P. Janila, K. Praveen, K. Singh,
786   S.K. Bera, M. Thudi, C. Kole, R.K. Varshney, M.K. Pandey, Industry perspective, genetics
787   and genomics of peanut blanchability, Plant Science (2025) 112473.

788   [35]   P.C. Barnes Jr, C.E. Holaday, J.L. Pearson, Device to measure ease of skin removal from
789   peanuts, J Food Sci 36 (1971) 405–407.

790   [36]   M.W. Hoover, A rotary air impact peanut blancher, Peanut Sci. 6 (1979) 84–87.

791   [37]   F.S. Wright, R.W. Mozingo, Laboratory device for peanut skin removal, Peanut Sci. 2
792   (1975) 11–15.

793   [38]   U. Singh, R. Sridhar, S.L. Dwivedi, S.N. Nigam, R. Jambunathan, Evaluation of Blanching
794   Quality in Groundnut (Arachis hypogaea L..), J Food Sci Technol 33 (1996) 211–214.

795   [39]   R.K. Varshney, S.N. Nayak, G.D. May, S.A. Jackson, Next-generation sequencing
796   technologies and their implications for crop genetics and breeding, Trends Biotechnol 27
797   (2009) 522–530.

798   [40]   M.K. Pandey, M. Roorkiwal, V.K. Singh, A. Ramalingam, H. Kudapa, M. Thudi, A.
799   Chitikineni, A. Rathore, R.K. Varshney, Emerging genomic tools for legume breeding:
800   current status and future prospects, Front Plant Sci 7 (2016) 455.

801   [41]   M.K. Pandey, A.K. Pandey, R. Kumar, C.V. Nwosu, B. Guo, G.C. Wright, R.S. Bhat, X.
802   Chen, S.K. Bera, M. Yuan, H. Jiang, I. Faye, T. Radhakrishnan, X. Wang, X. Liang, B.
803   Liao, X. Zhang, R.K. Varshney, W. Zhuang, Translational genomics for achieving higher
804   genetic gains in groundnut, Theor. Appl. Genet. 133 (2020) 1679–1702.

805   [42]   X. Liu, M. Huang, B. Fan, E.S. Buckler, Z. Zhang, Iterative usage of fixed and random
806   effect models for powerful and efficient genome-wide association studies, PLoS Genet 12
807   (2016) e1005767.

808   [43]   Q. Wang, F. Tian, Y. Pan, E.S. Buckler, Z. Zhang, A SUPER powerful method for genome
809   wide association study, PLoS One 9 (2014) e107684.

[44] M. Huang, X. Liu, Y. Zhou, R.M. Summers, Z. Zhang, BLINK: a package for the next level
of genome-wide association studies with both individuals and markers in the millions,
Gigascience 8 (2019) giy154.

[45] B. Song, W. Ning, D. Wei, M. Jiang, K. Zhu, X. Wang, D. Edwards, D.A. Odeny, S. Cheng,
Plant genome resequencing and population genomics: Current status and future prospects,
Mol Plant 16 (2023) 1252–1268.

[46] Q. Zhao, Q. Feng, H. Lu, Y. Li, A. Wang, Q. Tian, Q. Zhan, Y. Lu, L. Zhang, T. Huang, Y.
Wang, D. Fan, Y. Zhao, Z. Wang, C. Zhou, J. Chen, C. Zhu, W. Li, Q. Weng, Q. Xu, Z.-X.
Wang, X. Wei, B. Han, X. Huang, Pan-genome analysis highlights the extent of genomic
variation in cultivated and wild rice, Nat Genet 50 (2018) 278–284.

[47] N. Wu, M. Yang, U. Gaur, H. Xu, Y. Yao, D. Li, Alpha-ketoglutarate: physiological
functions and applications, Biomol Ther (Seoul) 24 (2016) 1.

[48] F.N. Xia, B. Zeng, H.S. Liu, H. Qi, L.J. Xie, L.J. Yu, Q.F. Chen, J.F. Li, Y.Q. Chen, L.
Jiang, S. Xiao, SINAT E3 ubiquitin ligases mediate FREE1 and VPS23A degradation to
modulate abscisic acid signaling, Plant Cell 32 (2020) 3290–3310.

[49] Y. Nakano, M. Yamaguchi, H. Endo, N.A. Rejab, M. Ohtani, NAC-MYB-based
transcriptional regulation of secondary cell wall biosynthesis in land plants, Front Plant Sci
6 (2015) 288.

[50] D. Sui, B. Wang, Transcriptome Analysis Reveals Complex Defensive Mechanisms in Salt-
Tolerant and Salt-Sensitive Shrub Willow Genotypes under Salinity Stress, Int J Genomics
2020 (2020) 6870157.

[51] J.K.C. Rose, J. Braam, S.C. Fry, K. Nishitani, The XTH family of enzymes involved in
xyloglucan endotransglucosylation and endohydrolysis: current perspectives and a new
unifying nomenclature, Plant Cell Physiol 43 (2002) 1421–1435.

[52] H. Koiwa, F. Li, M.G. McCully, I. Mendoza, N. Koizumi, Y. Manabe, Y. Nakagawa, J.
Zhu, A. Rus, J.M. Pardo, R.A. Bressan, P.M. Hasegawa, The STT3a subunit isoform of the
Arabidopsis oligosaccharyltransferase controls adaptive responses to salt/osmotic stress,
Plant Cell 15 (2003) 2273–2284.

[53]   Y. Cai, Lipid synthesis and beyond: SAD fatty acid desaturases contribute to seed
development, (2020).

Table. 1: Marker trait association identified candidate genes and significant STAs for groundnut blanchability using 58 K
"Axiom_*Arachis*" array

Trait	SNP ID	Chr	Position	*p* value	Model	SNP effect	PVE (%)	Gene ID	Annotations
BLCB_50R, BLCB_50_PR_R	AX_147256471	B07	118262617	1.05E-07	a, c	13.72	24.87	*Araip.V0BSS*	NRAMP metal ion transporter 6
BLCB_50PR	AX_147234396	A09	114692575	4.28E-06	c	11.99	5.08	*Aradu.G1YNF*	fatty acid desaturase 2
BLCB_50PR	AX_147247942	B04	105437967	1.40E-06	c	-12.43	6.10	*Araip.LL89K*	ethylene-responsive transcription factor
BLCB_50PR	AX_147247969	B04	108209828	8.80E-06	c	11.97	5.32	*Araip.BZ4NA*	isocitrate dehydrogenase
BLCB_50PR, BLCB_200PR	AX_147234469	A09	115432582	3.79E-07	c	12.44	32.24	*Aradu.N9ZS9*	Transcription factor jumonji (jmjC) domain-containing protein
BLCB_50PR	AX_147231354	A08	41497800	6.13E-06	c	-21.86	1.68	*Aradu.YDH5Z*	5-methylthioadenosine/ S-adenosylhomocysteine deaminase
BLCB_50PR	AX_147257272	B08	3609752	3.01E-06	c	-21.88	17.13	*Araip.2B9XL*	4-hydroxy-3-methylbut-2-enyl diphosphate synthase
BLCB_50PR	AX_147258765	B08	110518175	3.01E-06	c	-21.88	14.18	*Araip.FRJ8B*	carotenoid cleavage dioxygenase 1

BLCB_200PR	AX_147216911	A03	44353475	2.14E-07	a, b, c	7.36	20.00	*Aradu.D66ST*	Helicase-like protein
BLCB_200PR	AX_147254562	B07	2964357	3.69E-06	a, b, c	-9.66	35.29	*Araip.GTG68*	DTW domain-containing protein
BLCB_200PR	AX_147260814	B09	26207513	2.30E-07	a, b, c	8.26	17.59	*Araip.1L939*	DNA repair protein UVH3
BLCB_200PR	AX_147222611	A05	71291610	2.95E-06	c	10.37	0.41	*Aradu.BRU54*	Epstein-Barr virus EBNA-1
BLCB_200PR	AX_147225399	A06	57729112	1.89E-06	c	10.96	1	*Aradu.G23Y9*	ubiquitin-protein ligase 7
BLCB_200PR	AX_147214749	A02	84589354	4.41E-06	c	-7.43	0.91	*Aradu.QS8M2*	WEB family protein
BLCB_200PR	AX_147218592	A03	133198789	4.94E-06	c	-7.26	3.45	*Aradu.4PJ68*	unknown protein
BLCB_200PR	AX_147247867	B04	101149286	4.36E-06	c	7.28	9.74	*Araip.UM7BZ*	DEAD-box ATP-dependent RNA helicase
BLCB_200PR	AX_147253251	B06	114586674	1.12E-06	c	16.11	32.81	*Araip.2H071*	xyloglucan endotransglucosylase/ hydrolase 5
BLCB_50_PR_R	AX_147207637	A01	230616	4.37E-06	a	6.65	5.82	*Aradu.0B4KA*	uncharacterized protein
BLCB_50_PR_R	AX_147216570	A03	29722463	5.45E-13	b, c	-9.48	25.49	*Aradu.K3EGK*	Unknown protein
BLCB_50_PR_R	AX_147227734	A07	12521563	2.91E-06	b	-6.16	17.60	*Aradu.V4VHC*	pumilio 2

BLCB_50_PR_R	AX_147222844	A05	85198860	5.08E-06	a, b, c	-6.83	23.26	*Aradu.B0E28*	Oligosaccharyl transferase STT3 subunit homolog
BLCB_50_PR_R	AX_147211857	A01	97179228	5.36E-07	c	12.42	1.46	*Aradu.C1KH2*	chromodomain-helicase-DNA-binding protein
BLCB_50_PR_R	AX_147219449	A04	10026356	4.46E-06	c	11.77	7.05	*Aradu.MI7GZ*	Unknown protein
BLCB_50_PR_R	AX_147224932	A06	12625218	7.29E-07	c	6.03	7.12	*Aradu.QJ7XC*	ubiquitin ligase SINAT3
BLCB_50_PR_R	AX_147239773	B02	163783	4.85E-06	c	11.41	2.12	*Araip.PG50R*	N-terminal nucleophile aminohydrolases (Ntn hydrolases) superfamily protein
BLCB_50_PR_R	AX_147244167	B03	25025384	4.23E-06	c	11.16	4.00	*Araip.Q36GV*	putative membrane-bound O-acyltransferase C24H6.01c
BLCB_50_PR_R	AX_147244335	B03	31216966	5.94E-09	c	-8.19	13	*Araip.80IK3*	metallothionein 3
BLCB_50_PR_R	AX_147256997	B08	1018858	6.85E-07	c	-6.49	1.24	*Araip.1D5HT*	charged multivesicular body protein

Legends: - a: FarmCPU, b: BLINK, c: SUPER, BLCB_50R (Blanchability for 50 gm rainy sample), BLCB_50PR (Blanchability for 50 gm post-rainy sample), BLCB_200PR (Blanchability for 200 gm post-rainy sample), BLCB_50_PR_R (Blanchability for mean of 50

845   gm sample rainy and post-rainy). Some SNPs are positively regulating the trait (SNP effect positive) while some shows negative

846   regulation (SNP effect negative).

Figure 1: Protocol for phenotyping blanchability in groundnut

Figure 2: Different phenotypic analysis of blanchability across groundnut minicore: A. Phenotypic variability for blanchability in groundnut minicore collection, **B.** Diverse agronomic types in groundnut revealed substantial variation for blanchability in minicore collection across different seasons and sample sizes, **C.** The raincloud plot depicts the blanchability (%) values in season 1 (Blanchability_50R, pink color) and season 2 (Blanchability_50PR and 200PR, green and orange color; respectively), displaying considerable phenotypic variation with a continuous distribution of accession across both the seasons.

Figure 3. The PCA plot to depict the phenotypic variation. The distribution of the phenotypic dataset observed for blanchability across different seasons and sample size. The variation has been observed more for blanchability in 50 R. First two PCs collectively showing the 93.3% of phenotypic variation.

Figure 4: Pearson correlation analysis of blanchability, aflatoxin and agronomic traits were evaluated across diverse environments post-rainy (PR) and rainy (R). The four traits including pods per plant (PPP), blanchability (BLCB), aflatoxin (AFTX), *A. flavus* infection (AFLV), days to maturity (DM), pod length (PLN), seed length (SDL), seed weight (SWT), seed width (SWD), and pod width (PDWD) were evaluated in rainy and post-rainy environment. The color reflect the strength of correlation and further cluster has been made. PPP (R) positively correlated with blanchability and hence clustered together and other agronomic traits and aflatoxin make separate clusters. Scale ranges from -0.5 to 1, where the green color represents the least correlation while the red color shows strong correlation between the traits.

Figure 5: PCA depicting phenotypic variability of 88.9% for groundnut minicore collection, and identified 4 different clusters (pink, green, blue, brown) based on groundnut sub-species, botanical variety and agronomic type. Cluster 1 (pink color): (46 genotypes): *fastigata*, vulgaris, spanish bunch; Cluster 2 (green color): (75 genotypes): hypogaea, hypogaea, virginia bunch, Cluster 3 (Blue): (27 genotypes): fastigata, fastigata, valencia bunch, Cluster4 (Brown) (19 genotypes): fastigata, fastigata, vulgaris, spanish bunch, respectively.

Figure 6: Manhattan plots representing the identified STAs associated with the blanchability on the basis SUPER model in GAPIT for A: Blanchability 50R (50 gm sample size in rainy season), B: Blanchability 50PR (50 gm sample size in post-rainy season), C: Blanchability 200PR (200 gm sample size in post-rainy season) and D: Blanchability 50PR_R (50 gm sample size in post-rainy and rainy season), respectively. In these Manhattan plots, the highlighted (pink) region shows the gene associated with the validated polymorphic markers on chromosome A01, A06, B04 and B07; respectively. The other dots represent the significant STAs, and the genes found to be associated with blanchability.

Figure 7: Development and validation of Kompetitive allele specific polymerase chain reaction (KASP) markers from potential candidate genes identified for blanchability. A. Validation panel includes 20 good blanchability genotypes and groundnut varieties (50-80%-blanchability), and five poor blanchability genotypes (1-10%-blanchability).Four single-nucleotide polymorphisms (SNPs) show clear homozygous clusters and heterozygous cluster for **B.** snpAH00551 (A01_230616) for uncharacterized protein (*Aradu.0B4KA*) gene, **C.** snpAH00554 (A06_57729112) for *ubiquitin-protein ligase* (*Aradu.G23Y9*) gene **D.** snpAH00558 (B04_108209828) for *isocitrate dehydrogenase* (*Araip.BZ4NA*) gene, **E.** snpAH00559 (B07_2964357) for *DTW domain-containing protein* (*Araip.GTG68*) gene.