X-chromosome upregulation operates on a gene-by-gene basis at RNA and protein levels

Ryan N. Allsop[1,2,*], Jeffrey Boeren[3,*], Beatrice F. Tan[3,*], Sarra Merzouk[3], Suresh Poovanthingal[2,4], Wilfred F.J. van IJcken[5], Jeroen A.A. Demmers[6], Hegias Mira Bontenbal[3], Cristina Gontan[3], Joost Gribnau[3,#], Vincent Pasque[1,2,#].

[1] Department of Development and Regeneration, University of Leuven (KU Leuven), Leuven, Belgium
[2] KU Leuven Institute for Single Cell Omics (LISCO), Leuven, Belgium
[3] Department of Developmental Biology, Erasmus University Medical Center, Rotterdam, The Netherlands
[4] VIB-KU Leuven Center for Brain & Disease Research, Leuven, Belgium
[5] Erasmus Center for Biomics, Erasmus University Medical Center, Rotterdam, the Netherlands
[6] Proteomics Center, Erasmus University Medical Center, Rotterdam, the Netherlands

* These authors contributed equally: Ryan N. Allsop, Jeffrey Boeren, Beatrice F. Tan
# These authors jointly supervised this work: Joost Gribnau, Vincent Pasque

ABSTRACT

Gene dosage compensation mechanisms are crucial for mammalian development. In mice, recent findings show that cells can sense the number of X chromosomes. Loss or inactivation of one of the two X chromosomes is compensated by upregulating the remaining active X chromosome, a process termed X-chromosome upregulation (XCU). However, how cells sense X-chromosome dosage and induce XCU remains unclear. Here, we show that heterozygous X chromosome fragment deletions in mouse pluripotent stem cells induces XCU *in trans*, and that compensation takes place at the mRNA and protein level. Furthermore, we found that inducing gene silencing *in cis* on autosomes induces gene dosage compensation *in trans*. This work provides significant insights into the molecular foundations of mammalian gene dosage compensation.

INTRODUCTION

Most mammals are diploid, with two copies of each chromosome per cell[1]. In some cases however, the number of chromosomes or segments of chromosomes can change. Changes in gene copy number, also known as changes in gene dosage, have been closely linked to human diseases[2,3]. Gene dosage compensation mechanisms attempt to resolve imbalances in gene dosage[4]. However, how gene dosage compensation is triggered in mammals remains unclear.

A paradigm to study gene dosage compensation mechanisms is the X chromosome. In placental mammals, the female sex is typically characterized by two X chromosomes, while the male sex typically has one X chromosome and one Y chromosome. This imbalance in X-linked gene dosage between sexes is thought to have driven the evolution of X-linked dosage compensation mechanisms[4–11]. In female mammals, one of the two X chromosomes expresses the long non-coding RNA *Xist* and is inactivated *in cis* (on the same chromosome) during early development, a process termed X-chromosome inactivation (XCI)[12,13,14,15]. As a result, there is only one active X chromosome in female somatic cells, while male XY somatic cells also have one active X chromosome. In contrast to the X chromosome, each autosome is present in two active copies. In 1963, Susumu Ohno hypothesized that a dosage compensation mechanism evolved to upregulate X-linked gene expression on the active X chromosome, to balance X-linked gene dosage with the gene dosage of diploid autosomes[9]. This process is referred to as X-chromosome upregulation (XCU)[4]. Recent studies using allele-resolution transcriptome analyses have confirmed that XCU takes place at the transcriptional level in mouse cells with a single active X chromosome, and that XCU is developmentally regulated[4,12,16–34].

Multiple lines of evidence using allele-resolution transcriptome analyses have indicated that mammalian cells can sense the number of X chromosomes they have, as well as the transcriptional status of the X chromosome, and compensate X-linked gene expression accordingly[16,18,20]. Before XCI, female diploid XX mouse pluripotent stem cells (PSCs) have two active X chromosomes (Xa), while male XY cells have only one. By consequence, female mouse PSCs have higher levels of X-linked gene products than male cells. However, recent studies showed that unlike previously thought, each active X chromosome in female XX mouse PSCs and naive epiblast are not upregulated, consistent with no requirement for dosage compensation between two active X chromosomes and diploid autosomes[16,20]. Furthermore, when XX embryonic fibroblasts, with one upregulated active (Xu) and one inactive X chromosome (Xi), are reprogrammed to induced pluripotent stem cells (iPSCs), reactivation of the Xi occurs concomitantly with loss of upregulation from the Xu[16]. A study in mouse iPSCs showed that when one of the two non-upregulated Xa chromosomes is lost in PSCs, XCU is induced on the remaining X chromosome to compensate gene dosage imbalance with diploid autosomes[16]. This observation was later confirmed in mouse embryonic stem cells (mESCs)[20]. Similarly, XCU is also observed in male XY mESCs[20]. In summary, when both X chromosomes are active, there is no XCU. However, when only one active X chromosome is present, XCU is triggered on the remaining active X chromosome.

Collectively, these results suggest that mammalian cells can sense the number of active X
chromosomes and compensate for the presence of a single active X chromosome by XCU.
However, how cells sense the number of active X chromosomes and trigger compensation
for gene dosage imbalances, and if they can sense when specific X chromosome regions or
genes are missing, remains unclear.

The molecular mechanisms controlling XCU in mammals are still unknown. Previous studies
have attempted to identify active histone modifications associated with XCU in mice,
however the results have been conflicting or unclear. One study reported enrichment of
histone 4 lysine 16 acetylation (H4K16ac) on the Xu, but analyzed cells with two active X
chromosomes, which we now know do not undergo XCU[29]. Another study reported
enrichment of histone 3 lysine 4 tri-methylation (H3K4me3) as well as histone 3 lysine 36 tri-
methylation (H3K36me3) on the Xu, however this enrichment was compared to autosomes,
rather than to the Xa[25]. Finally, a recent study found no enrichment of any common histone
modifications on the Xu compared to the Xa[20]. Therefore, it remains unclear whether histone
modifications are enriched on the mammalian Xu.

Changes in gene dosage of autosomes is a hallmark of cancer and early human IVF
embryos[35–39]. Studies in human cancer cells and human embryos have suggested that gene
dosage compensation can occur on autosomes at both the transcriptional and translational
levels[35,40]. However, it is unknown if these effects can also be induced by gene silencing on
an autosome, as opposed to genetic loss.

In this study, we sought to investigate how the X chromosome and autosomes sense and
compensate for changes in gene dosage in mammals. We use CRISPR-Cas9 to generate
large heterozygous X-linked deletions in female XX mESCs, followed by allele and
molecule-resolution single-cell and bulk transcriptomics, as well as quantitative proteomics
to measure the impact of the deletions on gene expression. We find evidence for dosage
compensation by increased transcription and protein levels specifically within the deleted
regions *in trans*. Furthermore, we investigated whether dosage compensation of autosomal
genes can occur without loss of genetic material, and found evidence for transcriptional
dosage compensation *in trans* (on the other chromosome) when one of two alleles is
inactivated *in cis*. In conclusion, this work establishes that mammalian cells can sense the
heterozygous loss or inactivation of specific fragments of chromosomes or genes and
compensate these gene dosage imbalances by upregulation at the transcriptional and
protein levels *in trans*.

107
RESULTS
Allele and molecule-resolution analysis of XCU in mouse PSCs
To investigate XCU, we used single-cell RNA-sequencing (scRNA-seq) in mouse PSCs with
a range of genotypes. The cell lines originate from a cross between two mouse subspecies
(*Mus musculus musculus* x *Mus musculus castaneus*, *Mus/Cast*). The cells were 1) female
iPSCs with two active X chromosomes ($X^{Cast}X^{Mus}$), 2) male ESCs ($X^{Mus}Y$) and 3) female
mESCs with X-chromosome monosomy of the paternal or maternal X chromosome (XO,

either $X^{Cast}O^{Mus}$ or $X^{Mus}O^{Cast}$) (**Figure 1a**, **Supplementary Fig. 1a**)[41]. Previous studies
indicated that both mESCs and iPSCs behave similarly with respect to X chromosome
dosage compensation properties[16,20,41–43]. We applied Smart-seq3xpress, a plate-based
scRNA-seq method providing full transcript coverage with the ability to count the number of
transcript molecules using unique molecular identifiers (UMIs)[44]. Transcripts were further
split by allele of origin based on a list of ~17.5 million single-nucleotide polymorphisms
(SNPs) unique to each subspecies[16,45,46], providing allele-specific resolution data.

To quantify gene expression from each X chromosome, we measured the mean normalized
expression (transcripts per million, TPM) of all expressed X-linked genes from each allele for
each cell line. This enabled us to identify which X-chromosome was lost in the XO cells
(**Supplementary Fig. 1b**). XX iPSCs and ESCs have been previously described to lose one
of the two X-chromosomes[41,43,47,48]. Consistent with this, after quality filtering, we obtained 6
cells which lost the *Cast* allele ($X^{Mus}O^{Cast}$), and 44 cells which lost the *Mus* allele ($X^{Cast}O^{Mus}$).
As expected, we observed significantly higher allelic expression levels when only one active
X chromosome is present, in $X^{Mus}Y$ and $X^{Mus}O^{Cast}$ cells (**Figure 1b**, **Supplementary Fig. 2a**).
When the *Mus* allele rather than the *Cast* X chromosome allele was lost, in $X^{Cast}O^{Mus}$ cells, X-
linked gene expression was increased on the *Cast* allele (**Figure 1c**). In summary, loss of
one X chromosome in XX mouse PSCs induces global compensation of gene expression on
the remaining monosomic X chromosome. Allelic X-to-autosome ratio (X:A) analysis
confirmed elevated X:A ratios in cells with a single X chromosome, while $X^{Cast}X^{Mus}$ cells
exhibited no increased X:A ratio, confirming previous findings[16,20] (**Figure 1d-e**,
Supplementary Fig. 2b). These findings confirm the presence of XCU in diploid cells with
a single X chromosome[16], and that mouse PSCs can sense the number of X chromosomes
and adapt gene expression by undergoing XCU when only one Xa is present. Consistent
with previous studies, the average expression from the *Mus* allele tended to be slightly
higher than the *Cast* allele, a slight bias most likely induced to mapping reads to the *Mus*
musculus musculus reference genome[16,18,49]. Together, these results confirm that
transcriptional dosage compensation by XCU occurs in mouse PSCs with a single active X
chromosome, but not with two active X chromosomes. Moreover, we can accurately
measure and identify XCU using a hybrid/polymorphic mouse PSC system combined with
allele-specific expression analysis using Smart-seq3xpress scRNA-seq.

Identification of genes subject to XCU
We next sought to characterize which X-linked genes are transcriptionally dosage
compensated in this system. Of the 405 expressed X-linked genes for which we obtained
allele-resolution expression data, 103 (~40%) were found to be significantly transcriptionally
upregulated in XO and XY cells (collectively $X^{monosomy}$) compared to XX cells (**Supplementary
Fig. 2c, Supplementary Table 1**). These results are in line with previous findings that
transcriptional XCU is specific to a subset of genes[17]. Both non-upregulated and
upregulated genes were found to be located across the X chromosome and not enriched to
a specific locus (**Supplementary Fig. 2d**). The mean allelic expression of the upregulated
genes was significantly higher in $X^{monosomy}$ cells than in XX cells (**Supplementary Fig. 2e-h**).

To characterize the expression levels of genes subject to XCU in cells that do not undergo
XCU, we compared in XX cells the mean expression of genes defined as upregulated and
non-upregulated. We found that genes that are upregulated in $X^{monosomy}$ are expressed at
significantly higher levels in XX iPSCs compared to genes that are not subject to
upregulation in $X^{monosomy}$ (**Figure 1f**). This is in line with previous findings that highly
expressed genes are also more likely to be dosage-sensitive[50], and that dosage
compensation mechanisms have evolved primarily to protect dosage-sensitive genes from
the harmful effects of gene dosage imbalances[51].

Interestingly, gene ontology analysis indicated that upregulated X-linked genes were
enriched for critical basic cellular functions such as regulation of chromatin organization
and positive regulation of nucleic-acid templated transcription, whereas non-upregulated
genes were enriched for specific developmental processes such as visual perception and
negative regulation of axonogenesis, processes that are irrelevant at this stage of
development (**Supplementary Fig. 2i**). Proteins encoded by upregulated genes were more
likely to have protein-protein interactions (PPIs) with transcription factors, which could be
due to feedback loops which allow upregulated genes to self-regulate their expression
(**Supplementary Fig. 2j**)[52]. Genes encoding proteins that form complexes have been
hypothesized to be particularly dosage-sensitive, potentially requiring gene dosage
compensation in order to maintain the correct stoichiometry for proper complex
function[51,53]. However, despite an apparent trend, our upregulated X-linked genes did not
contain a significantly higher proportion of genes which code for proteins in complexes
when compared to non-upregulated X-linked genes (**Supplementary Fig. 2k**). Examples of
important complexes containing subunits encoded by upregulated genes include the MLL1,
MSL, NSL, and DNA polymerase complexes, which contain subunits respectively encoded
by *Hcfc1*, *Msl3*, *Ogt*, and *Pola1* genes.

Proteins which form condensates have been proposed to be particularly dosage-sensitive[54].
However, we found no significant difference in the proportion of proteins forming
condensates between upregulated and non-upregulated genes, indicating that
transcriptionally upregulated X-linked genes may not be enriched for genes encoding
proteins known to form condensates (**Supplementary Fig. 2l**). Given the challenges in
studying condensates, it remains possible that not all condensate-forming proteins have
been identified, and future discoveries may revise this conclusion[55]. Furthermore, we
hypothesized that genes which escape XCI, and thus retain expression from both alleles,
require dosage compensation in $X^{monosomy}$ cells. We observed a non-significant trend
towards an increased proportion of upregulated genes that escape XCI (**Supplementary
Fig. 2m**), potentially due to the limited statistical power from the small escapee gene set
(n=38)[56]. Nearly all orthologs of the upregulated X-linked genes were associated with
diseases in the OMIM database, however, this proportion was not significantly higher than
that of the non-upregulated genes (**Supplementary Fig. 2n**). Taken together, these results
suggest that upregulated genes may be dosage-sensitive, and this may be linked to high
levels of gene expression, involvement in protein complexes, and important cellular
function.

Histone post-translational modifications and XCU

To explore a potential link between histone post-translational modifications (hPTMs) and XCU, we analyzed hPTMs in a system that enables conditional induction of XCU in mouse XX ESCs[57]. In the study of Zylicz et al.[57], a doxycycline inducible promoter was inserted upstream of the endogenous *Xist* gene on the *Mus* allele in polymorphic ($X^{Cast}X^{Mus}$) mESCs[57]. Addition of doxycycline for 24 hours activates *Xist* on the *Mus* allele, inducing XCI *in cis*, and triggers XCU on the opposite active X^{Cast} chromosome *in trans*[20]. Therefore, this experimental system enables us to examine hPTMs during XCU. To assess the enrichment of various hPTMs, we examined native ChIP-seq data with allele resolution for several hPTMs (H4ac, H3K27ac, H3K4me1, H3K4me3, H3K9ac, H3K27me3, H2AK119ub) at 0, 4, 12, and 24 hours of XCI and XCU induction with allele resolution (**Figure 1g**). As expected, during XCI, there was an increase in repressive marks H3K27me3 and H2AK119ub at the promoter region and across the gene body of X-linked genes on the Xi (**Supplementary Fig. 3a-c**). By contrast, on the Xa during XCU, there was no increase in the repressive mark H3K27me3 and minor increase in H2AK119ub in the promoter regions (**Figure 1h, Supplementary Fig. 3d**). Across the gene body, there was a significant decrease in H3K27me3 and a small but protracted increase in H2AK119ub on the Xa during XCU (**Supplementary Fig. 3d-e**). Additionally, active marks decreased in the promoter region of genes on the Xi (**Supplementary Fig. 4a-b**). However, on the Xa, H4ac, H3K27ac, H3K4me1, and H3K4me3 remained high, while H3K9ac significantly increased during XCU (**Figure 1i, Supplementary Fig. 4c**). These results indicate that while active marks become depleted during XCI, a subset of active hPTMs become more enriched during XCU, or remain high (**Supplementary Fig. 4d**). Our results implicate these hPTMs in XCU in mammals in contrast to the recent view that they are not enriched on the Xu[20,25,29,41].

In addition, we found that before XCU, all active marks studied are more highly enriched on genes that will become upregulated during XCU versus those that do not, in line with the higher expression levels of these genes before XCU (**Figure 1i, Supplementary Fig. 4c**). Unlike a previous study[20], we find evidence for an increase in several active hPTMs during XCU. This could be because the changes we detected are restricted to the promoter region. A previous study did propose increased hPTM on the Xu in XX mESCs[25], but given we now know XX mESCs have no XCU, the conclusions from this study must be revisited. By contrast, strong evidence has been provided for a role of H4K16ac in XCU in *Drosophila melanogaster* where the male-specific lethal (MSL) complex deposits H4K16ac on the Xu[58]. To investigate if mammals and *Drosophila* both acquired mechanisms to enrich H4K16ac on the Xu, we investigated whether H4K16ac is enriched on the Xu in mice. A previous study analyzed acetylated H4 (H4ac) in mESCs but did not identify any enrichment on the Xu, the pan-H4ac detection potentially masking H4K16ac enrichment[57]. We therefore performed H4K16ac ChIP-seq in both XX and XO mESCs, which showed no significant increase in H4K16ac on the Xu compared to the Xa (**Supplementary Fig. 4e-f**). In summary, we found evidence for enrichment of active hPTMs during XCU in mammals similar to *Drosophila*. However, different sets of histone marks appear to be enriched on the Xu in different species, H4K16ac in *Drosophila*[58], and H3K9ac in mice.

No evidence for an XCU center

To gain insights into how cells sense the number of active X chromosomes they carry and subsequently induce XCU, we hypothesized the existence of a dedicated regulatory locus on the X chromosome, which we term the XCU center (Xuc). By analogy to the well-characterized X-inactivation center (Xic)[8,14,59], the Xuc is defined as an X-linked locus whose loss or inactivation on one allele leads to chromosome-wide upregulation of gene expression *in trans* on the other X chromosome (**Figure 2a**). We propose that the presence of two active copies of the Xuc prevents XCU, whereas having only one active copy is necessary to induce XCU. Specifically, when both Xuc copies are active (two Xa), no XCU occurs; when one Xuc is active and the other is inactive or absent (one Xa), XCU is induced on the Xa. To test the Xuc hypothesis, we investigated the impact of deleting large segments on one of the two X chromosomes in XX mESCs on gene expression. We wondered if deleting a large segment or region on one of the two X chromosomes would be sufficient to induce chromosome-wide XCU *in trans*.

We used XX mESCs with large heterozygous segmental deletions on one of the two X chromosomes. The cell lines were 1) Deletion 1: a 49 Mb deletion between *Tfe3* and *Zic3*, 2) Deletion 2: a 15 Mb deletion between *Zic3* and *Dusp9*, and 3) Deletion 3: a 15 Mb deletion of the region between *Dusp9* and *Nr0b1* (**Figure 2b**, **Supplementary Table 2**)[41]. We also included $X^{Cast}X^{Mus}$ iPSCs, $X^{Cast}O^{Mus}$ iPSCs, $X^{Mus}O^{Cast}$ iPSCs and $X^{Mus}Y$ ESCs as controls and performed Smart-seq3xpress scRNA-seq as above. After quality filtering, we obtained allele-resolution expression values for 405 X-linked and 12,016 autosomal genes in 325 cells. ScRNA-seq data confirmed the genotype of the cells (**Figure 2c**). All cell lines expressed pluripotency markers, indicating that cells maintained a pluripotent stem cell identity (**Supplementary Fig. 5a**). Correlation analysis confirmed that the heterozygous deletions did not affect the transcriptome, as indicated by the strong correlation between all cell lines (**Supplementary Fig. 5b**). In summary, we generated allele-resolution scRNA-seq data for polymorphic PSCs with distinct heterozygous deletions of the X chromosome.

We first asked if deleting a large segment on one of the two active X chromosomes results in induction of global XCU *in trans*. Allele-specific gene expression analysis showed that none of the three deletions examined induced full global XCU *in trans* (**Supplementary Fig. 5c-f**). However, we noted a slight increase in gene expression *in trans* in two cell lines, Deletion 1 and 3, but the magnitude of the increase was low. To understand where this increase may come from, we first analyzed gene expression specifically outside of the deleted segments *in trans* (**Figure 2d**). While full loss of one of the two X chromosomes induces XCU, none of the segmental deletions induced full XCU *in trans* for genes located outside of the deletion segments (**Figure 2e-g, Supplementary Fig. 5g-j**).

To determine if the conclusions can be extended to other regions of the X chromosome, we proceeded to test deletions that collectively span the entire X chromosome. We generated 13 new individual large segmental heterozygous deletions of the X chromosome in $X^{Cast}X^{Mus}$ mESCs (**Figure 3a**, **Supplementary Fig. 6a**, **Supplementary Table 2**). This included ten deletions, labeled A to K, as well as three larger deletions spanning multiple regions (AB, C-G, I-TEL). Deleted segments were chosen to respect syntenic segments and TAD

boundaries[60] as much as possible (**Supplementary Fig. 6b**). These deletions were
generated in a *Rnf12*[+/-] parental cell line, which exhibits highly similar expression to wild
type XX mPSCs as expected (**Supplementary Fig. 6c**), and thus served as the reference
sample for these analyses. We generated bulk RNA-seq data from two independent clones
from most deleted lines. We used $X^{Cast}X^{Mus}$, $X^{Mus}O^{Cast}$, and $X^{Cast}O^{Mus}$ mESCs as controls. After
quality filtering, we obtained expression values for 447 X-linked and 11,203 autosomal
genes, enabling us to investigate X-linked gene dosage compensation. Allele-resolution
analysis of X-linked gene expression confirmed the expected genotype of the cells (**Figure**
3b). Pluripotency gene expression was maintained (**Supplementary Fig. 6d**). The
heterozygous deletions did not affect the global transcriptome, as confirmed by a
correlation analysis between all sequenced samples (**Supplementary Fig. 6e**). These
results confirm the successful generation of a high-quality dataset of deletions spanning the
entire X chromosome, suitable for testing the Xuc hypothesis.

As expected, XX cells had higher X-linked gene expression levels compared to XO cells
(**Supplementary Fig. 6f**). However, when gene expression is normalized per X
chromosome, XO cells showed XCU (**Figure 3c**). In contrast, none of the 13 new deletions
induced global full XCU for genes located outside of the deleted regions. At the gene level,
genes upregulated in $X^{Cast}O^{Mus}$ cells remained expressed at lower levels outside the
deletions across all deletion lines compared to $X^{Cast}O^{Mus}$ cells (**Figure 3d**). None of the
deletions caused significant expression changes relative to $X^{Cast}X^{Mus}$, with the exception of
deletion AB. This likely represents an outlier, as deletion of region A or B alone had no
significant effect, and the magnitude of change for AB was smaller than in $X^{Cast}O^{Mus}$ cells. In
summary, deletion of large segments on one of the two X chromosomes is not sufficient to
induce full XCU outside of the deleted regions. Collectively, these results do not support the
hypothesis that any of the segments deleted act as an Xuc with the ability to induce global
XCU of the X chromosome *in trans*.

Cells sense large segmental deletions and compensate gene expression *in trans*
To test if large segmental deletions are compensated within the deleted regions *in trans*, we
examined XCU specifically within deleted segments (**Figure 4a**). Remarkably, there was a
strong induction of XCU specifically within deleted segments *in trans* (**Figure 4b-d**,
Supplementary Fig. 7a-c). Moreover, the level of XCU was similar to that of XO cells in
most cell lines. We made similar observations for the 13 segmental deletions of the X
chromosome, in which XCU was induced within deleted segments *in trans* (**Figure 4e**,
Supplementary Fig. 7d). To quantify XCU at the gene level, we examined expression
levels of genes upregulated in $X^{Cast}O^{Mus}$ cells relative to $X^{Cast}X^{Mus}$. These genes were
consistently upregulated across the deletion lines, reaching expression levels similar to
those in $X^{Cast}O^{Mus}$ cells (**Figure 4f**). Allele-specific analysis confirmed upregulation of the
Cast allele in $X^{Cast}O^{Mus}$ cells (**Supplementary Fig. 7e**). These results demonstrate that
mammalian cells can sense when specific segments of one of the two X chromosomes are
genetically deleted and compensate for gene dosage imbalances by transcriptional
upregulation specifically *in trans*.

Next, we aimed to define if the upregulated genes within the deleted regions in the deleted
lines are the same as the upregulated genes in XO cells. To address this, we asked if genes
upregulated in XO cells were also upregulated when deleted in cells with large
heterozygous deletions. We found that 64 of the 65 upregulated genes located within
deletions 1-3 were also upregulated in each of the deletions *in trans* (**Figure 4g**). Likewise,
genes upregulated in cells with deletions were also upregulated in cells with $X^{monosomy}$
(**Supplementary Fig. 7e**). This high degree of overlap suggests a shared transcriptional
response to X chromosome dosage loss.

These results favor a model in which specific regions or even individual genes can sense
gene dosage and even gene expression states and adapt X-linked gene expression
accordingly by XCU. We investigated whether deletion of a single allele of a gene would
also result in transcriptional dosage compensation of that gene *in trans*. In other words, we
asked if one allele of a gene can sense and respond to deletion of the other allele *in trans*.
We focused on a single gene, *Pdzd4*, that is subject to XCU, and generated mESCs which
contain a heterozygous deletion of the promoter and gene body of *Pdzd4* on the *Mus* allele
(**Supplementary Fig. 7g**). Allele-specific quantitative PCR analysis indicated upregulation
in trans for three of the four clones heterozygous for *Pdzd4* (**Figure 4h**). These results
suggest that *Pdzd4* is transcriptionally compensated *in trans* when one of the two alleles of
Pdzd4 is deleted. We conclude that heterozygous loss of a gene body and corresponding
promoter on the X chromosome can be sensed and compensated via increased
transcription in *trans*.

Autosomal gene dosage compensation by transcriptional upregulation *in trans*
So far we have provided evidence for transcriptional compensation of monosomies of the X
chromosome *in trans*, consistent with the evolution of dosage compensation mechanisms
on the X chromosome by XCU[4–11]. Next, we aimed to determine if dosage compensation by
transcriptional upregulation *in trans* can take place on autosomes. To explore autosomal
gene dosage compensation, we used a bulk RNA-seq dataset of *Mus/Cast* mESCs with
inducible autosomal gene silencing *in cis*[61]. In this system, addition of dox activates an
inducible *Xist* transgene inserted on the *Cast* allele of chromosome 3, resulting in the
transcriptional downregulation of many genes *in cis* on that allele (**Figure 5a,**
Supplementary Fig. 8a)[61]. This experimental setting enabled us to ask if reduction of
autosomal gene expression *in cis* induces dosage compensation *in trans*. We confirmed
reduced gene expression *in cis* and identified a ~80 Mb region with a high density of genes
that decrease expression on the *Cast* allele after *Xist* expression for 24 hours (**Figure 5b-c**).
To assess gene-by-gene compensation, we first examined genes located outside of the 80-
160 Mb region *in trans*, on the *Mus* allele, and did not detect changes in gene expression
(**Figure 5d**). By contrast, inducing reduced gene expression of chromosome 3 *in cis*
induced significant transcriptional upregulation of the corresponding *Mus* allele *in trans* in
the 80-160 Mb region (**Figure 5e**). Contrary to our findings on XCU, the baseline expression
level prior to addition of doxycycline was significantly lower for upregulated genes

compared to non-upregulated genes (**Supplementary Fig. 8b**), potentially due to the fact
that the expression of highly expressed genes is not sufficiently reduced in this
experimental system. Upregulated chromosome 3 genes were not enriched for genes which
encode proteins in complexes, or proteins which form condensates (**Supplementary Fig.**
8c-d). In summary, we found that decreasing expression *in cis* resulted in upregulation *in*
trans for several genes on chromosome 3 after 24 hours of doxycycline. Therefore, these
results suggest that reducing expression of autosomal genes *in cis* can be compensated by
transcriptional upregulation *in trans*.

Gene dosage compensation at the protein level
XCU has been described to occur at the transcriptional level in mouse embryos and ESCs
but its occurrence at the protein level in mouse PSCs remains to be established[16,17,20,62]. To
determine if XCU takes place at the protein level in XO and XY ESCs, we performed
quantitative proteomics in $X^{Cast}X^{Mus}$, $X^{Mus}Y$, and $X^{Cast}O^{Mus}$ ESCs. We also included two ESC
lines with large heterozygous segmental deletions (Deletions E and K), which showed high
percentages of transcriptionally upregulated genes located within the deleted region
(**Supplementary Table 2**). We identified 8,050 proteins, of which 4,446 proteins were
detected in all samples.

We next analyzed X-linked protein levels and found them to be more abundant in $X^{Cast}X^{Mus}$
ESCs compared to $X^{Mus}Y$ and $X^{Cast}O^{Mus}$ cells, in agreement with increased X-linked gene
expression in these cells due to the presence of two active X chromosomes
(**Supplementary Fig. 9a**). Since mass spectrometry results do not provide allelic
resolution, we normalized protein abundances for the X-chromosome copy number. On a
per-X-chromosome basis, XCU was detected at the protein level in $X^{Mus}Y$ and $X^{Cast}O^{Mus}$
ESCs compared with $X^{Cast}X^{Mus}$ ESCs (**Figure 6a**). Indeed, ESCs with one active X
chromosome showed more than half of X-linked protein abundances, suggesting XCU at
the protein level (**Supplementary Fig. 9a**). The increase was also seen for the X:A ratio
(**Supplementary Fig. 9b**). Therefore, we confirm that XCU takes place at the protein level
in $X^{Mus}Y$ and $X^{Cast}O^{Mus}$ ESCs to compensate for the absence of one of the two active X
chromosomes. These results show that the single active X chromosome of mESCs is
upregulated at the protein level compared to each X chromosome in XX mESCs.

We next examined which proteins are subject to upregulation in ESCs with only one active
X chromosome. Focusing on the 159 detected X-linked proteins, 45 proteins (38%) were
significantly upregulated in both $X^{Mus}Y$ and $X^{Cast}O^{Mus}$ cells, indicating that some but not all
expressed proteins are compensated by XCU (**Figure 6b-c, Supplementary Fig. 9c-d,**
Supplementary Table 3). Compared to autosomes, the proportion of upregulated proteins
was highly enriched on the X chromosome (**Figure 6d**). At the protein level, upregulated X-
linked proteins were enriched for proteins that are part of complexes, suggesting evolution
selected dosage compensation mechanisms for these proteins (**Figure 6e**). Taken together,
our data suggest that a subset of X-linked genes are compensated by XCU at the protein
level in $X^{Mus}Y$ and $X^{Cast}O^{Mus}$ mESCs.

To determine if the same genes are compensated at the RNA and protein level, we
integrated the proteomics and transcriptomics data and compared genes only present in
both RNA and protein data. We found that of the 159 coding genes covered in both
datasets, a majority, 103 (60%), were more abundant in XO samples compared to the copy-
number corrected abundance in XX (**Figure 6f**). Specifically, 47 (28%) showed both
increased RNA and protein levels, whereas 56 (35%) genes were increased at the protein
level only, suggesting exclusive post-transcriptional dosage compensation of these genes.
A small proportion of genes were increased at the RNA level only (8%), while 27% of genes
did not show compensation at the RNA or protein level. Despite the limited sensitivity of
proteomics compared to transcriptomics and the commonly reported RNA-protein
discordance[63], we observed a positive correlation in upregulation (r=0.37). These results
indicate that the contribution of RNA and protein regulation to dosage compensation by
XCU may vary for different X-linked genes in mESCs, in agreement with transcription and
protein measurements in the brain[62]. Our results reveal that there are multiple layers of gene
and protein regulation involved in dosage compensation by XCU of individual X-linked
genes in ESCs.

Next, we examined dosage compensation at the protein level in two heterozygous
segmental deletion lines E and K. To assess XCU, we analyzed protein abundance
normalized to copy number, based on information on genetic regions with one or two
copies. Outside the deleted regions, there was no XCU at the protein level (**Figure 6g**,
Supplementary Fig. 9e). In contrast, within the deleted regions, we detected considerable
XCU at the protein level (**Figure 6h**, **Supplementary Fig. 9f**). The upregulation status of
individual proteins that are upregulated in $X^{Mus}Y$ and $X^{Cast}O^{Mus}$ lines was evaluated in the
deleted ESC lines (**Figure 6i**). Proteins encoded outside the deleted regions remained
unchanged, while several proteins within deleted regions were upregulated. In summary,
our results reveal the existence of different dosage compensation mechanisms in mouse
PSCs, with genes that tend to be compensated at the protein, RNA or both levels, and to
varying degrees. Together, we conclude that ESCs can compensate for mono-allelic gene
expression by XCU at both RNA and protein levels.

DISCUSSION
In this study, we investigated dosage compensation and sensing in PSCs. We discovered
that dosage of monosomically expressed X-linked and autosomal regions is compensated
at both the mRNA and protein levels. In addition, we found that dosage compensation does
not require one or more specific regions on the X chromosome involved in dosage sensing
and XCU. In contrast, our findings indicate that XCU is regulated at a gene-by-gene level
and involves upregulation of expression at the transcriptional and protein levels.

These results support previous findings by others and us related to elastic tuning of dosage
compensation with one allele becoming upregulated upon inactivation or loss of the other
allele during development[16,20]. In addition, our studies are in line with reports showing that
XCU is not global, as not all X-linked genes are transcriptionally upregulated[17]. Our data do
indicate that a majority of proteins detected by mass spectrometry display XCU at the

protein level. This finding may involve an over-interpretation of the percentage of genes
subject to XCU as our findings indicate that XCU genes are expressed at a higher level and
will therefore be more easily detected. On the contrary, our study only examined PSCs and
different cell types or cell states may require dosage compensation of different genes[17,62,64].
We conclude that the mammalian genome is more adaptive and robust than previously
suggested.

Our studies and others also indicate that different dosage compensation mechanisms or
different combinations of mechanisms are in place to regulate gene dosage compensation.
These mechanisms include transcriptional burst and translation initiation frequency as well
as mRNA and protein stability[19,26,62]. We hypothesize that self-regulating feedback loops
and enhancers may play a role in sensing changes in gene dosage, as inactivation or
deletion of one allele could drive differential transcription factor binding *in trans*. This could
then also be related to the increased transcriptional burst frequency which has been
reported with XCU[16,19,20]. Furthermore, in line with the increased transcriptional burst
frequency reported, we found an increased enrichment of one histone modification,
H4K9ac, associated with active transcription at the transcription start sites of genes subject
to XCU. These results show that H4K9ac is associated with XCU, however, whether this
mark is instructive or sufficient to induce dosage compensation remains to be tested.
Notably, H4K16ac was not significantly enriched indicating that mammalian cells employ
distinct mechanisms compared to *Drosophila* to upregulate expression of X-linked genes.
As our study indicates that XCU is regulated on a gene-by-gene basis, identification of the
mechanisms employed and interactions involved to regulate XCU requires further
investigation at the gene level.

Studies in humans and primates have suggested that XCU also occurs in cells with a single
active X chromosome[27,28,35,62]. We hypothesize that the gene-specific dosage sensing and
subsequent compensation observed in mouse cells also occurs in human stem cells and in
development. Furthermore, it has been suggested that XCU is widespread in placental
mammals[4], so we would expect gene-specific dosage compensation to occur for many
genes on the human X chromosome, however additional studies would be required to
confirm this. Widespread dosage compensation of X-linked genes could explain why
monosomy of the X chromosome (Turner syndrome) is the only viable monosomy in
humans and mice[65,66]. Studying X-linked dosage compensation in early human development
is difficult without allele-resolution data, and due to the low coverage of SNPs between
human alleles, allele-resolution data is typically only available for a few genes.

Our findings have implications for human disorders caused by haploinsufficiency. First,
dosage compensation of autosomes may have evolved to cope with heterozygous
mutations, monoallelic silencing, microdeletions or aneuploidies and their own effects.
Second, dosage compensation may enable survival, but is not sufficient for restoring a
wildtype phenotype and rescuing haploinsufficiency and as such may enable diseased
states.

Consistent with previous research in both human cancers and embryos, our findings
provide strong evidence for dosage compensation of autosomal genes[3,67]. This suggests
that the mammalian genome possesses an inherent ability to compensate for gene dosage
imbalances resulting from genomic loss or decreased expression, a capacity that extends
beyond the X chromosome. This intrinsic dosage compensation mechanism may have been
a key factor in the evolution of X-chromosome inactivation (XCI). As X-linked genes were
progressively lost from the Y chromosome during the evolution of mammalian sex
chromosomes[68], it is hypothesized that increasing selective pressure drove the
development of dosage compensation mechanisms for X-linked genes. This likely led to the
gradual incorporation of X-linked genes into the XCI process[8,34,69]. The gradual nature of this
evolution may explain the gene-by-gene regulation observed in XCU.

In conclusion, our study supports the hypothesis of gene-by-gene regulation of X-linked
gene dosage compensation involving multiple levels of regulation in a locus-specific
manner to sense and compensate for variations in X-linked gene dosage.

METHODS
Generation of cell lines
XX female polymorphic mESCs (129/Sv-Cast/EiJ) with heterozygous deletions located
between *Tfe3* and *Zic3*, *Zic3* and *Dusp9*, as well as between *Dusp9* and *Nr0b1* were
obtained from Song et al., 2019[41]. XO mouse iPSCs used in scRNA-seq were generated by
passaging XX-GFP female mouse iPSCs (129/Sv-Cast/EiJ, obtained from Talon et al.,
2021[16]) to passage 14 before FACS for GFP negative cells (XO iPSCs) and collection for
single-cell RNA-sequencing (see below). XX female iPSCs (129/Sv-Cast/EiJ) were derived
as described in Talon at al., 2021[16]. XY male mESCs (C57BL/6J-Cast/EiJ) were obtained
from Chen et al., 2016[70]. Both ESC and iPSC lines were used in this study, due to
availability and experimental design constraints, however, evidence indicates that mouse
iPSCs and ESCs behave equivalently with respect to X chromosome sensing and XCU[16,20].

For all newly generated lines, gRNAs (**Supplementary Table 4**) were integrated in the
pSpCas9(BB)-2A-Puro V2.0 PX459 vector (Addgene #62988), enabling transient puromycin
selection. Complementary oligonucleotides containing the gRNA sequence and compatible
BbsI overhangs were annealed and ligated in PX459. To generate homology arm vectors,
~500–1000 bp 5' and 3' homology arms were amplified by overhang PCR from mouse
genomic DNA with compatible restriction enzyme sequences. PCR products were purified
from gel using the QIAquick® Gel Extraction Kit (Qiagen, 28506). Homology arm PCR
products were digested with the indicated restriction enzymes and subsequently purified and
ligated into compatible sites in the pCR™2.1-TOPO™ vector (Invitrogen, K450002). A
neomycin resistance cassette flanked by LoxP sites was integrated in-between the 5' and
3' homology arms in an EcoRI restriction site for generating large X-chromosomal deletions.
The cassette was retained in all deletion lines, as Cre-mediated excision was not required
for downstream experiments.
531

Two independent *Rnf12*⁺ᐟ⁻ cell lines were generated in F1 2–1 hybrid (129/Sv-Cast/Ei)
mESCs. Although the heterozygous *Rnf12* mutation was not essential for the deletion
experiments described here, it induces complete skewing upon XCI, which was relevant for
a parallel study. The first *Rnf12*⁺ᐟ⁻ cell line was generated by introducing PX459 vectors
containing gRNAs targeting intron 2 and the 3'UTR. This *Rnf12*⁺ᐟ⁻ cell line served as the
parental cell line for generating deletions A through I and K, as well as the *Rnf12*⁺ᐟ⁻*Pdzd4*⁺ᐟ⁻
lines. Attempts to induce deletion J in this line failed due to chromosomal instability and
subsequent X chromosome loss. Therefore, a second *Rnf12*⁺ᐟ⁻ cell line was generated by
introducing PX459 vectors with alternative gRNAs. This line served as the parental
background for del J and I-TEL.

The first *Rnf12*⁺ᐟ⁻ was generated by transfecting gRNAs by electroporation in Gene Pulser
0.2 cm-gap cuvettes (Biorad, 1652086) at 118 kV, 1200 µF and ∞Ω in a Gene Pulser xCell
electroporation system (Biorad). 24h post-transfection, the medium was refreshed and
supplemented with 1µgmL−1 puromycin (Sigma-Aldrich, P8833) for 36h. Drug-resistant
clones were characterized by PCR spanning the deletion. All other cell lines were generated
by transfecting the plasmids with Lipofectamine 2000 (Invitrogen, 11668019) in 6 × 10^5
mESCs dissociated into single-cells. Each transfection contained 2 µg of the homology arm
vector and 1 µg of PX459 containing gRNAs targeting the 5' or 3' end of the deletion. 24h
post-transfection, the medium was refreshed and supplemented with 1µgmL−1 puromycin
for 48h followed by refreshing with medium supplemented with 285µgmL−1 Geneticin
(G418)(Gibco, 11811-031) for 4-6 days. No Geneticin/neomycin selection was performed for
generating the *Rnf12*⁺ᐟ⁻ and the *Rnf12*⁺ᐟ⁻*Pdzd4*⁺ᐟ⁻ cell lines. Drug-resistant clones were
characterized by PCR targeting genotype-specific length polymorphisms inside and outside
the targeted deletion (**Supplementary Table 5**). CRISPR-Cas9 targeting of X-linked regions
results in a population of cells losing part of the X chromosome extending beyond the
targeted regions. These deletion clones were identified and characterized by PCR (Del AB,
C-G and I-TEL). Deletion clones were identified and characterized by PCR and the deletion
boundaries were determined using allele-specific RNA sequencing data. Another population
of cells lost the entirety of one of the X chromosomes. These XO mESC lines were included
in the bulk RNA-seq analysis as treatment-matched XO controls.

562
Cell culture
Mouse embryonic fibroblasts (MEFs) were cultured in MEF medium [DMEM (Gibco, 10829-
018) supplemented with 10% (v/v) fetal bovine serum (FBS, Gibco, 10500-064), 1% (v/v)
penicillin/streptomycin (P/S, Gibco, 15140122), 1% (v/v) Glutamax (Gibco, 35050038), 1%
(v/v) non-essential amino acids (NEAA, Gibco, 11140035), and 0.008% (v/v) beta-
mercaptoethanol (Gibco, 31350010)]. Mouse iPSCs and mESCs were grown on Mitomycin-
treated MEFs in mESC medium [KnockOut DMEM (Gibco, 10829-018) supplemented with

15% FBS (Gibco, 10500-064), 1% (v/v) penicillin/streptomycin (P/S, Gibco, 15140122), 1%
(v/v) Glutamax (Gibco, 35050038), 1% (v/v) non-essential amino acids (NEAA, Gibco,
11140035), 0.008% (v/v) beta-mercaptoethanol (Gibco, 31350010), and mouse LIF
(1000U/mL, home-made) at 37°C under 5% CO2. *Rnf12+/-*, deletion lines A to K and
matched controls were grown on irradiated MEFs in adjusted mESC medium [DMEM
(Gibco, 11995065), 15% FBS (Capricorn Scientific, FBS-12A), 1% (v/v)
penicillin/streptomycin (Sigma-Aldrich, P0781), 1% (v/v) NEAA (Lonza, BE13-114E),
0.1 mM beta-mercaptoethanol and LIF (5000 U/ml, home-made)]. mESCs and iPSCs were
passaged every two days using trypsin dissociation.

579
Cell sorting
Cells were dissociated using trypsin digestion and subsequently washed in incubation
buffer (1x PBS, 0.5% BSA, 2mM EDTA) and filtered through a Falcon 40um Cell Strainer
(Corning, 352340). 1μg SSEA1-PE antibody (R&D Systems, FAB2155P) was added per 5
million cells. Cells were incubated for 30 min at 4°C. Cell death exclusion was performed by
adding DAPI (Sigma-Aldrich, D9542) prior to sorting. Sorting was performed on a BD
FACSAria Fusion (BD Biosciences) and performed by expert operators at the VIB FACS
core.

588
589 **Single-cell RNA-seq library preparation**
590 Single-cell RNA-seq libraries were prepared by following a previously published protocol
591 (Smartseq3xpress)[44], with minor modifications. Library quality was checked using an Agilent
592 Tapestation as well as an Agilent 2100 Bioanalyzer. Library concentration was checked with
593 a Qubit fluorometer (Qubit dsDNA HS Assay).

594
595 **Single-cell RNA-seq sequencing**
596 Single-cell RNA-seq libraries were sequenced in paired-end mode (150bp) by BGI
597 Genomics (Shenzhen, China) on a DNBSEQ-G400 sequencer (MGI Tech).

598
Single-cell RNA-seq analysis
Single-cell RNA-seq data was obtained as fastq files from BGI Genomics. The zUMIs[71]
pipeline was used to map reads using STAR[72] (v2.7.1a) with the options '--
clip3pAdapterSeq CTGTCTCTTATACACATCT'. Reads containing barcodes which
corresponded to XX, XO, Deletion 1, Deletion 2, or Deletion 3 cells were mapped to an
GRC38.p6 mouse reference genome N-masked for SNPs between the 129/SvJ and Cast
strains and SNPs between Bl6 and 129. Reads containing barcodes which corresponded to
XY cells were mapped to an GRC38.p6 mouse reference genome N-masked for SNPs
between the Bl6 and Cast strains. Reads were filtered such that reads containing at least 3
bases with Phred quality score less than 20 were omitted. Allele-resolution counts were
generated using a publicly available script 'get_variant_overlap_CAST.R'
(https://github.com/sandberg-lab/Smart-seq3/)[73] and a custom list of SNPs between

129/SvJ and Cast, with overlapping SNPs between 129/SvJ and Bl6, as well as overlapping
SNPs between Cast and Bl6 omitted. Allelic exon UMI counts were used to generate Seurat
[74] (v4.2.1) objects, which were then filtered such that cells with less than 10,000 UMIs for
either allele were omitted. After filtering, we obtained 69 $X^{Cast}X^{Mus}$, 44 $X^{Cast}O^{Mus}$, 6 $X^{Mus}O^{Cast}$,
61 $X^{Mus}Y$, 52 Deletion 1, 51 Deletion 2, and 41 Deletion 3 cells. Read counts were
normalized to TPM (transcripts per million). Normalized allele resolution UMI counts were
used for all analysis on the scRNA-seq dataset. Mean normalized gene expression was
calculated for genes between the 10th and 90th percentile of expression per cell for each
gene set of interest (all X-linked genes, inside/outside each deletion, inside/outside
inactivated region), and for all (other) autosomal genes. X-to-autosome ratio was calculated
by dividing the mean expression of genes in each subset by the median expression of
autosomal genes.

Characterization of X-chromosome status
The mean expression of X-linked genes from each allele was considered for each cell. If the
mean allelic expression was greater than two for the *Cast* allele and less than two for the
Mus allele, then the cell was defined as $X^{Cast}O^{Mus}$, and vice versa. One cell was omitted due
to ambiguous X-status.

Characterization of pluripotency status
The pluripotency marker genes *Essrb*, *Nanog*, and *Pou5f1* were used to assess
pluripotency status. High expression levels ($\log_2$(TPM) > 2.5) of each marker gene was
considered to be sufficient to conclude the cells had not differentiated and remained
pluripotent.

Identification of XCU genes
The expression values of each X-linked genes from the active allele of cells with a single
active X chromosome (X^{Cast} expression from $X^{Cast}O^{Mus}$, X^{Mus} expression from $X^{Mus}O^{Cast}$ and
$X^{Mus}Y$) was tested to be greater than half of the total allelic expression from XX cells via
Student's T.test. Genes with $p < 0.05$ and fold change > 1.2 were defined as being
transcriptionally upregulated.

Characteristics of XCU genes
A list of known escapee genes was downloaded from Berletch *et al.* 2015[56] and overlapped
with the list of upregulated genes in $X^{monosomy}$ and non-upregulated genes. The EMBL-EBI
protein complex database[75] was used to compare the proportion of genes which code for
proteins in complexes between upregulated and non-upregulated genes. The CD-CODE
database was used to determine the proportion of transcriptionally upregulated X-linked
genes which code for proteins that form condensates[76]. Human orthologs of upregulated X-
linked genes were identified using the BioMart tool from Ensembl[77]. This list of orthologs
was then overlapped with a list of known disease causing genes from the OMIM

database[78]. Significant differences between the lists of genes were calculated using a
Student's t-test when comparing expression and a Chi-square test when comparing
proportions.

The lists of upregulated and non-upregulated genes were examined for gene ontology (GO)
biological processes (BP) and protein-protein interaction (PPI) enrichment using Enrichr[79].
Significant terms (p-value < 0.05) were selected for each database and gene list, and the
top 10 terms for each category were visualized in barplots.

Bulk RNA-seq
Cells were cultured at least two passages before harvesting. mESCs were depleted of MEFs
by pre-plating dissociated cells for 45 minutes at 37°C in non-gelatinized plates. The
supernatant containing mESCs was spun down for 5 min at 300 × g. Cell pellets were lysed
and RNA was extracted using the ReliaPrep™ RNA Miniprep kit (Promega, Z6012) with
increased DNase treatment (25 min at room temperature). MEF depletion was validated on
the genomic DNA of part of the cell pellet using PCRs targeting genotype-specific length
polymorphisms and SNPs.

Sequencing libraries were prepared using the Truseq stranded mRNA library preparation
method from Illumina. The libraries of the XO cells were subsequently sequenced on an
Illumina NextSeq2000 sequencer resulting in single read clusters of 50 bases in length. The
libraries of all other cell lines were sequenced on a HiSeq2500 sequencer resulting in
paired-end clusters of 50 bases.

Bulk RNA-seq analysis
The SNPs in the 129/Sv and Cast/Ei lines were downloaded from the Sanger Institute (v.5
SNP142)[80]. These were used as input for SNPsplit[81] (v0.3.4), to construct an N-masked
reference genome based on mm10 in which all SNPs between 129/Sv and Cast/Ei are
masked. The reads were trimmed and aligned to the N-masked reference genome using
TrimGalore[82] (v0.6.7) and HISAT2[83] (v2.2.1), respectively. SNPsplit was then used to assign
the reads to either the 129/Sv or Cast/Ei BAM file based on the best alignment or to an
unassigned BAM file if mapping to a region without allele-specific SNPs (--paired for the
paired-end samples). The allele-specific and unassigned BAM files were merged into a
composite BAM file using Samtools[84] (v1.10). The number of mapped reads per gene was
counted for both alleles separately and for the composite BAM file using featureCounts[85]
(v2.0.6) based on the gene annotation from Ensembl v98 (-t exon -s 2). The raw counts
were normalized to transcripts per million (TPM).

Validation of bulk RNA-seq samples
All bulk RNA-seq samples were generated in a *Rnf12*$^{+/-}$ parental cell line. To assess a
potential bias, we performed differential expression analysis between WT XX and *Rnf12*$^{+/-}$
XX mESCs using DESeq2[86] (v1.30.1) with default settings. Results were plotted in a scatter
plot showing mean TPM in both cell lines, in which all differential expressed genes

693 (adjusted p-value < 0.05, $\log_2$(fold change) < -1 or > 1) are highlighted. The deleted regions
694 in all samples were validated by calculating the ratio of each sample to the mean of the
695 parental $Rnf12^{+/-}$ samples. One deletion A clone and both deletion G clones were excluded
696 due to a suspected recombination event rather than deletion. To ensure comparability
697 across the remaining samples, we assessed the expression levels of the pluripotency genes
698 *Essrb*, *Nanog* and *Pou5f1* by plotting their TPM values for all samples. Additionally, global
699 expression patterns were evaluated by calculating the Pearson correlation coefficients for
700 TPMs of all autosomal genes, which were visualized in a heatmap.

702 **XCU in bulk RNA-seq data**
703 To maximize statistical power, we analyzed non-allelic TPMs of all X-linked genes within
704 and outside each deletion region. Only expressed genes were selected by filtering for TPM
705 > 5 in $Rnf12^{+/-}$ samples. Deletion I was excluded from the plots showing expression within
706 each deletion region due to insufficient expressed genes. For each genomic region, mean
707 TPM values were calculated. These levels were compared to the mean XX level (from
708 $Rnf12^{+/-}$) and 50% of the mean XX level, representing the expected expression level for a
709 single X chromosome without upregulation.

710 To account for copy number differences, gene expression was normalized to copy number-
711 corrected values by dividing TPM values by the number of copies present, based on the
712 genomic locations of each deletion. Mean copy number-corrected TPMs were then
713 calculated for genes within and outside deletion regions and compared to the mean copy
714 number-corrected TPM in XX samples.

715 To validate allelic upregulation detected in non-allelic TPM values, we selected X-linked
716 genes with a fold change > 1.2 in copy number-corrected TPMs between XO and XX cells.
717 For this subset of upregulated genes, allelic *Cast* expression data was filtered, and the $\log_2$
718 fold change between XO and XX cells was calculated. The allelic upregulation status in XO
719 cells for each gene, annotated by overlapping deletion region, was visualized in a heatmap.
720 The level of upregulation in the deletion samples was assessed by analyzing the copy
721 number-corrected expression of genes upregulated in XO cells. For each gene, the $\log_2$ fold
722 change relative to the mean expression in XX cells was calculated. These values were
723 plotted separately for genes located inside and outside each deletion, with median values
724 per sample indicated. For visualization, extreme values were clipped at -1 and 4. Statistical
725 significance was assessed using two-sided paired t-tests on the mean expression per
726 genotype, with Bonferroni correction for multiple comparisons. No statistical testing was
727 performed for genes located within the deletions due to the limited number of genes per
728 deletion region.

730 **Autosomal upregulation analysis**
731 We reanalyzed a bulk RNA-seq dataset from GSE119602 using the same approach as
732 described in 'Bulk RNA-seq analysis'. Inactive genes were defined as genes whose fold
733 change was less than 0.15 when comparing samples with doxycycline to samples without

doxycycline. The inactive region was defined as the region from 80Mb to the 3' telomere.
Upregulated genes were defined as genes with a fold change greater than 0.5 when
comparing samples with doxycycline to samples without doxycycline. Characteristics of the
upregulated genes were evaluated as described in 'Characteristics of XCU genes'. Mean
normalized gene expression within/outside the inactive region for each allele was calculated
using genes with expression between the 5th and 95th percentile for each sample.

RT-qPCR
mESCs were pre-plated as described. Cell pellets were lysed and RNA was extracted using
the ReliaPrep™ RNA Miniprep kit (Promega, Z6012) with increased DNase treatment (25
min at room temperature). 500 ng RNA was reverse transcribed using Superscript III
(Invitrogen, 18080093) and random hexamers (Invitrogen, N8080127) with the addition of
RNaseOUT (Invitrogen, 10777019). cDNAs were diluted 1:1 in water. RT-qPCRs were
performed in triplicate in the GoTaq qPCR Master Mix (Promega, A6002) in a CFX384
Touch Real-Time PCR Detection System (BioRad, LJB22YE8Z). *Rplp0* and *H2afz* were
used as normalization controls. Allele-specific primer pairs were optimized using pure
gDNA of 129/Sv and Cast/Ei mice. All primers are included in (**Supplementary Table 6**).

Proteomics
Proteomics experiments were performed on $X^{Cast}X^{Mus}$ (*Rnf12*$^{+/-}$), $X^{Mus}Y$ (F1 2-3), and $X^{Cast}O^{Mus}$
(*Rnf12*$^{+}$) ESCs. Additionally, two deletion cell lines (Deletion E and K in *Rnf12*$^{+/-}$) were
included, selected based on their single-region deletions and a high proportion of
transcriptionally upregulated genes.

For global proteome analysis of whole cell extracts, cells were lysed in 100 mM Tris/HCl,
pH 8.2, containing 1 % sodium deoxycholate (SDC) using sonication in a Bioruptor Pico
(Diagenode). Protein concentrations were measured using the BCA assay (ThermoFisher
Scientific). 100 µg protein was reduced in lysis buffer with 5 mM dithiothreitol and alkylated
with 10 mM iodoacetamide. Next, proteins were digested with 2.5 µg trypsin (1:40
enzyme:substrate ratio) overnight at 37 °C. After digestion, peptides were acidified with
trifluoroacetic acid (TFA) to a final concentration of 0.5 % and centrifuged at 10,000 g for 10
min to spin down the precipitated SDC. Peptides in the supernatant were desalted on a 50
mg C18 Sep-Pak Vac cartridge (Waters). After washing the cartridge with 0.1 % TFA,
peptides were eluted with 50 % acetonitrile and dried in a Speedvac centrifuge. Peptides
were then analyzed by nanoflow LC-MS/MS as described below.

Nanoflow LC-MS/MS was performed on an Vanquish Neo LC system coupled to an
Orbitrap Exploris 480 mass spectrometer (both ThermoFisher Scientific), operating in
positive mode and equipped with a nanospray source. Peptide mixtures were trapped on a
PepMap trapping column (2 cm × 100 µm, Thermo, 164750) at a flow rate of 1 µl/min.
Peptide separation was performed on ReproSil C18 reversed phase column (Dr Maisch
GmbH; column dimensions 25 cm × 75 µm, packed in-house) using a linear gradient from 0
to 80% B (A = 0.1% FA; B = 80% (v/v) AcN, 0.1 % FA) in 120 min and at a constant flow

rate of 250 nl/min. The column eluent was directly sprayed into the ESI source of the mass
spectrometer.

For data independent acquisition (DIA), all spectra were recorded at a resolution of 120,000
for full scans in the scan range from 350–1100 m/z. The maximum injection time was set to
50 ms (AGC target: 4E5). For MS2 acquisition, the mass range was set to 336–1391 m/z
with variable isolation windows ranging from 7–82 m/z with a window overlap of 1 m/z. The
orbitrap resolution for MS2 scans was set to 30,000. The maximum injection time was at 54
ms (AGC target: 5E4; normalized AGC target: 100 %).

784 Proteomics analysis
DIA raw data files were analyzed with the Spectronaut Pulsar X software package (version
17.0.221202, www.biognosys.com), using directDIA for DIA analysis including MaxLFQ as
the LFQ method and Spectronaut's IDPicker algorithm for protein inference. The Q-value
cutoff at precursor and protein level was set to 0.01. All imputation of missing values was
disabled.

Proteins detected at least once in each sample were retained for further analysis. The mean
abundance of all X-linked proteins in each XX, XO and XY sample was plotted with
confidence intervals using seaborn pointplot. To avoid a detection bias of low-abundance
proteins between X-linked (1n in XO and XY cells) and autosomal proteins (2n), X:A ratios
were calculated using only proteins with a mean abundance greater than 12,000,000 in the
XX samples (2n for all proteins). The X:A ratio was defined as the abundance of each X-
linked protein divided by the mean autosomal abundance in the same sample, and
visualized using seaborn pointplot.

Since proteomics data lacks allele-specific resolution, we corrected the data for the copy
number to identify significantly upregulated proteins. The abundances in XX samples were
divided by 2 to account for two alleles, while abundances of XO and XY samples remained
unchanged. Moreover, autosomal abundances were divided by 2 in all samples. Differential
expression analyses were performed between XO and XX and between XY and XX using a
Tuckey's HSD test. Upregulated proteins were selected based on the criteria of $\log_2$(fold
change) > 0 and p-value < 0.05 in both XY and XO comparisons. The results for all X-linked
proteins were visualized using volcano plots, and the number of upregulated proteins per
chromosome were displayed in a barplot. To assess the overlap in upregulation between
XO and XY cells, gene lists identified in each analysis were compared using a Venn
diagram. In addition, a scatter plot was generated to visualize the correlation between cell
lines, showing the $\log_2$ fold change in copy number-corrected abundances for XO vs XX
and XY vs XX. The correlation between transcriptomic and proteomic upregulation was
explored using a scatter plot showing the $\log_2$ fold change of the copy number-corrected
abundances in XO versus XX and the copy number-corrected transcript expression in XO
versus XX.

Samples with deletions E and K were analyzed by labeling proteins according to the
genomic location of the gene encoding the protein (inside or outside the deleted region).
For each region separately, the mean normalized protein abundance was plotted,
comparing levels outside and inside the deletion to the XX reference level and 50% of the
XX reference level, respectively. Abundances were copy number-corrected by halving
abundances for two-copy proteins. These corrected abundances were visualized in a
heatmap, focusing on proteins significantly upregulated in XO and XY samples.

ChIP-seq
mESCs were depleted of MEFs 16 hours before the harvest by pre-plating dissociated cells
for 45 minutes at 37°C in non-gelatinized plates. In total, 1×10^8 cells were crosslinked by
the addition of 1% formaldehyde (Sigma-Aldrich, F1635-25ML) to the dish and incubated at
room temperature for 10 minutes on a rocking platform. Formaldehyde was quenched by
adding 125 mM glycine for 5 minutes. Subsequently, cells were washed once with $1 \times$ PBS
before scraping in $1 \times$ PBS supplemented with $1\times$ protease inhibitors (Roche,
11836170001). Cells were pelleted, resuspended and incubated for 10 minutes at 4°C in 1
$\times$ LB1 (50 mM HEPES pH 8.0, 140 mM NaCL, 1mM EDTA, 10% glycerol, 0.5% NP-40,
0.25% Triton X-100, $1\times$ protease inhibitors), in $1 \times$ LB2 (10 mM Tris-HCl pH 8.0, 200 mM
NaCl, 1 mM EDTA, 0.5 mM EGTA,$1\times$ protease inhibitors) and in $1\times$ LB3 (10 mM Tris-HCl pH
8.0, 100 mM NaCl, 1 mM EDTA, 0.5 mM EGTA, 0.1% Sodium Deoxycholate, 0.5% N-
laurolylsarcosine, $1\times$ protease inhibitors). DNA was sonicated to an average fragment
length of 100-300 bp using a Bioruptor Pico (Diagenode, B01060001) for 10 cycles of 30
sec on and 30 sec off. 1/20th volume of 20% Triton X-100 was added to the sonicated
chromatin, and incubated for 10 minutes at 4°C before centrifugation for 10 min at 4500 × g
10 at 4°C to remove debris. 10 µl chromatin was used to measure the chromatin
concentration by nanodrop and check the sonication efficiency. Sheared chromatin was
flash frozen in liquid nitrogen and stored at -80°C until further use. For H4K16ac ChIPs, 30
µg of chromatin was diluted to 500 µl in ChIP dilution buffer (16.7 mM Tris-HCl pH 8.0, 167
mM NaCl, 1.2 mM EDTA, 1.1% Triton X-100, 0.01% SDS, $1\times$ protease inhibitors) and
incubated while rotating overnight at 4°C with 2 µg H4K16ac antibody (Sigma-Aldrich, 07-
329, lot: 3772263) or IgG (Sigma-Aldrich, 12-370). 30 µl of Magnetic beads (Life
Technologies, 10004D) were washed once in ChIP dilution buffer, added to each sample
and incubated while rotating for 4 hours at 4°C. Beads were washed 3× in Low salt buffer
(20 mM Tris-HCl pH 8.0, 150 mM NaCl, 2 mM EDTA, 1% Triton X-100, 0.1% SDS, 1×
protease inhibitors), 1× in High salt buffer (20 mM Tris-HCl pH 8.0, 500 mM NaCl, 2 mM
EDTA, 1% Triton X-100, 0.1% SDS, 1× protease inhibitors), 1× in LiCl buffer (10 mM Tris-
HCl pH 8.0, 250 mM LiCl, 1 mM EDTA, 1% NP-40, 1% Sodium Deoxycholate, 1× protease
inhibitors) and rinsed once in TE (10 mM Tris-HCl pH 8.0, 1 mM EDTA), 10min rotating at
4°C each wash. Chromatin was eluted in 200µL Elution Buffer (1% SDS, 0.1M NaHCO3)

with 5µL protease K (10 mgmL−1) and 2µL RNAse A (10 mgmL−1) while shaking at
1,000 rpm on an orbital thermal-mixer for 2h at 37°C followed by 65°C overnight. DNA was
purified using the Zymo ChIP DNA clean concentrator kit (Zymo Research, D5202)
according to the manufacturer's instructions and eluted in 15µL H2O. 0.1 µL DNA was used
per qPCR to validate ChIP enrichment.

The ChIP-seq libraries were prepared with the ChIPThruPLEX method (Takara Bio,
R400675). These libraries were subsequently sequenced on an Illumina NextSeq2000
sequencer. Paired-end clusters were generated of 50 bases in length.
861

ChIP-seq analysis
The ChIP-seq samples were analyzed allele-specifically similar to the bulk RNA-seq
analysis. The reads were trimmed using Trim Galore and mapped to the N-masked
reference genome using Bowtie2[87] (v2.5.0). Bam files were split by allele using SNPsplit and
the resulting allele-specific bam files were normalized to counts per ten million (CP10M)
using the bamCoverage tool (-bs 1 --normalizeUsing CPM) from deepTools[88] (v3.5.5).
Moreover, allele-specific tracks from GSE116480 were downloaded, including CP10M-
normalized data for H3K17me3, H3K27ac, H4ac, H2AK119Ub, H3K4me1, H3K4me3 and
H3K9ac after 0, 4, 12 and 24 hours of doxycycline.

Enrichment of all ChIP-seq tracks was plotted for each allele separately on the genomic
regions of upregulated and not upregulated genes, selected based on the scRNA-seq
analysis. Mean enrichment per bin was calculated using deepTools computeMatrix (scale-
regions -bs 100 --averageTypeBins mean --missingDataAsZero -m 3000 --upstream 1500 -
-downstream 1500). Profile plots were generated using custom Python (v3.9.16) scripts.
The matrix file from deepTools was loaded, and average enrichment profiles per bin were
plotted using seaborn line plot, including standard error bands.

To evaluate temporal patterns and assess statistical significance, we calculated the mean
signal in two genomic regions: (1) the promoter region, defined as 1 kb upstream of the TSS
to the TSS, and (2) the gene body, defined as the region spanning from TSS to TES. For
each region, bed files were generated containing the corresponding genomic coordinates
for all genes. These bed files were used as input for deepTools multiBigwigSummary, along
with all allele-specific bigwig files, to calculate the average scores in these regions.
Changes over time were visualized with line plots, showing the mean CP10M score with
standard errors for upregulated and not upregulated genes separately.

Statistical testing was performed for each gene set separately. Significant differences
between 0 and 24 hours were calculated using two-sided paired t-tests, with P-values
corrected for multiple testing with the Benjamini-Hochberg approach. These adjusted P-
values were added next to each gene set in the ChIP-seq line plots. Additionally,
differences between upregulated and not upregulated genes were tested using two-sided
Student's t-tests, also corrected for multiple testing. The resulting significance was added

above the corresponding regions in the profile plots.

Genome browser overviews were created to show ChIP-seq enrichment on the active
upregulated allele at the loci of three upregulated genes. Tracks were selected based on
significant differences in enrichment between 0 and 24 hours, as well as between
upregulated and non-upregulated genes at 24 hours. CP10M-normalized allele-specific
tracks for the significantly enriched marks were visualized using IGV v2.19.4[89].
898
899   **Statistics and reproducibility**
900   No statistical method was used to predetermine sample size. One cell was excluded from
901   the scRNA-seq analysis due to ambiguous allelic expression from the X chromosome. One
902   deletion A clone and both deletion G clones were excluded from RNA-seq analysis due to
903   suspected recombination events or partial XO genotype, rather than the intended deletion.

904
DATA AVAILABILITY
The RNA-seq and ChIP-seq data generated in this study have been deposited in the NCBI
Gene Expression Omnibus (GEO) database under accession code GSE282792
(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE282792). Moreover, RNA-seq
and ChIP-seq datasets from GSE119602 and GSE116480 were reanalyzed.
The mass spectrometry proteomics data have been deposited to the ProteomeXchange
Consortium via the PRIDE partner repository with the dataset identifier PXD057399
(https://proteomecentral.proteomexchange.org/cgi/GetDataset?ID=PXD057399).
Source data are provided with this paper.

914
915   **CODE AVAILABILITY**
916   Code is available on the Pasque Lab GitHub account and can be accessed via the following
917   link: https://github.com/pasquelab/Mice-XCU   (ref.   )

REFERENCES

1. Maloy, S. Diploidy. in *Brenner's Encyclopedia of Genetics* 322–323 (Elsevier, 2013).
2. Storchova, Z. *Aneuploidy in Health and Disease.* (BoD – Books on Demand, 2012).
3. De Rossi, S. & Bianchi, F. *Aneuploidy: Etiology, Disorders, and Risk Factors.* (Nova Science Publishers, 2012).
4. Cecalev, D., Viçoso, B. & Galupa, R. Compensation of gene dosage on the mammalian X. *Development* **151**, (2024).
5. Disteche, C. M. Dosage compensation of the sex chromosomes and autosomes. *Semin. Cell Dev. Biol.* **56**, 9–18 (2016).
6. Brockdorff, N. & Turner, B. M. Dosage compensation in mammals. *Cold Spring Harb. Perspect. Biol.* **7**, a019406 (2015).
7. Gu, L. & Walters, J. R. Evolution of Sex Chromosome Dosage Compensation in Animals: A Beautiful Theory, Undermined by Facts and Bedeviled by Details. *Genome Biol. Evol.* **9**, 2461–2476 (2017).
8. Gribnau, J. & Grootegoed, J. A. Origin and evolution of X chromosome inactivation. *Curr. Opin. Cell Biol.* **24**, 397–404 (2012).
9. Ohno, S. *Sex Chromosomes and Sex-Linked Genes.* (Springer Verlag, 1967).
10. Graves, J. A., Disteche, C. M. & Toder, R. Gene dosage in the evolution and function of mammalian sex chromosomes. *Cytogenet. Cell Genet.* **80**, 94–103 (1998).
11. Beukeboom, L. W. & Perrin, N. *The Evolution of Sex Determination.* (Oxford University Press, USA, 2014).
12. Mattimoe, T. & Payer, B. The compleX balancing act of controlling X-chromosome dosage and how it impacts mammalian germline development. *Biochem. J* **480**, 521–537 (2023).
13. Lyon, M. F. Gene action in the X-chromosome of the mouse (Mus musculus L.). *Nature* **190**, 372–373 (1961).
14. Brockdorff, N. *et al.* Conservation of position and exclusive expression of mouse Xist from the inactive X chromosome. *Nature* **351**, 329–331 (1991).
15. Penny, G. D., Kay, G. F., Sheardown, S. A., Rastan, S. & Brockdorff, N. Requirement for Xist in X chromosome inactivation. *Nature* **379**, 131–137 (1996).
16. Talon, I. *et al.* Enhanced chromatin accessibility contributes to X chromosome dosage compensation in mammals. *Genome Biol* **22**, 302 (2021).
17. Naik, H. C., Hari, K., Chandel, D., Jolly, M. K. & Gayen, S. Single-cell analysis reveals X upregulation is not global in pre-gastrulation embryos. *iScience* **25**, 104465 (2022).
18. Naik, H. C. *et al.* Lineage-specific dynamics of erasure of X-upregulation during inactive-X reactivation. *bioRxiv* 2020.12.23.424181 (2023) doi:10.1101/2020.12.23.424181.
19. Larsson, A. J. M., Coucoravas, C., Sandberg, R. & Reinius, B. X-chromosome upregulation is driven by increased burst frequency. *Nat. Struct. Mol. Biol.* **26**, 963–969 (2019).
20. Lentini, A. *et al.* Elastic dosage compensation by X-chromosome upregulation. *Nat Commun* **13**, 1854 (2022).
21. Deng, X. *et al.* Evidence for compensatory upregulation of expressed X-linked genes in

mammals, Caenorhabditis elegans and Drosophila melanogaster. *Nat. Genet.* **43**,
1179–1185 (2011).

22. Lin, H. *et al.* Relative overexpression of X-linked genes in mouse embryonic stem cells
is consistent with Ohno's hypothesis. *Nat. Genet.* **43**, 1169–70; author reply 1171–2
(2011).

23. Li, X. *et al.* Dosage compensation in the process of inactivation/reactivation during
both germ cell development and early embryogenesis in mouse. *Sci. Rep.* **7**, 3729
(2017).

24. Sangrithi, M. N. *et al.* Non-Canonical and Sexually Dimorphic X Dosage Compensation
States in the Mouse and Human Germline. *Dev. Cell* **40**, 289–301.e3 (2017).

25. Yildirim, E., Sadreyev, R. I., Pinter, S. F. & Lee, J. T. X-chromosome hyperactivation in
mammals via nonlinear relationships between chromatin states and transcription. *Nat*
Struct Mol Biol **19**, 56–61 (2011).

26. Rücklé, C. *et al.* RNA stability controlled by mA methylation contributes to X-to-
autosome dosage compensation in mammals. *Nat. Struct. Mol. Biol.* **30**, 1207–1215
(2023).

27. Okamoto, I. *et al.* The X chromosome dosage compensation program during the
development of cynomolgus monkeys. *Science* **374**, eabd8887 (2021).

28. Okamoto, I. *et al.* Eutherian mammals use diverse strategies to initiate X-chromosome
inactivation during development. *Nature* **472**, 370–374 (2011).

29. Deng, X. *et al.* Mammalian X upregulation is associated with enhanced transcription
initiation, RNA half-life, and MOF-mediated H4K16 acetylation. *Dev. Cell* **25**, 55–68
(2013).

30. Gupta, V. *et al.* Global analysis of X-chromosome dosage compensation. *J. Biol.* **5**, 3
(2006).

31. Lin, H. *et al.* Dosage compensation in the mouse balances up-regulation and silencing
of X-linked genes. *PLoS Biol.* **5**, e326 (2007).

32. Johnston, C. M. *et al.* Large-scale population study of human cell lines indicates that
dosage compensation is virtually complete. *PLoS Genet.* **4**, e9 (2008).

33. Moreira de Mello, J. C., Fernandes, G. R., Vibranovski, M. D. & Pereira, L. V. Early X
chromosome inactivation during human preimplantation development revealed by
single-cell RNA-sequencing. *Sci. Rep.* **7**, 10794 (2017).

34. Nguyen, D. K. & Disteche, C. M. Dosage compensation of the active X chromosome in
mammals. *Nat. Genet.* **38**, 47–53 (2006).

35. Fernandez Gallardo, E. *et al.* A multi-omics genome-and-transcriptome single-cell atlas
of human preimplantation embryogenesis reveals the cellular and molecular impact of
chromosome instability. (2023) doi:10.1101/2023.03.08.530586.

36. Vanneste, E. *et al.* Chromosome instability is common in human cleavage-stage
embryos. *Nat. Med.* **15**, 577–583 (2009).

37. Mantzouratou, A. & Delhanty, J. D. A. Aneuploidy in the human cleavage stage embryo.
Cytogenet. Genome Res. **133**, 141–148 (2011).

38. Kojima, S. & Cimini, D. Aneuploidy and gene expression: is there dosage
compensation? *Epigenomics* **11**, 1827–1837 (2019).

39.   Bravo-Estupiñan, D. M. *et al.* Gene dosage compensation: Origins, criteria to identify
compensated genes, and mechanisms including sensor loops as an emerging
systems-level property in cancer. *Cancer Med.* **12**, 22130–22155 (2023).
40.   Schukken, K. M. & Sheltzer, J. M. Extensive protein dosage compensation in aneuploid
human cancers. *Genome Res.* **32**, 1254–1270 (2022).
41.   Song, J. *et al.* X-Chromosome Dosage Modulates Multiple Molecular and Cellular
Properties of Mouse Pluripotent Stem Cells Independently of Global DNA Methylation
Levels. *Stem Cell Reports* **12**, 333–350 (2019).
42.   Schulz, E. G. *et al.* The two active X chromosomes in female ESCs block exit from the
pluripotent state by modulating the ESC signaling network. *Cell Stem Cell* **14**, 203–216
(2014).
43.   Choi, J. *et al.* DUSP9 Modulates DNA Hypomethylation in Female Mouse Pluripotent
Stem Cells. *Cell Stem Cell* **20**, 706–719.e7 (2017).
44.   Hagemann-Jensen, M., Ziegenhain, C. & Sandberg, R. Scalable single-cell RNA
sequencing from full transcripts with Smart-seq3xpress. *Nat Biotechnol* **40**, 1452–1457
(2022).
45.   Keane, T. M. *et al.* Mouse genomic variation and its effect on phenotypes and gene
regulation. *Nature* **477**, 289–294 (2011).
46.   Doran, A. G. *et al.* Deep genome sequencing and variation analysis of 13 inbred mouse
strains defines candidate phenotypic alleles, private variation and homozygous
truncating mutations. *Genome Biol* **17**, 167 (2016).
47.   Pasque, V. *et al.* X Chromosome Dosage Influences DNA Methylation Dynamics during
Reprogramming to Mouse iPSCs. *Stem Cell Reports* **10**, 1537–1550 (2018).
48.   Zvetkova, I. *et al.* Global hypomethylation of the genome in XX embryonic stem cells.
Nat Genet **37**, 1274–1279 (2005).
49.   Janiszewski, A. *et al.* Dynamic reversal of random X-Chromosome inactivation during
iPSC reprogramming. *Genome Res.* **29**, 1659–1672 (2019).
50.   Bird, K. A., Pires, J. C., VanBuren, R., Xiong, Z. & Edger, P. P. Dosage-sensitivity
shapes how genes transcriptionally respond to allopolyploidy and homoeologous
exchange in resynthesized Brassica napus. *Genetics* **225**, (2023).
51.   Birchler, J. A. & Veitia, R. A. The gene balance hypothesis: implications for gene
regulation, quantitative traits and evolution. *New Phytol.* **186**, 54–62 (2010).
52.   Xie, Z. *et al.* Gene Set Knowledge Discovery with Enrichr. *Curr Protoc* **1**, e90 (2021).
53.   Pessia, E., Makino, T., Bailly-Bechet, M., McLysaght, A. & Marais, G. A. B. Mammalian
X chromosome inactivation evolved as a dosage-compensation mechanism for
dosage-sensitive genes on the X chromosome. *Proc. Natl. Acad. Sci. U. S. A.* **109**,
5346–5351 (2012).
54.   Yang, L. *et al.* Phase separation as a possible mechanism for dosage sensitivity.
Genome Biol. **25**, 17 (2024).
55.   Stortz, M., Presman, D. M. & Levi, V. Transcriptional condensates: a blessing or a curse
for gene regulation? *Commun Biol* **7**, 187 (2024).
56.   Berletch, J. B. *et al.* Identification of genes escaping X inactivation by allelic expression
analysis in a novel hybrid mouse model. *Data Brief* **5**, 761–769 (2015).

57. Żylicz, J. J. *et al.* The Implication of Early Chromatin Changes in X Chromosome
Inactivation. *Cell* **176**, 182–197.e23 (2019).

58. Gelbart, M. E., Larschan, E., Peng, S., Park, P. J. & Kuroda, M. I. Drosophila MSL
complex globally acetylates H4K16 on the male X chromosome for dosage
compensation. *Nat Struct Mol Biol* **16**, 825–832 (2009).

59. Galupa, R. & Heard, E. X-Chromosome Inactivation: A Crossroads Between
Chromosome Architecture and Gene Regulation. *Annu Rev Genet* **52**, 535–566 (2018).

60. Dixon, J. R. *et al.* Topological domains in mammalian genomes identified by analysis of
chromatin interactions. *Nature* **485**, 376–380 (2012).

61. Nesterova, T. B. *et al.* Systematic allelic analysis defines the interplay of key pathways
in X chromosome inactivation. *Nat. Commun.* **10**, 3129 (2019).

62. Wang, Z.-Y. *et al.* Transcriptome and translatome co-evolution in mammals. *Nature*
588, 642–647 (2020).

63. Liu, Y., Beyer, A. & Aebersold, R. On the Dependency of Cellular Protein Levels on
mRNA Abundance. *Cell* **165**, 535–550 (2016).

64. Lyu, Q. *et al.* A small proportion of X-linked genes contribute to X chromosome
upregulation in early embryos via BRD4-mediated transcriptional activation. *Curr. Biol.*
32, 4397–4410.e5 (2022).

65. Yang, S. Diagnostic and therapeutic considerations in Turner syndrome. *Ann Pediatr*
Endocrinol Metab **22**, 226–230 (2017).

66. Ashworth, A., Rastan, S., Lovell-Badge, R. & Kay, G. X-chromosome inactivation may
explain the difference in viability of XO humans and mice. *Nature* **351**, 406–408 (1991).

67. Tang, Y. A. *et al.* Efficiency of Xist-mediated silencing on autosomes is linked to
chromosomal domain organisation. *Epigenetics Chromatin* **3**, 10 (2010).

68. Graves, J. A. M. Sex chromosome specialization and degeneration in mammals. *Cell*
124, 901–914 (2006).

69. Charlesworth, B. The evolution of chromosomal sex determination and dosage
compensation. *Curr. Biol.* **6**, 149–162 (1996).

70. Chen, G. *et al.* Single-cell analyses of X Chromosome inactivation dynamics and
pluripotency during differentiation. *Genome Res* **26**, 1342–1354 (2016).

71. Parekh, S., Ziegenhain, C., Vieth, B., Enard, W. & Hellmann, I. zUMIs - A fast and
flexible pipeline to process RNA sequencing data with UMIs. *Gigascience* **7**, (2018).

72. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21
(2013).

73. Hagemann-Jensen, M. *et al.* Single-cell RNA counting at allele and isoform resolution
using Smart-seq3. *Nat. Biotechnol.* **38**, 708–714 (2020).

74. Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573–
3587.e29 (2021).

75. Meldal, B. H. M. *et al.* Complex Portal 2018: extended content and enhanced
visualization tools for macromolecular complexes. *Nucleic Acids Res.* **47**, D550–D558
(2019).

76. Rostam, N. *et al.* CD-CODE: crowdsourcing condensate database and encyclopedia.
Nat. Methods **20**, 673–676 (2023).

77.   Dyer, S. C. *et al.* Ensembl 2025. *Nucleic Acids Res* **53**, D948–D957 (2025).

78.   McKusick, V. A. *Mendelian Inheritance in Man: A Catalog of Human Genes and Genetic*
Disorders. (JHU Press, 1998).

79.   Chen, E. Y. *et al.* Enrichr: interactive and collaborative HTML5 gene list enrichment
analysis tool. *BMC Bioinformatics* **14**, 128 (2013).

80.   Sherry, S. T. *et al.* dbSNP: the NCBI database of genetic variation. *Nucleic Acids Res.*
29, 308–311 (2001).

81.   Krueger, F. & Andrews, S. R. SNPsplit: Allele-specific splitting of alignments between
genomes with known SNP genotypes. *F1000Res.* **5**, 1479 (2016).

82.   Krueger, F., James, F., Ewels, P., Afyounian, E. & Schuster-Boeckler, B.
FelixKrueger/TrimGalore: v0.6.7 - DOI via Zenodo. (Zenodo, 2021).
doi:10.5281/ZENODO.5127899.

83.   Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. *Nat. Biotechnol.* **37**, 907–
915 (2019).

84.   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,
2078–2079 (2009).

85.   Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program
for assigning sequence reads to genomic features. *Bioinformatics* **30**, 923–930 (2014).

86.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

87.   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat.*
Methods **9**, 357–359 (2012).

88.   Ramírez, F., Dündar, F., Diehl, S., Grüning, B. A. & Manke, T. deepTools: a flexible
platform for exploring deep-sequencing data. *Nucleic Acids Res* **42**, W187–91 (2014).

89.   Robinson, J. T. *et al.* Integrative genomics viewer. *Nat Biotechnol* **29**, 24–26 (2011).

90.   ryan-allsop, Pasque lab. pasquelab/Mice-XCU: X-chromosome upregulation
operates on a gene-by-gene basis at RNA and protein levels. *Zenodo*
https://doi.org/10.5281/zenodo.16760254 (2025).

1119

1120

ACKNOWLEDGMENTS
We thank Cheryl Maduro and Büşra Göynük for technical support in generating the cell
lines, the VIB FACS Core for their assistance in sorting cells, the KU Leuven Genomics Core
for assistance with sequencing, as well as the Vlaams Supercomputer Centrum for their
computational resources. We thank Björn Reinius and our colleagues for many helpful
discussions. The V.P. laboratory is supported by the Research Foundation-Flanders (FWO
grants G0C9320N and G0B4420N to V.P.), KU Leuven Research Fund (C1 grant
C14/21/119 to V.P.), and Pandarome project 40007487 (G0I7822N) (funded by the FWO
and F.R.S.-FNRS) under the Excellence of Science (EOS) program. R.N.A is supported by
FWO fellowship (11L0724N). J.B. is supported by the Oncode Institute. S.M. is supported
by ZonMW-Top subsidy (40008129815046).

1132
AUTHOR CONTRIBUTIONS
Conceptualization: V.P., R.N.A., J.G., B.F.T, J.B., H.M.B., C.G.
Data curation: R.N.A., B.F.T., J.B., S.M.
Formal analysis: R.N.A., B.F.T., J.B., S.P., W.F.J.I, J.A.A.D
Funding acquisition: V.P., J.G.
Investigation: V.P., R.N.A., J.G., B.F.T, J.B., H.M.B., C.G.
Methodology: R.N.A., B.F.T., J.B., S.P., W.F.J.I, J.A.A.D
Project administration: V.P., J.G.
Resources: V.P., J.G., J.A.A.D
Software: V.P., J.G., J.A.A.D
Supervision: V.P., J.G., H.M.B., C.G
Validation: R.N.A., B.F.T., J.B.
Visualization: R.N.A., J.B., B.F.T
Writing, reviewing and editing of the manuscript: V.P., R.N.A., J.G., B.F.T, J.B., H.M.B.,
C.G.

1148
1149 **COMPETING INTERESTS**
1150 KU Leuven in Belgium has filed an international patent application (PCT/EP2023/073949)
1151 for methods to create ExM cells from naive human PSCs. V.P. is one of the inventors of this
1152 patent. The remaining authors declare no competing interests.

FIGURE LEGENDS

Figure 1. Transcriptional dosage compensation of X-linked genes in mouse pluripotent stem cell lines with X-chromosome monosomy.

a. Schematic depiction of cells with different sex chromosome complements (XX, XY, and XO) and the corresponding expected upregulation status of the X chromosome (gray: not upregulated, green: upregulated).

b, c. Violin plots of the mean normalized gene expression from the *Mus* (**b**) and *Cast* (**c**) allele of all expressed X-linked genes in $X^{Cast}X^{Mus}$ (green), $X^{Cast}O^{Mus}$ (blue), $X^{Mus}O^{Cast}$ (orange), and $X^{Mus}Y$ (yellow) cells. Dashed line indicates the mean value of XX cells, serving as a reference line for expected expression without upregulation. Each dot represents a cell. P-values were calculated via the Tukey HSD test (two-sided, with Tukey-Kramer correction for multiple comparisons).

d, e. Violin plots of the mean X:A ratio of the *Mus* (**d**) and *Cast* (**e**) allele in $X^{Cast}X^{Mus}$, $X^{Cast}O^{Mus}$, $X^{Mus}O^{Cast}$, and $X^{Mus}Y$ cells. The dashed line indicates the mean value of XX cells, serving as a reference line for expected expression without upregulation. Each dot represents a cell. P-values were calculated using the Tukey HSD test (two-sided, with Tukey-Kramer correction for multiple comparisons).

f. Normalized expression in $X^{Cast}X^{Mus}$ iPSCs of genes annotated as upregulated (in XO and XY cells, n=103, green) or non-upregulated (n=168, gray). P-value was calculated using a two-sided Student's t-test.

g. Schematic overview of the experimental set-up used to generate allele-specific ChIP-seq data[57].

h-i. Line plots showing the promoter enrichment of repressive (**h**) and active (**i**) histone modifications on the active *Cast* allele during XCI inactivation, and thus XCU induction. Promoter enrichment was quantified as the mean counts per 10 million (CP10M) in the 1 kb region upstream of the TSS (as shown for the 24 hour time point in **Supplementary Fig. 3d and 4c**). Mean enrichment is plotted for the promoters of upregulated (green) and non-upregulated (gray) genes separately, with shaded areas indicating the standard error. P-values comparing enrichment at 0 and 24 hours were calculated using two-sided paired t-tests, corrected for multiple testing with the Benjamini-Hochberg method, and shown to the right of each gene set.

Figure 2. Large heterozygous segmental deletions do not result in global XCU *in trans*.
a. Schematic depiction of the XCU center hypothesis. Gray denotes an X chromosome that
is not upregulated, while green indicates an upregulated X chromosome.
b. Schematic overview of the deletion cell lines used in the scRNA-seq experiment and their
respective sex chromosome karyotypes.
c. Heatmap of all scRNA-seq data for X-linked genes, ordered by chromosomal position,
grouped by cell line, and colored by log-transformed normalized expression values for each
allele. The position of genes *Tfe3*, *Zic3*, *Dusp9*, and *Nr0b1* are highlighted, as these genes
denote the start and end positions of each deletion.
d. Schematic depiction of the XCU center hypothesis, such that deletion of a region of one
X chromosome induces transcriptional upregulation for genes located outside of the
deletion region *in trans*.
e-g. Violin plots separated by cell line showing the mean gene expression from the non-
deleted allele for the genes located outside of the regions of Deletion 1 (**e**, *Cast* allele),
Deletion 2 (**f**, *Cast* allele) and Deletion 3 (**g**, *Mus* allele). The dashed line indicates the mean
value of XX cells, serving as a reference line for expected expression without upregulation.
Each dot represents a cell. P-values calculated via the Tukey HSD test (two-sided, with
Tukey-Kramer correction for multiple comparisons, ns - not significant).

Figure 3. The X chromosome does not contain an XCU center.

a. Schematic overview of the deletions generated for the bulk RNA-seq experiment, as well
as deletions 1, 2, and 3. All deletions were introduced in hybrid (*Mus/Cast*) ESC lines with a
heterozygous *Rnf12* deletion on the *Mus* allele. Deletion G was found to be a recombination
event rather than a deletion, and has been omitted from all subsequent analyses.

b. Heatmap of all X-linked genes ordered by chromosomal position, grouped by cell line,
and colored by log-transformed normalized expression values for each allele. The expected
deleted region for each deletion line is highlighted on the right side of the heatmap.

c. Scatter plots separated by deletion showing the mean copy number-corrected
expression levels of genes located outside each deletion for $X^{Cast}X^{Mus}$, $X^{Cast}O^{Mus}$, and cells
with the corresponding deletion. Each dot represents a biological replicate (independent
clones). The dashed line represents the average level from the XX samples, serving as a
reference line for expected expression without upregulation. Note that the gene sets differ
between panels, resulting in varying expression levels in $X^{Cast}X^{Mus}$ and $X^{Cast}O^{Mus}$ across
panels. **d.** Scatter plots separated by deletion showing the $\log_2$ fold change in copy
number-corrected expression levels of replicates of $X^{Cast}X^{Mus}$, $X^{Cast}O^{Mus}$, and cells with the
corresponding deletion, relative to the mean $X^{Cast}X^{Mus}$ levels per gene. Only genes
upregulated in the $X^{Cast}O^{Mus}$ cells and located outside each respective deletion are shown
(n=110, 115, 93, 117, 112, 73, 113, 119, 118, 122, 95, 116, 116). Each dot represents a
gene, and the solid line indicates the median $\log_2$ fold change. P-values comparing mean
expression between genotypes were calculated using two-sided paired t-tests, and
corrected for multiple testing with the Bonferroni method (ns - not significant).

Figure 4. Dosage compensation by transcriptional upregulation occurs *in trans* for
genes within regions of X-chromosome monosomy.
a. Schematic depiction of gene-by-gene regulation of dosage compensation. Gray denotes
an X chromosome that is not upregulated, while green indicates an upregulated X
chromosome.
b-d. Violin plots separated by cell line showing the mean gene expression from the non-
deleted allele for the genes located inside the regions of Deletion 1 (**b**, *Cast* allele), Deletion
2 (**c**, *Cast* allele) and Deletion 3 (**d**, *Mus* allele). The dashed line indicates the mean value of
XX cells, serving as a reference line for expected expression without upregulation. Each dot
represents a cell. P-values were calculated via the Tukey HSD test (two-sided, with Tukey-
Kramer correction for multiple comparisons).
e. Scatter plots separated by deletion showing the mean copy number-corrected
expression levels of genes located within each deletion for $X^{Cast}X^{Mus}$, $X^{Cast}O^{Mus}$, and cells with
the corresponding deletion. Each dot represents a biological replicate (independent clones).
The dashed line indicates the average level from the XX samples, serving as a reference line
for expected expression without upregulation. Deletion I did not contain sufficient
expressed genes and has been omitted.
f. Scatter plots separated by deletion showing the $\log_2$ fold change in copy number-
corrected expression levels of replicates of $X^{Cast}X^{Mus}$, $X^{Cast}O^{Mus}$, and cells with the
corresponding deletion, relative to the mean $X^{Cast}X^{Mus}$ levels per gene. Only genes
upregulated in the $X^{Cast}O^{Mus}$ cells and located within each respective deletion are shown
(n=15, 10, 32, 8, 13, 52, 12, 6, 7, 3, 30, 9, 9). Each dot represents a gene, and the solid line
indicates the median $\log_2$ fold change.
g. Heatmap showing the fold change in expression of upregulated genes identified from the
allelic single-cell analysis between $X^{monosomy}$ and $X^{Cast}X^{Mus}$. Fold change is calculated between
Cast expression from either $X^{Cast}O^{Mus}$, Deletion 1, or Deletion 2 cells and total allelic
expression divided by 2 from XX cells, or *Mus* expression from $X^{Mus}O^{Cast}$, $X^{Mus}Y$, or Deletion
3 cells and total allelic expression divided by 2 from $X^{Cast}X^{Mus}$ cells.
h. Bar plot of the relative RNA expression of the gene *Pdzd4* in XX wild type ($X^{Cast}X^{Mus}$, n=3),
Rnf12$^{+/-}$ parental line (*Rnf12*$^{+/-}$, n=3), XO ($X^{Cast}O^{Mus}$, n=2), and 4 different *Rnf12*$^{+/-}$*Pdzd4*$^{+/-}$
clones for both the *Mus* (blue) and *Cast* (red) allele. Average expression ± standard
deviation, each dot represents a biological replicate (for XO and *Rnf12*$^{+/-}$*Pdzd4*$^{+/-}$ from
independent clones).

Figure 5. Dosage compensation is induced by transcriptional upregulation *in trans*
within regions of monosomic expression on chromosome 3.
a. Schematic depiction of the location of the *Xist* transgene on chromosome 3[61]. After 24
hours of dox treatment, *Xist* coats the *Cast* chromosome 3, reducing gene expression.
b. Overview of the genomic location of genes showing reduced expression on the Cast
allele of chromosome 3 (black lines), as well as the location of the left border (red line) of the
region defined as being inactive (green area) and the site of the *Xist* transgene (blue line).
c-e. Plot of median normalized expression from the *Cast* (**c**) or *Mus* (**d**, **e**) allele of genes
inside (**c**, **e**) or outside (**d**) the region with reduced gene expression for both control (-dox)
and treated (+dox) conditions. Dashed lines indicate the mean levels per condition. P-
values were calculated via the two-sided Student's t-test (ns - not significant). Each dot
represents a biological replicate.

Figure 6. Protein-level dosage compensation is induced within regions of monosomic
expression on the X chromosome.
a. Copy number-corrected protein abundance for all X-linked proteins in $X^{Cast}X^{Mus}$, $X^{Mus}Y$ and
$X^{Cast}O^{Mus}$ ESCs. The mean (of n=127 proteins) per biological replicate from a single clone is
depicted with a dot, with the corresponding 95% confidence intervals shown as error bars.
The dashed line indicates the mean value of $X^{Cast}X^{Mus}$ cells, serving as a reference line for
expected expression without upregulation.
b, c. Volcano plots displaying the differential protein expression analysis of X-linked
proteins between $X^{Mus}Y$ and $X^{Cast}X^{Mus}$ samples (**b**) and between $X^{Cast}O^{Mus}$ and XX samples (**c**)
(n=3). Each dot represents a protein. P-values were calculated from copy number-corrected
protein quantifications using Tukey's HSD test (two-sided, with Tukey-Kramer correction for
multiple comparisons). Significantly up- and downregulated proteins are highlighted in red
and blue, respectively ($\alpha < 0.05$, $\log_2$(fold change) > 0.5 or < -0.5, respectively).
d. Barplot displaying the proportion of upregulated genes per chromosome. The X
chromosome is highlighted in green.
e. Barplot showing the proportion of complex-forming proteins in the upregulated (green)
and non-upregulated (gray) protein lists. P-values were calculated via the Chi-square test.
f. Scatter plot showing the correlation between upregulation at RNA and protein level. Each
dot represents a gene/protein. For both levels, the fold change between $X^{Cast}O^{Mus}$ and half of
the $X^{Cast}X^{Mus}$ level is shown for each gene. Genes are colored according to their upregulation
status at both (orange), either (blue or green) or neither (gray) levels. Shaded area indicates
95% confidence interval.
g, h. Scatter plots separated by region showing the mean copy number-corrected
abundance levels of proteins encoded by genes located outside (**g**) and within (**h**) region E
and K for $X^{Cast}X^{Mus}$, $X^{Mus}Y$, $X^{Cast}O^{Mus}$ ESCs and ESCs with the corresponding deletion. Each
dot represents a biological replicate from a single clone. The dashed line represents the
average abundance levels from the $X^{Cast}X^{Mus}$ samples.
i. Heatmap showing the z-scores of copy-number corrected protein quantifications for X-
linked proteins upregulated in both $X^{Cast}Y$ and $X^{Cast}O^{Mus}$ samples compared to $X^{Cast}X^{Mus}$.
Proteins are ordered by genomic locations, with deletions E and K highlighted in light blue
and pink, respectively. Proteins expected to be upregulated are highlighted with a box.

a

X chromosome deletion

XX → XX ← inside *in trans*?

☐ Not upregulated
☐ Upregulated

b X−linked genes inside deletion 1 *Cast* allele

c X−linked genes inside deletion 2 *Cast* allele

d X−linked genes inside deletion 3 *Mus* allele

e Genes inside deletion region

A B AB C D C-G E F H I-TEL J K

f Upregulated X-linked genes inside deletion region

A B AB C D C-G E
F H I I-TEL J K

g
Deletion 1   Deletion 2   Deletion 3

$\log_2\left(\frac{X^{Monosomy}}{X^{Total}/2}\right)$

h *Pdzd4*
Mus Cast

a

tetO
Xist

+dox

Nesterova *et al.*, 2019

Dosage
compensation
by upregulation?

Xist

Mus *Cast*
chr3

Mus *Cast*
chr3

b

Inactivated genes on chr3 *Cast* allele upon addition of dox

Centromere

tetO *Xist*

Gene
density

80-160Mb
region

0bp   50mbp   80mbp100mbp   150mbp

Genomic position

c   **d**   **e**

Cast allele   *Mus* allele   *Mus* allele
80-160 Mb region   Outside 80-160 Mb region   80-160 Mb region

Mean gene expression (TPM)

1.84e-3   ns   0.0286

-dox   +dox   -dox   +dox   -dox   +dox