Evolution of thyroglobulin: an integrated view of its origin and complexity from a structural perspective.

Mauricio Gomes Pio[1,2], Carina M. Rivolta[1,2] and Héctor M. Targovnik[1*]

[1] **Universidad de Buenos Aires. Facultad de Farmacia y Bioquímica. Departamento de Microbiología, Inmunología, Biotecnología y Genética/Cátedra de Genética. Buenos Aires, Argentina.**

[2] **CONICET-Universidad de Buenos Aires. Instituto de Inmunología, Genética y Metabolismo (INIGEM). Buenos Aires, Argentina.**

Short title: Thyroglobulin evolution.

Key words: Evolution, Thyroglobulin, Nidogen 2, Sea lamprey, Bioinformatic tools

*Address correspondence and requests for reprints to:

Dr. Héctor M. Targovnik

Universidad de Buenos Aires. Facultad de Farmacia y Bioquímica. Departamento de Microbiología, Inmunología, Biotecnología y Genética/Cátedra de Genética, Av. Córdoba 2351, Cuarto Piso, Sala 5, C1120AAR - Buenos Aires, Argentina.

Tel.: 54-11-5950-8805   E-mail:htargovn@ffyb.uba.ar

Abstract

This study provides an extensive bioinformatics evaluation of the origin and structural complexity of thyroglobulin (TG). We investigate the structural and evolutionary conservation of TG in *Petromyzon marinus* (sea lamprey) by reconstructing its complete TG sequence and comparing it to human TG. Using genomic data from NCBI, Uniprot, and Ensembl, we assembled a 2831-amino-acid sequence (TGPM), identifying key TG modules, including 11 TG type 1 modules, three TG type 2 modules, five TG type 3 modules, as well as the linker, hinge, and ChEL domains. Sequence alignment with human TG revealed strong conservation of hormonogenic tyrosines and cysteine residues, underscoring their structural significance in both species. We generated a PDB model for TGPM using the Swiss Model program for homology modeling. The full-length sea lamprey TG was subsequently aligned with full-length human TG through the Matchmaker command in UCSF ChimeraX, revealing a consistent structural overlap across all four regions (I, II, III, IV), highlighting their evolutionary conservation. Additionally, we identified a second sea lamprey TG transcript, TGPM[1746], which begins with the TG type 1-10 module and extends to the carboxyl terminus, exhibiting high homology with TGPM except in region III.

We also present an evolutionary model proposing that nidogen, a protein containing TG type 1 modules, may have been the precursor to TG. Through genetic rearrangements and duplication events, a proto-TG likely emerged, influenced by environmental factors such as ionizing radiation. Structural modifications led to the loss of nidogen's globular regions, resulting in a secretory proto-TG containing three TG type 1 modules. Successive duplications expanded the TG structure, ultimately forming TG 11 type 1 modules, closely resembling the ancestral TG in *Petromyzon marinus*. The final evolutionary phase involved the incorporation of TG type 3 and TG type 2 modules, followed by the fusion of the ChEL domain, which enhanced TG secretion and thyroid hormone biosynthesis. Our findings confirm that the TG complexation process is

fully established in lampreys and has remained virtually unchanged throughout vertebrate evolution.

Introduction

Thyroid hormone (TH) biosynthesis played a crucial role in evolution, modulating gene expression across various tissues and regulating complex biological processes. These include the transition from larva to adult, cell differentiation, energy metabolism, seasonal physiological changes, and the development of the central nervous system. The significant role of TH biosynthesis in vertebrates contributed to their remarkable diversification and increasing complexity over time, enabling them to acquire unique traits. The emergence of an endocrine system capable of concentrating iodide from seawater was essential for TH production, providing metabolic support for the development of a more complex brain, the formation of a skeletal structure, and the regulation of body temperature.

During the evolution of chordates—including the subphyla Cephalochordata, Urochordata (or Tunicata), and Vertebrata (or Craniata)—thyroid hormone formation preceded the morphological differentiation of thyroid cells and their organization into follicles [Suzuki and Kondo, 1971; Suzuki et al., 1975]. The storage of thyroid hormones bound to thyroglobulin (TG) within the follicular lumen is a characteristic feature of vertebrates. Cephalochordates, urochordates, and vertebrates share a common ancestor that likely lived around 550 million years ago [Putnam et al., 2008]. Vertebrates are classified into gnathostomes and cyclostomes (also known as agnathans or jawless fish), which diverged early in vertebrate evolution before the emergence of hinged jaws, a defining trait of gnathostomes [Schimeld et al., 2012].

The only surviving lineages of cyclostomes today fall into two categories: lampreys (also known as Hyperoartia) and hagfishes (also known as Myxini) [Xu et al., 2016]. Cyclostomes are believed to have first appeared during the Ordovician, reached their peak in the Silurian and Devonian, and then gradually declined toward extinction in the late Devonian [Xu et al., 2016]. Recently, using fossil calibrations, Marlétaz et al. [2024] estimated that the divergence between lampreys and hagfishes occurred in the late Ordovician, approximately 449 million years ago, while the split

between cyclostomes and gnathostomes took place near the Cambrian-Ordovician boundary, around 493 million years ago.

The oldest known lamprey fossils date back to the late Devonian period, approximately 360 million years ago, and their morphology has remained virtually unchanged over hundreds of millions of years of evolution [Xu et al., 2016]. The most extensively studied modern lamprey species include *Petromyzon marinus* (sea lamprey), *Lethenteron japonicum*, *Lampetra fluviatilis*, and *Lampetra planeri*. Lampreys produce TG 18-19S before larval metamorphosis, meaning before the thyroid follicles observed in adults are fully organized [Monaco et al., 1978]. In contrast, hagfishes contain iodinated glycoproteins of a non-TG nature, distinguished by differences in molecular size, solubility, amino acid composition, and carbohydrate and iodoamino acid content and structure [Ohmiya et al., 1989]. However, the possibility that the amino acid sequence of the hagfish iodoprotein may correspond to or be related to a TG domain cannot be ruled out.

The monomeric human TG preprotein consists of a polypeptide comprising 2,768 amino acids (NCBI: NP_003226.4; Uniprot: P01266) [- Holzer et al., 2016; Malthiéry and Lissitzky, 1987; van de Graaf et al., 2001]. The classical model of TG's primary structure is organized into four distinct regions (I, II, III, and IV) [Citterio et al., 2021; Holzer et al., 2016; Malthiéry and Lissitzky, 1987; Mercken et al., 1985; Molina et al., 1996; Parma et al., 1987; van de Graaf et al., 2001]. Region I contains 10 of the 11 TG type 1 repeats, along with two modules known as the linker and hinge, plus an amino-terminal T_4-forming site. Region II includes three TG type 2 repeats and the 11th TG type 1 repeat, while Region III encompasses all five TG type 3 repeats. The fourth region consists of a domain homologous to esterases, known as the cholinesterase-like (ChEL) domain [Park and Arvan, 2004; Swillens et al., 1986], along with a carboxi-terminal T_3-forming site.

The TG type 1 domain first appeared at the base of metazoan evolution [Novinec et al., 2006; Takagi et al., 2022; Yurchenco and Wadsworth, 2004]. Several proteins containing TG type 1 domains emerged during the early stages of life development, including nidogen-1 (NID-1), nidogen

(NID-2), SMOC and SMOC-like, testican and testican-like, the invariant chain, IGFBP, and trop. Homology between esterases and TG was first identified by comparing the primary structures of *Torpedo californica* acetylcholinesterase and bovine TG [Schumacher et al., 1986], suggesting that both evolved from a common ancestral gene. Tandem duplication events in the amino-terminal and amino-central regions, followed by fusion with an acetylcholinesterase-like (ChEL) domain, enabled TG to acquire new folding, secretion, and TH biosynthesis capabilities [Holzer et al., 2016].

In this study, we present a comprehensive bioinformatics analysis of the potential origin and structural complexity of TG. The primary objectives were to examine the organization and architecture of full-length TG and its alternative transcripts in a primitive jawless vertebrate fish, sea lamprey, to hypothesize the origin of TG from a protein present at the base of vertebrate evolution that contains TG type 1 modules, such as NID-2, and to develop a hypothetical model outlining the evolutionary events that shaped the different TG domains.

Materials and Methods

Amino acid sequence alignments using Clustal Omega, T-Coffee, EMBOSS Needle and Blat Search Genome.

The TG amino acid sequence homology between multiple sequence were performed using the multiple sequence alignments Clustal Omega with its visualization tool MView (https://www.ebi.ac.uk/Tools/msa/clustalo/) and T-Coffee, (https://tcoffee.crg.eu/). T-Coffee is a versatile multiple sequence alignment tool suitable for aligning various types of biological sequences, including nucleic acids and proteins. More than just an aligner, T-Coffee acts as a framework that integrates different alignment methods with additional structural, evolutionary, or experimental data to enhance accuracy and meaningful alignments. It employs a library generated from a combination of pairwise alignments to construct multiple alignments. The coloured graphical output visually represents the level of consistency between the final alignment and the library used by T-Coffee, where the main score—the overall consistency value—ranges from 0 to 100. A score of 100 indicates complete agreement between the alignment and its reference library. The alignment file uses a color scheme to denote consistency, from blue (low-consistency regions), which are seldom biologically meaningful, to red (high-consistency regions)**,** which reflect strong agreement.

The amino acid sequence homology between sequence pairs was assessed using EMBOSS Needle (https://www.ebi.ac.uk/Tools/psa/emboss_needle/) and Blat Search Genome (https://genome-euro.ucsc.edu/cgi-bin/hgBlat).

SignalP 6.0: Signal Peptide Prediction and Cleavage Site Identification

The SignalP 6.0 server predicts the presence of signal peptides and determines their cleavage sites in proteins (https://services.healthtech.dtu.dk/services/SignalP-6.0/). It can distinguish between five types of signal peptides: (1) Sec/SPI, standard secretory signal peptides transported by the Sec translocon and cleaved by Signal Peptidase I; (2) Sec/SPII, lipoprotein signal peptides

transported by the Sec translocon and cleaved by Signal Peptidase II; (3) Tat/SPI, Tat signal peptides transported by the Tat translocon and cleaved by Signal Peptidase I; (4) Tat/SPII, Tat lipoprotein signal peptides transported by the Tat translocon and cleaved by Signal Peptidase II and (5) Sec/SPIII, Pilin and pilin-like signal peptides transported by the Sec translocon and cleaved by Signal Peptidase III.

Modeling the three-dimensional atomic structure of proteins

The three-dimensional atomic structure of *Homo sapiens* TG has been characterized using a composite cryo-electron microscopy (cryo-EM) density map (PDB: 6SCJ, resolution: 3.60 Å; https://www.rcsb.org/structure/6SCJ; Coscia et al., 2020). Coscia et al., [2020] proposes a new model of the primary structure of the human TG organized in five regions: N-Terminal Domain (NTD), Core, Flap, Arm and C-Terminal Domain (CTD). However, the TG models presented by Adaixo et al. [2022] and Kim et al. [2021] adhere to the general framework of the classical model for the TG monomer. Therefore, we adapted the Coscias et al., [2020] model align with the classical model. Spacers 1, 2, and 3 are located in regions II, III, and IV, respectively. The UCSF Chimera program was used to visualize the 3D model of the *Homo sapiens* TG (Pettersen et al., 2021; 2004; UCSF Resource for Biocomputing, Visualization, and Informatics at the University of California, San Francisco, https://www.cgl.ucsf.edu/chimera/).

The PDB for TG type 1-1 and TG type 1-2 from *Homo sapiens* NID-2, as well as of TGPM[2475] (Uniprot: S4R814_PETMA), TGPM[567] (NCBI: XP_032817835.1), TGPM[1746] (NCBI: XP_032817730.1) and full-lenght assembled TG (TGPM) from *Petromyzon marinus,* were generated using homology modeling with Swiss Model (https://swissmodel.expasy.org/interactive) due to the unresolved crystallographic data.

The three-dimensional structures of NID-2 from *Petromyzon marinus* and *Homo sapiens*, as well as TG type 1-1 and TG type 1-2 modules from NID-1, NID-2, and TG from *Petromyzon marinus*,

Mus musculus, and *Homo sapiens*, were generated using the AlphaFold server (https://alphafoldserver.com/). pLDDT (predicted Local Distance Difference Test) provides a per-atom confidence estimate on a 0-100 scale, where a higher value indicate greater reliability. Regions with low pLDDT scores may correspond to unstructured or highly flexible areas.

Structural matching using the Matchmaker command in UCSF ChimeraX.

Structural matching (structural superposition) of different PDBs was performed using the Matchmaker command in UCSF ChimeraX. This command first generates a pairwise sequence alignment and then refines the alignment by optimizing residue pairs at the atomic level. The RMSD (Root-Mean-Square Deviation) value in Matchmaker quantifies the average distance between atoms—typically backbone atoms—of superimposed molecules, measured in Ångströms (Å). In globular protein conformation studies, three-dimensional structural similarity is commonly assessed by calculating the RMSD of Cα atomic coordinates after optimal rigid-body superimposition. A cutoff value of 2 Å is used, where lower RMSD values indicate higher-quality superimposition, with values approaching 0 Å representing near-perfect alignment.

Results

Assembly of full-length thyroglobulin from sea lamprey and comparative domain analysis.

To obtain the full-length TG from *Petromyzon marinus* and analyze its structural organization, we retrieved three peptides from the NCBI, Uniprot, and Ensembl databases, each containing sequences characteristic of TG architecture. A 2475 amino-acids sequence (Supplementary Figure S1a), referred to here as TGPM^{2475}, is indexed in the Uniprot database (S4R814_PETMA) and extracted from the *Petromyzon marinus* genome assembly available on ENSEMBL (ENSPMAG 00000001187, ENSPMAT00000001350.1), located on scaffold GL476337: 707,134-808,438 forward strand [Smith et al., 2013]. Alignment with human TG (NP_003226.4) using EMBOSS Needle reveals that TGPM^{2475} contains eleven TG type 1 modules, three TG type 2 modules and five TG type 3 modules, linker and hinge domains, and spacers 1, 2 and 3, along with only the first 256 amino acids of the ChEL domain. However, TGPM^{2475} lacks the signal peptide, the amino-terminal hormonogenic acceptor tyrosine required for T$_4$ formation, the remaining amino acids of the ChEL domain, and the carboxy-terminal hormonogenic site necessary for T$_3$ formation. Our upstream analysis to the first amino acid in TGPM^{2475}, corresponding to the first cysteine of the TG type 1-1 module in scaffold GL476337, allowed us to identify the first 11 amino acids of exon 2 (YSDSKTSLASA). Among these, three amino acids correspond to the first three amino acids of the TG type 1-1 module (ASA) and the amino-terminal hormongenic acceptor site (Y) (Supplementary Figure S2a,b). The glutamic acid residue (gag codon) overlaps the splice site between exons 1 and 2. In upstream exon 1, following the 687-nt intron, the first 25 amino-terminal amino acids of sea lamprey TG were identified (Supplementary Figure S2a,b). The SignalP 6.0 program predicts that this sequence contains a signal peptide with a cleavage site between positions 24 and 25 (MRTSPLLPATTTLYLVLWIGTISA) with a probability of 0.981653 (Supplementary Figure S2c).

A second sequence of 1746 amino acids, corresponding to *Petromyzon marinus* TG was identified in the NCBI database and is referred to here as TGPM^{1746} (*Petromyzon marinus* isolate kPetMar1 chromosome 27, kPetMar1.pri, whole genome shotgun sequence, NCBI reference sequence: NC_046095, XP_032817730.1, Supplementary Figure S3a). Pairwise sequence alignment of TGPM^{2475} and TGPM^{1746} using the EMBOSS Needle program revealed that TGPM^{1746} begins at TG type 1-10 and, notably, contains the complete ChEL domain and the carboxy-terminal hormogenic site required for T$_3$ formation—which are partially and entirely absent in TGPM^{2475}, respectively.

The pairwise sequence alignment of TGPM^{1746} and the scaffold GL476337 using the Blat Search Genome, with assembly WUGSC 7.0/petMar2, revealed alignement with the initial 258 amino acids of the ChEL domain (positions 1172-1429 of TGPM^{1746}) and the last 115 carboxy-terminal amino acids (positions 1632-1746 of TGPM^{1746}), including the final 68 amino acids (positions 1632-1699 TGPM^{1746}) of the ChEL domain and the carboxy-terminal hormonogenic site required for T$_3$ formation (tyrosine^{1745}). However, no alignments was observed between residues 1430-1631 of TGPM^{1746} (202 amino acids) with scaffold GL476337, as this region is absent from the genome data (positions 811252-833817). This absence creates a gap in the ChEL domain of the *Petromyzon marinus* TG sequence previously published by Holzer et al. [2017]. A 26-amino-acids was identified immediately preceding the TG type 1-10 of TGPM^{1746}. This sequence begins with methionine and contains 3 alanines at positions 13, 14 and 15, as well as two cysteines at positions 8 and 19. (MPSRVRACVRAPAAATFLCSALPSPG). The SignalP 6.0 program predicted that TGPM^{1746} contains a signal peptide, with a probability of 0.4996 (Supplementary Figure S3b).

To construct the full-length *Petromyzon marinus* TG sequence, we integrated: 36 upstream amino acids of the TGPM^{2475} identified in the scaffold GL476337, 204 downstream amino acids of the TGPM^{2475} identified in NC_046095 (TGPM^{1746}), 202 of which correspond to the missing region in scaffold GL476337 and the 115 carboxy-terminal amino acids identified in the scaffold

GL476337. This assembly results in the full-length sea lamprey TG (TGPM), comprising 2831 amino acids. Figure 1 presents the alignment between TGPM and *Human sapiens* TG (TGHS) using EMBOSS Needle, highlighting the location of the different domains, cysteines and hormononegic tyrosines. Although the primary structures of sea lamprey TG and human TG diverge (identity: 35.1%, similarity: 50.5%), the hormonogenic tyrosines and over 90% of the cysteine residues are highly conserved at identical positions. (Figure 1).

In summary, according to our results the sea lamprey TG gene is a single-copy gene, approximately 121 kb long, divided into 53 exons. Its mRNA encodes 8493 nt, corresponding to coding sequences (Supplementary Figure S4). The sea lamprey monomeric TG preprotein consists of 24-amino-acid leader peptide followed by a 2807-amino-acid polypeptide.

Remarkably, the comparison of TG sequences between *Petromyzon marinus* (TGPM) and two other lamprey species, *Lampetra fluviatilis* (TGLF) and *Lampetra planeri* (TGLP), using Clustal Omega with its visualization tool Mview, reveals a high degree of sequence conservation among the three species. The sequence identity is 86.9% between TGPM and TGLF and 86.7% between TGPM and TGPLP (Supplementary Figure S5). This finding confirms that our strategy for obtaining the full-length TG of *Petromyzon marinus* was accurate. Additionally, it suggests that the divergence between these three lamprey lineages occurred relatively recently. A comparative analysis of TG organization in TGPM, TGLF and TGLP demonstrates that TG in each species contains all regions found in mammalian TG (Supplementary Figure S5). The four TG regions, along with their domains and modules, are similar in size to human TG, with strict conservation of cysteines residues. Furthermore, the amino-terminal and carboxyl-terminal hormonogenic sites are highly conserved across all three species (Figure 1, Supplementary Figure S5).

A comparative sequence analysis between positions 1189 and 2831 of TGPM and the full length of TGPM[1746], conducted using the EMBOSS Needle program, revealed five major differences at the following positions: 1388 (hinge domain), 1673 to 1710 (TG type 3a1), 1831 to 1850 (TG type 3b1),

1881 (TG type 3b1) and 2233 to 2235 (Spacer 3) of the TGPM (see Supplementary Figure S6).

Interestingly, in the latter region, TGPM[1746] exhibits a 39 amino acid sequence containing 8 cysteines, resembling the TG type 3b modules (SLGTSCAARCGLYSPGQPCQCNRECQRFGDCCPDAAVCL).

In addition to TG[1746] a second sequence of 567 amino acids, related to *Petromyzon marinus* TG, was identified in the NCBI database and is referred to here as TGPM[567] (XP_032817835.1, Supplementary Figure S7a). Pairwise sequence alignment of TGPM and TGPM[567] using the EMBOSS Needle program revealed that TGPM[567], in addition to linker domain, contains the TG type 1-4, TG type 1-5, TG type 1-6 modules and the first 42 amino acids of TG type 7. 51 amino acids at the amino-terminal end and 22 amino acids at the carboxy-terminal end of TGPM[567] are not represented in the TGPM (Supplementary Figure S8). The amino-terminal 51-amino-acid sequence begins with methionine and includes three alanines, at positions 13, 14 and 15, as well as one cysteine at position 51 (MELLPLGEKDDDAAALTAPALAPSVQEAPVFRLLRRRFLALRLLLSGSFRC). However, the SignalP 6.0 program does not predict this sequence to contain a signal peptide (Supplementary Figure S7b).

+

To analyze the homology profile among the 11 TG type 1 modules from TGPM and TGHS we performed an alignment using the T-Coffee program. The analysis revealed a high total consistency value of 75 points, indicating strong sequence conservation (Supplementary Figure S9). Among the 11 of TG type 1 modules, the six cysteine residues are highly conserved across both species. Figure 2 presents the classical alignment of the 11 TG type 1 modules, highlighting the six conserved cysteines. Most TG type 1 repeats exhibit similar sizes, except for TG Type 1-3, TG Type 1-7, and TG Type 1-8, which contain long sequence insertions. Remarkably, these insertions are highly conserved in both species. In TGPM, TG Type 1-3 exhibits a 40-amino-acids insertion in the middle

of the module and an additional 46-amino-acids insertion at the end of the module (40 and 45 amino acids, respectively, in human TG). Similarly, TG Type 1-7 contains a 128-amino-acids insertion in the middle of the module (123 in human TG), while TG Type 1-8 features a 100-amino-acids insertion at the beginning of the module, which is identical in human TG (Figure 2). A pairwise comparison of the individual modules between TGHS and TGPM, conducted using the EMBOSS Needle program, revealed an average identity of 37.94% and an average similarity of 51.24% (Table 1, Supplementary Figure S10).

The alignment of the three TG type 2 modules from TGPM and TGHS, performed using the T-Coffee program, yielded a total consistency value of 94, indicating strong sequence conservation (Supplementary Figure 11). This value is higher than that obtained in the the TG type 1 module comparison, suggesting greater homology. Both cysteines residues are fully conserved between the TG type 2 modules of TGPM and TGHS (Figure 3). A pairwise comparison of the individual modules, conducted using the EMBOSS Needle program, revealed an average identity of 45.70% and an average similarity of 55.86% (Table 1, Supplementary Figure S12).

The alignment of the five TG type 3 modules from TGPM and TGHS performed using T-Coffee program yielded a total consistency value of 69, indicating moderate sequence conservation (Supplementary Figure S13). This value is lower than that obtained in the TG type 1 modules and TG type 2 modules comparisons, suggesting lower homology. All six cysteine residues are fully conserved in the TG type 3a modules (Figure 4), as are all eight cysteines in the TG type 3b modules of TGPM and TGHS, except for the eighth cysteine in TG type 3b1 and TG type 3b2 which the T-Coffee program failed to align (Supplementary Figure S13). A pairwise comparison of the individual modules, conducted using the EMBOSS Needle program, revealed an average identity of 30.26% and an average similarity of 48.58% (Table 1, Supplementary Figure S14). In all cases, the six cysteines of the TG type 3a modules and the eight cysteines of the TG type 3b modules are conserved, except for the eighth cysteine in the TG type 3b1 module, which EMBOSS Needle also

failed to align, consistent with the T-Coffee results. Additionally, an extra cysteine was identified at position 1682 in the TG type 3a1 module of TGPM (Supplementary Figure S14).

Finally, the comparative analysis of the ChEL domain between TGPM and TGHS conducted using the EMBOSS Needle program revealed an average identity of 41.70% and an average similarity of 57.40% between the two species (Table 1). Six of the seven cysteines located in the ChEL domain of TGHS are conserved in TGPM (Figure 1).

Comparative 3D structural analysis of sea lamprey thyroglobulins and human thyroglobulin using the Matchmaker tool.

To compare the 3D structure of each of sea lamprey TG isoforms with human TG, as well as among the sea lamprey TG isoforms between themselves, we generated PDB for TGPM[567], TGPM[1746] and TGPM[2475], since PDBs were not available. We used the Swiss Model program to perform homology modeling for the sequence of these three proteins (TGPM[567]: NCBI, XP_032817835.1, Supplementary Figure S7c; NCBI, TG[1746]: XP_032817730.1, Supplementary Figure S3c; TG[2475]: Uniprot, S4R814_PETMA, Supplementary Figure S1b), using on PDB 7N4Y (resolution: 2.61Å) of *Bos Taurus* TG as a reference [https://www.rcsb.org/structure/7N4Y; Kim et al., 2021].

The structural comparison between sea lamprey TGPM[576] or TGPM[1746] or TGPM[2475] monomer (chain B) and human TG monomer (PDB: 6scj, chain A) was conducted using the Matchmaker command in the UCSF ChimeraX program. The sea lamprey TGPM[567] was aligned with TGHS with an RMSD of 0.200 Å across 396 overlapping atom pairs, with a sequence alignment score: 1067 (Figure 5a1, a2). TGPM[567] corresponds to region I of TGHS, overlapping amino-acids 266-771. The assembly of sea lamprey TGPM[1746] and TGHS resulted in an RMSD of 1.076 Å, based on 806 overlapping atom pairs, with a sequence alignment score of 2879.4 (Figura 5b1, b2). Similarly, the assembly of TGPM[2475] and TGHS yielded an RMSD of 1.219 Å, across 1271 overlapping atom

pairs, with a sequence alignment score of 4688.8 (Figure 5c1, c2). A consistent overlap was observed across the four regions (I, II, III, IV) of sea lamprey TGPM[1746] and TGPM[2475], aligning with the four regions of TGHS. This confirms the homology between the proteins, extending from the TG type 1-10 module to the carboxy-terminal end in TGPM[1746] and from the TG type 1-1 module to the first 256 amino acids of the ChEL domain in TGPM[2475].

Next, we compared the three-dimensional structure of TGPM[567] (chain A) and TGPM[1746](chain A), TGPM[567] (chain A) and TGPM[2475] (chain B), and TGPM[1746] (chain A) and TGPM[2475] (chain A). The Matchmaker tool confirmed the homogeneous overlap of the complexes, with RMSD values of 0.868, 1.176 and 0.295 Å for TGPM[567] and TGPM[1746] (42 overlapping atom pairs, sequence alignment score: 163.6, Figure 6a1, a2), for TGPM[567] and TGPM[2475] 351 overlapping atom pairs, sequence alignment score: 2407.7, Figure 6b1, b2) and for TGPM[1746] and TGPM[2475] (1246 overlapping atom pairs, sequence alignment score: 6249, Figure 6c1, c2) parings, respectively.

Finally, we generated a PDB for TGPM using the Swiss Model program for homology modeling (Figure 7, Supplementary Video 1), using on PDB 7N4Y of *Bos Taurus* TG as a reference. The full-length sea lamprey TG (TGPM, chain B) was then aligned with TGHS (chain A), yielding an RMSD of 1.191 Å across 1449 overlapping atom pairs, with a sequence alignment score: 5212.9) (Figure 7, Supplementary Video 2, Supplementary Video 3). As observed previously with TGPM[1746] and TGPM[2475], the alignment between TGPM and TGHS exhibits a homogeneous overlap across the four regions (I, II, III, IV) of sea lamprey TG, corresponding to the four analogous regions in human TG.

Comparative analysis of TG type 1 and TG type 2 modules between nidogen-2 and thyroglobulin.

As observed earlier, many proteins incorporate the TG type 1 module in their structure. Among them, nidogen stands out, as it is present at the base of vertebrates. Given that our extensive analysis demonstrates that cyclostomes, such as the sea lamprey, already exhibit a modular architecture

highly similar to human TG, we propose that nidogen may serve as the precursor of TG in its complexation process.

NID-1 and NID-2 are homologous sulfated monomeric glycoproteins found in the basal lamina, already observed in cyclostomes and amphioxus, two extensively studied protovertebrates [Holland and Short, 2010; Novinec et al., 2006; Takagi et al., 2022; Yurchenco and Wadsworth, 2004]. In mammals, both proteins exhibit a modular structure consisting of three globular regions (G1, G2 and G3), separated by a a short flexible linker region between G1 and G2 and a rod-like domain between G2 and G3 [Bechtel et al 2012, Töpfer and Holz, 2024] (Figure 8a). The G1 globular region contains the NIDO domain (nidogen-like domain), while the G2 globular region consists of the nidogen G2 beta-barrel domain and one EGF-like domain (epidermal growth factor-like domain). The rod-like domain is composed of four EGF-like domains and one TG type 1 domain in NID-1 (Uniprot: P14543, human) or two TG type 1 domain in NID-2 (Uniprot: Q14112, human) of mammals (Figure 8a). However, sea lamprey NID-2, which is related to a primitive nidogen, contains only a single TG type 1 module (Uniprot: A0AAJ7X4A4). Notably, invertebrates such as Drosophila and C. elegans do not possess TG type 1 domains. In NID-1, the G3 globular region consists of four LDL-receptor class B repeats and an additional EGF-like domain, whereas in NID-2, it contains five LY modules [Bechtel et al 2012, Töpfer and Holz, 2024]. Figure 8a presents the three-dimensional atomic structure of sea lamprey NID-2 (which contains only one TG type 1 module) and human NID-2 (which possesses two TG type 1 modules). These structures were generated using with AlphaFold server, as their crystallography remains unresolved.

To analyze the homology profile among the of TG type 1 modules of *Petromyzon marinus*, *Mus musculus* and *Homo sapiens*, including NID-1, NID-2, TGPM and TGHS, we performed an alignment using the T-Coffee program and 3D structure modeling using AlphaFold server. The T-Coffee analysis yielded a high total consistency value of 89 points, indicating strong sequence conservation (Supplementary Figure S15). The six cysteine residues were well conserved across all

type 1 TG repeats analyzed. Additionally, AlphaFold modeling of TG type 1-1 and TG type 1-2 modules from nidogens and thyroglobulins in sea lamprey, mouse, and human revealed similar secondary structures, characterized by one α-helix and two β-sheets, with a very high plDDT score (Figure 8b). Figure 9 illustrates the alignment of all TG type 1 modules around the six conserved cysteines. Notably, the structural conservation of these modules is high in both nidogens and TGs. The primary structure alignment, performed using EMBOSS Needle, reveals a sequence divergence between sea lamprey nidogen and human nidogen, with an identity of 37.0% and a similarity of 48.1%—values comparable to those observed in the TG comparison between both species (see above). The SignalP 6.0 program predicts that NID-2 sequence in both sea lamprey and human contains a signal peptide, with a cleavage site between positions 36 and 37 (MGMSRAVARARSPAGYAALPAALLCLAAALLGCARG) and a probability of 0.951473. In human NID-2, the cleavage site is predicted between positions 30 and 31 (MEGDRVAGRPVLSSLPVLLLLPLLMLRAAA) with a probability of 0.943032 (data not shown).

Structural pairing Analysis of TG type 1-1 and TG type 1-2 from human NID-2 and human TG using the Matchmaker tool.

To support the hypothesis that nidogen may have been the ancestral protein leading to the first proto-TG, we compared the 3D structures of TG type 1-1 and TG type 1-2 from human NID-2 and human TG. Since PDB for TG type 1-1 and TG type 1-2 from human NID-2 were not available, we generated them using the Swiss Model program for the homology modeling, based on the human sequence indexada en Uniprot (Q14112) and the 1NPE *mus musculus* PDB model, which is the only available PDB for nidogen.

The structural comparison between TG type 1-1 and TG type 1-2 from human NID-2 monomer (chain A) and human TG monomer (PDB: 6scj, chain A) was conducted using the Matchmaker

command of the UCSF ChimeraX program. The TG type 1 modules from human NID-2 were aligned with human TG, resulting in a sequence alignment score of 267.2 (Figure 10). Interestingly, the TG type 1-1 module of NID-2 precisely matches the TG type 1-1 module of human TG (Figure 10). In contrast, the TG type 1-2 module of NID-2 does not align with human TG; likely due to the greater complexity of its alpha helix, which may interfere with structural alignment. A similar result was observed when TG type 1-1 and TG type 1-2 from human NID-2 (chain A)—obtained through homology modeling—were alignd with full human NID-2 (1.cif chain A), generated using AlphaFold server. The alignement resulted in a sequence alignment score of 774.5 (data not shown).

Evolutionary model of the thyroglobulin complexation process.

In our evolutionary model, we propose that a protein with TG type 1 modules, such as nidogen, may have gave rise to TG. Nidogen could have undergone duplication events and genetic material loss, leading to the emergence of a proto-TG, a process facilitated by a favorable genomic and genetic rearrangement environment. Among the most significant catalysts for this process was ionizing radiation, particularly ultraviolet (UV) radiation, as early Cambrian atmospheric oxygen levels were estimated to be only about 10% of present levels [Sperling et al., 2015], and the ozone layer was not yet fully formed. Within nidogen, the TG type 1 module may have duplicated, resulting in one of its isoforms acquiring two additional domains, ultimately forming a structure composed of three TG type 1 modules (Figure 11a). Unlike nidogen, TG lacks the three globular regions (G1, G2 and G3) and none of the associated modules (NIDO, nidogen G2 beta-barrel, EGF-like, LDL-receptor class B). The elimination of the domains in G1 and G2, as well as EGF-like modules, may have resulted from recombination events (Figure 11b). The loss of the G1 domain could have been partial, preserving the signal peptide. Meanwhile, the loss of LDL-receptor class B repeats may be explained by molecular mechanisms leading to a premature termination codon before the G3 globular domain (Figure 11c). As a result, the first secretory proto-TG likely emerged,

containing three TG type 1 modules (Figure 11d), resembling the p.Arg296Ter mutant associated with HC. In addition to the three existing motifs, the protein may have undergone several variations, leading to the emergence of specific tyrosine residues. These residues facilitated the organification of iodine, enabling the synthesize T_4 through the interation of an acceptor tyrosine and a donor tyrosine (Figure 11e). It is plausible that a short amino-terminal portion of TG, containing a single hormonal site and exhibiting a secretory profile, was sufficient for the survival of primitive vertebrates on the earth's surface, exerting physiological effects of HT. This hypothesis is reinforced by extensive documentation of mild hypothyroidism phenotype in patients homozygous for the p.Arg296Ter variant [Caputo et al., 2007; Citterio et al., 2013; Pardo et al., 2009; Rivolta et al., 2005; Siffo et al., 2018; van de Graaf et al., 1999; Zou et al., 2018].

Following the formation of this first TG prototype, a new duplication event likely led to the emergence of a proto-TG containing four TG type 1 modules (Figure 11f), resembling the p.A340Ter mutant, which exibits an active secretory profile, as described by Lee et al., 2011. Subsequent tandem duplication events of TG type 1 modules continued to shape the TG structure, ultimately resulting in a protein with 11 repeats (Figure 11g). This marks another evolutionary stage in TG's development, bringing it closer to its definitive structure, as observed in the sea lamprey, positioned at the base of vertebrate evolution. Insertion of genetic material after modules 4 and 10 would lead to the formation of the non-repetitive linker and hinge domains, respectively (Figure 11h).

Our structural analysis suggests a relationship between the TG type 1 and TG type 3-a modules, as both contain of six cysteines. This finding supports the hypothesis that TG type 3 modules may have originated from the duplication of the TG type 1-11 module (Figure 12i). Through tandem duplications, the TG type 3-a1 module gave rise to the TG type 3-b1, TG type 3-a2 and TG type 3-a3 domains (Figure 12j). Further gene modulations involving partial deletions led to the differentiation of TG type 3-a2 and TG type 3-a3 from TG type 3-a1, while TG type 3-b1 underwent

an insertion event, and the addition of two extra cysteines due to the duplication of cysteine 3 and the presence an additional cysteine in the 3-b1 module (Figure 12k). The structural similarity between the TG type 3-b1 and TG type 3-b2 modules is due to the conservation of eight cysteines in both. This suggests that the duplication of the TG type 3-b1 module with its insertion between TG type a-2 and TG type a-3, led to the emergence of the TG type 3-b2 module (Figure 12l). TG type 3-b2 undergoes a deletion, reducing its length to 140 amino acids in sea lamprey TG (Figure 4), compared 172 amino acids in TG type 3-b1 and 134 out of 169 amino acids in human TG (Figure 4). Thus, TG incorporates the five modules that constitute its region III (Figure 12ll). These processes considerably reduced the sequence conservation patterns in region III due to its moderate evolutionary pressure, in contrast to TG type 1 repetitive modules, whose antiproteolytic activity is crucial for protein preservation.

The TG Type 2 modules likely originated from the insertion of a transposable DNA fragment between the hinge and TG Type 1-11 domains (Figure 12m), followed by two duplications events (Figure 12n). To investigate the origins of this module, the TG type 2-1, TG type 2-2 and TG type 2-3 modules were individually compared against the human genome ad other organisms using the BLAST tool (NCBI). The analysis revealed no homology with any genomic elements apart from the TG Type 2 domain itself. However, Lee et al. [2011] classified it as a distant member of the GCC2/GCC3 domain superfamily, despide the lack of significant homology. The higher conservation of TG Type 2 modules compared to TG type 1 and TG type 3 modules, as indicated by the sea lamprey-human alignment (Figure 3, Table 1), supports to the hypothesis that the TG Type 2 region emerged after the formation of the TG type 3 modules.

The final—and perhaps most significant—structural modulation event in TG was the incorporation of the ChEL domain and its fusion with regions I, II, and III (Figure 12o and p). The ChEL domain functions as an intramolecular chaperone, enhancing TG's secretory potential [Lee et al., 2008]. The higher conservation of the ChEL domain compared to TG Type 1 and TG Type 3

modules, as indicated by the sea lamprey-human alignment (Table 1), supports the hypothesis that the ChEL domain emerged after the formation of TG Type 1 and TG Type 3 modules. With the incorporation of the ChEL domain, TG is now secreted more efficiently and has an enhanced ability to increase T_4 concentration in the blood. Additionally, TG has acquired the carboxy-terminal hormonogenic site necessary for T_3 formation. The secretion of TH in its active form, T_3, along with the increased levels of T_4, may have enhanced all functions associated with TH. As previously noted, in the TG of the *Petromyzon marinus*, the complexation process is already fully established.

Discussion

Protein diversity arises through a series of processes, including mutation, fixation, and selection, which can be neutral, deleterious, or advantageous. Domain duplication, gene fusion, and recombination are fundamental evolutionary mechanisms that shape proteins and contribute to proteome complexity [Vogel et al., 2005]. Protein evolution is influenced not only by selection for structure and function but also by the genomic location of coding genes, their expression patterns, their role within biological networks, and their resistance to translation errors [Pál et al., 2006]. Many proteins consist of combinations of multiple domains from diverse origins, enabling the modification of pre-existing proteins or the development of new architectures. Over time, these processes have led to the emergence of novel biological functions, increasing the complexity of multicellular organisms [Patthy, 2003; Vogel et al., 2004].

Interestingly, domain combinations within a protein can be represented as a network, where nodes correspond to functional domains from different superfamilies. The structural organization of TG exemplifies genetic evolution through intragenic duplication events involving TG type 1, TG type 2, and TG type 3 modules, as well as gene fusion, as observed with the ChEL domain (Figures 11 and 12). While the number of possible domain combinations is virtually unlimited, only a select few persist due to strong evolutionary selection. However, the reduction in protein diversity appears to be linked more to the rate of protein production than to overall protein abundance [Pál et al., 2006]. Additionally, radiation—particularly UV radiation—can act as a non-biological catalyst, driving successive cycles of DNA damage and subsequent repair through homologous recombination. This process likely accelerated the rapid emergence and increasing complexity of genomes, leading to the formation of intricate proteins such as TG within a relatively short evolutionary timeframe. All the modulating events proposed in our model occurred randomly, within a brief lapse of time, governed by the fundamental forces of natural selection and genetic

drift, with radiation as persistent physical stressor to DNA that possibly served as a catalyst, accelerating these evolutionary processes.

Using accumulated genomic data, we obtained the full-length cDNA of sea lamprey TG, TGPM (Figure 1), and compared its regions and domains with human TG. Notably, a second lamprey TG transcript, TGPM[1746] (Supplementary Figure S6), was identified in the NCBI database. This transcript begins with the TG type 1-10 module and extends to the carboxy-terminal end, exhibiting high homology with TGPM, except in region III, where the TG type 3 repeats show significant variability, and in spacer 3, they resemble the TG type 3b modules. Additionally, this transcript features an amino-terminal sequence compatible with a signal peptide and was likely generated through alternative splicing from a novel promoter.

Although the primary structures of sea lamprey and human TG diverge, with 35.1% identity and 50.5% similarity, all domains—both repetitive (TG type 1, TG type 2, and TG type 3) and non-repetitive (linker, hinge, and ChEL)—as well as the amino- and carboxy-terminal hormonogenic sites, remain strictly conserved in sea lamprey (Figure 1). Given the lower identity and similarity of TG type 1 modules compared to the ChEL domain, we propose that these two domains emerged at different stages of TG evolution, under different environmental conditions. TG type 1 modules likely arose earlier, in more primitive periods under stressful environmental conditions, which drove more modulated events leading to sequence variability. In contrast, the more conserved amino acid profile of the ChEL domain suggests that its fusion with the first proto-TG occurred during a period of reduced genomic rearrangement and lower environmental stress. The structural complexity of genetic traits enabled vertebrates to colonize terrestrial environments and become the dominant species on earth. The fusion of TG with the ChEL domain may have marked a pivotal turning point in vertebrate evolution.

The TG type 1 module predates the emergence of TG in the earliest vertebrates and is present in several extracellular proteins, suggesting its likely involvement in the evolution of multicellular

organisms by facilitating cell-cell and cell-environment interactions [Novinec et al., 2006]. A total of 170 protein sequences have been identified containing 333 TG type 1 modules [Novinec et al., 2006]. Several proteins with TG type 1 domains, such as nidogen—proposed by us as a precursor to TG type 1 modules—play crucial physiological roles, including partial or complete inhibition of protein degradation [Fowlkes et al., 1997]. The TG type 1 module can also regulate specific degradation events in proteins that incorporate this module into their structure [Fineschi et al., 1995]. The evolutionary success of the TG type 1 module is likely attributed to its ability to reversibly bind proteases, offering protection to proteins with more resilient TG type 1 modules, enabling them to endure and thrive in proteolytic environments during the evolution and establishment of multicellular life [Molina, 1996; Novinec et al., 2006]. In a multicellular environment, close cellular interactions create a microenvironment more exposed to proteolytic enzymes released by neighboring cells during natural metabolic processes. The process of TH release itself involves the transport of TG to lysosomes [Ericson, 1981; Rousset and Mornex, 1991]. TG degradation is a slow process, requiring several hours [Brix and Herzog, 1994], and involves proteolytic cleavage by thyroid proteases [Dunn, 1991]. The eleven TG type 1 repeats may regulate this process in thyrocytes through selective and reversible inhibition of endosomal and/or lysosomal proteases, playing a vital role in the physiology of TH release [Molina, 1996].

In the TG type 1-3, TG type 1-7, and TG type 1-8 modules, insertions in human TG are highly conserved in sea lamprey TG (Figure 2), likely originating from an intron exonization event, as suggested by Parma et al. [1987] for the central insertion of TG type 1-7, or from random DNA insertion mechanisms. While both possibilities must be considered, it is also plausible that gene rearrangements and recombination-repair processes played a key role in promoting this insertion pattern, alongside tandem duplication and gene modulation events that shaped these modules throughout their evolutionary trajectory. These findings lead us to hypothesize that these TG type 1 repeats share a common ancestral domain that underwent extensive intragenic rearrangements,

preserving only structural elements essential for their physiological function, such as the six cysteines required for disulfide bond formation. This strongly suggests that these insertions emerged during or prior to lamprey evolution and have remained unchanged in vertebrates.

Proteins require a stable three-dimensional structure to function properly. This stability ensures that a sufficient number of active proteins are available in the cell [Pál et al., 2006]. Each TG monomer contains 122 cysteine residues, accounting for 4.44% of the total amino acids [Citterio et al., 2021; Holzer et al., 2016; Malthiery and Lissitzky, 1987; Mercken et al., 1985; van de Graaf et al., 2001]. Newly synthesized TG folds through the formation of up to 60 intrachain disulfide bonds. As noted above, TG type 1, TG type 2, and TG type 3 domains are cysteine-rich repeat modules that are covalently linked by intrachain disulfide bonds, playing a crucial role in shaping the tertiary structure of TG. The TG type 1 domains from human and bovine TG are further categorized into two subgroups: type 1A and type 1B. Type 1A repeats (TG type 1–1 to TG type 1–8 and TG type 1–10) contain six cysteines residues, whereas type 1B repeats (TG type 1–9 and TG type 1–11) include only four cysteines residues [Molina et al., 1996]. Conserved intradomain disulfide bonding patterns—Cys1–Cys2, Cys3–Cys4, and Cys5–Cys6—are consistently observed in all type 1A repeats [Molina et al., 1996]. Additionally, TG type 1B, TG type 2, TG type 3 repeats, and the six Cys residues of the ChEL domain also undergo intrachain disulfide bonding. Remarkably, over 90% of the cysteine residues are conserved between sea lamprey and human TG, underscoring their critical role in the structural and functional integrity of TG.

In conclusion, we have identified and characterized the full-length cDNA of sea lamprey TG, analyzing its structural conservation throughout evolution. Additionally, we detected a second TG transcript in sea lamprey, likely originating from an alternative splicing process. Based on our findings, we propose a model for the TG complexation process, beginning with the emergence of TG type 1 modules from nidogen, followed by the incorporation of TG type 3 and subsequently TG type 2 modules, leading to fusion with the ChEL domain as the final event in this process. Finally,

our data confirm that the TG complexation process is fully established in the lamprey and has remained unchanged throughout vertebrate evolution.

Acknowledgements

Conflict of Interest Statement

The authors have no conflicts of interest to declare.

Funding Sources

This study was funded by grants from the Fondo para la Investigación Científica y Tecnológica (FONCyT-ANPCyT-MINCyT, PICT-2018-02146 to H.M.T.), CONICET (PIP 2021-11220200102976CO to C.M.R.) and Universidad de Buenos Aires (UBACyT 2020-20020190100050BA to C.M.R.).

Credit Author Statement

M.G.P. contributed to bioinformatic analysis and performing structural modeling analysis. C.M.R. contributed to the acquisition of funds and the design of the study. HMT contributed to bioinformatics alignement analysis, funding acquisition, study conception and design, and writing of the article. All authors critically reviewed and participated in manuscript revision and approved the final draft.

Data Availability Statement

Data and material are available from the authors upon request.

References

Adaixo, R., Steiner, E.M., Righetto, R.D., Schmidt, A., Stahlberg, H., Taylor, N.M.I., 2022. Cryo-EM structure of native human thyroglobulin. Nat. Commun. 13, 61. doi:10.1038/s41467-021-27693-8.

Bechtel, M., Keller, M.V., Bloch, W., Sasaki, T., Boukamp, P., Zaucke, F., Paulsson, M., Nischt, R., 2012. Different domains in nidogen-1 and nidogen-2 drive basement membrane formation in skin organotypic cocultures. FASEB J. 26, 3637-3648. doi:10.1096/fj.11-194597

Brix, K., Herzog, V., 1994. Extrathyroidal release of thyroid hormones from thyroglobulin by J774 mouse macrophages. J. Clin. Invest. 93, 1388-1396. doi: 10.1172/JCI117115

Caputo, M., Rivolta, C.M., Esperante, S.A., Gruñeiro-Papendieck, L., Chiesa, A., Pellizas, C.G., González-Sarmiento, R., Targovnik, H.M., 2007. Congenital hypothyroidism with goitre caused by new mutations in the thyroglobulin gene. Clin. Endocrinol. 67, 351–357. doi: 10.1111/j.1365-2265.2007.02889.x

Citterio, C.E., Machiavelli, G.A., Miras, M.B., Gruñeiro-Papendieck, L., Lachlan, K., Sobrero, G., Chiesa, A., Walker, J., Muñoz, L., Testa, G., Belforte, F.S., Gonzalez- Sarmiento, R., Rivolta, C.M., Targovnik, H.M., 2013. New insights into thyroglobulin gene: molecular analysis of seven novel mutations associated with goiter and hypothyroidism. Mol. Cell. Endocrinol. 365, 277–291. doi: 10.1016/j.mce.2012.11.002

Citterio, C.E., Rivolta, C.M., Targovnik, H.M., 2021. Structure and genetic variants of thyroglobulin: pathophysiological implications. Mol. Cell. Endocrinol. 508, 111227. https://doi.org/10.1016/j.mce.2021.111227

Coscia, F., Taler-Verčič, A., Chang, V.T., Sinn, L., O'Reilly, F.J., Izoré, T., Renko, M., Berger, I., Rappsilber, J., Turk, D., Löwe, J., 2020. The structure of human thyroglobulin. Nature 578, 627-630. doi:10.1038/s41586-020-1995-4.

Dunn, A.D., Cruntchfield, H.E., Dunn, J.T., 1991. Proteolytic processing of thyroglobulin by extracts of thyroid lysosomes. Endocrinology 128: 3073-3080. doi: 10.1210/endo-128-6-3073

Ericson, L., 1981. Exocytosis and endocytosis in the thyroid follicle cell. Mol. Cell. Endocrinol. 22, 1-24. doi: 10.1016/0303-7207(81)90098-8

Fineschi, B., Arneson, L.S., Naujokas, M.F., Miller, J., 1995. Proteolysis of major histocompatibility complex class II-associated invariant chain is regulated by the alternatively spliced gene product, p41. Proc. Natl. Acad. Sci. USA 92, 10257-10261. doi: 10.1073/pnas.92.22.10257

Fowlkes, J.L., Thrailkill, K.M., George-Nascimento, C., Rosenberg, C.K., Serra, D.M., 1997. Heparin-binding, highly basic regions within the thyroglobulin type-1 repeat of insulin-like growth factor (IGF)-binding proteins (IGFBPs) -3, -5, and -6 inhibit IGFBP-4 degradation. Endocrinology 138, 2280-2285. doi: 10.1210/endo.138.6.5182

Holzer, G., Morishita, Y., Fini, J.-B., Lorin, T., Gillet, B., Hughes, S., Tohmé, M., Deléage, G., Demeneix, B., Arvan, P., Laudet, V., 2016. Thyroglobulin represents a novel molecular architecture of vertebrates. J. Biol. Chem. 291, 16553-16566. doi: 10.1074/jbc.M116.719047

Holland, L.Z., Short, S., 2010. Alternative splicing in development and function of chordate endocrine systems: a focus on Pax genes. Integr. Comp. Biol. 50, 22-34. doi: 10.1093/icb/icq048

Kim, K., Kopylov, M., Bobe, D., Kelley, K., Eng, E.T., Arvan, P., Clarke, O.B., 2021. The structure of natively iodinated bovine thyroglobulin. Acta Crystallogr. D Struct. Biol. 77, 1451-1459. doi:10.1107/S2059798321010056.

Lee, J., Di Jeso, B., Arvan, P., 2008. The cholinesterase-like domain of thyroglobulin functions as an intramolecular chaperone. J. Clin. Invest. 118, 2950-2958. doi: 10.1172/JCI35164

Lee, J., Di Jeso, B., Arvan, P., 2011. Maturation of thyroglobulin protein región I. J. Biol. Chem. 286, 33045-33052. doi: 10.1074/jbc.M111.281337

Malthiéry, Y., Lissitzky, S., 1987. Primary structure of human thyroglobulin deduced from the sequence of its 8448-base complementary DNA. Eur. J. Biochem. 165, 491-498. doi:10.1111/j.1432-1033.1987.tb11466.x

Marlétaz, F., Timoshevskaya, N., Timoshevskiy, V.A., Parey, E., Simakov, O., Gavriouchkina, 0D., Suzuki, M., Kubokawa, K., Brenner, S., Smith, J.J., Rokhsar, D.S., 2024. The hagfish genome and the evolution of vertebrates. Nature 627, 811-820. doi: 10.1038/s41586-024-07070-3

Mercken, L., Simons, M.-J., Swillens, S., Massaer, M., Vassart, G., 1985. Primary structure of bovine thyroglobulin deduced from the sequence of its 8,431-base complementary DNA. Nature 316, 647-651. doi:10.1038/316647a0

Molina, F., Bouanani, M., Pau, B., Granier, C., 1996. Characterization of the type-1 repeat from thyroglobulin, a cysteine-rich module found in proteins from different families. Eur. J. Biochem. 240, 125-133. doi:10.1111/j.1432-1033.1996.0125h.x

Monaco, F., Andreoli, M., La Posta, A., Roche, J., 1978. Thyroglobulin biosynthesis in a larval (ammocoete) and adult freshwater lamprey (Lampetra planeri Bl.). Comp. Biochem. Physiol. B 60, 87-91. doi: 10.1016/0305-0491(78)90033-0

Novinec, M., Kordiš, D., Turk, V., Lenarčič, B. 2006. Diversity and evolution of the thyroglobulin type-1 domain superfamily. Mol. Biol. Evol. 23, 744-755. doi: 10.1093/molbev/msj082

Ohmiya, Y., Suzuki, S., Kondo, Y., 1989. Isolation and characterization of hagfish thyroid iodoprotein by its non-thyroglobulin nature, very high iodine and carbohydrate contents and low hormone/iodine ratio. Eur J Biochem 182, 11-8. doi: 10.1111/j.1432-1033.1989.tb14794.x.

Pál C, Papp, B., Lercher, M.J., 2006. An integrated view of protein evolution. Nat Rev Genet. 7, 337-348. doi: 10.1038/nrg1838

Pardo, V., Vono-Toniolo, J., Rubio, I.G.S., Knobel, M., Possato, R.F., Targovnik, H.M., Kopp, P., Medeiros-Neto, G., 2009. The p.A2215D thyroglobulin gene mutation leads to deficient synthesis and secretion of the mutated protein and congenital hypothyroidism with wide phenotype variation. J. Clin. Endocrinol. Metab. 94, 2938–2944. doi: 10.1210/jc.2009-0150

Park, Y.-n., Arvan, P., 2004.The acetylcholinesterase homology region is essential for normal conformational maturation and secretion of thyroglobulin. J. Biol. Chem. 279, 17085-17089. doi:10.1074/jbc.M314042200.

Parma, J., Christophe, D., Pohl, V., Vassart, G., 1987. Structural organization of the 5' region of the thyroglobulin gene. Evidence for intron loss and "exonization" during evolution. J. Mol. Biol. 196, 769-779. doi:10.1016/0022-2836(87)90403-7

Patthy, L, 2003. Modular assembly of genes and the evolution of new functions. Genetica 118, 217–231.

Pettersen, E.F., Goddard, T.D., Huang, C.C., Couch, G.S., Greenblatt, D.M., Meng, E.C., Ferrin, T.E., 2004. UCSF Chimera-a visualization system for exploratory research and analysis. J. Comput. Chem. 25, 1605-1612. doi: 10.1002/jcc.20084

Pettersen, E.F., Goddard, T.D., Huang, C.C., Meng, E.C., Couch, G.S., Croll, T.I., Morris, J.H., Ferrin, T.E., 2021. UCSF ChimeraX: Structure visualization for researchers, educators, and developers. Protein Sci. 30, 70-82. doi: 10.1002/pro.3943

Putnam, N.H., Butts, T., Ferrier, D.E., Furlong, R.F., Hellsten, U., Kawashima, T., Robinson-Rechavi, M., Shoguchi, E., Terry, A., Yu, J.K., Benito-Gutiérrez, E.L., Dubchak, I., Garcia-Fernàndez, J., Gibson-Brown, J.J., Grigoriev, I.V., Horton, A.C., de Jong, P.J., Jurka, J., Kapitonov, V.V., Kohara, Y., Kuroki, Y., Lindquist, E., Lucas, S., Osoegawa, K., Pennacchio, L.A., Salamov, A.A., Satou, Y., Sauka-Spengler, T., Schmutz, J., Shin-I, T., Toyoda, A., Bronner-Fraser, M., Fujiyama, A., Holland, L.Z., Holland, P.W., Satoh, N., Rokhsar, D.S., 2008. The amphioxus genome and the evolution of the chordate karyotype. Nature 453, 1064-1072. doi: 10.1038/nature06967

Rivolta, C.M., Moya, C.M., Gutnisky, V.J., Varela, V., Miralles-García, J.M., González- Sarmiento, R., Targovnik, H.M., 2005. A new case of congenital goiter with hypothyroidism due to a homozygous p.R277X mutation in the exon 7 of the thyroglobulin gene: a mutational hot spot

could explain the recurrence of this mutation. J. Clin. Endocrinol. Metab. 90, 3766–3770. doi: 10.1210/jc.2005-0278

Rousset, B., Mornex, R., 1991. The thyroid hormone secretory pathway--current dogmas and alternative hypotheses. Mol. Cell. Endocrinol. 78, C89-C93. doi: 10.1016/0303-7207(91)90176-s

Schumacher, M., Camp, S., Maulet, Y., Newton M., MacPhee-Quigley K., Taylor, S. S, Friedmann, T., Taylor, P., 1986. Primary structure of *Torpedo californica* acetylcholinesterase deduced from its cDNA sequence. Nature 319, 407-409. doi: 10.1038/319407a0

Shimeld, S.M., Donoghue, P.C., 2012. Evolutionary crossroads in developmental biology: cyclostomes (lamprey and hagfish). Development 139, 2091-2099. doi: 10.1242/dev.074716

Siffo, S., Adrover, E., Citterio, C.E., Miras, M.B., Balbi, V.A., Chiesa, A., Weill, J., Sobrero, G., González, V.G., Papendieck, P., Martinez, E.B., González-Sarmiento, R., Rivolta, C.M., Targovnik, H.M., 2018. Molecular analysis of thyroglobulin mutations found in patients with goiter and hypothyroidism. Mol. Cell. Endocrinol. 473, 1–16. doi: 10.1016/j.mce.2017.12.009

Smith, J.J., Kuraku, S., Holt, C., Sauka-Spengler, T., Jiang, N., Campbell, M.S., Yandell, M.D., Manousaki, T., Meyer, A., Bloom, O.E., Morgan, J.R., Buxbaum, J.D., Sachidanandam, R., Sims, C., Garruss, A.S., Cook, M., Krumlauf, R., Wiedemann, L.M., Sower, S.A., Decatur, W.A., Hall, J.A., Amemiya, C.T., Saha, N.R., Buckley, K.M., Rast, J.P., Das, S., Hirano, M., McCurley, N., Guo, P., Rohner, N., Tabin, C.J., Piccinelli, P., Elgar, G., Ruffier, M., Aken, B.L., Searle, S.M., Muffato, M., Pignatelli, M., Herrero, J., Jones, M., Brown, C.T., Chung-Davidson, Y.W., Nanlohy, K.G., Libants, S.V., Yeh, C.Y., McCauley, D.W., Langeland, J.A., Pancer, Z., Fritzsch, B., de Jong, P.J., Zhu, B., Fulton, L.L., Theising, B., Flicek P, Bronner, M.E., Warren, W.C., Clifton, S.W., Wilson, R.K., Li, W., 2013. Sequencing of the sea lamprey (Petromyzon marinus) genome provides insights into vertebrate evolution. Nat. Genet. 45, 415-421, 421e1-2. doi: 10.1038/ng.2568

Sperling, E.A., Knoll, A.H., Girguis, P.R., 2015. The ecological physiology of earth's second oxygen revolution. Annu. Rev. Ecol. Evol. Syst. 46, 215-235. doi: 10.1146/annurev-ecolsys-110512-135808

Suzuki, S., Kondo, Y., 1971. Demonstration of thyroglobulin-like iodinated proteins in the branchial sac of tunicates. Gen. Comp. Endocrinol. 17, 402-406. doi: 10.1016/0016-6480(71)90149-3

Suzuki, S., Gorbman, A., Rolland, M., Montfort, M.F., Lissitzky, S., 1975. Thyroglobulins of cyclostomes and an elasmobranch. Gen. Comp. Endocrinol. 26, 56-69. doi: 10.1016/0016-6480(75)90215-4

Swillens, S., Ludgate, M., Mercken, L., Dumont, J.E., Vassart, G., 1986. Analysis of sequence and structure homologies between thyroglobulin and acetylcholinesterase: possible functional and clinical significance. Biochem. Biophys. Res. Commun. 137, 142-148. doi:10.1016/0006-291x(86)91187-3.

Takagi, W., Sugahara, F., Higuchi, S., Kusakabe, R., Pascual-Anaya, J., Sato, I., Oisi, Y., Ogawa, N., Miyanishi, H., Adachi, N., Hyodo, S., Kuratani, S., 2022. Thyroid and endostyle development in cyclostomes provides new insights into the evolutionary history of vertebrates. BMC Biol. 20, 76. doi: 10.1186/s12915-022-01282-7

Töpfer, U., Holz, A., 2024. Nidogen in development and disease. Front. Cell. Dev. Biol. 12:1380542. doi: 10.3389/fcell.2024.1380542

van de Graaf, S.A.R., Ris-Stalpers, C., Veenboer, G.J.M., Cammenga, M., Santos, C., Targovnik, H.M., de Vijlder, J.J.M., Medeiros-Neto, G., 1999. A premature stopcodon in thyroglobulin mRNA results in familial goiter and moderate hypothyroidism. J. Clin. Endocrinol. Metab. 84, 2537–2542. doi: 10.1210/jcem.84.7.5862

van de Graaf, S.A.R., Ris-Stalpers, C., Pauws, E., Mendive, F.M., Targovnik, H.M., de Vijlder, J.J.M., 2001. Up to date with human thyroglobulin. J. Endocrinol. 170, 307-321. doi: 10.1677/joe.0.1700307

Vogel, C., M. Bashton, N. D. Kerrison, C. Chothia, and S. A. Teichmann, 2004. Structure, function and evolution of multidomain proteins. Curr. Opin. Struct. Biol. 14,208–216. doi: 10.1016/j.sbi.2004.03.011

Vogel. C., Teichmann, S.A., Pereira-Leal, J., 2005. The Relationship Between Domain Duplication and Recombination. J. Mol. Biol. 346, 355-365. doi: 10.1016/j.jmb.2004.11.050

Xu, Y., Zhu, S.-W., Li, Q.-W., 2016. Lamprey: a model for vertebrate evolutionary Research. Zool Res 37, 263-269. doi: 10.13918/j.issn.2095-8137.2016.5.263

Yurchenco, P.D., Wadsworth, W.G., 2004. Assembly and tissue functions of early embryonic laminins and netrins. Curr. Opin. Cell. Biol. 16, 572-579. doi: 10.1016/j.ceb.2004.07.013

Zou, M., Alzahrani, A.S., Al-Odaib, A., Alqahtani, M.A., Babiker, O., Al-Rijjal, R.A., BinEssa, H.A., Kattan, W.E., Al-Enezi, A.F., Al Qarni, A., Al-Faham, M.S.A., Baitei, E. Y., Alsagheir, A., Meyer, B.F., Shi, Y., 2018. Molecular analysis of congenital hypothyroidism in Saudi Arabia: SLC26A7 mutation is a novel defect in thyroid dyshormonogenesis. J. Clin. Endocrinol. Metab. 103, 1889–1898. doi: 10.1210/jc.2017-02202

Table 1: Comparative analysis of the TG type 1, TG type 2, and TG type 3 modules, along with the ChEL domain, of full-length thyroglobulin from *Petromyzon marinus* (TGPM) and *Homo sapiens* (TGHS), performed using the EMBOSS Needle program.

	Identity %	Similarity %	Identity average %	Similarity average %
TG type 1-1 **TGPM** /TG type 1-1 **TGHS**	41.3	57.1	**37.94**	**51.24**
TG type 1-2 **TGPM** /TG type 1-2 **TGHS**	61.8	70.6		
TG type 1-3 **TGPM** /TG type 1-3 **TGHS**	31.4	47.9		
TG type 1-4 **TGPM** /TG type 1-4 **TGHS**	44.4	57.1		
TG type 1-5 **TGPM** /TG type 1-5 **TGHS**	42.1	57.9		
TG type 1-6 **TGPM** /TG type 1-6 **TGHS**	42.6	54.4		
TG type 1-7 **TGPM** /TG type 1-7 **TGHS**	34.7	48.8		
TG type 1-8 **TGPM** /TG type 1-8 **TGHS**	37.2	53.2		
TG type 1-9 **TGPM** /TG type 1-9 **TGHS**	26.3	44.7		
TG type 1-10 **TGPM** /TG type 1-10 **TGHS**	38.8	50.7		
TG type 1-11 **TGPM** /TG type 1-11 **TGHS**	16.7	21.2		
TG type 2-1 **TGPM** /TG type 2-1 **TGHS**	42.9	50.0	**45.70**	**55.86**
TG type 2-2 **TGPM** /TG type 2-2 **TGHS**	47.1	58.8		
TG type 2-3 **TGPM** /TG type 2-3 **TGHS**	47.1	58.8		
TG type 3-a1 **TGPM** /TG type 3-a1 **TGHS**	32.5	48.0	**30.26**	**48.58**
TG type 3-b1 **TGPM** /TG type 3-b1 **TGHS**	31.2	45.7		
TG type 3-a2 **TGPM** /TG type 3-a2 **TGHS**	27.8	53.7		
TG type 3-b2 **TGPM** /TG type 3-b2 **TGHS**	31.0	41.3		
TG type 3-a3 **TGPM** /TG type 3-a3 **TGHS**	28.8	54.2		
ChEL **TGPM** /ChEL **TGHS**	41.7	57.4	**41.7**	**57.4**

```
                        signal peptide                              TG Type 1-1
                   ┌──────────────────────────┐                  ┌──────────────
TGPM    1  MRTSPLLPATTTLYLVLWIGTISALEYSDSKTSLASACELLRQQAMEDGR  50
                .:.:|...|.|..:|: :.:..||......| .||.|:.|.....
TGHS    1  --MALVLEIFTLLASICWV-SANIFEYQVDAQPL-RPCELQRETAFLKQA  46
                                                              TG Type 1-2
TGPM   51  LHVPQCSPHGAFRPLQCDAAGEPCWCVDAAGEELPGTRRAEGPPPSCLSF 100
            .:||||:..|:|::.||...|.||.|.|.:|:.   .|.:.|||
TGHS   47  DYVPQCAEDGSFQTVQCQNDGRSCWCVGANGSEVLGSRQ-PGRPVACLSF  95

TGPM  101  CQLHRQRVLLSGYLNGTRALHLPQCTQDGRYRPAQVD-SSGQGWCVDADG 149
           |||.:|::|||||.|....:|||...|.|.|    ...|.|||||:|
TGHS   96  CQLQKQQILLSGYINSTDTSYLPQCQDSGDYAPVQCDVQQVQCWCVDAEG 145
                                  TG Type 1-3
TGPM  150  MEVYGTRQLGAPLSCPSPCQVSVRRAVRASTPGSPPPQCDRDEQRVLATQ 199
           ||||||||.|.||..|.|::..||.:....|  |||..||.  |...:.|
TGHS  146  MEVYGTRQLGRPKRCPRSCEIRNRRLLHGVGDKS-PPQCSA-EGEFMPVQ 193

TGPM  200  CQLIGTAGRSTLSLLDAFSMHPEAFRSLAEFRARLPGVHAYCYCADPQGR 249
           |:.:.|......|:.:.:.|:||.:.:|:.|.|.|..|:|||.|||
TGHS  194  CKFVNTTDMMIFDLVHSYNRFPDAFVTFSSFQRRFPEVSGYCHCADSQGR 243

TGPM  250  ELPETGDRGVRSFLFVAYFSGHGLNHEVVGSVLGKTRHEKFLKVQDRLPG 299
           ||.|||...:...:...|:|..|......|.:...::||.||..:.|
TGHS  244  ELAETGLELLLDEIYDTIFAGLDLPSTFTETTLYRILQRRFLAVQSVISG 293
                      TG Type 1-4
TGPM  300  HIVRPTVCEVERSLAERLGGRGFAPVCSQDGSYIPTQC--HGPTCWCVDT 347
           ....||.|||||||..|...  |..:.|.|.:|.|...||  .||.|||||.
TGHS  294  RFRCPTKCEVERFTATSF-GHPYVPSCRRNGDYQAVQCQTEGP-CWCVDA 341
                                   Linker
TGPM  348  RGDEVFGSRVRDSVPNCGSSSSPCQLERQLALSHLFLGPSAPFID-DLIS 396
           :|.|:.|:|.:....|:|....|  |...|||.|||.|:.|.|..|..  ||.|
TGHS  342  QGKEMHGTRQQGEPPSCAEGQS-CASERQQALSRLYFGTSGYFSQHDLFS 390

TGPM  397  ------ALPRDGSVSFSESPSQEMQHELWLHWRALREQLLDGG-LGEILS 439
           |.||....:..|:               ::|...|.|   .::..
TGHS  391  SPEKRWASPRVARFATSCPPT-------------IKELFVDSGLLRPMVE 427

TGPM  440  QASRLFPASGQASRLLGDLVGGIFHTQELALTAAGANVQPGRIAEMLFGD 489
           ..|:.|..|       ..||.:.:..||.::.||...|.....|.|:.:.|||.
TGHS  428  GQSQQFSVS---ENLLKEAIRAIFPSRGLARLALQFTTNPKRLQQNLFGG 474

TGPM  490  RFLKNLKFFNFTGAVGGRGTFNFSKVFGQVGLTGMYDGANFEELARLFAP 539
           :||.|:..||..:||:|.|||||||.|.|.:||....:.|.|:||.:..:.
TGHS  475  KFLVNVGQFNLSGALGTRGTFNFSQFFQQLGLASFLNGGRQEDLAKPLSV 524

TGPM  540  GEDSYLTRGSSNFSRESFNFDQSIDDPFARATNLERNRNVLGFVVTLLED 589
           |.||..:|.....::.....:.:.....|...||:.|:|.|.|.|:.:|||.
TGHS  525  GLDSNSSTGTPEAAKKDGTMNKPTVGSFGFEINLQENQNALKFLASLLEL 574

TGPM  590  ARFY----------GLVRDITPLIGLSGGDVGLNALTFLRMAEEDRPTQ 628
           ..|.           .:.||:       |||           ..|.
TGHS  575  PEFLLFLQHAISVPEDVARDL--------GDV-------------METV 602
                      TG Type 1-5
TGPM  629  NAADAADGPPASRPFVARCGEDGGAYEPVQCHGAACWCVDAHGREIAGTR 678
           .:.:..:..|  .|.||.||.:  |:||.|||....||||:.:.|:|:.|:|
TGHS  603  LSSQTCEQTP-ERLFVPSCTTE-GSYEDVQCFSGECWCVNSWGKELPGSR 650
                          TG Type 1-6
TGPM  679  AVGGAPPRCPSACEGERARALLARRGRPVGTPLFVPECDGAGDYRPVQCS 728
           ..||  .|||||:.||.:|||.......:|.:.||||.|....|.:.||||
TGHS  651  VRGG-QPRCPTDCEKQRARMQSLMGSQPAGSTLFVPACTSEGHFLPVQCF 699
                                   TG Type 1-7
TGPM  729  GDRCFCVGAGGAEVPGTGRPLGDPVTCPTPCQAAAASELLQQLRHLGPVL 778
           ...|:||.|.|..:|||....:|.|.|..||||||.|.:.:..|:..:.|
TGHS  700  NSECYCVDAEGQAIPGTRSAIGKPKKCPTPCQLQSEQAFLRTVQAL---L 746

TGPM  779  RGDAPSSLPPL---YVPRCDARGGWRPVQCDGAGQQIREFVDAWTKDEGN 825
           ...  |.|||   |:|:|:.||.|.:|:|:.:.|.:.
TGHS  747  SNS--SMLPTLSDTYIPQCSTDGQWRQVQCNGPPEQVFELYQRW-EAQNK 793
```

```
                            ___________________________________________________
TGPM      826  ASKATLSDLRELLARVRGAGGGGGGGGGKGLRSFLSFLYDSGRQDMFPELS   875
               ....|.:.|...:...|.|..|          ....|:..||::|:||:||.||
TGHS      794  GQDLTPAKLLVKIMSYREAASG------NFSLFIQSLYEAGQQDVFPVLS       837

TGPM      876  LYPSLRGPPEVLNFSGSSGR-------FLRNPEVVWKILTENASFVYTGD       918
               .||||:..|      .:...|:        .|..|.:.|:||....|.|.|.
TGHS      838  QYPSLQDVP----LAALEGKRPQPRENILLEPYLFWQILNGQLS-QYPGS       882
                                                          TG Type 1-8
TGPM      919  YAEFSGRYADFESRLCWCVDAAGHEIDGTRTVSPGQLPKCPGACHLAAAD       968
               |::||...|.|:.||||||.||.|::|.|: .|.:||.|||.|.|...
TGHS      883  YSDFSTPLAHFDLRNCWCVDEAGQELEGMRS-EPSKLPTCPGSCEEAKLR       931

TGPM      969  VNRYLQQADLLIGSAGAS--SAGEGVAFGRGLAFTEDELLGSPLGLGDVK      1016
               |.:.::::.:.::.::.:|  ..||...::|:....:: |||.| .|...:
TGHS      932  VLQFIRETEEIVSASNSSRFPLGESFLVAKGIRLRNED-LGLP-PLFPPR       979

TGPM     1017  GEVAAVLAGGTGYAVRLAAQAAMHFHWRRRFFGPGTIGEGVFNGFDPYMP      1066
               ...|.....|:.||:|||||:..|:.||||....:.|........||||
TGHS      980  EAFAEQFLRGSDYAIRLAAQSTLSFYQRRRFSPDDSAGASALLRSGPYMP      1029
                                                           TG Type 1-9
TGPM     1067  QCTEPGGWEPAQCVVRAGAGFCWCVDAAGEFVAGSLVARPRRRPQCATPC      1116
               ||...|.|||.||  .||.|.|||||..|.|.|:.|||.||..:.|||.|.|
TGHS     1030  QCDAFGSWEPVQC--HAGTGHCWCVDEKGGFIPGSLTARSLQIPQCPTTC      1077

TGPM     1117  QRARAEALLTGWKSLGSVENGSSIEVLTKHTPACTPSRKYKKCHYLLRPN      1166
               :::|...||:.||...|.||.|.:.    .|||..:.:|.:........
TGHS     1078  EKSRTSGLLSSWKQARSQENPSPKDLF---VPACLETGEYARLQASGAGT      1124
                                               TG Type 1-10
TGPM     1167  WWPSPSYSYTPTHPSPTTNAESPSLCELRRRQALLRNAGLGSVPECDSR-      1215
               |...|: |.....|...:.::|.|||.|.||||.|....|...|
TGHS     1125  WCVDPA-SGEELRPGSSSSAQCPSLCNVLKSGVLSRRVSPGYVPACRAED      1173
                                                         HINGE
TGPM     1216  GDYAARQC--ADGTCWCAGAGGEEIPDTRRHAGEPGVVCHEPRCELPQDE      1263
               |.:..|| |.|:||||..||.....|||.:|.| .|..|:||.||..:.
TGHS     1174  GGFSPVQCDQAQGSCWCVMDSGEEVPGTRVTGGQP--ACESPRCPLPFNA      1221

TGPM     1264  RRAGRWALVCEDGAAATAAAGLQRCLLACLRGFAHTLDGAGHDGGAPREF      1313
               .......::||..:.|..|.:| :|:|.|.|.:.         .|...
TGHS     1222  SEVVGGTILCETISGPTGSA-MQQCQLLCRQGSWSVF--------PPGPL      1262

TGPM     1314  QCNATSGEWIGSSPQPDACQEIHPWQAVQLSISFTLRFPEEKMCNPEYEG      1363
               .|:..||....|||.||.||....||:.|...|.|:..|.:||...:|..
TGHS     1263  ICSLESGRWESQLPQPRACQRPQLWQTIQTQGHFQLQLPPGKMCSADYAD      1312

TGPM     1364  LLESFNSFLLADLTARGFCHLSNGVGSDAVA--VCDAASVRVRCQDALRL      1411
               ||::|..|:|.|:|||||||||||.:.......|: ||:.:||:|.|....||
TGHS     1313  LLQTFQVFILDELTARGFCQIQVKTFGTLVSIPVCNNSSVQVGCLTRERL      1362

TGPM     1412  LVNVTWRSSLSRLPVQSYPDLHNIERAFVAH--AQRFAQLLDDRGYRLAL      1459
               .|||||:|.|.|..:||.|.||||:|||.|.  ..||..|:....:|.|
TGHS     1363  GVNVTWKSRLEDIPVASLPDLHDIERALVGKDLLGRFTDLIQSGSFQLHL      1412
                                                        TG Type 2-1
TGPM     1460  NGRDYSSEGAARFRKDELYDTSPAFRVACEPGYVRI-------AQGCAA      1501
               :.:.:.:| ..||.:.:.:.|||....|..|.|:.::      ..||.
TGHS     1413  DSKTFPAE-TIRFLQGDHFGTSPRTWFGCSEGFYQVLTSEASQDGLGCVK      1461
                         TG Type 2-2           TG Type 2-3       Spacer 1 TG Type 1-11
TGPM     1502  CPRGSLWRDGGCALCPADHYQDEAGQTACTPCPADTGTLTAGAAAAFQCR      1551
               ||.||..:|..|.|.||..||::||.||.||||.||.|||....|::|||.:....|.
TGHS     1462  CPEGSYSQDEECIPCPVGFYQEQAGSLACVPCPVGRTTISAGAFSQTHCV      1511

TGPM     1552  TPCER-----EALRRGQLGAGER----LELYC-DPEG----LYEGEKPAK      1587
               |.|:| :.:.:.:...|...   .:.:.|.  :|.|.|:
TGHS     1512  TDCQRNEAGLQCDQNGQYRASQKDRGSGKAFCVDGEGRRLPWWETEAPLE      1561
                       Spacer 2                              TG Type 3-a1
TGPM     1588  GGQDEVRRMFEEVPSEDLVLGAVDAVALRIRTETG--PVQEVEPRCLQEC      1635
               ..|..:.::||:||...:.|.:.|.|||:||..|. ||.: ||.:
TGHS     1562  DSQCLMMQKFEKVPESKVIFDANAPVAVRSKVPDSEFPVMQ----CLTDC      1607
```

```
                                ----------------------------------------------------
TGPM    1636 LKEERCDFFVFSTQGDVSSCHFYNGSRESYQCEKAPEEKSALGISLCVVF 1685
             .::|.|.||..||.....||.||..:..:.:::.||||.|..|
TGHS    1608 TEDEACSFFTVSTTEPEISCDFYAWTSDNVACMTSDQKRDALGNSKATSF 1657

TGPM    1686 IAIKCFVRGVGFHVSSASGRFYFHAGHELPSWDGQLYRPTGFGNAASGAY 1735
             .:::|.|: |..| ...|....|.....:....||||..|||..||.|
TGHS    1658 GSLRCQVK-VRSH-GQDSPAVYLKKGQGSTTTLQKRFEPTGFQNMLSGLY 1705
                                                   TG Type 3-b1
                                              ----------------------------------------
TGPM    1736 RTLALPAGTTTLPDAHLYCRSACSAQACCDGFLLKELPPGVLVCALLSGP 1785
             .:...|....|.||||.||..||...|||||.|.::.|..::|.|||.|
TGHS    1706 NPIVFSASGANLTDAHLFCLLACDRDLCCDGFVLTQVQGGAIICGLLSSP 1755

TGPM    1786 TVLTCRAPWWPRTADEYGDGECSGLRVHKPSRTFGFSLGGVHYNASNPPS 1835
             :||.|....|....:.:.:.:.|.|:..:.|......||..:..|.:|
TGHS    1756 SVLLCNVKDWMDPSEAWANATCPGVTYDQESHQVILRLGDQEFIKSLTP- 1804

TGPM    1836 AVASTAAAAAAASQVIYLWAGSEAGSRDVCLHTKAYT-PQPGEPAGAADV 1884
             :..|.........|| |||..|:.|||..:..:..| |:|..|..|...
TGHS    1805 -LEGTQDTFTNFQQV-YLWKDSDMGSRPESMGCRKDTVPRPASPTEAGLT 1852
                                                  TG Type 3-a2
                                         --------------------------------------
TGPM    1885 SGRFSTLRPEEVTLQPEMRVAQQNFWLFRRAFSPPQQATSWCLRRC-RRD 1933
             :..||.:...:|.:......::.|..|||:..|| .|||..|||.|| :.
TGHS    1853 TELFSPVDLNQVIVNGNQSLSSQKHWLFKHLFS-AQQANLWCLSRCVQEH 1901

TGPM    1934 ALCKAAAVGDGPAGWLECLLYPDTQSCGTSQELLERPARAPSCASLLPRI 1983
             :.|:.|.:.:.:....|.||||:.|.| .:::| :.|..|..:||::
TGHS    1902 SFCQLAEITESASLYFTCTLYPEAQVC---DDIME--SNAQGCRLILPQM 1946
                                                         TG Type 3-b2
                                              -------------------------
TGPM    1984 DGTLYRKRDGAEGPVRRLYRRTAFRSEIGVLLRSVANLTGMALSDGFSRC 2033
             ...|:||:...::....:.|..:.:...:...::.::.:|:.::.|||.|
TGHS    1947 PKALFRKKVILEDKVKNFYTRLPFQKLMGISIRNKVPMSEKSISNGFFEC 1996

TGPM    2034 ERTCDEDPCCEGFAFLRRALSSAGEALLQCATHTGPGVQLCGRESPMGA- 2082
             ||.||.||.||.||.||..:...|| :.|:|:.|.|:|:|.|:..|:. ||
TGHS    1997 ERRCDADPCCTGFGFLNVSQLKGGE--VTCLTLNSLGIQMCSEENG-GAW 2043

TGPM    2083 -TFGCPPPGVDTASSLFGWYRAQARSSPQTAQLCPAVTLPPAPTKGAADG 2131
             ...|..|.::..:.:....||||.:....:.:|:...|:..:.:.
TGHS    2044 RILDCGSPDIEVHTYPFGWYQKPIAQN-NAPSFCPLVVLPSLTEKVSLDS 2092
                                                          TG Type 3-a3
                                              -----------------------------------
TGPM    2132 FQSLDVSTAAVDPTISAFDVVVLLGPGAGGSPRAAPPSD-----EWCLAV 2176
             :|||.:|:..|||:|.|||.:.||.:     ..|..|: :.||:.
TGHS    2093 WQSLALSSVVVDPSIRHFDVAHV---------STAATSNFSAVRDLCLSE 2133
                                              --------------------------------
TGPM    2177 CQRSPWCASVGVRHLGDSTRCVLHAETLACGYGARGGRHCRLGVLEPPAR 2226
             |.:...|....:...:.:|.::||.:.:|:.|.|.:.| |:::||.:.|....
TGHS    2134 CSQHEACLITTLQTQPGAVRCMFYADTQSCTHSLQ-GQNCRLLLREEATH 2182
              -- -  Spacer 3                                ChEL
                                                       ==================================
TGPM    2227 LYRKKAPSPGGGQGPVRPVVPLGS-GQLQGASRTVEVDSQVKVIDVFLGV 2275
             :||||...|....:.| |.||:: |:|.|.|.:.::|.:.:.|:|.||||
TGHS    2183 IYRKPGISLLSYEASV-PSVPISTHGRLLGRSQAIQVGTSWKQVDQFLGV 2231
             ==================================================
TGPM    2276 PYAAPPLDANRFSGPQPPRPLAVNETWDADRYKTRPDCLQPDGQRGTS-- 2323
             ||||||...|| |.|.||....:|||. |.|..|.||..:...||
TGHS    2232 PYAAPPLAERRF---QAPEPLNWTGSWDAS--KPRASCWQPGTRTSTSPG 2276
             ==================================================
TGPM    2324 LSEDCLYLNIFVPKVKPHNASVLVFFHG---GDNSFGGSAQGPLDPSYLA 2370
             :||||||||||:|:|:....|||||||||.  .:|.|.|..| :|.|:||
TGHS    2277 VSEDCLYLNVFIPQNVAPNASVLVFFHNTMDREESEGWPA---IDGSFLA 2323
             ==================================================
TGPM    2371 ALGDIIVVTANYRLGLFGFLSTGDEAAAGNWGLLDQQAALRWVRDHAALF 2420
             |:|::|||||.||.|.|||||.:|..|.|||||||||.|||.||..:|..|
TGHS    2324 AVGNLIVVTASYRVGVFGFLSSGSGEVSGNWGLLDQVAALTWVQTHIRGF 2373
             ==================================================
TGPM    2421 GGSAGAVTVAGDRSSADNVGLHLV---APGSRGLFQRAILMGGSVLSPSA 2467
             ||....|::|.||..||..||..|:   |..|: ||:||.||||:|||.|
TGHS    2374 GGDPRRVSLAADRGGADVASIHLLTARATNSQ-LFRRAVLMGGSALSPAA 2422
```

Figure 1. Comparative analysis of full-length thyroglobulins. A comparative analysis of the full-length thyroglobulin from *Petromyzon marinus* (TGPM, 2831 amino acids) and *Homo sapiens* (TGHS, 2768 amino acids) was conducted using the EMBOSS Needle program. Amino acids are represented using single-letter codes, with signal peptides enclosed in boxes. The linker, hinge and ChEL domain, along with eleven TG type 1 modules, three TG type 2 modules, five TG type 3 modules, and spacers 1, 2 and 3 are displayed. The cysteines are highlighted in yellow, while the amino-terminal hormonogenic sites (TGPM: Y^{27} and Y^{153}, TGHS: Y^{24} and Y^{149}) and carboxyl-terminal hormonogenic sites (TGPM: Y^{2830}, TGHS: Y^{2766}) are marked in red. Additionally, the 202 amino acids missing in the GL476337 scaffold are shaded gray.

Figure 2. Alignment of eleven TG type 1 modules. The alignment of the eleven TG type 1 modules of full-length thyroglobulin from *Petromyzon marinus* (TGPM) and *Homo sapiens* (TGHS) was conducted according to Parma y col. [1987], Malthíery y col. [1987], Molina y col. [1996], van Graaf y col., [2001]. The six cysteines are conserved across all modules, except in TG type 1-2, TG type 1-3, TG type 1-5, TG type 1-9 and TG type 1-11 in TGPM, and in TG type 1-9 and TG type 1-11 in TGHS. Additionally, TG type 1-3, TG type 1-7 and TG type 1-8 contain long inserts, which are underlined in the alignment.

TG11PM: TG type 1-1 TGPM, TG12PM: TG type 1-2 TGPM, TG13PM: TG type 1-3 TGPM, TG14PM: TG type 1-4 TGPM, TG15PM: TG type 1-5 TGPM, TG16PM: TG type 1-6 TGPM, TG17PM: TG type 1-7 TGPM, TG18PM: TG type 1-8 TGPM, TG19PM: TG type 1-9 TGPM, TG110PM: TG type 1-10 TGPM, TG111PM: TG type 1-11 TGPM.

TG11HS: TG type 1-1 TGHS, TG12HS: TG type 1-2 TGHS, TG13HS: TG type 1-3 TGHS, TG14HS: TG type 1-4 TGHS, TG15HS: TG type 1-5 TGHS, TG16HS: TG type 1-6 TGHS, TG17HS: TG type 1-7 TGHS, TG18HS: TG type 1-8 TGHS, TG19HS: TG type 1-9 TGHS, TG110HS: TG type 1-10 TGHS, TG111HS: TG type 1-11 TGHS.

Figure 3. Alignment of three TG type 2 modules. The alignment of the three TG type 2 modules of full-length thyroglobulin from *Petromyzon marinus* (TGPM) and *Homo sapiens* (TGHS) was conducted according to Parma et al. (1987), Malthiéry et al. (1987), Molina et al. (1996), and van Graaf et al. (2001). The two cysteines are conserved across all modules.

TG21PM: TG type 2-1 TGPM, TG22PM: TG type 2-2 TGPM, TG23PM: TG type 2-3 TGPM.

TG21HS: TG type 2-1 TGHS, TG22HS: TG type 2-2 TGHS, TG23HS: TG type 2-3 TGHS.

Figure 4. Alignment of five TG type 3 modules. The alignment of the five TG type 3 modules of full-length thyroglobulin from *Petromyzon marinus* (TGPM)and *Homo sapiens* (TGHS) was conducted according to Parma y col. [1987], Malthíery y col. [1987], Molina y col. [1996], van Graaf y col., [2001].   All six cysteine are conserved in the TG type 3-a modules, while all eight cysteines are preserved in the TG type 3-b modules of TGPM and TGHS. TG type 3-a1 in TGPM contains an extra cysteine.

TG3a1PM: TG type 3-a1 TGPM, TG3b1PM: TG type 3-b1 TGPM, TG3a2PM: TG type 3-a2 TGPM, TG3b1PM: TG type 3-b2 TGPM, TG3a3PM: TG type 3-a3 TGPM.

TG3a1HS: TG type 3-a1 TGHS, TG3b1HS: TG type 3-b1 TGHS, TG3a2HS: TG type 3-a2 TGHS, TG3b2HS: TG type 3-b2 TGHS, TG3a3HS: TG type 3-a3 TGHS.

TGPM^567

Region I

TGPM^1746

Region I   Region II   Region III   Region IV

TGPM^2475

Region I   Region II   Region III   Region IV

TGHS

Region I   Region II   Region III   Region IV

MatchMaker

Tertiary structure view

a1   b1   c1

Surface view

a2   b2   c2

Figure 5. Structural alignment of thyroglobulins from *Petromyzon marinus* and *Homo sapiens.* The structural alignment of **thyroglobulins** TGPM[567], TGPM[1747], and TGPM[2475] from *Petromyzon marinus* with **Homo sapiens thyroglobulin (TGHS)** is presented. The **top section** displays the **homology modeling** of TGPM[567], TGPM[1747], and TGPM[2475]. The **middle section** shows the **TGHS model**. The **bottom section** illustrates the **superposition of thyroglobulin monomers** (TGPM[567] **with TGHS,** TGPM[1747] **with TGHS, and** TGPM[2475] **with TGHS**) across the **four regions (I, II, III, IV)** of the **classical model,** using the **Matchmaker command** in **UCSF ChimeraX** (https://www.cgl.ucsf.edu/chimera/). Additionally, in **a1, b1, and c1**, the **superposition of monomers** is displayed **without the surface**, while in **a2, b2, and c2**, it is shown **with the surface**.

TGPM567   **TGPM1746**   **TGPM2475**

Region I

Region I   Region II   Region III   Region IV

Region I   Region II   Region III   Region IV

MatchMaker

Tertiary structure view

a1   b1   c1

Surface view

a2   b2   c2

Figure 6. Structural alignment of thyroglobulins from *Petromyzon marinus*. The structural alignment of thyroglobulins TGPM567, TGPM1747, and TGPM2475 from *Petromyzon marinus* is presented. The top section displays the homology modeling of TGPM567, TGPM1747, and TGPM2475. The middle and bottom sections illustrate the superposition of thyroglobulin monomers, including TGPM567 with TGPM1747, TGPM567 with TGPM2475, and TGPM1747 with TGPM2475, across the four regions (I, II, III, IV) of the classical model, using the Matchmaker command in UCSF ChimeraX (https://www.cgl.ucsf.edu/chimera/). Additionally, in a1, b1, and c1, the superposition of monomers is displayed without the surface, while in a2, b2, and c2, it is shown with the surface.

Figure 7. Structural alignment of full-length thyroglobulins from *Petromyzon marinus* and *Homo sapiens.* The structural alignment of full-length **thyroglobulins** from *Petromyzon marinus* **(TGPM)** and *Homo sapiens* **(TGHS)** is presented. The **top section** displays the **homology modeling** of **TGPM and TGHS.** The **bottom section** illustrates the **superposition of thyroglobulin monomers (TGPM with TGHS)** across the **four regions (I, II, III, IV)** of the **classical model**, using the **Matchmaker command** in **UCSF ChimeraX** (https://www.cgl.ucsf.edu/chimera/). Additionally, the **left side** shows the **superposition of monomers without the surface**, while the **right side** displays the **superposition with the surface**.

a

NID-2 PM

NID-2 HS

SP | **G1** | **linker** | **G2** | **rod-like** | **G3**

SP | Nido | EGF | Nidogen G2 beta-barrel | EGF | EF(| EGF EGF | TG1 | LD LL DL LL DL LL DL LL DL

b

NID-1 TG type 1-1 PM

NID-1 TG type 1-1 MM

NID-1 TG type 1-1 HS

NID-2 TG type 1-1 PM

NID-2 TG type 1-1 MM

NID-2 TG type 1-1 HS

NID-2 TG type 1-2 MM

NID-2 TG type 1-2 HS

TG TG type 1-1 PM

TG TG type 1-2 PM

TG TG type 1-1 HS

TG TG type 1-2 HS

Very high (pIDDT>90)   Confident (90>pIDDT)   Low (70>pIDDT<50)   Very low (pIDDT<50)

Figure 8. Three-dimensional atomic structure of nidogen 2, and TG type 1 modules using AlphaFold server. (a) The 3D structure of nidogen 2 (NID-2) from *Petromyzon marinus* and *Homo sapiens*. Arrows indicate the TG type 1 modules. **(b)** The modeled structure of TG type 1-1 and TG type 1-2 modules of nidogen 1 (NID-1), nidogen 2 (NID-2), and thyroglobulin (TG) from *Petromyzon marinus* (PM), *Mus musculus* (MM), and *Homo sapiens* (HS). The pLDDT (predicted Local Distance Difference Test) provides a per-atom confidence estimate on a 0-100 scale, where a higher value indicate greater reliability. Regions with low pLDDT scores may correspond to unstructured or highly flexible areas.

```
TG11N1PM   KST │CE│ 25aa │YVPQC│ 3aa │GTFLPVQC│ 6aa │CWCVD│ 2aa GQE 2aa GTR 8aa │C│
TG11N1MM   KTR │CQ│ 23aa │FVPQC│ 3aa │GHYVPTQC│ 6aa │CWCVD│ 2aa GRE 2aa GSR 8aa │C│
TG11N1HS   KTR │CQ│ 23aa │FVPEC│ 3aa │GHYAPTQC│ 6aa │CWCVD│ 2aa GRE 2aa GTR 8aa │C│
TG11N2PM   RTP │CE│ 22aa │YVPQC│ 3aa │GAFRPVQC│ 6aa │CWCVT│ 2aa GQE 2aa GTR 8aa │C│
TG11N2MM   LKP │CE│ 17aa │HIPQC│ 3aa │GNFVPLQC│ 6aa │CWCVD│ 2aa GHE 2aa GTQ 9aa │C│
TG11N2HS   LTP │CE│ 17aa │HIPQC│ 3aa │GNFLPLQC│ 6aa │CWCVD│ 2aa GHE 2aa GTQ 9aa │C│
TG11PM     ASA │CE│ 12aa │HVPQC│ 3aa │GAFRPLQC│ 6aa │CWCVD│ 2aa GEE 2aa GTR 8aa │C│
TG11HS     LRP │CE│ 12aa │YVPQC│ 3aa │GSFQTVQC│ 6aa │CWCVG│ 2aa GSE 2aa GSR 7aa │C│
TG12N2MM   RTV │CE│ 18aa │YVPQC│ 3aa │GHFIPLQC│ 6aa │CWCVD│ 2aa GRE 2aa GTR 8aa │C│
TG12N2HS   PTI │CE│ 18aa │YVPQC│ 3aa │GHFIPLQC│ 6aa │CWCVD│ 2aa GRE 2aa GTR 8aa │C│
TG12PM     LSF │CQ│ 18aa │HLPQC│ 3aa │GRYRPAQV│ 5aa │GWCVD│ 2aa GME 2aa GTR 7aa │C│
TG12HS     LSF │CQ│ 18aa │YLPQC│ 3aa │GDYAPVQC│ 6aa │CWCVD│ 2aa GME 2aa GTR 7aa │C│
```

Figure 9. Alignment of TG type 1 modules across species. The alignment of TG type 1 modules from *Petromyzon marinus* (PM), *Mus musculus* (MM), and *Homo sapiens* (HS)—including nidogen 1 (NID-1) and nidogen 2 (NID-2), as well as full-length thyroglobulins from *Petromyzon marinus* (TGPM) and *Homo sapiens* (TGHS)— was conducted according to Parma et al. (1987), Malthiéry et al. (1987), Molina et al. (1996), and van Graaf et al. (2001). The six cysteines are conserved across all modules.
TG11N1PM: PM NID-1 TG type 1-1, TG11N1MM: MM NID-1 TG type 1-1, TG11N1HS: HS NID-1 TG type 1-1, TG11N2PM: PM NID-2 TG type 1-1, TG11N2MM: MM NID-2 TG type 1-1, TG11N2HS: HS NID-2 TG type 1-1, TG11PM: TGPM TG type 1-1, TG11HS: TGHS TG type 1-1, TG12N2MM: MM NID-2 TG type 1-2, TG12N2HS: HS NID-2, TG type 1-2, TG12PM: TGPM TG type 1-2, TG12HS: TGHS TG type 1-2.

Figure 10. Structural alignment of TG type 1-1 modules. The structural alignment of TG type 1-1 module from *Homo sapiens* nidogen 2 (NID-2 HS) and thyroglobulin (TGHS) was performed using the Matchmaker command in UCSF ChimeraX (https://www.cgl.ucsf.edu/chimera/). Interestingly, the TG type 1-1 module of NID-2 HS exactly matches the TG type 1-1 module of TGHS (dotted circle). In contrast, the TG type 1-2 module of NID-2 HS does not align with TG type 1-2 module of TGHS. The four regions (I, II, III, IV) of the classical model of TGHS are displayed.

Figure 11. Flow chart of thyroglobulin complexation (stages a, b, c, d, e, f, g, and h). The flow chart illustrates the complexation of thyroglobulin, tracing its progression from nidogen to the acquisition of eleven TG type 1 modules and the insertion of non-repetitive linker and hinge domains.

Figure 12. Flow chart of thyroglobulin complexation (stages i, j, k, d, I, II, m, n and p). The flow chart illustrates the complexation of thyroglobulin, tracing its progression from the acquisition of five TG type 3 modules and three TG type 2 modules to the fusion of the ChEL domain. The four regions (I, II, III, IV) of the classical model of TGHS are displayed in botton.