Visual Working Memory Guides Attention Rhythmically

Jiachen Lu[a,b,c,1,*]; Yaochun Cai[b,c,d,e,1]; Xilin Zhang[b,c,d,e*]

[a]Research Center of Adolescent Psychology and Behavior, School of Education, Guangzhou University, Guangzhou, Guangdong 510006, China

[b]Key Laboratory of Brain, Cognition and Education Sciences, Ministry of Education, South China Normal University, 510631 Guangzhou, China

[c]School of Psychology, South China Normal University, 510631 Guangzhou, China.

[d]Center for Studies of Psychological Application, Guangdong Key Laboratory of Mental Health and Cognitive Science, South China Normal University, 510631 Guangzhou, China

[e]Philosophy and Social Science Laboratory of Reading and Development in Children and Adolescents, Ministry of Education, South China Normal University, 510631 Guangzhou, China

[1]These authors contributed equally

*Correspondence: ljcps@gzhu.edu.cn (J. L.); xlzhang@m.scnu.edu.cn (X. Z.)

Key Laboratory of Brain, Cognition and Education Sciences, Ministry of Education, School of Psychology, South China Normal University, Guangzhou, Guangdong, China

19   Tel: + (86)186-8283-9781

20   Website: https://xlzhanglab.scnu.edu.cn/

Abstract

How does internal representation held in visual working memory (VWM), known as the attentional template, guide attention? A longstanding debate concerns whether only one (Single-Item-Template theory) or multiple (Multiple-Item-Template theory) items serve as attentional templates simultaneously. Here we propose a Rhythmic-Item-Template hypothesis, successfully reconciling these seemingly contradictory theories. Using the classical VWM-guided attention task, we found that two VWM items alternately dominate behavioral guidance in theta-rhythmic (4–8 Hz), with anti-correlated activation states in time, and more importantly, this rhythmic oscillation was not driven by the retro-cue processing. Neural recordings revealed that occipital alpha-oscillation (8–14 Hz) governed item-specific prioritization and its amplitude closely tracked subjects' behavioral guidance, while frontal theta-oscillations phase-led and coupled with occipital alpha-oscillations during the item transition. Our Rhythmic-Item-Template results not only resolve previous Single-Item-Template versus Multiple-Item-Template debate but also advance our understanding of how distributed brain rhythms coordinate flexible resource allocation in multi-item memory systems.

Keywords: visual working memory, attention, rhythmic-item-template, theta-oscillation, alpha-oscillation

Introduction

The capacity of visual working memory (VWM) to guide attention through multiple internal templates remains contested(Frătescu, Van Moorselaar, & Mathôt, 2019). While VWM can store several items(Constantinidis & Klingberg, 2016; Cowan, 2001; Oberauer, Farrell, Jarrold, & Lewandowsky, 2016), behavioral studies conflict: some suggest only one item dominantly guides attention at any moment(Büsel, Pomper, & Ansorge, 2019; Olivers, Peters, Houtkamp, & Roelfsema, 2011), whereas others report parallel guidance by multiple templates(Kerzel & Witzel, 2019; Kristjánsson & Kristjánsson, 2018). Hollingworth and Beck(Hollingworth & Beck, 2016) revealed that distractors matching either of two target colors elicited equivalent attentional capture, supporting dual templates. However, attenuated capture magnitude compared to single-item conditions exposed a critical paradox: if multiple templates coexist, why does their behavioral efficacy diminish? Three hypotheses emerge: (1) transient dominance of a single item suppresses others, (2) independent but weakened template influences, or (3) rhythmic alternation between items via theta-band oscillations (4 – 8 Hz). Critically, the third hypothesis posits that limited attentional resources are dynamically allocated through cyclical prioritization rather than static competition—a mechanism aligning with the oscillatory nature of neural processes(Fiebelkorn, Pinsk, & Kastner, 2018; Rufin VanRullen, 2016).

This oscillatory framework is rooted in attention's discrete temporal dynamics. When monitoring two locations, human attention rhythmically samples each location at 4–

10 Hz, with asynchronous peaks between sites — a "sequential attentional

spotlight"(Landau & Fries, 2012). Neural recordings demonstrate that alpha-band

suppression (8–14 Hz), reflecting attentional engagement, alternates between objects

every ~200 ms (theta rhythm(Jia, Liu, Fang, & Luo, 2017)). These findings imply a

conserved theta-rhythmic mechanism for resolving representational competition,

whether for external stimuli or internal VWM representations. Recent studies extend

this to VWM: multi-item retention exhibits 6 – 10 Hz oscillatory patterns(Chota, Leto,

van Zantwijk, & Van der Stigchel, 2022; Peters, Kaiser, Rahm, & Bledowski, 2021;

Pomper & Ansorge, 2021), yet debates persist about whether these rhythms reflect

true prioritization dynamics or spatiotemporal confounds (Huang, Jia, Han, & Luo,

2018; Liu, Dolan, Kurth-Nelson, & Behrens, 2019).

Moreover, Previous evidence for rhythmic memory processing primarily stems from

behavioral measures, lacking direct neural evidence of stimulus-specific modulation.

While alpha oscillations (8 – 14 Hz) in visual cortex track VWM item prioritization(I.

E. De Vries, Van Driel, & Olivers, 2017; Pavlov & Kotchoubey, 2022; Wolff, Jochim,

Akyurek, & Stokes, 2017), they likely enhance signal-to-noise ratios through selective

neural recruitment rather than directly governing task goals(Benedek, Schickel, Jauk,

Fink, & Neubauer, 2014; Klimesch, 2012; Peylo, Hilla, & Sauseng, 2021). Conversely,

frontal theta oscillations (4 – 8 Hz) coordinate goal-directed behaviors and task

switching(Helfrich, Huang, Wilson, & Knight, 2017; Josselyn & Tonegawa, 2020) ,

exerting top-down control over sensory regions via phase synchronization(Berger et

al., 2019; van Driel, Gunseli, Meeter, & Olivers, 2017). Critically, fronto-posterior

network coupling mediates control shifts in VWM: increased frontal theta power

predicts contralateral occipital alpha suppression during priority transitions(Audrain et

al., 2020; I. E. J. de Vries, Slagter, & Olivers, 2020; I. E. J. de Vries, van Driel,

Karacaoglu, & Olivers, 2018). This supports a mechanistic hypothesis: frontal theta

oscillations rhythmically drive spatially distributed alpha oscillations through

cross-frequency coupling, dynamically activating multiple VWM items in

theta-rhythmic cycles.

To test this theoretical proposition, we designed and conducted three experiments

using a memory - search paradigm that effectively rules out strategic control(Soto,

Humphreys, & Rotshtein, 2007). Experiment 1, with dense temporal sampling,

revealed 7 Hz anti - phasic oscillations in attentional capture between two VWM

items. Experiment 2, by implementing individualized temporal analysis to eliminate

retrospective cueing artifacts, replicated the 7 Hz behavioral rhythm. To further

explore the neural mechanisms and comprehensively validate our theoretical

hypothesis, we carried out Experiment 3, in which electroencephalography (EEG) was

employed to record and analyze brain electrical activity with high temporal resolution.

EEG results demonstrated that activation in the occipital alpha band was closely

associated with behavioral performance and exhibited significant phase coupling with

prefrontal theta oscillations. Additionally, during the memory maintenance phase, the

coupling strength of alpha - theta between the bilateral visual cortex and the prefrontal

cortex alternated in a theta - rhythmic pattern in leading. These results collectively

establish frontally driven theta - alpha coupling as a mechanism for rhythmically

allocating attentional access to multiple VWM items, reconciling the divergence

between single - and dual - template theories through a dynamic, oscillation - based

model.

Study overview

Experiment 1: theta-rhythmic oscillation of visual working memory

25 participants engaged in Experiment 1. As shown in Figure 1C, subjects were asked

to remember two memory items that appeared simultaneously. During the memory

retention interval, a cue randomly instructed to one of the items, directing internal

attention towards the cued stimulus. To examine the time-course of VWM, the

response interface appeared at various time intervals following the cue, with a high

temporal resolution (stimulus onset asynchrony, SOA: starting at 233 ms and

increasing in increments of 33 ms up to 833 ms). The response interface required

participants to complete either a search task (80% of trials) or a recall task (20% of

trials). In the search task, participants were asked to search a target square (with a gap

at the top/bottom) while disregarding a distractor square (with a gap on the left/right).

To ensure the search target and the memory item did not share the same spatial

location, the target and distractor were positioned vertically relative to the

participants' point of gaze. Only one of the target and distractor matched the color of

memory items, or neither did. The attentional capture effect was measured as the

difference in response time between the distractor matching the memory items and the

target matching the memory items. In the recall task, participants were asked to

determine whether a presented color corresponded to one of the memory items,

thereby assessing the accuracy of item retention.

Figure 1A: Hypothetical models illustrating the activation patterns of two memory

items during the memory retention phase. Hypothesis 1 posits a single template,

Hypothesis 2 suggests multiple templates, and Hypothesis 3 proposes dynamic

templates. B: Behavioral results, showing the attentional capture effect size

(calculated as the difference in response time between the distractor matching the

memory items and the target matching the memory items) and memory accuracy for

the two memory items. C: Experimental procedure. A cue stimulus randomly indicated

one of the two memory items. In 80% of the trials, participants performed a search

task to identify the item with a gap facing upward or downward. In 20% of the trials,

participants performed a recall task to determine whether the probed item matched

one of the two memory colors.

Initially, we calculated the recall accuracy for both memory items, as depicted in

Figure 1B. The recall accuracy for cued items ($t(24) = 91.60$; $p < 0.001$, Cohen's d =

37.39) and uncued items ($t(24) = 70.74$; $p < 0.001$, Cohen's d = 28.88) significantly

exceeded the chance level (0.5), demonstrating effective memory retention for both

item types. Subsequently, we examined the attentional capture effects associated with

these memory items. The analysis revealed that the magnitude of the attentional

capture was significantly above zero for both cued items ($t(24) = 6.16$; $p < 0.001$,

Cohen's d = 2.51) and uncued items ($t(24) = 8.02$; $p < 0.001$, Cohen's d = 3.27),

indicating that both items effectively captured attention. These findings align with

previous studies(Hollingworth & Beck, 2016), confirming that human brain is capable

of retaining multiple memory items simultaneously, with each item significantly

influencing attentional processes.

Figure 2. A: Line graph. The attentional capture effect size for the cued and uncued

items across different time intervals (SOA), calculated as the difference in response

time between the invalid and valid conditions. B: Spectrum plot. The red line

represents the amplitude of the values from A at different frequencies; the gray line

indicates the 95th percentile from the permutation test; *: p < 0.05. C: Phase-locking

value (PLV). The red line shows the PLV values at 7 Hz for the cued and uncued items,

representing the average phase difference across all participants; gray circles

indicate the 0-95 percentile range from the permutation test; blue hollow circles

represent the phase differences for individual participants between the two items.

With respect to the central question of our study, we found clear evidence for memory

item sampling as a preferred template. As shown in Figure 2A, we examined the

attentional capture effects of cued and uncued items separately for different temporal

SOA conditions. It was found that the capture effects of these two items alternated in

dominance rather than remaining stable ($F(1, 19) = 4.67$, $p < 0.001$, $\eta2 = 0.16$).

Specifically, the capture effect of cued items was significantly greater than that of

uncued items at SOAs of 233ms ($t(24) = 2.72$, $p = 0.03$, Cohen's $d = 1.11$), 633ms

($t(24) = 2.37$, $p = 0.03$, Cohen's $d = 0.97$) and 800ms ($t(24) = 3.53$, $p = 0.002$, Cohen's

$d = 1.44$), while the capture effect of uncued items was significantly greater than that

of cued items at SOAs of 300ms ($t(24) = 2.97$, $p = 0.007$, Cohen's $d = 1.21$), 333ms

($t(24) = 2.14$, $p = 0.04$, Cohen's $d = 0.87$), 400ms ($t(24)= 2.49$, $p = 0.02$, Cohen's $d = $

1.02), 433ms ($t(24)=2.37$, $p = 0.03$, Cohen's $d = 0.97$) and 567ms ($t(24)=2.72$, $p = $

0.02, Cohen's $d = 1.11$).

In addition, as shown in Figure 2B, the Fourier transformation was applied to the time

course of the item-based benefit to determine the temporal frequency of these

fluctuations. The item-based benefit was calculated as the difference in capture effects

between the cued and uncued item for equidistant time intervals ranging from 200 ms

to 833 ms. The amplitude spectrum analysis of frequencies ranging from 1 Hz to 15

Hz (Figure 2B) revealed a significant peak at 7 Hz ($p < 0.05$, FDR corrected). This

finding implies that the rhythmic fluctuations in attentional capture associated with

VWM are likely driven by an oscillatory mechanism within the theta frequency range,

supporting the proposed sampling frequency in VWM.

Consistent with the hypothesis that only a single memory item serves as a prioritized

template at any given moment, the speed at which cued and uncued items are

accessed should exhibit a negative correlation, as demonstrated in previous research

on visuospatial and object-based attention(Fiebelkorn, Saalmann, & Kastner, 2013;

Landau & Fries, 2012). To investigate the possibility that attention within VWM

alternates between items, we analyzed the Fourier coefficients at 7 Hz corresponding

to the attentional capture effects for both cued and uncued items and subsequently

calculated their phase-locking values (PLV(Fiebelkorn et al., 2013)). As shown in

Figure 2C, the phase-locking between the 7 Hz rhythms of the two items was found to

be significant (PLV = 0.402, p < 0 .05), displaying an average phase angle difference

of 118° (95% CI from 48° to 188°). This anti-phase relationship between the

oscillations supports the notion that internal attention alternately samples items in

working memory.

Experiment 2: Behavioral oscillation without cueing

17 participants took part in Experiment 2, which closely replicated the design of

Experiment 1 with one critical modification: no retro-cue was presented after the two

memory items. Instead, the probe appeared directly after one of three fixed

stimulus-onset asynchronies (SOAs: 233 : 33 : 860 ms; see Figure 3A). This

adjustment was implemented to rule out the possibility that working memory

oscillations were driven by cue processing rather than intrinsic maintenance

mechanisms.

Figure 3. A: Experimental procedure. Two memory items are presented

simultaneously without any post-cue prompts. In 80% of the trials, participants

performed a search task to identify the item with a gap facing upward or downward.

In 20% of the trials, participants performed a recall task to determine whether the

probed item matched one of the two memory colors. B: Behavioral results, showing

the attentional capture effect size and memory accuracy for the two memory items.

As illustrated in Figure 3B, the recall accuracy for both the left memory item ($t(16) =$

53.40, $p < 0.001$, Cohen's $d = 13.35$) and the right memory item ($t(16) = 51.10$, $p <$

0.001, Cohen's $d = 12.78$) significantly exceeded the chance level (0.5), suggesting

effective retention of the memory items by participants. Furthermore, our analysis of

the attentional capture effect of both memory items revealed that the effect size was

notably higher than zero for both the left item ($t(16) = 3.24$, $p = 0.005$, Cohen's $d =$

0.81) and the right item ($t(16) = 3.21$, $p = 0.005$, Cohen's $d = 0.80$), demonstrating

that both memory items were effective in capturing attention.

Figure 4. The blue line represents the average across all participants of the Fourier

transforms of the differences in capture effects between left and right memory items

for individual participants; The red line represents the group average of medians from

1000 permutations with Fourier transforms per participant; *: p < 0.05.

Since no retro-cue was presented in Experiment 2, the data lacked a fixed temporal

reference point across participants. Thus, unlike in Experiment 1 (where capture

effects were averaged before Fourier analysis), we performed Fourier transforms

individually for each participant before grand-averaging. Specifically, we computed

the item-based benefit for each participant by subtracting the capture effect of the

right memory item from that of the left item, evaluated at equidistant time intervals

(233 – 833 ms). For each participant, we conducted amplitude spectrum analysis (1–

15 Hz) on these time series. The resulting spectra were then grand-averaged across

participants (blue line, Figure 4). To assess whether spectral power exceeded chance

238   levels, we generated a null distribution by (1) randomly shuffling each participant's

239   time series, (2) recomputing the Fourier transform, and (3) repeating this procedure

240   1,000 times. The median amplitude of these surrogate data served as the chance

241   baseline. A paired-samples T-test comparing the original amplitudes against this null

242   distribution revealed significant oscillatory power at 7 Hz (p < 0.05, corrected).

Experiment 3: Neural mechanisms of visual working memory oscillations

In Experiment 3, 27 participants were recruited, with 3 participants excluded due to

excessive artifacts in EEG data. This experiment differed from Experiment 1 in that

the search task and the cue were presented at a fixed interval of 1500 ms, while the

interval between the memory task and the probe varied randomly between 200 ms and

2000 ms (see Figure 5B). All other aspects of the experiment remained consistent with

those of Experiment 1. Additionally, EEG data were recorded simultaneously from

participants.

Figure 5. A: Behavioral results, showing the attentional capture effect size and

memory accuracy for the two memory items. B: Experimental procedure. A cue

stimulus randomly indicated one of the two memory items. In 80% of the trials,

participants performed a search task to identify the item with a gap facing upward or

downward. In 20% of the trials, participants performed a recall task to determine

whether the probed item matched one of the two memory colors.

First, we calculated the recall accuracy for both memory items. As shown in Figure

5A, cued (t(23) = 45.70, p < 0.001, Cohen's d = 19.06) and uncued items (t(23) =

23.56, p < 0.001, Cohen's d = 9.82) were both recalled correctly at a significantly

greater level than the chance level (0.5), indicating successful retention of both types

of items. Additionally, we analyzed the attentional capture effects of the two

memory items. The findings revealed that the attentional capture effects for cued (t(23)

= 6.45, p < 0.001, Cohen's d = 2.69) and uncued items (t(23) = 5.61, p < 0.001,

Cohen's d = 2.34) were significantly greater than zero, indicating that both types of

memory items effectively captured attention.

Figure 6. A: Time-frequency results of the contralateral occipital lobe (PO7/8) for

cued items, showing significant alpha band activation (8-14 Hz) during the memory

retention phase. B: Time-frequency results of the contralateral occipital electrodes

(PO7/8) for uncued items, with a similar activation pattern to that observed for cued

items.

Subsequently, an individual time-frequency analysis was conducted on the EEG data

during the memory retention phase before the search task. This analysis delved into

the temporal and spectral dynamics across frequencies from 1 to 30 Hz and time

intervals from -500 to 2000 ms for each participant. Consistent with prior

studies(Rufin VanRullen & Macdonald, 2012), (I. E. De Vries et al., 2017),

pronounced alpha band activity (8 - 14 Hz) was observed in regions contralateral

(PO7/8) to the memorized items (see Figure 6).

Figure 7. The attentional capture effect size of cued items was positively correlated

with the mean amplitude of the alpha band activity during the last 200 ms of the

memory retention phase, which occurs 200 ms before the onset of the search target.

To evaluate the impact of alpha band activity on attentional capture, correlation

coefficients between behavioral performance and alpha power were calculated.

Specifically, the correlation between the alpha power contralateral to the items 200

milliseconds before the onset of the search task and the capture effect of the item was

analyzed. As shown in Figure 7, a significant positive correlation was found for the

cued item (r = 0.43, p = 0.03) , indicating that stronger alpha band activity correlated

with an enhanced capture effect. These findings imply that alpha activation patterns

prior to the onset of search stimulus can predict the effectiveness of VWM in guiding

the search. Essentially, the alpha band may reflect the priority state within VWM,

aligning with findings from previous research(I. E. De Vries et al., 2017; I. E. J. de

Vries, van Driel, & Olivers, 2019).

Figure 8. Time-frequency results, showing the difference in contralateral electrode

activity (PO7/PO8) for same-color and same-location items under cued and uncued

conditions.

Analysis of power differences between cued and uncued items unveiled a distinct

pattern of alpha inhibition followed by rebound activation within the first 200 ms, as

illustrated in Figure 8. Remarkably, this alternation in alpha band power persisted

throughout the memory retention phase and showed a significant rhythm of

approximately 4 Hz (Figure 10A, p < 0 .05, FDR corrected).

Figure 9. A: Topographical maps. Cross-frequency coupling between the alpha phase

at the white electrode points and the theta phase at other electrode points (the left two

maps), and the theta phase at the white electrode points and the alpha phase at other

electrode points (the right map). B: Functional connectivity map. Cross-frequency

coupling between the alpha phase at the contralateral electrodes (PO7/PO8) of the

two memory items and the theta phase at the frontal electrode (Pz).

Notably, both this study and prior research have identified that the alpha band decodes

the preferred memory template. Intriguingly, an observation common to our study and

earlier work is that the alternating frequency of this preferred template aligns with the

theta rhythm range (4-7 Hz). To explore the potential impact of theta oscillations in

different brain regions on alpha oscillations in regions contralateral to the memorized

items, we employed the weighted phase lag index (wPLI) to measure the interaction

between alpha oscillations at PO7 and PO8 and theta oscillations in other brain

regions. The findings, illustrated in Figure 9A, indicated significant coupling between

PO7/8 and the prefrontal regions. Additionally, wPLI calculations revealed significant

coupling for theta oscillations in the prefrontal cortex (Fz) and alpha oscillations in

the visual regions. Fascinatingly, by assessing the alpha-theta coupling strength

between the posterior regions contralateral to the two items (PO7/8) and the prefrontal

regions (Fz) on a moment-to-moment basis during the memory retention period, we

found that the coupling strength of the two items alternated in lead. To further define

the temporal frequency characteristics of these alternations, Fourier transformation

was applied to the differences in wPLI associated with the two items. An amplitude

spectrum analysis across frequencies from 1 Hz to 50 Hz (Figure 10B) identified a

significant peak at 4 Hz ($p < 0.05$, FDR corrected), underscoring the rhythmic

interconnection between these neural activities.

Figure 10. A: Spectrum results. Power spectrum of the difference in contralateral

alpha power between the two memory items during the retention phase across 0-50

Hz. B: Spectrum results. Power spectrum of the difference in cross-frequency coupling

334 *(wPLI) between the alpha phase at contralateral electrodes (PO7/PO8) and the theta*

335 *phase at the frontal electrode (Fz) across 0-50 Hz during the retention phase. The*

336 *gray shaded area represents the 0-95th percentile range from the permutation test; *:*

337 *$p < 0.05$.*

Discussion

This study employed the function of VWM as a preferred template to automatically capture attention and integrated EEG techniques to systematically investigate the mechanisms underlying the representation of two memory items that occurred simultaneously or in close temporal proximity during the memory retention phase. Our findings revealed several key insights. Firstly, we observed that the ability of the memory items to capture attention alternated over time, displaying opposite trends (Experiment 1). This indicates that the preference for the memory items alternated rather than occurring simultaneously. Power analysis showed alternating rhythms within the 4-7 Hz range (Experiment 1-3), corresponding to theta oscillations. Secondly, during the memory retention phase, the strength of alpha band oscillations, induced by cued and uncued items, alternated in dominance throughout the retention period. Thirdly, we observed strong rhythmic alternations in the coupling strength between the alpha band of both posterior regions and the theta band of the prefrontal region. The coupling exhibited a leading frequency of approximately 4 Hz. In conclusion, our findings suggest that memory is maintained by alternately processing multiple items within the theta rhythm, rather than continuously entangling a single object. Furthermore, the posterior alpha oscillations reflect the prioritization of memory items and exhibit a close association with theta oscillations in the prefrontal cortex.

Traditional research on working memory has often adopted a static viewpoint,

positing that its performance remains constant throughout the retention phase. This

perspective, however, is challenged by the findings of the present study, which

investigates the ability of two memory items to capture attention under various

Stimulus-Onset Asynchrony (SOA) conditions. Interestingly, the study reveals that the

attentional capture abilities of these memory items alternate in dominance. Spectral

analysis of the SOA function identified significant rhythmic fluctuations at 7 Hz, with

opposing trends. This pattern suggests that, in scenarios requiring the maintenance of

two working memory items for a task, these items alternately serve as the dominant

template during the retention phase.

This discovery aligns with the existing literature on the rhythmic processing of

external stimuli. Specifically, it has been well-documented that attention oscillates

within the theta rhythm (4-8 Hz) across various task types, including those based on

spatial attention (Jensen & Mazaheri, 2010; Jiang, He, & Zhang, 2024), feature

attention(Re, Inbar, Richter, & Landau, 2019), and object attention(Fiebelkorn et al.,

2018; Fiebelkorn et al., 2013; Helfrich et al., 2018). Moreover, rhythmic attentional

modulation has been observed in auditory attention(Ho, Burr, Alais, & Morrone, 2019;

Ho, Leung, Burr, Alais, & Morrone, 2017) , in the effects of top-down predictions on

brightness perception(Han & VanRullen, 2017), and in the process of visual feature

binding(Nakayama & Motoyoshi, 2019). These findings collectively highlight that

attention operates within a theta rhythm, extending beyond external stimuli to include

the internal processing of memory information, such as that involved in working

memory(Peters et al., 2021; Pomper & Ansorge, 2021).

The concept of a dynamic template mechanism emerges as a novel resolution to the

ongoing debate between the single-template and multi-template hypotheses in

working memory research(Frătescu et al., 2019; Olivers et al., 2011; van Moorselaar,

Theeuwes, & Olivers, 2014). This mechanism posits that, when maintaining multiple

memory items, an individual employs a singular, dynamically shifting template to

represent these items. This flexible approach not only aligns with the observed

phenomenon of multiple memory items capturing attention, but also provides a new

perspective on the discourse on how working memory operates, bridging a critical gap

in the existing literature.

The current study reveals that during the memory retention phase, the dominance of

posterior lateralized alpha oscillation alternates between two memory items and

significantly correlates with behavioral outcomes. This finding aligns with extensive

evidence suggesting that alpha oscillations are crucial in modulating visual processing,

going beyond mere sensory involvement to encompass functional roles in sensory

gating and top-down control of preparatory activity(Chen et al., 2023; Gazzaley &

Nobre, 2012; Jensen & Mazaheri, 2010; Klimesch, 2012). Notably, alpha band

activity modulation is essential for prioritizing both incoming sensory data and stored

memory content(D. Gresch, Behnke, van Ede, Nobre, & Boettcher, 2025; Van Diepen,

Foxe, & Mazaheri, 2019). A seminal example is a study in which subjects memorized

two sequentially presented items, demonstrating the capability of posterior lateralized

alpha oscillations to differentiate the memory item relevant to the current task,

thereby elucidating the role of oscillations in managing the priority of concurrent

task-relevant items (I. E. J. de Vries et al., 2019).

It's critical to acknowledge that the investigation of posterior alpha oscillations in

working memory, including the design of Experiment 3 in this study's, often involves

spatial differentiation of items(I. E. J. de Vries et al., 2019). Such spatial segregation

allows alpha power modulations to be attributed to specific items through

lateralization, achieved by presenting items in distinct hemifields. This

methodological choice, driven by the spatial specificity of neurons encoding working

memory items, serves as a pragmatic experimental strategy rather than implying that

memory item prioritization solely depends on alpha oscillation lateralization (Foster,

Bsales, Jaffe, & Awh, 2017; Daniela Gresch, Boettcher, Gohil, van Ede, & Nobre,

2024). For example, de Vries et al.(I. E. De Vries et al., 2017) discovered that

posterior, rather than lateralized, alpha oscillations could delineate memory items

related to forthcoming tasks.

The findings of this study, revealing strong phase coupling between posterior alpha

oscillations and theta oscillations in the prefrontal lobe, contribute to the expanding

literature on the intricate interplay between different brain oscillation frequencies and

their roles in cognitive functions. The association of prioritization processes with theta

oscillations in the frontal cortex is well-documented(I. E. J. de Vries et al., 2020; I. E.

J. de Vries et al., 2019; Quentin et al., 2019; Wallis, Stokes, Cousijn, Woolrich, &

Nobre, 2015), underscoring the critical role of frontal cortex in orchestrating

goal-directed behavior(Miller & Cohen, 2001), managing multiple

objectives(Mansouri, Koechlin, Rosa, & Buckley, 2017), and facilitating task

switching(Monsell, 2003).

Particularly during working memory tasks, the frontal cortices play a critical role in

processing and maintaining abstract goal-related representations and task-specific

information. This is supported by the evidence of mixed selectivity in frontal

neurons(Fusi, Miller, & Rigotti, 2016) and their coordination of sensory areas based

on this information(Christophel, Klink, Spitzer, Roelfsema, & Haynes, 2017;

Sreenivasan, Curtis, & D'Esposito, 2014). For instance, fMRI studies have highlighted

enhanced connectivity between frontal regions and posterior task-related sensory

areas specific to working memory representations and those prioritized by

maintenance cues (retrospective cues(Nelissen, Stokes, Nobre, & Rushworth, 2013;

Van Ede & Nobre, 2023)), with this increased connectivity correlating with improved

performance of cued memory representations(Gazzaley & Nobre, 2012; Kuo, Yeh,

Chen, & D'Esposito, 2011).

Such evidence underscores the ability of the frontal cortex to selectively coordinate

the activation of visual cortical working memory representations relevant to the

current task. As outlined in a review by de Vries et al.(I. E. J. de Vries et al., 2020),

alpha modulation facilitates the flexible tracking of changes in the state of VWM.

This top-down control over the current state of VWM is accomplished through

prefrontal theta oscillations, which in turn orchestrate alpha oscillations via a broad

spectrum of cross-frequency interactions, highlighting a sophisticated mechanism of

cognitive control and memory prioritization.

In conclusion, our study aligns with the evolving understanding that oscillatory neural

mechanisms underpin the complex interplay between attention and working memory.

It builds on the premise that external attentional processes are intricately linked with

neural oscillations, while also highlighting the crucial role of internal attention in

modulating working memory representations through rhythmic activity. Our findings

support a framework in which internal attention is mediated by a dynamic interaction

between prefrontal theta oscillations, serving as a top-down control mechanism, and

posterior alpha oscillations, which are instrumental in the selective inhibition or

facilitation of memory items. This synthesis not only challenges conventional static

views of working memory but also proposes a refined model of cognitive processing,

where memory maintenance and attentional prioritization are orchestrated within a

rhythmic neural symphony.

Materials and methods

Participants

A total of 25 participants (9 males, aged 22 ± 1.9 years) took part in Experiment 1, 24 subjects (10 males, aged 20 ± 1.8 years) participated in Experiment 2, and 27 subjects (12 males, aged 21 ± 2.0 years) were involved in Experiment 3. All participants had normal or corrected-to-normal vision and no history of psychiatric or neurological disorders. The experiments were conducted in accordance with the Declaration of Helsinki and received ethical approval from the Research Ethics Committee at South China Normal University (approval date: 2021-04-01). Before the start of the experiment, all participants provided written informed consent. Participants completed the experiment in a dark, quiet and isolated room, with their heads fixed on a head rest and their eyes looking directly at the centre of the screen at a distance of 57 cm from the display.

Stimuli

The same stimuli were used for all three experiments. The memory stimuli were a square in 6 colours: cyan (RGB: 5, 200, 200), red (RGB: 200, 80, 40), yellow (RGB: 200, 200, 5), blue (RGB: 5, 200, 200), purple (RGB: 112, 48, 160), and green (RGB: 50, 200, 50), each with a visual angle of 1.2 × 1.2°. The search stimuli were colored outlined squares (1.2 × 1.2°, 0.2° line thickness), with a 1.2° gap on the top, bottom, left side, or right side. The color of the search stimulus was consistent with that of the

memory stimulus. The cue stimulus is a black square with side length 1.2°. The shape

and color of the detection stimulus are consistent with those of the memory stimuli.

The experiments were programmed using the Psychtoolbox in Matlab 2019b. All

stimuli were presented on a 17-inch CRT monitor with a resolution of 1024×768, a

refresh rate of 60 Hz, and a gray background (RGB: 128, 128, 128).

Tasks

Experiment 1

Each trial began with a 500 ms fixation point, followed by a 1000 ms memory array.

In the memory array, two differently colored memory stimuli were presented to the

left and right at a distance of 3° from the centre of the screen. After a 250ms fixation

point, the cue array was presented for 50 ms, with the cue appearing randomly on the

left or right at a distance of 3° from the centre of the screen. After a pseudo-random

SOA (200 ms - 833 ms), the search array (80% of trials) or the memory detection

array (20% of trials) was presented. In the search array, two search stimuli were

presented directly above or below the centre of the screen at a distance of 6° for 2000

ms, one with a gap facing up or down (target) and the other with a gap facing left or

right (distractor). Participants were asked to respond by pressing the "A" key (gap

facing up) or the "Z" key (gap facing down). In the memory detection array, a

detection stimulus was presented at the centre of the screen for 2000 ms. Participants

were asked to press the "N" key (belong) or the "M" key (not belong) to determine

whether the detection stimulus matched either of the memory stimuli in the memory

array, regardless of the cue. After responding, participants moved on to the next trial,

with a 1000ms interval between trials. To familiarize participants with the task, 15

practice trials were conducted before the formal experiment. Participants completed

30 blocks over four days within one week, with each block containing 150 trials (120

trials for search tasks and 30 trials for memory detection tasks).

Experiment 2

Each trial began with a 500 ms fixation point, followed by a 1000 ms memory array.

In the memory array, two differently colored memory stimuli were presented to the

left and right at a distance of 3° from the centre of the screen. After a pseudo-random

SOA (200 ms - 833 ms), a search array was presented in 80% of the trials, while a

memory detection array was presented in 20% of the trials after a random blank

interval of 200 ms to 2000 ms. The task for the participants was consistent with

Experiment 1. Participants completed 30 blocks over four days within one week, with

each block containing 150 trials (120 trials for search tasks and 30 trials for memory

detection tasks).

Experiment 3

The task for the participants was consistent with Experiment 1. However, in

Experiment 2, high temporal precision EEG technology was utilized, eliminating the

need for dense SOA manipulation to investigate dynamic features. Additionally, to

allow for a longer memory retention time window during analysis, the interval

between the cue array and the search array was fixed at 2000 ms. Furthermore, to

encourage active maintenance of the memory content during the retention phase, the

interval between the cue array and the memory detection array was randomly varied

between 200 ms and 2000 ms. Participants were required to complete 4 blocks, with

each block consisting of 150 trials.

EEG recordings and preprocessing

EEG data was recorded using a cap with 64 electrodes arranged according to the

international 10–20 system (Brain Products, Munich, Germany). The frontal electrode

FCz was utilized as the online reference point, and the AFz electrode was employed

as the ground. All electrodes were amplified using a 0.01–70 Hz online band-pass

filter and continuously sampled at a rate of 1000 Hz per channel.

The offline continuous EEG data was preprocessed using EEGLAB, an open-source

toolbox within the MATLAB environment. Initially, re-reference was conducted by

using the bilateral mastoids TP9 and TP10. Next, all EEG signals underwent a 0.1 Hz

high-pass filter, a 30 Hz low-pass filter, and a 50 Hz notch filter. Subsequently,

independent component analysis (ICA) was applied to each participant;s data to

eliminate components related to eye movements and artifacts. The remaining

components after this process were then projected back into the channel space. We

extracted data from -500 ms to 2000 ms relative to cue stimulus presentation in

Experiment 2.

Data analysis

Behavioral performance analysis

We utilized the difference in reaction times (ΔRT) between the invalid trials (where

the color of the memory item matched that of the interfering item) and the valid trials

(where the color of the memory item matched that of the target item) extracted from

the search array as the capture effects.

Subsequently, we employed a one-sample t-test to separately examine whether the

memory accuracy rates of the two memory items in the three experiments were

significantly higher than the guessing level (50%), and whether the capture effects of

the two memory items were significantly greater than 0. In this process, the memory

accuracy rates and capture effects in Experiment 1 and Experiment 3 were the results

of averaging all the SOA (stimulus onset asynchrony) conditions.

For Experiment 1, we concentrated on exploring how the capture effect of the two

memory items on attention evolved over time. To accomplish this, we carried out a

repeated-measures analysis of variance (ANOVA) with a 2 (cued item vs. uncued item)

$\times$ 20 (all SOAs) design. Post-hoc comparisons were conducted using paired-sample

t-tests to thoroughly investigate the potential changes in the capture effect of the

memory items on attention.

In order to obtain the frequency spectrum of memory-based attention capture over

time, we analyzed the difference in capture effects between the cued and uncued items

across all SOAs at evenly spaced temporal intervals. Subsequently, we performed a

fast Fourier transform (FFT) to estimate the spectral composition, which yielded

power values across 14 frequency bins ranging from 1 Hz to 14 Hz. Regarding the

phase relationship of 7-Hz oscillations between the cued and uncued items, we

conducted separate Fourier transforms for these conditions as previously described.

Then, we calculated the angular difference between the phase angles of the 7-Hz

oscillations for each condition. This angular difference was projected onto the unit

circle in the complex plane and averaged across all participants. The length and angle

of the resulting vector represented the phase-locking value (PLV (Lachaux, Rodriguez,

Martinerie, & Varela, 1999)) and the mean phase difference. To evaluate the statistical

significance, a non-parametric approach was adopted to estimate the probability of the

observed data under the null hypothesis. In 1000 permutation samples, each

time-course was shuffled before the analysis, generating one mean amplitude

spectrum for the memory-based capture effect, and one mean phase difference

between the cued and uncued conditions for each permutation sample. To control the

false discovery rate at 5%, the individual frequency p-values in the amplitude

spectrum were adjusted according to the number of frequency bins, following the

method of Benjamini and Hochberg(Benjamini & Hochberg, 1995).

For Experiment 2, since Experiment 2 adopted a modified approach due to the

absence of retro-cues. In this experiment, we compared the left and right memory

items instead of the cued/uncued items. The item-based attentional benefit was

calculated for each participant by subtracting the right-item capture effect from the

left-item effect at equidistant intervals (233-833 ms). Individual FFTs (1-15 Hz) were

performed on these difference scores before conducting a grand average across all

participants. To determine the significance thresholds, we generated null distributions

by randomly shuffling each participant's time series 1000 times, recomputing the

FFTs for each permutation, and using the median amplitude as the chance baseline. A

paired-samples t-test comparing the original amplitudes against this null distribution

revealed significant 7-Hz oscillatory power ($p < 0.05$, corrected), as depicted in

Figure 4.

Time-frequency decomposition

We used the short-time Fourier transform (STFT) function in Matlab 2019b to Fourier

transform the baseline-corrected segments to obtain the power information, where the

frequency was set to 30 frequency points of equal length from 1 to 30 Hz and the

sliding window was a 200 ms hanning window. The frequency-specific power at each

time point was calculated as the square of the amplitude of the complex signal

resulting from the convolution, determined by the sum of the squares of the real and

imaginary parts. Based on findings from previous studies indicating that memory

stimuli can be characterized by alpha oscillations in the contralateral occipital lobes(I.

E. De Vries et al., 2017), we therefore extracted alpha band data (8-14 Hz) from PO7

and PO8. We separately averaged the power of the contralateral side for the cued and

uncued items. As illustrated in Figure 8, to compare the power induced by cued items

with that induced by uncued items during the memory retention phase, we extracted

the contralateral power for each color during memory retention. The power whe

n cued was subtracted from the power when uncued. Finally, we averaged the results

across all memory colors and participants.

Interregional connectivity

We computed the weighted phase lag index (wPLI) between the alpha band phase at

electrodes PO7 and PO8 and the theta band phase at other electrode. The wPLI

assesses phase-based functional connectivity by measuring the degree of phase

clustering between sites(Cohen, 2014) , while mitigating the influence of random

phase lag, thereby controlling for volume conduction artifacts(Stam, Nolte, &

Daffertshofer, 2007). The results revealed that both electrodes PO7 and PO8 exhibited

the strongest phase coupling with electrodes in the prefrontal region. Following this,

we calculated the wPLI between the theta band at electrode Fz in the prefrontal region

and the alpha band at other electrodes to confirm the specificity of cross-frequency

coupling between the prefrontal regions and visual cortex. Finally we extracted the

wPLI between the electrodes PO7/8 in the contralateral visual region relative to the

cued and uncued items respectively, and the Fz electrode at the forehead.

The correlation between neural activation and behavior

As shown in Figure S1, a clear activation of the alpha band was observed during the

memory retention phase. To investigate whether this activation was related to

behavioral performance, we calculated the alpha activation strength in the 200 ms

preceding the search task. Specifically, we computed the average amplitude within the

8-14 Hz frequency range during the 1800-2000 ms time window of the retention

phase, contralateral to the cue. This was then correlated with the capture effect size of

the cued items using Pearson's correlation, as shown in Figure S2.

Frequency spectrum analysis

As shown in Figure 7, the alpha power induced by cued and uncued items alternated

in dominance during the memory retention phase. To quantify this alternation rhythm,

we employed a spectral analysis. Specifically, we averaged the power difference

between the cued and uncued items within the 8-14 Hz range during the memory

retention phase. The data were then downsampled to 100 Hz, using a 10 ms window

for averaging. The resulting one-dimensional time-series data for the 0-2000 ms

memory retention phase were subjected to amplitude spectrum analysis over

frequencies ranging from 1 Hz to 50 Hz. A permutation test was used to assess

significance. In the permutation test, we randomly shuffled the time series and

performed Fourier transformations, repeating this process 1000 times. If the power of

the original time series ranked within the top 5% of these random permutations, it was

considered a significant non-random event. Similarly, we analyzed the difference in

the weighted phase-lag index (wPLI) between the contralateral sides of the two items

and the prefrontal area during the retention phase, using amplitude spectrum analysis

of frequencies ranging from 1 Hz to 50 Hz. In Experiment 3, we averaged the

predicted difference values over the visual temporal dimension and performed

amplitude spectrum analysis of the memory time series over frequencies ranging from

1 Hz to 50 Hz, followed by a permutation test.

Acknowledgements

We acknowledge the subjects for their contribution to this study. This work was

supported by National Natural Science Foundation of China (32271099), Research

Center for Brain Cognition and Human Development of Guangdong Province

(2024B0303390003), Striving for the First-Class, Improving Weak Links and

Highlighting Features (SIH) Key Discipline for Psychology in South China Normal

University, and National Outstanding Youth Science Fund Project of National Natural

Science Foundation of China (32022032).

Author contributions

J. L., Conceptualization, Formal analysis, Investigation, Visualization, Methodology,

Writing original draft; Y. C., Investigation, Visualization, Methodology, Writing

original draft; X. Z., Conceptualization, Resources, Supervision, Funding acquisition,

Investigation, Methodology, Writing original draft, Project administration.

Ethics

All experiments were carried out in accordance with the Declaration of Helsinki. All

participants provided written informed consent prior to the start of the experiment,

which was approved by the Research Ethics Committee at South China Normal

University (2021-04-01).

Data and Code Availability

All analyses were conducted using custom code in MATLAB and the EEGlab toolbox for EEG data analysis. The code and processed data used for the final analyses are available at https://osf.io/34cex/. Raw data for this study can be requested from the Lead Contact, and the authors confirm that all reasonable requests will be fulfilled.

Reference

Audrain, S. P., Urbain, C. M., Yuk, V., Leung, R. C., Wong, S. M., & Taylor, M. J. (2020). Frequency-specific neural synchrony in autism during memory encoding, maintenance and recognition. *Brain Commun, 2*(2), fcaa094. doi:10.1093/braincomms/fcaa094

Benedek, M., Schickel, R. J., Jauk, E., Fink, A., & Neubauer, A. C. (2014). Alpha power increases in right parietal cortex reflects focused internal attention. *Neuropsychologia, 56*(100), 393-400. doi:10.1016/j.neuropsychologia.2014.02.010

Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal statistical society: series B, 57*(1), 289-300.

Berger, B., Griesmayr, B., Minarik, T., Biel, A. L., Pinal, D., Sterr, A., & Sauseng, P. (2019). Dynamic regulation of interregional cortical communication by slow brain oscillations during working memory. *Nat Commun, 10*(1), 4242. doi:10.1038/s41467-019-12057-0

Büsel, C., Pomper, U., & Ansorge, U. (2019). Capture of attention by target-similar cues during dual-color search reflects reactive control among top-down selected attentional control settings. *Psychonomic Bulletin Review 26*, 531-537.

Chen, X., Ma, R., Zhang, W., Zeng, G. Q., Wu, Q., Yimiti, A., . . . Meng, X. (2023). Alpha oscillatory activity is causally linked to working memory retention. *PLoS Biology, 21*(2), e3001999.

Chota, S., Leto, C., van Zantwijk, L., & Van der Stigchel, S. (2022). Attention rhythmically samples multi-feature objects in working memory. *Sci Rep, 12*(1), 14703. doi:10.1038/s41598-022-18819-z

Christophel, T. B., Klink, P. C., Spitzer, B., Roelfsema, P. R., & Haynes, J. D. (2017). The Distributed Nature of Working Memory. *Trends Cogn Sci, 21*(2), 111-124. doi:10.1016/j.tics.2016.12.007

Cohen, M. X. (2014). *Analyzing neural time series data: theory and practice*: MIT press.

Constantinidis, C., & Klingberg, T. (2016). The neuroscience of working memory capacity and training. *Nat Rev Neurosci, 17*(7), 438-449. doi:10.1038/nrn.2016.43

Cowan, N. (2001). The magical number 4 in short-term memory: a reconsideration of mental

storage capacity. *Behav Brain Sci, 24*(1), 87-114; discussion 114-185. doi:10.1017/s0140525x01003922

De Vries, I. E., Van Driel, J., & Olivers, C. N. (2017). Posterior α EEG dynamics dissociate current from future goals in working memory-guided visual search. *Journal of Neuroscience37*(6), 1591-1603.

de Vries, I. E. J., Slagter, H. A., & Olivers, C. N. L. (2020). Oscillatory Control over Representational States in Working Memory. *Trends Cogn Sci, 24*(2), 150-162. doi:10.1016/j.tics.2019.11.006

de Vries, I. E. J., van Driel, J., Karacaoglu, M., & Olivers, C. N. L. (2018). Priority Switches in Visual Working Memory are Supported by Frontal Delta and Posterior Alpha Interactions. *Cereb Cortex, 28*(11), 4090-4104. doi:10.1093/cercor/bhy223

de Vries, I. E. J., van Driel, J., & Olivers, C. N. L. (2019). Decoding the status of working memory representations in preparation of visual selection. *Neuroimage, 191*, 549-559. doi:10.1016/j.neuroimage.2019.02.069

Fiebelkorn, I. C., Pinsk, M. A., & Kastner, S. (2018). A dynamic interplay within the frontoparietal network underlies rhythmic spatial attention. *Neuron, 99*(4), 842-853. e848.

Fiebelkorn, I. C., Saalmann, Y. B., & Kastner, S. (2013). Rhythmic sampling within and between objects despite sustained attention at a cued location. *Current biology, 23*(24), 2553-2558.

Foster, J. J., Bsales, E. M., Jaffe, R. J., & Awh, E. (2017). Alpha-Band Activity Reveals Spontaneous Representations of Spatial Position in Visual Working Memory. *Curr Biol, 27*(20), 3216-3223 e3216. doi:10.1016/j.cub.2017.09.031

Frătescu, M., Van Moorselaar, D., & Mathôt, S. (2019). Can you have multiple attentional templates? Large-scale replications of Van Moorselaar, Theeuwes, and Olivers (2014) and Hollingworth and Beck (2016). *Attention, Perception, Psychophysics, 81*, 2700-2709.

Fusi, S., Miller, E. K., & Rigotti, M. (2016). Why neurons mix: high dimensionality for higher cognition. *Curr Opin Neurobiol, 37*, 66-74. doi:10.1016/j.conb.2016.01.010

Gazzaley, A., & Nobre, A. C. (2012). Top-down modulation: bridging selective attention and working memory. *Trends Cogn Sci, 16*(2), 129-135. doi:10.1016/j.tics.2011.11.014

Gresch, D., Behnke, L., van Ede, F., Nobre, A. C., & Boettcher, S. E. P. (2025). Neural dynamics of reselecting visual and motor contents in working memory after external interference. *J Neurosci, 45*(18). doi:10.1523/JNEUROSCI.2347-24.2025

Gresch, D., Boettcher, S. E., Gohil, C., van Ede, F., & Nobre, A. C. (2024). Neural dynamics of shifting attention between perception and working-memory contents. *Proceedings of the National Academy of Sciences, 121*(47), e2406061121.

Han, B., & VanRullen, R. (2017). The rhythms of predictive coding? Pre-stimulus phase modulates the influence of shape perception on luminance judgments. *Sci Rep, 7*(1), 43573. doi:10.1038/srep43573

Helfrich, R. F., Fiebelkorn, I. C., Szczepanski, S. M., Lin, J. J., Parvizi, J., Knight, R. T., & Kastner, S. (2018). Neural Mechanisms of Sustained Attention Are Rhythmic. *Neuron, 99*(4), 854-865 e855. doi:10.1016/j.neuron.2018.07.032

Helfrich, R. F., Huang, M., Wilson, G., & Knight, R. T. (2017). Prefrontal cortex modulates

posterior alpha oscillations during top-down guided visual perception. *Proc Natl Acad Sci U S A, 114*(35), 9457-9462. doi:10.1073/pnas.1705965114

Ho, H. T., Burr, D. C., Alais, D., & Morrone, M. C. (2019). Auditory perceptual history is propagated through alpha oscillations. *Current biology, 29*(24), 4208-4217. e4203.

Ho, H. T., Leung, J., Burr, D. C., Alais, D., & Morrone, M. C. (2017). Auditory sensitivity and decision criteria oscillate at different frequencies separately for the two ears. *Current Biology, 27*(23), 3643-3649. e3643.

Hollingworth, A., & Beck, V. M. (2016). Memory-based attention capture when multiple items are maintained in visual working memory. *Journal of experimental psychology: human perception performance, 42*(7), 911.

Huang, Q., Jia, J., Han, Q., & Luo, H. (2018). Fast-backward replay of sequentially memorized items in humans. *Elife, 7*, e35164. doi:10.7554/eLife.35164

Jensen, O., & Mazaheri, A. (2010). Shaping functional architecture by oscillatory alpha activity: gating by inhibition. *Front Hum Neurosci, 4*, 186. doi:10.3389/fnhum.2010.00186

Jia, J., Liu, L., Fang, F., & Luo, H. (2017). Sequential sampling of visual objects during sustained attention. *PLoS Biol, 15*(6), e2001903. doi:10.1371/journal.pbio.2001903

Jiang, Y., He, S., & Zhang, J. (2024). The adaptive flexibility of rhythmic attentional sampling in attending to multiple targets. *J Exp Psychol Gen, 153*(1), 26-37. doi:10.1037/xge0001468

Josselyn, S. A., & Tonegawa, S. (2020). Memory engrams: Recalling the past and imagining the future. *Science, 367*(6473), eaaw4325. doi:10.1126/science.aaw4325

Kerzel, D., & Witzel, C. (2019). The allocation of resources in visual working memory and multiple attentional templates. *J Exp Psychol Hum Percept Perform, 45*(5), 645-658. doi:10.1037/xhp0000637

Klimesch, W. (2012). alpha-band oscillations, attention, and controlled access to stored information. *Trends Cogn Sci, 16*(12), 606-617. doi:10.1016/j.tics.2012.10.007

Kristjánsson, T., & Kristjánsson, Á. (2018). Foraging through multiple target categories reveals the flexibility of visual working memory. *Journal of experimental psychology: human perception performance, 183*, 108-115.

Kuo, B. C., Yeh, Y. Y., Chen, A. J., & D'Esposito, M. (2011). Functional connectivity during top-down modulation of visual short-term memory representations. *Neuropsychologia, 49*(6), 1589-1596. doi:10.1016/j.neuropsychologia.2010.12.043

Lachaux, J. P., Rodriguez, E., Martinerie, J., & Varela, F. J. (1999). Measuring phase synchrony in brain signals. *Hum Brain Mapp, 8*(4), 194-208. doi:10.1002/(sici)1097-0193(1999)8:4<194::aid-hbm4>3.0.co;2-c

Landau, A. N., & Fries, P. (2012). Attention samples stimuli rhythmically. *Curr Biol, 22*(11), 1000-1004. doi:10.1016/j.cub.2012.03.054

Liu, Y., Dolan, R. J., Kurth-Nelson, Z., & Behrens, T. E. J. (2019). Human Replay Spontaneously Reorganizes Experience. *Cell, 178*(3), 640-652 e614. doi:10.1016/j.cell.2019.06.012

Mansouri, F. A., Koechlin, E., Rosa, M. G., & Buckley, M. (2017). Managing competing goals—a key role for the frontopolar cortex. *Nature Reviews Neuroscience, 18*(11), 645-657.

Miller, E. K., & Cohen, J. D. (2001). An integrative theory of prefrontal cortex function. *Annu*
Rev Neurosci, 24(1), 167-202. doi:10.1146/annurev.neuro.24.1.167

Monsell, S. (2003). Task switching. *Trends Cogn Sci, 7*(3), 134-140.
doi:10.1016/s1364-6613(03)00028-7

Nakayama, R., & Motoyoshi, I. (2019). Attention Periodically Binds Visual Features As
Single Events Depending on Neural Oscillations Phase-Locked to Action. *J Neurosci,*
39(21), 4153-4161. doi:10.1523/JNEUROSCI.2494-18.2019

Nelissen, N., Stokes, M., Nobre, A. C., & Rushworth, M. F. (2013). Frontal and parietal
cortical interactions with distributed visual representations during selective attention
and action selection. *J Neurosci, 33*(42), 16443-16458.
doi:10.1523/JNEUROSCI.2625-13.2013

Oberauer, K., Farrell, S., Jarrold, C., & Lewandowsky, S. (2016). What limits working
memory capacity? *Psychological bulletin, 142*(7), 758.

Olivers, C. N., Peters, J., Houtkamp, R., & Roelfsema, P. R. (2011). Different states in visual
working memory: When it guides attention and when it does not. *Trends in cognitive*
sciences, 15(7), 327-334.

Pavlov, Y. G., & Kotchoubey, B. (2022). Oscillatory brain activity and maintenance of verbal
and visual working memory: A systematic review. *Psychophysiology, 59*(5), e13735.
doi:10.1111/psyp.13735

Peters, B., Kaiser, J., Rahm, B., & Bledowski, C. (2021). Object-based attention prioritizes
working memory contents at a theta rhythm. *journal of Experimental Psychology:*
General, 150(6), 1250.

Peylo, C., Hilla, Y., & Sauseng, P. (2021). Cause or consequence? Alpha oscillations in
visuospatial attention. *Trends Neurosci, 44*(9), 705-713.
doi:10.1016/j.tins.2021.05.004

Pomper, U., & Ansorge, U. (2021). Theta-Rhythmic Oscillation of Working Memory
Performance. *Psychol Sci, 32*(11), 1801-1810. doi:10.1177/09567976211013045

Quentin, R., King, J. R., Sallard, E., Fishman, N., Thompson, R., Buch, E. R., & Cohen, L. G.
(2019). Differential Brain Mechanisms of Selection and Maintenance of Information
during Working Memory. *J Neurosci, 39*(19), 3728-3740.
doi:10.1523/JNEUROSCI.2764-18.2019

Re, D., Inbar, M., Richter, C. G., & Landau, A. N. (2019). Feature-Based Attention Samples
Stimuli Rhythmically. *Curr Biol, 29*(4), 693-699 e694. doi:10.1016/j.cub.2019.01.010

Soto, D., Humphreys, G. W., & Rotshtein, P. (2007). Dissociating the neural mechanisms of
memory-based guidance of visual selection. *Proc Natl Acad Sci U S A, 104*(43),
17186-17191. doi:10.1073/pnas.0703706104

Sreenivasan, K. K., Curtis, C. E., & D'Esposito, M. (2014). Revisiting the role of persistent
neural activity during working memory. *Trends Cogn Sci, 18*(2), 82-89.
doi:10.1016/j.tics.2013.12.001

Stam, C. J., Nolte, G., & Daffertshofer, A. (2007). Phase lag index: assessment of functional
connectivity from multi channel EEG and MEG with diminished bias from common
sources. *Hum Brain Mapp, 28*(11), 1178-1193. doi:10.1002/hbm.20346

Van Diepen, R. M., Foxe, J. J., & Mazaheri, A. (2019). The functional role of alpha-band
activity in attentional processing: the current zeitgeist and future outlook. *Curr Opin*

Psychol, 29, 229-238. doi:10.1016/j.copsyc.2019.03.015

van Driel, J., Gunseli, E., Meeter, M., & Olivers, C. N. (2017). Local and interregional alpha
EEG dynamics dissociate between memory for search and memory for recognition.
Neuroimage, 149, 114-128.

Van Ede, F., & Nobre, A. C. (2023). Turning attention inside out: How working memory
serves behavior. *Annual review of psychology, 74*(1), 137-165.

van Moorselaar, D., Theeuwes, J., & Olivers, C. N. (2014). In competition for the attentional
template: Can multiple items within visual working memory guide attention? *Journal*
of Experimental Psychology: Human Perception Performance, 40(4), 1450.

VanRullen, R. (2016). How to evaluate phase differences between trial groups in ongoing
electrophysiological signals. *Frontiers in neuroscience, 10*, 426.

VanRullen, R., & Macdonald, J. S. (2012). Perceptual echoes at 10 Hz in the human brain.
Current biology, 22(11), 995-999.

Wallis, G., Stokes, M., Cousijn, H., Woolrich, M., & Nobre, A. C. (2015). Frontoparietal and
Cingulo-opercular Networks Play Dissociable Roles in Control of Working Memory.
J Cogn Neurosci, 27(10), 2019-2034. doi:10.1162/jocn_a_00838

Wolff, M. J., Jochim, J., Akyurek, E. G., & Stokes, M. G. (2017). Dynamic hidden states
underlying working-memory-guided behavior. *Nat Neurosci, 20*(6), 864-871.
doi:10.1038/nn.4546

852