Strong negative reinforcement interferes with visual learning in a solitary pollinator

Anupama Nayak Manel[1,2], James Jonathan Foster[1,3], Anna Stöckl[1,4]

[1] Department of Biology, University of Konstanz, Konstanz, Germany

[2] International Max Planck Research School for Quantitative Behaviour, Ecology and Evolution (IMPRS-QBEE), Max Planck Institute of Animal Behaviour, Radolfzell, Germany

[3] Centre for the Advanced Study of Collective Behaviour, University of Konstanz, Konstanz, Germany

[4] Zukunftskolleg, University of Konstanz, Konstanz, Germany

ABSTRACT

Visual learning in insects can be strongly influenced by the underlying context and conditions. Improved colour learning has been demonstrated with differential conditioning using reward-aversion paradigms in eusocial insects like wasps, bumblebees, and honeybees, where substances that induce strong aversion, such as quinine, enable more accurate learning. Although the associations created during learning are intrinsically linked to the reinforcements used, few studies have compared the role of negative reinforcements in shaping these associations, and their relevance across insect groups with different life histories.

In our study, we compared the effects of aversive substances for learning in the hummingbird hawkmoth (*Macroglossum stellatarum)*, a solitary pollinator that relies on vision for foraging. Linking to previous learning studies in insects, we combined a sugar-rewarded target with a distractor that was paired with either quinine, salt, citric acid, water, or was presented with no aversive substance. Learning was assessed by conditioning hawkmoths to invert a strong colour preference between perceptually close or distant colour pairs, to provide tasks with disparate challenges. Contrary to results from eusocial insect species, hawkmoths trained with quinine were worse at switching preferences between similar colours compared to training with citric acid or appetitive-only differential conditioning. Learning success was associated with the animals' foraging plasticity, where negative reinforcements were found to

32   suppress exploration during foraging. Furthermore, we show that quinine interfered with

33   sucrose perception, potentially impairing target acquisition during conditioning. Our results

34   provide insights into the impact of negative reinforcements on solitary foragers and highlight

35   the role of sensory ecology and life history in shaping learning outcomes.

INTRODUCTION

Visual learning in insects can be highly plastic, with behavioural outcomes varying in response to experiential details such as learning durations (Stach & Giurfa, 2005), or the context in which learning takes place (Avarguès-Weber & Giurfa, 2014; Buatois et al., 2017; Dyer & Chittka, 2004; Dyer & Howard, 2023; Giurfa, 2004; Howard et al., 2019; Kelber, 2010a; Matsumoto & Mizunami, 2002). In particular, the conditions in which stimuli are presented are known to alter the speed and accuracy of learning depending on whether the animal experiences absolute conditioning to a single target, or differential conditioning in the presence of a rewarded target stimulus along with a distractor stimulus associated with a neutral or negative feedback (Avarguès-Weber & Giurfa, 2014). Differential conditioning enables narrower colour discrimination and pattern recognition where absolute conditioning is less effective (Dyer & Chittka, 2004; Giurfa, 2004). Such improved learning performance has been attributed to the reward and punishment dynamic of differential conditioning, which could motivate animals to be more attentive to the task due to the adverse consequences of incorrect choices (Avarguès-Weber & Giurfa, 2014). Moreover, differential learning provides a richer set of information to enable selective association, since both, the positively associated target and negatively associated distractor stimuli can be learnt and distinguished simultaneously.

Since the strength of association with the target and distractor is a key aspect of differential learning, the reinforcement type paired with the distractor plays a critical role in the conditioning process. Across insect groups, various substances, such as water, salt solution, or bitter alkaloids (Hymenoptera: Josens et al., 2009; Lacombrade et al., 2023; Orthoptera: Matsumoto, 2022; Lepidoptera: Rodrigues, 2016; Young et al., 2024) have been used as negative reinforcements. Alternatively, appetitive-only conditioning, where the target is rewarded while the distractor is not associated with any substance, has also been used for insect training (Goyret et al., 2008; Kelber, 2002). Previously, in eusocial Hymenoptera, colour discrimination between perceptually similar– and thus harder to learn– stimuli was shown to improve by using the aversive alkaloid quinine as a negative reinforcer, compared to using water (Avarguès-Weber et al., 2010; Rodríguez-Gironés et al., 2013), or with no negative reinforcement (Dyer & Howard, 2023). However, even within Hymenoptera, the use of quinine as a negative reinforcement remains contested since aside from being toxic to the insects (Ayestaran et al., 2010), its efficacy is also context-dependent: although it improved learning in untethered honeybees, it had no effect on learning in harnessed bees, which sometimes even ingested the toxic bitter compound (Ayestaran et al., 2010; de Brito Sanchez et al., 2015).

The wide disparity of learning conditions, in addition to the limited evidence for assessing their suitability across insect models, makes it a challenge to interpret learning outcomes across species. To understand which learning strategies are shared across insects, and which are specific adaptations to a species' lifestyle and environment, it becomes crucial to understand how reinforcing substances might present disparate qualities for species with different sensory systems and life histories. For instance, phytophagous insects specialised on certain host plants may have different tolerances for specific secondary compounds produced by the plant. The fly, *Drosophila sechellia*, a specialist adapted to the toxic fruits of *Morinda citrifolia*, shows reduced aversion to the bitter compounds rejected by other Drosophilidae (Reisenman et al., 2023). Similarly, frugivorous butterflies were shown to be more receptive to amino acids in their diets than nectarivorous butterflies (Ravenscraft & Boggs, 2016). Such differences, even between closely related species, question whether there is a ubiquitous role of appetitive and aversive substances in insect learning.

To assess how different negative reinforcements influence visual discrimination learning in a solitary insect, we differentially conditioned the day-active hummingbird hawkmoth, *Macroglossum stellatarum (Fig. 1)*. *M. stellatarum* are generalist nectar feeders that rely on vision for locating and interacting with flowers (Balkenius, Rosén, et al., 2006; Kelber, 1997; A. Stöckl et al., 2016). They represent a large group of solitary insects that occupy a key pollinator niche (Johnson et al., 2017; Singh et al., 2025), sharing many habitats and floral hosts with eusocial pollinators. Both behavioural and neural correlates of visual foraging in this species have been extensively studied (Balkenius, Kelber, et al., 2006; Kannegieser et al., 2024; Kelber, 2002; A. Stöckl et al., 2016; A. L. Stöckl et al., 2017). Previously, they have been found to make rapid visual associations via absolute conditioning. However, their learning outcomes were limited by their strong preferences for certain pattern and colour features (Kelber, 2002). To test whether conditioning with negative reinforcements could improve their visual association learning, we differentially conditioned the hawkmoths using substances in the taste categories -sour, salt, bitter, and water (neutral) – as negative reinforcements. These negative reinforcements were compared with an appetitive-only condition. Hawkmoths were conditioned to invert a strong colour preference in two visual discrimination tasks: one with perceptually close and another with perceptually distant colours. With this preference inversion paradigm, we addressed whether the type of negative reinforcement altered learning outcomes, and how the effects of reinforcement varied with the challenge of the visual discrimination task.

Fig. 1 **Comparing negative reinforcement types for colour discrimination learning in**

the diurnal hawkmoth, *Macroglossum stellatarum*. a. Learning was compared across

differential conditioning protocols with various treatment types (-) using aversive substances

common in insect training, as well as an appetitive-only condition with no aversive

substance. Asterisks indicate treatments used only in the distant colour discrimination task.

b. Colour preference inversion was compared across two discrimination tasks with stimuli

that were perceptually distant (blue-yellow) and close (blue-cyan). Relative reflectance of the

colours used in the experiment (see Methods for details on measurements, and Fig. S1 for

distances modelled in the hawkmoths' colour space). **c.** The experimental paradigm was

divided into 3 phases: moths were pretrained to artificial flowers of a single colour in groups

and then individually tested for a preferred colour in a dual choice assay. Following this, their

preferred colour was paired with a single treatment type (-) as a negative reinforcer and the

unpreferred colour was rewarded with sugar (+). The figure depicts the distant colour

discrimination task as an example, with yellow as the moth's preferred colour. Durations for

each phase are depicted as white boxes representing a single day.

Materials and Methods

Animals

Hummingbird hawkmoth (*Macroglossum stellatarum*) populations were maintained indoors at the University of Konstanz, Germany at 23 °C and 40-60% humidity with a 14:10 h light:dark cycle. Populations were supplemented with wild-caught moths when available. Caterpillars were raised on their native host plant *Gallium* sp.. Adults were fed in flight cages (60 cm × 60 cm × 60 cm) with 20% sucrose-water solution from gravity feeders. Multiple feeders in blue and yellow colours were provided simultaneously to discourage colour associations.

Animals used in the experiment were 3-6 weeks of age. Moths were placed in plastic vials (8 cm x 3.5 cm length x diameter) and kept in the dark when not participating in the experiments. The vials were marked to track individual identity across the experiment.

Experimental Setup

Experiments were performed in a flight cage of 60 cm × 60 cm × 60 cm dimension at 23 °C. The cage was lit from above with four 60 cm long daylight-like fluorescent tubes (Osram L 18 W/965 Biolux Tageslicht G13). Lights were connected to an electrical ballast (GloMat 2 × 40 W, Hagen), to increase the flicker frequency of the fluorescent tubes above 25 kHz, beyond the resolvable range of hawkmoths (Stöckl et al., 2019). A USB camera fixed to the upper corner of the cage was used to record the experiments. It was controlled by the ContaCam software (ContaCam v9.0.9beta4, https://www.contaware.com) to record videos at 30 fps.

Artificial flowers were constructed by pasting coloured discs (38 mm diameter) on cylindrical plastic pedestals (8 cm x 3.5 cm) wrapped with unbleached paper. Discs in Blue (CMYK 91:66:0:0), Yellow (CMYK 0:0:100:0), and Cyan (CMYK 91:26:0:0) were designed in the vector graphics software Inkscape (Inkscape v1.3 2023, https://inkscape.org). The colours were laser printed (imageRUNNER ADVANCE DX C3926i, Canon) on unbleached paper ('Classic White', Steinbeis) and laminated with matte foil (S-PP525-22 matt, Peach, PRT GmbH). To provide optic flow during flight, matte-laminated dead-leaves patterns (Lee et al., 2001) were used on the floor of the cage. During the test and conditioning phase of the experiments, the flowers were placed on a dead-leaves pattern covered cardboard platform (30 cm × 50 cm × 50 cm) higher off the ground of the cage to decrease the flight height and thereby get increased feeding responses.

Irradiance measurements of the flower colours, background, and lighting were taken using a
calibrated Miniature Spectrometer (OceanHDX, Ocean Optics Inc., Dunedin, FL, USA)
connected to a UV-VIS optical fibre (P400-2-UV-Vis). Relative reflectance of the stimuli was
then calculated with respect to a 99% white standard (Spectralon diffuse reflectance
standards, Labspehere Inc., USA). All spectral measurements were processed and
visualized in RStudio (Rstudio Team, v2023.09.1) using the pavo 2.2.0 package (Maia et al.
2019) (*Fig. S1a*).

To map the colour distances, quantum catches for the photoreceptors were calculated by
integrating the sensitivity peaks of *M. stellatarum* at 349, 440, and 521 nm (Telles et al.,
2014), the cage illumination, and irradiance measures of the stimuli (Renoult et al., 2017).
The 'sensmodel' function from the pavo package was used to set the sensitivity values for
the hawkmoths, which were then input into the 'vismodel' function to calculate photoreceptor
stimulation based on von-Kries colour corrections (Maia et al., 2019). Relative chromatic and
achromatic distance between the colours were calculated as noise-weighted Euclidean
distances measured in Just Noticeable Differences (JNDs) (Table S1, *Fig. S1b*), which is a
contrast threshold that uses the RNL model (Vorobyev & Osorio, 1998). Relative
photoreceptor densities were assigned as 1:1:7 for the small, medium, and long wavelength
receptors (Kelber et al., 2003).

173 Experimental Procedure

Hawkmoths were differentially conditioned to invert a strong colour preference using various
treatment types as the negative reinforcement. By using quinine hydrochloride (120 mM),
citric acid (1.04 M), and sodium chloride (3 M) solution, distilled water, and no substance
(appetitive-only condition), we combined treatments from insect literature (Avarguès-Weber
et al., 2010; Kelber, 2002) to compare a variety of negative reinforcements. We assessed
learning in 2 visual discrimination tasks associated with disparate learning challenges (Dyer
& Chittka, 2004; Giurfa, 2004). For the easier task we used blue and yellow which are further
apart in hawkmoths' colour space (distant colours), and for the harder discrimination task we
used blue and cyan (close colours) (*Fig. 1b, see Fig. S1b for colour spaces*) as the
conditioned stimuli (CS). Individual hawkmoths experienced a single treatment and
discrimination type with the full paradigm divided into 3 phases – Pretraining, Preference
test, and Preference inversion *(Fig. 1c).*

In all but the pretraining phase, animals were tested individually in the flight cages. Food
(20% sucrose solution) was only available during the trials and for the remainder of time,
animals were kept in vials under dark conditions. Artificial flowers within the flight cage were

randomly arranged at least 5 cm away from each other, and their positions were changed
within and between trials to avoid place learning. The negative reinforcement substances and
sugar rewards were pipetted out at several points on the surface of the flower as small
droplets (20-30 µl), requiring the moths to make multiple choices before satiation. The flower
surfaces were cleaned between all trials. A trial commenced when the moth began its flight,
and during the trial every new approach to an artificial flower that ended with a proboscis
contact to the surface was scored as a choice. Repeated contacts on the same surface while
hovering or repositioning around the flower were not included as additional choices. All trials
were video documented.

Pretraining

To habituate the moths to the artificial flowers, groups of 5-8 moths were pretrained to the
setup by sequentially switching their regular gravity feeders with artificial flowers across 3
days. On the first day the moths had one feeder and 2 artificial flowers at different elevations,
and by the third day only 3 artificial flowers were available to feed from. Each group was
pretrained to flowers of a single randomly chosen colour. Either yellow or blue was used for
the distant colour experiments, and either blue or cyan were used for the close colour
experiments. During this phase, the moths could feed ad libitum on 20% sucrose from the
surface of the flowers for 10-12 hours each day. To ensure the flowers always contained
sucrose during visits, they were removed from the cage in the evenings. On the evening of
the third day, moths were packed into individual vials and kept in a dark box for 48 hours to
motivate feeding during the next phase.

Preference test

Moths were tested for a preferred feeding colour between the two colours assigned to their
discrimination type, blue and yellow (distant colours), or blue and cyan (close colours). The
identity and sex of the moth were noted, along with the time of start of each experiment. Six
artificial flowers, 3 of each colour, were arranged in the flight cage with 20% sucrose drops
available on all the flowers. All choices were rewarded to prevent the animals from forming
negative associations with their preferred colour, so the animals could feed during the trial
rather than in a non-experimental setting. Flower choices were scored by the experimenter
across 5 minutes of flight. Only animals with a relative colour preference, i.e. at least 60% of
choices on one of the colours, were used in the following preference inversion phase.

Preference inversion

Hawkmoths were differentially conditioned to their unpreferred colour as determined by the
preference test. For the distant colour experiments, the artificial flowers with the unpreferred
colour (CS+) were paired with a reward of 20% sucrose while the preferred colour (CS-) was
associated either with citric acid (n=18), quinine hydrochloride solution (n=16), sodium

chloride solution (n=18), distilled water (n=14), or, in the case of the appetitive-only
conditioning, no substance (n=14).

For the close colour (blue-cyan) discrimination condition, only citric acid (n=13), quinine
(n=15), and no substance (n=14) were used, as these treatment types emerged as most
relevant from the distant colour experiments.

The moths were differentially conditioned in 4 trials, one per day, during which they had
access to the artificial flowers (3 x CS+ and 3 x CS-) for 8 minutes of flight. This duration was
determined to be a suitable interval for most moths to approach the flowers and make
sufficient choices until satiation based on pilot experiments. Only sessions in which moths
made at least one choice were included as trial days.

Control experiment

For the appetitive-only conditioning, only the CS+ was presented with droplets. To assess
whether the animals associated the droplets rather than the colour as a rewarding cue, we
altered the preference inversion phase with a test on the fourth day where all the flower
surfaces were bare *(Fig. S2a)*. For this test, choices were recorded for 5 minutes of flight,
rather than 8 minutes, to account for changes in preference associated with finding no
rewards on the CS+. This control was conducted for the appetitive-only condition (n=9) with
the close colour setup where the results showed a significant relation between learning and
treatment type.

Data Analysis

Data was analysed separately for the distant colour and close colour discrimination
experiments. Moths that participated for at least 3 days of differential conditioning were
included in the analysis. All statistical analyses were conducted in R (R Core Team, 2021),
and plots were created with the ggplot2 package (Wickham, 2011).

To compare the number of individuals that succeeded in the preference inversion task
between treatments, we assigned a success criterion to moths that shifted their colour
preference in relation to the 60% preference criterion in the preliminary test, i.e., individuals
with greater than 40% CS+ choices (the initially non-preferred colour) on the fourth day. A
pairwise Fisher's exact test with Holm corrections for multiple testing was then implemented
to compare the numbers of successful and unsuccessful moths between treatments *(Fig.*
S3b).

Choice rates

Generalized linear-mixed models (GLMM) fitted with the 'glmer' function from the lme4
package were used to test if the correct choice rate varied significantly between treatment
types, and across the differential conditioning trials. Choice data from the differential
conditioning was fitted to a binomial family with treatment type, day, preferred colour, sex,
and the choice rate from the preference test as fixed effects, and individual ID as the random
effect. Residuals from the model were simulated using the 'simulateResiduals' function from
the DHARMa package (Hartig, 2018) to validate model accuracy. Stepwise reduction of the
full model was conducted with likelihood ratio tests (LRT) via the 'drop1' (Crawley, 2002)
function, and the random effect structure of the final model was selected based on Akaike
information criterion (AIC) comparisons. The simplest adequate model was formulated as

$$(correct\ choices, incorrect\ choices) \sim Day * Treatment * Preferred\ colour$$
$$+ Pref. test\ choice\ rate + (1|Animal\ ID)$$

Preferred colour as a predictor was removed from the close colour experiment model where
only blue was the CS-. The 'Anova' function from the car package (Fox and Weisberg, 2019)
with Wald's Chi-square test was used to determine the significance of the fixed effect
variables on the choices. Post-hoc comparisons were then conducted to identify pairwise
differences using a Sidak (Šidák, 1967) correction for multiple comparisons with the
emmeans package (Lenth et al. 2019) *(Fig. 2,3)*.

Data from the control experiment between close colour stimuli was similarly fit to a binomial
GLMM and post-hoc comparisons were performed to estimate choice rate differences
between days *(Fig. S2)*. Correlation between the choices on the third and fourth day was
calculated with the 'cor.test' function using Spearman's method to assess whether moths
made comparable choices in the test with no droplets.

Learning curves

To model learning curves, we used a GLMM incorporating a psychometric function (see
Kirwan and Nilsson, 2019) using Bayesian estimation. The learning curves were modelled on
data from the successful individuals (>40% correct choices on day 4), to visualize how the
treatment type affected learning dynamics *(Fig. 4)*. The model was fitted with the Stan
language (Carpenter et al., 2017; http://mc-stan.org/) via the brms package (v2.6.0; Bürkner,
2018; https://cran.r-project.org/web/packages/brms/) within R.

A logistic regression model modified from Olsson et al. (Olsson et al. 2018) was used, where
the initial guess rate and lapse in learning were assigned as the baseline and upper
asymptote of the curve respectively. This psychometric formula was used to estimate the
learning curve's inflection point (Inflex) and 80% of the curve's rise range (Width) to compare

between treatment types. To restrict the estimates of the baseline and lapse rate between
zero and one, they were estimated on a logit scale {y=ln[x/(1−x)]}, and the curve width was
estimated on the natural logarithmic scale. Choices from each day were set as the response
variable and fit to a binomial family with an identity link. Informative priors for the baseline
were estimated close to 0 based on the animals' unambiguous choices during the preliminary
test. The learning lapse was set near 0 for the distant colour experiments and around 0.25 for
the close colour experiments where fewer animals reached a 100% choice rate. Strong priors
were also set for the threshold and width of the curve with a distribution centred on the first
conditioning day for the distant colour experiments and the third day for the close colour
experiments, based on visualising the choice data. Preferred colour, treatment type, choice
rate from the preference test, and their interactions were included as fixed effects, and
individual ID was set as the random effect to obtain threshold-width values specific to each
condition and individual.

A model of best fit was assessed by dropping predictor variables from the full model and
comparing all possible models using leave-one-out cross validation via the 'LOO' function
(Vehtari et al., 2017). The model of best fit was chosen as the one with the highest ELPD
(Estimated Log Predictive Density) value. The estimates were calculated from the final model
with predictors assigned as

$$Baseline \sim logit\,(Pref.test\;choice\;rate) \; + \; (1|Animal\;ID)$$

$$Lapse \sim 1 \; + \; (1|Animal\;ID)$$

$$Inflex \sim Treatment \; * \; Preferred\;color \; + \; (1|Animal\;ID)$$

$$Width \sim Treatment \; * \; Preferred\;color \; + \; (1|Animal\;ID)$$

308 Preferred colour was excluded from the close colour experiments with only blue as the CS-,
309 and based on the data from this discrimination task, the predictors for lapse were adjusted to

$$Lapse \sim Treatment \; + \; (1|Animal\;ID)$$

Parameters were estimated using four Markov–Chain Monte Carlo (MCMC) chains running
for 2000 iterations post-warmup, and model convergence was checked and adjusted by
replacing the default priors with more informative priors when necessary (*Table S2*). The final
estimates were drawn from models where all chains converged (R-hat convergence
diagnostic < 1.01). Learning curves predicted for each individual were also used to visually
validate predictions.

Colour switching rates

As a proxy for learning dynamics during the differential conditioning, switching rates were
calculated for each individual. Successive choices of the same colour were marked a '0'
switch, while choosing different colours in successive flower contacts was marked a '1'
switch. The switching rates were then calculated as the sum of switches divided by the
number of possible switches across the total choices made by an individual during the
differential conditioning period. To examine the variation of switching rates with
discrimination and treatment type, a GLM including the treatment types common to both
discrimination tasks was modelled using the 'glmmTMB' function (Brooks et al., 2017). The
model was fit to a betabinomial family with the interaction of treatment and discrimination
type, and the number of feeding counts as predictors. After stepwise model reduction, the
simplest adequate formula was structured as,

$(Total\ switch\ counts, (Total\ choices - 1) - Total\ switch\ counts) \sim Treatment\ type\ *$
$Discrimination\ type$

The 'Anova' function with Wald's Chi-square test was used to determine the significance of
the fixed variables on the switching rates. Post-hoc comparisons were then conducted to
identify pairwise differences using Sidak adjustments for multiple-testing corrections *(Fig.*
5b).

To test whether switching between colours was linked to the animals' correct choice rates, we
fitted correlations between the switching index and the choices rates on the fourth day using
a Spearman's tests with the 'cor.test' function *(Fig. 5c, d).*

Proboscis contact duration

To quantify the duration of proboscis contacts with the treatment substances – citric acid,
quinine, water, salt, and the sucrose solution, we annotated videos from the preference
inversion phase using BORIS software (v8.22.6, https://www.boris.unito.it/) (Friard & Gamba,
2016). Videos from the first day, during which animals selected both the CS+ and CS-, were
collected and scored in a blind assessment where treatment type and the rewarded colour
were undisclosed. Contact durations were recorded as time between the first to last moment
of contact with a droplet. Only videos where the proboscis contacts were clearly observed
were used. To minimise the effects of satiation or habituation on durations, not more than 10
contacts were scored. Only animals with at least 5 contacts with both the CS+ and CS- were
included in the final analysis.

A generalised linear mixed model (GLMM) was fitted to a gamma family with a log link with
the contact durations as the dependent variable and stimulus type (CS+ or CS-), treatment

type, discrimination type, and preferred colour as the predictors, and animal ID as the
random effect. Estimates were drawn from the simplest adequate model,

$$Contact\_Duration \sim Stimulus\ type * Treatment + (1|Animal\ ID)$$

352 Pairwise comparisons for the treatment types were estimated with Sidak corrections for
353 multiple testing.

Results

To compare the impact of various negative reinforcements on learning outcome, hawkmoths
were trained to invert a colour preference in a discrimination task with either stimuli of
perceptually distant colours (blue-yellow) or close colours (blue-cyan). Accordingly,
comparisons between the different treatment types, i.e., appetitive-only conditioning, or citric
acid, quinine, salt, or water as negative reinforcements were interpreted within their
respective discrimination tasks.

Since in all but the appetitive-only conditioning, both stimuli had droplets on the surface, we
performed a control with the no substance treatment to ensure that the droplets did not aid
associations to the CS+. As there were no significant differences in choice rates between the
third day of differential conditioning with only sugar rewards and the subsequent test without
droplets on either stimulus (*Fig. S2b*), we concluded that the moths did not rely on the
presence of droplets for their choices. A moderate positive correlation (Spearman's $\rho=0.61$,
$p=0.08$; *Fig. S2c*) between the choices from the third day and the test supports the
conclusion that the animals maintained their associations across days despite the absence of
droplets on the flowers. Thus, we conclude that all choices made in these experiments were
colour driven.

Negative reinforcement type did not alter visual discrimination learning of distant colours

Fig. 2 **Preference inversion in the distant colour experiments.** Correct choice rates represent feeding on the preferred colour during the preference test (grey highlight), and on the unpreferred colour (CS+) during the preference inversion phase. Choice rates of individuals are denoted by circles with the size representing the number of choices made in a day (see legend). Statistical results (GLMM, Fig. S4a) of differences in choices rates between days within a treatment type are indicated on the treatment-specific coloured bars where days with no differences *(p>0.05)* share the same letter. Pairwise comparisons of treatments within a day were not significant.

In the distant colour experiments, moths pretrained to blue or yellow artificial flowers selected their pretrained colour *(Fig. S3a)* with a high choice rate during the preference test (*Fig. 2 Preference test*).

In the preference inversion phase, moths significantly increased their correct choice rate across the differential conditioning period (GLMM Day - X^2= 570.94, df= 3, p< 0.001; *Fig. 2*). A success criterion was assigned to animals with a 40% baseline of correct choices on day 4, indicating a shift from their initial colour preference of 60%. This criterion was met by 71.05% (54/76) of the conditioned animals with a high median choice rate of 100% *(Fig. 2 Day 4)*. The treatment type did not significantly influence the number of conditioned animals that achieved the success criteria (*Fig. S3c, Table S3*).

The moths' preferred colour influenced their choice rate (GLMM Day: Preferred colour -
X^2= 11.62, df= 3, p= 0.009; *Fig. S4b*), with animals conditioned to yellow (CS+) against their
blue preference having a greater correct choice rate on the second and third day. This
difference disappeared on the last day. Generally, moths with a higher choice rate for their
preferred colour during the preference test performed more poorly on the first day of
differential conditioning in the inversion phase (Spearman's test rho= -0.45, p< 0.002; *Fig.*
S4d). However, this correlation was not significant by the last day of the conditioning,
showing that their initial colour preference did not impact the final learning outcome. We
found no interaction between the treatment type and the moths' preferred colour.

There was also a significant difference in the choices rates across days between treatment
types (Day: Treatment - X^2= 40.5465, df= 12, p< 0.001; *Fig. 2*) but within a day there were no
significant differences between the treatments. Despite this, we found a general trend of citric
acid and appetitive-only conditioned individuals more consistently achieving higher correct
choice rates. Quinine appeared to motivate learning more slowly while water and salt shared
similar trends with highly variable success rates.

In conclusion, with perceptually distant colours, the negative reinforcement type did not
distinctly alter learning success on individual days, as the hawkmoths readily inverted their
preferences in all conditions.

414 Negative reinforcement type altered visual discrimination
415 learning for close colours

To pose a more challenging learning context, in which the effect of different aversive
treatments might be observed more clearly, we conducted the experiment using stimuli closer
in the hawkmoths' colour space (*Fig. S1b*), with treatments that emerged as most effective
from the distant colour experiments - citric acid, quinine, and the appetitive-only conditioning.

Fig. 3 **Preference inversion in the close colour experiments.** *Correct choice rates*
represent feeding on the preferred colour during the preference test (grey highlight), and on
the unpreferred colour (CS+) during the preference inversion phase. Choice rates of
individuals are denoted by circles with the size representing the number of choices made in a
day (see legend). Statistical results (GLMM, Fig. S4c) of differences in choices rates between
days within a treatment type are indicated on the treatment-specific coloured bars where
days with no differences (p>0.05) share the same letter. Significant differences between
treatment types within a day are shown *with black asterisks* (*** p < 0.001, **< 0.01, *< 0.05, .
< 0.1).

In the analysis of the close colour experiments, we only included cyan as the CS+, since
moths pretrained to both cyan or blue, selected blue flowers during their preference test (*Fig.*
S3a). In this discrimination task, fewer moths shifted their colour preference in comparison
to the distant colour experiments, with only 52% (22/42) of animals reaching the success
criterion. Although the moths modified their foraging choices across the duration of the
differential conditioning (GLMM Day - X^2= 47.51, df= 3, p< 0.001; *Fig. 3*), their median choice
rate for the CS+ (cyan) by the end of the differential conditioning reached only 51.04% (*Fig. 3*
Day 4), distinctly lower than their counterparts in the distant colour experiments. While
successfully conditioned animals in the distant colour experiments decisively chose the CS+,

most moths in the close colour experiments did not make as strong an association with the
rewarded target.

Interestingly, the treatment type influenced the proportion of animals that reached the
success criterion, which was significantly lower when conditioned with quinine in comparison
to the appetitive-only condition (Fisher's exact test: quinine/none- p= 0.01; *Fig. S3b; Table*
S3). The treatment type also influenced the variation in choice rates across the differential
conditioning period (GLMM Day: Treatment - X^2= 14.862, df= 6, p= 0.021; *Fig. 3*). In contrast
to the distant colour experiments, there were significant differences in choice rates between
treatment types within each day. Groups with no substance or citric acid on the CS- had a
significantly higher correct choice rate on each day than the group treated with quinine (*Fig.*
3). There were no significant differences between the no substance and citric acid groups
within days. Since animals in the close colour experiments were trained only with cyan as the
CS+, the effect of preferred colour on choice rates could not be assessed.

Thus, with perceptually close colours, where hawkmoths did not learn to invert their colour
preference to the same degree as with distant colours, there were distinct effects of the
negative reinforcement type on learning: the highest learning success was achieved with the
appetitive-only and citric acid treatment, and the lowest with quinine.

Learning and foraging dynamics during differential conditioning

To investigate the impact of the treatment types on learning dynamics, psychometric learning curves were modelled to the choice data from the successful animals.

Fig. 4 **Learning dynamics across treatment types for each discrimination task. a-b**. Psychometric learning curves modelled for data from successful individuals in the preference inversion phase (see Methods for details). Points depict the correct choice rates of individuals, and colours represent the respective treatment types. Curves are extrapolated to the 0th day to visualize learning predictions for the baseline (initial choice rate). The dotted line depicts the 40% choice criterion which was considered successful learning. **c.** Inflection points of the learning curves estimated for the treatment types for each discrimination task model (close colour and distant colour). Upper and lower box limits depict the Standard error, and the whiskers depict the 95% Credible Intervals. **d.** Widths of the learning curves estimated for the treatment types for each discrimination type model. Blue-yellow and blue-cyan circle pairs depict the discrimination task corresponding to the data.

The learning curves modelling the distant colour experiments had steep sigmoidal shapes plateauing at a 100% choice rate within the second day of the differential conditioning (*Fig. 4a*). The learning curves of the close colour experiments were flatter and reached their plateau later in the conditioning period at around a 75% choice rate (*Fig. 4b*). Across treatment types, there was a similar trend in the blue-yellow and blue-cyan experiments for the time taken to arrive at the inflection point, the halfway mark between the lower and upper

asymptote of the curves. For both discrimination types, colour choices shifted early in the
preference inversion phase (*Fig. 4c*), with the inflection point shifted relatively later for
quinine, indicating longer conditioning times, while appetitive-only conditioning required the
shortest time. This was the case in both colour tasks, but more pronounced with close
colours. The widths, which represent the time range during which 80% of changes in the
responses occur, were narrow in the distant colour experiments, as represented by the steep
learning slopes, and much wider for the close colour experiments where target acquisition
occurred at a slower pace (*Fig. 4d*). There were no differences in widths between the
treatment types within a discrimination task, suggesting that the treatment type did not
significantly change the speed of target acquisition but rather, influenced the time taken by
the animals to sample the CS+.

Fig. 5 **Exploration dynamics during differential conditioning. a.** Example choice sequence of individual moths in the appetitive-only conditioning from the distant colour (left) and close colour experiment (right). **b.** Switching rates of individuals calculated for choices across the colour inversion phase. A value of 1 represents colour switching with every choice, a value of 0 represents no switching during the entire inversion training. Estimates with 95% confidence intervals (in black) are drawn from the GLMM (see Methods). Switching rate of individual moths are represented as points with the size of the circle depicting the number of choices made by the individual in the colour inversion phase (see legend). *Significant differences* are shown with black asterisks (*** p < 0.001, **< 0.01, *< 0.05). **c-d.** Correlation of switching rates with the choice rate on day 4 of preference inversion. Points represent individual moths, and colours indicate their treatment type. Results of a Spearman's rank correlation coefficients (rho) are shown in the graph. Blue-yellow and blue-cyan colour pairs depict the discrimination task corresponding to the data.

Successful animals in the distant colour experiments rapidly and reliably switched to feeding on the CS+ after a few rewarded encounters. However, in the close colour experiments, most successful individuals distributed their feeding choices between both CS+ and CS- rather than only visiting the rewarded target (*Fig. 5a*). To quantify the likelihood of switching between a preferred and unpreferred colour across both colour discrimination tasks, we calculated the animals' switching rates, where a value of 1 represented animals that switched colour with every choice, and a value of 0 represented those that exclusively chose one colour during the entire inversion training phase. These measures differed significantly between the treatment types depending on the discrimination tasks within which the data

was nested (GLM Treatment type: Discrimination type - X^2= 8.93, df= 2, p= 0.01). The
measures indicate a stronger colour fidelity in the distant colour experiment in comparison to
the close colour experiments (GLM Discrimination type - X^2= 42.09, df= 1, p< 0.001).
Switching rates varied significantly between the treatment types only in the close colour
experiments, with appetitive-only conditioning promoting the greatest number of colour
switches and quinine the lowest *(Fig. 5b).* We correlated switching rates to task success to
assess whether a strong fidelity to their preferred colour could prevent the hawkmoths from
sampling the CS+ and forming reward associations. A positive correlation was found between
the choice rate on the fourth day of preference inversion and the animal's switching rate in
the close colour experiments (Spearman's test rho= -0.48, p= 0.002; *Fig. 5d*), but there was
no link between switching and success for the distant colour experiments (*Fig. 5c*).
Interestingly, we found no correlation between the number of choices made during the
conditioning period and the success rates, indicating that their switching rates rather than the
number of choices predicted success.

Thus, in addition to learning success, the learning dynamics, both in terms of how fast
hawkmoths learned, and how they explored the stimuli, were impacted by the discrimination
task and the aversive treatment type. With close colour pairs, appetitive-only conditioning
promoted the fastest learning, with the highest number of switches between the two stimuli,
while the opposite was true for quinine.

All treatment types were aversive, but only quinine impacted the positive reinforcement

To quantify whether the moths did, in fact, react aversively upon probing the treatment substances compared to sucrose, and whether there were differences in aversive strengths between the substances, we analysed proboscis contact durations with the droplets across discrimination tasks and treatment types (except appetitive-only, in which we did not present droplets on the CS-). Contact durations were significantly longer for sucrose in comparison to all treatment substances (GLMM Stimulus type - X^2= 226.46, df= 1, p< 0.001; *Fig. 6*) but we found no significant differences in contact durations between the treatment substances. Thus, while the treatment substances were perceived as aversive, there were no treatment-specific differences in feeding responses to indicate disparate aversive strengths. However, the type of negative reinforcement influenced the contacts with sucrose (GLMM Stimulus type: Treatment type - X^2= 29.42, df= 3, p=< 0.001; *Fig. 6*), as quinine-treated animals had significantly shorter feeding durations than animals in all other treatments.

Fig. 6 **Contact duration with sucrose and the different negative reinforcements.** Log transformed contact durations with the droplets on the CS+ (sucrose) and CS- (treatment substances). Shaded points show all the scored contact data for each treatment type. Estimates and their 95% confidence intervals (in black) are drawn from the GLMM (see Methods, Table S6). Differences between treatment types are indicated with letters, where treatments with no difference (p>0.05) share the same letter. Significant difference between the stimulus types is indicated by asterisks (*** p < 0.001).

Discussion

In this study we demonstrate how the underlying context and conditions can modify learning outcomes by comparing the role of various negative reinforcements for visual discrimination in a solitary insect pollinator. By differentially conditioning the hummingbird hawkmoth, a generalist nectarivore, in two colour discrimination tasks, we show that quinine, a commonly used negative reinforcement, produces worse target association in comparison with other substances, such as citric acid, and with appetitive-only conditioning. We link these differences in learning to the effect of the reinforcing substance on modifying reward association and foraging dynamics in freely flying animals, subsequently affecting the individual's accuracy and speed of learning.

Learning asymmetry and innate preferences

Learning dynamics between the close and distant colour experiments varied greatly, with animals making rapid and reliable associations to the rewarded blue or yellow but failing to make similar associations to the cyan flowers. These results correspond to previously reported colour discrimination results in honeybees and bumblebees (Avarguès-Weber et al., 2010; Dyer & Chittka, 2004) where discrimination thresholds for perceptually close colours do not reach the same criteria as those of distinct colours, even with strongly reinforced training protocols.

In our experiments, the differences between the colour discrimination tasks are unlikely to be explained by the challenge of perceptually distinguishing the colours, as *M. stellatarum* can distinguish spectrally similar colours differing by as low as 1-2 nm in the blue 400 and 480 nm range, a spectral resolution even higher than in honeybees (Telles et al., 2016), and can also use achromatic cues to discriminate stimuli (Kelber, 2005; van der Kooi & Kelber, 2022). The cyan and blue stimuli we used had both achromatic and chromatic differences that were large enough to be exploited by the hawkmoths, and correspondingly, they made strongly biased colour choices during their initial preference test, which would not be possible if they were unable to discriminate between the colours.

Rather, the differences in choice rates between the distant and close experiments could have emerged from the moths' strong innate colour bias, since individuals show untrained preferences for blue or yellow (Kelber, 1996, 2002). The cyan stimulus used in this setup was less preferred in comparison, as evidenced by the results of the pretraining where even after being absolutely conditioned to cyan for 3 days, moths chose to feed on blue when given a choice. While the change from their initial choice rates suggests that learning did take place

in both discrimination tasks, new reward associations with the cyan flowers might have been
created without losing their initial preference for blue. As a result, at the end of the differential
conditioning period, most moths chose equally between cyan and blue rather than
associating with a single colour as with the blue-yellow stimuli. Innate colour preferences
could have also led to the asymmetric learning acquisitions in the distant colour experiments
between the groups with blue and yellow as their CS+. In this species, a larger proportion of
animals have a spontaneous preference for blue over yellow (Kelber 1997). Subsequently,
more moths with an innate blue preference would have been pretrained through absolute
conditioning to yellow artificial flowers than the other way around. Thus more animals with
blue as their CS+ would have been 're-learning' a colour, a task requiring more cognitive
flexibility and considered more challenging (Izquierdo et al., 2017; Raine & Chittka, 2012;
Young et al., 2024).

Another possible explanation for the decreased target acquisition in the close colour
experiments could emerge from the moths allocating their attention to the salient blue
stimulus and not forming immediate associations to the rewarded target. Such phenomena
as blocking or overshadowing of learning due to the presence of a stimulus with high valence
in the same environment have been suggested in previous learning studies (Balkenius &
Kelber, 2006) in this species.

An interesting aspect of the hawkmoths' colour associations is demonstrated by their
foraging dynamics, where we found a low colour fidelity in the close colour setup, whereas
animals rarely switched between the yellow and blue colours. This could indicate a
categorical grouping of colours, analogous to human-defined groups, where colours across a
range of chromatic and achromatic properties are grouped into the same class, such as
blues, greens, yellows, etc.. Seen through the lens of foraging, this would suggest that
hawkmoths are less prone to switching between flowers in different colour categories, but
when their preferred colour is unavailable or unrewarded, might sample instead from flowers
with similar colours. While categorical colour perception has been uniquely linked to primates
(Kelber, 2016), generalization of colours by insects, where behavioural responses do not
correlate to the colours' distance in their colour space (Benard & Giurfa, 2008; Troje, 1993),
might show that similar processes are utilized.

Tasks requiring discrimination between close colours have been described as 'challenging'
due to the lower learning or target acquisition levels and have subsequently been used to
demonstrate the benefit of strong negative reinforcements for learning in insects (Avarguès-
Weber et al., 2010; Dyer & Chittka, 2004; Dyer & Howard, 2023; Giurfa, 2004). However,
what is viewed as a learning deficit might instead be the animal's strategy to benefit from
exploring within a colour category rather than making strong associations to a particular

colour within a category and restricting foraging flexibility. Here, we interpret the results of
discrimination learning as an interaction of their innate preferences, which are known to be
reliably recalled even after training or hibernation periods (Gumbert, 2000; Kelber, 2010b;
Menzel, 1968), and the functional basis of colour recognition for hawkmoths while foraging.

Tastes as negative reinforcements in hawkmoths

Colour discrimination learning was influenced by the type of negative reinforcement used
during the differential conditioning, which points to a disparate negative valence associated
with the treatment substances. The 5-way comparison of treatments in the distant colour
experiment, albeit statistically inconclusive, produced citric acid, quinine, and appetitive-only
conditioning as the more reliable training conditions, suggesting that these substances were
perceived differently from at least water and salt solution. The next series of experiments,
where moths were trained to discriminate close colours, demonstrate that citric acid and
quinine were also perceived differently from each other. This points strongly to aversion
being generated not only due to the mismatch of the substance to the expectation of a sugar
reward, but also due to the specific substances being perceived uniquely. Such differences in
perception could emerge from a negative sensory experience on sampling the substance,
such as that caused by an aversive taste or viscosity (Josens & Farina, 2001). The
hawkmoth's proboscis have sensilla with gustatory and mechanosensory receptors which
they use to assess food quality (Kelber, 2003). Studies on the gustatory coding of a closely
related hawkmoth, *Manduca sexta*, suggest that tastes in this species are represented in a
tastant-dependent manner rather than in the categories - bitter, salt, sour, sugar, umami-
typically known in other animals (Reiter et al., 2015). In another moth, *Heliothis virescens*, it
was shown that two bitter substances, one of which was quinine, produced separate
behavioural and neural responses (Kvello et al., 2010). We did not find differences in the
proboscis contact durations between the treatment substances which could have been due
to the limitation of visually scoring feeding from the videos, where finer differences in contact
durations might not have been captured. It is also possible that all substances were aversive
beyond a threshold at which the moths show highly stereotyped behaviour of quickly
retracting their proboscis. However, the differences in learning outcomes as well as previous
studies on moth gustation suggest that the negative reinforcements were perceived as
unique gustatory sensations.

Post-ingestional physiological effects, such as malaise, or a degradation of the reward quality
of sugar from contacting the treatment substances could have also played a role in learning.
In our experiments, contact durations with the treatment substances were significantly lower

than those with the sucrose rewards and typically lasted less than a second, making it
unlikely that they ingested toxic quantities of the substance. Instead, upon contacting quinine
on the CS-, moths made much shorter contacts with sucrose and were observed to
sometimes be repelled on contacting the sucrose droplets. This strongly evidences the
inhibition of sugar receptors by bitter substances that has been reported in many insects (De
Brito Sanchez et al., 2005; Dethier & Bowdan, 1992; Jørgensen et al., 2007; Liscia & Solari,
2000; Meunier et al., 2003). Thus, using quinine as a negative reinforcement could have
prevented animals from making accurate associations to the rewarded stimulus. Additionally,
we also found a lower switching rate for quinine reinforced animals with the close colour
stimuli which could have been due to the lack of motivation to diverge from their preferred
colour since the alternate was also not perceived as rewarding.

669 Learning visual discrimination

Previous work on insects suggests that differential conditioning using strongly aversive
substances improves target recognition (Avarguès-Weber et al., 2010; Dyer & Chittka, 2004;
Dyer & Howard, 2023; Rodríguez-Gironés et al., 2013) as a consequence of the punishment
associated with making incorrect choices, along with the positive conditioning to the
rewarded reinforcement. The faster and stronger associations thus formed are discussed as
emerging from the punished individual being more attentive to the task (Chittka et al., 2003)
and retaining more information associated with the stimuli (Giurfa, 2004). Although learning
studies in Lepidoptera have demonstrated that feeding behaviour can be modified in
response to non-sugar compounds (Broadhead & Raguso, 2021; Rodrigues et al., 2010), in
our experiments, moths treated with strong negative reinforcements such as concentrated
quinine or salt solution did not perform better than the animals that only encountered an
empty flower. Rather, finding their preferred colour sugar-empty motivated the animals to
sample the alternative colour, resulting in better colour associations with the rewarded target.
Interestingly, we found no correlation between the number of foraging attempts and the
learning acquisition, but rather, in the present discrimination context, the animal's tendency to
explore foraging alternatives and encounter the rewarded target predicted task success

Hawkmoths are solitary insects making few and therefore more restricted feeding choices in
comparison to many eusocial foragers that make multiple foraging bouts to feed the colony.
Moreover, insects within a eusocial lifestyle may display more exploratory behaviour since
they may also supplement their nutrition and fitness through the hive (Hendriksma et al.,
2019), whereas risk aversion could be a strong limiting factor in the foraging behaviour of
solitary pollinators. In the case of the hummingbird hawkmoth, we found that conditioning

with an aversive stimulus diminished their motivation to explore foraging alternatives in the
vicinity of the stimulus, instead of providing association cues to aid visual discrimination.
Since hawkmoths form feeding associations in a highly reward-motivated manner (Balkenius
et al., 2008; Kelber, 1996), with the ability to discriminate between sugar types and learn
colours associated with their preferred sugars (Kelber, 2003), training protocols with
disparate reward associations might motivate learning, while also preventing any loss in
performance caused by harsh punishments. Such a conditioning method would also
represent a naturalistic foraging situation for this species, as their longer lifespans and
migration activities (Kelber, 2010b) require them to adapt to new environments with different
resources and select among the most energy efficient food source.

The contrast in learning outcomes between eusocial and solitary pollinator species in
response to negative reinforcements suggests that aversion does not pervasively enhance
associations across insect groups. This highlights the necessity of interpreting learning and
foraging behaviour not only through the animals' sensory ecology, but also as a factor of their
life history.

708 Conclusions

This study highlights that visual learning in insects can be strongly influenced by the
underlying context and conditions. Contrary to findings in eusocial insects where learning
success has been linked to differential conditioning with strong negative reinforcements,
hummingbird hawkmoths showed poorer target association when trained with quinine,
compared to citric acid or appetitive-only conditioning with no aversive substance. In addition
to interfering with sucrose perception and thus hindering reward association, we demonstrate
that quinine was not conducive to the hawkmoths' exploratory behaviour during foraging.
These results underscore the role of foraging dynamics and an animal's sensory ecology in
shaping learning outcomes.

719 Acknowledgements

720 We thank Sophie Fetter for assistance with scoring videos, and Samira Giger and Lucas
721 Heger for helping with pilot studies to establish methods. We are also grateful to Valerie
722 Kuklovsky for providing useful input, and to all the students involved with animal care during
723 this study.

Competing interests

The authors declare no competing or financial interests.

Author contributions

Conceptualization: A.N.M., A.S.; Methodology: A.N.M., A.S., J.J.F.; Validation: A.N.M., A.S.; Formal analysis: A.N.M, J.J.F.; Investigation: A.N.M.; Resources: A.S.; Data curation: A.N.M.; Writing - original draft: A.N.M.; Writing - review & editing: A.N.M., A.S., J.J.F.; Visualization: A.N.M.; Supervision: A.S.; Funding acquisition: A.S

Funding

We acknowledge funding to A.S. by the Bavarian Academy of Sciences and Humanities, the Zukunftskolleg Konstanz, and the Emmy Noether Programme of the DFG (STO 1255/4-1).

Supplementary Material

Figures S1 to S4. Tables S1 to S6.

References

Avarguès-Weber, A., & Giurfa, M. (2014). Cognitive components of color vision in honey bees: How conditioning variables modulate color learning and discrimination. *Journal of Comparative Physiology A*, *200*(6), 449–461. https://doi.org/10.1007/s00359-014-0909-z

Avarguès-Weber, A., Sanchez, M. G. de B., Giurfa, M., & Dyer, A. G. (2010). Aversive Reinforcement Improves Visual Discrimination Learning in Free-Flying Honeybees. *PLOS ONE*, *5*(10), e15370. https://doi.org/10.1371/journal.pone.0015370

Ayestaran, A., Giurfa, M., & Sanchez, M. G. de B. (2010). Toxic but Drank: Gustatory Aversive Compounds Induce Post-ingestional Malaise in Harnessed Honeybees. *PLOS ONE*, *5*(10), e15000. https://doi.org/10.1371/journal.pone.0015000

Balkenius, A., & Kelber, A. (2006). Colour preferences influences odour learning in the hawkmoth, Macroglossum stellatarum. *Naturwissenschaften*, *93*(5), 255–258. https://doi.org/10.1007/s00114-006-0099-9

Balkenius, A., Kelber, A., & Balkenius, C. (2006). Modelling Multi-modal Learning in a

Hawkmoth. In S. Nolfi, G. Baldassarre, R. Calabretta, J. C. T. Hallam, D. Marocco, J.-

A. Meyer, O. Miglino, & D. Parisi (Eds.), *From Animals to Animats 9* (Vol. 4095, pp.

422–433). Springer Berlin Heidelberg. https://doi.org/10.1007/11840541_35

Balkenius, A., Kelber, A., & Balkenius, C. (2008). How Do Hawkmoths Learn Multimodal

Stimuli? A Comparison of Three Models. *Adaptive Behavior*, *16*(6), 349–360.

https://doi.org/10.1177/1059712308092955

Balkenius, A., Rosén, W., & Kelber, A. (2006). The relative importance of olfaction and vision

in a diurnal and a nocturnal hawkmoth. *Journal of Comparative Physiology A*, *192*(4),

431–437. https://doi.org/10.1007/s00359-005-0081-6

Benard, J., & Giurfa, M. (2008). The cognitive implications of asymmetric color generalization

in honeybees. *Animal Cognition*, *11*(2), 283–293. https://doi.org/10.1007/s10071-007-

0112-5

Broadhead, G. T., & Raguso, R. A. (2021). Associative learning of non-sugar nectar

components: Amino acids modify nectar preference in a hawkmoth. *Journal of*

Experimental Biology, *224*(12), jeb234633. https://doi.org/10.1242/jeb.234633

Brooks, M. E., Kristensen, K., Benthem, K. J. van, Magnusson, A., Berg, C. W., Nielsen, A.,

Skaug, H. J., Mächler, M., & Bolker, B. M. (2017). *Modeling zero-inflated count data*

with glmmTMB (p. 132753). bioRxiv. https://doi.org/10.1101/132753

Buatois, A., Pichot, C., Schultheiss, P., Sandoz, J.-C., Lazzari, C. R., Chittka, L., Avarguès-

Weber, A., & Giurfa, M. (2017). Associative visual learning by tethered bees in a

controlled visual environment. *Scientific Reports*, *7*(1), 12903.

https://doi.org/10.1038/s41598-017-12631-w

De Brito Sanchez, M. G., Giurfa, M., De Paula Mota, T. R., & Gauthier, M. (2005).

Electrophysiological and behavioural characterization of gustatory responses to

antennal 'bitter' taste in honeybees. *European Journal of Neuroscience*, *22*(12),

3161–3170. https://doi.org/10.1111/j.1460-9568.2005.04516.x

de Brito Sanchez, M. G., Serre, M., Avarguès-Weber, A., Dyer, A. G., & Giurfa, M. (2015).

Learning context modulates aversive taste strength in honey bees. *Journal of*

Experimental Biology, *218*(6), 949–959. https://doi.org/10.1242/jeb.117333

Dethier, V. G., & Bowdan, E. (1992). Effects of alkaloids on feeding by Phormia reginaconfirm

the critical role of sensory inhibition. *Physiological Entomology*, *17*(4), 325–330.

https://doi.org/10.1111/j.1365-3032.1992.tb01030.x

Dyer, A. G., & Chittka, L. (2004). Fine colour discrimination requires differential conditioning

in bumblebees. *Naturwissenschaften*, *91*(5), 224–227.

https://doi.org/10.1007/s00114-004-0508-x

Dyer, A. G., & Howard, S. R. (2023). Aversive reinforcement improves visual discrimination

learning in free-flying wasps (Vespula vulgaris). *Behavioral Ecology and Sociobiology*,

77(9), 101. https://doi.org/10.1007/s00265-023-03375-x

Friard, O., & Gamba, M. (2016). BORIS: A free, versatile open-source event-logging software

for video/audio coding and live observations. *Methods in Ecology and Evolution*,

7(11), 1325–1330. https://doi.org/10.1111/2041-210X.12584

Giurfa, M. (2004). Conditioning procedure and color discrimination in the honeybee Apis

mellifera. *Naturwissenschaften*, *91*(5), 228–231. https://doi.org/10.1007/s00114-004-

0530-z

Gumbert, A. (2000). Color choices by bumble bees (Bombus terrestris): Innate preferences

and generalization after learning. *Behavioral Ecology and Sociobiology*, *48*(1), 36–43.

https://doi.org/10.1007/s002650000213

Hendriksma, H. P., Toth, A. L., & Shafir, S. (2019). Individual and Colony Level Foraging

Decisions of Bumble Bees and Honey Bees in Relation to Balancing of Nutrient

Needs. *Frontiers in Ecology and Evolution*, *7*.

https://doi.org/10.3389/fevo.2019.00177

Howard, S. R., Avarguès-Weber, A., Garcia, J. E., Greentree, A. D., & Dyer, A. G. (2019).

Surpassing the subitizing threshold: Appetitive–aversive conditioning improves

805 discrimination of numerosities in honeybees. *Journal of Experimental Biology*,

806 *222*(19), jeb205658. https://doi.org/10.1242/jeb.205658

807 Izquierdo, A., Brigman, J. L., Radke, A. K., Rudebeck, P. H., & Holmes, A. (2017). The neural

808 basis of reversal learning: An updated perspective. *Neuroscience*, *345*, 12–26.

809 https://doi.org/10.1016/j.neuroscience.2016.03.021

810 Johnson, S. D., Moré, M., Amorim, F. W., Haber, W. A., Frankie, G. W., Stanley, D. A.,

811 Coccuci, A. A., & Raguso, R. A. (2017). The long and the short of it: A global analysis

812 of hawkmoth pollination niches and interaction networks. *Functional Ecology*, *31*(1),

813 101–115. https://doi.org/10.1111/1365-2435.12753

814 Jørgensen, K., Stranden, M., Sandoz, J.-C., Menzel, R., & Mustaparta, H. (2007). Effects of

815 two bitter substances on olfactory conditioning in the moth Heliothis virescens.

816 *Journal of Experimental Biology*, *210*(14), 2563–2573.

817 https://doi.org/10.1242/jeb.004283

818 Josens, R. B., & Farina, W. M. (2001). Nectar feeding by the hovering hawk moth

819 Macroglossum stellatarum: Intake rate as a function of viscosity and concentration of

820 sucrose solutions. *Journal of Comparative Physiology A*, *187*(8), 661–665.

821 https://doi.org/10.1007/s00359-001-0238-x

822 Kannegieser, S., Kraft, N., Haan, A., & Stöckl, A. (2024). Visual guidance fine-tunes probing

823 movements of an insect appendage. *Proceedings of the National Academy of*

824 *Sciences*, *121*(6), e2306937121. https://doi.org/10.1073/pnas.2306937121

825 Kelber, A. (1996). Colour Learning in the Hawkmoth Macroglossum Stellatarum. *Journal of*

826 *Experimental Biology*, *199*(5), 1127–1131. https://doi.org/10.1242/jeb.199.5.1127

827 Kelber, A. (1997). Innate Preferences for Flower Features in the Hawkmoth Macroglossum

828 Stellatarum. *Journal of Experimental Biology*, *200*(4), 827–836.

829 https://doi.org/10.1242/jeb.200.4.827

830 Kelber, A. (2002). Pattern discrimination in a hawkmoth: Innate preferences, learning

831 performance and ecology. *Proceedings of the Royal Society of London. Series B:*

832 *Biological Sciences*, *269*(1509), 2573–2577. https://doi.org/10.1098/rspb.2002.2201

Kelber, A. (2003). Sugar preferences and feeding strategies in the hawkmoth Macroglossum

stellatarum. *Journal of Comparative Physiology A*, *189*(9), 661–666.

https://doi.org/10.1007/s00359-003-0440-0

Kelber, A. (2005). Alternative use of chromatic and achromatic cues in a hawkmoth.

Proceedings of the Royal Society B: Biological Sciences, *272*(1577), 2143–2147.

https://doi.org/10.1098/rspb.2005.3207

Kelber, A. (2010a). What a hawkmoth remembers after hibernation depends on innate

preferences and conditioning situation. *Behavioral Ecology*, *21*(5), 1093–1097.

https://doi.org/10.1093/beheco/arq115

Kelber, A. (2010b). What a hawkmoth remembers after hibernation depends on innate

preferences and conditioning situation. *Behavioral Ecology*, *21*(5), 1093–1097.

https://doi.org/10.1093/beheco/arq115

Kelber, A. (2016). Colour in the eye of the beholder: Receptor sensitivities and neural circuits

underlying colour opponency and colour perception. *Current Opinion in Neurobiology*,

41, 106–112. https://doi.org/10.1016/j.conb.2016.09.007

Kvello, P., Jørgensen, K., & Mustaparta, H. (2010). Central Gustatory Neurons Integrate

Taste Quality Information From Four Appendages in the Moth Heliothis virescens.

Journal of Neurophysiology, *103*(6), 2965–2981.

https://doi.org/10.1152/jn.00985.2009

Lee, A. B., Mumford, D., & Huang, J. (2001). Occlusion Models for Natural Images: A

Statistical Study of a Scale-Invariant Dead Leaves Model. *International Journal of*

Computer Vision, *41*(1), 35–59. https://doi.org/10.1023/A:1011109015675

Liscia, A., & Solari, P. (2000). Bitter taste recognition in the blowfly: Electrophysiological and

behavioral evidence. *Physiology & Behavior*, *70*(1), 61–65.

https://doi.org/10.1016/S0031-9384(00)00249-3

Matsumoto, Y., & Mizunami, M. (2002). Temporal determinants of long-term retention of

olfactory memory in the cricket Gryllus bimaculatus. *Journal of Experimental Biology*,

205(10), 1429–1437. https://doi.org/10.1242/jeb.205.10.1429

Menzel, R. (1968). Das Gedächtnis der Honigbiene für Spektralfarben. *Zeitschrift für*

vergleichende Physiologie, *60*(1), 82–102. https://doi.org/10.1007/BF00737097

Meunier, N., Marion-Poll, F., Rospars, J.-P., & Tanimura, T. (2003). Peripheral coding of bitter

taste in Drosophila. *Journal of Neurobiology*, *56*(2), 139–152.

https://doi.org/10.1002/neu.10235

Raine, N. E., & Chittka, L. (2012). No Trade-Off between Learning Speed and Associative

Flexibility in Bumblebees: A Reversal Learning Test with Multiple Colonies. *PLOS*

ONE, *7*(9), e45096. https://doi.org/10.1371/journal.pone.0045096

Ravenscraft, A., & Boggs, C. L. (2016). Nutrient acquisition across a dietary shift: Fruit

feeding butterflies crave amino acids, nectivores seek salt. *Oecologia*, *181*(1), 1–12.

https://doi.org/10.1007/s00442-015-3403-6

Reisenman, C. E., Wong, J., Vedagarbha, N., Livelo, C., & Scott, K. (2023). Taste

adaptations associated with host specialization in the specialist Drosophila sechellia.

Journal of Experimental Biology, *226*(3), jeb244641.

https://doi.org/10.1242/jeb.244641

Reiter, S., Campillo Rodriguez, C., Sun, K., & Stopfer, M. (2015). Spatiotemporal Coding of

Individual Chemicals by the Gustatory System. *The Journal of Neuroscience*, *35*(35),

12309–12321. https://doi.org/10.1523/JNEUROSCI.3802-14.2015

Renoult, J. P., Kelber, A., & Schaefer, H. M. (2017). Colour spaces in ecology and

evolutionary biology. *Biological Reviews*, *92*(1), 292–315.

https://doi.org/10.1111/brv.12230

Rodrigues, D., Goodner, B. W., & Weiss, M. R. (2010). Reversal Learning and Risk-Averse

Foraging Behavior in the Monarch Butterfly, Danaus plexippus (Lepidoptera:

Nymphalidae). *Ethology*, *116*(3), 270–280. https://doi.org/10.1111/j.1439-

0310.2009.01737.x

Rodríguez-Gironés, M. A., Trillo, A., & Corcobado, G. (2013). Long Term Effects of Aversive

Reinforcement on Colour Discrimination Learning in Free-Flying Bumblebees. *PLOS*

ONE, *8*(8), e71551. https://doi.org/10.1371/journal.pone.0071551

Šidák, Z. (1967). Rectangular Confidence Regions for the Means of Multivariate Normal

Distributions. *Journal of the American Statistical Association*, *62*(318), 626–633.

https://doi.org/10.1080/01621459.1967.10482935

Singh, N., Mitra, D., Lenka, R., Chatterjee, P., Basumatary, S. K., & Tripathi, S. (2025).

Among hawkmoths (Sphingidae, Lepidoptera), Macroglossinae dominate pollen

transportation in central and East Himalaya (North-East India). *Arthropod-Plant*

Interactions, *19*(1), 20. https://doi.org/10.1007/s11829-024-10127-9

Stach, S., & Giurfa, M. (2005). The influence of training length on generalization of visual

feature assemblies in honeybees. *Behavioural Brain Research*, *161*(1), 8–17.

https://doi.org/10.1016/j.bbr.2005.02.008

Stöckl, A., Grittner, R., & Pfeiffer, K. (2019). The role of lateral optic flow cues in hawkmoth

flight control. *Journal of Experimental Biology*, *222*(13), jeb199406.

https://doi.org/10.1242/jeb.199406

Stöckl, A., Heinze, S., Charalabidis, A., el Jundi, B., Warrant, E., & Kelber, A. (2016).

Differential investment in visual and olfactory brain areas reflects behavioural choices

in hawk moths. *Scientific Reports*, *6*(1), 26041. https://doi.org/10.1038/srep26041

Stöckl, A. L., O'Carroll, D., & Warrant, E. J. (2017). Higher-order neural processing tunes

motion neurons to visual ecology in three species of hawkmoths. *Proceedings of the*

Royal Society B: Biological Sciences, *284*(1857), 20170880.

https://doi.org/10.1098/rspb.2017.0880

Telles, F. J., Kelber, A., & Rodríguez-Gironés, M. A. (2016). Wavelength discrimination in the

hummingbird hawkmoth Macroglossum stellatarum. *Journal of Experimental Biology*,

219(4), 553–560. https://doi.org/10.1242/jeb.130484

Troje, N. (1993). Spectral Categories in the Learning Behaviour of Blowflies. *Zeitschrift Für*

Naturforschung C, *48*(1–2), 96–104. https://doi.org/10.1515/znc-1993-1-218

van der Kooi, C. J., & Kelber, A. (2022). Achromatic Cues Are Important for Flower Visibility

to Hawkmoths and Other Insects. *Frontiers in Ecology and Evolution*, *10*.

https://www.frontiersin.org/articles/10.3389/fevo.2022.819436

Vehtari, A., Gelman, A., & Gabry, J. (2017). Practical Bayesian model evaluation using leave-

one-out cross-validation and WAIC. *Statistics and Computing*, *27*(5), 1413–1432.

https://doi.org/10.1007/s11222-016-9696-4

Vorobyev, M., & Osorio, D. (1998). Receptor noise as a determinant of colour thresholds.

Proceedings of the Royal Society of London. Series B: Biological Sciences,

265(1394), 351–358. https://doi.org/10.1098/rspb.1998.0302

Young, F. J., Melo-Flórez, L., McMillan, W. O., & Montgomery, S. H. (2024). Reversal

learning of visual cues in Heliconiini butterflies. *Animal Behaviour*, *208*, 69–77.

https://doi.org/10.1016/j.anbehav.2023.11.016