1 Adaptative and ancient co-evolution of integrons with

2 *Xanthomonas* genomes

Elena Colombi[1,2 *], Timothy M. Ghaly[1,2], Vaheesan Rajabal[1,2], Liam D.H. Elbourne[1,2], Michael

Gillings[1] & Sasha Tetu[1,2]

[1] School of Natural Sciences, Macquarie University, Sydney, NSW, 2109, Australia

[2] ARC Centre of Excellence in Synthetic Biology, Sydney, NSW, 2109, Australia

*Corresponding author: elena.colombi@mq.edu.au

Keywords: genome evolution, horizontal gene transfer, niche adaptation, plant-pathogen, mobile genetic

elements

ABSTRACT

Integrons are genetic elements that facilitate gene acquisition. They have been extensively

studied in clinical bacteria, but their evolutionary role in phytopathogens remains

underexplored. Here, we analysed complete genomes of *Xanthomonas* species to

investigate the origin, distribution, and functional dynamics of integrons in this genus. We

found that 93% of genomes harboured integrons. The integron-integrase gene *intI* was

predominantly located downstream of *ilvD*, indicating an ancestral acquisition of integrons,

predating diversification within the genus. Phylogenetic analyses support vertical

inheritance of *intI*, with the exception of rare horizontal gene transfer events, notably in *X.*

arboricola. Despite their widespread presence, full-length *intI* genes and active integron

platforms are only retained in some species, especially *X. campestris*, which shows high

integron gene cassette variability and functional integron activity. In contrast, species such

as *X. cissicola* and *X. phaseoli* exhibit widespread *inI* inactivation, likely occurring early in

their divergence, leading to more stable cassette arrays and conserved integron-associated

phenotypes. The number and diversity of genes within cassette arrays varied significantly by

species and, to a lesser extent, by the ecological context of plant host cultivation. While

most cassettes encoded proteins without a known function, those with annotated roles

were associated with stress response mechanism, competitive exclusion, and plant-

associated functions. Together, our findings demonstrate that integrons in *Xanthomonas*

likely originated from a single ancient acquisition event, preceding genus-wide speciation,

and have co-evolved with *Xanthomonas* pathovars as they adapted to distinct plant hosts.

IMPACT STATEMENT

This study provides the first comprehensive genus-wide analysis of integron evolution and

dynamics in *Xanthomonas*, a globally distributed and versatile plant pathogen. Integrons are

genetic platforms that allow horizontal transfer of genes. They are well characterised in
human pathogens where they mediate transfer of antibiotic resistance genes, but little is
known about their role in plant associated bacteria. We showed that in *Xanthomonas* the
integron platform was ancestrally acquired, yet integrons have undergone repeated lineage-
specific inactivation events. Despite widespread erosion of integron activity, some species
such as *X. campestris* maintain robust and functionally diverse integrons that continue to
shape genome plasticity. Notably, high cassette diversity, combined with the presence of
rare and often uncharacterized genes within these arrays (some potentially involved in
environmental sensing or host interaction) suggest that integrons may serve as reservoirs of
adaptive potential. Our findings reshape current views of integron function beyond
antibiotic resistance and highlight their long-term role in microbial evolution, niche
adaptation, and genome innovation in plant-associated bacteria.
50

51   **DATA SUMMARY**
52   The authors confirm all supporting data, code and protocols have been provided within the
53   article or through supplementary data files.
54

INTRODUCTION

Integrons are genetic elements that facilitate horizontal gene transfer in Bacteria (Stokes &
Hall, 1989), and Archaea (Ghaly *et al.*, 2022). They are ancient structures that have been
involved in the evolution of bacterial genomes for hundreds of millions of years (Rowe-
Magnus & Mazel, 2001). Functional integrons are composed of an integron-integrase gene
(*intI*), an integron recombination site (*attI*), a promoter that drives the expression of the
integrase (P_{int}) and generally also have a second promoter (P_c), oriented for expression of
gene cassette array proteins (Hall & Collis, 1995, Mazel *et al.*, 1998, Escudero *et al.*, 2018).

Gene cassettes are mobile, non-replicating elements, which generally consist of an open
reading frame (ORF) and an *attC* site. They exist as circular molecules when excised from a
cassette array. Although cassettes generally consist of a single ORF, cassettes that have two
ORFs or ORF-less cassettes with promoter activity have been observed (Michael & Labbate,
2010, Tansirichaiya *et al.*, 2019, Blanco *et al.*, 2024). The tyrosine recombinase IntI can
capture circular cassettes and integrate them into the genome by mediating the
recombination between *attC* and *attI* sites, forming arrays of 1 to 200+ gene cassettes. As
well as mediating the insertion of cassettes, IntI can also excise cassettes from an array.
Cassette arrays lacking an integrase are called CALIN (clusters of *attC* sites lacking integron-
integrases) (Cury *et al.*, 2016), and can be mobilized *in trans* by a functional IntI.

Technically, integrons themselves are not mobile, whereas their gene cassette cargo is.
Integron activity can facilitate horizontal gene cassette transfer between prokaryotes,
contributing to the rapid evolution and genetic diversity of Bacteria and Archaea (Escudero

et al., 2018). Because DNA damage and nutrient starvation induce activity of the integron
integrase via bacterial SOS and stringent responses (Guerin *et al.*, 2009, Cambray *et al.*,
2011, Strugeon *et al.*, 2016), acquisition of novel genetic functions is preferentially triggered
during periods of environmental challenge. This provides a mechanism for adaptive
innovation when survival pressures are highest.

Integrons have been extensively studied because of their role in the accumulation and
dissemination of antibiotic resistance genes in clinical isolates (Gillings *et al.*, 2008, Hipólito
et al., 2023). However, integrons are also common in other environments such as marine
environments (Pereira *et al.*, 2020), soil (Ghaly *et al.*, 2019), and the plant rhizoplane
(Rajabal *et al.*, 2024), (Qi *et al.*, 2024). In these environments, the majority of gene cassettes
remain functionally uncharacterised, but predicted functions encode extensive functional
diversity including traits involved in microbe-host interactions (Gillings, 2014, Escudero *et*
al., 2018, Ghaly *et al.*, 2020).

Xanthomonas is a genus of Gram-negative bacteria in the class Gammaproteobacteria that
contains pathogenic strains able to cause disease in more than 400 different plants. In
common with other genera of plant pathogens, *Xanthomonas* also contains strains
associated with plants, that do not cause disease symptoms (Vauterin *et al.*, 1996, Essakhi *et*
al., 2015, Garita-Cambronero *et al.*, 2017). *Xanthomonas* is a large genus that encompasses
more than 35 species, which are subdivided into pathovars (Timilsina *et al.*, 2020).

Integrons have been described in some *Xanthomonas* isolates (Gillings *et al.*, 2005,
Barionovi & Scortichini, 2006, Barionovi & Scortichini, 2008). Using PCR with site-specific
primers annealing to the proximal part of *intI* and to the *attI* region, Gillings and colleagues
screened 32 *Xanthomonas* strains representing 12 pathovars, and found that all the strains
analysed had an integron. In all cases, *intI* was integrated downstream of the acid
dehydratase gene, *ilvD*, however, the majority of integrase genes were predicted to be
inactivated by frameshifts, stop codons, or large deletions. Groups of strains with the same
deletions or stop codons/frameshifts in *intI* usually contained identical arrays of genes.
There was no evidence of integrons generating diversity within pathovars but strong
evidence for diversity between pathovars. With some minor exceptions, individual
pathovars had distinct proximal gene cassette arrays, and every cassette identified was
found only in one pathovar.

Complete genome sequences for hundreds of *Xanthomonas* isolates are now available,
allowing for a systematic and detailed investigation of integrons in these important plant-
associated bacteria. Here, we have examined the presence of integrons in all publicly
available complete genomes of *Xanthomonas*, significantly expanding our knowledge of
these elements, and showing how integrons have been a dynamic component during
Xanthomonas genome evolution.

METHODS

Xanthomonas classification and phylogeny

Complete genomes of *Xanthomonas* were downloaded from the National Center for
Biotechnology Information (NCBI) Database (NCBI txid: 338) (708 genomes, last accessed
September 2024). Taxonomic classifications of the genomes was based on the Genome
Taxonomy Database (GTDB) (Parks *et al.*, 2022) release 2.2.0 using GTDB-Tk v2.4.0
(Chaumeil *et al.*, 2022) (Table S1). The command classify_wf was used with default settings.
GTDB-Tk first aligns 120 single-copy phylogenetic marker genes, then classifies each genome
based on its placement into domain-specific reference trees (built from 47,899 prokaryote
genomes), its relative evolutionary divergence, and average nucleotide identity to reference
genomes in the GTDB. Two genomes (GCA_041228075.1 and GCA_002240395.1) did not
belong to the *Xanthomonas* genus (*Luteibacter* and *Luteimonas*_D, respectively) and were
removed from the dataset. The GTDB-Tk classification system does not strictly adhere to the
traditional nomenclature of *Xanthomonas* strains, sometimes reassigning species. We use
GTDB-Tk classification in order to use a standardised microbial taxonomy method based on
genome phylogeny, and not to attempt to re-classify or rename isolates.

Phylogeny of the *Xanthomonas* genus was inferred using on the alignment of the 120
marker genes produced with the GTDB-Tk command classify_wf using GTDB-Tk command
infer (default setting with --gamma) which uses FastTree 2.1 (Price *et al.*, 2010) to build an
approximately-maximum-likelihood phylogenetic tree. The tree was rooted with
GCA_041228075.1 (*Luteibacter*) then the tip was removed with the R package ape 5.8-1
(Paradis & Schliep, 2019). Single-species phylogeny was built using Realphy (Bertels *et al.*,
2014) which maps genomes to a series of reference genomes via bowtie2 2.0.0-beta7
(Langmead & Salzberg, 2012). From these, multiple sequence alignments are constructed
and used to infer phylogenetic trees via PhyML 3.1 (Langmead & Salzberg, 2012). Trees are
visualised using the ggtree 3.16 (Yu *et al.*, 2017) and ggtreeExtra 2.3 (Xu *et al.*, 2021) R
packages.

Analysis of integrons

Integrons were identified with IntegronFinder 2.0.5 (Néron *et al.*, 2022) (parameters: --local-
max --gbk --promoter-attI) (Table S1). Integron sequences with 5 kb up and downstream
were extracted (fasta files deposited at https://github.com/EC-
MQ/Xanthomonas_integrons) and annotated with Bakta (Schwengers *et al.*, 2021).
Taxonomic origins of integron gene cassette recombination sites (*attC*s) were predicted
using the script attC-taxa.sh (Ghaly *et al.*, 2021, Ghaly *et al.*, 2022). The *attC*-taxa pipeline
(https://github.com/timghaly/attC-taxa) can detect *attC* sites that are conserved among one
of 11 bacterial taxa, comprising: Alteromonadales, Methylococcales, Oceanospirillales,

Pseudomonadales, Vibrionales, Xanthomonadales, Acidobacteria, Cyanobacteria,
Deltaproteobacteria, Planctomycetes and Spirochaetes.

IntI sequences identified by Integron Finder were aligned using PRANK (Löytynoja, 2013) run
with default settings, the alignment was stripped of gaps with the goalign 'clean sites'
command (Lemoine & Gascuel, 2021) and the tree inferred with FastTree 2.1 (Price *et al.*,
2010) using the generalized time-reversible model (--gtr).

We used multiple approaches to annotate the cassette-encoded proteins. Proteins were
functionally annotated with eggNOG-mapper v2 (Huerta-Cepas *et al.*, 2019, Cantalapiedra *et*
al., 2021), executed in DIAMOND (Buchfink *et al.*, 2015) mode. Additionally, Foldseek (Van
Kempen *et al.*, 2024) was used to create a protein structural database with the ProstT5
protein language model (Heinzinger *et al.*, 2024). The database was then used to perform
structural alignment against the SwissProt database using a Foldseek search. Foldseek
convertalis was used to convert the alignment database in a tab-delimited output, only
functional annotations against the Swiss-Prot database with a coverage of 80% and an e-
value < 0.001 were considered.

To identify genes involved in functions of ecological relevance (trace gas oxidation, carbon
cycling, nitrogen cycling, sulphur cycling, phosphorus cycling, iron cycling, plant-microbe
interactions, and osmotic stress tolerance) we used the *EcoFoldDB-annotate* pipeline (Ghaly
et al., 2025).

Plant growth-promoting traits were identified using the Plant Growth-Promoting Traits
Prediction tool (PGPT-Pred) (Ashrafi *et al.*, 2022). SignalP V6.0 (Almagro Armenteros *et al.*,
2019) was used for signal peptide predictions as a marker for identifying transmembrane or
secreted gene products. DefenseFinder (Tesson *et al.*, 2022) was used to detect known anti-
phage systems.

Unique gene orthologues encoded by the cassettes and their distributions were identified
with Proteinortho 6.0.22 (Klemm *et al.*, 2023) run with the -singles option.

To detect genes encoding type III secretion effectors (T3SE), we used a database of 66
previously characterized 66 *Xanthomonas* T3SE retrieved from the EuroXanth platform
(Costa *et al.*, 2024) (Biessy *et al.*, 2025). T3SE in the cassettes were identified with
Proteinortho v6.0.22 (Klemm *et al.*, 2023) by querying the curated effectors database
against the amino acid sequences of the cassette-encoded genes.

Horizontal gene transfer events between species

Alfy 1.0.5 (Domazet-Lošo & Haubold, 2011) was used to guide the detection of the
movement of cassettes between integron arrays. Alfy was run with the -M option, and
selecting only matches with a *P* value <0.05 within a sliding window of 100⬚bp. Only arrays

sharing closest homology (recombination) with at least 10% of their sequence with an array

harboured by a different species were considered. Horizontal gene transfer (HGT) events of

cassettes were further filtered based on average nucleotide identity. FastANI 1.33 (Jain *et*

al., 2018) was run between the arrays and the whole genomes of the strains. To be

considered a genuine HGT event, ANI identity between arrays needs to be >96% and the ANI

between the genomes harbouring them to be <95%, with at least a 2% ANI difference

between cassette and genome. Cytoscape (Shannon *et al.*, 2003) was then used to visualise

the numbers of cassettes transferred between *Xanthomonas* species.

RESULTS AND DISCUSSION

The *Xanthomonas* integron platform is ancestral

Xanthomonas complete genomes classified based on the Genome Taxonomy Database

(GTDB) as "*Xanthomonas*" (n=629) and "*Xanthomonas_A*" (n=78), known also as

Xanthomonas group 2 and 1, respectively (Pena *et al.*, 2024) were analysed for integron

features. We retained the genomes of *Xanthomonas_A,* even though they are classified as a

different genus by GTDB-Tk, because from a plant pathology perspective they are still

considered *Xanthomonas*, and encompass *X. translucens,* an important causal agent of

diseases in cereal crops and forage grasses (Sapkota *et al.*, 2020).

Using IntegronFinder 2.0.5 (Néron *et al.*, 2022), we identified integrons, In0 (integron-

integrases that lack cassettes), or CALINs (clusters of *attC* sites lacking integron-integrases)

in 93% of the genomes (657 genomes) (Figure 1). *X. albilineans* (n=6, group 1) and *X.*

fragariae (n=6, group 2) were the only species with more than 2 genome sequences

available that did not harbour either an *intI* or *attC* sites. However, integrons have been

previously detected in some isolates of *X. fragariae* via PCR amplification (Barionovi &

Scortichini, 2006).

Figure 1 – Distribution of integron platforms in *Xanthomonas* groups 1 and 2. The maximum likelihood phylogenetic tree is based on a concatenated alignment of 120 top-ranked marker proteins. The tree was

rooted at the midpoint. Tips are labelled with species names assigned with Genome Taxonomy Database
phylogeny. Integrons were classified as present if the isolates carried a functional integron, *intI* only or an
attC array only. Scale bar indicates substitutions per site.

Because integrons are widespread in *Xanthomonas*, our first question was whether the
acquisition of the integron module was ancestral. The integron-integrase encoding gene *intI*
was integrated downstream *ilvD* in all genomes, with the two exceptions being isolates of *X.*
oryzae and *X. cucurbitae*. Of the 187 *oryzae* isolates examined, *X. oryzae* pv. *oryzicola* (26
genomes) harbour a truncated *intI* (561 bp) that was not integrated downstream *ilvD*.
However, the integron locus (*intI* and cassette array) is surrounded by transposases, which
could indicate that the integron was moved via transposition. In all seven *X. cucurbitae*
genomes, *intI* was not integrated downstream *ilvD*. However, downstream of *ilvD* there are
two genes without recognizable *attCs* which are encoded in the other integrons (including a
DUF1488 encoding gene which occurs in multiple species).

The different genomic locations of *intI* could be explained by either within-genome
transposition or recombination (*intI* was initially located downstream *ilvD* and then moved),
or by an independent acquisition of *intI* from another source which was integrated in a
different locus. To test this, we constructed the phylogeny of all the *Xanthomonas intIs*
(Figure S1). If *intIs* located elsewhere from *ilvD* were acquired independently, they would
likely have a different evolutionary history, and form separate clades in the *intI* phylogeny.

The phylogeny of *intI* clusters this gene within species, generally being independent of the
integration locus. *X. oryzae intIs* cluster together, and the *X. cucurbitae intIs* form a clade
with pv. *phaseoli* (Figure S1). The only notable exception is *X. arboricola*. *X. arboricola*
strains can carry one of two variant *intIs*. In two strains, *intI* clusters closest to the *X.*
campestris intIs and are truncated and likely not functional. Phylogenetically close *X.*
arboricola strains do not harbour an *intI* at all. The other *X. arboricola* strains carry *intI*
(either full-length or with an early stop codon) that are more distantly related to all other
intIs. These more distantly related *intI* could have been acquired from a source outside
Xanthomonas. We compared the *ilvD-intI* locus sequence in two representative *arboricola*
strains harbouring the two *intI* variants (GCA_018141705.1 and GCA_905367745.1). While
the *ilvD* genes share 97.1% average nucleotide identity (ANI), the *intIs* share 68.2% ANI,
suggesting a recombination event.

We used two full length *intI* variants (one belonging to *X. arboricola*, one to *X. campestris*) as
a query in a BLASTn search on complete genomes in NCBI excluding *Xanthomonas*
(taxid:338). The two closest *intIs* belonged to *Lysobacter* sp. CECT 30171 (locus tag:
LYB30171_00805) and *[Pseudomonas] boreopolis* strain GO2 (locus tag: M3M27_18655)
(both genera in the Xanthodomonaceae family). Curiously, in isolate GO2, *intI* was also
located downstream of *ilvD*. We constructed a phylogenetic tree using *Xanthomonas intI*

sequences longer than 900 bp and included the *intIs* of CECT 30171 and GO2, using *Vibrio*
sp. SCSIO 43136 *intI* as an outgroup (Figure 2). The *intI* variants in *X. arboricola* were more
closely related to the *intI* in GO2 than the other *Xanthomonas intIs*. *ilvD* phylogeny instead
clusters together all *X. arboricola* isolates within the *Xanthomonas* group 2, and GO2 as
basal both to *Xanthomonas* group 1 and 2 isolates. This suggests that, in those *arboricola*
strains, *intI* was acquired via horizontal gene transfer (HGT) from another member of the
Xanthodomonaceae family.

Figure 2 – Phylogeny of *intI*. Sequences of *intI* longer than 900 bp were aligned with PRANK, stripped of
gaps and used to infer the tree with FastTree. *IntI* from *Vibrio* sp. SCSIO 43136 (locus tag: J4N39_08275)
was used as an outgroup. The scale bar indicates substitutions per site. Only bootstrap values 0.80 are
shown. Group 1 *Xanthomonas* are indicated on the side of the plot.

ilvDs of *Xanthomanas* group 1 form a separate group in comparison to group 2 (Figure S2),
while the group 1 *intIs* are nested within the most common *Xanthomanas intIs*. Genetic
exchange between a restricted number of strains of group 2 and the entire group 1 clade
has been reported (Pena *et al.*, 2024), therefore it is plausible that horizontal transfer of an

intI gene from a group 2 *Xanthomonas* species occurred during the early evolutionary
history of group 1 *Xanthomonas*.

Together, this suggests that the acquisition of the integron platform was ancestral to the
Xanthomonas genus, and then *intI* diversified with the species. The exceptions are some *X.*
arboricola strains that could have recruited *intI* from another source, while group 1
Xanthomonas could have acquired *intI* from group 2 *Xanthomonas* in one single gene flow
event.

Additionally, approximately 50% of genomes harbor CALINs, here defined as clusters of at
least two *attC* sites, in loci other than the canonical *ilvD* region and in the absence of a
proximal *intI* gene. In genomes where *intI* is retained, this gene is consistently integrated
downstream of *ilvD,* while these CALINs show spatial separation, being located in other
regions of the genome. Among the species represented by more than 10 genomes, CALINs
not adjacent to *ilvD* were identified at seven distinct chromosomal loci. In *X. arboricola*,
campestris, *cissicola*, *euvesicatoria*, and *hortorum*, CALINs consistently occurred adjacent to
lamG and/or *secF*. In *X. cissicola*, additional CALINs were located near *recD* and a gene
encoding a NAD(P)/FAD-dependent oxidoreductase. In *X. oryzae*, two strains possessed
CALINs adjacent to an outer membrane protein gene, while in *X. translucens*, CALINs were
observed at two distinct genomic positions. One *X. campestris* strain (GCA_028749605.1)
harboured a CALIN on a conjugative plasmid. The relatively conserved positioning of CALINs
across species supports the hypothesis of ancestral integron activity. This contrasts with
experimental data from *Escherichia coli*, where cassette libraries integrated via *attG* sites
(consensus recombination sequences that are not *attC* or *attI*) exhibited a broad distribution
across numerous genomic loci, indicating that *de novo* integration is typically less site-
specific (Loot *et al.*, 2024).

Acquisition of the integron platform is ancestral but its activity is progressively lost

To investigate integron integrase activity and the complement of integron gene cassettes in
Xanthomonas genomes, we focused on species which had more than 10 genomes in our
dataset: *campestris* (n=111) (Figure 3), *cissicola* (n= 124) (Figure 4), *arboricola* (n=41) (Figure
S3), *translucens* (n=46) (Figure S4), *euvesicatoria* (n=46) (Figure S5), *oryzae* (n=161) (Figure
S6), *phaseoli* (n=23) (Figure S7), *hortorum* (n=12) (Figure S8). A detailed description of the
integrons in these eight species is reported in the Supplementary Results Section.

Although integron associated sequences are common in *Xanthomonas* groups 1 and 2, most
strains appear to have lost integron integrase activity, and the distribution of functional
integrons appears to be species specific. However, among the species analysed, *X.*
campestris stands out for maintaining a largely functional integron platform (Figure 3). A
high proportion (85.6%) of its genomes carry a full-length *intI* gene, and integrons are

consistently flanked by extensive cassette arrays (ranging from 3 to 29 *attCs*, averaging 14.3
per genome). Cassette composition is highly variable even among strains of the same
pathovar, suggesting that, in *X. campestris*, integron activity drives rapid diversification
within genetically related groups. This robust activity is also evident in the gene content:
216 orthologous cassette-encoded genes were identified, 41.1% of which are singletons
(observed in only one instance in this dataset). Despite this extensive variability, a gene
encoding a DUF1488 domain-containing protein is present in the cassettes of 81.5% of
genomes, while *symE* (a toxin-encoding gene) is found in 56.8% of the cassette arrays. This
level of integron retention suggests that active integrons continue to play a key role in the
adaptive evolution of *X. campestris*.

Figure 3 – Integrons in *X. campestris*. The phylogeny of *X. campestris* was built using Realphy, using
GCA_000007145.1 as a reference genome. The tree was rooted with GCA_000972745 (*X. arboricola*) then
the tip was removed from the tree. Tip labels show pathovars. The scale bar indicates substitutions per
site. A) Indicates the presence of a complete integron or *attC* sites in the *ilvD* locus, B) indicates the
presence of *intI*, C) indicates the presence of a CALINs (cluster of *attC* sites) not in the ilvD locus. D)
indicates whether IntI is predicted to be functional (full length). E) indicates the number of *attC* sites
(number of cassettes) in the genome. Panel F) represents the distribution of orthologous genes among all
cassettes carried by the corresponding isolate inferred with Proteinortho. Orthologous genes are listed in
decreasing order based on the number of strains within the species that carry them.

X. arboricola exhibited a modest preservation of the *intI* gene with 7% of strains retaining a
full-length *intI*. In the remaining genomes, *intI* is either deleted or contains an early stop

codon. The *X. arboricola* phylogeny divides strains into two clades, clade A which includes
pathovars that cause diseases in *Prunus*, *Juglans*, and *Corylus* spp. (pvs. *pruni*, *juglandis*, and
corylina, respectively), and clade B which comprises many isolates lacking metadata (Figure
S3). Among clade A isolates, none possess a full-length *intI*. In contrast, clade B strains that
carry *intI*, either full-length or truncated (accounting for 58.8% of clade B isolates), harbor a
variant likely acquired via HGT, suggesting the acquisition event dated prior to the
diversification of this clade. Strains of clade B also carried more cassettes and a broader
array of genes compared to clade A.

X. translucens (Figure S4) and *X. euvesicatoria* (Figure S5) emphasize the dynamic nature of
integrons in *Xanthomonas*. Both species exhibit high frequencies of full-length *intI* (22.7%
and 45.6% respectively) and variability in cassette content is high (44.6% and 50%
respectively).

X. oryzae includes pathogens of rice (pv. *oryzae* and pv. *oryzicola*), and pathogens of a
pervasive weed species that grows along rivers and canals surrounding rice paddies (pv.
leersiae). However, only X11-5A, a weakly pathogenic strain (Triplett *et al.*, 2011) distantly
related to the above-mentioned pathovars carries a full-length *intI*. Divergent cassette
compositions were observed among the pathovars (Figure S6). This supports the hypothesis
that integron activity in *X. oryzae* was ancestrally active, followed by progressive
inactivation concomitant with pathovar diversification.

None of the *X. cissicola* (Figure 4) and X. *phaseoli* (Figure S7) isolates retain a full-length,
functional *intI*. However, 28.7% and 38% of the genes carried by the cassettes in these
species, respectively, appear in one isolate only (singletons). *X. cissicola* comprises isolates
commonly known as *X. citri*, an important pathogen able to infect many plants including
Citrus (pv. *citri*). *X. citri* pv. *citri* includes three recognized pathotypes: A, A*, and AwA. All
pv. *citri* genomes analysed here, which include both A and AwA pathotypes, harbor a CALIN
element adjacent to *secD* and lack the integrase gene *intI*. All the CALINs in the pv. *citri*
contain the same two gene cassettes, which are not present in any other genome analysed
in this study, suggesting that the integron platform was inactivated prior to pathotype
diversification (Figure 4). A recent genomic study of 95 pv. *citri* strains estimated that the
diversification of these pathotypes occurred 1,730 to 5,663 years ago (Patané *et al.*, 2019).
Given that citrus domestication is thought to have occurred at least 2,000 years ago (Rao *et*
al., 2021), the inactivation event likely predates the domestication of the host genus. One of
the two cassettes encodes a protein homologous to members of the late embryogenesis
abundant (LEA) protein family, which is known to confer protection against water deficit in
bacteria (Raga-Carbajal *et al.*, 2022). This indicates that the cassette's function could be
oriented toward general environmental stress rather than mediating specific interactions
with the plant host.

Figure 4 – Integrons in *X. cissicola*. The phylogeny of *X. cissicola* was built using Realphy, using
GCA_000007165.1 as a reference genome. The tree was rooted with GCA_000009165.1 (*X. euvesicatoria*)
then the tip was removed from the tree. Tip labels show pathovars, which were assigned from the
literature search. The scale bar indicates substitutions per site. A) Indicates the presence of a complete
integron or *attC* sites in the *ilvD* locus, B) indicates the presence of *intI*, C) indicates the presence of a
CALINs (cluster of *attC* sites) not in the ilvD locus. D) indicates the number of *attC* sites (number of
cassettes) in the genome. Panel E) represents the distribution of orthologous genes among the all the
cassettes carried by the corresponding isolate inferred with Proteinortho. Orthologous genes are given in
decreasing order based on the number of strains within the species that carry them.

Gene cassette abundance is species specific.

We investigated whether variation in integron cassette array size, quantified by the number
of *attC* sites per genome, is primarily shaped by the host genetic background (i.e.,
Xanthomonas species) or by the ecological context of isolation, categorized by cultivation
type (crop, non-crop, crop tree, or ornamental host plants). We used only the species that
had more than 10 genomes in our dataset, additionally, we excluded genomes lacking
information on the host of isolation, and a strain isolated from mud. We performed a
Kruskal-Wallis test to assess if there were statistically significant differences in *attC* number
associated with bacterial species and the type of plant the strains were isolated from. The

results showed significant variation in *attC* count across both factors (Figure S9). Regarding
the cultivation type, the average mean was 12.1 *attC*s for ornamental, 9.77 for non-crop,
6.01 for crop and 3.05 for tree crop. Dunn's test with Bonferroni adjustment showed
statistically significant (p<0.001) differences between strains isolated from crops and non-
crops, between crops and ornamentals, non-crop and tree crops and between ornamentals
and tree-crops. Dunn's test identified three distinct statistical groupings among the
Xanthomonas species, with *X. oryzae* and *X. phaseoli* forming the group with the fewest
cassettes, and *X. campestris* having the highest number.

To assess the contribution of species identity and cultivation type on variation in cassette
number (number of *attC* sites) across samples, we performed a PERMANOVA analysis. The
results indicated that cultivation explained 8.78% of the variation in number of *attC* sites (p-
value = 0.001), suggesting that the type of cultivation has a significant, though relatively
small, effect on the observed variation. In contrast, the species the isolate belong to
accounted for a much larger proportion, explaining 48.17% of the variation (p-value =
0.001). This highlights the significant role of taxonomy at the species level in shaping the
number of *attC* sites, with its influence being far more pronounced than that of cultivation.

Cassette arrays in *Xanthomonas* harbor diverse but poorly characterized gene functions

Prediction of the taxonomic origins of integron *attC*s was performed with a classification
tool developed for *attC*s, which uses both sequence and structural homology information
(Ghaly *et al.*, 2021), to identify matches to known *attC* 'types'. The method classified all *attC*
sites in the *Xanthomonas* strains analysed as originating within the *Xanthomonadales*.
Therefore, there is no evidence of long-distance acquisition of cassettes, and the cassettes
appear to circulate at least within the *Xanthomonadales* family.

Arrays greatly varied greatly in length across genomes, with a maximum of 46 cassettes in a
single genome (X. *arboricola_F* GCA_040182365.1) and 33 cassettes in a single array (X.
arboricola GCA_041475745.1 and GCA_041475785.1) (Table S1). Despite the majority of the
isolates carrying inactive integron modules, with *intI* deleted or truncated, the cassette
arrays appear to have been extremely dynamic before the inactivation of the integron
integrase.

Overall, the 706 *Xanthomonas* genomes carried 1004 cassettes arrays, with a total of 4,773
cassettes, within these, Proteinortho (Klemm *et al.*, 2023) identified 1,087 different
orthologous genes. Of the 1,087 orthologous genes identified by Proteinortho, 476 (43.8%)
were present as singletons, unique to one isolate.

Of the 1,087 orthologous genes, only 11.9% could be classified into a known Clusters of
Orthologous Groups of proteins (COG) category (Galperin *et al.*, 2025). This

underrepresentation of known COGs among cassette proteins is well known (Rowe-Magnus
et al., 2003, Mazel, 2006, Boucher *et al.*, 2007, Ghaly *et al.*, 2024, Rajabal *et al.*, 2024). Of
the most represented functions, excluding unknown and recombination and repair, 1.47%
of the protein had a putative role in transcription, 0.9% in amino acid transport and
metabolism, and 0.55% in defence mechanisms. To gain further insight into possible
cassette functions, we performed both sequence-based homology searching against
eggNOG 5.0 (Huerta-Cepas *et al.*, 2019), and protein structural homology searching against
the Swiss-Prot database. From this combined approach, putative functions could only be
assigned to 6.8% of the proteins. Of these annotated cassette-encoded proteins, 47.7%
were classified either as transposase, integrases or insertion sequences (4.8% of the total).
The presence of insertion sequences and transposases within gene cassettes (Ghaly *et al.*,
2024), and targeting *attC* sites (Tetu & Holmes, 2008, Post & Hall, 2009), has been
frequently observed in past integron studies.

Signal peptides were identified in 11.6% of cassette-encoded proteins, using signalP6.0
(Teufel *et al.*, 2022). Again, transmembrane and secreted proteins are commonly encoded
by gene cassettes in Bacteria (Rowe-Magnus *et al.*, 2003, Ghaly *et al.*, 2023) and are
hypothesized to help facilitate interactions with their broader environment. Additionally,
PGPT-Pred (Ashrafi *et al.*, 2022) predicted 6.7% of cassette-encoded proteins to have plant
growth promoting traits such as biofertilization, plant signal production and stress control.
However, the most common category, at 4.9% of the total proteins, were categorised to
have competitive exclusion functions. These included antimicrobial resistance and
detoxification functions, and toxin-antitoxin systems, which in a plant pathogenic context
could contribute to niche adaptation by enhancing survival in a competitive microbial
environment. One gene was classified by the EcoFoldDB (Ghaly *et al.*, 2025) pipeline as
involved in spermidine production, which can play an important role in plant growth and
stress (Dunn & Becerra-Rivera, 2023).

DefenseFinder (Tesson *et al.*, 2022) identified 20 orthologous genes (1.8%) as genes
involved in defences against phages. Twelve different defence systems were detected, these
included restriction-modification and abortive infection systems, but also the newly
described Kiwa (Zhang *et al.*, 2023) and Shedu defence systems (Loeff *et al.*, 2025).

Using a database of previously characterized type three secretion system effectors (T3SEs)
retrieved from the EuroXanth platform (Costa *et al.*, 2024, Biessy *et al.*, 2025), we identified
two T3SEs: AvrBs3 (a Transcription Activator-Like effector, TALe) in African-like pv. *oryzae*
(Schepler-Luu *et al.*, 2023), and XopAF2 in three *X. vasicola* isolates. T3SEs are protein
secreted via the type III secretion system (T3SS) that suppress or induce plant defences.
TALes are a particular type of T3SE, once inside the host cell, they translocate to the nucleus
where their unique domain of tandemly arranged 34-amino acid repeats mediates binding
to specific promoter elements. Several TALes are known to induce host SWEET sucrose
uniporter genes, thereby facilitating sucrose efflux from xylem parenchyma into the

apoplasm at the infection sites. TALes may enhance disease by targeting susceptibility
genes, or may trigger a resistance response, and are therefore important for pathogen host
range and virulence (Schepler-Luu *et al.*, 2023).

Given that the majority of cassette-encoded proteins are of unknown function and often
occur as singletons (observed only in one strain), the phenotypic impact of these cassettes
remains unclear. It is uncertain whether they confer a selective advantage or are maintained
in the arrays through neutral processes. A cassette that is conserved across multiple species
is more likely to encode a beneficial function. The most widespread cassette, identified in 13
species, encodes *symE* and the non-coding small RNA *sRNA-Xcc1*. *sRNA-Xcc1* is known to be
activated by regulatory elements of the T3SS (Chen *et al.*, 2011), suggesting a potential role
in plant host colonization. In *E. coli*, *symE* forms part of the SymE-SymR toxin-antitoxin
system, where SymE induces nucleoid condensation, disrupts DNA replication and
transcription, and causes double-strand DNA breaks (Thompson *et al.*, 2022). Regulation in
E. coli occurs via the cis-encoded small RNA *SymR* (Kawano *et al.*, 2007), whereas in the
integron cassettes, *sRNA-Xcc1* is located upstream of *symE*, suggesting a divergent
regulatory mechanism. Notably, unlike most integron-associated genes oriented to be
transcribed from the *Pc* promoter (typically located within *intI*), both *sRNA-Xcc1* and *symE*
are oriented in the opposite direction and not under *Pc* control.

Excluding transposases, the second most recurrent cassette, shared among 8 species,
encoded a protein of unknown function. The third most recurrent, present in 7 species,
included two hypothetical proteins and a gene conferring resistance to bleomycin.
Bleomycin resistance genes have been previously identified as enriched in gene cassettes
relative to total metagenomes in environmental samples, as shown by both cassette-
targeted amplicon sequencing and shotgun metagenomics (Ghaly *et al.*, 2023). One of the
recurring cassettes is an "empty" cassette which can occur multiple times in the same array.
This cassette could have promoter activity as it was previously suggested in other species
(Michael & Labbate, 2010, Tansirichaiya *et al.*, 2019, Blanco *et al.*, 2024).

Detection of inter-species horizontal transfer of cassette arrays

To assess interspecies cassette gene sharing, we examined pairs of species that most
frequently share orthologous genes. As expected, species harbouring a greater number of
cassettes tend to share more orthologous genes with others. For example, *X. arboricola* and
X. campestris shared the highest number of orthologous genes (21), with *X. campestris*
sharing the most overall (57 orthologous genes across multiple species).

Orthologous genes encoded in the cassettes are shared between *Xanthomonas* species and
even between group 1 and 2 *Xanthomonas*. We then looked for more recent evidence of
HGT of cassettes present in multiple species. We used Alfy v1.0.5 (Domazet-Lošo & Haubold,

2011) to identify putative regions subject to HGT, selecting those based on the difference in
ANI between the cassette and the genomes harbouring them. With this approach, we
detected the movement of 95 cassettes between 38 pairs of species (Figure 5). The highest
flux of cassettes (10) was between *X. arboricola* and *euroxanthea,* and between *X.*
campestris and *X. hortorum.* However, *X. campestris* is the species with the highest flux of
cassettes compared to any other species (36). One of the cassettes that showed evidence of
horizontal transfer was that carrying *symE* and *sRNA-Xcc1.*

Figure 5 – Horizontal transfer of cassettes among *Xanthomonas* species. Cytoscape network illustrating
transfer of cassettes between species. Nodes represent *Xanthomonas* species, species containing more
than 10 genomes in the dataset are coloured as in Figure 1 (tip labels). Line width between species is
proportional to the number of cassettes moved between species.

CONCLUSIONS

Our comprehensive analysis of 706 complete genomes reveals that the acquisition of the
integron platform is likely an ancestral event in the evolution of the *Xanthomonas* genus.
The widespread conservation of the *intI* gene downstream of *ilvD* across species, along with
intI phylogeny, strongly supports the early acquisition of this mobile genetic element. Only
in *X. arboricola* clade B, the "original" *intI* appears to have been substituted via HGT with an
intI from another member of the Xanthodomonaceae family.

Despite the ancestral acquisition, our results show that integron activity, defined by the
presence of full length, potentially functional *intI* genes, and large and diverse cassette
arrays, has been progressively lost in many *Xanthomonas* lineages. This inactivation appears
to be species-specific and often predates major evolutionary or ecological transitions, such
as pathovar diversification, such as in *X. cissicola* and *X. oryzae*. Nevertheless, in species
such as *X. campestris*, integrons remain robustly functional and continue to contribute to
genomic diversification, as evidenced by the presence of full length *intI*, high cassette
variability, and the of strain specific gene cassettes in cassette arrays. Selective pressures
and ecological niches may favor the retention of an active integron system in this species
and not in the others, for reasons not clear at this time.

The phenotypic impact of the large pool of gene cassettes residing in *Xanthomonas*
genomes remains largely unresolved. Nearly half of orthologous genes carried by gene
cassettes occurred only once, contributing to strain level genetic variability. These rare
genes may represent a reservoir of adaptive potential, but could also be selectively neutral
or even transient. Cassettes in *Vibrio* chromosomal integron platforms have been suggested
to have been selected to be as neutral as possible (Blanco *et al.*, 2024). The frequent
occurrence of hypothetical proteins, including highly conserved cassettes ORFs, suggests
important functional roles that remain undiscovered. For instance, a cassette carrying the
DUF1488 encoding gene is present, a few times in multiple copies, in 16.7% of the genomes
across 6 species, sometimes with multiple copies within the same integron array. Among
cassette-encoded genes with predicted putative functions, many have predicted functions
linked to environmental interaction and thus have the potential to contribute to niche
adaptation. For example, the most widespread cassette in our dataset encodes *symE* and
the small coding sRNA-Xcc1, known to be activated by T3SS regulatory proteins (Chen *et al.*,
2011), which may influence bacterial growth during plant colonization. In *Xanthomonas*, we
found a fairly small fraction of cassette genes to be classified as involved defence
mechanisms, despite the emerging role of integron platforms in encoding anti-phage genes
(Darracq *et al.*, 2025, Getz *et al.*, 2025, Kieffer *et al.*, 2025). However, this may be an
underestimate, as work on large sedentary chromosomal integrons in *Vibrio cholerae*
showed that 20% of their cassettes with no predicted function encode for anti-phage
defence systems, despite these genes not being detected as such by bioinformatic tools
(Darracq *et al.*, 2025).

Importantly, our analysis revealed evidence of interspecies horizontal transfer of cassettes.
For both *X. campestris* and *X. arboricola* there is evidence of recent horizontal transfer of
integron-associated cassettes with other *Xanthomonas* species. Interestingly, however,
despite their high levels of cassette acquisition, there is little evidence of recent and direct
cassette exchange between these two species. Instead, *X. campestris* shows the highest
degree of gene sharing with *X. hortorum*, a species more closely related to *X. arboricola*
than to *X. campestris*. Conversely, the species with which *X. arboricola* shares the greatest
number of cassettes is *X. euroxanthea*, its closest phylogenetic neighbour among the

genomes analysed. These patterns suggest that both phylogenetic proximity and ecological
overlap, such as co-occurrence in the phyllosphere or endosphere during mixed infections,
have contributed to the dynamics of cassette exchange. The shared habitat provides
repeated opportunities for interspecies contact and genetic exchange, likely promoting the
horizontal dissemination of integron cassettes across the genus.

In summary, our analysis indicates that integron platforms were likely acquired early in the
evolution of the *Xanthomonas* genus and have since followed divergent evolutionary
trajectories. While many lineages show signs of integron inactivation, others, such as *X.*
campestris, retain active systems contributing to genomic diversification. The gene cassettes
carried by these integrons, including many with unknown or potentially adaptive functions,
add to strain-level variability and may influence niche adaptation. Patterns of horizontal
gene transfer suggest that both phylogenetic relatedness and ecological overlap shape
cassette exchange across species, highlighting integrons as dynamic elements in the
evolutionary landscape of *Xanthomonas*.

576 *Funding information:* This work is supported by funding from the ARC Centre of Excellence in Synthetic
577 Biology (CE200100029).
578 *Conflicts of interest:* The authors declare that there is no conflict of interest.

REFERENCES

Almagro Armenteros JJ, Tsirigos KD, Sønderby CK, Petersen TN, Winther O, Brunak S, von
Heijne G & Nielsen H (2019) SignalP 5.0 improves signal peptide predictions using deep
neural networks. *Nature biotechnology* **37**: 420-423.

Ashrafi S, Kuzmanović N, Patz S, Lohwasser U, Bunk B, Spröer C, Lorenz M, Elhady A, Frühling
A & Neumann-Schaal M (2022) Two new Rhizobiales species isolated from root nodules of
common sainfoin (*Onobrychis viciifolia*) show different plant colonization strategies.
Microbiology Spectrum **10**: e01099-01022.

Barionovi D & Scortichini M (2006) Assessment of integron gene cassette arrays in strains of
Xanthomonas fragariae and *X. arboricola* pvs. *fragariae* and *pruni*. *Journal of Plant*
Pathology 279-284.

Barionovi D & Scortichini M (2008) Integron variability in *Xanthomonas arboricola* pv.
juglandis and *Xanthomonas arboricola* pv. *pruni* strains. *FEMS microbiology letters* **288**: 19-
24.

Bertels F, Silander OK, Pachkov M, Rainey PB & Van Nimwegen E (2014) Automated
reconstruction of whole-genome phylogenies from short-sequence reads. *Molecular biology*
and evolution **31**: 1077-1088.

Biessy A, Cadieux M, Ciotola M, McDuff F, Soufiane B, Laforest M & Filion M (2025)
Characterization of a plant-pathogenic T3SS-lacking *Xanthomonas* strain isolated from
common ragweed. *Plant Pathology* **74**: 308-319.

Blanco P, Hipólito A, García-Pastor L, Trigo da Roza F, Toribio-Celestino L, Ortega AC, Vergara
E, San Millán Á & Escudero JA (2024) Identification of promoter activity in gene-less
cassettes from *Vibrionaceae* superintegrons. *Nucleic acids research* **52**: 2961-2976.

Boucher Y, Labbate M, Koenig JE & Stokes H (2007) Integrons: mobilizable platforms that
promote genetic diversity in bacteria. *Trends in microbiology* **15**: 301-309.

Buchfink B, Xie C & Huson DH (2015) Fast and sensitive protein alignment using DIAMOND.
Nature methods **12**: 59-60.

Cambray G, Sanchez-Alberola N, Campoy S, Guerin É, Da Re S, González-Zorn B, Ploy M-C,
Barbé J, Mazel D & Erill I (2011) Prevalence of SOS-mediated control of integron integrase
expression as an adaptive trait of chromosomal and mobile integrons. *Mobile DNA* **2**: 6.

Cantalapiedra CP, Hernández-Plaza A, Letunic I, Bork P & Huerta-Cepas J (2021) eggNOG-
mapper v2: functional annotation, orthology assignments, and domain prediction at the
metagenomic scale. *Molecular biology and evolution* **38**: 5825-5829.

Chaumeil P-A, Mussig AJ, Hugenholtz P & Parks DH (2022) GTDB-Tk v2: memory friendly
classification with the genome taxonomy database. *Bioinformatics* **38**: 5315-5316.

Chen X-L, Tang D-J, Jiang R-P, He Y-Q, Jiang B-L, Lu G-T & Tang J-L (2011) *sRNA-Xcc1*, an
integron-encoded transposon-and plasmid-transferred trans-acting sRNA, is under the
positive control of the key virulence regulators HrpG and HrpX of *Xanthomonas campestris*
pathovar *campestris*. *RNA biology* **8**: 947-953.

Costa J, Pothier JF, Bosis E, Boch J, Kölliker R & Koebnik R (2024) A community-curated
DokuWiki resource on diagnostics, diversity, pathogenicity, and genetic control of
xanthomonads. *Molecular Plant-Microbe Interactions* **37**: 347-353.

Cury J, Jové T, Touchon M, Néron B & Rocha EP (2016) Identification and analysis of
integrons and cassette arrays in bacterial genomes. *Nucleic acids research* **44**: 4539-4550.

Darracq B, Littner E, Brunie M, Bos J, Kaminski PA, Depardieu F, Slesak W, Debatisse K,
Touchon M & Bernheim A (2025) Sedentary chromosomal integrons as biobanks of bacterial
antiphage defense systems. *Science* **388**: eads0768.

Domazet-Lošo M & Haubold B (2011) Alignment-free detection of local similarity among
viral and bacterial genomes. *Bioinformatics* **27**: 1466-1472.

Dunn MF & Becerra-Rivera VA (2023) The biosynthesis and functions of polyamines in the
interaction of plant growth-promoting rhizobacteria with plants. *Plants* **12**: 2671.

Escudero JA, Loot C & Mazel D (2018) Integrons as adaptive devices. *Molecular mechanisms*
of microbial evolution 199-239.

Essakhi S, Cesbron S, Fischer-Le Saux M, Bonneau S, Jacques M-A & Manceau C (2015)
Phylogenetic and variable-number tandem-repeat analyses identify nonpathogenic
Xanthomonas arboricola lineages lacking the canonical type III secretion system. *Applied*
and Environmental Microbiology **81**: 5395-5410.

Galperin MY, Vera Alvarez R, Karamycheva S, Makarova KS, Wolf YI, Landsman D & Koonin
EV (2025) COG database update 2024. *Nucleic Acids Research* **53**: D356-D363.

Garita-Cambronero J, Palacio-Bielsa A, López MM & Cubero J (2017) Pan-genomic analysis
permits differentiation of virulent and non-virulent strains of *Xanthomonas arboricola* that
cohabit *Prunus* spp. and elucidate bacterial virulence factors. *Frontiers in microbiology* **8**:
573.

Getz LJ, Fairburn SR, Vivian Liu Y, Qian AL & Maxwell KL (2025) Integrons are anti-phage
defence libraries in *Vibrio parahaemolyticus*. *Nature Microbiology* 1-10.

Ghaly TM, Tetu SG & Gillings MR (2021) Predicting the taxonomic and environmental
sources of integron gene cassettes using structural and sequence homology of attC sites.
Communications biology **4**: 946.

Ghaly TM, Geoghegan JL, Alroy J & Gillings MR (2019) High diversity and rapid spatial
turnover of integron gene cassettes in soil. *Environmental Microbiology* **21**: 1567-1574.

Ghaly TM, Geoghegan JL, Tetu SG & Gillings MR (2020) The peril and promise of integrons:
beyond antibiotic resistance. *Trends in microbiology* **28**: 455-464.

Ghaly TM, Gillings MR, Rajabal V, Paulsen IT & Tetu SG (2024) Horizontal gene transfer in
plant microbiomes: integrons as hotspots for cross-species gene exchange. *Frontiers in*
Microbiology **15**: 1338026.

Ghaly TM, Rajabal V, Russel D, Colombi E & Tetu SG (2025) EcoFoldDB: Protein structure-
guided functional profiling of ecologically relevant microbial traits at the metagenome scale.
bioRxiv 2025.2004. 2002.646905.

Ghaly TM, Tetu SG, Penesyan A, Qi Q, Rajabal V & Gillings MR (2022) Discovery of integrons
in Archaea: Platforms for cross-domain gene transfer. *Science Advances* **8**: eabq6376.

Ghaly TM, Penesyan A, Pritchard A, Qi Q, Rajabal V, Tetu SG & Gillings MR (2022) Methods
for the targeted sequencing and analysis of integrons and their gene cassettes from complex
microbial communities. *Microbial genomics* **8**: 000788.

Ghaly TM, Rajabal V, Penesyan A, Coleman NV, Paulsen IT, Gillings MR & Tetu SG (2023)
Functional enrichment of integrons: Facilitators of antimicrobial resistance and niche
adaptation. *Iscience* **26**.

Gillings M, Boucher Y, Labbate M, Holmes A, Krishnan S, Holley M & Stokes H (2008) The
evolution of class 1 integrons and the rise of antibiotic resistance. *Journal of bacteriology*
190: 5095-5100.

Gillings MR (2014) Integrons: past, present, and future. *Microbiology and molecular biology*
reviews **78**: 257-277.

Gillings MR, Holley MP, Stokes H & Holmes AJ (2005) Integrons in *Xanthomonas*: a source of
species genome diversity. *Proceedings of the National Academy of Sciences* **102**: 4419-4424.

Guerin É, Cambray G, Sanchez-Alberola N, Campoy S, Erill I, Da Re S, Gonzalez-Zorn B, Barbé
J, Ploy M-C & Mazel D (2009) The SOS response controls integron recombination. *Science*
324: 1034-1034.

Hall RM & Collis CM (1995) Mobile gene cassettes and integrons: capture and spread of
genes by site-specific recombination. *Molecular microbiology* **15**: 593-600.

Heinzinger M, Weissenow K, Sanchez JG, Henkel A, Mirdita M, Steinegger M & Rost B (2024)
Bilingual language model for protein sequence and structure. *NAR Genomics and*
Bioinformatics **6**: lqae150.

Hipólito A, García-Pastor L, Vergara E, Jové T & Escudero JA (2023) Profile and resistance
levels of 136 integron resistance genes. *npj Antimicrobials and Resistance* **1**: 13.

Huerta-Cepas J, Szklarczyk D, Heller D, Hernández-Plaza A, Forslund SK, Cook H, Mende DR,
Letunic I, Rattei T & Jensen LJ (2019) eggNOG 5.0: a hierarchical, functionally and
phylogenetically annotated orthology resource based on 5090 organisms and 2502 viruses.
Nucleic acids research **47**: D309-D314.

Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT & Aluru S (2018) High throughput
ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nature*
communications **9**: 5114.

Kawano M, Aravind á & Storz G (2007) An antisense RNA controls synthesis of an
SOS-induced toxin evolved from an antitoxin. *Molecular microbiology* **64**: 738-754.

Kieffer N, Hipólito A, Ortiz-Miravalles L, Blanco P, Delobelle T, Vizuete P, Ojeda FM, Jové T,
Jurenas D & García-Quintanilla M (2025) Mobile integrons encode phage defense systems.
Science **388**: eads0915.

Klemm P, Stadler PF & Lechner M (2023) Proteinortho6: pseudo-reciprocal best alignment
heuristic for graph-based detection of (co-) orthologs. *Frontiers in bioinformatics* **3**:
1322477.

Langmead B & Salzberg SL (2012) Fast gapped-read alignment with Bowtie 2. *Nature*
methods **9**: 357-359.

Lemoine F & Gascuel O (2021) Gotree/Goalign: toolkit and Go API to facilitate the
development of phylogenetic workflows. *NAR Genomics and Bioinformatics* **3**: lqab075.

Loeff L, Walter A, Rosalen GT & Jinek M (2025) DNA end sensing and cleavage by the Shedu
anti-phage defense system. *Cell* **188**: 721-733. e717.

Loot C, Millot GA, Richard E, Littner E, Vit C, Lemoine F, Néron B, Cury J, Darracq B & Niault T
(2024) Integron cassettes integrate into bacterial genomes via widespread non-classical attG
sites. *Nature Microbiology* **9**: 228-240.

Löytynoja A (2013) Phylogeny-aware alignment with PRANK. *Multiple sequence alignment*
*methods,*p.^pp. 155-170. Springer.

Mazel D (2006) Integrons: agents of bacterial evolution. *Nature Reviews Microbiology* **4**:
608-620.

Mazel D, Dychinco B, Webb VA & Davies J (1998) A distinctive class of integron in the *Vibrio*
cholerae genome. *Science* **280**: 605-608.

Michael CA & Labbate M (2010) Gene cassette transcription in a large integron-associated
array. *BMC genetics* **11**: 1-13.

Néron B, Littner E, Haudiquet M, Perrin A, Cury J & Rocha EP (2022) IntegronFinder 2.0:
identification and analysis of integrons across bacteria, with a focus on antibiotic resistance
in Klebsiella. *Microorganisms* **10**: 700.

Paradis E & Schliep K (2019) ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics* **35**: 526-528.

Parks DH, Chuvochina M, Rinke C, Mussig AJ, Chaumeil P-A & Hugenholtz P (2022) GTDB: an
ongoing census of bacterial and archaeal diversity through a phylogenetically consistent,
rank normalized and complete genome-based taxonomy. *Nucleic acids research* **50**: D785-
D794.

Patané JS, Martins J, Rangel LT, Belasque J, Digiampietri LA, Facincani AP, Ferreira RM,
Jaciani FJ, Zhang Y & Varani AM (2019) Origin and diversification of *Xanthomonas citri* subsp.
citri pathotypes revealed by inclusive phylogenomic, dating, and biogeographic analyses.
BMC genomics **20**: 1-23.

Pena MM, Bhandari R, Bowers RM, Weis K, Newberry E, Wagner N, Pupko T, Jones JB,
Woyke T & Vinatzer BA (2024) Genetic and functional diversity help explain pathogenic,
weakly pathogenic, and commensal lifestyles in the genus *Xanthomonas*. *Genome Biology*
and Evolution **16**: evae074.

Pereira BM, Österlund T, Eriksson KM, Backhaus T, Axelson-Fisk M & Kristiansson E (2020) A
comprehensive survey of integron-associated genes present in metagenomes. *BMC*
genomics **21**: 1-14.

Post V & Hall RM (2009) Insertion sequences in the IS 1111 family that target the attC
recombination sites of integron-associated gene cassettes. *FEMS microbiology letters* **290**:
182-187.

Price MN, Dehal PS & Arkin AP (2010) FastTree 2–approximately maximum-likelihood trees
for large alignments. *PloS one* **5**: e9490.

Qi Q, Ghaly TM, Rajabal V, Russell DH, Gillings MR & Tetu SG (2024) Vegetable phylloplane
microbiomes harbour class 1 integrons in novel bacterial hosts and drive the spread of
chlorite resistance. *Science of the Total Environment* **954**: 176348.

Raga-Carbajal E, Espin G, Ayala M, Rodríguez-Salazar J & Pardo-López L (2022) Evaluation of
a bacterial group 1 LEA protein as an enzyme protectant from stress-induced inactivation.
Applied Microbiology and Biotechnology **106**: 5551-5562.

Rajabal V, Ghaly TM, Egidi E, Ke M, Penesyan A, Qi Q, Gillings MR & Tetu SG (2024) Exploring
the role of mobile genetic elements in shaping plant–bacterial interactions for sustainable
agriculture and ecosystem health. *Plants, People, Planet* **6**: 408-420.

Rao MJ, Zuo H & Xu Q (2021) Genomic insights into citrus domestication and its important
agronomic traits. *Plant Communications* **2**.

Rowe-Magnus DA & Mazel D (2001) Integrons: natural tools for bacterial genome evolution.
Current opinion in microbiology **4**: 565-569.

Rowe-Magnus DA, Guerout A-M, Biskri L, Bouige P & Mazel D (2003) Comparative analysis of
superintegrons: engineering extensive genetic diversity in the Vibrionaceae. *Genome*
research **13**: 428-442.

Sapkota S, Mergoum M & Liu Z (2020) The translucens group of *Xanthomonas translucens*:
complicated and important pathogens causing bacterial leaf streak on cereals. *Molecular*
plant pathology **21**: 291-302.

Schepler-Luu V, Sciallano C, Stiebner M, Ji C, Boulard G, Diallo A, Auguy F, Char SN, Arra Y &
Schenstnyi K (2023) Genome editing of an African elite rice variety confers resistance against
endemic and emerging *Xanthomonas oryzae* pv. *oryzae* strains. *Elife* **12**: e84864.

Schwengers O, Jelonek L, Dieckmann MA, Beyvers S, Blom J & Goesmann A (2021) Bakta:
rapid and standardized annotation of bacterial genomes via alignment-free sequence
identification. *Microbial genomics* **7**: 000685.

Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B &
Ideker T (2003) Cytoscape: a software environment for integrated models of biomolecular
interaction networks. *Genome research* **13**: 2498-2504.

Stokes Ht & Hall R (1989) A novel family of potentially mobile DNA elements encoding
site-specific gene-integration functions: integrons. *Molecular microbiology* **3**: 1669-1683.

Strugeon E, Tilloy V, Ploy M-C & Da Re S (2016) The stringent response promotes antibiotic
resistance dissemination by regulating integron integrase expression in biofilms. *MBio* **7**:
10.1128/mbio. 00868-00816.

Tansirichaiya S, Mullany P & Roberts AP (2019) Promoter activity of ORF-less gene cassettes
isolated from the oral metagenome. *Scientific Reports* **9**: 8388.

Tesson F, Hervé A, Mordret E, Touchon M, d'Humières C, Cury J & Bernheim A (2022)
Systematic and quantitative view of the antiviral arsenal of prokaryotes. *Nature*
communications **13**: 2561.

Tetu SG & Holmes AJ (2008) A family of insertion sequences that impacts integrons by
specific targeting of gene cassette recombination sites, the IS 1111-attC group. *Journal of*
bacteriology **190**: 4959-4970.

Teufel F, Almagro Armenteros JJ, Johansen AR, Gíslason MH, Pihl SI, Tsirigos KD, Winther O,
Brunak S, von Heijne G & Nielsen H (2022) SignalP 6.0 predicts all five types of signal
peptides using protein language models. *Nature biotechnology* **40**: 1023-1025.

Thompson MK, Nocedal I, Culviner PH, Zhang T, Gozzi KR & Laub MT (2022) *Escherichia coli*
SymE is a DNA-binding protein that can condense the nucleoid. *Molecular microbiology* **117**:
851-870.

Timilsina S, Potnis N, Newberry EA, Liyanapathiranage P, Iruegas-Bocardo F, White FF, Goss
EM & Jones JB (2020) *Xanthomonas* diversity, virulence and plant–pathogen interactions.
Nature Reviews Microbiology **18**: 415-427.

Triplett LR, Hamilton J, Buell C, Tisserat N, Verdier V, Zink F & Leach J (2011) Genomic
analysis of *Xanthomonas oryzae i*solates from rice grown in the United States reveals
substantial divergence from known X. oryzae pathovars. *Applied and Environmental*
Microbiology **77**: 3930-3937.

Van Kempen M, Kim SS, Tumescheit C, Mirdita M, Lee J, Gilchrist CL, Söding J & Steinegger
M (2024) Fast and accurate protein structure search with Foldseek. *Nature biotechnology*
42: 243-246.

Vauterin L, Yang P, Alvarez A, Takikawa Y, Roth DA, Vidaver AK, Stall RE, Kersters K & Swings
J (1996) Identification of non-pathogenic *Xanthomonas* strains associated with plants.
Systematic and applied microbiology **19**: 96-105.

Xu S, Dai Z, Guo P, Fu X, Liu S, Zhou L, Tang W, Feng T, Chen M & Zhan L (2021) ggtreeExtra:
compact visualization of richly annotated phylogenetic data. *Molecular biology and*
evolution **38**: 4039-4042.

Yu G, Smith DK, Zhu H, Guan Y & Lam TTY (2017) ggtree: an R package for visualization and
annotation of phylogenetic trees with their covariates and other associated data. *Methods*
in Ecology and Evolution **8**: 28-36.

Zhang Z, Todeschini TC, Wu Y, Kogay R, Naji A, Rodriguez JC, Mondi R, Kaganovich D, Taylor
DW & Bravo JP (2023) Kiwa is a bacterial membrane-embedded defence supercomplex
activated by phage-induced membrane changes. *bioRxiv* 2023.2002. 2026.530102.

FIGURE AND TABLE LEGENDS

Figure 1 – Distribution of integron platforms in *Xanthomonas* groups 1 and 2. The
maximum likelihood phylogenetic tree is based on a concatenated alignment of 120 top-
ranked marker proteins. The tree was rooted at the midpoint. Tips are labelled with species
names assigned with Genome Taxonomy Database phylogeny. Integrons were classified as
present if the isolates carried a functional integron, *intI* only or an *attC* array only. Scale bar
indicates substitutions per site.

Figure 2 – Phylogeny of *intI*. Sequences of *intI* longer than 900 bp were aligned with PRANK,
stripped of gaps and used to infer the tree with FastTree. *IntI* from *Vibrio* sp. SCSIO 43136
(locus tag: J4N39_08275) was used as an outgroup. The scale bar indicates substitutions per
site. Only bootstrap values 0.80 are shown. Group 1 *Xanthomonas* are indicated on the side
of the plot.

Figure 3 – Integrons in *X. campestris*. The phylogeny of *X. campestris* was built using
Realphy, using GCA_000007145.1 as a reference genome. The tree was rooted with
GCA_000972745 (*X. arboricola*) then the tip was removed from the tree. Tip labels show
pathovars. The scale bar indicates substitutions per site. A) Indicates the presence of a
complete integron or *attC* sites in the *ilvD* locus, B) indicates the presence of *intI*, C)
indicates the presence of a CALINs (cluster of *attC* sites) not in the ilvD locus. D) indicates
whether IntI is predicted to be functional (full length). E) indicates the number of *attC* sites
(number of cassettes) in the genome. Panel F) represents the distribution of orthologous
genes among all cassettes carried by the corresponding isolate inferred with Proteinortho.
Orthologous genes are listed in decreasing order based on the number of strains within the
species that carry them.

Figure 4 – Integrons in *X. cissicola*. The phylogeny of *X. cissicola* was built using Realphy,
using GCA_000007165.1 as a reference genome. The tree was rooted with
GCA_000009165.1 (*X. euvesicatoria*) then the tip was removed from the tree. Tip labels
show pathovars, which were assigned from the literature search. The scale bar indicates
substitutions per site. A) Indicates the presence of a complete integron or *attC* sites in the
ilvD locus, B) indicates the presence of *intI*, C) indicates the presence of a CALINs (cluster of
attC sites) not in the ilvD locus. D) indicates the number of *attC* sites (number of cassettes)

in the genome. Panel E) represents the distribution of orthologous genes among the all the
cassettes carried by the corresponding isolate inferred with Proteinortho. Orthologous
genes are given in decreasing order based on the number of strains within the species that
carry them.

Figure 5 – Horizontal transfer of cassettes among *Xanthomonas* species. Cytoscape
network illustrating transfer of cassettes between species. Nodes represent *Xanthomonas*
species, species containing more than 10 genomes in the dataset are coloured as in Figure 1
(tip labels). Line width between species is proportional to the number of cassettes moved
between species.

Figure S1 – Phylogeny of *Xanthomonas intI*. A maximum likelihood phylogenetic tree of all
intI in *Xanthomonas*. IntI from *Vibrio* sp. SCSIO 43136 (locus tag: J4N39_08275) was used as
an outgroup. The tree was inferred with FastTree and rooted with *Vibrio* sp. SCSIO 43136,
the scale bar indicates substitutions per site. * indicates the *intI*s are not integrated adjacent
to *ilvD*.

Figure S2 – Phylogeny of *ilvD*. The maximum-likelihood phylogenetic tree of all *ilvD* in
Xanthomonas. *ilvD* from *Vibrio* sp. SCSIO 43136 was used as an outgroup. The tree was
inferred with FastTree and rooted at midpoint, the scale bar indicates substitutions per site.

Figure S3 – Integrons in *X. arboricola*. The phylogeny of *X. arboricola* was built using
Realphy, using GCA_000972745.1 as a reference genome. The tree was rooted with
GCA_001908755.1 (*X. hortorum*) then the tip was removed from the tree. Tip labels show
pathovars, which were assigned via literature search. The red triangle indicates the probable
acquisition of the *intI* variant. The scale bar indicates substitutions per site. A) Indicates the
presence of a complete integron or *attC* sites in the *ilvD* locus, B) indicates the presence of
intI, C) indicates the presence of a CALINs (cluster of *attC* sites) not in the ilvD locus. D)
indicates whether IntI is predicted to be functional (full length). E) indicates the number of
attC sites (number of cassettes) in the genome. Panel F) represents the distribution of
orthologous genes among the all cassettes carried by the corresponding isolate inferred
with Proteinortho. Orthologous genes are given in decreasing order based on the number of
strains within the species that carry them.

Figure S4 – Integrons in *X. translucens*. The phylogeny of *X. translucens* was built using
Realphy, using GCA_001021935.1 as reference genome. The tree was rooted with
GCA_000007145.1 (*X. campestris*) then the tip removed from the tree. Tip labels show
pathovars, which were assigned via literature search. The scale bar indicates substitutions
per site. A) Indicates the presence of a complete integron or *attC* sites in the *ilvD* locus, B)
inidicates the presence of *intI*, C) indicate the presence of a CALINs (cluster of *attC* sites) not
in the ilvD locus. D) indicates whether IntI is predicted to be functional (full length). E)
indicates the number of *attC* sites (number of cassettes) in the genome. Panel F) represents
the distribution of orthologous genes among the all the cassettes carried by the
corresponding isolate interfered with Proteinortho. Orthologous genes are ordered in
decreasing order based on the number of strains within the species that carry them.

Figure S5 – Integrons in *X. euvesicatoria*. The phylogeny of *X. euvesicatoria* was built using
Realphy, using GCA_000009165.1 as reference genome. The tree was rooted with
GCA_002759095.2 (*X. phaseoli*) then the tip removed from the tree. The scale bar indicates
substitutions per site. A) Indicates the presence of a complete integron or *attC* sites in the
ilvD locus, B) inidicates the presence of *intI*, C) indicate the presence of a CALINs (cluster of
attC sites) not in the ilvD locus. D) indicates whether IntI is predicted to be functional (full
length). E) indicates the number of *attC* sites (number of cassettes) in the genome. Panel F)
represents the distribution of orthologous genes among the all the cassettes carried by the
corresponding isolate interfered with Proteinortho. Orthologous genes are ordered in
decreasing order based on the number of strains within the species that carry them.

Figure S6 – Integrons in *X. oryzae*. The phylogeny of *X. oryzae* was built using Realphy, using
GCA_000007385.1 as reference genome. The tree was rooted with GCA_002759095.2 (*X.*
phaseoli) then the tip removed from the tree. Tip labels show pathovars, which were
assigned via literature search. The scale bar indicates substitutions per site. A) Indicates the
presence of a complete integron or *attC* sites in the *ilvD* locus, B) inidicates the presence of
intI, C) indicate the presence of a CALINs (cluster of *attC* sites) not in the ilvD locus. D)
indicates whether IntI is predicted to be functional (full length). E) indicates the number of
attC sites (number of cassettes) in the genome. Panel F) represents the distribution of
orthologous genes among the all the cassettes carried by the corresponding isolate
interfered with Proteinortho. Orthologous genes are ordered in decreasing order based on
the number of strains within the species that carry them.

Figure S7 – Integrons in *X. phaseoli*. The phylogeny of *X. phaseoli* was built using Realphy,
using GCA_008639345.1 as reference genome. The tree was rooted with GCA_000009165.1
(*X. euvesicatoria*) then the tip removed from the tree. Tip labels show pathovars, which
were assigned via literature search. The scale bar indicates substitutions per site. A)
Indicates the presence of a complete integron or *attC* sites in the *ilvD* locus, B) inidicates the
presence of *intI*, C) indicate the presence of a CALINs (cluster of *attC* sites) not in the ilvD
locus. D) indicates whether IntI is predicted to be functional (full length). E) indicates the
number of *attC* sites (number of cassettes) in the genome. Panel F) represents the
distribution of orthologous genes among the all the cassettes carried by the corresponding
isolate interfered with Proteinortho. Orthologous genes are ordered in decreasing order
based on the number of strains within the species that carry them.

Figure S8 – Integrons in *X. hortorum*. The phylogeny of *X. hortorum* was built using Realphy,
using GCA_002285515.1 as reference genome. The tree was rooted with GCA_905367715.1
(*X. arboricola*) then the tip removed from the tree. Tip labels show pathovars, which were
assigned via literature search. The scale bar indicates substitutions per site. A) Indicates the
presence of a complete integron or *attC* sites in the *ilvD* locus, B) indicates the presence of
intI, C) indicate the presence of a CALINs (cluster of *attC* sites) not in the *ilvD* locus. D)
indicates whether IntI is predicted to be functional (full length). E) indicates the number of
attC sites (number of cassettes) in the genome. Panel F) represents the distribution of
orthologous genes among the all the cassettes carried by the corresponding isolate
interfered with Proteinortho. Orthologous genes are ordered in decreasing order based on
the number of strains within the species that carry them.

Figure S9 – Integron gene cassette counts in *Xanthomonas*. Boxplot of the number of *attC*
sites per genome grouped by plant of isolation and by species (only species with more than
10 genomes are shown). Letters indicate statistically different groups (Dunn's test with
Bonferroni adjustment, p<0.001).

Table S1 – Integrons and number of cassettes in the *Xanthomonas* genomes used in this
study

929