Title:
Multimodal single-cell analyses reveal distinct fusion-regulated transcriptional programs in Ewing sarcoma.

Authors:
Olivia G. Waltner[1], [†]April A. Apfelbaum[2], [†]Emma D. Wrenn[2], Shruti S. Bhise[1], Sami B. Kanaan[1], Rula Green Gladden[1], Mark A. Mendoza[1], Roger Volden[3], Zev Kronenberg[3], Anand Patel[4], Michael Dyer[4], Jay F. Sarthy[2,5], [††]Elizabeth R. Lawlor[2,5*], and [††]Scott N. Furlan[1,5*]

Affiliations:
[1]Translational Science and Therapeutics Division, Fred Hutch Cancer Research Center, Seattle, Washington.
[2]Ben Towne Center for Childhood Cancer and Blood Disorders Research, Seattle Children's Research Institute, Seattle, Washington.
[3]Pacific Biosciences, Menlo Park, California.
[4]St. Jude Children's Research Hospital, Memphis, Tennessee,
[5]Department of Pediatrics, University of Washington, Seattle, Washington.

Author List Footnotes:
[†]These authors contributed equally
[††]These authors contributed equally

*Correspondence:
Elizabeth R. Lawlor, MD PhD
Ben Towne Center for Childhood Cancer Research
Seattle Children's Research Institute
1920 Terry Ave, Seattle, Washington, 98101 USA
Email: beth.lawlor@seattlechildrens.org

Scott N. Furlan, MD (Lead)
Fred Hutchinson Cancer Research Center
1100 Fairview Ave N, Seattle, Washington, 98109 USA
Email: sfurlan@fredhutch.org

Abstract

Ewing sarcoma (EwS) is a fusion-driven malignancy, peaking in adolescence. Although EwS tumors are driven uniquely by EWS::FLI1 and related fusions, patient outcomes vary greatly. If and how tumor plasticity of EWS::FLI1-regulated transcriptional signatures contribute to disease progression is not known. We utilized a single-cell co-assay of RNA and ATAC sequencing to identify gene regulatory networks in EwS. By comprehensively characterizing regulatory elements across cell lines, we identified multiple unique modules of gene regulation. Differential usage and prevalence of these modules was evident across cell lines, associated with distinct epigenetic and transcriptomic signatures, and in specific cases, modifiable by exogenous TGF-β. Evaluating these modules in transcriptomic data from patients confirmed their enrichment across and within EwS primary tumors. We conclude that multiple gene regulatory networks co-exist but are differentially abundant within EwS tumors providing insight into how heterogenous transcriptional networks may contribute to clinical outcomes.

Introduction

Ewing sarcoma (EwS) is an aggressive, fusion-driven bone and soft tissue tumor encountered most in adolescents and young adults. Over a third of patients, including nearly all with metastatic or recurrent disease, do not survive[2,3]. Clinical presentation of EwS is heterogeneous, with 80% of localized cases presenting in the bone, and 20% arising in extraosseous locations. Histologically, EwS tumors are highly undifferentiated, exhibiting phenotypic features of both mesenchymal and neural crest progenitors. Although the cell of origin is controversial, EwS tumors are presumed to be derived from these lineages[4-7]. Chromosomal translocations that create fusions between the N-terminal domain of *EWSR1* and the DNA-binding domain of ETS transcription factor (TF) family members are the primary oncogenic lesion in EwS[2]. The resulting EWS::ETS chimeric oncoprotein reprograms gene regulatory networks through diverse mechanisms, including modifying chromatin state through its function as a pioneer factor at GGAA microsatellite repeats, creating *de novo* enhancers[8,9]. Apart from their fusions, EwS tumors are relatively bereft of mutations[2,10]. Neither additional genetic alterations, nor differences in fusion breakpoint and binding partner heterogeneity[11] adequately explain the diverse clinical outcomes that are observed in patients who receive identical standard-of-care therapies. Advances in transcriptomic and epigenomic analyses have in recent years demonstrated the profound contributions of tumor cell plasticity and cell state heterogeneity to tumor metastasis, relapse, and therapy resistance[12,13]. EwS tumors demonstrate marked inter- and intra-tumoral heterogeneity in their epigenetic states[14,15] and, as such, these differences may be key determinants of disease progression and clinical outcome.

Recent studies by our group and others have used single cell genomics to study cellular heterogeneity within EwS tumors[15-24]. These studies reveal that EwS tumors coopt distinct developmental programs to promote proliferation or migratory behavior[16,17,19,21,22,25,26]. In addition, accruing evidence shows that cooperativity between TFs, EWS::FLI1, and diverse chromatin states may underlie these various oncogenic programs[21,27-35]. We hypothesized that joint single-cell transcriptional and chromatin profiling technology would reveal new insight into the cell-intrinsic epigenetic and transcriptional heterogeneity of EwS cell states. Using this approach, we have identified a set of diverse gene regulatory networks that are differentially influenced by the EWS::FLI1 fusion and TGF-β in the tumor microenvironment. These distinctive networks vary both across and within tumors from EwS patients providing evidence that a diversity of gene regulatory programs may contribute to the heterogeneity of this disease.

Fig.1. Multiomic analyses reveal distinct signatures between non-EwS and EwS cells. (A) UMAP embedding of single cell RNA sequencing and ATAC sequencing of EwS and non EwS cells. (B) chromVAR deviation scores of EWS::FLI motif enrichment across EwS and non-EwS cells (C) ChromVAR deviation scores of EWS::FLI binding site enrichment, separated by regions with or without GGAA microsatellites (μSAT; GGAA repeat ≥ 4), grouped by cell line. (D) Fraction of cells in each cell line with detection of EWS::FLI fusion transcript. (E) Differentially expressed genes (FDR ≤ 0.01 & Log2FC ≥ 1.25) across EwS/non-EwS landscape. (F) Differentially accessible peaks (FDR ≤ 0.01 & Log2FC ≥ 1.25) of EwS and non EwS cells with row dendrogram showing phenotypic clusters. Row dendrogram shows EwS cell clusters highlighted in bold and red.

Results

Multiomic analyses reveal distinct chromatin accessibility states across EwS cell lines

Cell lines are one of few available models used to interrogate EwS biology given the rarity of the disease and the lack of appropriately faithful animal models[36]. One drawback of cell line use is potential loss of intra-tumoral heterogeneity compared to the original tumor due to serial passaging. We reasoned that profiling an array of EwS cell lines would capture the diversity of gene regulatory networks that may underlie the clinical and transcriptional heterogeneity of this disease. Because of the well-described direct influence of the EwS fusion on chromatin remodeling [35], we utilized an assay capable of pairing single-cell RNA transcript counts with a chromatin assay for transposase-accessible chromatin (ATAC) (herein referred to as 'multiomic' sequencing) on 9 EwS and 5 non-EwS cell lines (Fig.1A, Fig.S1A). We recovered 30,105 profiles after filtering for poor quality cells (Fig.1A, Fig.S1A, Fig.S1B). Multiomic single-cell profiles self-assembled after dimensionality reduction according to cell line identity (Fig.1A).

First, to identify the extent of transcriptional heterogeneity of validated direct EWS::FLI1 target genes[20] (EwS TGs) in individual cells, we calculated expression of EwS TGs at the single-cell level, subtracted by the aggregated expression of control gene sets[37]. As expected, RNA expression of these EwS TGs was higher in EwS compared to non-EwS cell lines (Fig. S1C). As these published EwS TGs were experimentally derived from the A673 cell line[20], it was unsurprising that A673 exhibited the highest average expression of these genes. To interrogate EwS TGs using chromatin-accessibility data, we

generated gene activity scores[38] which model accessibility of regulatory elements in the vicinity of a given gene. These scores confirmed that chromatin accessibility of EwS TGs was higher in EwS cell lines compared to non-EwS cells (Fig. S1D). To compare EWS::FLI1 motif enrichment across cells, we calculated single-cell ChromVAR deviation scores[38] of GGAA-microsatellites (four or more consecutive GGAA repeats; abbrev. 'GGAA- μSat') in accessible regions. As anticipated, this analysis revealed low enrichment in non-EwS cell lines. Unexpectedly, stark differences in EWS::FLI1 motif enrichment were observed across EwS cell lines (Fig. 1B). In some cell lines accessibility of GGAA- μSats was similar across single cells (e.g. TC71, RDES) while others showed a broader range of accessibility (e.g. SKNMC, TC32, PDX305). CHLA10 cells showed a clear bimodal distribution of EWS::FLI1 motif accessibility (Fig. 1B). This hints that within these cell lines, heterogeneity exists among individual cells with respect to accessibility of GGAA- μSats. Next, we collated a list of publicly available EWS::FLI1 fusion binding sites[39] with or with a GGAA μSat present. We visualized the enrichment of EWS::FLI1 sites across all cell lines and discovered distinct EwS cell-line specific usage (Fig. 1C & Fig. S1D).

Given the differences in EWS::FLI1 motif and binding site enrichment across EwS cell lines, we hypothesized that EwS cell lines have different levels of EWS::FLI1 transcript. It has been reported that lower levels of EWS::FLI1 gene expression and activity induce mesenchymal phenotypes in EwS cell lines[40]. To address this, we designed an approach to enrich EWS::FLI1 fusion transcript from the RNA components of our multiomic capture using biotinylated hybridization probes that tiled the EWSR1 and FLI1 gene loci. We then performed long-read sequencing on the product and counted fusion reads using novel fusion detection software designed specifically for long-reads (see Methods for details). As expected, EWS::FLI1 fusion transcripts were detected in EWS cells (Fig.1D). Scant non-ES cells showed low levels of the fusion, likely from PCR artifacts of low-complexity products. Overall, however, we observed that EwS cell lines exhibit a high degree of variability in detectible EWS::FLI1 fusion transcript (Fig. 1D & Fig. S1E). Generally, the proportion of cells with detectible transcript correlated with values obtained using single-cell qPCR to assess fusion expression[16]. These findings are generally congruent with motif enrichment in that high fusion-expressing cell lines (e.g. A4573, SKNMC, and RDES) exhibit high EWS::FLI1 motif enrichment while cell lines bearing lower levels of fusion (e.g. PDX305 and TC32) exhibit lower EWS::FLI1 chromatin signature (Fig.1B & Fig. 1D). By contrast, other cell lines displayed inverse trends of fusion transcript to chromatin enrichment (e.g. A673 and TC71).

To profile the global chromatin and transcriptional heterogeneity using an unbiased approach, we calculated top differentially expressed (DE) genes and differentially accessible (DA) regions across EwS and non EwS cells. As with the EwS TG-focused analysis, this approach showed that top DE/DA features readily distinguished EwS cells from non-EwS cells (Fig. 1E-F). We observed self-aggregation of EwS cells into distinct clusters. Specifically, in terms of top DA regions, CHLA9, CHLA10, TC32, and PDX305 cells were more similar to each other than other EwS cell lines (Fig. 1F). Altogether, these data show that EwS cell lines exhibit heterogeneous gene-regulatory profiles that segregate into groups that display distinct trends in accessibility, potentially reflecting unique underlying gene regulatory networks that are active in discrete cell contexts.

EwS cells utilize gene regulatory programs that drive inter-tumoral heterogeneity

Next, to evaluate how cis-regulatory gene networks differ across EwS cells, we correlated transcript expression to regions of accessibility using the joint measurements of transcript count and accessibility in each cell (so-called "peak-to-gene linkages")[38]. Hierarchical clustering of these cis-regulatory networks with a k parameter equal to 3 clearly separated CHLA9, TC32, and PDX305 into one distinct network module and SKNMC, A4573, RDES, and TC71 into another (Fig. 2A), recapitulating what we observed using top DE/DA features. Cell subsets of CHLA10 separated into either cluster 2 or 3 (Fig. 2A) but more cells had cumulative expression of module 2 genes (Fig. 2B). We then assigned the 9 cell lines into three groups based on their dominant module signature (Fig. 2A-B), resulting in assignment of A673 cells as group 1, CHLA9, CHLA10, TC32, and PDX305 cells as group 2, and SKNMC, A4573, TC71, and RDES as group 3 (Fig. 2A). Enrichment analysis using Gene Ontology and Hallmark terms confirmed that the genes associated with these modules have functional associations with distinct cell behaviors (Fig. 2C). While there was some degree of overlap, module 2 showed a relative enrichment in GO terms related to mesenchymal identity, and modules 1 and 3 had enriched neuronal identity markers. Genes involved in oxidative phosphorylation and targets of mTOR were uniquely enriched in module 1, possibly due to the presence of a BRAF[V600E] mutation in A673 cells given the reported effects of this mutation on the mTOR pathway[41-43].

Next, we investigated if gene modules might harbor distinct TF binding sites. We performed differential motif enrichment using HOMER[44] and found a strikingly well-delineated, and module-specific enrichment of specific TF families across the cis-regulatory networks (Fig. 2D). The AP-1 family of motifs (FOS, FOSL2, JUN, AP1), which has been suggested to form ternary complexes with EWS::FLI1[45], enriched in modules 1 and 2, but not in module 3 (Fig. 2D). Zinc Finger (ZF) TFs (KLF4, KLF5, MAZ, SP1) were most enriched in modules 1 and 3 with limited evidence for activity in regulation of module 2 (Fig. 2D). Module 2 peaks uniquely exhibited motif enrichment for developmental regulators from the Homeobox family of TFs (HOXD13, HOXA9) and downstream transcriptional regulators of the Hippo and Wnt signal transduction pathways (TEAD1, TEAD4). To

Fig. 2 EwS cells utilize distinct gene regulatory modules

(A) Heatmap of unsupervised kmean clusters of peak-to-gene links (PGLs) across 9 EwS cell lines. Left-peaks, right-genes, top- colors denote cell lines. Right of heatmap-UMAP of cell lines labeled by subgroup. Group 1 labelled with navy/dashed lines, group 2 in orange and group 3 in light blue. (B) Violin plot of modules 1-3 gene expression and peak accessibility across EwS cell lines. (C) Universal enrichment analysis of module gene (top 300 filtered on peak-to-gene correlation) sets using both MSigDB Hallmark pathways and Gene Ontologies: Biological Process. The top 7 terms filtered by lowest p-value (those greater than 0.05 are not plotted) and count are shown. More significant terms are darker blue. The number of genes in each module that overlap with a gene set term is indicated by the count metric—represented as dot size. (D) HOMER de novo motif enrichment for each module peak set. Results depict the top significantly enriched motifs in each module peak set, and enrichments in other modules. (E) TFs with high expression and motif enrichment in entire single-cell data set of EwS cell lines. Statistically significant TFs correlation to gene expression > 0.5 and a maximum inter EwS subgroup difference in motif deviation z-score (denoted max TF motif delta) > 1.9 highlighted in red. (F) ChromVAR motif deviation scores and gene expression of highly enriched TFs, parsed by EwS subgroup. (G) Western blot of selected TFs across EwS cell lines. GAPDH = loading control.

corroborate these findings, we performed an analysis which correlates, at the single-cell level, TF binding motifs in accessible regions with gene expression (Fig. 2E). This showed that TEAD1, HOXD13, and AP-1 TFs were highly enriched in group 2 cells (Fig. 2E-F). Group 3 lacked motif enrichment for AP-1 TFs but showing robust enrichment in ALX1/4, RUNX3, CEBPB and EN2 TFs (Fig. 2E-F). We assessed protein levels of selected TF regulators by Western blots to evaluate their heterogeneity across the cell lines. The protein abundance of these TFs exhibited substantial variability among the cells tested (Fig. 2G). Together, these multimodal analyses of single cells uncovered at least two dominant gene regulatory circuits. One of these (module 2) displays mesenchymal/EMT-like functionality that is likely to be mediated by high activity of TEAD, AP-1, and HOXD13

TFs. The other (module 3) has enriched neural program functionality most likely mediated by ZF TFs and neural and neural crest development TFs such as RUNX3 and HAND2.

EwS gene regulatory modules are differentially regulated by EWS::FLI1

We next sought to address whether the identified gene regulatory modules are directly or indirectly modulated by EWS::FLI1. We first measured EWS::FLI1, EWS::ERG (the second most common fusion in EwS tumors), and wild-type FLI1 motif enrichment and found all to be enriched in all three modules (Fig. 3A) suggesting that EWS::FLI1 is likely to regulate expression of linked genes across all cells irrespective of the dominant module. However, by annotating these sites using published EWS::FLI1 binding sites (*39*) and segregating into those with or without a GGAA microsatellite (GGAA- µSat), we observed enrichment of GGAA-µSats only in module 3 peaks (Fig. 3B). Thus, this analysis suggests that EWS::FLI1 binding is likely to directly influence gene regulatory circuits and its impact on transcriptional state depends on GGAA patterning, as has previously been suggested[8].

To begin to address the impact of EWS::FLI1 activity on cis-regulatory modules, we sourced data from published lists of EWS::FLI1 "activated" and "repressed" genes and performed gene set enrichment[34] (Fig. 3C). Overall, EWS::FLI activated genes were enriched in all modules, whereas repressed genes were enriched only in module 2 (Fig.

Fig.3. EwS gene regulatory modules are differentially regulated by EWS::FLI1. (B) HOMER enrichment of EWS/ETS motifs (EWS::FLI, pink; FLI::ETS, gray; EWS::ERG, black) in module peak sets. (B) HOMER enrichment of EWS::FLI binding sites, separated by regions with (orange) or without (dark purple) GGAA microsatellites. (C) Universal enrichment analysis of MSigDB EWS::FLI1 and Ewing sarcoma terms in module gene sets. (D) UMAP embeddings of shNS and shFLI1 samples. EwS subgroup identity of cell lines represented by dashed boxes. Group 1 (A673) labelled with navy/dashed lines, group 2 (CHLA10, PDX305) in orange and group 3 (TC71) in light blue. (E) Expression of EwS target gene expression across shNS vs. shFLI1 samples. (F-H) Module gene set expression across knockdown samples.

3C). When combined with the lack of GGAA-uSat enrichment in module 2 (Fig. 3B), these observations are consistent with previous reports that repressed loci usually do not have long stretches of GGAA repeats[8,46]. Altogether, these data reveal distinct and polarized fusion-mediated regulatory events across EwS cell lines that differentially depend on GGAA repeats, suggesting that the diversity of fusion function is dependent on the motif landscape at fusion binding sites.

To directly address how different gene modules are affected by EWS::FLI1, we performed single-cell multiomic sequencing on selected cell lines from each EwS subgroup after shRNA-mediated knockdown of the fusion (Fig. 3D). Following quality control (Fig. S2A), we recovered 11,614 cells expressing negative control

shRNA (A673 shNS, CHLA10 shNS, PDX305 shNS, TC71 shNS) and 12,466 cells expressing shRNA specific to the FLI1 transcript (A673 shFLI1, CHLA10 shFLI1, PDX305 shFLI1, TC71 shFLI1). All shFLI1 cells showed decreased expression of EwS TGs compared to shNS cells (Fig. 3E), providing evidence that the knockdown effectively induced loss of EWS::FLI1 activity. EWS::FLI1 knockdown had specific, predictable, and reproducible effects on all cell lines. Importantly the same stratification of shNS cell lines into groups with specific module enrichment was observed, indicating that lentiviral transduction and selection did not alter dominant gene programs (Fig. 3F-H, Fig. S2B-D). In addition, each module was affected by EWS::FLI1 knockdown in a manner consistent with predictions based on module gene sets. Specifically, EWS::FLI1 knockdown resulted in decreased expression of module 1 and 3 genes (Fig. 3F,H, Fig. S2B,D) and increased expression of module 2 genes (Fig. 3G, Fig. S2C). These data confirm that EWS::FLI1 activates cis-regulatory networks found in modules 1 and 3 but represses module 2. The results also support our initial hypothesis that EWS::FLI1-regulated gene modules differ among commonly used EwS cell lines. They further show that specific and reproducible patterns of transcriptomic and phenotypic signatures dominate each cell line and these predictably respond to fusion knockdown.

CHLA10 exhibits intra-tumoral heterogeneity of gene regulatory modules

Most EwS cell lines in this study are dominated by a single gene-regulatory program (modules 1, 2, or 3). A notable exception is CHLA10, which contains mutually exclusive sub-programs defined by differences in both EWS::FLI1 motif activity and module utilization (Fig. 1B, Fig. 2A). This distinction is particularly striking because CHLA10 and CHLA9 originated from tumors from the same patient, with CHLA10 being derived post-therapy and following disease relapse. To further investigate this intratumoral heterogeneity, we performed unsupervised Louvain clustering[47] on EwS cell lines (Fig. S3A) and computed the Pearson correlation coefficient of the most highly variable regions of accessibility. As expected, CHLA9, TC32, and PDX305 cells grouped together based on similarity (Fig. S3B). In contrast, CHLA10 segregated into two clusters (C22 &

Fig. 4. CHLA10 is a microcosm of EWS::FLI1 mediated heterogeneity
(A) UMAP of CHLA10 colored by cluster, C22 dark purple, C23 light purple. (B) UMAP of module peak accessibility and gene set expression in CHLA10 cells. (C) Differential motif enrichment between C22 and C23, TFs with -log10(p-value) > 250 highlighted. (D) Universal enrichment of GO: Biological Process terms in differentially expressed genes between C22 and C23, terms associated with TGF-β in bold. (E) Universal enrichment of MSigDB Ewing Sarcoma terms in differentially expressed genes between C22 and C23. (F) Pathway activity scores across C22 and C23, TGF-β highlighted in bold.

C23) that each showed greater transcriptional similarity to other cell lines than to each another (Fig 4A, Fig. S3B), highlighting the extent of its internal heterogeneity.

Cell-cycle effects contributed to but could not fully explain CHLA10 heterogeneity (Fig. S3C-D). Instead, plotting peak accessibility and transcript abundance of each module revealed that C22 cells enriched for module 3 while C23 cells relied on module 2 (Fig. 4B). Motif enrichment of differentially accessible peaks further

underscored the nature of the split: RUNX motifs were enriched in C22 cells whereas AP-1, TEAD, and HOX TF motifs predominated in C23 cells (Fig. 4C). Gene-set enrichment mirrored these findings, C22 expressed neural programs whereas C23 cells were enriched for EMT, migration, and TGF-β signaling gene sets (Fig.S3D-E). Pathway-activity inference analysis between cell clusters[48] further implicated TGF-β as a dominant upstream driver in C23 (Fig. 4F).

Collectively, the isogenic model CHLA9/CHLA10 encapsulates the spectrum of transcriptional states seen across the other EwS cell lines and pinpoints TGF-β signaling as key mediator of the module 2 transcriptional state.

Sub-populations of EwS cell lines differentially respond to TGF-β stimulus

Prior reports have shown that TGF-β promotes pro-angiogenic extracellular-matrix (ECM) gene expression in EwS[49]. To interrogate the impact of TGF-β signaling on the newly defined transcriptional states in EwS, we performed multiomic profiling on A673 (module 1-dominant), CHLA10 (module 2 dominant), and TC71 (module 3 dominant) after treatment with recombinant TGF-β1 for 24 hours and compared them to vehicle-treated controls. After quality control filtering 19,611 cells were retained for analysis (Fig. S4A). Dimensionality reduction revealed substantial shifts in both gene expression and chromatin accessibility in A673 and CHLA10 following TGF-β treatment, whereas TC71 profiles remained relatively stable (Fig. 5A-

Fig. 5 Sub-populations of EwS cell lines differentially respond to TGF-β stimulus
(A-C) UMAP of A673, CHLA10, and TC71 cells colored by condition (vehicle control or TGF-β1 treatment) – *column 1*; and cluster – *column 2*; Fraction of treated versus control cells in each cluster with highest and lowest indicated by arrow - *column 3*; Expression of fusion regulated module 2 gene set, highest and lowest TGF-β clusters highlighted with dashed lines – *column 4*. (D-F) Correlation between TF expression and chromVAR TF deviation scores of EwS TFs across clusters, highest and lowest TGF-β clusters highlighted by arrows. (G) Western blot of active (pS465/467) and total SMAD2 and FOSL2 protein in TC71 and CHL10 cells after short (30 minute) and long term (24 hr) TGF-β1 treatment durations. (H) H&E staining of 3 CHLA10 xenografts in NSG mice generated by subcutaneous or tail vein injection (*upper left*). Those regions of interest (ROIs) subjected to whole human transcriptome digital spatial profiling are highlighted. Heatmap of module gene expression across ROIs (*bottom left*). (I) Scatter plot of average expression of Gene Ontology: Biological Process (GO:BP) TGF-β response genes and extracellular matrix (ECM) organization genes compared to *FOSL2* expression in ROIs. Pearson R and p-value highlighted (*top left*).

C). In A673 and CHLA10, TGF-β treatment resulted in a clear polarization of cells towards module 2 usage (Fig. 5A-C), while expression patterns of modules 1 and 3 remained consistent with prior observations (Fig. S4B).

The muted response of the TC71 cell line (module-3 dominant) to the influence of TGF-β (Fig. 5A-C, Fig. S4B) suggests that pre-existing transcriptional states may condition the sensitivity to TGF-β signaling.

We next examined TF expression and motif accessibility in clusters exhibiting the most pronounced or lowest enrichment of TGF-β-treated cells (indicated by arrows in Fig. 5A-C). Correlation analyses revealed a reciprocal enrichment pattern, where FOSL2 was upregulated in TGF-β responsive clusters, while RUNX3 was enriched in non-responsive clusters. Western blot validation confirmed TGF-β-induced FOSL2 expression specifically in CHLA10 but not in the relatively insensitive TC71 cells, despite observed SMAD2 phosphorylation (Fig. 5G).

Our recent whole human transcriptome digital spatial profiling (DSP) study of CHLA10 tumor xenografts (Nanostring GeoMx) showed that transcriptionally distinct tumor cell subpopulations are spatially distributed[22]. We therefore interrogated these data to determine if the identified gene regulatory modules in CHLA10 cells were randomly or spatially distributed across tumors in vivo. We first selected genes from modules 1 & 3 that were significantly upregulated (average $\log_2$ fold change > 0.25) by EWS::FLI1 as well as module 2 genes that were significantly repressed by the fusion (average $\log_2$ fold change < -0.25) and applied these fusion-regulated gene sets to the DSP data. Intriguingly, regions of interest (ROIs) that were captured from tumor borders (ROIs 2, 6, 8 & 10) were enriched with tumor cells that displayed heightened module-2 expression, whereas densely packed tumor cores (ROIs 1, 3, 4, 5, 7, 11) favored modules 1 and 3 (Fig. 5H). Module 3 was also highly favored in subcutaneous compared to tail vein-derived disseminated tumors (Fig. 5H). Cells captured specifically from the border of a metastatic retroperitoneal tumor (ROIs 9 & 10) expressed the highest levels of *FOSL2*, TGF-β-response genes, and ECM-organization programs (Fig. 5I).

In summary, these findings highlight TGF-β-driven activation of module 2 gene expression in EwS cells via AP-1 transcription factors, particularly FOSL2. Furthermore, spatial analysis suggests module 2 programs are enriched at invasive tumor regions, underscoring the role of TGF-β signaling in mediating EwS cell heterogeneity and tumor progression dynamics.

Patient tumors show heterogeneous usage of EwS cis gene regulatory modules

We next sought to determine whether the gene-regulatory networks defined in our cell line panel are similarly deployed across the landscape of early passage patient-derived-xenograft (PDX) and primary patient biopsies. To determine how the three regulatory gene programs manifest in these tumors, we used the fusion-regulated gene sets as defined above. In a series of publicly available PDX tumor profiles[20], module 1 was prominent, while modules 2 and 3 (Fig. 6A) showed more modest and heterogeneous enrichment. On the other hand, in six primary EwS tumor samples[50] (Fig. 6B, Fig. S4D), profiled by single-nucleus RNA-seq, every case contained discrete subpopulations that preferentially expressed either module 2 or module 3. In contrast to the PDX data, module 1-positive cells were comparatively rare in primary tumors.

We next asked whether these single-cell patterns could be detected in bulk transcriptome data that were generated from three independent patient datasets[1,10,51]. To achieve this we employed a deconvolution method[52] that identifies the overlap of signals from bulk measurements with a single-cell transcriptomic reference and captures any proportion of the bulk signal not represented by single-cell data in an 'intercept' term (schematized in Fig. 6C). We first validated the method on bulk RNA sequencing data taken from many of the same EwS cell lines we profiled in our experiments and found a high level of concordance with little signal captured in the intercept term (Fig. S4E). We then tested the transcriptomic overlap of EwS subgroups in 85 non-clinically annotated patient tumors that were profiled by bulk RNA sequencing in two independent cohorts (PRJNA253759, PRJNA261990). Group 2 signals were evident in nearly all tumors and group 3 cell signals were also present in more than half of tumors, supporting the confirming heterogeneity of gene network usage and cell state in primary tumors (Fig. 6D). Tumors with abundant group 2 signal also displayed significantly higher expression of TGF-β response genes and FOSL2 expression (Fig. 6E), linking our in-vitro observation that TGF-β drives module 2 activation to patient tumors in the clinical setting.

Finally, we examined microarray data from a series of 46 clinically annotated primary tumors (GSE61357) that were uniformly treated and for whom outcome data were available[1]. First, we observed significant correlation between expression of FOSL2, TGF-β response and module 2 expression (Fig. 6F). Furthermore, high expression of fusion-repressed module 2, and particularly of its key effector *FOSL2*, identified patients with markedly worse event-free survival, whereas modules 1 and 3 had little prognostic value (Fig. 6G-H, Fig. S4F). Module 2, linked to TGF-β responsiveness and FOSL2 activity, emerges as the predominant—and

clinically adverse—program in primary EwS, underscoring its potential as both a biomarker of aggressive disease and a therapeutic target.

Fig.6 Human tumor data show usage of EWS::FLI1 gene regulatory modules. (A) UMAP embedding of five EwS PDX samples (*left*) UMAP of fusion regulated module gene sets in PDX samples (*right*) (B) UMAP embedding of single nuclei sequencing of 6 patient tumor samples (left). Patient tumor scRNA expression of fusion regulated modules (*right*). (C) Schematic depicting deconvolution method (D) Heatmap EwS subgroup signal in bulk EwS patient tumor RNA-seq. The intercept row represents remaining signal (bottom). Patient tumor samples annotated by GEO BioProject. (E) Scatter plot of bulk RNA patient samples by group 2 signature and GO: BP TGF-β response genes and colored log10 FOSL2 expression. Pearson R and p-value highlighted (*top left*). (E) Scatter plot of microarray patient samples by module 2 expression and GO: BP TGF-β response genes and colored FOSL2 expression. Pearson R and p-value highlighted (*top left*). (G) Survival curves of patient microarray data[1] (n = 46), high expression of module 2 gene set in red (top 15% expressing samples, n = 7), low expression in blue (bottom 85%, n = 39) (H) Survival curves of high expression of *FOSL2* samples (top 15%, n = 7, bottom 50%, n=39).

Discussion

Single-cell genomics has made it possible to chart, with unprecedented resolution, the transcriptional circuitry that underlies intratumoral diversity[53,54]. In EwS, a cancer that has very few genetic mutations beyond its canonical chromosomal translocation[20-22,40] and notable clinical heterogeneity, this diversity likely arises from epigenetic plasticity[55,56]. Our study enhances this understanding by defining two distinct transcriptional and chromatin accessibility states consistently observed across cell lines, patient-derived xenografts (PDXs), and primary tumor samples.

The first transcriptional state, associated with module 3 genes, is enriched for canonical GGAA-microsatellite enhancers, high EWS::FLI1 expression, and neural-crest transcription factors such as TFAP2A and RUNX3. In contrast, the second state associated with module 2 genes lacks GGAA repeats, is suppressed by EWS::FLI1, and involves activation by TEAD and AP-1 family members together with the homeobox factor HOXD13—all elements previously linked to epithelial-to-mesenchymal transition (EMT), invasion, and a low-fusion state[57-59].

The mutually antagonistic architecture of these modules is consistent with earlier observations that EWS::FLI1-repressed loci tend to be short-repeat or single ETS sites that recruit additional repressor complexes such as ETV6[8,60,61]. Further supporting this, recent data[35] demonstrate EWS::FLI1's role in shaping cell identity by controlling promoter-enhancer looping and 3D chromatin networks. Loss of EWS::FLI1 was shown to alter 3D chromatin structure, activating enhancers enriched for AP-1 motifs and facilitating mesenchymal gene expression, consistent with our findings.

Analysis of three independent patient cohorts confirmed module 2 and module 3 signatures in most primary tumors. Clinically, elevated expression of module 2 genes, notably FOSL2, correlated with inferior event-free survival in a prospectively treated cohort of patients with localized EwS. This association nominates module 2 expression as a potential biomarker of aggressive disease and links TGF-β signaling to EMT-like transcriptional states and poor prognosis.

Another key insight from our study is the link between cells expressing high levels of module 2 genes and heightened responsiveness to TGF-β. These findings argue that external cues can re-wire specific subsets of EwS cells towards a more extreme mesenchymal/EMT-like state in the absence of genetic change. This is in line with emerging data that cell plasticity in EwS is influenced by a TGF-β-induced transition of EwS cells to cancer-associated fibroblast-like states[22], which appear to be "unlocked" by TGFB2 response[62]. Thus, therapeutic strategies that disrupt these microenvironmental signals—or directly target downstream effectors such as FOSL2—may limit metastatic spread by constraining module 2 activation. Moreover, by promoting a fibroblast-like phenotype and dampening inflammatory signaling, TGF-β may also contribute to immune evasion and the establishment of an immunosuppressive tumor microenvironment[23,24,63].

Our study is constrained by the limited number of primary single-cell datasets currently available; larger, clinically annotated cohorts will be required to validate the prognostic utility of module balance and to map temporal dynamics during therapy and relapse. Moreover, while CHLA9/CHLA10 serves as a useful isogenic model of plasticity, early-passage or organoid systems may preserve an even richer landscape of states. Finally, dissecting the full spectrum of extrinsic factors—beyond TGF-β—that tip the balance between modules will be key to developing microenvironment-targeted interventions.

In summary, our study provides novel insights into the TF regulatory networks that interact with the EWS::FLI1 fusion protein to sustain distinct cellular states and facilitate dynamic transitions between these states. Specifically, we identified core gene network modules and key TFs, notably FOSL2 and other module 2 TFs, which antagonize fusion-mediated repression, promoting a mesenchymal and EMT-like state associated with more aggressive disease phenotypes. Furthermore, our findings highlight cell-extrinsic TGF-β as a critical factor capable of activating FOSL2 and driving the aggressive module 2 program in vivo. By uncovering these regulatory mechanisms and their functional implications, we offer a foundation for developing therapeutic strategies aimed at limiting metastasis, overcoming treatment resistance, and ultimately improving clinical outcomes for patients with Ewing sarcoma.

Acknowledgments

The authors thank the Fred Hutchinson Scientific Computing Group and Genomics Core staff. Grant and gift support for this work is gratefully acknowledged and was provided by the following sources: Fred Hutch Translational Data Science Integrated Research Center Award (to O.G. Waltner), NIH/NCI 1L40CA264534-01 (to S.N. Furlan), 1Million4Anna Foundation (to S.N. Furlan and E.R. Lawlor), NIH/NCI R01 CA215981 (to E.R. Lawlor), F31CA247104 (to A.A. Apfelbaum); research supported by the 2022 AACR-QuadW Foundation Sarcoma Research Fellowship in Memory of Willie Tichenor, Grant Number 22–40–37-WREN (to E.D. Wrenn); Sam Day Foundation (to E.R. Lawlor). Fred Hutch Scientific Computing, NIH grants S10-OD-020069 and S10-OD-028685, and the Fred Hutch/University of Washington/Seattle Children's Cancer Center Consortium NCI Cancer Consortium Grant P30 CA01570.

Authors' Contributions

O.G. Waltner: Conceptualization, data curation, software, formal analysis, visualization, writing–original draft, writing–review and editing. **A. Apfelbaum:** Conceptualization, data generation and curation, formal analysis, visualization, writing–original draft, writing–review and editing. **E.D. Wrenn:** Conceptualization, data generation and curation, formal analysis, visualization, writing–original draft, writing–review and editing. **M. Mendoza, S. Bhise, and S.B. Kanaan:** Data generation and curation. **R. Green-Gladden:** Manuscript writing–review and editing. **R. Volden and Z Kronenberg:** Formal analysis, software, writing–review and editing. **A. Patel and M. Dyer:** Data generation, writing–original draft, writing–review and editing. **J.R. Sarthy:** Conceptualization, supervision, writing–review and editing. **E.R. Lawlor:** Conceptualization, supervision, funding acquisition, project administration, writing–review and editing. **S.N. Furlan:** Conceptualization, software, formal analysis, supervision, funding acquisition, project administration, writing–review and editing.

Declaration of Interests

S.N. Furlan reports grants from the American Cancer Society, 1Million 4 Anna Foundation, and an NIH Extramural Loan Repayment Program for Pediatric Research. E.R. Lawlor reports grants from the National Cancer Institute, 1Million 4 Anna, and Sam Day Foundation during the conduct of the study as well as grants from Cure Childhood Cancer Foundation outside the submitted work. No disclosures were reported by the other authors.

Methods
Cell Lines and PDX

The EwS cell lines A673, SKNMC, CHLA10, CHLA9, TC32, A4573, RDES, and TC71 were obtained from ATCC and COG (https://www.childrensoncologygroup.org/) cell lines repositories. The osteosarcoma cell line U2OS and rhabdomyosarcoma cell line RD were also obtained from ATCC. 5064L and 240L MSCs were obtained from Dr. Darwin Prockop. H7 MSCs were generously gifted from Dr. Alejandro Sweet-Cordero. PDX305 was generated from a recurrent metastatic tumor, as previously described[22]. A673, TC32, A4573, TC71, and SKNMC cell lines were maintained in RPMI 1640 media (Gibco) supplemented with 10% FBS (Atlas Biologicals) and 2 mmol/L-glutamine (Life Technologies). RDES was maintained in RPMI 1640 media supplemented with 15% FBS. Isogenic cell lines, CHLA10 and CHLA9, were maintained in IMDM media (Fisher) supplemented with 20% FBS, 2 mmol/L-glutamine, and 1X Insulin-Transferrin-Selenium (Gibco). PDX305 was maintained in RPMI 1640 media supplemented with 10% FBS, 2 mmol/L-glutamine, and 1% Antimycotic-Antibiotic (Gibco). U2OS was maintained in McCoy's 5A media supplemented with 10% FBS and 2mmol/L-glutamine and RD was maintained in DMEM with 10% FBS. All hMSC lines were maintained in MEM-alpha media supplemented with 20% FBS, 2 mmol/L-glutamine, and 1% Antimycotic-Antibiotic. Cells were cultured at 37°C with 5% CO_2. Cells were all confirmed to be mycoplasma free and identities subject to STR-confirmation every 6 months.

Lentivirus Production and Genetic knockdown

pLKO.1 shNS (SHC002) and shFLI1 (TRCN0000005322) were used for FLI1 knockdown multiome experiments (Sigma). Knockdown plasmids were co-transfected with pCD/NL-BH*DDD (addgene, #17531) and pMD2.G (addgene, #12259) plasmids into 293FT packaging cells using PEI. 24 hours post-transfection, sodium butyrate was added to plates for 6 hours. Viral supernatant was collected 48 hours after transfection, filtered through 40 μM filters, and added the cells for 24-hours. Ewing sarcoma cells were selected by supplementing the media with 1.5 μg/mL puromycin (Gibco) for 48 hours before collection at 96 hours.

TGF-β treatment prior to single cell sequencing

TC71, A673, and CHLA10 cells were plated at 500,000 viable cells per 10cm plate. 48 hours later recombinant human TGFβ1 (R&D 7754-BH-005, final dose 10 ng/mL in sterile 4 mM HCl and 0.1% BSA) or vehicle control was added. 24 hours later cells were trypsinized and >$1x10^6$ viable cells were subjected to single cell library preparation and sequencing (details below).

Western blots

Cells were washed with ice cold PBS, scraped and centrifuged, then lysed with RIPA buffer (Fisher Scientific) supplemented with protease and phosphatase inhibitors (Sigma) and sonicated (ten 30 second cycles). For TGF-β treated samples, recombinant TGFβ1 (10 ng/mL) or vehicle control (4 mM HCl, 0.1% BSA) were included in the media for either 30 minutes or 24hrs before lysis. Western blots were performed using the Bio-Rad Mini-PROTEAN Tetra System using 40 ug of lysate and 4-15% Mini-PROTEAN TGX polyacrylamide gels. Following transfer, nitrocellulose membranes were blocked in Odyssey Blocking Buffer (LI-COR) for 1 hour. Membranes were washed and incubated rotating overnight at 4°C with primary antibodies (FOSL2 CST #19967, HAND2 R&D AF3876, RUNX3 CST #13089, TEAD1 CST #12292, pSMAD2 S465/467 CST #3108, SMAD2 CST #3103, GAPDH Invitrogen AM4300, Vinculin CST #13901). Membranes were then washed four times in TBST for 5 minutes each and incubated with secondary antibodies (LI-COR IRDye 700CW or 800CW; 1:10,000) for 1 hour. After 4 additional 5 minute TBST washes membranes were scanned on a LI-COR Odyssey scanner.

Digital Spatial Profiling

Q3 normalized gene counts from GeoMx Digital Spatial Profiling (NanoString) of EwS cell line xenograft tumors formed by CHLA10 injection were obtained from previous work[22]. This processed data was used to query fusion regulated module gene sets with respect to tumor microenvironment and injection site. Gene sets values for each ROI were calculated by converting the DSP Q3 counts matrix into a Seurat object and using the AddModuleScore function.

Single-cell analysis

Cell and library prep

One million EwS cells were resuspended in PBS containing 2% BSA and nuclei were prepared according to manufacturer's specifications. Cell lines in triplicate were pooled 1:1:1 at 1000 cells/uL and libraries were generated using the Chromium Next GEM Single- Cell Multiome ATAC+Gene Expression Kit following the manufacturer's protocol (CG000338 Rev C). The resulting libraries were sequenced using a NovaSeq 6000 targeting a depth of 25K reads per cell (ATAC libraries) and 20K reads per cell (RNA libraries).

Single-cell multiome data analysis

Sequencing reads were demultiplexed and processed using the cellranger ARC pipeline. Cells with poor quality ATAC fragments and multiplets were filtered using ArchR's standard workflow. Cells passing the following were kept for later analysis: percent mitochondrial RNA reads < 15%; 3 < log10(RNA/UMI counts) <= 4.5. Souporcell[64] was used to identify cell lines and remove additional multiplets. Top RNA markers from scRNA data were compared to previously hashed CITEseq profiles[22] to match genotype to cell line (GSE236289: GSM7527501, GSM7527502, GSM7527503). ArchR[38], Signac[65], and Seurat[47] were used for downstream analysis. Dimensionality reduction was performed using addIterativeLSI() for both tile and gene expression matrices. An embedding was created by combineDims() and then addUMAP() with the mulitomic reduction. Group coverages and peaks calls were made using cell line identities. The motif matrix was built using addMotifDeviations(), addBdgPeaks(), and addDeviationsMatrix().

Average gene expression and gene activity scores were computed using addModuleScore(). EWS::FLI1 deviations were computed using the JASPAR2020 motif database. Transcription factor analyses used the cisBP database[66]. Highly utilized TFs were inferred using correlateMatrices(). All heatmaps were made with ComplexHeatmap. Marker peaks and genes were calculated using getMarkerFeatures() and getMarkersHeatmap(). Cis-regulatory elements were identified using addPeak2GeneLinks(). Universal gene set enrichment analysis was performed using clusterProfiler using the function enricher(), with MSigDB[67] Hallmark and gene ontology references. Motif enrichment analysis on kmean peak data was performed using HOMER. The most enriched motifs were identified using findMotifsGenome.pl and motif enrichment was computed and visualized using annotatePeaks.pl.. EWS:ETS binding sites were sourced from previously published data[68], and analyzed using HOMER and chromVAR with the same workflow as above.

Cells treated with TGF-β agonist and captured for multiomic sequencing were demultiplexed and processed using the same workflow as above. QC metrics used were identical to those above. A similar method of identifying each cell line using genetic demultiplexing was employed. Signac(64) was used to perform dimensionality reduction using weighted nearest neighbors. Motif matrices were constructed using RunChromVAR(), using cisBP[66] as a database.

EWS::FLI1 long read sequencing
Starting material was 10x Genomics-generated full-length barcoded cDNA, leftover from what was used to create the Illumina-compatible libraries in the multiome downstream steps. Because 10x Genomics single-cell cDNA preparations typically contain a template switch oligo (TSO) priming artifact (i.e., TSO sequence at both ends of the cDNA molecule) instead of the correct structure, TSO artifact depletion was performed by using biotin-labeled primers for the streptavidin/biotin magnetic selection of molecules with the correct structure, as previously described[69]. To capture the gene transcripts encoding the EWS::FLI1 fusions, an enrichment was performed using xGen™ Hybridization and Wash v2 kit magnetic beads and reagents (IDT #10010351), as described in the manufacturer's protocol, in addition to using 500 ng of cDNA starting material and using biotinylated capture probes designed with high specificity, tiling along exons 5, 6, and 7 of the *EWSR1* transcript, and exon 9 of the *FLI1* transcript. Following hybridization and magnetic capture of EWS::FLI1 molecules, an on-bead PCR was performed to amplify captured molecules, using standard 10x Genomic cDNA amplification protocol (described in 10x Genomics document #CG000315_Rev_B, Step 2.2) with the exception of a number of PCR cycles of 16 and an extension time at 72°C of 3 min. Following this, 500 ng of captured cDNA was DNA-repaired, A-tailed, SMRTbell adaptor-ligated, nuclease-treated, and SMRTbell-cleanup purified using the PacBio SMRTbell Prep kit 3.0 (PacBio #102-141-700) as per manufacturer's protocol. Libraries were sequenced on a PacBio Sequel IIe with 2 hours pre-extension time and 30 hours movie time.
Long-read computational processing pipeline consisted of retrieving error-corrected circular consensus sequence (CCS) BAM files generated by the on-board PacBio Sequel IIe (Pacific Biosciences of California, Inc.). CCS data were then processed by the programs in the IsoSeq3 suite of computational tools: 1) *Lima* to remove sequencing primers, 2) *Tag* to extract UMI and CB sequences, 3) *Refine* to remove polyA sequences, 4) *Correct* to error-correct CBs, and 5) *Dedup* to remove PCR duplicates, 4) *Pbmm2* to align reads. All the IsoSeq3 pipeline steps were used with default settings. Mapped BAMs were then used for pbfusion to detect fusions. Barcodes from long read molecules were extracted from mapped BAMs and merged with data of pbfusion.breakpoints.groups.bed files to count fusion transcript in single cells.

External Data
The 78 direct EwS TGs were sourced from a previously published dataset[20]. Bulk RNAseq patient tumor samples were sourced from dbGaP under accession numbers PRJNA253759 and PRJNA261990. FASTQ files were aligned to reference genome (GRCh38) and using the piperna (https://github.com/furlan-lab/piperna) workflow which uses STAR[70] aligner and generates a count matrix using the Bioconductor summarize overlap function (https://www.bioconductor.org) with its default settings. Microarray data was accessed from GSE63157[1]. RNA probes were annotated using pd.huex.1.0.st.v2, and normalized using oligo.

scRNAseq Patient Tumors
Patient tumor biopsies were sourced and sequenced at St. Jude's Children's Hospital [50]. 10X reads were demultiplexed and processed using the cellranger pipeline. Ambient RNA was filtered out using cellbender. Cells with poor RNA measurements were filtered using the following thresholds: Dimensionality reduction and clustering was performed using standard workflow. Surrounding tissues cells were identified by using FindAllMarkers on Seurat clusters. Tumor enrichment was validated by choosing cells that express *CD99*, *NR0B1*, and EwS TGs. Identified tumor cells underwent normalization and dimensionality reduction with the same workflow. Average gene set scores were calculated using AddModuleScore().

Deconvolution
Bulk RNAseq patient data were deconvoluted using cellSignalAnalysis (https://github.com/constantAmateur/cellSignalAnalysis). EwS subgroups 0-2 were used as transcriptomic references.

References

Volchenboum, S. L. *et al.* Gene Expression Profiling of Ewing Sarcoma Tumors Reveals the Prognostic Importance of Tumor-Stromal Interactions: A Report from the Children's Oncology Group. *J Pathol Clin Res* **1**, 83-94 (2015). https://doi.org/10.1002/cjp2.9

Riggi, N., Suva, M. L. & Stamenkovic, I. Ewing's Sarcoma. *N Engl J Med* **384**, 154-164 (2021). https://doi.org/10.1056/NEJMra2028910

Grunewald, T. G. P. *et al.* Ewing sarcoma. *Nat Rev Dis Primers* **4**, 5 (2018). https://doi.org/10.1038/s41572-018-0003-x

Tirode, F. *et al.* Mesenchymal stem cell features of Ewing tumors. *Cancer Cell* **11**, 421-429 (2007). https://doi.org/10.1016/j.ccr.2007.02.027

Riggi, N., Suva, M. L. & Stamenkovic, I. Ewing's sarcoma origin: from duel to duality. *Expert Rev Anticancer Ther* **9**, 1025-1030 (2009). https://doi.org/10.1586/era.09.81

Lin, P. P., Wang, Y. & Lozano, G. Mesenchymal Stem Cells and the Origin of Ewing's Sarcoma. *Sarcoma* **2011** (2011). https://doi.org/10.1155/2011/276463

Miller, H. E. *et al.* Reconstruction of Ewing Sarcoma Developmental Context from Mass-Scale Transcriptomics Reveals Characteristics of EWSR1-FLI1 Permissibility. *Cancers (Basel)* **12** (2020). https://doi.org/10.3390/cancers12040948

Riggi, N. *et al.* EWS-FLI1 utilizes divergent chromatin remodeling mechanisms to directly activate or repress enhancer elements in Ewing sarcoma. *Cancer Cell* **26**, 668-681 (2014). https://doi.org/10.1016/j.ccell.2014.10.004

Boulay, G. *et al.* Cancer-Specific Retargeting of BAF Complexes by a Prion-like Domain. *Cell* **171**, 163-178 e119 (2017). https://doi.org/10.1016/j.cell.2017.07.036

Crompton, B. D. *et al.* The genomic landscape of pediatric Ewing sarcoma. *Cancer Discov* **4**, 1326-1341 (2014). https://doi.org/10.1158/2159-8290.CD-13-1037

Zoubek, A. *et al.* Does expression of different EWS chimeric transcripts define clinically distinct risk groups of Ewing tumor patients? *J Clin Oncol* **14**, 1245-1251 (1996). https://doi.org/10.1200/JCO.1996.14.4.1245

Davies, A., Zoubeidi, A., Beltran, H. & Selth, L. A. The Transcriptional and Epigenetic Landscape of Cancer Cell Lineage Plasticity. *Cancer Discov* **13**, 1771-1788 (2023). https://doi.org/10.1158/2159-8290.CD-23-0225

Hanahan, D. Hallmarks of Cancer: New Dimensions. *Cancer Discov* **12**, 31-46 (2022). https://doi.org/10.1158/2159-8290.CD-21-1059

Tomazou, E. M. *et al.* Epigenome mapping reveals distinct modes of gene regulation and widespread enhancer reprogramming by the oncogenic fusion protein EWS-FLI1. *Cell Rep* **10**, 1082-1095 (2015). https://doi.org/10.1016/j.celrep.2015.01.042

Sheffield, N. C. *et al.* DNA methylation heterogeneity defines a disease spectrum in Ewing sarcoma. *Nat Med* **23**, 386-395 (2017). https://doi.org/10.1038/nm.4273

Franzetti, G. A. *et al.* Cell-to-cell heterogeneity of EWSR1-FLI1 activity determines proliferation/migration choices in Ewing sarcoma cells. *Oncogene* **36**, 3505-3514 (2017). https://doi.org/10.1038/onc.2016.498

Adane, B. *et al.* STAG2 loss rewires oncogenic and developmental programs to promote metastasis in Ewing sarcoma. *Cancer Cell* **39**, 827-844 e810 (2021). https://doi.org/10.1016/j.ccell.2021.05.007

Pedersen, E. A. *et al.* Activation of Wnt/beta-Catenin in Ewing Sarcoma Cells Antagonizes EWS/ETS Function and Promotes Phenotypic Transition to More Metastatic Cell States. *Cancer Res* **76**, 5040-5053 (2016). https://doi.org/10.1158/0008-5472.CAN-15-3422

Surdez, D. *et al.* STAG2 mutations alter CTCF-anchored loop extrusion, reduce cis-regulatory interactions and EWSR1-FLI1 activity in Ewing sarcoma. *Cancer Cell* **39**, 810-826 e819 (2021). https://doi.org/10.1016/j.ccell.2021.04.001

Aynaud, M. M. *et al.* Transcriptional Programs Define Intratumoral Heterogeneity of Ewing Sarcoma at Single-Cell Resolution. *Cell Rep* **30**, 1767-1779 e1766 (2020). https://doi.org/10.1016/j.celrep.2020.01.049

Apfelbaum, A. A. *et al.* EWS::FLI1 and HOXD13 Control Tumor Cell Plasticity in Ewing Sarcoma. *Clin Cancer Res* **28**, 4466-4478 (2022). https://doi.org/10.1158/1078-0432.CCR-22-0384

Wrenn, E. D. *et al.* Cancer-associated fibroblast-like tumor cells remodel the Ewing sarcoma tumor microenvironment. *Clin Cancer Res* (2023). https://doi.org/10.1158/1078-0432.CCR-23-1111

Visser, L. L. *et al.* Ewing Sarcoma Single-cell Transcriptome Analysis Reveals Functionally Impaired Antigen-presenting Cells. *Cancer Res Commun* **3**, 2158-2169 (2023). https://doi.org/10.1158/2767-9764.CRC-23-0027

Goodspeed, A. *et al.* Single-Cell RNA Sequencing of Ewing Sarcoma Tumors Demonstrates Transcriptional Heterogeneity and Clonal Evolution. *Clin Cancer Res* **31**, 2010-2023 (2025). https://doi.org/10.1158/1078-0432.CCR-24-2040

Chaturvedi, A., Hoffman, L. M., Welm, A. L., Lessnick, S. L. & Beckerle, M. C. The EWS/FLI Oncogene Drives Changes in Cellular Morphology, Adhesion, and Migration in Ewing Sarcoma. *Genes Cancer* **3**, 102-116 (2012). https://doi.org/10.1177/1947601912457024

Chaturvedi, A. *et al.* Molecular dissection of the mechanism by which EWS/FLI expression compromises actin cytoskeletal integrity and cell adhesion in Ewing sarcoma. *Mol Biol Cell* **25**, 2695-2709 (2014). https://doi.org/10.1091/mbc.E14-01-0007

Smith, R. *et al.* Expression profiling of EWS/FLI identifies NKX2.2 as a critical target gene in Ewing's sarcoma. *Cancer Cell* **9**, 405-416 (2006). https://doi.org/10.1016/j.ccr.2006.04.004

Bilke, S. *et al.* Oncogenic ETS fusions deregulate E2F3 target genes in Ewing sarcoma and prostate cancer. *Genome Res* **23**, 1797-1809 (2013). https://doi.org/10.1101/gr.151340.112

Bledsoe, K. L. *et al.* RUNX3 facilitates growth of Ewing sarcoma cells. *J Cell Physiol* **229**, 2049-2056 (2014). https://doi.org/10.1002/jcp.24663

Garcia-Garcia, L. *et al.* The Transcription Factor FEZF1, a Direct Target of EWSR1-FLI1 in Ewing Sarcoma Cells, Regulates the Expression of Neural-Specific Genes. *Cancers (Basel)* **13** (2021). https://doi.org/10.3390/cancers13225668

Lin, L. *et al.* Super-enhancer-associated MEIS1 promotes transcriptional dysregulation in Ewing sarcoma in co-operation with EWS-FLI1. *Nucleic Acids Res* **47**, 1255-1267 (2019). https://doi.org/10.1093/nar/gky1207

Marques Howarth, M. *et al.* Long noncoding RNA EWSAT1-mediated gene repression facilitates Ewing sarcoma oncogenesis. *J Clin Invest* **124**, 5275-5290 (2014). https://doi.org/10.1172/JCI72124

Shi, X. *et al.* EWS-FLI1 regulates and cooperates with core regulatory circuitry in Ewing sarcoma. *Nucleic Acids Res* **48**, 11434-11451 (2020). https://doi.org/10.1093/nar/gkaa901

Kinsey, M., Smith, R. & Lessnick, S. L. NR0B1 is required for the oncogenic phenotype mediated by EWS/FLI in Ewing's sarcoma. *Mol Cancer Res* **4**, 851-859 (2006). https://doi.org/10.1158/1541-7786.MCR-06-0090

Sanalkumar, R. *et al.* Highly connected 3D chromatin networks established by an oncogenic fusion protein shape tumor cell identity. *Sci Adv* **9**, eabo3789 (2023). https://doi.org/10.1126/sciadv.abo3789

Minas, T. Z. *et al.* Combined experience of six independent laboratories attempting to create an Ewing sarcoma mouse model. *Oncotarget* **8**, 34141-34163 (2017). https://doi.org/10.18632/oncotarget.9388

Tirosh, I. *et al.* Dissecting the multicellular ecosystem of metastatic melanoma by single-cell RNA-seq. *Science* **352**, 189-196 (2016). https://doi.org/10.1126/science.aad0501

Granja, J. M. *et al.* ArchR is a scalable software package for integrative single-cell chromatin accessibility analysis. *Nat Genet* **53**, 403-411 (2021). https://doi.org/10.1038/s41588-021-00790-6

Orth, M. F. *et al.* Systematic multi-omics cell line profiling uncovers principles of Ewing sarcoma fusion oncogene-mediated gene regulation. *Cell Rep* **41**, 111761 (2022). https://doi.org/10.1016/j.celrep.2022.111761

Apfelbaum, A. A., Wrenn, E. D. & Lawlor, E. R. The importance of fusion protein activity in Ewing sarcoma and the cell intrinsic and extrinsic factors that regulate it: A review. *Front Oncol* **12**, 1044707 (2022). https://doi.org/10.3389/fonc.2022.1044707

Faustino, A. *et al.* mTOR pathway overactivation in BRAF mutated papillary thyroid carcinoma. *J Clin Endocrinol Metab* **97**, E1139-1149 (2012). https://doi.org/10.1210/jc.2011-2748

Kakkar, A. *et al.* Alterations in BRAF gene, and enhanced mTOR and MAPK signaling in dysembryoplastic neuroepithelial tumors (DNTs). *Epilepsy Res* **127**, 141-151 (2016). https://doi.org/10.1016/j.eplepsyres.2016.08.028

Prabowo, A. S. *et al.* BRAF V600E mutation is associated with mTOR signaling activation in glioneuronal tumors. *Brain Pathol* **24**, 52-66 (2014). https://doi.org/10.1111/bpa.12081

Heinz, S. *et al.* Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Mol Cell* **38**, 576-589 (2010). https://doi.org/10.1016/j.molcel.2010.05.004

Kim, S., Denny, C. T. & Wisdom, R. Cooperative DNA binding with AP-1 proteins is required for transformation by EWS-Ets fusion proteins. *Mol Cell Biol* **26**, 2467-2478 (2006). https://doi.org/10.1128/MCB.26.7.2467-2478.2006

Johnson, K. M., Taslim, C., Saund, R. S. & Lessnick, S. L. Identification of two types of GGAA-microsatellites and their roles in EWS/FLI binding and gene regulation in Ewing sarcoma. *PLoS One* **12**, e0186275 (2017). https://doi.org/10.1371/journal.pone.0186275

Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902 e1821 (2019). https://doi.org/10.1016/j.cell.2019.05.031

Badia, I. M. P. *et al.* decoupleR: ensemble of computational methods to infer biological activities from omics data. *Bioinform Adv* **2**, vbac016 (2022). https://doi.org/10.1093/bioadv/vbac016

Hawkins, A. G. *et al.* Wnt/beta-catenin-activated Ewing sarcoma cells promote the angiogenic switch. *JCI Insight* **5** (2020). https://doi.org/10.1172/jci.insight.135188

The Single-cell Pediatric Cancer Atlas: Data portal and open-source tools for single-cell transcriptomics of pediatric tumors. *bioRxiv* (2024).

Brohl, A. S. *et al.* The genomic landscape of the Ewing Sarcoma family of tumors reveals recurrent STAG2 mutation. *PLoS Genet* **10**, e1004475 (2014). https://doi.org/10.1371/journal.pgen.1004475

Young, M. D. *et al.* Single cell derived mRNA signals across human kidney tumors. *Nat Commun* **12**, 3896 (2021). https://doi.org/10.1038/s41467-021-23949-5

Gavish, A. *et al.* Hallmarks of transcriptional intratumour heterogeneity across a thousand tumours. *Nature* **618**, 598-606 (2023). https://doi.org/10.1038/s41586-023-06130-4

Marusyk, A., Almendro, V. & Polyak, K. Intra-tumour heterogeneity: a looking glass for cancer? *Nat Rev Cancer* **12**, 323-334 (2012). https://doi.org/10.1038/nrc3261

van Groningen, T. *et al.* Neuroblastoma is composed of two super-enhancer-associated differentiation states. *Nat Genet* **49**, 1261-1266 (2017). https://doi.org/10.1038/ng.3899

Boeva, V. *et al.* Heterogeneity of neuroblastoma cell identity defined by transcriptional circuitries. *Nat Genet* **49**, 1408-1413 (2017). https://doi.org/10.1038/ng.3921

Zhang, H. *et al.* TEAD transcription factors mediate the function of TAZ in cell growth and epithelial-mesenchymal transition. *J Biol Chem* **284**, 13355-13362 (2009). https://doi.org/10.1074/jbc.M900843200

Katschnig, A. M. *et al.* EWS-FLI1 perturbs MRTFB/YAP-1/TEAD target gene regulation inhibiting cytoskeletal autoregulatory feedback in Ewing sarcoma. *Oncogene* **36**, 5995-6005 (2017). https://doi.org/10.1038/onc.2017.202

Croushore, E. E., Stipp, C. S. & Gordon, D. J. EWS-FLI1 and Activator Protein-1 (AP-1) Reciprocally Regulate Extracellular-Matrix Proteins in Ewing sarcoma Cells. *Int J Mol Sci* **25** (2024). https://doi.org/10.3390/ijms25168595

Sankar, S. *et al.* Mechanism and relevance of EWS/FLI-mediated transcriptional repression in Ewing sarcoma. *Oncogene* **32**, 5089-5100 (2013). https://doi.org/10.1038/onc.2012.525

Lu, D. Y. *et al.* The ETS transcription factor ETV6 constrains the transcriptional activity of EWS-FLI to promote Ewing sarcoma. *Nat Cell Biol* **25**, 285-297 (2023). https://doi.org/10.1038/s41556-022-01059-8

Wrenn, E. D. *et al.* Autocrine TGFβ2 enforces a transcriptionally hybrid cell state in Ewing sarcoma. *bioRxiv* (2025).

Kumar, R. R. *et al.* Decoding the immune landscape in Ewing sarcoma pathogenesis: The role of tumor infiltrating immune cells and immune milieu. *J Bone Oncol* **52**, 100678 (2025). https://doi.org/10.1016/j.jbo.2025.100678

Heaton, H. *et al.* Souporcell: robust clustering of single-cell RNA-seq data by genotype without reference genotypes. *Nat Methods* **17**, 615-620 (2020). https://doi.org/10.1038/s41592-020-0820-1

Stuart, T., Srivastava, A., Madad, S., Lareau, C. A. & Satija, R. Single-cell chromatin state analysis with Signac. *Nat Methods* **18**, 1333-1341 (2021). https://doi.org/10.1038/s41592-021-01282-5

Weirauch, M. T. *et al.* Determination and inference of eukaryotic transcription factor sequence specificity. *Cell* **158**, 1431-1443 (2014). https://doi.org/10.1016/j.cell.2014.08.009

Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* **102**, 15545-15550 (2005). https://doi.org/10.1073/pnas.0506580102

Theisen, E. R. *et al.* Chromatin profiling reveals relocalization of lysine-specific demethylase 1 by an oncogenic fusion protein. *Epigenetics* **16**, 405-424 (2021). https://doi.org/10.1080/15592294.2020.1805678

Al'Khafaji, A. M. *et al.* High-throughput RNA isoform sequencing using programmed cDNA concatenation. *Nat Biotechnol* **42**, 582-586 (2024). https://doi.org/10.1038/s41587-023-01815-7

Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013). https://doi.org/10.1093/bioinformatics/bts635

Supplemental Figures

A

Cell line/PDX name	Cancer/lineage type	age;sex	Fusion type	Tumor site
A673	Ewing sarcoma	15y;F	EWS::FLI1;type 1	Muscle
A4573	Ewing sarcoma	17y;F	EWS::FLI1;type 3	Pleural effusion
CHLA9 (isogenic with CHLA10)	Ewing sarcoma	14y;F	EWS::FLI1;type 1	Thoracic lymph node
CHLA10 (isogenic with CHLA9)	Ewing sarcoma	14y;F	EWS::FLI1;type 1	Thoracic lymph node
PDX305	Ewing sarcoma	17y;M	EWS::FLI1;type 1	Abdominal soft-tissue metastasis
RDES	Ewing sarcoma	19y;M	EWS::FLI1;type 2	Humerus
SKNMC	Ewing sarcoma	14y;F	EWS::FLI1;type 1	Retroorbital metastasis
TC32	Ewing sarcoma	2.7y;F	EWS::FLI1;type 1	Ileum and adjacent soft tissue
TC71	Ewing sarcoma	22y;M	EWS::FLI1;type 1	Humerus
RD	rhabdomyosarcoma	7w;F	N/A	Pelvic
U2OS	osteosarcoma	15y;F	N/A	Tibia
5046L	mesenchymal stem cell	27y;M	N/A	Bone marrow aspirate
240L	mesenchymal stem cell	24y;M	N/A	Bone marrow aspirate
H7	mesenchymal stem cell	13y;M	N/A	Bone marrow aspirate

Figure S1

Supp. Fig.S1. Multiomic analyses reveal distinct signatures between non-EwS and EwS cells. (A) Table of all 14 cell lines used for the multiome sequencing. (B) Quality control metrics across cell lines. (C) Gene set expression and activity score of EwS target genes. (D) UMAP of all Ewing and Non-Ewing cell lines colored by chromVAR deviation scores of EWS::FLI1 binding sites, parsed by regions with or without GGAA microsatellites (µSAT; GGAA repeat ≥ 4). (E) UMAP of EwS and non-EwS cells highlighted (red) if EWS::FLI1 fusion transcript was detected by long-read sequencing.

Supp. Fig.S2. EwS gene regulatory modules are differentially regulated by EWS::FLI1. (A) Quality control metrics of knockdown samples. (B-D) UMAP embedding of gene expression scores of module gene sets. Cell lines most enriched for a module gene set are highlighted.

Supp. Fig. S3 CHLA10 is a microcosm of EWS::FLI1 mediated heterogeneity
(A) UMAP of 9 EwS cell lines colored and labeled by cluster (B) Pearson correlation heatmap of EwS clusters. The rows are labeled by cluster ID, the columns are labeled by cell line identity. Cell lines sharing epigenetic signatures highlighted in bold and red boxes. (C) *MKI67* expression across EwS cell lines. (D) Proportion of cells in G1, G2M, and S phase within EwS clusters.

A

Number of genes

Log10 RNA UMI

TSS enrichment

Log10 ATAC fragments

B

Fusion regulated **module 1** gene expression

A673   CHLA10   TC71

Fusion regulated **module 3** gene expression

A673   CHLA10   TC71

C

Tumor ID	SJID	Age	Sex	Primary Site	Sample Site	Sample	Fusion Status
1	SJEWS001321D2	10	M	tibia	tibia	primary	EWSR1-FLI1
2	SJEWS049193D1	11	M	ilium	lung	metastasis	EWSR1-FLI1
3	SJEWS063826D1	17	M	rib	axilla	metastasis	EWSR1-FLI1
4	SJEWS063829D1	8	F	femur	femur	primary	EWSR1-FLI1
5A	SJEWS063834A1	14	F	nd	chest wall	metastasis	EWSR1-FLI1
5B	SJEWS063834A2	14	F	nd	lung	metastasis	EWSR1-FLI1

D

E

Fusion regulated **module 1** gene expression $p = 0.5152$

Fusion regulated **module 3** gene expression $p = 0.7308$

Supp. Fig. S4. Sub-populations of EwS cell lines differentially respond to TGF-β stimulus
(A) Quality control metrics of vehicle control and TGF-β treated cell lines. (B) UMAP colored by gene expression of fusion regulated modules 1 & 3 across cell lines. (C) Clinical details of patient samples subjected to single-cell sequencing. (D) Heatmap depicting similarity scores of publicly available bulk EwS cell line data with multiome scRNA-seq data. (E) Survival curves of patient microarray data[1] (n = 46), high expression of module genes 1 and 3 set in red (top 15% expressing samples, n = 7), low expression in blue (bottom 85%, n = 39)