ASC contributes to sustained neuroinflammation and mild cognitive impairment after closed-head injury

Tao Li*[1], Sergio Castro-Gomez*[#1,2], Pablo Botella Lucena[3], Ana Vieira-Saecker[1], Sarah Schwartz[1], Yushuang Deng[4], Yingying Ding[1], Verena Stein[1], Eicke Latz[5,6,7,8], Michael T. Heneka[# 3,7]

1. Institute of Physiology II, University Hospital Bonn, University of Bonn, Bonn, Germany
2. Center for Neurology, Clinic of Parkinson, Sleep and Movement Disorders, University Hospital Bonn, University of Bonn, Bonn, Germany
3. Luxembourg Centre for Systems Biomedicine (LCSB), University of Luxembourg, Esch-sur-Alzette/Belvaux, Luxembourg
4. German Center for Neurodegenerative Diseases (DZNE), Bonn, Germany
5. Institute of Innate Immunity, University Hospital Bonn, Bonn, Germany
6. Centre of Molecular Inflammation Research, Norwegian University of Science and Technology, Trondheim, Norway
7. Division of Infectious Diseases and Immunology, University of Massachusetts Medical School, Worcester, USA
8. Deutsches Rheuma-Forschungszentrum (DRFZ), Berlin, Germany

***Authorship notes:** TL and SC-G contributed equally to this work.

[#] **Correspondence**: michael.heneka@uni.lu, sergio.castro-gomez@ukbonn.de

Competing interests: M.T.H. is a scientific advisory board member at Alector the Dementia Discovery Fund, and Muna Therapeutics and has received honoraria for oral presentations from Pfizer, Novartis, Roche, Abbvie, and Biogen.

Abstract

Mild brain traumatic brain injury (mTBI) from closed-head injuries (CHI) can lead to prevalent neuropsychiatric disorders, including an increased risk for neurodegenerative diseases and dementia. Inflammasomes are molecular complexes crucial for neuroinflammation and secondary damage after trauma, however their role in mild CHI is poorly understood. In this study, we investigate the cellular expression of inflammasome-related genes and their functional significance in CHI models. Single-cell RNA sequencing of cortical tissue revealed selective expression of *Pycard* (PYD and CARD domain containing), also known as *Asc* (apoptosis-associated speck-like protein containing a caspase recruitment domain), which encodes the inflammasome adaptor ASC, predominantly in microglial clusters. Sustained upregulation of inflammasome-related proteins persisted up to 21 days in a model for mTBI, with this pattern significantly reduced in *Asc*[-/-] mice. Importantly, mild cognitive impairment induced after mild CHI was largely abrogated in *Asc*[-/-] mice. These findings suggest that ASC, as the primary inflammasome adaptor, plays a critical role in sustaining neuroinflammation and contributes to cognitive deficits after mild CHI. This study provides insights into the molecular neuroinflammatory mechanisms underlying CHI, potentially informing future therapeutic strategies.

Introduction

Traumatic brain Injury (TBI) represents a significant global health challenge and is considered a silent epidemic, affecting individuals across all demographics and age groups. Every year, TBI affects around 1.5 million people in both the European Union and the United States. In the EU, these injuries are linked to about 57,000 deaths and 1.5 million hospital admissions (1), while in the U.S. they cause roughly 50,000 deaths, 230,000 hospitalizations with survival, and 80,000–90,000 new cases of long-term disability (2). The majority of cases are clinically classified as mild TBI (mTBI), typically resulting from closed-head injuries (CHI) caused by falls, traffic accidents, violence, contact sports, or military actions (3, 4). Although mTBI symptoms can recover within days or weeks, up to 20% of individuals experience persistent physical, cognitive, and behavioral impairments that lead to a reduced quality of daily life and an elevated risk of neuropsychiatric disorders, including depression and dementia (5–7). Despite its prevalence, the underlying pathophysiology of mTBI resulted from CHI remains poorly understood, posing great challenges for early clinical diagnosis and timely intervention.

Reactive microglia and astrocytes play crucial roles in the innate immune response and secondary damage following TBI. Prolonged activation of these cells impairs debris clearance, contributing to neuroinflammation, exacerbating neuronal dysfunction and damage, and promoting abnormal protein aggregation (8–10). Advances in imaging techniques have allowed the *in vivo* visualization of glial reactivity after CHI, emphasizing the involvement of microglia and astrocytes in the inflammatory cascade that contributes to the development of post-traumatic disorders (11–13). A key element in initiating and maintaining innate immune responses of the brain is inflammasome activation and signaling. Inflammasomes are oligomeric protein scaffolds rapidly activated by some damage-associated molecular patters (DAMPs), and organized as a tripartite complex: a cytosolic danger sensor such as NOD-like receptor family pyrin domain containing 3 (NLRP3), a unique adaptor protein called apoptosis-associated speck-like protein containing a CARD domain (ASC), and a proteolytic effector caspase-1 (CASP1) (14). ASC, an intracellular adaptor protein common to almost all inflammasomes, oligomerizes into highly cross-linked macromolecular assemblies, forming visible ASC aggregates. The fibrillar ASC acts as a molecular platform to recruit pro-CASP1, which optimizes signal transduction via proximity-induced CASP1 activation. Activated CASP1 catalyzes the maturation of pro-forms of Interleukin-1β (IL-1β), IL-18

and mediates the proteolytic activation of Gasdermin D (GSDMD). Oligomerized GSDMD forms pores in

the plasma membrane, allowing the release of IL-1β, IL-18 and other proinflammatory cytokines. The

detection of tissue ASC aggregates is considered a typical hallmark of inflammasome activation (15).

In humans, some inflammasome proteins are detected in blood and cerebrospinal fluid after TBI, and

circulating levels of some proteins such as ASC and CASP1 have been proposed as biomarkers for

determining injury severity (16–18). Furthermore, inflammasome activators, including NLRP1 (19), NLRP3

(20, 21), and absent in melanoma 2 (AIM2) (22), presumably contribute to the innate immune response

and functional disability in mouse models of TBI. these studies primarily focused on inflammasome

activation in models of moderate to severe TBI, such as the commonly used Controlled Cortical Impact

(CCI), in which ASC has been hypothesized to be dispensable (23) . At the same time, their contribution to

mild CHI remains largely unexplored. In the present study, we hypothesized that ASC inflammasomes

ensembles contribute actively to the secondary damage following CHI. We aimed to investigate their role

in neuroinflammatory responses and subsequent cognitive functions by analyzing a well-established

single-cell RNA sequencing (scRNA-seq) dataset of cortical cells from mice subjected to a diffuse TBI

model (26) and by using an additional CHI mouse model that induces mild cognitive impairment (27). Our

scRNA-seq data analysis reveals a subacute expression of *Asc* (also known as *Pycard*) in cortical cells

from mice subjected to midline fluid percussion injury (mFPI), particularly in microglia subpopulation. In

concordance with this analysis, we observed sustained upregulation of inflammasome-related proteins,

including NLRP3, ASC, CASP-1, and IL-1β, following CHI, persisting up to 21-days post injury (dpi). This

expression pattern was significantly reduced in *Asc*$^{-/-}$ mice. Moreover, *Asc*$^{-/-}$ mice were protected from the

mild cognitive impairment induced by our CHI model, as assessed by the Morris Water Maze (MWM). We

also observe ASC-dependent microglial and astrocytic reactivity and interactions that persists over time,

accompanied by the upregulation and aggregation of the ASC protein. Our findings identify ASC, the

primary inflammasome adaptor, as a key molecule in sustaining neuroinflammation and contributing to

cognitive deficits after CHI.

Results

Subacute Asc (Pycard) expression in microglia subpopulations after a model for CHI. To examine the cell expression of gene coding for inflammasome-related proteins after CHI, our first approach was to adopt a single-cell genomic analysis. We chose to re-examine a publicly available scRNA-seq dataset from dissociated cortical cells 7 days (7 days post injury, 7 dpi) followed mFPI (26). mFPI mimics several relevant features of CHI including diffused trauma without localized lesion or hemorrhages, and closely models mild to moderate TBI (28). After quality control, we analyzed 14688 control cells and 10,745 cells in the mFPI group (Figure 1A and supplemental Figure 1A). The complex multidimensional dataset was transformed and visualized using the nonlinear dimensionality reduction technique Uniform Manifold Approximation and Projection (UMAP) (Figure 1A, 1B, 1E). Examination of cellular transcriptional identities at the single-cell level uncovered 22 distinct cellular clusters. Cellular subpopulation labels were assigned via automated cell type annotation per media sample using SingleR (29) (Supplemental Figure 1B). Cells identified as microglia were specially enriched in this dataset (Figure 1B). A differential expression gene analysis of this subset of cells showed a typical disease-associated microglia signature after TBI, with overexpression of type-1 interferon genes (such as *Ifi27l2a, Ifitm3, Ifit3, rf7*) and *Apoe,* and reduction of homeostatic genes such as *tmem119* or *tgfb1* (Figure 1C).

We then evaluated, for each predicted cellular subpopulation, the expression of genes annotated in pathways related to inflammasome activation (pathcards.genecards.org (30), Supplemental Fig 1C). Notably, we found *Asc* (or *pycard,* gene coding for the adaptor protein ASC), as one of the genes with wider expression in clusters represented by microglia, macrophage, and other peripheral infiltrating immune cells (Figure 1D-E and Supplemental Figure 1C, 1D). These clusters show a more reduced percentage of cells expression genes coding for inflammasome effectors such *Gsdmd, Casp1, il1b* or *il18* (Figure 1D and E). Surprisingly, we found a very low percentage of cells expressing genes coding for inflammasome activators such as *Nlrc4, Nlrc5, Nlrp1b, Nlrp3 or Aim2* (Supplemental Fig 1C). These results suggest that the subacute and prolonged expression of the adaptor protein ASC, especially in microglia, may play a central role in the sustained innate immune response and pathophysiology of TBI related to CHI.

ASC contributes to the sustained expression and processing of inflammasome mediators following mild

CHI. To unveil the pathophysiological role of inflammasomes and specifically of ASC as main

inflammasome adaptor and abundantly expressed gene in microglia after CHI, we adopted an

electromagnetic CHI procedure as main model for mTBI (31). In this model, a controlled cortical impact

injury is delivered directly to the midline surface of the skull using a 5 mm diameter tip at a specified depth

and velocity (Supplemental Figure 2A). We found mild motor, reflex or reaction deficits at 1- or 7 dpi when

the mice were evaluated using the Revised Neurobehavioral Severity Scale (NSS-R) for rodents (32)

(Supplemental Figure 2B). To examine the protein expression of inflammasome-related proteins after CHI,

we performed immunoblot analysis of cortices from WT and $Asc^{-/-}$ mice that underwent either sham or CHI

surgery. We analyzed and semi-quantified the protein levels of NLRP3, ASC, CASP1, and IL-1β at 1, 7,

and 21 dpi (Figure 2A-I). Our results revealed sustained upregulation of these proteins up to 21 days after

CHI in WT mice. Notably, mice with a constitutive genetic deficiency of the *pycard* gene ($Asc^{-/-}$) showed a

very significant deficit in the expression and cleavage of CASP1 (Figure 2D-E) and of IL-1β over time

(Figure 2G) fallowing CHI. ELISA measurements further demonstrated significantly lower soluble levels of

IL-1β and TNF-α in $Asc^{-/-}$ mice (Figure 2H-I), particularly at later time points. We additionally evaluated the

levels of cleavage of Caspase-8 (CASP8), a proposed non-canonical enzyme involved in IL-1β processing

(33) (Supplemental Figure 2C and D), and Caspase-3 (CASP3), as indirect measurement of apoptotic

pathways (34) (Supplemental Figure 2C and E). Consisting with the reduction in soluble IL-1β, $Asc^{-/-}$

cortices showed significantly reduced levels of cleaved CASP8 up to 21 days post CHI (Supplemental

Figure 2D), which suggest parallel activation of canonical and non-canonical pathways of IL-1β

processing, both of which appear largely dependent on ASC. Additionally, the cleavage of CASP3 was

significantly reduced in $Asc^{-/-}$ mice (Supplemental Figure 2E), suggesting a strong neuroprotective effect

and a complex crosstalk involving the activation of multiple types of cell death influenced by

inflammasome signaling (35).

To identify the cellular sources of IL-1β following mild CHI, we conducted immunohistochemical staining

for IL-1β, ionized calcium binding adaptor molecule 1 (Iba1) and Glial fibrillary acidic protein (GFAP) on

cortices of WT mice in sham, 7 and 21 dpi groups (Supplemental Figure 3a-d). At 7 days post-mTBI, IL-1β

signal was found to be colocalized with or in the vicinity of GFAP[+] cells (presumably reactive astrocytes)

and with Iba1[+] cells (likely microglia) (Supplemental Figure 1A-D). Interestingly, at 21 dpi, the majority of

IL-1β was detected in Iba1$^+$ cells (microglia or infiltrating macrophages) suggesting a very active role of

microglia on the TBI-dependent upregulation and secretion of this cytokine (Supplemental Figure 3C-D).

Our findings thus emphasize the crucial role of inflammasome signaling, with ASC as the common adaptor

protein, in sustaining the expression and processing of neuroinflammatory mediator after CHI. Moreover,

we identified Iba1$^+$ cells, likely microglia, or infiltrating macrophages, as the primary cellular sources of IL-

1β, especially at later time points post-injury.

ASC shapes the morphology of Iba1$^+$ cells following CHI. Microglia exhibit distinct adaptive morphologies

in response to mechanical stimuli, and their morphology is considered a proxy for their function (36). To

investigate microglia architecture after CHI, we performed immunofluorescence staining for Iba1 and

performed skeleton analysis (37) to examine cell morphology in cortices of WT and *Asc$^{-/-}$* mice at 1, 7,

and 21 dpi (Figure 3A–B). First, we observed that *Asc$^{-/-}$* mice consistently exhibited lower Iba1+ cell

counts compared with WT mice at 7 dpi (Figure 3C), suggesting a decreased microgliosis in the

subacute CHI phase. Next, we quantified the normalized changes in the morphology of Iba1+ cells based

on cell count. The morphological alterations, including the number of branches per cell, endpoints per

cell, and branch length per cell, significantly decreased after injury in WT mice, particularly at 21 dpi. In

contrast, *Asc$^{-/-}$* mice showed a significant preservation of these morphological features at 21 dpi

compared to WT mice (Figure 3D-F). In conclusion, our observations suggest a wave of microgliosis in

the subacute phase and revealed a progressive debranching of Iba1+ cells over time following injury.

Notably, the lack of ASC demonstrated protection against these sustained cellular morphological

alterations, particularly at later time points.

ASC regulates the reactivity of GFAP$^+$ cells following CHI. Previous studies have provided evidence that

morphological changes in astrocytes are intricately linked to microglia activation during inflammation-

induced responses (38). To examine reactive astrocytes and their morphological alterations following mild

CHI, we performed a morphological analysis of GFAP$^+$ cells in contusional layers 1 and 2/3 of the

retrosplenial and primary motor cortices, using the same method as for Iba1+ cells. (Figure 3H-I). We

initially observed a noteworthy increase in the count of GFAP$^+$ cells at 1, 7 and 21 dpi in WT mice (Figure

3j), confirming increased reactive astrogliosis in our model. *Asc* deficiency exerted a genotype-related

effect in the cell count similar to what we observed in Iba1$^+$ cells, showing significantly lower numbers,

especially at 7 dpi (Figure 3j). Subsequently, we quantified normalized morphological changes in GFAP$^+$

cells based on cell counts. The branches per cell, endpoints per cell, and length of branches per cell were

significantly decreased after CHI, particularly at 21 dpi. In contrast, *Asc$^{-/-}$* mice show preserved

morphological features at 21 dpi compared to WT mice (Figure 3K–M). Interestingly, at 21 dpi, we

observed clusters of aggregated GFAP+ cells, indicating spatial reorganization post-injury. Similar

morphological changes were evident in both microglia and reactive astrocytes after injury, suggesting a

potential correlation or interactions between these two glial cell populations.

Iba$^+$ and GFAP$^+$ cell interactions are modulated by ASC following CHI. To investigate the interaction

between reactive astrocytes and microglia in the contusional cortex following injury, we performed

surface–surface colocalization analysis of the contact areas after 3D reconstruction of confocal images

from GFAP- and Iba1-immunostained tissue (Figure 4A). We observed a significant increase in contact

area between reactive astrocytes and microglia at 7 dpi, reaching a peak at 21 dpi in WT mice (Figure

4B). Notably, the contact area was significantly reduced at 7 and 21 dpi in *Asc$^{-/-}$* mice (Figure 4B) which

may be explained by the deficient microglia activation and reactive astrogliosis observed in our previous

analysis. These findings, together with our expression analysis, suggest that inflammasome ensembles

and neuroinflammatory processing are essential for sustained microglial and astrocytic responses and

cellular interactions after mild CHI.

Genetic deficiency in the Asc gene provides protection against mild cognitive impairments following.

Having demonstrated that *Asc* is predominantly expressed by microglia in the subacute phase of CHI,

contributing to the sustained inflammasome activation, as well as chronic microglial and astrocytic

reactivity following CHI, we next sought to investigate whether the genetic removal of the main

inflammasome adaptor protein ASC influences cognitive outcomes after mild CHI. In humans, a

substantial number of individuals present transient or persistent cognitive impairment following mTBI,

including deficits in memory recall, procedural tasks, executive functions, attention and emotional

symptoms such as heightened anxiety (39). To investigate how ASC contribute to these neurobehavioral

phenotypes, we performed Morris Water Maze (MWM) as main behavioral task in WT and *Asc$^{-/-}$* mice after

CHI. MWM was conducted from days 10 to 20 after CHI (10-20 dpi) and consisted in a spatial learning

phase (10 to 15 dpi), a 24-hour long-term spatial memory test (16 dpi), and a cued learning phase as

control procedure between 17 and 20 dpi (Figure 5A). In comparison to the WT sham group, WT mice

subjected to CHI learned the task with comparable daily reduction in path lengths towards the hidden

platform (Figure 5B). However, CHI induces significantly prolonged swimming time near the walls

(thigmotaxis) during the learning phase in WT injured mice (Figure 5D). We observed similar tendency in

thigmotaxis during the cued version of the task (Figure 5H). This procedural deficit was largely reduced in

$Asc^{-/-}$ mice (Figure 5D-E and 5H-I). Additionally, $Asc^{-/-}$ mice demonstrated preserved long-term spatial

memory 24 hours after the learning phase, as visually suggested by the density plots (heat maps) of swim

paths (Figure 5J). This finding was further supported by a significantly greater percentage of time spent

exploring the target quadrant (Figure 5K), as well as more frequent crossings of the virtual target platform

(Figure 5I) compared to the corresponding measures in another quadrant or at other virtual platforms

(Figure 5L) during the probe trial performed at 16 dpi. Furthermore, $Asc^{-/-}$ mice displayed slightly higher

velocity in the last day of cued learning (Supplemental Figure 4A and B). These data show that CHI

induces mild cognitive impairments, mainly as procedural and memory deficits in the MWM, which are

mostly dependent on ASC expression.

NLRP3 regulates ASC aggregation and distribution following CHI. The NLRP3 is widely recognized as one

of the most relevant inflammasome activator in the context of TBI, with growing evidence indicating its

central role in mediating neuroinflammation and secondary brain damage (40). To explore the involvement

NLRP3 in the aggregation of ASC as the main inflammasome adaptor in post-CHI, we conducted

immunohistochemical staining of the contusional cortical tissue for Iba1, GFAP and ASC at 7 and 21 dpi in

in WT and $Nlrp3^{-/-}$ mice (Figure 6A). We used a KO-validated monoclonal antibody for ASC detection

(Supplemental Figure 5A-Bb). 3D surface reconstruction of Iba1, GFAP and ASC immunosignal analysis

were subsequently conducted (Figure 6A and Supplemental Figure 5C). We detected a significant

increase in the total number of fluorescent ASC aggregates, especially in Iba1$^+$ cells (likely microglia) at 7

dpi similar to the mRNA expression of *Asc* observed in the scRNA-Seq data analysis following mFPI

(Figure 1B, Figure 6B and Supplemental Figure 5D). The number of ASC aggregates within Iba1$^+$ cells

followed the total ASC aggregates pattern (Figure 6B and Supplemental Figure 5C-D). Notably, $Nlrp3^{-/-}$

injured mice showed comparable ASC aggregates counts, both overall and within Iba1$^+$ cells, compared to

WT mice (Figure 6b) but significantly lower aggregates in GFAP$^+$ cells (presumably reactive astrocytes,

Figure 6c). Interestingly, at 21 dpi, the number of ASC aggregates outside Iba1$^+$ and GFAP$^+$ cells (extra-

glial ASC) significantly increased in both WT and *Nlrp3$^{-/-}$* mice (Figure 6d). However, this ensembles count

was significantly lower in *Nlrp3$^{-/-}$* mice compared to WT controls (Figure 6d). Additionally, as ASC

aggregates numbers increased within microglia, their corresponding volumes also increased. To further

analyze ASC aggregation, we categorized ASC aggregates by volume (1–20 μm^3, 20–30 μm^3, 30–40

μm^3) and quantified the number of ensembles in each category (Supplemental Figure 5E-F and Figure

6E). ASC aggregates within a range of 30–40 μm^3 were significantly decreased in the *Nlrp3$^{-/-}$* mice at 21

dpi (Figure 6E). Collectively, our findings demonstrate a remarkable upregulation of ASC aggregates

outside of Iba1$^+$ and GFAP$^+$ cells, along with marked ASC aggregation at 21 dpi. *Nlrp3$^{-/-}$* mice displayed

reduced ASC aggregation within Iba1$^+$ cells and fewer extra-glia ASC aggregates compared to WT group

at 21 dpi. These observations highlight the essential role of NLRP3 not only in promoting inflammasome

activation after brain injury but also in facilitating significant ASC aggregation, as previously suggested in

in vitro studies (41).

Interaction between reactive astrocytes and microglia correlates with ASC distribution following CHI. To

further investigate the relationship between reactive astrocytes and microglia interactions and ASC

distribution following CHI, we conducted correlation analysis between ASC aggregates and glial contact

areas after 3D reconstruction (Figure 6 F–I). Notably, we observed a strong positive correlation between

the cell contact area and the number of extra-glia ASC aggregates *(r = 0.86, p < 0.0001)* (Figure 6g) as

well as ASC aggregates within GFAP$^+$ cells *(r = 0.93, p < 0.0001)* (Figure 6I), in WT animals after CHI.

Although lower correlations coefficients were found between the reactive astrocytes and microglia contact

areas and the number of ASC aggregates in microglia *(r = 0.73, p < 0.0001)* (Figure 6h) and the overall

count of ASC aggregates *(r = 0.82, p < 0.0001)* (Figure 6f), these correlations remained significantly

positive (Figure 6F–I). Interestingly, the genetic deficiency of the *Nlrp3* gene significantly reduced the

correlation between the contact area and the number of extra-glia ASC aggregates *(z = 1.95, p < 0.05)*

(Figure 6G) as well as ASC in GFAP$^+$ cells *(z = 2.54, p < 0.01)* (Figure 6I). In summary, these results

indicate that ASC aggregation after CHI is partly dependent on NLRP3 and additionally reactive astrocytes

and microglia contacts are influenced by ASC expression, distribution, and aggregation, especially the

ASC aggregates located outside of Iba1$^+$ and GFAP$^+$ cells.

Discussion

In this study, we identify *Asc* as a key gen expressed in monocytes, macrophages, and especially in

microglia populations during the subacute phase of CHI based on scRNA-seq data from mouse cortical

cells after mFPI. Similarly, in our mild CHI mouse model, sustained ASC protein expression was

confirmed, driving reactivity of microglia and astrocytes, and innate immune responses that contribute to

neurobehavioral deficits following mTBI. Our findings not only complement and expand previous

preclinical and clinical studies on moderate to severe TBI models (42, 43), which highlight the role of

inflammasome cascades and their critical involvement in regulating secondary inflammatory responses

after TBI, but also demonstrate for the first time that the main adaptor protein ASC is a key element in

neuroinflammatory responses and cognitive impairment following mTBI. In contrast to several studies on

moderate-to-severe TBI models, where the expression of inflammasome-related proteins typically peaks

within the initial week following injury and declines over time (21, 44, 45), we found a progressive and

sustained increase in inflammasome-related proteins specially after the first 7 days following mild CHI.

This temporal pattern aligns with our observations of sustained morphological changes of reactive

astrocytes and microglia during a later phase (21 dpi). These discrepancies between models and TBI

types suggest that inflammasome engagement in innate immune responses and tissue repair varies

depending on injury severity. Moreover, the prolonged pro-inflammatory characteristics exhibited by

activated microglia and reactive astrocytes indicate that both glial cell types are capable of propagating

inflammation through inflammasome-related pathways post-injury. Interestingly, our investigation

demonstrated that astrocytes display more significant reactive morphology at 21 dpi compared to 7 dpi.

This may result from the potential further induction of neurotoxic astrocytes by microglial inflammasome in

response to sustained tissue damage, as previously reported in stress models (46). Furthermore, these

findings align with existing evidence suggesting that distinct glial cell types respond differentially to brain

injury in a temporal sequence (37, 47–49).

Additionally, we observed a widespread upregulation of inflammasome components following CHI,

particularly at 21 dpi, with a significant increase in the cleaved forms of CASP1, CASP8 and IL-1β. Our

results suggest the involvement of several ASC-dependent inflammasomes in the inflammatory pathways

after CHI. This is evident by the pronounced reduction in CASP1, CASP3 and CASP8 levels and

morphological changes of microglia and reactive astrocytes observed in *Asc*[-/-] mice. Most notably, our

behavioral evaluation during the late subacute phases shows that anxiety-related behaviors (thigmotaxis),

mild procedural learning deficits and long-term memory impairments were significantly diminished in

animals with genetic deficiency on the *Asc* gene. Furthermore, our analysis established a significant

positive correlation between reactive astrocyte and activated microglia interaction and the number of ASC

aggregates. Notably, a potential connection has been identified between extracellular ASC and cellular

propagation of inflammatory mediators, including IL-1β and TNF-α following inflammasome activation (50,

51). These cytokines not only promote the aggregation of glia and infiltrated cells around the trauma

epicenter (52, 53), but may also influence cell communication far from the lesion site through extracellular

ASC. ASC possess prion-like properties (51) and could propagate to neighboring glial cells, thereby

triggering the activation of the ASC-mediated inflammasome cascade as previously shown in the context

of Alzheimer's disease- related amyloidosis (54, 55). These observations suggest a pivotal role of ASC in

sustaining neuroinflammation and glial reactivity following CHI.

Our findings, however, contrast with a previous report on moderate CCI, where *Asc*[-/-] mice showed no

differences in motor recovery or volume lesions (56). This inconsistency may suggest that, in the context

of low-grade chronic neuroinflammation induced by mild, uncontrolled stimuli, such as a single concussion

event, the expression, oligomerization, and propagation of ASC may occur gradually compared to the

high-grade neuroinflammation triggered by severe injuries. Similarly, ASC-dependent inflammasome

activation and pathological protein aggregation have been observed in several neurodegenerative

disorders characterized by sustained chronic low-grade neuroinflammation (57), such as Alzheimer's and

Parkinson's disease, where ASC plays a decisive role in the aggregating and further propagating Amyloid

β (54) and α-Synuclein (58). Considering our findings, ASC may govern the immune molecular

mechanism that triggers neurodegeneration and increase the risk of developing neurodegenerative

conditions following TBI (59). Nonetheless, future investigations are still required to establish a

comprehensive understanding of the specific temporal participation of inflammasomes across varying

injuries severity and the risk factor for neurodegeneration.

There are certain limitations that should be acknowledged in our study. Focusing on the expression of
inflammasome mediators and cell morphological characteristics may not fully capture the complex
interactions with other compensatory mechanisms within the affected neuronal networks or the further
anti-inflammatory modulation of secondary immune responses. Additionally, although our scRNA-Seq
analysis determined specific cellular clusters upregulating *Asc* in the subacute phase of a CHI model, our
study using conventional knockout line cannot dissect precisely the role of different peripheral and cellular
actors in the immunological process following CHI. Thus, there is an imperative need to investigate the
neuroinflammatory evolution of the specific neural, glial and peripheral cellular subtracts over extended
post-traumatic periods.

Our findings pave the way to delve into specific pharmacological agents that can effectively modulate ASC
expression, activity and aggregation and assess their impact on neurological outcomes after TBI.
Considering our findings, pharmacological interventions targeting ASC may help to mitigate
neuroinflammation and potentially promote neuroprotection, enhancing recovery of injury and preventing
further neurodegeneration and disability.

340 **Methods**

Ethical compliance. We followed the European guidelines for animal research, conformed to the
requirements of the German Animal Welfare Act and received approval from North Rhine-Westphalia
State Agency for Nature, Environment and Consumer Protection under folder number 81-
02.04.2019.A026 and 2024-791-Grundantrag.

Single-cell RNA-sequencing data analysis. Raw FASTQ files and count matrices are publicly available in
the NCBI GEO database under accession number GSE160763 (26). Single-cell RNA-seq data were
analyzed using Seurat v5.1.0 in R v4.3.2 (60). Briefly, cells with over 5000 detected genes, UMI counts
under 12,500, and less than 1% mitochondrial gene content were retained, followed by log-transformation
of the data. We selected the top 4000 highly variable genes for principal component analysis (PCA) and
used the top 20 principal components as input for UMAP dimensionality reduction. Cell cycle states were
determined using Seurat's CellCycleScoring (61) function with marker genes from Tirosh *et al.* (60). Batch
correction was performed using Harmony to integrate datasets across samples (62). Clusters were

identified at a resolution of 0.5. Cellular identities were assigned via automated cell type annotation using

SingleR (29), which assigns cellular identity for single cell transcriptomes by comparison to reference data

sets of pure cell types sequenced by microarray or RNA-sequencing (RNA-seq). Annotation relied on a

reference dataset of 358 mouse RNA-seq samples annotated to 18 major cell types (63). Figures were

generated using Tidyverse(64), SCpubr(65), and Enhanced Volcano (66).

Mice. In this study, we used male and female mice aged 6–7 months with a C57BL/6 mixed genetic

background (C57BL/6N-J) from our own animal facility (House of the Experimental Medicine) at the

University Hospital Bonn in German. Wild-type (WT), *Nlrp3$^{-/-}$*, and *Asc$^{-/-}$* mice were initially sourced from

Millennium Pharmaceuticals as previously published (67). All mice were backcrossed with C57BL/6N mice

and maintained under pathogen-free, standardized conditions with a 12:12-hour light/dark cycle and ad

libitum access to food and water. Behavioral experiments were conducted during the dark phase of the

light/dark cycle.

Electromagnetic controlled closed-head injury model of mild traumatic brain injury. To model mild TBI in

mice**,** we adapted the electromagnetic controlled Closed-Head Model previously described (31)

(Supplemental Figure 2A). Briefly, mice were anesthetized with 5% isoflurane during induction, and 1.5-

2% isoflurane mixed with 100% oxygen (0.5- 1 l/min) during surgery and fixed in a stereotaxic frame

(Stoelting, Dublin, Ireland). A 1-mL latex pipette bulb filled with water was placed under the head to

distribute the impact force. A midline sagittal scalp incision was made, and a single controlled midline skull

impact was delivered at coordinates 0.0 mm mediolateral and -1.5 mm anteroposterior, with a velocity of 5

m/s, dwell time of 100 ms, and impact depth of 1.0 mm, using a stereotaxic electromagnetic impactor with

a 5.0-mm steel tip (Stereotaxic Impactor, Leica Biosystems, Germany). Sham mice underwent identical

surgical procedures without impact injury. In the post-operative phase, mice received carprofen (5 mg/kg)

subcutaneously once daily and tramadol (1 mg/mL) in drinking water for three days.

Morris water maze (MWM). The evaluation of spatial learning and memory was conducted between 10

and 20 dpi. A circular pool with a diameter of 1 m was filled with white opacified water maintained at a

temperature of 21–23 °C. The pool was dimly lit (approximately 40 lx) and surrounded by a white curtain.

Asymmetrically placed distal cues were positioned on the pool wall to serve as spatial references. The

pool was virtually divided into four quadrants, one of which contained a hidden platform (15 cm in

diameter) submerged 1.5 cm below the water surface. Mice were trained to locate the platform using the

distal cues for orientation. Training consisted of four trials per day for five consecutive days. During each

trial, mice were placed into the water facing the pool wall in a quasi-randomized order to prevent them

from developing fixed strategies. Mice were given 60 s to find the platform; if they failed to locate it within

this time, they were manually guided to it. Once on the platform, mice were allowed to remain there for 15

s before the next trial began. After completing all four trials for the day, mice were dried and returned to

their home cages. The integrated time or distance traveled during these trials was analyzed. A spatial

probe trial was conducted 24 hours after the final training session (16 dpi). For this test, the hidden

platform was removed, and mice were allowed to swim freely for 60 s. The drop position was in the

quadrant opposite where the platform had been located, with each mouse facing the wall at the center of

that quadrant. One day after the spatial probe trial, a visually cued learning phase began. During this

phase, the platform was flagged for visibility and relocated to a different quadrant, while all distal cues

were removed. Mice underwent three trials per day for three consecutive days (17–20 dpi) to learn to

locate the flagged platform. All mouse movements were recorded and tracked using Noldus EthoVision

software (Wageningen, Netherlands).

Tissue preparation. Mice were deeply anesthetized using ketamine (100 mg/kg) and xylazine (20 mg/kg)

and transcardially perfused with at least 30 ml of ice-cold PBS at designated time points (Sham, 1, 7, and

21 dpi). Following perfusion, brains were carefully dissected. One hemisphere was snap-frozen in liquid

nitrogen and stored at -80°C until biochemical analyses, while the other hemisphere was fixed in 4%

paraformaldehyde overnight. Coronal brain sections (40 μm thick) were obtained using a Leica VT1000S

vibratome (Leica Microsystems Inc.) from the region spanning bregma -0.45 to -1.85 mm. Sections were

collected in PBS at intervals of 300 μm for various staining procedures.

Primary antibodies. For immunoblotting, following primary antibodies were used: Anti-Caspase 1 from

AdipoGen (USA, mAb Casper-1, mouse anti-mouse, 1:1000), anti-ASC from Cell Signaling (Frankfurt,

Germany, D2W8U, rabbit anti-mouse, 1:1500), anti-NLRP3 from Cell Signaling (Frankfurt, Germany,

D4D8T, rabbit anti-mouse, 1:500), anti-IL-1β from Cell Signaling (Frankfurt, Germany, E7V2A, rabbit anti-

mouse, 1:1000), anti-Caspase 3 from Cell Signaling (Frankfurt, Germany, D3E9, rabbit anti-mouse,

1:1000), and anti-Caspase 8 from Cell Signaling (Frankfurt, Germany, D5B2, rabbit anti-mouse, 1:1000).

For immunohistochemical analysis, the antibodies included anti-Iba1 from Wako Chemicals (Neuss,

Germany, rabbit anti-mouse, 1:1000), anti-IBA1 from Abcam (Cambridge, UK, ab289874, goat anti-

mouse, 1:1000), anti-GFAP from ThermoFisher (Waltham, USA, 2.2B10, rat anti-mouse, 1:300), anti-IL-1β

from R&D Systems (Minneapolis, USA, AF-401-SP, goat anti-mouse, 1:500), and anti-P53 from Abcam

(Cambridge, UK, ab131442, rabbit anti-mouse, 1:500).

Tissue protein extraction and immunoblotting. After thawing, brain tissue was homogenized in PBS

containing 1 mM EDTA, 1 mM EGTA, and protease inhibitor cocktail Halt™ (Thermo Scientific, Waltham,

Massachusetts, U.S.). The homogenates were extracted in RIPA buffer [25 mM Tris–HCl (pH 7.5), 150

mM NaCl, 1% Nonidet P-40, 0.5% NaDOC, 0.1% SDS] and centrifuged at 20000 ×g for 30 minutes. RIPA

fractions were separated using NuPAGE® electrophoresis gels (Thermo Scientific, Waltham,

Massachusetts, USA) and immunoblotted with primary antibodies, followed by incubation with the

corresponding secondary antibodies. Immunoreactivity was detected with an Odyssey CLx Imaging

System (LI-COR, Bad Homburg, Germany), and images were analyzed using ImageJ (NIH, Bethesda,

USA).

ELISA pro-inflammatory cytokine panel. The levels of IL-1β and TNF-α in RIPA fractions were measured

using the V-PLEX Plus Mouse Pro-Inflammatory Panel 1 (Meso Scale Discovery, Rockville, USA),

following the manufacturer's protocol. Samples were diluted 1:1 on the plate with the reagent diluent

provided in the kit. Signals were detected using a QuickPlex SQ 120 plate reader (Meso Scale Discovery,

Rockville, USA).

Immunohistochemistry. Brain sections were washed three times for 5 min with PBS and incubated in

citrate buffer for 5 min at 95°C. After antigen retrieval, the sections were cooled down at room temperature

and washed with PBS and in 0,5% Triton X-100 B (PBS-T), then blocked for 1 h with BSA 1% in PBS-T

and incubated overnight with the primary antibodies. Next, the sections were washed three times for 5 min

in PBS-T, incubated with appropriate secondary antibodies (1:1000) for 60 min, and following with three

times with PBS washing for 5 min. The tissue was mounted using ProLong™ Gold Antifade Mountant with

DNA Stain DAPI (Thermo Scientific, Waltham, Massachusetts, U.S.). Three coronal brain sections per

animal (between bregma −1.00 and −1.85) were selected. Confocal z-stack images of the contusion were

primarily acquired from layers 1 and 2/3 of the retrosplenial and primary motor cortex. Each z-stack

consisted of 10 optical slices (c × z × t = 10), with one frame captured per acquisition. Imaging parameters

were standardized across all samples: the pixel size was set to 0.1559814 μm in both the x and y axes,

and the voxel depth (z-step) was maintained at 2.0 μm.

Skeleton analysis of microglia and reactive astrocytes. Morphological changes in microglia and reactive

astrocytes were quantitatively analyzed using a skeleton analysis method (68). Images were imported into

ImageJ software (Bethesda, Maryland, USA), converted to grayscale, and adjusted for brightness and

contrast to enhance cell process visibility. After obtaining binary images of microglia and reactive

astrocytes, we used the plugin skeletonize to obtain skeletons, representing the medial axis of cell

processes. Morphological parameters measured included: Process Branches, defined as number of

branching points. Endpoints, defined as number of terminal points. Process Lengths, defined as total

length of all processes. The data were normalized by dividing the measurements by the number of cells in

each image.

Contact analysis between microglia and reactive astrocytes. Imaris Image Analysis Software (version 9.5,

Oxford Instruments, Oxon, England) was used to analyze the contact area between microglial and

astrocytic surfaces. 3D reconstructions were created using the tool surfaces by setting intensity thresholds

to differentiate cell surfaces from the background. Adjustments were made to ensure accurate cell

boundary representation. The Surface-Surface Contact X-Tension identified and analyzed overlapping

regions between microglial and astrocytic surfaces, generating a new surface to represent the contact

area. The contact area (μm^2) was calculated based on the number of overlapping voxels within the 3D

reconstructions.

Analysis of ASC aggregates. The cortical impact area was examined using a confocal LSM900

microscope (Zeiss, Oberkochen, Baden-Württemberg, Germany). Images were captured under consistent

confocal settings and processed with Imaris Image Analysis Software (Oxford Instruments, Oxon,

England) to identify ASC aggregates. Standardized settings were maintained across samples, including

pinhole size, laser intensity, digital gain, and offset. Z-stacks were collected to obtain the complete 3D

structure of cells and ASC aggregates. Image preprocessing included adjusting brightness and contrast,

reducing noise, and subtracting background to enhance aggregate visibility. Imaris Image Analysis

Software was used to create 3D reconstructions of microglia, astrocytes, and ASC aggregates. The

"Surfaces" tool generated accurate 3D models by setting intensity thresholds. The Spots tool identified

and quantified ASC aggregates based on fluorescence intensity and size, with thresholds set using

positive (mCherry-ASC mice) and negative (*Asc*[-/-] mice) controls. The Spots Close to Surface tool

analyzed the spatial distribution of ASC aggregates relative to microglia or astrocytes, indicating potential

inflammasome activation.

IL-1β analysis. Co-immunostainings for IL-1β, Iba1 and GFAP were performed at 7 and 21 dpi. Three-

dimensional reconstructions of microglia, reactive astrocytes, neuronal nuclei, and IL-1β signals were

generated using Imaris Image Analysis Software (Oxford Instruments, Oxon, UK). The "Spots" tool

quantified IL-1β signals in the 3D space by analyzing fluorescence intensity and particle size thresholds.

To assess spatial relationships, the "Spots Close to Surface" tool measured the proximity of IL-1β signals

to cellular surfaces, evaluating potential associations with microglia, astrocytes, or neurons. Quantitative

metrics, including the number of IL-1β spots adjacent to specific cell types, were measured for statistical

analysis.

Statistics and reproducibility. Data other than scRNA-seq were analyzed using GraphPad Prism Software

version 9.0 or later (GraphPad Software, Inc., La Jolla, CA, USA) and are presented as mean ± SEM.

After assessing the distribution, statistical comparisons were performed using either a two-tailed t-test or

one- or two-way ANOVA with Bonferroni's or Tukey's post hoc test to identify group differences. Morris

Water Maze (MWM) learning data were analyzed using repeated-measures two-way ANOVA with Tukey's

post hoc test. Correlation analyses were performed using non-parametric Spearman correlations.

Differences between Spearman correlation coefficients were assessed using the standard Fisher's z-

transformation and subsequent comparison(69). Results were considered statistically significant at $p <$

0.05.

Data availability. The data that supports the findings of this study are available on request from the

corresponding authors upon reasonable request.

*Code availability. p*Code used in analyzing scRNA-seq data is available at

https://gitlab.lcsb.uni.lu/pablo.botellalucena/rna_single_seq.

Author contributions

493   Conceptualization: SCG, MTH. Methodology: ASV, PBL, SCG, TL, SS, YD. Investigation: ASV, PBL,

494   SCG, TL, SS, YD. Funding acquisition: SCG, MTH. Supervision: SCG, MTH. Writing of original draft:

495   SCG, TL, YD. Writing, review, and editing: MTH, SCG, PB, TL, YD, YD, VS, EL

Acknowledgements

We thank Ildikó Rácz and Christina Ising for their help in obtaining approval from the local ethics committee for the animal experiments; Sean-Patrick Hermann for assistance with editing the final draft; Juan I. Muñoz, Paula Martorell, and Nadia Villacampa for valuable discussions; and the DZNE Light Microscopy Facility for support with imaging and analysis. PBL is supported by the Fonds National de la Recherche within the PEARL program (FNR/16745220). SCG is supported by the Alzheimer Forschung Initiative e.V (Grant 21060), the Hertie Network of Excellence in Clinical Neuroscience (Grant 2021-1A-12), the Bonn Neuroscience Clinician Scientist Program (Neuro-aCSis), and the DFG (EXC2151–390873048). TL was supported by Department of Biotherapy Cancer Center and State Key Laboratory of Biotherapy, West China Hospital, Sichuan University, Chengdu, China. Y.D. is supported by China Scholarship Council (CSC, File No. 202308310126). Y.D. is supported by China Scholarship Council (CSC, File No. 201908510162).

References

1. Maas AIR, et al. Traumatic brain injury: integrated approaches to improve prevention, clinical care, and research. *Lancet Neurol*. 2017;16(12):987–1048.

2. Thurman DJ, et al. Traumatic Brain Injury in the United States: A Public Health Perspective: *J Head Trauma Rehabil*. 1999;14(6):602–615.

3. Cassidy JD, et al. Incidence, risk factors and prevention of mild traumatic brain injury: results of the who collaborating centre task force on mild traumatic brain injury. *J Rehabil Med*. 2004;36(0):28–60.

4. Naumenko Y, et al. Mild traumatic brain injury as a pathological process. *Heliyon*. 2023;9(7):e18342.

5. Rathbone ATL, et al. A review of the neuro- and systemic inflammatory responses in post concussion symptoms: Introduction of the "post-inflammatory brain syndrome" PIBS. *Brain Behav Immun*. 2015;46:1–16.

6. Ahman S, et al. Long-term follow-up of patients with mild traumatic brain injury: a mixed-method study. *J Rehabil Med*. 2013;45(8):758–764.

7. Déry J, et al. Prognostic factors for persistent symptoms in adults with mild traumatic brain injury: an overview of systematic reviews. *Syst Rev*. 2023;12(1):127.

8. Burda JE, Bernstein AM, Sofroniew MV. Astrocyte roles in traumatic brain injury. *Exp Neurol.* 2016;275 Pt 3(0 3):305–315.

9. Loane DJ, Kumar A. Microglia in the TBI brain: The good, the bad, and the dysregulated. *Exp Neurol.* 2016;275 Pt 3(0 3):316–327.

10. Simon DW, et al. The far-reaching scope of neuroinflammation after traumatic brain injury. *Nat Rev Neurol.* 2017;13(9):572.

11. Perez-Polo JR, et al. Inflammatory consequences in a rodent model of mild traumatic brain injury. *J Neurotrauma.* 2013;30(9):727–740.

12. Weil ZM, Gaier KR, Karelina K. Injury timing alters metabolic, inflammatory and functional outcomes following repeated mild traumatic brain injury. *Neurobiol Dis.* 2014;70:108–116.

13. Lv W, et al. mTBI-Induced Systemic Vascular Dysfunction in a Mouse mTBI Model. *Brain Sci.* 2022;12(2):232.

14. Ravichandran KA, Heneka MT. Inflammasomes in neurological disorders - mechanisms and therapeutic potential. *Nat Rev Neurol.* 2024;20(2):67–83.

15. Franklin BS, Latz E, Schmidt FI. The intra- and extracellular functions of ASC specks. *Immunol Rev.* 2018;281(1):74–87.

16. Pérez-Bárcena J, et al. Serum Caspase-1 as an Independent Prognostic Factor in Traumatic Brain Injured Patients. *Neurocrit Care.* 2022;36(2):527–535.

17. Kerr N, et al. Inflammasome proteins as biomarkers of traumatic brain injury. *PloS One.* 2018;13(12):e0210128.

18. Eagle SR, et al. Impact of obese body mass index on inflammasome blood biomarkers and neurocognitive performance following traumatic brain injury with Glasgow coma scale 13 to 15. *J Neurol Sci.* 2024;464:123159.

19. de Rivero Vaccari JP, et al. Therapeutic neutralization of the NLRP1 inflammasome reduces the innate immune response and improves histopathology after traumatic brain injury. *J Cereb Blood Flow Metab Off J Int Soc Cereb Blood Flow Metab.* 2009;29(7):1251–1261.

20. Irrera N, et al. Lack of the Nlrp3 Inflammasome Improves Mice Recovery Following Traumatic Brain Injury. *Front Pharmacol.* 2017;8:459.

21. Liu H-D, et al. Expression of the NLRP3 inflammasome in cerebral cortex after traumatic brain injury in a rat

model. *Neurochem Res.* 2013;38(10):2072–2083.

22. Adamczak SE, et al. Pyroptotic neuronal cell death mediated by the AIM2 inflammasome. *J Cereb Blood Flow*

Metab Off J Int Soc Cereb Blood Flow Metab. 2014;34(4):621–629.

23. Brickler T, et al. Nonessential Role for the NLRP1 Inflammasome Complex in a Murine Model of Traumatic Brain

Injury. *Mediators Inflamm.* 2016;2016:1–11.

24. Vincent JC, et al. IL-1R1 signaling in TBI: assessing chronic impacts and neuroinflammatory dynamics in a mouse

model of mild closed-head injury. *J Neuroinflammation.* 2023;20(1):248.

25. Bachstetter AD, et al. Attenuation of traumatic brain injury-induced cognitive impairment in mice by targeting

increased cytokine levels with a small molecule experimental therapeutic. *J Neuroinflammation.* 2015;12:69.

26. Witcher KG, et al. Traumatic Brain Injury Causes Chronic Cortical Inflammation and Neuronal Dysfunction

Mediated by Microglia. *J Neurosci Off J Soc Neurosci.* 2021;41(7):1597–1616.

27. Macheda T, Roberts K, Bachstetter AD. Electromagnetic Controlled Closed-Head Model of Mild Traumatic Brain

Injury in Mice. *J Vis Exp.* 2022;(187):64556.

28. Lifshitz J, et al. Clinical relevance of midline fluid percussion brain injury: Acute deficits, chronic morbidities and

the utility of biomarkers. *Brain Inj.* 2016;30(11):1293–1301.

29. Aran D, et al. Reference-based analysis of lung single-cell sequencing reveals a transitional profibrotic

macrophage. *Nat Immunol.* 2019;20(2):163–172.

30. Belinky F, et al. PathCards: multi-source consolidation of human biological pathways. *Database J Biol Databases*

Curation. 2015;2015:bav006.

31. Macheda T, Roberts K, Bachstetter AD. Electromagnetic Controlled Closed-Head Model of Mild Traumatic Brain

Injury in Mice. *J Vis Exp JoVE.* [published online ahead of print: September 28, 2022];(187).

https://doi.org/10.3791/64556.

32. Yarnell AM, et al. The Revised Neurobehavioral Severity Scale (NSS-R) for Rodents. *Curr Protoc Neurosci.*

2016;75:9.52.1-9.52.16.

33. Antonopoulos C, et al. Caspase-8 as an Effector and Regulator of NLRP3 Inflammasome Signaling. *J Biol Chem*.

2015;290(33):20167–20184.

34. Clark RS, et al. Caspase-3 mediated neuronal death after traumatic brain injury in rats. *J Neurochem*.

2000;74(2):740–753.

35. Bertheloot D, Latz E, Franklin BS. Necroptosis, pyroptosis and apoptosis: an intricate game of cell death. *Cell Mol*

Immunol. 2021;18(5):1106–1121.

36. Green TRF, Rowe RK. Quantifying microglial morphology: an insight into function. *Clin Exp Immunol*.

2024;216(3):221–229.

37. Morrison H, et al. Quantitative microglia analyses reveal diverse morphologic responses in the rat cortex after

diffuse brain injury. *Sci Rep*. 2017;7(1):13211.

38. Liddelow SA, et al. Neurotoxic reactive astrocytes are induced by activated microglia. *Nature*.

2017;541(7638):481–487.

39. McCauley SR, et al. Patterns of Early Emotional and Neuropsychological Sequelae after Mild Traumatic Brain

Injury. *J Neurotrauma*. 2014;31(10):914–925.

40. Chakraborty R, Tabassum H, Parvez S. NLRP3 inflammasome in traumatic brain injury: Its implication in the

disease pathophysiology and potential as a therapeutic target. *Life Sci*. 2023;314:121352.

41. Hochheiser IV, et al. Directionality of PYD filament growth determined by the transition of NLRP3 nucleation seeds

to ASC elongation. *Sci Adv*. 2022.

42. Hu X, et al. Role of Pyroptosis in Traumatic Brain and Spinal Cord Injuries. *Int J Biol Sci*. 2020;16(12):2042–2050.

43. Mortezaee K, et al. Inflammasome: Its role in traumatic brain and spinal cord injury. *J Cell Physiol*.

2018;233(7):5160–5169.

44. Dempsey C, et al. Inhibiting the NLRP3 inflammasome with MCC950 promotes non-phlogistic clearance of

amyloid-β and cognitive function in APP/PS1 mice. *Brain Behav Immun*. 2017;61:306–316.

45. Xu X, et al. Selective NLRP3 inflammasome inhibitor reduces neuroinflammation and improves long-term

neurological outcomes in a murine model of traumatic brain injury. *Neurobiol Dis*. 2018;117:15–27.

46. Li S, et al. Microglial NLRP3 inflammasome activates neurotoxic astrocytes in depression-like mice. *Cell Rep.*
2022;41(4):111532.

47. Witcher KG, et al. Traumatic brain injury-induced neuronal damage in the somatosensory cortex causes formation
of rod-shaped microglia that promote astrogliosis and persistent neuroinflammation. *Glia.* 2018;66(12):2719–2736.

48. Susarla BTS, et al. Temporal Patterns of Cortical Proliferation of Glial Cell Populations after Traumatic Brain Injury
in Mice. *ASN Neuro.* 2014;6(3):AN20130034.

49. Nespoli E, et al. Glial cells react to closed head injury in a distinct and spatiotemporally orchestrated manner. *Sci*
Rep. 2024;14(1):2441.

50. Franklin BS, Latz E, Schmidt FI. The intra- and extracellular functions of ASC specks. *Immunol Rev.*
2018;281(1):74–87.

51. Franklin BS, et al. The adaptor ASC has extracellular and "prionoid" activities that propagate inflammation. *Nat*
Immunol. 2014;15(8):727–737.

52. Sofroniew MV. Molecular dissection of reactive astrogliosis and glial scar formation. *Trends Neurosci.*
2009;32(12):638–647.

53. Wang H, et al. Portrait of glial scar in neurological diseases. *Int J Immunopathol Pharmacol.*
2018;31:2058738418801406.

54. Venegas C, et al. Microglia-derived ASC specks cross-seed amyloid-β in Alzheimer's disease. *Nature.*
2017;552(7685):355–361.

55. Friker LL, et al. β-Amyloid Clustering around ASC Fibrils Boosts Its Toxicity in Microglia. *Cell Rep.*
2020;30(11):3743-3754.e6.

56. Brickler T, et al. Nonessential Role for the NLRP1 Inflammasome Complex in a Murine Model of Traumatic Brain
Injury. *Mediators Inflamm.* 2016;2016:6373506.

57. Castro-Gomez S, Heneka MT. Innate immune activation in neurodegenerative diseases. *Immunity.*
2024;57(4):790–814.

58. Zheng R, et al. ASC specks exacerbate α☐synuclein pathology via amplifying NLRP3 inflammasome activities. *J*

Neuroinflammation. 2023;20(1):26.

59. Graham A, et al. Mild Traumatic Brain Injuries and Future Risk of Developing Alzheimer's Disease: Systematic

Review and Meta-Analysis. *J Alzheimers Dis*. 2022;87(3):969–979.

60. Tirosh I, et al. Dissecting the multicellular ecosystem of metastatic melanoma by single-cell RNA-seq. *Science*.

2016;352(6282):189–196.

61. Watson S, et al. Modification of Seurat v4 for the Development of a Phase Assignment Tool Able to Distinguish

between G2 and Mitotic Cells. *Int J Mol Sci*. 2024;25(9):4589.

62. Sun N, et al. Human microglial state dynamics in Alzheimer's disease progression. *Cell*. 2023;186(20):4386-

4403.e29.

63. Benayoun BA, et al. Remodeling of epigenome and transcriptome landscapes with aging in mice reveals

widespread induction of inflammatory responses. *Genome Res*. 2019;29(4):697–709.

64. Wickham H, et al. Welcome to the Tidyverse. *J Open Source Softw*. 2019;4(43):1686.

65. Blanco-Carmona E. Generating publication ready visualizations for Single Cell transcriptomics using SCpubr

[preprint]. 2022. https://doi.org/10.1101/2022.02.28.482303.

66. Blighe K, Rana S, Lewis M. EnhancedVolcano: Publication-ready volcano plots with enhanced colouring and

labeling [Internet]. 2018. https://github.com/kevinblighe/EnhancedVolcano.

67. Kanneganti T-D, et al. Bacterial RNA and small antiviral compounds activate caspase-1 through cryopyrin/Nalp3.

Nature. 2006;440(7081):233–236.

68. Young K, Morrison H. Quantifying Microglia Morphology from Photomicrographs of Immunohistochemistry

Prepared Tissue Using ImageJ. *J Vis Exp*. 2018;(136):57648.

69. Lenhard W, Lenhard A. Testing the Significance of Correlations [preprint]. 2014.

https://doi.org/10.13140/RG.2.1.2954.1367.

Figure 1: Subacute *Asc (Pycard)* expression in microglia after a model for CHI.

A-B. Single-cell RNA sequencing (scRNA-seq) analysis of brain suspended cells from mice 7 days after
mFPI. **A.** UMAP visualization showing clustering of different cell populations, with mFPI and control
samples overlaid. **B.** Annotated cell types, including microglia, macrophages, and other immune and
neural cells after cluster identification. **C.** Volcano plot depicting differentially expressed genes in microglia

of control and mFPI samples. **D.** Dot plot expression analysis showing percent of expressed key

inflammasome-related genes across various brain cell types, highlighting *Asc* (*Pycard*) expression in

microglia, Natural Killers (NK) cells, macrophages and monocytes. **E.** UMAP projection of cells colored

relative to the expression of canonical inflammasome-related genes *Nlrp3, Asc* (*Pycard*) *Casp1, Il1b, Il18*

and Gsdmd.

  Figure 2: **ASC contributes to the upregulation of inflammasome mediators following CHI.**

A

A. Immunoblot detection and **B-G.** densitometric semi-quantification of inflammasome-related proteins NLRP3, ASC, Caspase-1 (CASP1), IL-1β at 1, 7, and 21 dpi in contusional cortices from WT and *Asc*^-/- mice following Sham or CHI. (left panels *n* = 4 per group and time point, **p* < 0.05, ***p* < 0.01, ****p* < 0.001, *****p* < 0.0001, Ordinary two-way ANOVA, Bonferroni's test. AU, arbitrary units. **H, I.** Levels of IL-1β and TNF-α in brain lysates from WT and *Asc*^-/- mice at 1, 7, and 21 dpi following CHI. *n* = 3 - 4 per group and time point, Left panels represent ordinary two-way ANOVA, Bonferroni's test, **p* < 0.05, ***p* < 0.01, ****p* < 0.001. Right panels in **B-I** represent integrated relative concentrations (Area Under Curve, AUC), ***p* < 0.01, ****p* < 0.001 by Student's t-test. dpi, days post-injury.

674   **Figure 3: Morphology of Iba1+ and GFAP+ cells are modulated by ASC following CHI.**

676
677

A. Immunohistochemistry of Iba1 (green, presumably microglia) in cortices from WT, and *Asc*-/- mice at 1,
7, and 21 dpi following CHI. Arrows indicate representative morphological changes. Scale bar, 20 μm. **B.**
Representative skeletonized Iba1+ cells following CHI. Scale bars, 10 μm. Quantitative analysis of **C.** Iba1+
cell counts; **D.** branch numbers per cell, **E.** total branch length per cell and **F.** endpoint numbers per cell at
1, 7, and 21 dpi in WT and *Asc*-/- mice subjected to sham or CHI. **H.** Immunohistochemistry of GFAP
(magenta, presumably astrocytes) in cortices from WT and *Asc*-/- mice at 1, 7, and 21 dpi following CHI.
Arrows indicate representative morphological changes. Scale bar, 20 μm. **I.** Representative skeletonized
GFAP+ cells following CHI. Scale bars, 10 μm. Quantitative analysis of **J.** GFAP+ cell counts; **K.** branch
numbers per cell, **L.** total branch length per cell and **M.** endpoint numbers per cell at 1, 7, and 21 dpi in

WT and $Asc^{-/-}$ mice subjected to sham or CHI. n = 12 slices from 4 mice per group per each time point, *p

< 0.05, ***p < 0.01, ***p < 0.001, ****p < 0.0001 by ordinary two-way ANOVA with Bonferroni's post hoc

test. Right panels show integrative AUC of each marker over time (dpi). *p < 0.05, **p < 0.01, ***p < 0.001,

****p < 0.0001 by Student's t-test.

Figure 4: Cellular interactions of microglia and reactive astrocytes are modulated by ASC following CHI.

A

A. Immunohistochemistry for Iba1 (green) and GFAP (magenta) in cortices of WT and $Asc^{-/-}$ mice at 7 and 21 dpi. Surface-surface colocalization analysis after 3D reconstruction was used to estimate contact areas between Iba1$^+$ cells and GFAP$^+$ cells. Rectangles highlight examples of glial surface-surface colocalization, shown in high magnification views. Scale bar, 20 µm. **B.** Quantification of contact areas between Iba1$^+$ cells and GFAP$^+$ cells at 7 and 21 dpi in WT, and $Asc^{-/-}$ mice subjected to Sham or CHI. n = 12 slices from 4 mice per group per each time point, $*p < 0.05$, $***p < 0.01$, $***p < 0.001$, $****p < 0.0001$ by ordinary two-way ANOVA with Bonferroni's post hoc test. Right panels show integrative AUC of each marker over time (dpi). $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$ by Student's t-test.

Figure 5: Genetic deficiency of the Asc gene protects against mild cognitive impairments following mild CHI.

A. Experimental timeline for cognitive assessment following CHI, including spatial learning phase (10-15 dpi), probe trial (16 dpi), and cued learning (17-20 dpi) in the Morris water maze. **B-E.** Morris water maze performance in WT and $Asc^{-/-}$ mice, showing path length and thigmotaxic across spatial **B-E** and **F-J** cued training days (n=10-11 mice per genotype $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$ by repeated measurements two-way ANOVA with Tukey's post hoc test). Integrative AUC for path length and thigmotaxic from spatial learning (**C** and **E**) and cued learning phase (**G** and **I**). **J.** Representative heatmaps mean swimming trajectories and **k.** quantification of percentage of time spent in target in comparison to another quadrants (a.o.) or **L.** platform crossing times in virtual target platform in comparison to another viral platform (a.o.) (n=10-10 mice per Genotype, $*p < 0.05$, $**p < 0.01$, $***p < 0.001$ by ordinary one-way and two-way ANOVA with Tukey's post-hoc tests). Data are presented as mean ± SEM.

Figure 6: NLRP3 regulates ASC aggregation and distribution following CHI.

A. Immunohistochemistry for Iba1 (green), GFAP (magenta) and ASC (red) in cortices of WT and $Nlrp3^{-/-}$

mice at 7 and 21 dpi. ASC localization within Iba1+ cells (yellow spots), $GFAP^+$ cells (blue spots), and

outside glial cells (red spots) were analyzed using IMARIS. Rectangles highlight examples of ASC

distribution shown in high magnification views. Scale bar, 20 μm. **B-E.** Quantification of total ASC

aggregates numbers (**B**), intracellular ASC aggregates within $Iba1^+$ cells (**C**), ASC aggregates outside glia

cells (**D**), and ASC aggregates with a volume range 30–40 um^3 (**E**) at 7 and 21 dpi in WT and $Nlrp3^{-/-}$ mice

subjected to Sham or CHI. n = 12 slices from 4 mice per group per each time point, $^*p < 0.05$, $^{***}p < 0.01$,

$^{***}p < 0.001$, $^{****}p < 0.0001$ by ordinary two-way ANOVA with Bonferroni's post hoc test. Right panels

show integrative AUC of each marker over time (dpi). $^*p < 0.05$, $^{**}p < 0.01$, $^{***}p < 0.001$, $^{****}p < 0.0001$ by

Student's t-test. **F-I.** Spearman's correlation analysis (r) with a linear regression model (black lines WT,

red lines $Nlrp3^{-/-}$) with an interaction term for ASC aggregates (total, within glial cells, and outside glial

cells) and glia cell contact areas. n = 36 slice from 12 mice per genotype, $^*p < 0.05$, $^{**}p < 0.01$, $^{***}p <$

0.001, $^{****}p < 0.0001$. Differences between WT and $Nlrp3^{-/-}$ Spearman Correlation Coefficients (r) was

performed using the standard Fisher's z-transformation and subsequent comparison(69) $^*p < 0.05$, $^{**}p <$

0.01.

Supplemental Figure 1: Single-Cell Transcriptomic Analysis of Brain Immune Cells Following TBI.

A. UMAP clustering of single-cell RNA sequencing (scRNA-seq) data, comparing control (red) and TBI (green) samples, illustrating shifts in immune cell populations post-TBI. **B.** Heatmap showing gene expression profiles across different brain cell types and cluster identification. **C.** Dot plot depicting the

756   expression of key inflammasome-related genes across different brain cell types. **D.** Dot plot depicting

757   gene expression profile of inflammasome-related genes in microglia after TBI.

Supplemental Figure 2: ASC modulates expression and cleavage of CASP-8 and CASP-3 following

CHI.

A. Schematic of injury model/location. Coronal view (right) shows tip positioned over the skull midline.

Horizontal view (left) shows impact site relative to bregma. **B.** No significant motor, reflex or reaction

deficits at 1- or 7-days post-intervention (dpi) when the mice were evaluated using the Revised

Neurobehavioral Severity Scale (NSS-R). (Two-way ANOVA followed by Bonferroni's post-hoc tests, n=11

mice per group per each time point, data are presented as the mean ± SEM) **c.** Representative

immunoblot images of CAPS 8 and CASP 3 at 1, 7, and 21 dpi in peri-contusional cortices of mice

subjected to Sham or CHI, comparing WT and *Asc $^{-/-}$* groups. **D-e.** Quantification of expressions of CASP

8 and CASP 3. The band intensity of a given target protein was normalized to the corresponding GAPDH

signal for each sample. Data were further normalized to the average of the corresponding WT group and

are presented as the mean ± SEM. Statistical significance was determined using two-way ANOVA

followed by Bonferroni's post-hoc tests, n = 4 mice per group per each time point, $*p < 0.05$, $**p < 0.01$,

$***p < 0.001$, $****p < 0.0001$ (left panels). Quantification of area under the curve (AUC) of each target over

time (dpi) in WT and Asc$^{-/-}$ mice is shown in right panels. Statistical analysis was performed using an

unpaired two-tailed t-test. Data are presented as the mean ± SEM. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$,

$****p < 0.0001$.

Supplemental Figure 3: The cellular source of IL-1 β Following mTBI.

A. Representative images of immunohistochemical staining of the contusional cortices for IL-1β (red), Iba1
(green) and GFAP (green) in mice following sham surgery and 7dpi and 21 dpi after CHI. **B.** 3D
reconstruction of IL-1β immunostaining at 7dpi and 21 dpi following CHI. IL-1β (yellow spot) localization

within Iba1$^+$ and GFAP$^+$ cells. Scale bar, 20 μm. **C-D.** IL-1β spot counts assessed in Iba1$^+$ and GFAP$^+$

cells at 7 and 21 dpi following CHI. Data are presented as the mean ± SEM. Statistical significance was

determined using two-way ANOVA followed by Bonferroni's post-hoc tests, n = 12 slice (4 mice) per group

per each time point, *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001.

Supplemental Figure 4: Swimming velocity during cued learning in the Morris water maze.

793
794
795 Swimming velocity during cued learning in Sham WT, CHI-injured WT **A.** CHI-injured *Asc*[-/-]. **B.**
796 Comparison of the integrated area under the curve of swimming velocity.
797

Supplemental Figure 5: ASC antibody validation and ASC aggregates analysis.

A-B. Validation of ASC antibody specificity using immunofluorescence staining and immunoblotting. **A.**

ASC signal is detected in WT mice but absent in *Asc*$^{-/-}$ mice. **B.** Precipitated supernatant from LPS-

stimulated primary microglia and cortical tissue from CCI brain-injured mice were used as positive

controls. **c.** Representative immunohistochemical images of Iba1 (green) and ASC (red) in peri-

contusional cortex of WT and *Nlrp3*$^{-/-}$ mice at different time points (sham, 7dpi, 21 dpi). ASC aggregates

were analyzed and 3D reconstructed using IMARIS. Scale bar, 20 μm. **D-F.** Quantification of ASC

aggregation and ASC aggregates volume. Two-way ANOVA with Bonferroni's post-hoc tests (*$p < 0.05$,

**$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$). n = 12 slices (4 mice) per group per each time point. Data shown

as mean ± SEM.

A

Sham

WT

Asc⁻/⁻

7 dpi

WT

Asc⁻/⁻

21dpi

WT

Asc⁻/⁻

IBA1 GFAP   Contact   Merge

B