ASC contributes to sustained glial reactivity and mild cognitive impairment after closed-head injury

Li T*[1], Castro-Gomez S*[#1,2], Botella Lucena P[3], Vieira-Saecker A[1], Schwartz S[1], Deng Y[4], Ding Y[1], Stein V[1], Latz E[5,6,7,8], Heneka MT[#3,7]

1. Institute of Physiology II, University Hospital Bonn, University of Bonn, Bonn, Germany
2. Center for Neurology, Clinic of Parkinson, Sleep and Movement Disorders, University Hospital Bonn, University of Bonn, Bonn, Germany.
3. Luxembourg Centre for Systems Biomedicine (LCSB), University of Luxembourg, Esch-sur-Alzette/Belvaux, Luxembourg
4. German Center for Neurodegenerative Diseases (DZNE), Bonn, Germany
5. Institute of Innate Immunity, University Hospital Bonn, Bonn, Germany
6. Centre of Molecular Inflammation Research, Norwegian University of Science and Technology, Trondheim, Norway
7. Division of Infectious Diseases and Immunology, University of Massachusetts Medical School, Worcester, USA
8. Deutsches Rheuma-Forschungszentrum (DRFZ), Berlin, Germany

* These authors contributed equally

[#] Corresponding authors: michael.heneka@uni.lu, sergio.castro-gomez@ukbonn.de

Abstract

Mild brain trauma from closed-head injuries (CHI) can lead to prevalent neuropsychiatric disorders, including an increased risk for neurodegenerative diseases and dementia. Inflammasomes are molecular complexes crucial for neuroinflammation and secondary damage after trauma, however their role in mild CHI is poorly understood. In this study, we investigate the cellular expression of inflammasome-related genes and their functional significance after a CHI models. Analyzing single-cell RNA sequencing data from cortical cells from a model of CHI, we found that *Pycard* (*Asc*), the gene encoding for a common inflammasome adaptor apoptosis-associated speck-like protein containing a caspase recruitment domain (ASC), is expressed particularly in microglial clusters. Sustained upregulation of inflammasome-related proteins persisted up to 21 days in a model for mild CHI, with this pattern significantly reduced in $Asc^{-/-}$ mice. Importantly, mild cognitive impairment induced after mild CHI was largely abrogated in $Asc^{-/-}$ mice. These findings suggest that ASC, as the primary inflammasome adaptor, plays a critical role in sustaining neuroinflammation and contributes to cognitive deficits after mild CHI. This study provides insights into the molecular neuroinflammatory mechanisms underlying CHI, potentially informing future therapeutic strategies.

Introduction

Traumatic brain Injury (TBI) represents a significant global health challenge and is considered a silent epidemic, affecting individuals across all demographics and age groups. In the European Union, it was estimated that TBI causes approximately 1.5 million hospital admissions and 57,000 deaths annually[1]. The majority of cases are clinically classified as mild TBI (mTBI), typically resulting from closed-head injuries (CHI) caused by falls, traffic accidents, violence, contact sports, or military actions[2,3]. Although mTBI symptoms can recover within days or weeks, up to 20% of individuals experience persistent physical, cognitive, and behavioral impairments that lead to a reduced quality of daily life and an elevated risk of neuropsychiatric disorders, including depression and dementia[4–6]. Despite its prevalence, the underlying pathophysiology of mTBI resulted from CHI remains poorly understood, posing great challenges for early clinical diagnosis and timely intervention.

Neuroglia play crucial roles in innate immune responses and secondary damage after TBI. Prolonged glial activation impairs debris clearance, contributing to neuroinflammation, exacerbating neuronal damage and abnormal protein aggregation[7–9]. Advances in imaging techniques have allowed the *in vivo* visualization of glial reactivity after CHI, emphasizing the involvement of microglia and astrocytes in the inflammatory cascade that contributes to the development of post-traumatic disorders[10–12]. A key element in initiating and maintaining innate immune responses of the brain is inflammasome signaling. Inflammasomes are oligomeric protein scaffolds rapidly activated by some damage-associated molecular patters (DAMPs), and organized as a tripartite complex: a cytosolic danger sensor such as NOD-like receptor family pyrin domain containing 3 (NLRP3), a unique adaptor protein called apoptosis-associated speck-like protein containing a CARD domain (ASC), and a proteolytic effector caspase-1 (CASP1)[13]. ASC, an intracellular adaptor protein common to almost all inflammasomes, oligomerizes into highly cross-linked macromolecular assemblies, forming visible ASC aggregates. The fibrillar ASC acts as a molecular platform to recruit pro-CASP1, which optimizes signal transduction via proximity-induced CASP1 activation. Activated CASP1 catalyzes the maturation of pro-forms of Interleukin-1β (IL-1β), IL-18 and mediates the proteolytic activation of Gasdermin D (GSDMD). Oligomerized GSDMD forms pores in the plasma membrane, allowing the release of IL-1β, IL-18 and other proinflammatory cytokines. The detection of tissue ASC aggregates is considered a typical hallmark of inflammasome activation[14].

In humans, some inflammasome proteins are detected in blood and cerebrospinal fluid after TBI, and circulating levels of some proteins such as ASC and CASP1 have been proposed as biomarkers for determining injury severity[15–17]. Furthermore, inflammasome activators, including NLRP1[18], NLRP3[19,20], and absent in melanoma 2 (AIM2)[21], presumably contribute to the innate immune response and functional disability in mouse models of TBI. However, the findings of some studies have been counterintuitive. For instance, a study using *Nlrp1b* and *Asc* deficient mice found that, although cortical IL-1β levels were significantly reduced following controlled cortical impact (CCI)—a standard open-head traumatic brain injury model—no notable differences were observed in either motor function impairments or lesion size compared to controls[22]. Noteworthy, these studies primarily focus on inflammasome activation in models of moderate to severe TBI, such as CCI. At the same time, their contribution to mild CHI remains largely

unexplored, with evidence limited to observations of dynamics in cytokine release[23,24]. In the present
study, we hypothesized that inflammasomes ensembles contribute actively to the secondary damage
following CHI. We aimed to investigate their role in neuroinflammatory responses and subsequent
cognitive functions by analyzing a well-established single-cell RNA sequencing (scRNA-seq) dataset of
cortical cells from mice subjected to a diffuse TBI model,[25] and by using an additional CHI mouse model
that induces mild cognitive impairment[26]. Our scRNA-seq data analysis reveals a subacute expression of
Asc (also known as *Pycard*) in cortical cells from mice subjected to midline fluid percussion injury (mFPI),
particularly in microglia subpopulation. In concordance with this analysis, we observed sustained
upregulation of inflammasome-related proteins, including NLRP3, ASC, CASP-1, and IL-1β, following CHI,
persisting up to 21-days post injury (dpi). This expression pattern was significantly reduced in $Asc^{-/-}$ mice.
Moreover, $Asc^{-/-}$ mice were protected from the mild cognitive impairment induced by our CHI model, as
assessed by the Morris Water Maze (MWM). We additionally observe microglia, and astrocytes reactivity
that sustains overtime together with the upregulation and aggregation of the ASC protein. These findings
suggest that ASC, as the primary inflammasome adaptor, plays a critical role in sustaining
neuroinflammation and contributes to the cognitive deficits after CHI.
90

Results

Subacute *Asc (Pycard)* expression in microglia after a model for CHI

To examine the cell expression of gene coding for inflammasome-related proteins after CHI, our first
approach was to adopt a single-cell genomic analysis. We chose to examine a publicly available scRNA-
seq dataset from dissociated cortical cells 7 days (7 days post injury, 7dpi) followed mFPI[25]. mFPI mimics
several relevant features of CHI including diffused trauma without localized lesion or hemorrhages, and
closely models mild to moderate TBI[27]. After quality control, we analyzed 14688 control cells and 10,745
cells in the mFPI group (Fig. 1b and Supplementary Fig. 1a). The complex multidimensional dataset was
transformed and visualized using the nonlinear dimensionality reduction technique Uniform Manifold
Approximation and Projection (UMAP) (Fig. 1a, 1b, 1e). Examination of cellular transcriptional identities at
the single-cell level uncovered 22 distinct cellular clusters. Cellular subpopulation labels were assigned via
automated cell type annotation per media sample using SingleR[28] (Supplementary Fig. 1b). Cells
identified as microglia were specially enriched in this dataset (Fig. 1b). A differential expression gene
analysis of this subset of cells showed a typical disease-associated microglia signature after TBI, with
overexpression of type-1 interferon genes (such as *Ifi27l2a, Ifitm3, Ifit3, rf7*) and *Apoe,* and reduction of
homeostatic genes such as *tmem119* or *tgfb1* (Fig. 1c).
We then evaluated, for each predicted cellular subpopulation, the expression of genes annotated in
pathways related to inflammasome activation (pathcards.genecards.org[29], Supplementary Fig 1c).
Notably, we found *Asc* (or *pycard,* gene coding for the adaptor protein ASC), as one of the genes with
wider expression in clusters represented by microglia, macrophage, and other peripheral infiltrating
immune cells (Fig. 1d-e and Supplementary Fig. 1c, 1d). These clusters show a more limited percentage
of cells expression genes coding for inflammasome effectors such *Gsdmd, Casp1, il1b* or *il18* (Fig. 1d and

e). Surprisingly, we found a very low percentage of cells expressing genes coding for inflammasome
activators such as *Nlrc4, Nlrc5, Nlrp1b, Nlrp3 or Aim2* (Supplementary Fig 1c). These results suggest that
the subacute and prolonged expression of the adaptor protein ASC, especially in microglia, may play a
central role in the sustained innate immune response and pathophysiology of TBI related to CHI.

ASC contributes to the sustained expression and processing of inflammasome mediators
following mild CHI
To unveil the pathophysiological role of inflammasomes and specifically of ASC as main inflammasome
adaptor and abundantly expressed gene in microglia after CHI, we adopted an electromagnetic CHI
procedure as main model for mTBI[30] In this model, a controlled cortical impact injury was delivered directly
to the midline surface of the skull using a 5 mm diameter tip at a specified depth and velocity
(Supplementary Fig. 2a). We found no significant motor, reflex or reaction deficits at 1- or 7 dpi when the
mice were evaluated using the Revised Neurobehavioral Severity Scale (NSS-R)[31] for rodents
(Supplementary Fig. 2b). To examine the protein expression of inflammasome-related proteins after CHI,
we performed immunoblot analysis of cortices from WT and $Asc^{-/-}$ mice that underwent either sham or CHI
surgery. We analyzed and semi-quantified the protein levels of NLRP3, ASC, CASP1, and IL-1β at 1, 7,
and 21 dpi (Fig. 2a-i). Our results revealed sustained upregulation of these proteins up to 21 days after
CHI in WT mice. Notably, $Asc^{-/-}$ mice showed a very significant deficit in the expression and cleavage of
CASP1 (Fig. 2d-e) and of IL-1β over time (Fig. 2g). ELISA measurements further demonstrated
significantly lower soluble levels of IL-1β and TNF-α in $Asc^{-/-}$ mice (Fig. 2h-i), particularly at later time
points. We additionally evaluated the levels of cleavage of Caspase-8 (CASP8), a proposed non-canonical
enzyme involved in IL-1β processing[32] (Supplementary Fig. 2c and d), and Caspase-3 (CASP3), as
indirect measurement of apoptotic pathways[33] (Supplementary Fig. 2c and e). Consisting with the soluble
IL-1β measurements, $Asc^{-/-}$ cortices showed significantly reduced levels of cleaved CASP8 up to 21 days
post CHI (Supplementary Fig. 2d), which suggest parallel activation of canonical and non-canonical
pathways of IL-1β processing, both of which appear largely dependent on ASC. Additionally, the cleavage
of CASP3 was significantly reduced in $Asc^{-/-}$ mice (Supplementary Fig. 2e), suggesting a stronger
neuroprotective effect and a complex crosstalk involving the activation of multiple types of cell death
influenced by inflammasome signaling[34].
To identify the cellular sources of IL-1β following mild CHI, we conducted immunohistochemical staining
for IL-1β, ionized calcium binding adaptor molecule 1 (Iba1), Glial fibrillary acidic protein (GFAP), and
NeuN on cortices of WT mice in sham, 7 and 21 dpi groups (Supplementary Fig. 3a-d). At 7 days post-
mTBI, approximately 60% of IL-1β signal was found to be colocalized with or in the vicinity of GFAP[+] cells
(presumably astrocytes), 30% with Iba1[+] cells (likely microglia), and 10% with NeuN[+] cells (likely neurons)
(Supplementary Fig. 1c). Interestingly, at 21 dpi, the majority of IL-1β (approximately 68%) was detected
in Iba1[+] cells (microglia or infiltrating macrophages), while around 23% was found in GFAP[+] cells and 9%
in the vicinity of NeuN[+] cells (Supplementary Fig. 3c). Our findings thus emphasize the crucial role of
inflammasome signaling, with ASC the common adaptor protein, in sustaining the expression and

processing of inflammatory mediator after CHI. Moreover, we identified Iba1$^+$ cells, likely microglia, or

infiltrating macrophages, as the primary cellular sources of IL-1β, especially at later time points post-injury.

ASC shapes the morphology of Iba1$^+$ cells following CHI

Microglia exhibit distinct adaptive morphologies in response to mechanical stimuli, and their morphology is

considered a proxy for their function[35]. To investigate microglia architecture after CHI, we performed

immunofluorescence staining for Iba1 and performed skeleton analysis[36] to examine cell morphology in

cortices of WT and $Asc^{-/-}$ mice at 1, 7, and 21 dpi (Fig. 3a–b). First, we observed that $Asc^{-/-}$ mice

consistently exhibited lower Iba1+ cell counts compared with WT mice at 7 dpi (Fig. 3c), suggesting a

decreased microgliosis in the subacute CHI phase. Next, we quantified the normalized changes in the

morphology of Iba1+ cells based on cell count. The morphological alterations, including the number of

branches per cell, endpoints per cell, and branch length per cell, significantly decreased after injury in WT

mice, particularly at 21 dpi. In contrast, $Asc^{-/-}$ mice showed a significant preservation of these

morphological features at 21 dpi compared to WT mice (Fig. 3d-f). In conclusion, our observations suggest

a wave of microgliosis in the subacute phase and revealed a progressive debranching of Iba1+ cells over

time following injury. Notably, the lack of ASC demonstrated protection against these sustained cellular

morphological alterations, particularly at later time points.

ASC regulates the reactivity of GFAP+ cells following CHI

Previous studies have provided evidence that morphological changes in astrocytes are intricately linked to

microglia activation during inflammation-induced responses[37]. To examine the morphological alterations in

astrocytes following mild CHI, we conducted a morphological analysis of GFAP-immunostained tissue,

applying the same method performed for Iba1+ cells (Fig. 3h-i). We initially observed a noteworthy

increase in the count of GFAP+ cells at 1, 7 and 21 dpi in WT mice (Fig. 3j). *Asc* deficiency exerted a

genotype-related effect in the cell count like what we observed in Iba1+ cells, showing significantly lower

numbers, especially at 7 dpi (Fig. 3j). Subsequently, we quantified normalized morphological changes in

GFAP+ cells based on cell counts. The branches per cell, endpoints per cell, and length of branches per

cell were significantly decreased after CHI, particularly at 21 dpi. In contrast, $Asc^{-/-}$ mice present preserved

astrocyte morphological features at 21 dpi compared to WT mice (Fig. 3k–m). Interestingly, at 21 dpi, we

observed clusters of aggregated GFAP+ cells, indicating spatial reorganization post-injury. Similar

morphological changes were evident in both microglia and astrocytes after injury, suggesting a potential

correlation or interactions between these two glial cell populations.

Glial cell interactions are modulated by ASC following CHI

To investigate the interaction between astrocytes and microglia in the contusional cortex following injury,

we performed surface–surface colocalization analysis of the contact areas after 3D reconstruction of

confocal images from GFAP- and Iba1-immunostained tissue (Fig. 4a). We observed a significant

increase in contact area between microglia and astrocytes at 7 dpi, reaching a peak at 21 dpi in WT mice

(Fig. 4b). Notably, the contact area was significantly reduced at 7 and 21 dpi in $Asc^{-/-}$ mice (Fig. 4b) which

may be explained by the deficient gliosis observed in our previous analysis. These findings, together with

our expression analysis, suggest that inflammasome ensembles and neuroinflammatory processing are

essential for sustained neuroglial responses and cellular interactions after mild CHI.

Constitutional *Asc* deletion protects against mild cognitive impairments after CHI

Having demonstrated that ASC is predominantly expressed by microglia in the subacute phase of CHI,

contributing to the sustained inflammasome activation, as well as chronic neuroglial reactivity following

CHI, we next sought to investigate whether the genetic removal of the main inflammasome adaptor protein

ASC influences cognitive outcomes after mild CHI. In humans, a substantial number of individuals present

transient or persistent cognitive impairment following mTBI, including deficits in memory recall, procedural

tasks, and executive functions[38]. To investigate how ASC contribute to these neurobehavioral phenotypes,

we performed Morris Water Maze (MWM) as main behavioral task in WT and $Asc^{-/-}$ mice after CHI. MWM

was conducted from days 10 to 20 after CHI (10-20 dpi) and consisted in a spatial learning phase (10 to

15 dpi), a 24-hour long-term spatial memory test (16 dpi), and a cued learning phase as control procedure

between 17 and 20 dpi (Fig. 5a). In comparison to the WT Sham group, WT mice subjected to CHI

learned the task with comparable daily reduction in path lengths towards the hidden platform (Fig. 5b).

However, CHI induces significant prolonged swimming time near the walls (thigmotaxis) during the

learning phase in WT injured mice (Fig. 5d). We observed similar tendency in thigmotaxis during the cued

version of the task (Fig. 5h). This procedural deficit was largely reduced in $Asc^{-/-}$ mice (Fig. 5d-e and 5h-i).

Additionally, $Asc^{-/-}$ mice demonstrated preserved long-term spatial memory 24 hours after the learning

phase, as visually suggested by the density plots (heat maps) of swim paths (Fig. 5j). This finding was

further supported by a significantly greater percentage of time spent exploring the target quadrant (Fig.

5k), as well as more frequent crossings of the virtual target platform (Fig. 5i) compared to the

corresponding measures in another quadrant or at other virtual platforms (Fig. 5l) during the probe trial

performed at 16 dpi. Furthermore, $Asc^{-/-}$ mice displayed slightly higher velocity in the last day of cued

learning (Supplementary Fig. 4a and b). These data show that CHI induces mild cognitive impairments,

mainly as procedural and memory deficits in the MWM, which are mostly dependent on ASC expression.

NLRP3 regulates ASC aggregation and distribution following CHI

The NLRP3 inflammasome is widely recognized as one of the most relevant in the context of TBI, with

growing evidence indicating its central role in mediating neuroinflammation and secondary brain

damage[39]. To explore the involvement NLRP3 in the aggregation of ASC as the main inflammasome

adaptor in post-CHI, we conducted immunohistochemical staining of the contusional cortical tissue for

Iba1, GFAP and ASC at 7 and 21 dpi in in WT and $Nlrp3^{-/-}$ mice (Fig. 6a). We used a KO-validated

monoclonal antibody for ASC detection (Supplementary Fig. 5a-b). 3D surface reconstruction of microglia

and ASC immunosignal analysis were subsequently conducted (Fig. 6a and Supplementary Fig. 5c). We

detected a significant increase in the total number of fluorescent ASC aggregates, especially in microglia

at 7 dpi similar to the mRNA expression of *Asc* observed in the scRNA-Seq data analysis following mFPI
(Fig. 1b, Fig. 6b and Supplementary Fig. 5d). The number of ASC aggregates within Iba1+ cells followed
the total ASC aggregates pattern (Fig. 6b and Supplementary Fig. 5c-d). Notably, *Nlrp3*$^{-/-}$ injured mice
showed comparable ASC aggregates counts, both overall and within Iba1+ cells, compared to WT mice
(Fig. 6b) but significantly lower aggregates in GFAP+ cells (presumably Astrocytes, Fig. 6c). Interestingly,
at 21 dpi, the number of ASC aggregates outside Iba1+ and GFAP+ cells (extra-glial ASC) significantly
increased in both WT and *Nlrp3*$^{-/-}$ mice (Fig. 6d). However, this ensembles count was significantly lower in
Nlrp3$^{-/-}$ mice compared to WT controls (Fig. 6d). Additionally, as ASC aggregates numbers increased
within microglia, their corresponding volumes also increased. To further analyze ASC aggregation, we
categorized ASC aggregates by volume (1–20 µm³, 20–30 µm³, 30–40 µm³) and quantified the number of
ensembles in each category (Supplementary Fig. 5e-f and Fig. 6e). ASC aggregates within a range of 30–
40 µm³ were significantly decreased in the *Nlrp3*$^{-/-}$ mice at 21 dpi (Fig. 6e). Collectively, our findings
demonstrate a remarkable upregulation of ASC aggregates outside of Iba1+ and GFAP+ cells, along with
marked ASC aggregation at 21 dpi. *Nlrp3*$^{-/-}$ mice displayed reduced ASC aggregation within Iba1+ cells
and fewer extra-glia ASC aggregates compared to WT group at 21 dpi. These observations highlight the
essential role of NLRP3 not only in promoting inflammasome activation after brain injury but also in
facilitating significant ASC aggregation, as previously suggested in *in vitro* studies[40].

Interaction between microglia and astrocytes correlates with ASC distribution following CHI
To further investigate the relationship between glial interactions and ASC distribution following CHI, we
conducted correlation analysis between ASC aggregates and glial contact areas after reconstruction (Fig.
6 f–i). Notably, we observed a strong positive correlation between the glial contact area and the number of
extra-glia ASC spots *(r = 0.86, p < 0.0001)* (Fig. 6g) as well as ASC aggregates within GFAP+ cells *(r =
0.93, p < 0.0001)* (Fig. 6i), in WT animals after CHI. Although lower correlations coefficients were found
between the glial contact areas and the number of ASC aggregates in microglia *(r = 0.73, p < 0.0001)*
(Fig. 6h) and the overall count of ASC aggregates *(r = 0.82, p < 0.0001)* (Fig. 6f), these correlations
remained significantly positive (Fig. 6f–i). Interestingly, the constitutional deficiency of the *Nlrp3* gene
significantly reduced the correlation between the glial contact area and the number of extra-glia ASC
aggregates *(z = 1.95, p < 0.05)* (Fig. 6g) as well as ASC in GFAP+ cells *(z = 2.54, p < 0.01)* (Fig. 6i). In
summary, these results indicate that ASC aggregation after CHI is partly dependent on NLRP3 and
additionally glial contact is influenced by ASC expression, distribution, and aggregation, especially the
ASC aggregates located outside Iba1^{+} and GFAP^{+} cells.

Discussion

In this study, we identify *Asc* as a key gen expressed in monocytes, macrophages, and especially in
microglia during the subacute phase of CHI based on scRNA-seq data from mouse cortical cells after
mFPI. Similarly, in our mild CHI mouse model, sustained ASC protein expression was confirmed, driving
glial reactivity and innate immune responses that contribute to neurobehavioral deficits following mTBI.

Our findings complement and expand previous preclinical and clinical studies on moderate to severe TBI

models[41,42]. These studies highlight the role of inflammasome cascades and their critical involvement in

regulating the secondary inflammatory responses after TBI. However, unlike moderate-to-severe TBI

models, where the expression of inflammasome-related proteins typically peaks within the initial week

following injury and declines over time[20,43,44], we found a progressive and sustained increase in

inflammasome-related proteins specially after the first 7 days following mild CHI. This temporal pattern

aligns with our observations of sustained morphological changes of microglia and astrocytes during a later

phase (21 dpi). These discrepancies suggest that inflammasome engagement in innate immune

responses and tissue repair varies depending on injury severity. Moreover, the prolonged pro-

inflammatory characteristics exhibited by microglia and astrocytes indicate both of these glia cells are

capable of propagating inflammation through the inflammasome-related pathways post-injury.

Interestingly, our investigation demonstrated that astrocytes display more significant inflammatory

morphology at 21 dpi compared to 7 dpi. This may result from the potential further induction of neurotoxic

astrocytes by microglial inflammasome in response to sustained tissue stress, as previously reported[45].

These findings align with existing evidence suggesting that distinct glial cell types respond differentially to

brain injury in a temporal sequence[36,46–48].

Additionally, we observed a widespread upregulation of inflammasome components following CHI,

particularly at 21 dpi, with a significant increase in the cleaved forms of CASP1, CASP8 and IL-1β. Our

results suggest the involvement of several ASC-dependent inflammasomes in the inflammatory pathways

after CHI. This is evident by the pronounced reduction in CASP1, CASP3 and CASP8 levels and glial

morphological changes observed in *Asc*[-/-] mice. Most notably, our behavioral evaluation during the late

subacute phases shows that mild procedural learning deficits and long-term memory impairments were

significantly diminished in animals with constitutional deletion on the *Asc* gene. Furthermore, our analysis

established a significant positive correlation between astrocyte-microglia interaction and the number of

ASC aggregates. Notably, a potential connection has been identified between extracellular ASC and

cellular propagation of inflammatory mediators, including IL-1β and TNF-α following inflammasome

activation[49,50]. These cytokines not only promote the aggregation of neuroglia and infiltrated cells around

the trauma epicenter[51,52], but may also influence glial interactions far from the lesion site through

extracellular ASC. ASC possess prion-like properties[50] and could propagate to neighboring glial cells,

thereby triggering the activation of the ASC-mediated inflammasome cascade as previously shown in the

context of Alzheimer's disease- related amyloidosis[53,54]. These observations suggest a pivotal role of ASC

in sustaining neuroinflammation and glial reactivity following CHI.

Our findings, however, contrast with a previous report on moderate controlled cortical impact, where *Asc*[-/-]

mice showed no differences in motor recovery or volume lesions[22]. This inconsistency may suggest that,

in the context of low-grade chronic neuroinflammation induced by mild, uncontrolled stimuli, such as a

single concussion event, the expression, oligomerization, and propagation of ASC may occur more

gradually compared to the high-grade neuroinflammation triggered by severe injuries. Similarly, ASC-

dependent inflammasome activation and pathological protein aggregation have been observed in several

neurodegenerative disorders characterized by sustained chronic low-grade neuroinflammation[55], such as
Alzheimer's and Parkinson's disease, where ASC plays a decisive role in the aggregating and further
propagating Amyloid β[53] and α-Synuclein[56]. Considering our findings, ASC may govern the immune
molecular mechanism that triggers neurodegeneration and increase the risk of developing
neurodegenerative conditions following traumatic brain injuries[57]. Nonetheless, future investigations are
still required to establish a comprehensive understanding of the specific temporal participation of
inflammasomes across varying injuries severity and the risk factor for neurodegeneration.

There are certain limitations that should be acknowledged in our study. Focusing on the expression of
inflammasome mediators and glial morphological characteristics may not fully capture the complex
interactions with other compensatory mechanisms within the affected neuronal networks or the further
anti-inflammatory modulation of secondary immune responses. Additionally, although our scRNA-Seq
analysis determined specific cellular clusters upregulating *Asc* in the subacute phase of a CHI model, our
study using constitutional knockouts cannot dissect precisely the role of different peripheral and cellular
actors in the immunological process following CHI. Thus, there is an imperative need to investigate the
inflammatory evolution of the specific neural and glial cellular subtracts over extended post-traumatic
periods.

Our findings pave the way to delve into specific pharmacological agents that can effectively modulate ASC
activity and aggregation and assess their impact on neurological outcomes after TBI. Considering the
pathological mechanisms described, pharmacological interventions targeting ASC may help to mitigate
neuroinflammation and potentially promote neuroprotection, enhancing recovery of injury and preventing
further neurodegeneration and disability.

Methods

Ethical compliance

327 We followed the European guidelines for animal research and received approval from North Rhine-
328 Westphalia State Agency for Nature, Environment and Consumer Protection under folder number 81-
329 02.04.2019.A026 and 2024-791-Grundantrag.

Single-cell RNA-sequencing data analysis

Raw FASTQ files and count matrices are publicly available in the NCBI GEO database under accession
number GSE160763[25]. Single-cell RNA-seq data were analyzed using Seurat v5.1.0 in R v4.3.2[58]. Briefly,
cells with over 5000 detected genes, UMI counts under 12,500, and less than 1% mitochondrial gene
content were retained, followed by log-transformation of the data. We selected the top 4000 highly
variable genes for principal component analysis (PCA) and used the top 20 principal components as input
for UMAP dimensionality reduction. Cell cycle states were determined using Seurat's CellCycleScoring[59]
function with marker genes from Tirosh et al.[58]. Batch correction was performed using Harmony to
integrate datasets across samples[60]. Clusters were identified at a resolution of 0.5. Cellular identities were
assigned via automated cell type annotation using SingleR[28], which assigns cellular identity for single cell

transcriptomes by comparison to reference data sets of pure cell types sequenced by microarray or RNA-
sequencing (RNA-seq). Annotation relied on a reference dataset of 358 mouse RNA-seq samples
annotated to 18 major cell types[61]. Figures were generated using Tidyverse[62], SCpubr[63], and
EnhancedVolcano[64].

Mice

In this study, we used male and female mice aged 6–7 months with a C57BL/6 mixed genetic background
(C57BL/6N-J) from our own animal facility (House of the Experimental Medicine) at the University Hospital
Bonn in German. Wild-type (WT), *Nlrp3*$^{-/-}$, and *Asc*$^{-/-}$ mice were initially sourced from Millennium
Pharmaceuticals as previously published[65]. All mice were backcrossed with C57BL/6N mice and
maintained under pathogen-free, standardized conditions with a 12:12-hour light/dark cycle and ad libitum
access to food and water. Behavioral experiments were conducted during the dark phase of the light/dark
cycle.

Electromagnetic controlled closed-head injury model of mild traumatic brain injury

To model mild TBI in mice**,** we adapted the electromagnetic controlled Closed-Head Model previously
described[30] (Supplementary Fig. 2a). Briefly, Mice were anesthetized with 5% isoflurane during induction,
and 1.5-2% isoflurane mixed with 100% oxygen (0.5- 1 l/min) during surgery and fixed in a stereotaxic
frame (Stoelting, Dublin, Ireland). A 1-mL latex pipette bulb filled with water was placed under the head to
distribute the impact force. A midline sagittal scalp incision was made, and a single controlled midline skull
impact was delivered at coordinates 0.0 mm mediolateral and -1.5 mm anteroposterior, with a velocity of 5
m/s, dwell time of 100 ms, and impact depth of 1.0 mm, using a stereotaxic electromagnetic impactor with
a 5.0-mm steel tip (Stereotaxic Impactor, Leica Biosystems, Germany). Sham mice underwent identical
surgical procedures without impact injury. In the post-operative phase, mice received carprofen (5 mg/kg)
subcutaneously once daily and tramadol (1 mg/mL) in drinking water for three days.

Morris water maze (MWM)

The evaluation of spatial learning and memory was conducted between 10 and 20 dpi. A circular pool with
a diameter of 1 m was filled with white opacified water maintained at a temperature of 21–23 °C. The pool
was dimly lit (approximately 40 lx) and surrounded by a white curtain. Asymmetrically placed distal cues
were positioned on the pool wall to serve as spatial references. The pool was virtually divided into four
quadrants, one of which contained a hidden platform (15 cm in diameter) submerged 1.5 cm below the
water surface. Mice were trained to locate the platform using the distal cues for orientation. Training
consisted of four trials per day for five consecutive days. During each trial, mice were placed into the water
facing the pool wall in a quasi-randomized order to prevent them from developing fixed strategies. Mice
were given 60 s to find the platform; if they failed to locate it within this time, they were manually guided to
it. Once on the platform, mice were allowed to remain there for 15 s before the next trial began. After
completing all four trials for the day, mice were dried and returned to their home cages. The integrated

time or distance traveled during these trials was analyzed. A spatial probe trial was conducted 24 hours
after the final training session (16 dpi). For this test, the hidden platform was removed, and mice were
allowed to swim freely for 60 s. The drop position was in the quadrant opposite where the platform had
been located, with each mouse facing the wall at the center of that quadrant. One day after the spatial
probe trial, a visually cued learning phase began. During this phase, the platform was flagged for visibility
and relocated to a different quadrant, while all distal cues were removed. Mice underwent three trials per
day for three consecutive days (17–20 dpi) to learn to locate the flagged platform. All mouse movements
were recorded and tracked using Noldus EthoVision software (Wageningen, Netherlands).
387
Tissue preparation and slicing
Mice were deeply anesthetized using ketamine (100 mg/kg) and xylazine (20 mg/kg) and transcardially
perfused with at least 30 ml of ice-cold PBS at designated time points (Sham, 1, 7, and 21 dpi). Following
perfusion, brains were carefully dissected. One hemisphere was snap-frozen in liquid nitrogen and stored
at -80°C until biochemical analyses, while the other hemisphere was fixed in 4% paraformaldehyde
overnight. Coronal brain sections (40 μm thick) were prepared using a Leica VT1000S vibratome (Leica
Microsystems Inc.) from the region spanning bregma -0.45 to -1.85 mm. Sections were collected in PBS at
intervals of 300 μm for various staining procedures.
396
Primary antibodies
For immunoblotting, following primary antibodies were used: Anti-Caspase 1 from AdipoGen (USA, mAb
Casper-1, mouse anti-mouse, 1:1000), anti-ASC from Cell Signaling (Frankfurt, Germany, D2W8U, rabbit
anti-mouse, 1:1500), anti-NLRP3 from Cell Signaling (Frankfurt, Germany, D4D8T, rabbit anti-mouse,
1:500), anti-IL-1β from Cell Signaling (Frankfurt, Germany, E7V2A, rabbit anti-mouse, 1:1000), anti-
Caspase 3 from Cell Signaling (Frankfurt, Germany, D3E9, rabbit anti-mouse, 1:1000), and anti-Caspase
8 from Cell Signaling (Frankfurt, Germany, D5B2, rabbit anti-mouse, 1:1000). For immunohistochemical
analysis, the antibodies included anti-Iba1 from Wako Chemicals (Neuss, Germany, rabbit anti-mouse,
1:1000), anti-IBA1 from Abcam (Cambridge, UK, ab289874, goat anti-mouse, 1:1000), anti-GFAP from
ThermoFisher (Waltham, USA, 2.2B10, rat anti-mouse, 1:300), anti-IL-1β from R&D Systems
(Minneapolis, USA, AF-401-SP, goat anti-mouse, 1:500), and anti-P53 from Abcam (Cambridge, UK,
ab131442, rabbit anti-mouse, 1:500).
409
Tissue protein extraction and immunoblotting
After thawing, brain tissue was homogenized in PBS containing 1 mM EDTA, 1 mM EGTA, and protease
inhibitor cocktail Halt™ (Thermo Scientific, Waltham, Massachusetts, U.S.). The homogenates were
extracted in RIPA buffer [25 mM Tris–HCl (pH 7.5), 150 mM NaCl, 1% Nonidet P-40, 0.5% NaDOC, 0.1%
SDS] and centrifuged at 20000 ×g for 30 minutes.
RIPA fractions were separated using NuPAGE® electrophoresis gels (Thermo Scientific, Waltham,
Massachusetts, USA) and immunoblotted with primary antibodies, followed by incubation with the

corresponding secondary antibodies. Immunoreactivity was detected with an Odyssey CLx Imaging
System (LI-COR, Bad Homburg, Germany), and images were analyzed using ImageJ (NIH, Bethesda,
USA).
420

ELISA pro-inflammatory cytokine panel

422 The levels of IL-1β and TNF-α in RIPA fractions were measured using the V-PLEX Plus Mouse Pro-
423 Inflammatory Panel 1 (Meso Scale Discovery, Rockville, USA), following the manufacturer's protocol.
424 Samples were diluted 1:1 on the plate with the reagent diluent provided in the kit. Signals were detected
425 using a QuickPlex SQ 120 plate reader (Meso Scale Discovery, Rockville, USA).
426

Immunohistochemistry

Brain sections were washed three times for 5 min with PBS and incubated in citrate buffer for 5 min at
95°C. After antigen retrieval, the sections were cooled down at room temperature and washed with PBS
and in 0,5% Triton X-100 B (PBS-T), then blocked for 1 h with BSA 1% in PBS-T, and incubated overnight
with the primary antibodies. Next, the sections were washed three times for 5 min in PBS-T, incubated
with appropriate secondary antibodies (1:1000) for 60 min, and following with three times with PBS
washing for 5 min. The tissue was mounted using ProLong™ Gold Antifade Mountant with DNA Stain
DAPI (Thermo Scientific, Waltham, Massachusetts, U.S.).
435

Skeleton analysis of microglia and astrocytes

Morphological changes in microglia and astrocytes were quantitatively analyzed using a skeleton analysis
method[66]. Images were imported into ImageJ software (Bethesda, Maryland, USA), converted to
grayscale, and adjusted for brightness and contrast to enhance cell process visibility. After obtaining
binary images of microglia and astrocytes, we used the skeletonize plugin to obtain skeletons,
representing the medial axis of cell processes. Morphological parameters measured included: Process
Branches, defined as number of branching points. Endpoints, defined as number of terminal points.
Process Lengths, defined as total length of all processes. The data were normalized by dividing the
measurements by the number of cells in each image.
445

Contact analysis between microglia and astrocytes

Imaris Image Analysis Software (Oxford Instruments, Oxon, England) was used to analyze the contact
area between microglial and astrocytic surfaces. 3D reconstructions were created using the tool surfaces
by setting intensity thresholds to differentiate cell surfaces from the background. Adjustments were made
to ensure accurate cell boundary representation. The Surface-Surface Contact X-Tension identified and
analyzed overlapping regions between microglial and astrocytic surfaces, generating a new surface to
represent the contact area. The contact area (μm^2) was calculated based on the number of overlapping
voxels within the 3D reconstructions.
454

Analysis of ASC aggregates

The cortical impact area was examined using a confocal LSM900 microscope (Zeiss, Oberkochen, Baden-
Württemberg, Germany). Images were captured under consistent confocal settings and processed with
Imaris Image Analysis Software (Oxford Instruments, Oxon, England) to identify ASC aggregates.
Standardized settings were maintained across samples, including pinhole size, laser intensity, digital gain,
and offset. Z-stacks were collected to obtain the complete 3D structure of cells and ASC aggregates.
Image preprocessing included adjusting brightness and contrast, reducing noise, and subtracting
background to enhance aggregate visibility. Imaris Image Analysis Software was used to create 3D
reconstructions of microglia, astrocytes, and ASC aggregates. The "Surfaces" tool generated accurate 3D
models by setting intensity thresholds. The Spots tool identified and quantified ASC aggregates based on
fluorescence intensity and size, with thresholds set using positive (mCherry-ASC mice) and negative (Asc^{-}
$^{/-}$ mice) controls. The Spots Close to Surface tool analyzed the spatial distribution of ASC aggregates
relative to microglia or astrocytes, indicating potential inflammasome activation.

IL-1β analysis

Co-immunostainings for IL-1β, Iba1, GFAP, and NeuN were performed at 7 and 21 dpi. Three-dimensional
reconstructions of microglia, astrocytes, neuronal nuclei, and IL-1β signals were generated using Imaris
Image Analysis Software (Oxford Instruments, Oxon, UK). The "Spots" tool quantified IL-1β signals in the
3D space by analyzing fluorescence intensity and particle size thresholds. To assess spatial relationships,
the "Spots Close to Surface" tool measured the proximity of IL-1β signals to cellular surfaces, evaluating
potential associations with microglia, astrocytes, or neurons. Quantitative metrics, including the number of
IL-1β spots adjacent to specific cell types, were measured for statistical analysis.

Statistics and reproducibility

Data other than scRNA-seq were analyzed using GraphPad Prism Software version 9.0 or later
(GraphPad Software, Inc., La Jolla, CA, USA) and are presented as mean ± SEM. After assessing the
distribution, statistical comparisons were performed using either a two-tailed t-test or one- or two-way
ANOVA with Bonferroni's or Tukey's post hoc test to identify group differences. Morris Water Maze
(MWM) learning data were analyzed using repeated-measures two-way ANOVA with Tukey's post hoc
test. Correlation analyses were performed using non-parametric Spearman correlations. Differences
between Spearman correlation coefficients were assessed using the standard Fisher's z-transformation
and subsequent comparison[67]. Results were considered statistically significant at $p < 0.05$.

Data availability

The data that supports the findings of this study are available on request from the corresponding authors
upon reasonable request.

Code availability

Code used in analyzing scRNA-seq data is available at

https://gitlab.lcsb.uni.lu/pablo.botellalucena/rna_single_seq

References

1. Maas, A. I. R. *et al.* Traumatic brain injury: integrated approaches to improve prevention, clinical care, and research. *Lancet Neurol.* **16**, 987–1048 (2017).

2. Cassidy, J. D. *et al.* Incidence, risk factors and prevention of mild traumatic brain injury: results of the who collaborating centre task force on mild traumatic brain injury. *J. Rehabil. Med.* **36**, 28–60 (2004).

3. Naumenko, Y., Yuryshinetz, I., Zabenko, Y. & Pivneva, T. Mild traumatic brain injury as a pathological process. *Heliyon* **9**, e18342 (2023).

4. Rathbone, A. T. L., Tharmaradinam, S., Jiang, S., Rathbone, M. P. & Kumbhare, D. A. A review of the neuro- and systemic inflammatory responses in post concussion symptoms: Introduction of the 'post-inflammatory brain syndrome' PIBS. *Brain. Behav. Immun.* **46**, 1–16 (2015).

5. Ahman, S., Saveman, B.-I., Styrke, J., Björnstig, U. & Stålnacke, B.-M. Long-term follow-up of patients with mild traumatic brain injury: a mixed-method study. *J. Rehabil. Med.* **45**, 758–764 (2013).

6. Déry, J., Ouellet, B., de Guise, É., Bussières, È.-L. & Lamontagne, M.-E. Prognostic factors for persistent symptoms in adults with mild traumatic brain injury: an overview of systematic reviews. *Syst. Rev.* **12**, 127 (2023).

7. Burda, J. E., Bernstein, A. M. & Sofroniew, M. V. Astrocyte roles in traumatic brain injury. *Exp. Neurol.* **275 Pt 3**, 305–315 (2016).

8. Loane, D. J. & Kumar, A. Microglia in the TBI brain: The good, the bad, and the dysregulated. *Exp. Neurol.* **275 Pt 3**, 316–327 (2016).

9. Simon, D. W. *et al.* The far-reaching scope of neuroinflammation after traumatic brain injury. *Nat. Rev. Neurol.* **13**, 572 (2017).

10. Perez-Polo, J. R. *et al.* Inflammatory consequences in a rodent model of mild traumatic brain injury. *J. Neurotrauma* **30**, 727–740 (2013).

11. Weil, Z. M., Gaier, K. R. & Karelina, K. Injury timing alters metabolic, inflammatory and functional outcomes following repeated mild traumatic brain injury. *Neurobiol. Dis.* **70**, 108–116 (2014).

12. Lv, W. *et al.* mTBI-Induced Systemic Vascular Dysfunction in a Mouse mTBI Model. *Brain Sci.* **12**, 232 (2022).

13. Ravichandran, K. A. & Heneka, M. T. Inflammasomes in neurological disorders - mechanisms and therapeutic potential. *Nat. Rev. Neurol.* **20**, 67–83 (2024).

14. Franklin, B. S., Latz, E. & Schmidt, F. I. The intra- and extracellular functions of ASC specks. *Immunol. Rev.* **281**, 74–87 (2018).

15. Pérez-Bárcena, J. *et al.* Serum Caspase-1 as an Independent Prognostic Factor in Traumatic Brain Injured Patients. *Neurocrit. Care* **36**, 527–535 (2022).

529 16. Kerr, N. *et al.* Inflammasome proteins as biomarkers of traumatic brain injury. *PloS One* **13**, e0210128
530   (2018).

531 17. Eagle, S. R. *et al.* Impact of obese body mass index on inflammasome blood biomarkers and
532   neurocognitive performance following traumatic brain injury with Glasgow coma scale 13 to 15. *J.*
533   *Neurol. Sci.* **464**, 123159 (2024).

534 18. de Rivero Vaccari, J. P. *et al.* Therapeutic neutralization of the NLRP1 inflammasome reduces the
535   innate immune response and improves histopathology after traumatic brain injury. *J. Cereb. Blood*
536   *Flow Metab. Off. J. Int. Soc. Cereb. Blood Flow Metab.* **29**, 1251–1261 (2009).

537 19. Irrera, N. *et al.* Lack of the Nlrp3 Inflammasome Improves Mice Recovery Following Traumatic Brain
538   Injury. *Front. Pharmacol.* **8**, 459 (2017).

539 20. Liu, H.-D. *et al.* Expression of the NLRP3 inflammasome in cerebral cortex after traumatic brain injury
540   in a rat model. *Neurochem. Res.* **38**, 2072–2083 (2013).

541 21. Adamczak, S. E. *et al.* Pyroptotic neuronal cell death mediated by the AIM2 inflammasome. *J. Cereb.*
542   *Blood Flow Metab. Off. J. Int. Soc. Cereb. Blood Flow Metab.* **34**, 621–629 (2014).

543 22. Brickler, T. *et al.* Nonessential Role for the NLRP1 Inflammasome Complex in a Murine Model of
544   Traumatic Brain Injury. *Mediators Inflamm.* **2016**, 6373506 (2016).

545 23. Vincent, J. C. *et al.* IL-1R1 signaling in TBI: assessing chronic impacts and neuroinflammatory
546   dynamics in a mouse model of mild closed-head injury. *J. Neuroinflammation* **20**, 248 (2023).

547 24. Bachstetter, A. D. *et al.* Attenuation of traumatic brain injury-induced cognitive impairment in mice by
548   targeting increased cytokine levels with a small molecule experimental therapeutic. *J.*
549   *Neuroinflammation* **12**, 69 (2015).

550 25. Witcher, K. G. *et al.* Traumatic Brain Injury Causes Chronic Cortical Inflammation and Neuronal
551   Dysfunction Mediated by Microglia. *J. Neurosci. Off. J. Soc. Neurosci.* **41**, 1597–1616 (2021).

552 26. Macheda, T., Roberts, K. & Bachstetter, A. D. Electromagnetic Controlled Closed-Head Model of Mild
553   Traumatic Brain Injury in Mice. *J. Vis. Exp.* 64556 (2022) doi:10.3791/64556-v.

554 27. Lifshitz, J. *et al.* Clinical relevance of midline fluid percussion brain injury: Acute deficits, chronic
555   morbidities and the utility of biomarkers. *Brain Inj.* **30**, 1293–1301 (2016).

556 28. Aran, D. *et al.* Reference-based analysis of lung single-cell sequencing reveals a transitional
557   profibrotic macrophage. *Nat. Immunol.* **20**, 163–172 (2019).

558 29. Belinky, F. *et al.* PathCards: multi-source consolidation of human biological pathways. *Database J.*
559   *Biol. Databases Curation* **2015**, bav006 (2015).

560 30. Macheda, T., Roberts, K. & Bachstetter, A. D. Electromagnetic Controlled Closed-Head Model of Mild
561   Traumatic Brain Injury in Mice. *J. Vis. Exp. JoVE* (2022) doi:10.3791/64556.

562 31. Yarnell, A. M. *et al.* The Revised Neurobehavioral Severity Scale (NSS-R) for Rodents. *Curr. Protoc.*
563   *Neurosci.* **75**, 9.52.1-9.52.16 (2016).

564 32. Antonopoulos, C. *et al.* Caspase-8 as an Effector and Regulator of NLRP3 Inflammasome Signaling.
565   *J. Biol. Chem.* **290**, 20167–20184 (2015).

33. Clark, R. S. *et al.* Caspase-3 mediated neuronal death after traumatic brain injury in rats. *J.*
Neurochem. **74**, 740–753 (2000).

34. Bertheloot, D., Latz, E. & Franklin, B. S. Necroptosis, pyroptosis and apoptosis: an intricate game of
cell death. *Cell. Mol. Immunol.* **18**, 1106–1121 (2021).

35. Green, T. R. F. & Rowe, R. K. Quantifying microglial morphology: an insight into function. *Clin. Exp.*
Immunol. **216**, 221–229 (2024).

36. Morrison, H., Young, K., Qureshi, M., Rowe, R. K. & Lifshitz, J. Quantitative microglia analyses reveal
diverse morphologic responses in the rat cortex after diffuse brain injury. *Sci. Rep.* **7**, 13211 (2017).

37. Liddelow, S. A. *et al.* Neurotoxic reactive astrocytes are induced by activated microglia. *Nature* **541**,
481–487 (2017).

38. McCauley, S. R. *et al.* Patterns of Early Emotional and Neuropsychological Sequelae after Mild
Traumatic Brain Injury. *J. Neurotrauma* **31**, 914–925 (2014).

39. Chakraborty, R., Tabassum, H. & Parvez, S. NLRP3 inflammasome in traumatic brain injury: Its
implication in the disease pathophysiology and potential as a therapeutic target. *Life Sci.* **314**, 121352
(2023).

40. Hochheiser, I. V. *et al.* Directionality of PYD filament growth determined by the transition of NLRP3
nucleation seeds to ASC elongation. *Sci. Adv.* (2022).

41. Hu, X. *et al.* Role of Pyroptosis in Traumatic Brain and Spinal Cord Injuries. *Int. J. Biol. Sci.* **16**, 2042–
2050 (2020).

42. Mortezaee, K., Khanlarkhani, N., Beyer, C. & Zendedel, A. Inflammasome: Its role in traumatic brain
and spinal cord injury. *J. Cell. Physiol.* **233**, 5160–5169 (2018).

43. Dempsey, C. *et al.* Inhibiting the NLRP3 inflammasome with MCC950 promotes non-phlogistic
clearance of amyloid-β and cognitive function in APP/PS1 mice. *Brain. Behav. Immun.* **61**, 306–316
(2017).

44. Xu, X. *et al.* Selective NLRP3 inflammasome inhibitor reduces neuroinflammation and improves long-
term neurological outcomes in a murine model of traumatic brain injury. *Neurobiol. Dis.* **117**, 15–27
(2018).

45. Li, S. *et al.* Microglial NLRP3 inflammasome activates neurotoxic astrocytes in depression-like mice.
Cell Rep. **41**, 111532 (2022).

46. Witcher, K. G. *et al.* Traumatic brain injury-induced neuronal damage in the somatosensory cortex
causes formation of rod-shaped microglia that promote astrogliosis and persistent neuroinflammation.
Glia **66**, 2719–2736 (2018).

47. Susarla, B. T. S., Villapol, S., Yi, J.-H., Geller, H. M. & Symes, A. J. Temporal Patterns of Cortical
Proliferation of Glial Cell Populations after Traumatic Brain Injury in Mice. *ASN Neuro* **6**, AN20130034
(2014).

48. Nespoli, E. *et al.* Glial cells react to closed head injury in a distinct and spatiotemporally orchestrated
manner. *Sci. Rep.* **14**, 2441 (2024).

49. Franklin, B. S., Latz, E. & Schmidt, F. I. The intra- and extracellular functions of ASC specks.
Immunol. Rev. **281**, 74–87 (2018).

50. Franklin, B. S. *et al.* The adaptor ASC has extracellular and 'prionoid' activities that propagate
inflammation. *Nat. Immunol.* **15**, 727–737 (2014).

51. Sofroniew, M. V. Molecular dissection of reactive astrogliosis and glial scar formation. *Trends*
Neurosci. **32**, 638–647 (2009).

52. Wang, H. *et al.* Portrait of glial scar in neurological diseases. *Int. J. Immunopathol. Pharmacol.* **31**,
2058738418801406 (2018).

53. Venegas, C. *et al.* Microglia-derived ASC specks cross-seed amyloid-β in Alzheimer's disease.
Nature **552**, 355–361 (2017).

54. Friker, L. L. *et al.* β-Amyloid Clustering around ASC Fibrils Boosts Its Toxicity in Microglia. *Cell Rep.*
30, 3743-3754.e6 (2020).

55. Castro-Gomez, S. & Heneka, M. T. Innate immune activation in neurodegenerative diseases.
Immunity **57**, 790–814 (2024).

56. Zheng, R. *et al.* ASC specks exacerbate α‐synuclein pathology via amplifying NLRP3 inflammasome
activities. *J. Neuroinflammation* **20**, 26 (2023).

57. Graham, A., Livingston, G., Purnell, L. & Huntley, J. Mild Traumatic Brain Injuries and Future Risk of
Developing Alzheimer's Disease: Systematic Review and Meta-Analysis. *J. Alzheimers Dis.* **87**, 969–
979 (2022).

58. Tirosh, I. *et al.* Dissecting the multicellular ecosystem of metastatic melanoma by single-cell RNA-seq.
Science **352**, 189–196 (2016).

59. Watson, S., Porter, H., Sudbery, I. & Thompson, R. Modification of Seurat v4 for the Development of
a Phase Assignment Tool Able to Distinguish between G2 and Mitotic Cells. *Int. J. Mol. Sci.* **25**, 4589
(2024).

60. Sun, N. *et al.* Human microglial state dynamics in Alzheimer's disease progression. *Cell* **186**, 4386-
4403.e29 (2023).

61. Benayoun, B. A. *et al.* Remodeling of epigenome and transcriptome landscapes with aging in mice
reveals widespread induction of inflammatory responses. *Genome Res.* **29**, 697–709 (2019).

62. Wickham, H. *et al.* Welcome to the Tidyverse. *J. Open Source Softw.* **4**, 1686 (2019).

63. Blanco-Carmona, E. Generating publication ready visualizations for Single Cell transcriptomics using
SCpubr. Preprint at https://doi.org/10.1101/2022.02.28.482303 (2022).

64. Blighe, K., Rana, S. & Lewis, M. EnhancedVolcano: Publication-ready volcano plots with enhanced
colouring and labeling. https://github.com/kevinblighe/EnhancedVolcano (2018).

65. Kanneganti, T.-D. *et al.* Bacterial RNA and small antiviral compounds activate caspase-1 through
cryopyrin/Nalp3. *Nature* **440**, 233–236 (2006).

66. Young, K. & Morrison, H. Quantifying Microglia Morphology from Photomicrographs of
Immunohistochemistry Prepared Tissue Using ImageJ. *J. Vis. Exp.* 57648 (2018) doi:10.3791/57648.

67.   Lenhard, W. & Lenhard, A. Testing the Significance of Correlations. Preprint at

https://doi.org/10.13140/RG.2.1.2954.1367 (2014).

Acknowledgements

We thank Ildikó Rácz and Christina Ising for their help in obtaining approval from the local ethics committee for the animal experiments; Sean-Patrick Hermann for assistance with editing the final draft; Juan I. Muñoz, Paula Martorell, and Nadia Villacampa for valuable discussions; and the DZNE Light Microscopy Facility for support with imaging and analysis.

Author contributions

Conceptualization: SCG, MTH

Methodology: ASV, PBL, SCG, TL, SS, YD

Investigation: ASV, PBL, SCG, TL, SS, YD

Funding acquisition: SCG, MTH

Supervision: SCG, MTH

Writing of original draft: SCG, TL, YD

Writing, review, and editing: MTH, SCG, PB, TL, YD, YD, VS, EL

Funding information

PBL is supported by the Fonds National de la Recherche within the PEARL program (FNR/16745220)

SCG is supported by the Alzheimer Forschung Initiative e.V (Grant 21060), the Hertie Network of Excellence in Clinical Neuroscience (Grant 2021-1A-12), the Bonn Neuroscience Clinician Scientist Program (Neuro-aCSis), and the DFG (EXC2151–390873048).

TL was supported by Department of Biotherapy Cancer Center and State Key Laboratory of Biotherapy, West China Hospital, Sichuan University, Chengdu, China

Y.D. is supported by China Scholarship Council (CSC, File No. 202308310126).

Y.D. is supported by China Scholarship Council (CSC, File No. 201908510162).

Competing interests

M.T.H. is a scientific advisory board member at Alector the Dementia Discovery Fund, and Muna Therapeutics and has received honoraria for oral presentations from Pfizer, Novartis, Roche, Abbvie, and Biogen.

Additional information

Supplementary information the online version contains supplementary material

Correspondence and requests for materials should be addressed to Sergio Castro-Gomez or Michael T. Heneka.

ORCID

TL: 0000-0001-6114-8461

SC-G: 0000-0002-1581-474X

PBL:0000-0002-9096-4633

YD: 0000-001-9644-7461

YD: 0000-0001-5915-5355 (Yushuang)

VS: 0000-0003-1222-4928

EL: 0000-0003-1488-5666

MTH: 0000-0003-4996-1630

Fig. 1: Subacute *Asc (Pycard)* expression in microglia after a model for CHI.

a-b. Single-cell RNA sequencing (scRNA-seq) analysis of brain suspended cells from mice 7 days after mFPI. **a.** UMAP visualization showing clustering of different cell populations, with mFPI and control samples overlaid. **b.** Annotated cell types, including microglia, macrophages, and other immune and neural cells after cluster identification. **c.** Volcano plot depicting differentially expressed genes in microglia of control and mFPI samples. **d.** Dot plot expression analysis showing percent of expressed key inflammasome-related genes across various brain cell types, highlighting *Asc* (*Pycard*) expression in microglia, Natural Killers (NK) cells, macrophages and monocytes. **e.** UMAP projection of cells colored relative to the expression of canonical inflammasome-related genes *Nlrp3*, *Asc* (*Pycard*) *Casp1, Il1b, Il18 and Gsdmd*.

Fig. 2: **ASC contributes to the upregulation of inflammasome mediators following CHI.**

a. Immunoblot detection and **b-g.** densitometric semi-quantification of inflammasome-related proteins NLRP3, ASC, Caspase-1 (CASP1), IL-1β at 1, 7, and 21 dpi in contusional cortices from WT and $Asc^{-/-}$ mice following Sham or CHI. (left panels $n = 4$ per group and time point, *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$, Ordinary two-way ANOVA, Bonferroni's test. AU, arbitrary units. **h, i.** Levels of IL-1β and TNF-α in brain lysates from WT and $Nlrp3^{-/-}$ mice at 1, 7, and 21 dpi following CHI. $n = 3 - 4$ per group and time point, Left panels represent ordinary two-way ANOVA, Bonferroni's test, *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$. Right panels in **b-i** represent integrated relative concentrations (Area Under Curve, AUC), **$p < 0.01$, ***$p < 0.001$ by Student's t-test. dpi, days post-injury.

Fig. 3: Gliosis and morphology of Iba1⁺ and GFAP+ cells are modulated by ASC following CHI.

a. Immunohistochemistry of Iba1 (green, presumably microglia) in cortices from WT, and $Asc^{-/-}$ mice at 1, 7, and 21 dpi following CHI. Arrows indicate representative morphological changes. Scale bar, 20 μm. **b.** Representative skeletonized Iba1+ cells following CHI. Scale bars, 10 μm. Quantitative analysis of **c.** Iba1+ cell counts; **d.** branch numbers per cell, **e.** total branch length per cell and **f.** endpoint numbers per cell at 1, 7, and 21 dpi in WT, $Nlrp3^{-/-}$ and $Asc^{-/-}$ mice subjected to sham or CHI. **h.** Immunohistochemistry of GFAP (magenta, presumably astrocytes) in cortices from WT and $Asc^{-/-}$ mice at 1, 7, and 21 dpi following CHI. Arrows indicate representative morphological changes. Scale bar, 20 μm. **i.** Representative skeletonized GFAP+ cells following CHI. Scale bars, 10 μm. Quantitative analysis of **j.** GFAP+ cell counts; **k.** branch numbers per cell, **l.** total branch length per cell and **m.** endpoint numbers per cell at 1, 7, and 21 dpi in WT and $Asc^{-/-}$ mice subjected to sham or CHI. $n = 12$ slices from 4 mice per group per each time point, *$p < 0.05$, ***$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$ by ordinary two-way ANOVA with Bonferroni's post hoc test. Right panels show integrative AUC of each marker over time (dpi). *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$ by Student's t-test.

Fig. 4: Glial cell interactions are modulated by ASC following CHI.

a. Immunohistochemistry for Iba1 (green) and GFAP (magenta) in cortices of WT and $Asc^{-/-}$ mice at 7 and
21 dpi. Surface-surface colocalization analysis after 3D reconstruction was used to estimate contact areas
between Iba1+ cells and GFAP+ cells. Rectangles highlight examples of glial surface-surface
colocalization, shown in high magnification views. Scale bar, 20 µm. **b.** Quantification of contact areas
between Iba1+ cells and GFAP+ cells at 7 and 21 dpi in WT, and $Asc^{-/-}$ mice subjected to Sham or CHI. *n*
= 12 slices from 4 mice per group per each time point, $*p < 0.05$, $***p < 0.01$, $***p < 0.001$, $****p < 0.0001$
by ordinary two-way ANOVA with Bonferroni's post hoc test. Right panels show integrative AUC of each
marker over time (dpi). $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$ by Student's t-test.
733
Fig. 5: Constitutional ASC Deletion Protects from Mild Cognitive Impairments After CHI.
a. Experimental timeline for cognitive assessment following CHI, including spatial learning phase (10-15
dpi), probe trial (16 dpi), and cued learning (17-20 dpi) in the Morris water maze. **b-e.** Morris water maze
performance in WT and $Asc^{-/-}$ mice, showing path length and thigmotaxic across spatial **b-e** and **f-j.** cued
training days (n=10-11 mice per genotype $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$ by repeated
measurements two-way ANOVA with Tukey's post hoc test). Integrative AUC for path length and
thigmotaxic from spatial learning (**c** and **e**) and cued learning phase (**g** and **i**). **j.** Representative heatmaps
mean swimming trajectories and **k.** quantification of percentage of time spent in target in comparison to
another quadrants (a.o.) or **l.** platform crossing times in virtual target platform in comparison to another
viral platform (a.o.) (n=10-10 mice per Genotype, $*p < 0.05$, $**p < 0.01$, $***p < 0.001$ by ordinary one-way
and two-way ANOVA with Tukey's post-hoc tests). Data are presented as mean ± SEM.
745
Fig. 6: NLRP3 regulates ASC aggregation and distribution following CHI.
a. Immunohistochemistry for Iba1 (green), GFAP (magenta) and ASC (red) in cortices of WT and $Nlrp3^{-/-}$
mice at 7 and 21 dpi. ASC localization within Iba1+ cells (yellow spots), GFAP+ cells (blue spots), and
outside glial cells (red spots) were analyzed using IMARIS. Rectangles highlight examples of ASC
distribution shown in high magnification views. Scale bar, 20 µm. **b-e.** Quantification of total ASC
aggregates numbers (**b**), intracellular ASC aggregates within Iba1+ cells (**c**), ASC aggregates outside glia
cells (**d**), and ASC aggregates with a volume range 30–40 um^3 (**e**) at 7 and 21 dpi in WT and $Nlrp3^{-/-}$ mice
subjected to Sham or CHI. *n* = 12 slices from 4 mice per group per each time point, $*p < 0.05$, $***p < 0.01$,
$***p < 0.001$, $****p < 0.0001$ by ordinary two-way ANOVA with Bonferroni's post hoc test. Right panels
show integrative AUC of each marker over time (dpi). $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$ by
Student's t-test. **f-i.** Spearman's correlation analysis (*r*) with a linear regression model (black lines WT, red
lines $Nlrp3^{-/-}$) with an interaction term for ASC aggregates (total, within glial cells, and outside glial cells)
and glia cell contact areas. n = 36 slice from 12 mice per genotype, $*p < 0.05$, $**p < 0.01$, $***p < 0.001$,
$****p < 0.0001$. Differences between WT and $Nlrp3^{-/-}$ Spearman Correlation Coefficients (*r*) was performed
using the standard Fisher's z-transformation and subsequent comparison[67] $*p < 0.05$, $**p < 0.01$.
761
Supplementary Fig. 1: Single-Cell Transcriptomic Analysis of Brain Immune Cells Following TBI.

a. UMAP clustering of single-cell RNA sequencing (scRNA-seq) data, comparing control (red) and TBI

(green) samples, illustrating shifts in immune cell populations post-TBI. **b.** Heatmap showing gene

expression profiles across different brain cell types and cluster identification. **c.** Dot plot depicting the

expression of key inflammasome-related genes across different brain cell types. **d.** Dot plot depicting gene

expression profile of inflammasome-related genes in microglia after TBI.

Supplementary Fig. 2: ASC modulates the expression and cleavage of CASP-8 and CASP-3

following CHI.

a. Schematic of injury model/location. Coronal view (right) shows tip positioned over the skull midline.

Horizontal view (left) shows impact site relative to bregma. **b.** No significant motor, reflex or reaction

deficits at 1- or 7-days post-intervention (dpi) when the mice were evaluated using the Revised

Neurobehavioral Severity Scale (NSS-R). **c.** Representative immunoblot images of CAPS 8 and CASP 3

at 1, 7, and 21 dpi in peri-contusional cortices of mice subjected to Sham or CHI, comparing WT and *Asc* $^{-}$

$^{/-}$ groups. **d-e.** Quantification of expression of CASP 8 and CASP 3. The band intensity of a given target

protein was normalized to the corresponding GAPDH signal for each sample. Data were further

normalized to the average of the corresponding WT group and are presented as the mean ± SEM.

Statistical significance was determined using two-way ANOVA followed by Bonferroni's post-hoc tests, n =

4 mice per group per each time point, $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$ (left panels).

Quantification of area under the curve (AUC) of each target over time (dpi) in WT and $Asc^{-/-}$ mice is shown

in right panels. Statistical analysis was performed using an unpaired two-tailed t-test. Data are presented

as the mean ± SEM. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$.

Supplementary Fig. 3: The cellular source of IL-1 β Following mTBI.

a. Representative images of immunohistochemical staining of the peri-contusional cortices for IL-1β (red),

Iba1 (green), GFAP (green), and NeuN (green) in sham operation, 7*dpi* and 21 *dpi* groups. **b.** IMARIS was

used for the IL-1β analysis at 7*dpi* and 21 *dpi*. IL-1β (yellow spot) localization within Iba1+, GFAP, and

NeuN cells. Scale bar, 20 μm. **c-d.** The percentage and number of IL-1β spots expression was assessed

in Iba1- positive cells, GFAP-positive cells, and NeuN-positive cells at both 7 dpi and 21 dpi. Data are

presented as the mean ± SEM. Statistical significance was determined using two-way ANOVA followed by

Bonferroni's post-hoc tests, n = 12 slice (4 mice) per group per each time point, $*p < 0.05$, $**p < 0.01$, $***p$

< 0.001, $****p < 0.0001$.

Supplementary Fig. 4: Swimming velocity during cued learning in the Morris water maze.

Swimming velocity during cued learning in Sham WT, CHI-injured WT **a.** CHI-injured $Asc^{-/-}$. **b.** Comparison

of the integrated area under the curve of swimming velocity.

Supplementary Fig. 5: ASC antibody validation and ASC aggregates analysis.

a-b. Validation of ASC antibody specificity using immunofluorescence staining and immunoblotting. **a.**

ASC signal is detected in WT mice but absent in *Asc*$^{-/-}$ mice. **b.** Precipitated supernatant from LPS-

stimulated primary microglia and cortical tissue from CCI brain-injured mice were used as positive

controls. **c.** Representative immunohistochemical images of Iba1 (green) and ASC (red) in peri-

contusional cortex of WT and *Nlrp3*$^{-/-}$ mice at different time points (sham, 7dpi, 21 dpi). ASC aggregates

were analyzed and 3D reconstructed using IMARIS. Scale bar, 20 μm. **d-f.** Quantification of ASC

aggregation and ASC aggregates volume. Two-way ANOVA with Bonferroni's post-hoc tests (*p < 0.05,

**p < 0.01, ***p < 0.001, ****p < 0.0001). n = 12 slices (4 mice) per group per each time point. Data shown

as mean ± SEM.

a

Sham

	IBA1 GFAP	Contact	Merge
WT			
Asc⁻/⁻			

7 dpi

WT			
Asc⁻/⁻			

21dpi

WT			
Asc⁻/⁻			

b

○ WT
○ Asc⁻/⁻

Contact Area (μm²)

****

****
****

Sham   7 dpi   21 dpi

AUC Contact Area (μm² x 1000 / dpi)²

****

WT   Asc⁻/⁻