**Photo-based individual identification is more reliable than visible implant elastomer tags or toe-tipping in young agile frogs**

Edina Nemesházi[1] [*], Nikolett Ujhegyi[2,3], Zsanett Mikó[2,4], Andrea Kásler[2,4,5], Vera Lente[6], Veronika Bókony[2]

*[1]Konrad Lorenz Institute of Ethology, Department of Interdisciplinary Life Sciences, University of Veterinary Medicine Vienna, Savoyenstrasse 1a, AT-1160 Vienna, Austria*

*[2]Department of Evolutionary Ecology, Plant Protection Institute, HUN-REN Centre for Agricultural Research, Nagykovácsi út 26-30, HU-1029 Budapest, Hungary*

*[3]Department of Wildlife Biology and Management, Institute for Wildlife Management and Nature Conservation, Hungarian University of Agriculture and Life Sciences, Páter Károly u. 1, HU-2100 Gödöllő, Hungary*

*[4]Department of Systematic Zoology and Ecology, ELTE Eötvös Loránd University, Pázmány Péter sétány 1/C, HU-1117 Budapest, Hungary*

*[5]Doctoral School of Biology, Institute of Biology, ELTE Eötvös Loránd University, Pázmány Péter sétány 1/C, HU-1117 Budapest, Hungary*

*[6]Department of Freshwater Fish Ecology, Institute of Aquaculture and Environmental Safety, Hungarian University of Agriculture and Life Sciences, Páter Károly u. 1, HU-2100 Gödöllő, Hungary*

*[*]e-mail: nemeshazi.edina@protonmail.com*

**Abstract**

In amphibian capture-recapture studies, commonly used individual-identification methods include toe clipping as well as less invasive alternatives of varying cost. Yet, choosing the best method for a study is challenging, because both the reliability for identification and the severity of adverse effects of a given mark type can greatly vary between species as well as life stages. We compared the reliability of three identification methods in young agile frogs (*Rana dalmatina*): clipping a single phalanx, injecting visible-implant-elastomer (VIE) tags (one of six colours) under the skin, and photo-based identification using natural colouration. Individuals were regularly photographed from the start of metamorphosis onwards, and were marked by the other two methods soon after all of them finished metamorphosis. Subsequently, we checked mark retention by each method multiple times for more than a year. Photo-based identification was far more reliable than the other two methods: 100% identification success post-metamorphosis in all checking events within small housing groups, and 98% with computer-assisted identification across all housing groups. Post-metamorphic body colouration remained largely stable, and the major patterns were already present at metamorphosis. Based on VIE colour alone, we could correctly identify the animals in 77% of the checks. Green tags were the easiest to recognize. VIE often broke into smaller parts, and those were sometimes found only outside the originally tagged body part (15%) or not found at all (10%). VIE was better retained in the legs than in the arms. Dissection revealed that migrating VIE pieces can accumulate in the internal organs, especially the kidneys. The clipped toe was successfully recognized only in 41% of checks. We conclude that photo-based identification is superior to both VIE-tagging and toe tipping in young agile frogs, and this method should be preferred in future studies seeking a cost-efficient yet reliable identification method in this species.

## 1. Introduction

Tracking individuals across time and space is key in research and conservation programs. There are various methods of individual identification, and choosing the right method for a particular goal is crucial for the success of the study. For example, the anatomical, physiological and behavioural features of a species may influence the suitability of a given identification method [1]. Different methods can significantly differ in costs in terms of money and time (e.g. expensive but easy-to-read microchips vs. inexpensive but time-consuming photography-based identification), hence limiting the options of projects on tight budget. Ethical considerations require to take into account the 3Rs principle (refinement, reduction and replacement) [2], and the ideal method should minimise adverse effects while maximising identification accuracy. Therefore, assessing the reliability of identification methods is important before starting long-term application in large numbers.

For many decades, toe clipping was the most conventional method of individual identification in amphibians. This is an inexpensive and fast method, but negative effects have been reported on various welfare-related endpoints such as stress hormone levels and behavioral changes as well as on long-term growth and survival [3], and clipped toes can reportedly regrow over time which limits identification success [4–6]. Clipping only the most distal phalanx (or just the toe pad in hylid frogs; both known as 'toe tipping') was proposed as an alternative to reduce the adverse effects, and this method has been used in different species [7,8]. Yet, studies on both the actual reduction of adverse effects and the reliability of toe tipping for individual identification are missing. Methods that are considered less invasive require the injection of extraneous material under the skin [1]. One such material is visible implant elastomer (VIE), a colourful silicone-based material that is usually considered harmless, with no effect on stress, movement or growth [9–16]. However, reports from several amphibian studies suggest that VIE may migrate under the skin, its colour may fade with

time, and the material may even be rejected by the body, leading to identification failures

[4,9,17–19]. Retainment of VIE tags may depend on the body part in which it was applied

[4,18,20]. Studies that compared the reliability of toe clipping and VIE tagging in anurans

came to contradicting conclusions [4–6,9], suggesting that the suitability of each method for

individual identification depends on the species and circumstances of the study.

In contrast to the above methods, photo-based identification relies on the distribution of

natural macroscopic colour patterns of the amphibian skin. Such patterns remain largely stable

at least during adult life [21,22], thus providing reliable alternative to artificial tagging

methods in various species [23–26]. By far, this method is the least invasive, and it requires

only a suitable photographing device. However, when the number of individuals is more than

a handful, photo-based identification is time-consuming and difficult to use on-the-spot,

because the individual needs to be compared to all the previously taken photographs. Finding

the matching images can be aided by various software, some of which are pricey though, and

their suitability may greatly differ for a given set of images [24,25].

Reliability of different individual identification methods may greatly vary between adult

and young life stages. Because juveniles have faster growth rates and potentially better

regeneration capabilities, and skin coloration may change with ontogeny, methods that are

reliable in adults may not necessarily work as well in young animals [27,28]. In this study, we

assessed the reliability of three low-cost identification methods in young agile frogs (*Rana*

*dalmatina*): toe tipping (clipping only the most distal phalanx), VIE tagging, and photo-based

identification. We applied all three methods on recently metamorphosed froglets and followed

the retention of individual markings for 488 days, including one winter hibernation period.

We kept the frogs in small housing groups, where the application of only one VIE tag and one

toe mark per individual was required for following the animals' true identities. This enabled

us to record mark retention by each method with high accuracy. Our study demonstrated that

natural marks were far more stable than the applied artificial marks, and photo-based

individual identification was always successful in young agile frogs, both when comparing

small and large numbers of individuals.

**2. Methods**

*2.1. Data collection*

We used 113 agile frogs for this study, which we raised from the egg for the purposes of a

long-term experiment. The eggs were collected on 15 March 2023 from 10 egg masses

spawned naturally in our study pond (47.551195°N, 18.926682°E). The embryos and tadpoles

were raised as described in [29]; after the third week of larval development until

metamorphosis they were kept in mesocosms as described in [30], in groups of six siblings.

When an animal reached the start of metamorphosis (appearance of forelimbs), we assigned

an individual identifier ('animal ID') to it, photographed it in shallow water (either by digital

camera or smart phone), and housed it in an individually-labelled plastic box until marking.

After marking (see below), we housed the froglets in groups of five or six siblings outdoors,

except for the winter during which they hibernated in a climate chamber [31]. Group sizes

decreased with mortality during the experiment, and animals that died before marking or were

excluded due to other reasons (S1 Appendix) were not counted in the N=113 for the present

study. Details on housing conditions as well as animal exclusions from the study are described

in S1 Appendix.

After all individuals had completed metamorphosis, we photographed them and marked

them with both toe tipping and VIE tagging (Table 1). At tagging, the froglets had 0.6 – 1.3 g

body mass and 19.6 – 26.4 mm snout-vent length (SVL; measured by Fiji version 1.54g [32]

based on photographs). Each froglet was anesthetized by immersing in a 1 g/L water bath of

tricaine-methanesulfonate (MS-222; a safe and humane anaesthetic widely used for

amphibians [33–35]) buffered to neutral pH with the same amount of disodium hydrogen

phosphate. When reflexes were lost (after ca. 2 minutes in the room-temperature solution), we

placed the animal on wet paper towel for marking. We used flame-sterilized sharp scissors to

remove one toe tip (the most distal phalanx) following a toe-clip coding system (Fig 1), such

that every frog in a housing box would have a different toe tip removed. Additionally, we

injected a VIE tag (Northwest Marine Technology, Inc., Shaw Island, WA, USA) of one out of

six colours under the skin on one out of three body parts either on the left or the right side

(Fig 1) following the manufacturer´s guidelines (but see S2 Appendix Fig S1 for signs of

differential viscosity between the applied tags). Building on previous reports on varying tag

retention [4,18,20], we chose body parts that are relatively far away from the lymph sinuses:

the upper arm, the shin, and the foot at the edge of the sole. Another reason for choosing these

three body parts was that they are relatively easy to see without picking up the animal, which

might facilitate individual identification in the field, although in our study the tags were

always examined with the animal being held in hand. We did not tag the toes or foot webs

because these were too small. Length of the applied VIE tags varied between ca. 3 and 9 mm,

and they were in general ca. twice as long in the shin than in the upper arm, because tag size

was limited by the short length of the latter. Because VIE tags have been reported to move

under the skin of frogs, we applied VIE in the same body part for all individuals within a

housing box; therefore, the original body part remained unambiguously known for each

individual, regardless of later tag relocations. However, the combination of tag colour and

side (left/right) was different for all individuals housed in the same box. After marking, the

animal was placed on its belly with legs unfolded to a loosely stretched posture for better

visibility, and the full dorsal body surface was photographed along with its 'animal ID' using

a Canon EOS R50 camera. The froglets were kept wet until they recovered from anaesthesia;

all animals survived the procedures and woke up after ca. half an hour.

**Table 1. Timing of procedures and number of individuals (N) throughout the study.**

| Procedure | Date | N | Photo taken | VIE tag check | Toe-clip check |
|---|---|---|---|---|---|
| Metamorphosis | 29 May-25 Jun 2023 | 112* | + | - | - |
| Marking | 5-6 Jul 2023 | 113 | + | - | - |
| Checking | 29-30 Aug 2023 | 113 | + | + | + |
| | 6-7 Nov 2023 | 110 | + | + | + |
| | 13 February 2024** | 110 | + | - | - |
| | 13 Mar 2024 | 109 | + | + | - |
| | 15 May 2024 | 109 | + | + | - |
| | 15 July 2024 | 107 | + | + | - |
| | 16 Sep 2024 | 100 | + | + | + |
| | 4 Nov 2024 | 100 | + | + | - |

* Note that due to an error, photo taken at metamorphosis was missing for one individual,

therefore such photos were available for only 112 out of those 113 animals that we marked

and monitored in this study.

** We only recorded the natural markings on this date. Therefore, we excluded these images

from the analyses comparing the reliability of the three identification methods within housing

groups, and only included them in the HotSpotter database.

**Fig 1. Individual identification by toe tipping and VIE tagging.** Each individual was

marked by the removal of one phalanx and the injection of one VIE colour into one of the

highlighted body parts (A). VIE codes are highlighted with blue frame. Each marking code

indicates the side of the body (L: left, R: right). Additionally, toe-clip codes indicate the limb

(F: forelimb, H: hind limb) and digit (according to proximity to the body), while VIE codes

indicate the body part (A: upper arm, S: shin, F: foot). The applied VIE colours and whether

or not they are fluorescent under ultraviolet lighting are indicated on the right. Example

photos show LF-4 combined with RS-red (B) and RF-4 combined with LF-green (C) just after

tagging. Exposure curves were applied for each image to facilitate pattern visibility.

We evaluated the reliability of each identification method for over at least 14 months after

marking, up until the start of the froglets' second winter hibernation (Table 1). Checking

events took place every two months, except for the duration of winter hibernation. During

each checking event, we removed all frogs from each housing box in a haphazard order, and

to each animal we assigned a number between 1 and the total number of frogs still alive

('sampling ID'). Two researchers tried to identify the clipped toe and the colour and location

of the VIE tag for each frog. They searched for VIE tags both with and without a UV-A torch

light, because most of the VIE colours we used were fluorescent (Fig 1). At the first checking

event (August 2023), researchers were allowed to discuss their decisions regarding VIE tags,

while across the rest of the checking events, the two researchers independently assessed both

VIE tags and toe-tip clips. They also recorded if their identifications of VIE colours and

clipped toes were confident or uncertain. Lastly, they guessed the 'animal ID' for each froglet

('on-site ID') based on a combination of the available data, including the VIE tags, clipped

toes, and exclusion (i.e. when markings were not detected on a single frog in the box, that frog

could be identified as the only missing 'animal ID' in that box). Note that the sole purpose of

the 'on-site ID' was to ensure that we could determine the true identity of each animal, that is

essential for comparing mark retentions (see also the next paragraph). Because toe-tip clips

were highly ambiguous during the first two checking events (see Results), after that we

stopped evaluating mark retention by toe-tipping, except for the penultimate checking event

when the froglets weighed 2.7 – 7.5 g with 30.6 – 44.5 mm SVL (Table 1).

At each checking event, the froglets were photographed together with their 'sampling ID'

while positioning themselves freely in clean water [26] either by a Sony Cyber-shot DSC-

HX200V under varying light conditions indoors, or by a Canon EOS R50 digital camera

equipped with circular polar filter in a custom-developed photo box (light: 5000K Matcheasy

LED strip, 5V, 3m). Later, we brightened the images by applying exposure curves in

RawTherapee (version 5.9; https://rawtherapee.com) to enhance the visibility of skin patterns.

Then, an experienced researcher who was blind to the 'on-site ID' compared the images of

frogs in a housing box to the photographs captured at marking of all froglets housed in the

same box, and assigned a 'photo-based ID' to each individual at each checking event. Each

photo was compared with all individuals photographed at marking from the same housing box

(regardless of being identified already, and being dead or alive), so exclusion was never used

during photo-identification. VIE tags were rarely visible on the images taken during the

checking events; for such cases, the researcher was instructed to ignore VIE tags during

photo-identification and only consider the skin patterns of the frogs.

To assess the agreement of photo-identification within the small housing groups between

different observers, a subset of the images was evaluated by a second, less experienced

researcher. For this purpose, the 20 housing boxes were randomly distributed across the seven

checking events, such that a similar number of frogs would be chosen from each event, and

each of the 113 frogs would be chosen only once. Additionally, to investigate whether the

photographs allow individual identification already at the start of metamorphosis, these

photographs were compared to the images captured in August 2023; this was done by the

experienced researcher for all frogs (N = 113), and by the less experienced researcher for a

subset of two frogs picked randomly from each housing box (N = 40). For all photo-

identifications, the observers also recorded if they were confident or uncertain. For the photos

taken at metamorphosis, the experienced researcher scored picture quality as good, medium,

or poor (Fig S2), based on the visibility of skin markings due to both image properties (e.g.

sometimes the animal was not in focus or partially hidden by reflections on the water) and the

posture of the animal (e.g. the tail sometimes partially covered the legs). These quality issues

occurred because, for logistical feasibility, the metamorphs were photographed by several

researchers, none of whom had experience with capturing high-quality pictures for individual

identification. All photos taken after marking had good quality, with all skin markings being

fully visible, as all these pictures were taken by a single, experienced researcher.

To evaluate if the natural markings enabled individual identification in a large group of

juvenile frogs as well (and not just among five or six animals as above), we performed

computer-assisted identification in HotSpotter [36] using images from two different dates per

individual. To avoid disturbing effects of the background [26], we used images from those

two occasions when the animals were photographed in water against a uniform white

background: November 2023 and February 2024. First, we added one photo of each

individual from November and assigned the 'animal ID' to each image in the 'name' column

in HotSpotter. Then, we added a photo of each animal from February without assigning any

ID to it. To maximise the size of this dataset, we used images from all frogs still alive (N=112

in November 2023 and N=111 in February 2024, including two animals that were excluded

from all other analyses due to having two VIE tags). First, we carried out a 'software-

performance test' by running 'queries' in HotSpotter for the 111 individuals photographed in

February, when we recorded the rank of the true match from November for each queried

image. Meanwhile, the author who created the database checked if VIE tags were visible on

the included images, and listed those individuals whose images from neither November nor

February featured visible VIE (N=91). Finally, we performed an 'observer test' focusing on

the latter 91 animals to assess if human observers could identify the matching images solely

based on the natural colour markings of the animals. In the 'observer test', images from

February were haphazardly assigned to one of two authors who had no previous experience in

photo-based identification of these frogs (45 images queried by one and 46 queried by the

other author). Their task was to assign the 'animal ID' to each queried image by evaluating

the first 20 potential matches listed by HotSpotter (i.e. rank ≤ 20; a threshold also used in

previous studies [26, 27, 37]).

Mortality post-marking was low (N = 11) and happened during the warmest times (Table

1; note that two further individuals were taken out of the study after 5 and 6 checking events,

respectively, for reasons not related to this study). The froglets that died after marking were

dissected and examined for VIE material in their internal organs using an APOMIC SHD200

digital microscope.

*2.2. Data analysis*

First, we investigated which method yielded the most unambiguous decisions regardless of

the correctness of identification. To this end, we categorized each identification attempt as

'unambiguous' in cases where only a single, confident guess was made, and 'ambiguous'

where the observers were uncertain or in conflict with each other, or they could not find the

marking at all. We used a generalized linear mixed-effects model with binomial error and logit

link, where the dependent variable was the binary ambiguity category. The fixed factor was

the type of method (toe-clip, VIE tag, photo), and the random factor was 'animal ID' (inferred

from 'photo-based ID' as explained below). Because this analysis showed that both VIE and

toe-tip markings were often ambiguous (see Results), we could not use the 'on-site ID' as the

"true" individual identifier. Therefore, we used the 'photo-based ID' as the "true"

identification of 'animal ID' in the second set of analyses, considering the following reasons.

During photo-identification (using the images taken at marking as reference), the observers

were never in conflict and always found a putative match, which always matched the ID

assigned based on the only tag colour that was always unambiguously recognizable when

found (see Results). In each case where the 'on-site ID' did not match the 'photo-based ID',

we double-checked if there was any error made during photo-identification; in all these cases

the photo belonging to the non-matching 'on-site ID' clearly showed a different skin pattern

compared to the image taken at marking, meaning that the 'photo-based ID' was correct and

the 'on-site ID' (based on VIE and toe-tip clips) was mistaken.

To analyse the success rate of identification by toe-tip clips and VIE tags, we used the

same model structure as above, but the binary categories of the dependent variable were

'successful' (i.e. the assigned ID was the same as the 'photo-based ID') versus 'failed' (i.e. the

assigned ID was conflicted, or not found, or did not match the 'photo-based ID'). For toe-tip

clips, the fixed effects were the digit clipped (regardless of side) and the checking event (we

used the latter as categorical variable to allow for identification success to change non-linearly

over time). For VIE tags, the fixed effects were tag colour, the tagged body part (regardless of

side), and checking event. We calculated the identification success by VIE tags in three

alternative ways. First, we considered all identifications successful when the colour of the tag

was correctly recognized, regardless of tag location. Second, we considered all identifications

successful when the location of the tag was correct regardless of tag colour; tag location was

recorded as correct when it was found in the body part where it had been injected into, and

incorrect if it was not found at all or found in other part(s) of the body (i.e. further than the

nearest joints from the initial location). Third, we considered all identifications successful

when both the colour and location of the tag was correct (i.e. the right colour was found in the

body part where it had been injected).

For all statistical analyses, we used the R 4.3.2 software, with function 'glmer' of

package 'lme4' for binomial models. We used the 'DHARMa' package for residual

diagnostics to check that the statistical requirements of our models were met. From each

model, we calculated post-hoc pairwise comparisons using the 'emmeans' package, correcting

the P-values for multiple testing with the false-discovery rate method [38].

**2.3. Ethics statement**

This study was approved by the Ethics Committee of the Plant Protection Institute and

licensed by the Environment Protection and Nature Conservation Department of the Pest

County Bureau of the Hungarian Government (PE-06/KTF/00754-8/2022, PE-06/KTF/00754-

9/2022, PE-06/KTF/00754-10/2022, PE/EA/295-7/2018, PE/EA/00270-6/2023).

## 3. Results

Ambiguity rate differed significantly between each of the three methods: it was highest for toe

tipping and lowest for photo-identification (Table 2; $P < 0.0001$ for all pairwise differences).

The nature of ambiguity also varied by method (Table 2). For toe-tip clips, the two observers

often made conflicting identifications or recorded their guesses as uncertain (Table 2). For

VIE tags, these kinds of ambiguity were less frequent, but sometimes the tags were not found

at all (Table 2). For photo-identification based on the images taken at marking, ambiguity was

rare and always occurred in the form of the observers being somewhat uncertain about their

choice (Table 2). When the images taken at metamorphosis were used as the basis for photo-

identification, the observers were often uncertain and sometimes found conflicting or no

matches at all (Table 2). The dataset containing decisions for each individual at each checking

event is available on Figshare [39].

**Table 2. Percentage of ambiguous and unambiguous identifications for each marking method, regardless of correctness of the identification.**

| Method | N[*] | Unambiguous | Ambiguous Uncertain[†] | Conflicted[‡] | Not found[§] |
|---|---|---|---|---|---|
| **Toe-tip clips** | **323** | **37.2%** | **22.3%** | **40.2%** | **0.3%** |
| Front 2 | 62 | 46.8% | 40.3% | 12.9% | 0 |
| Front 3 | 53 | 20.8% | 49.1% | 28.3% | 1.9% |
| Front 4 | 56 | 35.7% | 44.6% | 19.6% | 0 |
| Hind 3 | 45 | 31.1% | 42.2% | 26.7% | 0 |
| Hind 4 | 54 | 31.5% | 42.6% | 25.9% | 0 |
| Hind 5 | 53 | 54.7% | 35.8% | 9.4% | 0 |
| **VIE tags** | **748** | **80.3%** | **4.3%** | **5.1%** | **10.3%** |
| Green | 122 | 97.5% | 0 | 0 | 2.5% |
| Yellow | 131 | 90.8% | 0.8% | 3.1% | 5.3% |
| Orange | 140 | 91.4% | 2.1% | 2.9% | 3.6% |
| Red | 128 | 60.9% | 10.2% | 7.8% | 21.1% |
| Pink | 117 | 74.4% | 8.5% | 13.7% | 3.4% |
| Purple | 110 | 63.6% | 4.5% | 3.6% | 28.2% |
| **Photos** compared to | | | | | |
| **Froglet marking** | **748** | **92.2%** | **7.8%** | **0** | **0** |
| Metamorphosis | 112 | 49.1% | 44.6% | 4.5% | 1.8% |

[*]Sample size (N) of identifications is the number of individuals × number of checking events. Note that for photos, the percentage of conflict was assessed on a subsample of N=113 based on the images taken at marking and N=40 based on the images taken at metamorphosis.

[†]Only one ID was assigned but with uncertainty.

[‡]More than one ID was deemed possible.

[§]No clipped toe or VIE tag or matching photo was found.

When the VIE tag was found, green was the only colour that was always recognized without ambiguity (Table 2). For the green-tagged individuals, every 'photo-based ID'

matched the guess based on the VIE tag (N = 119). For other tag colours, whenever there was

a mismatch between the 'photo-based ID' and the guess based on the VIE tag (N = 92 out of

552 individual checks where the tag was found), the photograph taken at marking always

matched the individual assigned by 'photo-based ID' and not the individual assigned by VIE

tag. Because of these reasons, we treated all identifications based on the images taken at

marking as correct, and we defined the success of the other methods based on the 'photo-

based ID'.

The overall success of identification by toe tipping was only 40.9%, varying from 15.1%

to 67.9% across digits (Fig 2) and from 35% during the first two checking events to 54% in

September before the second hibernation (Fig 2). For VIE tags, identification based on colour

was correct in 77.4% of cases, with significantly higher success for green, yellow, and orange

than for red, pink and purple tags (Fig 3). However, out of the 671 individual checks where

the tag was found, in 102 cases it was not located in the body part where it had been injected

(15.2%). In 77 cases (10.3%) involving 36 individuals the tag was not found at all, but in 16

of those individuals it was found again 1-4 checking events later (including 8 individuals in

which the tag disappeared and was found again more than once). Dissection of the froglets

that died or were taken out of the study revealed that, in 5 out of 13 animals, some of the VIE

material moved into the internal organs (Fig S3); this happened from all three body parts

where the tag had been injected. We recorded tag dislocation including complete

disappearance in 179 out of 748 individual checks (23.9%), and it was significantly more

frequent in the arms than in the legs (Fig 3), for the purple and red tags than for other colours

(Fig 3), and at later checking events (Fig 4). Altogether, when both the colour and the location

of the VIE tags were considered, identification success was 65.4% overall (Fig 3), decreasing

from 78.8% to 51% from the first to the last checking event (Fig 4).

**Fig 2. Identification success for marking by toe tipping, depending on the digit clipped**

**(A) and the timing of checking the markings (B).** Letters at the top of bar plots indicate

statistically significant differences (groups with different letters differ at $P < 0.05$ after

correcting for false-discovery rate).

**Fig**

**3. Identification success calculated in three different ways for marking by VIE tagging,**

**depending on tag colour (panels A, C and E) and tagged body part (panels B, D and F).**

355 Letters at the top of bar plots indicate statistically significant differences (groups with

356 different letters differ at P < 0.01 after correcting for false-discovery rate).

358
359 **Fig 4. dentification success calculated in three different ways for marking by VIE**

 **tagging, depending on the timing of checking the markings.** Letters at the top of bar plots

indicate statistically significant differences (groups with different letters differ at P < 0.01

after correcting for false-discovery rate).

When comparing each of 111 queried photos from February 2024 with all the other

images in the 'software-performance test', the image taken of the same animal three months

earlier was identified as the most likely match (rank=1) by HotSpotter for 75.7% of the

queried images, 92.8% received a rank ≤10, and 98.2% received a rank ≤ 20. Considering

only the results of those 91 individuals whose photos did not feature any signs of VIE, 73.6%

received rank=1, 92.3% received rank ≤ 10, and 98.9% received rank ≤ 20; the only

individual with rank > 20 received rank = 48. In the 'observer test', the 'animal ID' was

correctly assigned to each of the 90 individuals with rank ≤ 20, and the observer dealing with

the one individual whose matching pair was not present among the rank ≤ 20 images correctly

recognized that there was no 'animal ID' to assign. With other words, the overall false

rejection rate was less than 2% (due to software performance), with a false acceptance rate of

0% by the observers. The dataset containing all software-performance and observer results,

and screenshots from HotSpotter are available on Figshare [39].

Based on the photographs captured at metamorphosis, individual identification was

highly successful (93.8%) within the small groups of co-housed animals despite the relatively

high uncertainty reported by the observers. However, 47.3% of the photos taken at

metamorphosis had some problems with image quality or the body posture of the animal or

both, and 42.0% had poor quality (i.e. the problems resulted in low visibility of the leg

patterns). Ambiguity rate decreased as photo quality improved from poor (70.2%) through

medium (37.7%) to good (33.3%; $\chi^2$ test: $\chi^2 = 12.17$, df = 2, P = 0.002). Identification success

was 91.5% for poor-quality photos, 94.3% for medium-quality photos, and 100% for good-

quality photos. The dataset containing image qualities and identification results of

metamorphs is available on Figshare [39].

**4. Discussion**

In our study, photographs of the natural pigment markings on the skin were clearly superior to

both toe tipping and VIE tagging for individual identification of agile frogs during the first

one and a half years of their post-metamorphic life. When image quality was good, the

observers had low ambiguity when assessing potential matches, and the matches they

assigned were always correct when we were able to assess this with a reliable alternative

marking (i.e. green tags that were unambiguously detected in our study). Although the skin

patterns sometimes changed slightly as the animals aged (S2 Appendix Fig S4), the observers

were always able to find the correct match from up to six candidate individuals in our study

system. Reliability of computer-assisted identification in the large dataset (92% with rank $\leq$

10 and 99% with rank $\leq$ 20) was comparable to previous reports on successful photo-based

identification in amphibians [24–27, 39]. This result, together with the fact that all four

observers (two focusing on the small groups and two on the HotSpotter database) always

identified the matching photos if those were present among the list of potential matches that

they evaluated, demonstrates that natural colour markings of agile frogs are unique already in

their first year of life. Substantially different body postures on matching images likely

constrained the performance of HotSpotter in our study. By adding multiple images per

individual [25,40], and applying optimized techniques for image capturing while paying more

attention to animal positioning and lighting conditions [26], performance of computer-assisted

photo-identification is expected to improve even further. Indeed, using the latter approach, we

found in a parallel study on adult agile frogs that 100% accuracy can be achieved with

computer-assisted photo-identification in datasets consisting of several hundred individuals

[26]. Furthermore, in the present study, the photographs taken at metamorphosis (i.e.

emergence of front limbs) also proved suitable for individual identification of the young frogs

2-3 months later, when photos of ≤ 6 co-housed siblings were compared. When picture quality

was good, photo-based identification was 100% successful regardless of the timing of taking

the reference photographs (i.e. at metamorphosis or at froglet marking), and it was fairly high

even with the poorest-quality pictures taken at metamorphosis (91.5%). This suggests that the

unique individual patterns of agile frogs are already recognizable at very early life stages and

remain largely stable throughout and after metamorphosis (S2 Appendix Fig S5), similarly to

what has been found in a few other anuran species [41,42]. The drawback of the photo-

identification method is that comparison of the individual to the reference photographs

requires the availability of those photographs (e.g. on a computer at hand) and non-negligible

amount of time that increases with the number of potential matches that need to be checked.

For example, in our study, on-site assessment of VIE tags and toe-tip clips took qualitatively

similar time to photographing, but the latter had to be followed by off-site image processing

and subsequent photo comparison. Nevertheless, the time needed for finding the right match

can be sped up by computer-assisted photo-identification [43], using suitable software

[24,25,40] and optimized photography methods [23,26]. Earlier studies that used computer-

assisted photo-identification in amphibians had widely variable success rates (Table S1 in

[26]), which might be due to variability in the photography settings and software applied or

the stability of skin markings.

Overall, the second-best method in our study was VIE tagging. In our system, tag colour

alone should have been enough for successful individual identification because the same

colour was rarely injected into more than one animal in the same box, and the tags rarely ever

moved from one side of the body to the other. However, we had difficulty with telling apart

three of the six colours (red, pink, and purple), probably because the skin of the tagged body

parts is reddish-brown (S2 Appendix Figs S4 and S5) and sometimes can be quite dark. In this

respect, body parts with lightly coloured skin would be preferable for VIE tagging; however,

the only such area on agile frogs is the belly and the ventral side of the thigh, where tag

retention has been typically low in other species [18,20]. Furthermore, sometimes we did not

even find the tag, despite that we were often able to locate the same tags during some of the

subsequent checking events. The reason for the latter is that the tags often broke into several

pieces, and the pieces migrated away from the injection point both towards and away from the

centre of the body, often leaving only small, easy-to-miss dots close to the skin surface. These

findings are similar to those reported from studies on other amphibian species [18,19,44]. Tag

dislocation may further reduce success in situations where tag location is part of the

individual identification code; for example, we found that tags injected into the shin can move

into the foot and *vice versa*. In our study, dislocation was more frequent for tags injected into

the upper arm compared to the shin and foot; however, this contrasts findings in other anuran

species [20,28,45] and might be explained by tag size rather than by location, because we

injected smaller tags into the upper arms due to their smaller size relative to the two target

sites on the legs. However, larger tags were more likely to migrate in other amphibians [46].

Altogether, in our study, tag migration resulted in the 77% identification success based solely

on tag colour dropping to 65% based on the combination of tag colour and location. This

overall low success rate would be unacceptable for many studies; however, VIE tags could be

still reliable for applications that do not require a large number of individual codes. For

example, our data suggest that young agile frogs could be successfully batch-marked with

green, yellow, or orange VIE for tracking cohorts.

Our data on tag migration raise concerns about the health risks associated with VIE

tagging. Regardless of initial tag location and colour, some of the VIE material moved into the

internal organs, mostly the kidneys, which agrees with findings on adult cane toads (*Rhinella*

*marina*; [47]). In the latter study, VIE in the kidney was associated with inflammation and

damage to the vasculature that progressed in severity over time. This is especially worrisome

in light of growing concerns about the ecotoxicological effects of microplastics [48]. Notably,

although growth and survival were reported to be unaffected by VIE tagging in various

amphibians, this has been studied over relatively short periods, rarely longer than a year [9–

15]. Therefore, caution is recommended with using VIE especially in free-living wild animals,

and further research is needed to validate its long-term safety.

To our knowledge, this is the first report on mark retention by toe tipping under

controlled experimental conditions, despite its reported usage in the field [7,8]. Removal of

only the most distal phalanx yielded unacceptably low identification success in our study,

ranging between 15.1% (front 3) and 67.9% (hind 5) depending on the marked toe. This may

be due to particularly high regeneration capacities in young juveniles, although 100% toe-clip

retention was observed in boreal chorus frogs (*Psuedacris maculata*) over one year after

metamorphosis [9]. However, the length of the clipped part was not clarified in the latter

study; studies that reported high reliability of toe-clipping typically removed more than one

phalanx per toe (e.g. [4,44]) whereas we removed only the most distal phalanx. Nevertheless,

even if identification success would be increased by more extensive clipping, negative

consequences of toe-clipping on survival can increase with the severity of mutilation (e.g. by

the number of toes clipped per animal, which inevitably increases when a large number of

individuals have to be marked uniquely; [9,49, 50]).  Our results suggest that the practice of

clipping just one phalanx per toe may lead to the data turning out useless, and the mutilation

having been done in vain. As pointed out by a recent systematic review on the welfare impacts

of toe-clipping amphibians [3], data to help decisions in this ethically loaded topic are still too

scarce and heterogeneous. While the debate is still going on and more data are being

collected, we recommend using less invasive methods for marking individuals whenever

feasible.

In conclusion, our study shows that photo-based individual identification, besides serving

refinement according the 3Rs [2], is superior to both VIE-tagging and toe tipping for

individual identification in young agile frogs. When the images were taken with care to ensure

good quality, photo-identification had 100% success over one and a half years in the housing

groups, and major individually unique patterns were recognizable already at metamorphosis.

Although our animals were kept in small groups, the high stability of skin patterns and the

success of the computer-assisted identification for 91 froglets suggest that similarly exquisite

performance may be achieved with larger numbers of individuals. In contrast, both VIE-

tagging and toe tipping showed lower identification success, and the emergence of VIE in

internal organs suggests health risks. We encourage the usage of the most reliable and least

invasive, even though arguably most time-consuming photo-based method in future studies on

agile frogs whenever possible.

**Acknowledgements**

We thank Árpád Ferincz for the VIE material, and Emese Balogh, Judit Baumann, Zoltán

Örkényi, Mihály B. Rusz and all colleagues and students at the Department of Evolutionary

Ecology for their help during the study.

**References**

1.     Murray DL, Fuller MR. A critical review of the effects of marking on the biology of

vertebrates. In: Boitani L, Fuller TK, editors. Research Techniques in Animal Ecology:

Controversies and Consequences. New York: Columbia University Press; 2000. pp.

15–64.

2.    Russell WMS, Burch RL. The principles of humane experimental technique. UFAW,

Wheathampstead, UK: Universities Federation for Animal Welfare; 1992. Available:

https://caat.jhsph.edu/the-principles-of-humane-experimental-technique/

3.    Zemanova MA, Martín RL, Leenaars CHC. The impact of toe-clipping on animal

welfare in amphibians: A systematic review. Glob Ecol Conserv. 2025;59: e03582.

doi:10.1016/j.gecco.2025.e03582

4.    Hoffmann K, McGarrity ME, Johnson SA. Technology meets tradition: A combined

VIE-C technique for individually marking anurans. Appl Herpetol. 2008;5: 265–280.

doi:10.1163/157075408785911002

5.    Campbell TS, Irvin P, Campbell KR, Hoffmann K, Dykes ME, Harding AJ, et al.

Evaluation of a new technique for marking anurans. Appl Herpetol. 2009;6: 247–256.

doi:10.1163/157075409X420042

6.    Davis TM, Ovaska K. Individual recognition of amphibians: effects of toe clipping and

fluorescent tagging on the salamander *Plethodon vehiculum*. J Herp. 2001;35: 217–225.

7.    Phillott AD, Skerratt LF, McDonald KR, Lemckert FL, Hines HB, Clarke JM, et al.

Toe-clipping as an acceptable method of identifying individual anurans in mark

recapture studies. Herpetol Rev. 2007;38: 305–308.

8.    Kenyon N, Phillott AD, Alford RA. Evaluation of the photographic identification

method (PIM) as a tool to identify adult *Litoria genimaculata* (Anura: Hylidae).

Herpetol Conserv Biol. 2009;4: 403–410.

9.    Swanson JE, Bailey LL, Muths E, Chris Funk W. Factors influencing survival and

mark retention in postmetamorphic boreal chorus frogs. Copeia. 2013; 670–675.

doi:10.1643/CH-12-129

10.    Sapsford SJ, Roznik EA, Alford RA, Schwarzkopf L. Visible implant elastomer

marking does not affect short-term movements or survival rates of the treefrog litoria

rheocola. Herpetologica. 2014;70: 23–33. doi:10.1655/HERPETOLOGICA-D-13-0004

11.   Fouilloux CA, Garcia-Costoya G, Rojas B. Visible implant elastomer (VIE) success in

early larval stages of a tropical amphibian species. PeerJ. 2020;8: e9630.

doi:10.7717/peerj.9630

12.   Knapp DD, Diaz L, Unger S, Anderson CN, Spear SF, Williams LA, et al. Long-term

retention, readability, and health effects of visible implant elastomer (VIE) and visible

implant alpha (VI Alpha) tags in larval eastern hellbenders (*Cryptobranchus*

*alleganiensis alleganiensis*). J Herpetol. 2023;57: 133–141. doi:10.1670/22-011

13.   Bainbridge L, Stockwell M, Valdez J, Klop-toker K, Clulow S, Clulow J, et al.

Tagging tadpoles: retention rates and impacts of visible implant elastomer (VIE) tags

from the larval to adult amphibian stages. Herpetol J. 2015;25: 133–140.

14.   Phillips CT, Fries JN. An Evaluation of Visible Implant Elastomer for Marking the

Federally Listed Fountain Darter and the San Marcos Salamander. North Am J Fish

Manag. 2009;29: 529–532. doi:10.1577/m08-138.1

15.   Oropeza-Sánchez MT, Sandoval-Comte A, García-Bañuelos P, Hernández-López P,

Pineda E. Use of visible implant elastomer and its effect on the survival of an

endangered minute salamander. Anim Biodivers Conserv. 2020;43: 187–190.

doi:10.32800/abc.2020.43.0187

16.   Narayan EJ, Gramapurohit NP. Urinary corticosterone metabolite responses to capture

and visual elastomer tagging in the Asian toad (*Duttaphrynus melanostictus*). Herpetol

J. 2019;29: 179–183. doi:10.33256/hj29.3.179183

17.   Schmidt K, Schwarzkopf L. Visible implant elastomer tagging and toe-clipping:

Effects of marking on locomotor performance of frogs and skinks. Herpetol J. 2010;20:

99–105.

18.   Brannelly LA, Chatfield MWH, Richards-Zawacki CL. Visual implant elastomer (VIE)

559            tags are an unreliable method of identification in adult anurans. Herpetol J. 2013;23:

560            125–129.

561  19.   Heemeyer JL, Homyack JA, Haas CA. Retention and readability of visible implant

562            elastomer marks in eastern red-backed salamanders (*Plethodon cinereus*). Herpetol

563            Rev. 2007;38: 425–428.

564  20.   Moosman DL, Moosman Jr. PR. Subcutaneous movements of visible implant

565            elastomers in wood frogs (*Rana sylvatica*). Herpetol Rev. 2006;37: 300–301.

566  21.   Rojas B, Lawrence JP, Márquez R. Amphibian Coloration: Proximate Mechanisms,

567            Function, and Evolution. In: Moreno-Rueda G, Comas M, editors. Evolutionary

568            Ecology of Amphibians. Boca Raton: CRC Press; 2023. pp. 219–258.

569            doi:10.1201/9781003093312-12

570  22.   Price AC, Weadick CJ, Shim J, Rodd FH. Pigments, patterns, and fish behavior.

571            Zebrafish. 2008;5: 297–307. doi:10.1089/zeb.2008.0551

572  23.   Bendik NF, Morrison TA, Gluesenkamp AG, Sanders MS, O'Donnell LJ. Computer-

573            assisted photo identification outperforms Visible Implant Elastomers in an endangered

574            salamander, Eurycea tonkawae. PLoS One. 2013;8: e59424.

575            doi:10.1371/journal.pone.0059424

576  24.   Burgstaller S, Gollmann G, Landler L. The green toad example: A comparison of

577            pattern recognition software. North West J Zool. 2021;17: 96–99.

578  25.   Matthé M, Sannolo M, Winiarski K, Spitzen - van der Sluijs A, Goedbloed D,

579            Steinfartz S, et al. Comparison of photo-matching algorithms commonly used for

580            photographic capture–recapture studies. Ecol Evol. 2017;7: 5861–5872.

581            doi:10.1002/ece3.3140

582  26.   Nemesházi E, Mikó Z, Ujhegyi N, Kásler A, Lehofer N, Bókony V. Put the frog in

583            water: simple methods for improving individual identification as demonstrated with the

agile frog. Prepr BioRxiv. 2025. doi:10.1101/2025.02.25.640084

27.  Aevarsson U, Graves A, Carter KC, Doherty-Bone TM, Kane D, Servini F, et al.

Individual identification of the lake Oku clawed frog (*Xenopus longipes*) using a

photographic identification technique. Herpetol Conserv Biol. 2022;17: 67–75.

28.  Holcomb J, Lawrence K, Bailey S, Dittmer D. Considerations for and use of visible

implant elastomers to track and estimate abundance of juvenile boreal toads (Anaxyrus

boreas boreas) in Utah, USA. Herpetlogical Rev. 2022;53: 579–585.

29.  Bókony V, Balogh E, Mikó Z, Kásler A, Örkényi Z, Ujhegyi N. Higher sex-reversal

rate of urban frogs in a common-garden experiment suggests adaptive microevolution.

Evol Appl. 2025;18: e70093. doi:10.1111/eva.70093

30.  Bókony V, Üveges B, Móricz ÁM, Hettyey A. Competition induces increased toxin

production in toad larvae without allelopathic effects on heterospecific tadpoles. Funct

Ecol. 2018;32: 667–675. doi:10.1111/1365-2435.12994

31.  Kásler A, Holly D, Herczeg D, Ujszegi J, Hettyey A. Chytridiomycosis and climate

change: exposure to *Batrachochytrium dendrobatidis* and mild winter conditions do not

increase mortality in juvenile agile frogs during hibernation. Anim Conserv. 2023;26:

654–662. doi:10.1111/acv.12851

32.  Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji:

an open-source platform for biological-image analysis. Nat Methods. 2012;9: 676–682.

doi:10.1038/nmeth.2019

33.  Mitchell MA. Anesthetic considerations for amphibians. J Exot Pet Med. 2009;18: 40–

49. doi:10.1053/j.jepm.2008.11.006

34.  Downes H. Tricaine anesthesia in Amphibia: a review. Bull Assoc Reptil Amphib Vet.

1995;5: 11–16. doi:10.5818/1076-3139.5.2.11

35.  Hadfield CA, Whitaker BR. Amphibian emergency medicine and care. Semin Avian

Exot Pet Med. 2005;14: 79–89. doi:10.1053/j.saep.2005.04.003

36.   Crall JP, Stewart C V., Berger-Wolf TY, Rubenstein DI, Sundaresan SR. HotSpotter-

Patterned species instance recognition. Proc IEEE Work Appl Comput Vis. 2013; 230–

237. doi:10.1109/WACV.2013.6475023

37.   Morrison TA, Keinath D, Estes-Zumpf W, Crall JP, Stewart C V. Individual

identification of the endangered Wyoming Toad *Anaxyrus baxteri* and implications for

monitoring species recovery. J Herpetol. 2016;50: 44–49. doi:10.1670/14-155

38.   Pike N. Using false discovery rates for multiple comparisons in ecology and evolution.

Methods Ecol Evol. 2011;2: 278–282. doi:10.1111/j.2041-210X.2010.00061.x

39.   Nemesházi E, Ujhegyi N, Mikó Z, Kásler A, Lente V, Bókony V. Dataset for: Photo-

based individual identification is more reliable than visible implant elastomer tags or

toe tipping in young agile frogs. Figshare. 2025. doi:10.6084/m9.figshare.29320697

40.   Patel NG, Das A. Shot the spots: a reliable field method for individual identification of

*Amolops formosus* (Anura, Ranidae). Herpetozoa. 2020;33: 7–15.

doi:10.3897/HERPETOZOA.33.E47279

41.   Kenyon N, Phillott AD, Ross AA. Temporal variation in dorsal patterns of juvenile

green-eyed tree frogs, *Litoria genimaculata* (Anura: Hylidae). Herpetol Conserv Biol.

2010;5: 126–131.

42.   Bardier C, Székely D, Augusto-Alves G, Matínez-Latorraca N, Schmidt BR,

Cruickshank SS. Performance of visual vs. software-assisted photo-identification in

mark-recapture studies: a case study examining different life stages of the Pacific

Horned Frog (*Ceratophrys stolzmanni*). Amphibia-Reptilia. 2020;42: 17–28.

doi:10.1163/15685381-BJA10025

43.   Davis HP, Vancompernolle M, Dickens J. Effectiveness and reliability of photographic

identification methods for identifying individuals of a cryptically patterned toad.

634   Herpetol Conserv Biol. 2020;15: 204–211.

635 44. Brannelly LA, Berger L, Skerratt LF. Comparison of three widely used marking

636   techniques for adult anuran species *Litoria verreauxii alpina*. Herpetol Conserv Biol.

637   2014;9: 428–435.

638 45. Lunghi E, Bruni G, Manenti R, Ficetola GF. Use of visible implant elastomer on two

639   amphibians orders (Anura and Caudata): data on efficiency and reliability. Atti X

640   Congr Naz Soc Herpetol Ital. 2015;15: 163–167.

641 46. Grant EHC. Visual implant elastomer mark retention through metamorphosis in

642   amphibian larvae. J Wildl Manage. 2008;72: 1247–1252. doi:10.2193/2007-183

643 47. Cabot ML, Troan B V., Heugten KA Van, Schnellbacher RW, Smith D, Ridgley F, et

644   al. Migration and histologic effects of visible implant elastomer (VIE) and passive

645   integrated transponder (PIT) tags in the marine toad (*Rhinella marina*). Animals.

646   2021;11: 3255. doi:10.3390/ani11113255

647 48. Li C, Busquets R, Campos LC. Assessment of microplastics in freshwater systems: a

648   review. Sci Total Environ. 2020;707: 135578. doi:10.1016/j.scitotenv.2019.135578

649 49. Grafe TU, Stewart MM, Lampert KP, Rödel M. Putting toe clipping into perspective: a

650   viable method for marking anurans. J Herpetol. 2011;45: 28–35.

651 50. Waddle JH, Rice KG, Mazzotti FJ, Percival HF. Modeling the effect of toe clipping on

652   treefrog survival: Beyond the return rate. J Herpetol. 2008;42: 467–473.

653   doi:10.1670/07-265.1

655 **Supporting Information**

656 **S1 Appendix. Details of rearing conditions and exclusions from the experiment.**

657 **S2 Appendix. Supplementary figures.**