Photo-based individual identification is more reliable than visible implant elastomer tags

or toe clipping in young agile frogs

Edina Nemesházi[1] [*], Nikolett Ujhegyi[2,3], Zsanett Mikó[2,4], Andrea Kásler[2,4,5], Vera Lente[6],

Veronika Bókony[2]

[1]*Konrad Lorenz Institute of Ethology, Department of Interdisciplinary Life Sciences,*

University of Veterinary Medicine Vienna, Savoyenstrasse 1a, AT-1160 Vienna, Austria

[2]*Department of Evolutionary Ecology, Plant Protection Institute, HUN-REN Centre for*

Agricultural Research, Nagykovácsi út 26-30, HU-1029 Budapest, Hungary

[3]*Department of Wildlife Biology and Management, Institute for Wildlife Management and*

Nature Conservation, Hungarian University of Agriculture and Life Sciences, Páter Károly u.

1, HU-2100 Gödöllő, Hungary

[4]*Department of Systematic Zoology and Ecology, ELTE Eötvös Loránd University, Pázmány*

Péter sétány 1/C, HU-1117 Budapest, Hungary

[5]*Doctoral School of Biology, Institute of Biology, ELTE Eötvös Loránd University, Pázmány*

Péter sétány 1/C, HU-1117 Budapest, Hungary

[6]*Department of Freshwater Fish Ecology, Institute of Aquaculture and Environmental Safety,*

Hungarian University of Agriculture and Life Sciences, Páter Károly u. 1, HU-2100 Gödöllő,

Hungary

*e-mail: nemeshazi.edina@protonmail.com

Author contributions

This study was conceptualized and led by E.N. All authors participated in data collection.

Statistical analyses were conducted by V.B., data visualization was performed by E.N. and

V.B. The first draft was written by E.N. and V.B.; all authors were involved in review and

editing.

Keywords:

amphibian individual identification, VIE tagging, toe removal, melanin-based patterns in

juvenile frogs, long-term tag retainment

Abstract

In amphibian capture-recapture studies, commonly used individual-identification methods include toe clipping as well as less invasive alternatives of varying cost. Yet, choosing the best method for a study is challenging, because both the reliability for identification and the severity of adverse effects of a given mark type can greatly vary between species as well as life stages. We compared the reliability of three identification methods in young agile frogs (*Rana dalmatina*): clipping a single toe tip, injecting visible-implant-elastomer (VIE) tags (one of six colours) under the skin, and photo-based identification using natural colouration. Individuals were regularly photographed from the start of metamorphosis onwards, and were marked by the other two methods soon after all of them finished metamorphosis. Subsequently, we checked the presence of both the artificial and the natural marks multiple times for more than a year. Photo-based identification was far more reliable than the other two methods: 100 % identification success in all checking events across the entire study. Post-metamorphic body colouration remained largely stable, and the major patterns were already present at metamorphosis. Based on VIE colour alone, we could correctly identify the animals in 77 % of the checks. Green tags were the easiest to recognize. VIE often broke into smaller parts, and those were sometimes found only outside the originally tagged body part (15 %), or not found at all (10%). VIE was better retained in the legs than in the arm. Dissection revealed that migrating VIE pieces can accumulate in the internal organs, especially the kidneys. The clipped toe was successfully recognized only in 41% of the checks. We conclude that photo-based identification is superior to both VIE-tagging and toe clipping in young agile frogs, and this method should be preferred in future studies seeking a cost-efficient yet reliable identification method in this species.

Introduction

Tracking individuals across time and space is key in research and conservation programs. There are various methods of individual identification, and choosing the right method for a particular goal is crucial for the success of the study. For example, the anatomical, physiological and behavioural features of a species may influence the suitability of a given identification method [1]. Different methods can significantly differ in costs in terms of money and time (e.g. expensive but easy-to-read microchips vs. inexpensive but time-consuming photography-based identification), hence limiting the options of projects on tight budget. Therefore, assessing the reliability of identification methods is important before starting long-term application in large numbers.

For many decades, toe clipping was the most conventional method of individual identification in amphibians. This is an inexpensive and fast method, but negative effects have been reported on various welfare-related endpoints as well as on long-term growth and survival [2], and toe tips can reportedly regrow over time which limits identification success [3–5]. However, clipping only the last phalanx and only a small number of toes may reduce the adverse effects [6]. Methods that are considered less invasive require the injection of extraneous material under the skin [1]. One such material is visible implant elastomer (VIE), a colourful silicone-based material that is usually considered harmless [7–13]. However, reports from several amphibian studies suggest that VIE may migrate under the skin, its colour may fade with time, and the material may even be rejected by the body, leading to identification failures [3,7,14–16]. Retainment of VIE tags may depend on the body part in which it was applied [3,15,17]. Studies that compared the reliability of toe clipping and VIE tagging in anurans came to contradicting conclusions [3–5,7], suggesting that the suitability of each method for individual identification depends on the species and circumstances of the study.

In contrast to the above methods, photo-based identification relies on the distribution of natural macroscopic colour patterns of the amphibian skin. Such patterns remain largely stable at least during adult life [18,19], thus providing reliable alternative to artificial tagging methods in various species [20–23]. By far, this method is the least invasive, and it requires only a suitable photographing device. However, when the number of individuals is more than a handful, photo-based identification is time-consuming and difficult to use on-the-spot, because the individual needs to be compared to all the previously taken photographs. Finding the matching images can be aided by various software, some of which are pricey though, and their suitability may greatly differ for a given set of images [21,22].

Most studies that assessed the reliability of individual identification methods in

amphibians focused on adults, and relatively little is known about their performance in

younger life stages. Because juveniles have faster growth rates and potentially better

regeneration capabilities, and skin coloration may change with ontogeny, methods that are

reliable in adults may not necessarily work as well in young animals [24,25]. In this study, we

assessed the reliability of the three least expensive identification methods in young agile frogs

(*Rana dalmatina*): toe clipping, VIE tagging, and photo-based identification. We applied all

three methods on recently metamorphosed froglets and followed the retention of individual

markings for 488 days, including one winter hibernation period.

Methods

Data collection

We used 113 agile frogs for this study, which we raised from the egg for the purposes of a

long-term experiment. The eggs were collected on 15 March 2023 from 10 egg masses

spawned naturally in our study pond (47.551195°N, 18.926682°E). The embryos and tadpoles

were raised as described in [26]; after the third week of larval development until

metamorphosis they were kept in mesocosms as described in [27], in groups of six siblings.

When an animal reached the start of metamorphosis (appearance of forelimbs), we assigned

an individual identifier ('animal ID') to it, photographed it in shallow water (either by digital

camera or smart phone), and housed it in an individually-labelled plastic box until marking.

After marking (see below), we housed the froglets in groups of five or six siblings outdoors,

except for the winter during which they hibernated in a climate chamber [28]. Group sizes

decreased with mortality during the experiment, and animals that died before marking or were

excluded due to other reasons (Appendix S1) were not counted in the N=113 for the present

study. Details on housing conditions as well as animal exclusions from the study are described

in Appendix S1.

After all individuals had completed metamorphosis, we photographed them and marked

them with both toe clipping and VIE tagging (Table 1). At tagging, the froglets had 0.6 – 1.3 g

body mass and 19.6 – 26.4 mm snout-vent length (SVL; measured by Fiji version 1.54g [29]

based on photographs). Each froglet was anesthetized by immersing in a 1 g/L water bath of

tricaine-methanesulfonate (MS-222) buffered to neutral pH with the same amount of

disodium hydrogen phosphate. When reflexes were lost (after ca. 2 minutes in the room-

temperature solution), we placed the animal on wet paper towel for marking. We used flame-

sterilized sharp scissors to remove one toe tip (the most distal phalanx) following a toe-clip

coding system (Fig 1), such that every frog in a housing box would have a different toe-tip

removed. Additionally, we injected a VIE tag (Northwest Marine Technology, Inc., Shaw

Island, WA, USA) of one out of six colours under the skin on one out of three body parts

either on the left or the right side (Fig 1) following the manufacturer´s guidelines (but see

Appendix S2 Fig S1 for signs of differential viscosity between the applied tags). Building on

previous reports on varying tag retention [3,15,17], we chose body parts that are relatively far

away from the lymph sinuses: the upper arm, the shin, and the foot at the edge of the sole.

Another reason for choosing these three body parts was that they are relatively easy to see

without picking up the animal, which might facilitate individual identification in the field,

although in our study the tags were always examined with the animal being held in hand. We

did not tag the toes or foot webs because these were too small. Length of the applied VIE tags

varied between c.a. 3 and 9 mm, and they were in general c.a. twice as long in the shin than in

the upper arm, because tag size was limited by the short length of the latter. Because VIE tags

have been reported to move under the skin of frogs, we applied VIE in the same body part for

all individuals within a housing box; therefore, the original body part remained

unambiguously known for each individual, regardless of later tag relocations. However, the

combination of tag colour and side (left/right) was different for all individuals housed in the

same box. After marking, the animal was placed on its belly with legs unfolded to a loosely

stretched posture for better visibility, and the full dorsal body surface was photographed along

with its 'animal ID' using a Canon EOS R50 camera. The froglets were kept wet until they

recovered from anaesthesia; all animals survived the procedures and woke up after ca. half an

hour.

We evaluated the reliability of each identification method for over at least 14 months after

marking, up until the start of the froglets' second winter hibernation (Table 1). Checking

events took place every two months, except for the duration of winter hibernation. During

each checking event, we removed all frogs from each housing box in a haphazard order, and

to each animal we assigned a number between 1 and the total number of frogs still alive

('sampling ID'). Two researchers tried to identify the clipped toe and the colour and location

of the VIE tag for each frog. They searched for VIE tags both with and without a UV-A torch

light, because most of the VIE colours we used were fluorescent (Fig 1). At the first checking

event (August 2023), researchers were allowed to discuss their decisions regarding VIE tags,

while across the rest of the checking events, the two researchers independently assessed both

VIE tags and toe-clips. They also recorded if their identifications of VIE colours and clipped

toes were confident or uncertain. Lastly, they guessed the 'animal ID' for each froglet ('on-

site ID') based on a combination of the available data, including the VIE tags, clipped toes,

and exclusion (i.e. when markings were not detected on a single frog in the box, that frog

could be identified as the only missing 'animal ID' in that box). Because toe-clips were highly

ambiguous during the first two checking events (see Results), after that we stopped evaluating

toe-clips, except for the penultimate checking event when the froglets weighed 2.7 – 7.5 g

with 30.6 – 44.5 mm SVL (Table 1).

At each checking event, the froglets were photographed together with their 'sampling ID'

while positioning themselves freely in clean water [23] either by a Sony Cyber-shot DSC-

HX200V under varying light conditions indoors, or by a Canon EOS R50 digital camera

equipped with circular polar filter in a custom-developed photo box (light: 5000K Matcheasy

LED strip, 5V, 3m). Later, we brightened the images by applying exposure curves in

RawTherapee (version 5.9; https://rawtherapee.com) to enhance the visibility of skin patterns.

Then, an experienced researcher who was blind to the 'on-site ID' compared the images of

frogs in a housing box to the photographs captured at marking of all froglets housed in the

same box, and assigned a 'photo-based ID' to each individual at each checking event. VIE

tags were rarely visible on the images taken during the checking events; for such cases, the

researcher was instructed to ignore VIE tags during photo-identification and only consider the

skin patterns of the frogs.

To assess the agreement of photo-identification between different observers, a subset of

the images was evaluated by a second, less experienced researcher. For this purpose, the 20

housing boxes were randomly distributed across the seven checking events, such that a similar

number of frogs would be chosen from each event, and each of the 113 frogs would be chosen

only once. Additionally, to investigate whether the photographs allow individual identification

already at the start of metamorphosis, these photographs were compared to the images

captured in August 2023; this was done by the experienced researcher for all frogs (N = 113),

and by the less experienced researcher for a subset of two frogs picked randomly from each

housing box (N = 40). For all photo-identifications, the observers also recorded if they were

confident or uncertain. For the photos taken at metamorphosis, the experienced researcher

scored picture quality as good, medium, or poor (Fig S2), based on the visibility of skin

markings due to both image properties (e.g. sometimes the animal was not in focus or

partially hidden by reflections on the water) and the posture of the animal (e.g. the tail

sometimes partially covered the legs). These quality issues occurred because, for logistical

feasibility, the metamorphs were photographed by several researchers, none of whom had

experience with capturing high-quality pictures for individual identification. All photos taken

after marking had good quality, with all skin markings being fully visible, as all these pictures
were taken by a single, experienced researcher.

Mortality post-marking was low (N = 11) and happened during the warmest times (Table
1; note that two further individuals were taken out of the study after 5 and 6 checking events,
respectively, for reasons not related to this study). The froglets that died after marking were
dissected and examined for VIE material in their internal organs using an APOMIC SHD200
digital microscope.

This study was approved by the Ethics Committee of the Plant Protection Institute and
licensed by the Environment Protection and Nature Conservation Department of the Pest
County Bureau of the Hungarian Government (PE-06/KTF/00754-8/2022, PE-06/KTF/00754-
9/2022, PE-06/KTF/00754-10/2022, PE/EA/295-7/2018, PE/EA/00270-6/2023).

Data analysis
First, we investigated which method yielded the most unambiguous decisions regardless of
the correctness of identification. To this end, we categorized each identification attempt as
'unambiguous' in cases where only a single, confident guess was made, and 'ambiguous'
where the observers were uncertain or in conflict with each other, or they could not find the
marking at all. We used a generalized linear mixed-effects model with binomial error and logit
link, where the dependent variable was the binary ambiguity category. The fixed factor was
the type of method (toe-clip, VIE tag, photo), and the random factor was 'animal ID' (inferred
from 'photo-based ID' as explained below). Because this analysis showed that both VIE and
toe-clip markings were often ambiguous (see Results), we could not use the 'on-site ID' as the
"true" individual identification. Therefore, we used the 'photo-based ID' as the "true"
identification of 'animal ID' in the second set of analyses, considering the following reasons.
During photo-identification (using the images taken at marking as reference), the observers
were never in conflict and always found a putative match, which always matched the ID
assigned based on the only tag colour that was always unambiguously recognizable when
found (see Results). In each case where the 'on-site ID' did not match the 'photo-based ID',
we double-checked that there was no error made during photo-identification; in all these cases
the photo belonging to the non-matching 'on-site ID' clearly showed a different skin pattern
compared to the image taken at marking, meaning that the 'photo-based ID' was correct and
the 'on-site ID' (based on VIE and toe-clips) was mistaken.

To analyse the success rate of identification by toe-clips and VIE tags, we used the same
model structure as above, but the binary categories of the dependent variable were

'successful' (i.e. the assigned ID was the same as the 'photo-based ID') versus 'failed' (i.e. the

assigned ID was conflicted, or not found, or did not match the 'photo-based ID'). For toe-

clips, the fixed effects were the digit clipped (regardless of side) and the checking event (we

used the latter as categorical variable to allow for identification success to change non-linearly

over time). For VIE tags, the fixed effects were tag colour, the tagged body part (regardless of

side), and checking event. We calculated the identification success by VIE tags in three

alternative ways. First, we considered all identifications successful when the colour of the tag

was correctly recognized, regardless of tag location. Second, we considered all identifications

successful when the location of the tag was correct regardless of tag colour; tag location was

recorded as correct when it was found in the body part where it had been injected into, and

incorrect if it was not found at all or found in other part(s) of the body (i.e. further than the

nearest joints from the initial location). Third, we considered all identifications successful

when both the colour and location of the tag was correct (i.e. the right colour was found in the

body part where it had been injected).

For all statistical analyses, we used the R 4.3.2 software, with function 'glmer' of

package 'lme4' for binomial models. We used the 'DHARMa' package for residual

diagnostics to check that the statistical requirements of our models were met. From each

model, we calculated post-hoc pairwise comparisons using the 'emmeans' package, correcting

the P-values for multiple testing with the false-discovery rate method [30].

Results

Ambiguity rate differed significantly between each of the three methods: it was highest for

toe-clips and lowest for photo-identification (Table 2; $P < 0.0001$ for all pairwise differences).

The nature of ambiguity also varied by method (Table 2). For toe-clips, the two observers

often made conflicting identifications or recorded their guesses as uncertain (Table 2). For

VIE tags, these kinds of ambiguity were less frequent, but sometimes the tags were not found

at all (Table 2). For photo-identification based on the images taken at marking, ambiguity was

rare and always occurred in the form of the observers being somewhat uncertain about their

choice (Table 2). When the images taken at metamorphosis were used as the basis for photo-

identification, the observers were often uncertain and sometimes found conflicting or no

matches at all (Table 2). The dataset containing decisions for each individual at each checking

event is available from the authors upon request (it will be publicly shared once the final

article is published).

When the VIE tag was found, green was the only colour that was always recognized

without ambiguity (Table 2). For the green-tagged individuals, every 'photo-based ID'

matched the guess based on the VIE tag (N = 119). For other tag colours, whenever there was

a mismatch between the 'photo-based ID' and the guess based on the VIE tag (N = 92 out of

552 individual checks where the tag was found), the photograph taken at marking always

matched the individual assigned by 'photo-based ID' and not the individual assigned by VIE

tag. Because of these reasons, we treated all identifications based on the images taken at

marking as correct, and we defined the success of the other methods based on the 'photo-

based ID'.

The overall success of identification by toe-clips was only 40.9 %, varying from 15.1 %

to 67.9 % across digits (Fig 2) and from 35 % during the first two checking events to 54 % in

September before the second hibernation (Fig 2). For VIE tags, identification based on colour

was correct in 77.4 % of cases, with significantly higher success for green, yellow, and orange

than for red, pink and purple tags (Fig 3). However, out of the 671 individual checks where

the tag was found, in 102 cases it was not located in the body part where it had been injected

(15.2 %). In 77 cases (10.3 %) involving 36 individuals the tag was not found at all, but in 16

of those individuals it was found again 1-4 checking events later (including 8 individuals in

which the tag disappeared and was found again more than once). Dissection of the froglets

that died or were taken out of the study revealed that, in 5 out of 13 animals, some of the VIE

material moved into the internal organs (Fig S3); this happened from all three body parts

where the tag had been injected. We recorded tag dislocation including complete

disappearance in 179 out of 748 individual checks (23.9 %), and it was significantly more

frequent in the arms than in the legs (Fig 3), for the purple and red tags than for other colours

(Fig 3), and at later checking events (Fig 4). Altogether, when both the colour and the location

of the VIE tags were considered, identification success was 65.4 % overall (Fig 3), decreasing

from 78.8 % to 51 % from the first to the last checking event (Fig 4).

Based on the photographs captured at metamorphosis, individual identification was

highly successful (93.8 %) despite the relatively high uncertainty reported by the observers.

However, 47.3 % of the photos taken at metamorphosis had some problems with image

quality or the body posture of the animal or both, and 42.0 % had poor quality (i.e. the

problems resulted in low visibility of the leg patterns). Ambiguity rate decreased as photo

quality improved from poor (70.2 %) through medium (37.7 %) to good (33.3 %; χ^2 test: $\chi^2 =$

12.17, df = 2, P = 0.002). Identification success was 91.5 % for poor-quality photos, 94.3 %

for medium-quality photos, and 100 % for good-quality photos.

Discussion

In our study, photographs of the melanin markings on the skin were clearly superior to both

toe-clipping and VIE tagging for individual identification of agile frogs during the first one

and a half year of their post-metamorphic life. When image quality was good, the observers

had low ambiguity when assessing potential matches, and the matches they assigned were

always correct when we were able to assess this with a reliable alternative marking (i.e. green

tags that were unambiguously detected in our study). Although the skin patterns sometimes

changed slightly as the animals aged (Appendix S2 Fig S4), the observers were always able to

find the correct match from up to six candidate individuals in our study system. Notably, we

found similarly exquisite performance of photo-based individual identification in a parallel

study on adult agile frogs with large sample sizes (up to several hundreds of individuals),

demonstrating that 100% accuracy can be achieved with computer-assisted photo-

identification when the best techniques are applied for image capturing, paying attention to

animal positioning and lighting conditions [23]. Furthermore, in the present study, the

photographs taken at metamorphosis (i.e. emergence of front limbs) also proved suitable for

individual identification of the young frogs 2-3 months later. When picture quality was good,

photo-based identification was 100% successful regardless of the timing of taking the

reference photographs (i.e. at metamorphosis or at froglet marking), and it was fairly high

even with the poorest-quality pictures taken at metamorphosis (91.5 %). This suggests that the

unique individual patterns of agile frogs are already recognizable at very early life stages and

remain largely stable throughout and after metamorphosis (Appendix S2 Fig S5), similarly to

what has been found in a few other anuran species [31,32]. The drawback of the photo-

identification method is that comparison of the individual to the reference photographs

requires the availability of those photographs (e.g. on a computer at hand) and non-negligible

amount of time that increases with the number of potential matches that need to be checked.

For example, in our study, on-site assessment of VIE tags and toe-clips took qualitatively

similar time to photographing, but the latter had to be followed by off-site image processing

and subsequent photo comparison. Nevertheless, the time needed for finding the right match

can be sped up by computer-assisted photo-identification [33] and optimized photography

methods [20,23]. Earlier studies that used computer-assisted photo-identification in

amphibians had widely variable success rates (Table S1 in [23]), which might be due to

variability in the photography settings and software applied or the stability of skin markings.

Overall, the second-best method in our study was VIE tagging. In our system, tag colour

alone should have been enough for successful individual identification because the same

colour was rarely injected into more than one animal in the same box, and the tags rarely ever

moved from one side of the body to the other. However, we had difficulty with telling apart

three of the six colours (red, pink, and purple), probably because the skin of the tagged body

parts is reddish-brown (Appendix S2 Fig S4 and Fig S5) and sometimes can be quite dark. In

this respect, body parts with lightly coloured skin would be preferable for VIE tagging;

however, the only such area on agile frogs is the belly and the ventral side of the thigh, where

tag retention has been typically low in other species [15,17]. Furthermore, sometimes we did

not even find the tag, despite that we were often able to locate the same tags during some of

the subsequent checking events. The reason for the latter is that the tags often broke into

several pieces, and the pieces migrated away from the injection point both towards and away

from the centre of the body, often leaving only small, easy-to-miss dots close to the skin

surface. These findings are similar to those reported from studies on other amphibian species

[15,16,34]. Tag dislocation may further reduce success in situations where tag location is part

of the individual identification code; for example, we found that tags injected into the shin can

move into the foot and *vice versa*. In our study, dislocation was more frequent for tags

injected into the upper arm compared to the shin and foot; however, this contrasts findings in

other anuran species [17,25,35] and might be explained by tag size rather than by location,

because we injected smaller tags into the upper arms due to their smaller size relative to the

two target sites on the legs. However, larger tags were more likely to migrate in other

amphibians [36]. Altogether, in our study, tag migration resulted in the 77 % identification

success based solely on tag colour dropping to 65 % based on the combination of tag colour

and location. This overall low success rate would be unacceptable for many studies; however,

VIE tags could be still reliable for applications that do not require a large number of

individual codes. For example, our data suggest that young agile frogs could be successfully

batch-marked with green, yellow, or orange VIE for tracking cohorts.

Our data on tag migration raise concerns about the health risks associated with VIE

tagging. Regardless of initial tag location and colour, some of the VIE material moved into the

internal organs, mostly the kidneys, which agrees with findings on adult cane toads (*Rhinella*

marina; [37]). In the latter study, VIE in the kidney was associated with inflammation and

damage to the vasculature that progressed in severity over time. This is especially worrisome

in light of growing concerns about the ecotoxicological effects of microplastics [38]. Notably,

although growth and survival were reported to be unaffected by VIE tagging in various

amphibians, this has been studied over relatively short periods, rarely longer than a year [7–

13]. Therefore, caution is recommended with using VIE especially in free-living wild animals,

and further research is needed to validate its long-term safety.

Finally, toe-clipping yielded unacceptably low identification success in our study. This

may be due to particularly high regeneration capacities in young juveniles, although 100 %

toe-clip retention was observed in boreal chorus frogs (*Psuedacris maculata*) over one year

after metamorphosis [7]. However, the length of the clipped part was not clarified in the latter

study; studies that reported high reliability of toe-clipping typically removed more than one

phalanx per toe (e.g. [3,34]) whereas we removed only the most distal phalanx. Nevertheless,

even if identification success would be increased by more destructive clipping, we do not

recommend this because the negative consequences of toe-clipping on survival increase with

the severity of mutilation (e.g. by the number of toes clipped per animal; [7,39]). For the same

reasons, animal welfare and ethics committees are increasingly unwilling to permit the

clipping of more than one phalanx, to minimise pain and lasting harm. Our results suggest,

however, that this practice may lead to the data turning out useless, and the mutilation having

been done in vain. As pointed out by a recent systematic review on the welfare impacts of toe-

clipping amphibians [2], data to help decisions in this ethically loaded topic are still too scarce

and heterogeneous. While the debate is still going on and more data are being collected, we

recommend using less invasive methods for marking individuals whenever feasible.

In conclusion, our study shows that photo-based individual identification is superior to

both VIE-tagging and toe-clipping for individual identification in young agile frogs. When the

images were taken with care to ensure good quality, photo-identification had 100% success

over one and a half year, and major individually unique patterns were recognizable already at

metamorphosis. Although our animals were kept in small groups, the high stability of skin

patterns means that similarly exquisite performance may be achieved with larger numbers of

individuals. In contrast, both VIE-tagging and toe-clipping showed lower identification

success, and the emergence of VIE in internal organs suggests health risks. We encourage the

usage of the most reliable and least invasive, even though arguably most time-consuming

photo-based method in future studies whenever possible.

Acknowledgements

This research was funded in part by the Austrian Science Fund (FWF) [grant ESP 239-B to
E.N.]. For open access purposes, the author has applied a CC BY public copyright license to
any author-accepted manuscript version arising from this submission. The other part of the
funding came from the National Research, Development and Innovation Office of Hungary
(NKFIH, grant K-135016 to V.B.). The authors were further supported by the New National
Excellence Program (ÚNKP-23-4 to A.K.) and University Excellence Scholarship Program
(EKÖP-24 -DOK-06 to E.B., EKÖP-MATE/2024/25/K to N.U., and EKÖP-24-4-II-ELTE-
364 to Z.M) of the Ministry for Culture and Innovation from the source of the National
Research, Development and Innovation Fund. The funders had no role in study design, data
collection and analysis, decision to publish, or preparation of the manuscript.
We thank Árpád Ferincz for the VIE material, and Emese Balogh, Judit Baumann, Zoltán
Örkényi, Mihály B. Rusz and all colleagues and students at the Department of Evolutionary
Ecology for their help during the study.

Table 1. Timing of procedures and number of individuals (N) throughout the study.

Procedure	Date	N	Photo taken	VIE tag check	Toe-clip check
Metamorphosis	29 May-25 Jun 2023	112*	+	-	-
Marking	5-6 Jul 2023	113	+	-	-
Checking	29-30 Aug 2023	113	+	+	+
	6-7 Nov 2023	110	+	+	+
	13 Mar 2024	109	+	+	-
	15 May 2024	109	+	+	-
	15 July 2024	107	+	+	-
	16 Sep 2024	100	+	+	+
	4 Nov 2024	100	+	+	-

410   * Note that due to an error, photo taken at metamorphosis was missing for one individual,
411   therefore such photos were available for only 112 out of those 113 animals that we marked
412   and monitored in this study.

Table 2. Percentage of ambiguous and unambiguous identifications for each marking method, regardless of correctness of the identification.

Method	N[*]	Unambiguous	Ambiguous Uncertain[†]	Conflicted[‡]	Not found[§]
Toe-clips	**323**	**37.2%**	**22.3%**	**40.2%**	**0.3%**
Front 2	62	46.8%	40.3%	12.9%	0
Front 3	53	20.8%	49.1%	28.3%	1.9%
Front 4	56	35.7%	44.6%	19.6%	0
Hind 3	45	31.1%	42.2%	26.7%	0
Hind 4	54	31.5%	42.6%	25.9%	0
Hind 5	53	54.7%	35.8%	9.4%	0
VIE tags	**748**	**80.3%**	**4.3%**	**5.1%**	**10.3%**
Green	122	97.5%	0	0	2.5%
Yellow	131	90.8%	0.8%	3.1%	5.3%
Orange	140	91.4%	2.1%	2.9%	3.6%
Red	128	60.9%	10.2%	7.8%	21.1%
Pink	117	74.4%	8.5%	13.7%	3.4%
Purple	110	63.6%	4.5%	3.6%	28.2%
Photos compared to					
Froglet marking	**748**	**92.2%**	**7.8%**	**0**	**0**
Metamorphosis	112	49.1%	44.6%	4.5%	1.8%

[*]Sample size (N) of identifications is the number of individuals × number of checking events. Note that for photos, the percentage of conflict was assessed on a subsample of N=113 based on the images taken at marking and N=40 based on the images taken at metamorphosis.

[†]Only one ID was assigned but with uncertainty.

[‡]More than one ID was deemed possible.

[§]No clipped toe or VIE tag or matching photo was found.

Fig 1. Individual identification by toe-clipping and VIE tagging. Each individual was marked by the removal of one toe tip and the injection of one VIE colour into one of the highlighted body parts (A). VIE codes are highlighted with blue frame. Each marking code indicates the side of the body (L: left, R: right). Additionally, toe-clip codes indicate the limb (F: forelimb, H: hind limb) and digit (according to proximity to the body), while VIE codes indicate the body part (A: upper arm, S: shin, F: foot). The applied VIE colours and whether or not they are fluorescent under ultraviolet lighting are indicated on the right. Example photos show LF-4 combined with RS-red (B) and RF-4 combined with LF-green (C) just after tagging. Exposure curves were applied for each image to facilitate pattern visibility

Fig 2. Identification success for marking by toe-clipping, depending on the digit clipped (A) and the timing of checking the markings (B). Letters at the top of bar plots indicate statistically significant differences (groups with different letters differ at P < 0.05 after correcting for false-discovery rate).

Fig 3. Identification success calculated in three different ways for marking by VIE tagging, depending on tag colour (panels A, C and E) and tagged body part (panels B, D and F). Letters at the top of bar plots indicate statistically significant differences (groups with different letters differ at P < 0.01 after correcting for false-discovery rate).

Fig 4. Identification success calculated in three different ways for marking by VIE tagging, depending on the timing of checking the markings. Letters at the top of bar plots indicate statistically significant differences (groups with different letters differ at P < 0.01 after correcting for false-discovery rate).

References

1. Murray DL, Fuller MR. A critical review of the effects of marking on the biology of

vertebrates. In: Boitani L, Fuller TK, editors. Research Techniques in Animal Ecology:

Controversies and Consequences. New York: Columbia University Press; 2000. pp.

15–64.

2. Zemanova MA, Martín RL, Leenaars CHC. The impact of toe-clipping on animal

welfare in amphibians: A systematic review. Glob Ecol Conserv. 2025;59: e03582.

doi:10.1016/j.gecco.2025.e03582

3. Hoffmann K, McGarrity ME, Johnson SA. Technology meets tradition: A combined

VIE-C technique for individually marking anurans. Appl Herpetol. 2008;5: 265–280.

doi:10.1163/157075408785911002

4. Campbell TS, Irvin P, Campbell KR, Hoffmann K, Dykes ME, Harding AJ, et al.

Evaluation of a new technique for marking anurans. Appl Herpetol. 2009;6: 247–256.

doi:10.1163/157075409X420042

5. Davis TM, Ovaska K. Individual recognition of amphibians: effects of toe clipping and

fluorescent tagging on the salamander *Plethodon vehiculum*. J Herp. 2001;35: 217–225.

6. Phillott AD, Skerratt LF, McDonald KR, Lemckert FL, Hines HB, Clarke JM, et al.

Toe-clipping as an acceptable method of identifying individual anurans in mark

recapture studies. Herpetol Rev. 2007;38: 305–308.

7. Swanson JE, Bailey LL, Muths E, Chris Funk W. Factors influencing survival and

mark retention in postmetamorphic boreal chorus frogs. Copeia. 2013; 670–675.

doi:10.1643/CH-12-129

8. Sapsford SJ, Roznik EA, Alford RA, Schwarzkopf L. Visible implant elastomer

marking does not affect short-term movements or survival rates of the treefrog litoria

rheocola. Herpetologica. 2014;70: 23–33. doi:10.1655/HERPETOLOGICA-D-13-0004

9. Fouilloux CA, Garcia-Costoya G, Rojas B. Visible implant elastomer (VIE) success in

early larval stages of a tropical amphibian species. PeerJ. 2020;8: e9630.

doi:10.7717/peerj.9630

10. Knapp DD, Diaz L, Unger S, Anderson CN, Spear SF, Williams LA, et al. Long-term

retention, readability, and health effects of visible implant elastomer (VIE) and visible

implant alpha (VI Alpha) tags in larval eastern hellbenders (*Cryptobranchus*

alleganiensis alleganiensis). J Herpetol. 2023;57: 133–141. doi:10.1670/22-011

11. Bainbridge L, Stockwell M, Valdez J, Klop-toker K, Clulow S, Clulow J, et al.

Tagging tadpoles: retention rates and impacts of visible implant elastomer (VIE) tags

from the larval to adult amphibian stages. Herpetol J. 2015;25: 133–140.

12.   Phillips CT, Fries JN. An Evaluation of Visible Implant Elastomer for Marking the

Federally Listed Fountain Darter and the San Marcos Salamander. North Am J Fish

Manag. 2009;29: 529–532. doi:10.1577/m08-138.1

13.   Oropeza-Sánchez MT, Sandoval-Comte A, García-Bañuelos P, Hernández-López P,

Pineda E. Use of visible implant elastomer and its effect on the survival of an

endangered minute salamander. Anim Biodivers Conserv. 2020;43: 187–190.

doi:10.32800/abc.2020.43.0187

14.   Schmidt K, Schwarzkopf L. Visible implant elastomer tagging and toe-clipping:

Effects of marking on locomotor performance of frogs and skinks. Herpetol J. 2010;20:

99–105.

15.   Brannelly LA, Chatfield MWH, Richards-Zawacki CL. Visual implant elastomer (VIE)

tags are an unreliable method of identification in adult anurans. Herpetol J. 2013;23:

125–129.

16.   Heemeyer JL, Homyack JA, Haas CA. Retention and readability of visible implant

elastomer marks in eastern red-backed salamanders (*Plethodon cinereus*). Herpetol

Rev. 2007;38: 425–428.

17.   Moosman DL, Moosman Jr. PR. Subcutaneous movements of visible implant

elastomers in wood frogs (*Rana sylvatica*). Herpetol Rev. 2006;37: 300–301.

18.   Rojas B, Lawrence JP, Márquez R. Amphibian Coloration: Proximate Mechanisms,

Function, and Evolution. In: Moreno-Rueda G, Comas M, editors. Evolutionary

Ecology of Amphibians. Boca Raton: CRC Press; 2023. pp. 219–258.

doi:10.1201/9781003093312-12

19.   Price AC, Weadick CJ, Shim J, Rodd FH. Pigments, patterns, and fish behavior.

Zebrafish. 2008;5: 297–307. doi:10.1089/zeb.2008.0551

20.   Bendik NF, Morrison TA, Gluesenkamp AG, Sanders MS, O'Donnell LJ. Computer-

assisted photo identification outperforms Visible Implant Elastomers in an endangered

salamander, Eurycea tonkawae. PLoS One. 2013;8: e59424.

doi:10.1371/journal.pone.0059424

21.   Burgstaller S, Gollmann G, Landler L. The green toad example: A comparison of

pattern recognition software. North West J Zool. 2021;17: 96–99.

22.   Matthé M, Sannolo M, Winiarski K, Spitzen - van der Sluijs A, Goedbloed D,

Steinfartz S, et al. Comparison of photo-matching algorithms commonly used for

photographic capture–recapture studies. Ecol Evol. 2017;7: 5861–5872.

doi:10.1002/ece3.3140

23. Nemesházi E, Mikó Z, Ujhegyi N, Kásler A, Lehofer N, Bókony V. Put the frog in

water: simple methods for improving individual identification as demonstrated with the

agile frog. Prepr BioRxiv. 2025. doi:10.1101/2025.02.25.640084

24. Aevarsson U, Graves A, Carter KC, Doherty-Bone TM, Kane D, Servini F, et al.

Individual identification of the lake Oku clawed frog (*Xenopus longipes*) using a

photographic identification technique. Herpetol Conserv Biol. 2022;17: 67–75.

25. Holcomb J, Lawrence K, Bailey S, Dittmer D. Considerations for and use of visible

implant elastomers to track and estimate abundance of juvenile boreal toads (Anaxyrus

boreas boreas) in Utah, USA. Herpetlogical Rev. 2022;53: 579–585.

26. Bókony V, Balogh E, Mikó Z, Kásler A, Örkényi Z, Ujhegyi N. Higher sex-reversal

rate of urban frogs in a common-garden experiment suggests adaptive microevolution.

Evol Appl. 2025;18: e70093. doi:10.1111/eva.70093

27. Bókony V, Üveges B, Móricz ÁM, Hettyey A. Competition induces increased toxin

production in toad larvae without allelopathic effects on heterospecific tadpoles. Funct

Ecol. 2018;32: 667–675. doi:10.1111/1365-2435.12994

28. Kásler A, Holly D, Herczeg D, Ujszegi J, Hettyey A. Chytridiomycosis and climate

change: exposure to *Batrachochytrium dendrobatidis* and mild winter conditions do not

increase mortality in juvenile agile frogs during hibernation. Anim Conserv. 2023;26:

654–662. doi:10.1111/acv.12851

29. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji:

an open-source platform for biological-image analysis. Nat Methods. 2012;9: 676–682.

doi:10.1038/nmeth.2019

30. Pike N. Using false discovery rates for multiple comparisons in ecology and evolution.

Methods Ecol Evol. 2011;2: 278–282. doi:10.1111/j.2041-210X.2010.00061.x

31. Kenyon N, Phillott AD, Ross AA. Temporal variation in dorsal patterns of juvenile

green-eyed tree frogs, *Litoria genimaculata* (Anura: Hylidae). Herpetol Conserv Biol.

2010;5: 126–131.

32. Bardier C, Székely D, Augusto-Alves G, Matínez-Latorraca N, Schmidt BR,

Cruickshank SS. Performance of visual vs. software-assisted photo-identification in

mark-recapture studies: a case study examining different life stages of the Pacific

Horned Frog (*Ceratophrys stolzmanni*). Amphibia-Reptilia. 2020;42: 17–28.

doi:10.1163/15685381-BJA10025

33. Davis HP, Vancompernolle M, Dickens J. Effectiveness and reliability of photographic

identification methods for identifying individuals of a cryptically patterned toad.

Herpetol Conserv Biol. 2020;15: 204–211.

34. Brannelly LA, Berger L, Skerratt LF. Comparison of three widely used marking

techniques for adult anuran species Litoria verreauxii alpina. Herpetol Conserv Biol.

2014;9: 428–435.

35. Lunghi E, Bruni G, Manenti R, Ficetola GF. Use of visible implant elastomer on two

amphibians orders (Anura and Caudata): data on efficiency and reliability. Atti X

Congr Naz Soc Herpetol Ital. 2015;15: 163–167.

36. Grant EHC. Visual implant elastomer mark retention through metamorphosis in

amphibian larvae. J Wildl Manage. 2008;72: 1247–1252. doi:10.2193/2007-183

37. Cabot ML, Troan B V., Heugten KA Van, Schnellbacher RW, Smith D, Ridgley F, et

al. Migration and histologic effects of visible implant elastomer (VIE) and passive

integrated transponder (PIT) tags in the marine toad (*Rhinella marina*). Animals.

2021;11: 3255. doi:10.3390/ani11113255

38. Li C, Busquets R, Campos LC. Assessment of microplastics in freshwater systems: a

review. Sci Total Environ. 2020;707: 135578. doi:10.1016/j.scitotenv.2019.135578

39. Grafe TU, Stewart MM, Lampert KP, Rödel M. Putting toe clipping into perspective: a

viable method for marking anurans. J Herpetol. 2018;45: 28–35.