Iterative *in vivo* cut'n'paste of chromosomal loci in *Escherichia coli* K-12 using

synthetic DNA

Authors: Sebastian Gude[1,*], Andreas B. Bertelsen[1], and Morten H. H. Nørholm[1,2]

[1] The Novo Nordisk Foundation Center for Biosustainability, Technical University of

Denmark, Kongens Lyngby, Denmark

[2] current address: Novo Nordisk A/S, Hørsholm, Denmark

*Corresponding author: Sebastian Gude, The Novo Nordisk Foundation Center for

Biosustainability, Technical University of Denmark, Kongens Lyngby, Denmark,

segu@biosustain.dtu.dk

Research area:

Biological sciences, microbiology

Keywords:

easy-to-use, *in vivo*, genome engineering, genome refactoring, CRISPR Cas9, λ Red

recombineering, homologous recombination, bacteria, *Escherichia coli*

Preprint:

bioRxiv: https://doi.org/10.1101/2025.06.19.660532

23   CC-BY-NC-ND 4.0 International license

<u>Abstract</u>

We have developed an easy-to-use *in vivo* method to iteratively relocate

functional chromosomal loci onto an episome in *Escherichia coli* by utilizing synthetic

DNA fragments. In this *in vivo* cut'n'paste procedure, "cutting" is executed by the RNA-

guided DNA endonuclease Cas9 and a set of guides, while "pasting" is facilitated by

the phage λ Red recombinase, which are all synthesized on easily curable helper

plasmids. To demonstrate the utility of *in vivo* cut'n'paste, we commercially obtained

synthetic DNA fragments containing locus-specific homology regions, antibiotic

marker cassettes, and standardized Cas9 target sequences, and successfully

relocated seven functional chromosomal loci. Scarless relocation mutants of the *trg*,

aer, *tsr*, *malHM*, *malQT*, *macB-nadA*, and *folA* chromosomal loci were obtained as

antibiotic-resistant isolates by combining Cas9 counterselection with the restoration of

an antibiotic marker cassette. The additional antibiotic marker cassettes and

standardized Cas9 target sequences present in the synthetic DNA fragments are

inherently eliminated upon completion of the procedure, enabling iterative processing

of chromosomal loci. *In vivo* cut'n'paste should be widely useful, especially in genome

(re-)engineering efforts of *E. coli* and other bacteria because the procedure can be

performed by non-specialists in unmodified wild-type cells.

<u>Significance</u>

Editing of genomes is crucial to understand the inner workings of bacteria and

to alter them for biotechnological applications. *In vivo* cut'n'paste, in analogy to the

frequently used keyboard shortcuts Ctrl-X and Ctrl-V which allow to move text, images,

or files on computers, makes it easy to relocate large chunks of the bacterial genome

without requiring skills to manipulate DNA outside of the bacterial cell. Like Ctrl-X and

Ctrl-V, *in vivo* cut'n'paste can be executed several times to iteratively edit multiple

genomic locations in the same bacterium. Since *in vivo* cut'n'paste entirely takes place

inside the bacterial cell, it is particularly well suited to edit hard-to-tackle genomic

features such as essential genes.

Introduction

Democratising access to methodological advancements, i.e., making them

accessible to a broad audience of non-specialists, is a key component of method

development. The introduction of the point-and-click graphical user interface by Apple

and Microsoft[1], the creation of the "batteries included" programming language

Python[2], and the development of the one-step inactivation of chromosomal genes

procedure[3] are a few prominent examples of such endeavours. Graphical user

interfaces remove the hurdle to know how to interact with a computer via a text

console. The extensive Python Standard Library supports nearly every aspect of

computer science, thus, freeing programmers from having to reimplement many

commonly required features over and over again. Similarly, the one-step inactivation

of chromosomal genes procedure enables researchers to generate gene knockouts

without the need to assemble integrative plasmids.

The ability to manipulate genomes is key for uncovering and leveraging the

potential of (micro-)biology. A milestone in our ability to edit bacterial genomes was the

development of the one-step inactivation of chromosomal genes procedure[3]. Among

many other scientific and engineering endeavours, it facilitated the creation of a

comprehensive single gene knockout library and the charting of essential genes in *E.*

coli[4], the study of bacterial motility[5,6], stringent response[7], the interplay of growth and

gene expression[8], two-component systems[9], evolution of antibiotic resistsance[10], and

chromosomal organisation[11] as well as the development of strains with reduced

genomes[12] and highly titratable induction systems[13] while also playing a crucial role in

metabolic engineering applications[14,15]. Though the one-step inactivation procedure[3]

was preceded by numerous other bacterial genome editing techniques[16–23], its ease

of use and broad applicability ensured its still continuing success. The required short

homology extensions to create genomic deletions are easily attached to an antibiotic

marker cassette via PCR, the recombineering machinery is expressed from an easily

curable plasmid, and the procedure does not require any prior modifications to the

bacterial genome[3]. Furthermore, the antibiotic marker cassette can subsequently be

eliminated via FLP-FRT recombination, resulting in an unmarked, yet not fully scarless,

mutant that is compatible with another round of genome editing[3].

With the rise of CRISPR-Cas technologies[24], λ Red recombineering was

successfully supplemented with sequence-specific RNA-guided counterselection[25–27].

Combining λ Red recombineering and CRISPR-Cas with intricate multi-layered

selection, counterselection, and colour screening enables large-scale genome editing

such as genome recoding[28,29], global genome rearrangement[30,31], and interspecies

gene transfer[32]. While these sophisticated large-scale CRISPR-Cas genome editing

techniques accomplish feats vastly beyond the capabilities of the one-step inactivation

of chromosomal genes procedure[3], they unfortunately fall short in one aspect. Namely,

they are relatively difficult to implement in non-specialist laboratories due to their

technical complexity, thus impeding the democratisation of large-scale genome

editing.

Building upon established large-scale CRISPR-Cas genome editing methods[28–

32], we present here *in vivo* cut'n'paste - an attempt to democratise large-scale genome

editing. *In vivo* cut'n'paste is an easy-to-use procedure to relocate chromosomal loci

onto an episome in wild-type cells that was designed to be performed by non-

specialists without expert skills in molecular biology. *In vivo* cut'n'paste solely relies on

commercially synthesized DNA fragments and a set of easily curable helper plasmids.

Leveraging CRISPR-Cas cutting and λ Red recombineering with commercially

synthesized DNA fragments completely removes the need for *in vitro* DNA

manipulations. Furthermore, easily curable helper plasmids and an antibiotic marker

recycling scheme inherently enable the iterative creation of multiple large-scale

genome edits. We successfully relocated seven functional genomic loci, including

metabolic and essential genes, in *E. coli* of up to 140 kb in size.

Results

In vivo cut'n'paste

In vivo cut'n'paste enables to iteratively relocate genomic loci from the native

chromosome onto an episome (Fig. 1A, Supp. Movie 1). In each round, a genomic

locus, defined by two pairs of tandem homology regions HR^A-HR^B and HR^C-HR^D is

excised from the native chromosome and subsequently integrated in between the

homology regions HR^E and HR^{TET} on an episome (Fig. 1B). The entire *in vivo*

122 cut'n'paste procedure occurs inside the cell without requiring any *in vitro* DNA

123 manipulations.

124   Each round of *in vivo* cut'n'paste consists of two phases: (1) pre-loading and

125 (2) relocation (Fig. 1B). During pre-loading, two commercially synthesized DNA

126 fragments, PRE^{epi} and PRE^{chr}, are sequentially integrated via homologous

127 recombination into the episome and the native chromosome, respectively, to equip

128 cells with the required modification for the subsequent relocation of the genomic locus

129 (Fig. 1C; see Supplementary Materials and Supp. Fig. 1-3 for a detailed description of

130 pre-loading fragment design choices). Both pre-loading events are accompanied by

131 positive antibiotic selection.

132   PRE^{epi} is flanked by the homology regions HR^E and HR^{TET}. Directly downstream

133 of HR^E, PRE^{epi} contains the inside homology region HR^B of the genomic locus. The

134 two outside homology regions HR^A and HR^D of the genomic locus are located next to

135 HR^B followed by a chloramphenicol marker cassette (Fig. 1C). Integration of PRE^{epi} at

136 HR^E and HR^{TET} into the episome truncates the tetracycline maker of the episome and,

137 thus, renders the cells sensitive to tetracycline yet resistant to chloramphenicol (Fig.

138 1B).

139   PRE^{chr} is flanked by the homology region HR^C and HR^D of the genomic locus,

140 enabling its integration directly adjacent to the genomic locus of interest in the native

141 chromosome (Fig. 1B). In PRE^{chr}, HR^C is directly followed by $5'TET^R$, encoding a short,

142 non-functional fragment of the tetracycline marker cassette, and HR^{TET}. A kanamycin

143 marker cassette completes PRE^{chr} (Fig. 1C). Notably, the pre-loading fragments can

144 be commercially obtained in their entirety without requiring further assembly.

145   Once pre-loading is completed, cells contain all features required for relocation

146 (Fig. 1B,C). Firstly, the genomic locus in the native chromosome now contains HR^B on

one side and HR^{TET} on its other side. Secondly, the episome also contains both HR^B

and HR^{TET}. Thirdly, a "repair band-aid", consisting of HR^A-HR^D, which can fix the

double-stranded break that will be created by the excision of the genomic locus from

the native chromosome, is present on the episome (see Supp. Movie 1 for a detailed,

step-by-step illustration of the pre-loading process). Optionally, additional features

may be introduced during pre-loading. For example, a terminator can be inserted in

between HR^C and 5'TET^R in EPI^{chr}, which will, once relocation is completed, partially

insulate the relocated locus from its new local environment on the episome.

Relocation is initiated by transforming the helper plasmid pCut (Fig. 1B). pCut

constitutively expresses standardized, functionally tested guides targeting constant

cut sites (CS^1, CS^2, and CS^3) included in the pre-loading fragments from a CRISPR

array, as well as a single guide (CS^L) targeting a locus-specific junction between HR^A

and HR^B (Fig. 1B,C). The sequence encoding the locus-specific single guide can

effectively be inserted as a third synthetic DNA fragment into the helper plasmid pCut

via I-CreI cutting counterselection-aided plasmid recombineering, thus completely

eliminating the need for *in vitro* DNA manipulations (see Supp. Fig. 4 for details

regarding I-CreI cutting counterselection-aided plasmid recombineering; a technical

description of the procedure is provided in the Methods). In total, RNA-guided Cas9

endonuclease activity now introduces double-stranded breaks into six targeted

locations across the episome and the native chromosome (Fig. 1B,C). These double-

stranded breaks release the genomic locus from the native chromosome and the

"repair band-aid" from the episome while simultaneously exposing the required

homology regions for integration of the genomic locus into the episome and for

integration of the "repair band-aid" into the native chromosome. Importantly,

integration of the genomic locus into the episome not only fixes its double-stranded

break but, additionally, leads to the restoration of the tetracycline marker cassette,

enabling positive selection of successful relocation events in the presence of

tetracycline (see Supp. Movie 1 for a detailed step-by-step illustration of the relocation

process).

The chloramphenicol and kanamycin marker cassettes, along with the constant

cut sites, which were all introduced during pre-loading as components of PRE^{epi} and

PRE^{chr}, are inherently lost during the relocation process. Finally, the helper plasmid

pCut encodes rhamnose-inducible, self-targeting guides enabling its straightforward

elimination from cells (see Methods for details), thus immediately enabling relocation

mutants to undergo additional rounds of *in vivo* cut'n'paste (Fig. 1B). We typically

tested four isolates by diagnostic PCR at each editing step (episome pre-loading,

native chromosome pre-loading, and relocation) of the *in vivo* cut'n'paste procedure.

An individual isolate possessing the desired genomic alternation was chosen for

further processing.

<u>Relocating a genomic locus</u>

In a first attempt, we employed *in vivo* cut'n'paste to relocate the *trg* locus from

its native genomic location in strain CNP^{empty} (*E. coli* K-12 MG1655 WT + EPI^{empty})

onto an empty episome (EPI^{empty}). Candidates of successful relocations, CNP^{trg} (*E.*

coli K-12 MG1655 Δ*trg* + EPI^{trg}), were obtained under tetracycline selection. After

curing the helper plasmids, we sent a CNP^{trg} isolate for short-read whole genome re-

sequencing (Supp. Table 1). Mapping reads of the CNP^{trg} isolate to the parental

CNP^{empty} reference depicted a depth of coverage of about 400x across the native

chromosome and the episome (Fig. 2A, middle row). No mutations relative to the

parental strain CNP[empty] were observed. Yet, notably, pairs of adjacent bases at three

genomic locations were not spanned by any reads. Namely, both edges of the *trg* locus

on the native chromosome (HR[A]-HR[B] and HR[C]-HR[D]) and one edge of HR[E] on the

episome lacked spanning reads (Fig. 2A, middle row). No lack of spanning reads at

these locations was observed when reads of the parental CNP[empty] strain were

mapped to the CNP[empty] reference (Fig. 2A, top row). The lack of spanning reads in

the CNP[trg] isolate suggested that though the *trg* locus was present, it was not located

in its native genomic location, and, additionally, that something had likely been inserted

into the episome next to HR[E]. We updated the reference map to include the desired

relocation of the *trg* locus based on our *in silico* design of the *in vivo* cut'n'paste

procedure. When the reads of the CNP[trg] isolate were mapped to this updated

reference, the lack of spanning reads disappeared (Fig. 2A, bottom row), suggesting

that the *trg* locus had indeed been relocated from the native chromosome onto the

episome. Long-read sequencing of an extracted episome further supported the

successful relocation of the *trg* locus (Supp. Table 2). In total, we observed eleven

reads that spanned from the backbone of the episome across the entire, mutation-free

trg locus.

The *trg* locus ($\approx$1.8 kb) contains a single gene encoding the transmembrane

Trg chemoreceptor. Trg mediates directed motile behaviour (i.e., chemotaxis) toward

D-ribose[33]. We assayed chemotactic colony expansion dynamics in glycerol minimal

medium soft agar plates in the presence and absence of low supplementations of D-

ribose as proxy for ribose sensing (see Methods for details). Initially, a small drop of

bacterial culture was spotted into the centre of the soft agar plates. Upon growing,

bacteria locally deplete nutrients and thus create gradients enabling chemotactic

sensing and expansion[34]. Expansion dynamics of *trg*-relocated CNP[trg] isolates (blue)

in glycerol minimal medium soft agar plates in the presence (sold lines) and absence

(dashed lines) of D-ribose were comparable to the behaviour of their ancestor

CNP^empty (black), clearly indicating that *trg* retained its functionality after relocation

onto the episome (Fig. 2B,C). When the episome EPI^trg was removed from the CNP^trg

isolates (red), the expansion dynamics in presence of D-ribose were reduced to a

slower aerotactic behaviour[35] as also observed in glycerol minimal medium in the

absence of D-ribose, indicating that the ability to sense D-ribose and utilizing its

concentration gradients for chemotactic expansion was lost along with the episome

(Fig. 2B,C). As an additional test, we separately transferred the episomes EPI^trg and

EPI^empty into a strain background in which the *trg* gene had been disrupted by a

kanamycin marker cassette (*E. coli* K-12 MG1655 Δtrg::KanR). As expected, faster

chemotactic expansion dynamics in presence of D-ribose were restored only when

EPI^trg, containing the relocated *trg* locus, was present (Fig. 2D).

Iteratively relocating genomic loci

In accordance with the design principles of the *in vivo* cut'n'paste procedure,

the additional antibiotic marker cassettes and constant cut sites acquired during pre-

loading were inherently lost, and the tetracycline marker cassette on the episome was

restored, upon completion of the *trg* relocation (Fig. 1B). Thus, CNP^trg was capable of

undergoing further rounds of *in vivo* cut'n'paste. We iteratively applied *in vivo*

cut'n'paste to relocate two additional genomic loci onto the episome using CNP^trg as

a starting point (Fig. 3A). First, the *aer* locus ($\approx$1.7 kb) encoding the aerotaxis receptor

Aer was relocated, resulting in CNP^trg,aer (*E. coli* K-12 MG1655 Δtrg Δaer + EPI^trg,aer).

Then, the *tsr* locus ($\approx$1.8 kb) encoding the chemoreceptor Tsr, mediating aerotaxis and

sensing of L-Serine, was relocated, resulting in CNP^{trg,aer,tsr} (*E. coli* K-12 MG1655 Δ*trg*

Δ*aer* Δ*tsr* + EPI^{trg,aer,tsr}).

The relocated loci were mutation-free (Supp. Table 2). The triple-relocated

CNP^{trg,aer,tsr} qualitatively retained the colony expansion behaviour of its ancestor

CNP^{empty} in presence of D-ribose, L-Serine, or in the absence of supplementations to

glycerol minimal medium (Fig. 3 B,C). Furthermore, separately transferring the

episomes EPI^{empty} (Fig. 3D), EPI^{trg} (Fig. 3 E), and EPI^{trg,aer} (Fig. 3F) into a Δ*trg* Δ*aer*

Δ*tsr* chromosomal background (i.e., a CNP^{trg,aer,tsr} derivative from which EPI^{trg,aer,tsr} had

been eliminated) partially restored the colony expansion behaviour in presence of D-

ribose (sensed via Trg), L-Serine (sensed via Tsr), and in absence of any

supplementations (energytaxis via Aer and Tsr) as expected[33,35].

In an independent attempt, using CNP^{empty} as a starting point, we successfully

iteratively relocated two genomic loci (≈8.1 kb and ≈7.9 kb) encoding 10 genes

involved in maltose utilisation onto an episome (Fig. 4A). Again, the relocated loci were

mutation-free. The resulting relocation mutants CNP^{malHM} (*E. coli* K-12 MG1655

Δ(*malH-malM*) + EPI^{malHM}) and CNP^{malHM,malQT} (*E. coli* K-12 MG1655 Δ(*malH-malM*)

Δ(*malQ-malT*) + EPI^{malHM,malQT}) retained their ability to uptake and ferment maltose as

indicated by producing a dark red phenotype when spotted and grown on maltose-

MacConkey/agar plates (Fig. 4B).

<u>Relocating a large genomic locus and an essential gene</u>

To illustrate the utility of *in vivo* cut'n'paste to process large genomic loci, we

attempted to relocate the *macB-nadA* locus (≈140 kb) containing 135 non-essential

genes from its native genomic location onto an episome. The relocation was

successful and mutation-free (Supp. Table 1).

As opposed to traditional sequential deletion and complementation procedures,

all genes, including those located inside the genomic locus that is edited, remain

present throughout the entire *in vivo* cut'n'paste process. Hence, *in vivo* cut'n'paste

should be well suited for manipulating hard-to-edit genomic features such as essential

genes. Indeed, we successfully relocated the essential gene *folA* onto an episome,

resulting in CNP^{folA} (*E. coli* K-12 MG1655 Δ*folA* + EPI^{folA}).

An episome containing an essential gene that is simultaneously absent from

the native chromosome is expected to be stably maintained even in the absence of

antibiotic selection because cells are unable to survive without the essential gene. As

a direct test, we removed the tetracycline antibiotic marker cassette from EPI^{folA} and

inserted a highly expressed green fluorescent reporter (GFP) in its place, resulting in

CNP$^{folA,TetR::gfp}$ (*E. coli* K-12 MG1655 Δ*folA* + EPI$^{folA,TetR::gfp}$). We then assayed the

retention of the antibiotic marker-free episome EPI$^{folA,TetR::gfp}$ by serial culturing of

CNP$^{folA,TetR::gfp}$ in LB medium without any antibiotic supplementation (see Methods for

details). Plating on non-selective LB agar plates was performed at each transfer. After

overnight incubation, plates were imaged in GFP and brightfield channels (Fig. 5,

Supp. Fig. 5). At each transfer, all fluorescent colonies visible in the GFP channel (Fig.

5A) had a corresponding colony in the associated brightfield image (Fig. 5B) and vice

versa. Thus, 100% of the bacterial population stably retained the antibiotic marker-free

episome EPI$^{folA,TetR::gfp}$ throughout ten serial passages in the absence of antibiotic

selection.

<u>Unintended side effects</u>

All relocated genomic loci were mutation-free. Yet, some of the relocation

mutants acquired background mutations in their native chromosome while undergoing

in vivo cut'n'paste (Supp. Table 7). In general, it remains unclear whether these

background mutations have any functional relevance or are merely coincidental

genetic drift.

<u>Discussion</u>

We developed *in vivo* cut'n'paste - an easy-to-use procedure to relocate

chromosomal loci onto an episome in *E. coli* that does not rely on any *in vitro* DNA

manipulations. We successfully relocated genomic loci encoding metabolic and

essential genes, generated a genomic alteration >100 kb, and performed up to three

consecutive edits in a single strain. All relocated loci, including the 140 kb *macB-nadA*

relocation, were mutation-free. Due to its inherent robustness, as each editing step is

accompanied by positive antibiotic selection, *in vivo* cut'n'paste is well-suited for

implementation in non-specialist laboratories or via unsupervised lab automation.

In vivo cut'n'paste can be leveraged to precisely characterise gene function.

One round of *in vivo* cut'n'paste effectively creates a complementation mutant in which

the gene of interest is relocated onto a single-copy extra-chromosomal replicon. Thus,

in vivo cut'n'paste inherently overcomes the limitations of traditional sequential "delete

and complement" procedures hampering the investigation of essential genes while

also circumventing any convoluting copy number effects present when using

traditional multi-copy plasmid backbones.

By iterating the *in vivo* cut'n'paste procedure, arbitrary modules of genes, like

those obtained here (Fig. 3 & 4), can be created. For example, genes that are co-

modulated or functionally connected but natively dispersed across the chromosome

can easily be clustered onto a single extra-chromosomal replicon[36]. Utilising the origin

of transfer (oriT) encoded on the episome, such modules can be transferred to other

strains or species via conjugation without the size-limitations imposed by

transformation[37], and, hence, potential future adaptations of *in vivo* cut'n'paste to other

bacterial species will only broaden its utility (see Supplementary Text).

Its ability to relocate genomic loci makes *in vivo* cut'n'paste particularly well-

suited for studying how genome architecture impacts cell functioning, a topic of which

we are only beginning to scratch the surface[38,39]. Ultimately, *in vivo* cut'n'paste can

facilitate alternative avenues to obtain minimal genomes via bottom-up approaches to

complement traditional, ineffectual top-down genome reduction attempts.

Moreover, relocations of essential genes, such as demonstrated here in

CNP[folA], are highly desirable in the context of large-scale fermentation applications in

green biotechnology as they allow to stably maintain transferable extra-chromosomal

replicons in production strains without incurring the metabolic costs and legal

requirements associated with continual antibiotic selection[40].

Akin to the graphical user interface in personal computing[1], *in vivo* cut'n'paste

does not enable any feature that is not already present in existing techniques. We also

do not claim that *in vivo* cut'n'paste is faster or more efficient compared to other large-

scale genome editing procedures. Instead, the main contribution of *in vivo* cut'n'paste

lies in its ease of use, as it does not require any DNA manipulations outside of the

bacterial cell, not even for the engineering of helper plasmids. While specialists in

molecular biology may view multi-fragment DNA assemblies and PCRs of large, and

potentially complex, DNA pieces as trivial undertakings, such tasks can appear as

insurmountable hurdles to researchers utilising bacteria in fields such as biophysics,

protein structure characterisation, evolutionary biology, or large-scale fermentation. *In*

vivo cut'n'paste is an attempt to democratise large-scale genome editing and, thus, to

open up the world of large-scale genome editing to non-specialists in molecular

biology. In essence, *in vivo* cut'n'paste solely requires (1) to decide which genomic

locus to relocate while identifying the homology regions HR^A-HR^D (see Fig. 1B), (2) to

complete the annotated pre-loading fragment genbank file templates (supplied in the

Supplementary Materials), (3) to commercially obtain standardised synthetic DNA

fragments, (4) to transform these synthetic DNA fragments and the easily curable

helper plasmids into the strain of interest while culturing in appropriate selective

conditions, (5) to cure helper plasmids, and, finally, (6) to send the mutants for whole

genome re-sequencing. Thus, *in vivo* cut'n'paste empowers experimenters to refactor

genomes, akin to the seven genomic edits demonstrated here, without requiring

specialized knowledge in molecular biology methodologies.

Materials and Methods

See "Methods" document.

Data availability

All data is publicly available (see Supp. Table 8 for details). All custom-written

scripts are supplied in the Supplementary Materials. Sequencing data was deposited

at NCBI (PRJNA1271273; see Supp. Table 1, 2, and 3 for assession numbers).

Sequence-verified, annotated maps of episomes and helper plasmids are supplied in

the Supplementary Materials. All *in vivo* cut'n'paste homology regions are listed in

Supp. Table 6. Strains are available for non-commercial use through the Belgian

Coordinated Collections of Microorganisms (https://bccm.belspo.be; see Supp. Table

1 for assession numbers). Helper plasmids are available for non-commercial use

through Addgene (https://addgene.org; see Supp. Table 3 for assession numbers).

Authors' contributions

SG, ABB, and MHHN developed the initial concept of *in vivo* cut'n'paste. SG

implemented, tested, and refined *in vivo* cut'n'paste iteratively. SG performed all

experiments, analysed the data, and wrote the manuscript. All authors read the final

version of the manuscript.

Conflicts of interest

The authors do not have any conflicts of interest to disclose.

Acknowledgement and funding information

The authors would like to acknowledge Valeria Stavila (The Novo Nordisk

Foundation Center for Biosustainability, Technical University of Denmark, Kongens

Lyngby, Denmark) for technical assistance. S.G. would like to acknowledge Alex T.

Nielsen (The Novo Nordisk Foundation Center for Biosustainability, Technical

University of Denmark, Kongens Lyngby, Denmark) for hosting the final period of

research activity in his laboratory. *E. coli* K-12 MG1655 WT was a kind gift of Tom Shimizu (AMOLF, Amsterdam, The Netherlands). The title and the abstract of this manuscript pay homage to the article "One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products". This work was supported by The Novo Nordisk Foundation.

References

1. Abbate, J. Getting small: a short history of the personal computer. *Proc. IEEE* **87**, 1695–1698 (1999).

2. Dubois, P. F. Guest Editor's Introduction: Python: Batteries Included. *Comput. Sci. Eng.* **9**, 7–9 (2007).

3. Datsenko, K. A. & Wanner, B. L. One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc. Natl. Acad. Sci.* **97**, 6640–6645 (2000).

4. Baba, T. *et al.* Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the Keio collection. *Mol. Syst. Biol.* **2**, 2006.0008 (2006).

5. Chapman, M. R. *et al.* Role of *Escherichia coli* Curli Operons in Directing Amyloid Fiber Formation. *Science* **295**, 851–855 (2002).

6. Smith, K. D. *et al.* Toll-like receptor 5 recognizes a conserved site on flagellin required for protofilament formation and bacterial motility. *Nat. Immunol.* **4**, 1247–1253 (2003).

7. Traxler, M. F. *et al.* The global, ppGpp-mediated stringent response to amino acid starvation in *Escherichia coli*. *Mol. Microbiol.* **68**, 1128–1148 (2008).

8. Klumpp, S., Zhang, Z. & Hwa, T. Growth Rate-Dependent Global Effects on Gene

Expression in Bacteria. *Cell* **139**, 1366–1375 (2009).

9. Zhou, L., Lei, X.-H., Bochner, B. R. & Wanner, B. L. Phenotype MicroArray Analysis of

Escherichia coli K-12 Mutants with Deletions of All Two-Component Systems. *J.*

Bacteriol. **185**, 4956–4972 (2003).

10.   Liu, J., Gefen, O., Ronin, I., Bar-Meir, M. & Balaban, N. Q. Effect of tolerance on

the evolution of antibiotic resistance under drug combinations. *Science* **367**, 200–

204 (2020).

11.   Lau, I. F. *et al.* Spatial and temporal organization of replicating *Escherichia coli*

chromosomes. *Mol. Microbiol*. **49**, 731–743 (2003).

12.   Kolisnychenko, V. *et al*. Engineering a Reduced *Escherichia coli* Genome.

Genome Res. **12**, 640–647 (2002).

13.   Meyer, A. J., Segall-Shapiro, T. H., Glassey, E., Zhang, J. & Voigt, C. A. Escherichia

coli "Marionette" strains with 12 highly optimized small-molecule sensors. *Nat.*

Chem. Biol. **15**, 196–204 (2019).

14.   Lee, K. H., Park, J. H., Kim, T. Y., Kim, H. U. & Lee, S. Y. Systems metabolic

engineering of *Escherichia coli* for L -threonine production. *Mol. Syst. Biol*. **3**, 149

(2007).

15.   Atsumi, S. *et al*. Metabolic engineering of Escherichia coli for 1-butanol

production. *Metab. Eng*. **10**, 305–311 (2008).

16.   Winans, S. C., Elledge, S. J., Krueger, J. H. & Walker, G. C. Site-directed insertion

and deletion mutagenesis with cloned fragments in Escherichia coli. *J. Bacteriol*. **161**,

1219–1221 (1985).

17.   Hamilton, C. M., Aldea, M., Washburn, B. K., Babitzke, P. & Kushner, S. R. New

444   method for generating deletions and gene replacements in Escherichia coli. *J.*

445   *Bacteriol.* **171**, 4617–4622 (1989).

446   18.   Russell, C. B., Thaler, D. S. & Dahlquist, F. W. Chromosomal Transformation of

447   Escherichia coli recD Strains with Linearized Plasmidst. *J BACTERIOL* **171**, (1989).

448   19.   Metcalf, W. W. *et al.* Conditionally Replicative and Conjugative Plasmids Carrying

449   lacZa for Cloning, Mutagenesis, and Allele Replacement in Bacteria. *Plasmid* **35**, 1–13

450   (1996).

451   20.   Link, A. J., Phillips, D. & Church, G. M. Methods for generating precise deletions

452   and insertions in the genome of wild-type Escherichia coli: application to open

453   reading frame characterization. *J. Bacteriol.* **179**, 6228–6237 (1997).

454   21.   Kato, C., Ohmiya, R. & Mizuno, T. A Rapid Method for Disrupting Genes in the

455   Escherichia coli Genome. *Biosci. Biotechnol. Biochem.* **62**, 1826–1829 (1998).

456   22.   Zhang, Y., Buchholz, F., Muyrers, J. P. P. & Stewart, A. F. A new logic for DNA

457   engineering using recombination in Escherichia coli. *Nat. Genet.* **20**, 123–128 (1998).

458   23.   Posfai, G., Kolisnychenko, V., Bereczki, Z. & Blattner, F. R. Markerless gene

459   replacement in Escherichia coli stimulated by a double-strand break in the

460   chromosome. *Nucleic Acids Res.* **27**, 4409–4415 (1999).

461   24.   Jinek, M. *et al.* A Programmable Dual-RNA–Guided DNA Endonuclease in

462   Adaptive Bacterial Immunity. *Science* **337**, 816–821 (2012).

463   25.   Jiang, W., Bikard, D., Cox, D., Zhang, F. & Marraffini, L. A. RNA-guided editing of

464   bacterial genomes using CRISPR-Cas systems. *Nat. Biotechnol.* **31**, 233–239 (2013).

465   26.   Pyne, M. E., Moo-Young, M., Chung, D. A. & Chou, C. P. Coupling the

466   CRISPR/Cas9 System with Lambda Red Recombineering Enables Simplified

Chromosomal Gene Replacement in Escherichia coli. *Appl. Environ. Microbiol.* **81**,

5103–5114 (2015).

27.   Jiang, Y. *et al.* Multigene Editing in the Escherichia coli Genome via the CRISPR-

Cas9 System. *Appl. Environ. Microbiol.* **81**, 2506–2514 (2015).

28.   Wang, K. *et al.* Defining synonymous codon compression schemes by genome

recoding. *Nature* **539**, 59–64 (2016).

29.   Zürcher, J. F. *et al.* Continuous synthesis of E. coli genome sections and Mb-

scale human DNA assembly. *Nature* **619**, 555–562 (2023).

30.   Wang, K., De La Torre, D., Robertson, W. E. & Chin, J. W. Programmed

chromosome fission and fusion enable precise large-scale genome rearrangement

and assembly. *Science* **365**, 922–926 (2019).

31.   Su, J. *et al.* A CRISPR-based chromosomal-separation technique for Escherichia

coli. *Microb. Cell Factories* **21**, 235 (2022).

32.   Teufel, M., Klein, C. A., Mager, M. & Sobetzko, P. A multifunctional system for

genome editing and large-scale interspecies gene transfer. *Nat. Commun.* **13**, 3430

(2022).

33.   Kondoh, H., Ball, C. B. & Adler, J. Identification of a methyl-accepting chemotaxis

protein for the ribose and galactose chemoreceptors of Escherichia coli. *Proc. Natl.*

Acad. Sci. **76**, 260–264 (1979).

34.   Adler, J. Chemotaxis in Bacteria. *Science* **153**, 708–716 (1966).

35.   Zhulin, I. B., Rowsell, E. H., Johnson, M. S. & Taylor, B. L. Glycerol elicits energy

taxis of Escherichia coli and Salmonella typhimurium. *J. Bacteriol.* **179**, 3196–3201

(1997).

36.   Tan, J. *et al.* Independent component analysis of E. coli's transcriptome reveals

the cellular processes that respond to heterologous gene expression. *Metab. Eng.* **61**,

360–368 (2020).

37.   Hanahan, D. Studies on transformation of Escherichia coli with plasmids. *J. Mol.*

Biol. **166**, 557–580 (1983).

38.   El Houdaigui, B. *et al.* Bacterial genome architecture shapes global

transcriptional regulation by DNA supercoiling. *Nucleic Acids Res.* **47**, 5648–5657

(2019).

39.   Dame, R. T., Rashid, F.-Z. M. & Grainger, D. C. Chromosome organization in

bacteria: mechanistic insights into genome structure and function. *Nat. Rev. Genet.*

21, 227–242 (2020).

40.   Sevillano, L., Díaz, M. & Santamaría, R. I. Development of an antibiotic marker-

free platform for heterologous protein production in Streptomyces. *Microb. Cell*

Factories **16**, 164 (2017).

<u>Figures and figure legends</u>

See "Figures" document.

<u>Supplementary Materials</u>

See "SupplementaryMaterials" file.

Iterative *in vivo* cut'n'paste of chromosomal loci in *Escherichia coli* K-12 using

synthetic DNA

<u>Figures and figure legends</u>

Figure 1: Iterative *in vivo* cut'n'paste

a) Graphical overview of *in vivo* cut'n'paste. **b)** Outline of the individual steps (pre-loading and relocation) of the *in vivo* cut'n'paste procedure. **c)** Features of commercially synthesized DNA fragments PRE^{epi} and PRE^{chr}. A more detailed description of the design choices is provided in the Supplementary Materials. Half-arrows indicate the relative orientations of features. Solid, downwards arrows mark constant cut sites. pRedCas9 and pCut: easily curable helper plasmids, HR: homology region, CS: cut site, TET^R: tetracycline marker cassette, CM^R: chloramphenicol marker cassette, KAN^R: kanamycin marker cassette, 5'TET^R: non-functional fragment of TET^R, Tet: tetracycline, Cm: chloramphenicol, Kan: kanamycin, R: resistant, S: susceptible, and bp: base pairs. Not to scale.

Figure 2: *In vivo* cut'n'paste of the *trg* locus

a) Number of mapped reads in 3.2 kb segments of the native chromosome and the

episome. Only reads mapping to a position and to both of its direct neighbors are

counted. Strain identities and reference maps as indicated within the panel. The *trg*

locus is highlighted in yellow. Triangles mark locations of homology regions. **b-d)**

Motility phenotypes in soft agar. Colony expansion as function of time in glycerol

minimal medium with (solid) or without (dashed) 100 μM D-ribose. **b-c)** CNP^{trg} (blue),

CNP^{trg} without EPI^{trg} (red), and CNP^{empty} (black). Three independent CNP^{trg} isolates

and a single replicate of CNP^{empty}. **d)** *E. coli* K-12 MG1655 Δtrg::KanR with EPI^{trg}

(blue), *E. coli* K-12 MG1655 Δtrg::KanR with EPI^{empty} (red), and CNP^{empty} (black). A

single replicate each.

Figure 3: Iterative *in vivo* cut'n'paste of three chemoreceptor loci

a) Illustration of iterative *in vivo* cut'n'paste of the *trg*, *aer*, and *tsr* chemoreceptor loci.

b-f) Motility phenotypes in soft agar. Colony expansion as function of time in glycerol

minimal medium with 100 μM D-ribose (blue), with 100 μM L-Serine (red), or without

any additions (black). Strain identities as defined within the individual panels.

Figure 4: Iterative *in vivo* cut'n'paste of two maltose-utilization loci

a) Illustration of iterative *in vivo* cut'n'paste of the *malHM* (i.e., *malEFGH* and *malKlamBmalM*) and *malQT* (i.e., *malPQ* and *malT*) maltose-utilization loci. **b)** Maltose utilization of CNP strains. Spotting of liquid culture onto maltose-MacConkey+Tet/agar. Imaged after 9 hours. Dark red disks: cells can utilize maltose, pale yellow disks with halo: cells cannot utilize maltose. The orange-reddish background depicts MacConkey/agar. I: CNP^{empty}, II: CNP^{malHM}, III: CNP^{malHM} ΔEPI^{malHM}::EPI^{empty}, IV: CNP^{malHM,malQT}, and V: CNP^{malHM,malQT} ΔEPI^{malHM,malQT}::EPI^{empty}. Scale bar: 10 mm.

Figure 5: Episome retention in absence of antibiotic selection

a) False-colored fluorescence (GFP; excitation: 455-485 nm, emission: 508-557 nm, exposure time: 15 msec) and **b)** brightfield images of CNP$^{folA,TetR::gfp}$ plated onto LB agar at transfers during serial culturing in LB without any antibiotic supplementation. Numerals within the panels indicate transfer numbers. Cultures reached comparable densities of OD$_{600}$ ≈4 at each transfer. Scale bar: 30 mm.

Supplementary Figure 1: Potential for "merging" of synthetic DNA fragments during

simultaneous pre-loading

Illustration of "merged" dimers of the synthetic DNA fragments PRE[epi] (yellow) and

PRE[chr] (blue) due to sequence homology at shared features. "Merging" of PRE[epi] and

PRE[chr] causes feature loss. Locations of "merging" are highlighted in bold and,

additionally, stated on the left of the illustrations. Definitions as defined in Fig. 1. Not

to scale.

Supplementary Figure 2: Potential integration events of synthetic DNA fragment

PRE^{epi} during sequential pre-loading

Illustration of integration events of the synthetic DNA fragment PRE^{epi} (yellow) due to

homology at pairs of shared features into **a)** the episome, **b)** the native chromosome,

or **c)** a native chromosome containing PRE^{chr} (blue). Integration events may cause

feature loss. Many pairs of shared features cannot undergo successful integration due

to their relative orientations (feature orientation mismatch) or relative locations (feature

order mismatch). Integration homologies are highlighted in bold and, additionally,

stated on the left side of the illustrations. Definitions as defined in Fig. 1. Not to scale.

A)

PRE^{chr} integration into chromosome

HR^C & HR^D

B)

PRE^{chr} integration into episome

impossible (no shared pairs of features)

C)

PRE^{chr} integration into pre-loaded episome

CS^1 & HR^D	impossible (feature order mismatch)
CS^1 & CS^2	impossible (feature orientation mismatch)
CS^1 & HR^{TET}	impossible (feature orientation mismatch)
HR^D & CS^2	impossible (feature orientation mismatch)
HR^D & HR^{TET}	impossible (feature orientation mismatch)
CS^2 & HR^{TET}	impossible (feature order mismatch)

Supplementary Figure 3: Potential integration of synthetic DNA fragment PRE^{chr}

during sequential pre-loading

Illustration of integration events of the synthetic DNA fragment PRE^{chr} (blue) due to

homology at pairs of shared features into **a)** the native chromosome, **b)** the episome,

or **c)** an episome containing PRE^{epi}. Most pairs of shared features cannot undergo

successful integration due to their relative orientations (feature orientation mismatch)

or relative locations (feature order mismatch). Homologies used for integration are

highlighted in bold and, additionally, stated on the left side of the illustrations.

Definitions as defined in Fig. 1. Not to scale.

Supplementary Figure 4: Insertion of locus-specific guide sequence into pCut_GFP

via I-CreI cutting counterselection-aided plasmid recombineering

a) Overview of I-CreI cutting counterselection-aided plasmid recombineering workflow.

b) Image of transformants (1:10^4 dilution) plated onto LB-agar containing ampicillin

[100 mg/L] (plate diameter: 90 mm). **c)** Agarose gel electrophoresis of diagnostic

PCRs across the engineering location in pCut (primers: diag_guide_A, diag_guide_B).

Non-green transformants (lanes 2-7), green transformants (lanes 8 and 9), and "gene

ruler [1 kb]" DNA ladder (lanes 1 and 10). Expected band sizes are indicated by black

triangles.

Supplementary Figure 5: *E. coli* K-12 MG1655 WT plated on LB agar

a) False-colored fluorescence (GFP; excitation: 455-485 nm, emission: 508-557 nm,

exposure time: 15 msec) and **b)** brightfield images of *E. coli* K-12 MG1655 WT plated

onto LB agar. Identical contrast settings as in Fig. 5. Scale bar: 30 mm.