The functional morphology of hawkmoth lamina monopolar cells reveals mechanisms of spatial processing in insect motion vision

Ronja Bigge[1*], Kentaro Arikawa[2], Anna Stöckl*[1]

[1] Department of Biology, University of Konstanz, Universitätsstr. 10, 78464 Konstanz, Germany

[2] Research Center for Integrative Evolutionary Science, SOKENDAI, Hayama, Kanagawa 240-0193, Japan

* corresponding authors: Ronja Bigge, ronja.bigge@uni-konstanz.de
                                     Anna Stöckl, anna.stoeckl@uni-konstanz.de

Abstract

Many animals strongly rely on their sense of vision, as it provides information about the natural world with particularly high dimensionality. In insects, the first visual processing stage of the brain, the lamina, plays an important role in parallel processing of this complex information. Its main relay neurons, lamina monopolar cells (LMCs), receive information directly from the photoreceptors and shape the contrast, luminance, spatial and temporal tuning of the insect visual system in a cell-type specific manner. One of their best-investigated downstream targets is the motion vision pathway. However, how LMC types that feed into motion processing delineate contrast and luminance is only known from fruit flies, while the contribution of LMCs to spatial processing has only been described in hawkmoths. Here, we provide a novel characterization of hawkmoths lamina monopolar cells, to integrate the contrast, luminance and spatial processing properties of LMCs in the motion pathway. We used serial block-face scanning electron microscopy to reconstruct the anatomical fine structure of LMCs in a focal lamina cartridge, including their pre- and post- synaptic connections. Combining our novel LMC classification with intracellular recordings, we further investigated the functional role of hawkmoth L1 and L2 in terms of contrast and spatial processing. We show that unlike in flies, L1 and L2, the main relay neurons to the motion pathway, process contrast and luminance information in a similar manner. We further demonstrated that the spatial processing properties of these cells are highly similar as well, and can be explained by the density and the distribution of their synapses across different lamina layers. Based on these findings, we propose that the different lamina layers support distinct connectivity and functional roles in spatial processing.

Introduction

The visual system acquires sensory information about the environment with particularly high dimensionality, providing luminance, contrast, colour, polarisation information, as well as multiple derived qualities, such as motion or depth information (Land & Nilsson, 2012). To extract relevant features for specific behavioural tasks, neural systems filter and categorize the visual input early on, to reduce its complexity (Niven & Laughlin, 2008). A key feature of such early processing are parallel pathways, to process features simultaneously (Macpherson et al., 2021). In vertebrates, for instance, retinal ganglion cells initialize parallel channels in the retina projecting ultimately to the visual cortex, where spatially defined, cell-specific connectivity recombines these inputs into new parallel outputs (Watanabe & Rodieck, 1989), reviewed in (Nassi & Callaway, 2009). Despite the structural differences in organization, the insect visual system employs very similar parallel processing strategies (Clark & Demb, 2016).

A prime area in the insect brain where parallel processing manifests both morphologically and physiologically is the lamina. As the first processing stage of visual information, it is retinotopically organized into repeated sets of neurons, the so-called cartridges. Each cartridge processes information from one functional "pixel" of the image, typically receiving information from the photoreceptor axons of a single ommatidium in the eye (with the exception of dipteran flies (Braitenberg, 1967; Kirschfeld, 1967)). Depending on species (Strausfeld, 2012), it contains 4-5 types of lamina monopolar cells (LMCs) each, of which processing the photoreceptor input. In forwarding this information to the next brain area, the medulla, they form the beginning of distinct visual pathways (Cajal & Sanchez, 1915; Strausfeld, 1970; Strausfeld & Blest, 1970).

A key pathway that lamina neurons provide input to is motion vision (Borst, 2014). In fruit flies, the two central LMCs in each cartridge, L1 and L2, initiate two parallel pathways with opposing contrast information: brightness increments and decrements, so-called ON (L1) and OFF (L2) components of the visual signal (Joesch et al., 2010). L1 and L2 are the primary postsynaptic partners of photoreceptors (Rivera-Alba et al., 2011), and have the largest axon diameters and similar branching patterns in the lamina proper (Fischbach & Dittrich, 1989), but project to different layers in the medulla (Meinertzhagen & O'Neil, 1991). Physiological investigations have shown that L1 and L2 have different contrast and luminance coding properties, where L1 encodes a larger luminance component of the visual signal, while L2 encodes primarily contrast (Ketkar et al., 2022, Ketkar et al., 2023).

In addition to luminance and contrast processing, the insect lamina has been implicated in spatial processing since the first descriptions of lamina monopolar cells (Cajal & Sanchez, 1915), because of a prominent morphological characteristic: LMC types in many insect species

extend lateral processes into neighboring cartridges, and are thus ideally suited to either
spatially inhibit or pool information from neighboring visual units. As these processes extend
much further in most nocturnal insects than in diurnal ones (Greiner et al., 2004a; Ribi, 1977;
Stöckl et al., 2016b; Strausfeld & Blest, 1970; Wolburg-Buchholz, 1979), they were
hypothesized to spatially integrate visual information, thereby enhancing the visual signal and
reducing associated noise in dim light (Greiner et al., 2005; Greiner et al., 2004b;
Meinertzhagen et al., 1983; Strausfeld & Blest, 1970; Strausfeld & Braitenberg, 1970; Warrant,
1999). Since the main relay neurons of photoreceptor information in the fruit fly lamina, L1-L3,
lack processes that extend into neighboring cartridges (Fischbach & Dittrich, 1989; Rivera-
Alba et al., 2011), the spatial integration function of LMCs has been mainly investigated in other
insects. Evidence for spatial summation was physiologically demonstrated in the so-called type
II neurons in the lepidopteran hummingbird hawkmoth, which integrate information from their
long processes in dim light, while mainly receiving input from short processes in bright
conditions (Stöckl et al., 2020). This dynamic adjustment of spatial tuning and visual sensitivity
is measurable in the responses of motion neurons, suggesting that the investigated LMC types
in hawkmoths contribute significantly to the motion vision pathway (Stöckl et al., 2017).

However, it is unclear how the spatial processing functions described in hawkmoths align with
the contrast and luminance processing contributing to motion vision in fruit flies, since
homology between lamina monopolar cells across these species has not been established. To
reconcile the contribution of LMCs to spatial, contrast and luminance processing in motion
vision across insects, we investigated the anatomical fine structure of all LMCs and
photoreceptors in a cartridge of the hummingbird hawkmoth down to the synaptic level. We
thereby established a novel classification of LMCs in hawkmoths, proposing homology to other
insects with lamina connectome data. We further characterized the contrast and luminance
coding properties of individual lamina neurons using intracellular recordings. Finally, we
investigated the LMC's spatial response properties physiologically, and propose how these
arise from the LMC's synaptic connectivity profiles.

Methods

Anatomy

Tissue preparation and image acquisition

The tissue preparation for serial block-face scanning electron microscopy (SBF-SEM) was performed as previously described in (Deerinck, 2010) and will briefly be summarized here. We dissected the brains out of the head capsule and pre-fixated them in 4% glutaraldehyde and 1% paraformaldehyde in 2mM $CaCl_2$ in 0.15M cacodylate buffer (pH 7.3, Ca-CB). The dissection and pre-fixation was performed at Würzburg University, Germany, and the following procedures at Sokendai, Japan. The brains were then washed in cacodylate with 2 mM $CaCl_2$, embedded in 35% gelatin-filled mold and cut into 150 µm thick sections using a vibratome. The sections were stored in Ca-CB overnight at 4 °C. The next day, the sections were treated with 2 % osmium tetroxide (OsO_4) and 1.5 % K-ferrocyanide in Ca-CB for 1 h on ice. Then, they were washed with Milli-Q water (5 x 3 minutes at room temperature). Subsequently, the sections were treated with 1 % thiocarbohydrazide in Milli-Q water (20 minutes at room temperature) and then, washed with Milli-Q water (5 x 3 minutes at room temperature). After that, the sections were transferred to 2 % osmium tetroxide (OsO_4) in Milli-Q water for 30 min at room temperature. They were then again washed with Milli-Q water (5x 3 min at RT) and subsequently transferred to 1 % uranyl acetate in Milli-Q at 4 °C overnight. The next day, the sections were transferred to Walton's lead aspartate solution for 30 min at 60 °C. Finally, the sections were then dehydrated in an increasing ethanol series (5 min each on ice), treated with acetone, and embedded in hard Spurr's resin (Polyscience, Inc.).

The serial images were obtained using SBF-SEM (Zeiss Gemini SEM300, Gatan 3 View). The samples were aligned such that the images were taken perpendicular to the cell axon's longitudinal axis. In total, about 3000 images were collected at a 70 nm interval in z-direction, covering 80 x 80 µm in x-y direction (10,000 x 10,000 pixels, pixel size = 8 nm). The images contain about 70 cartridges in total and cover the cells from the cell body layer, along the full extent of the lamina to the first optic chiasm.

Image Analysis

3D segmentation

We registered the image stack using TrakEM2 (Cardona et al., 2012) in Fiji (Schindelin et al., 2012). We then obtained a detailed volumetric reconstruction of the fine morphology of the different types of lamina monopolar cells, as well as the photoreceptors manually using 3Dmod (3dmod Version 4.9.12, University of Colorado). For the 3D segmentation, we selected all neurons in one central cartridge in the image stack.

To confirm that the detailed morphology of neurons in the focal cartridge extended to other
cartridges, we used Catmaid (Version 2021.12.21, CATMAID Developers) to assess the
number and types of lamina monopolar cells and photoreceptors, and their locations within
multiple neighboring cartridges. Further, we created skeleton reconstructions of 5 pairs of L1
and L2 lamina cells to quantify the variation in the lengths and extension of their lateral
processes across cartridges.

Synaptic connectivity

We identified all synaptic connections of the LMCs in the focal cartridge manually in 3dmod. A
synaptic connection was classified as an aggregation of vesicles next to the neuron's
membrane in one cell, and there were 1 or more cells attached to that membrane close to the
vesicle cluster (see Fig. 1E). Within these synapses we distinguished pre- and postsynaptic
connections, as well as synaptic connection between the LMC and a photoreceptor or with
another cell type. On L4, where we found the synapse distribution along the processes and
the spread of the processes to be highly consistent, we localized the synapses on a subset of
3 processes and interpolated the total number based on these.

Physiology

Electrophysiological recordings

To perform intracellular electrophysiological recordings, the moths were fixated: their wings
were cut off and their thorax and abdomen were restrained by wrapping the animals in tape
such that only the head remained free. To reduce movement further, any dental wax was
applied on the thorax to minimize thorax movements, and the head-thorax area to fixate the
head. In addition, the proboscis was extended and fixated in front of the head, and the
antennae were fixated frontally on the head. Subsequently, all scales were removed from the
head capsule and a hole was cut, extending from the dorsal rim area of the left eye to the base
of the left antenna.

We used borosilicate glass electrodes (Sutter Instrument Borosilicate glass with filament,
BF150-75-10) to impale the lamina neurons. The electrodes were pulled either using a P-97
micropipette puller with a 2.5 trough filament (Sutter instruments) and a P-1000 micropipette
puller with 2.5 box filament (Sutter instruments). When recording, the electrode tips were filled
with Neurobiotin (Vector Laboratories, Burlingame, USA, 4% dissolved in 1M KCl) and
backfilled with 1M KCl for a subset of the experiments and filled with Alexa Fluor 568 Hydrazide
(ThermoFisher Scientific) and backfilled with 200mM KCl for another set of recordings. The
resistances of the pulled electrodes ranged from 80 MOhm to 120 MOhm for Neurobiotin
injections but were considerably higher for A568 recordings, with resistances up to 400 MOhm.
After recording the neuronal responses, the neurons were dye injected by applying a constant

iontophoretic current. For Neurobiotin, a depolarizing current of 1.3 nA was applied for 2-3 minutes, while for Alexa568, a hyperpolarizing current of 2.5 nA was applied for 3-5 minutes.

The electrodes were controlled by a manual micromanipulator attached to a piezo control box. The recorded neural responses were pre-amplified by a headstage and then amplified by a bridge amplifier (BA-03X, NPI electronic GmbH, Tamm, Germany). Signals were monitored with an oscilloscope and an audio monitor and then digitized by using a data acquisition card (National Instruments, USB 6221). The stimuli and data acquisition were generated and controlled by a custom written Matlab script (Matlab2015a, Mathworks) used with psychtoolbox.

Histology

The brains containing dye injected neurons were dissected out of the head capsule and transferred to a petri dish containing phosphate buffer, in which the trachea and any other loose tissue were removed from the brain. The retina of the left optic lobe was cut off with fine scissors to fully expose the lamina. Neurobiotin-injected brains were fixated at 4 °C overnight in neurobiotin-fixative (4 % paraformaldehyde, 0.25 % glutaraldehyde, 0.2 % saturated picric acid in 0.1 M phosphate buffer). After fixation, brains were washed 5 times for 15 minutes in 0.1 M PBS (containing 0.145 M NaCl) and then incubated with Streptavidin conjugated Alexa Fluor 568 (ThermoFisher Scienfitic) 1:1000 in PBT (0.1 M PBS containing 0.3 % Triton X-100) for 3 days in darkness at 4 °C. After the incubation, the brains were washed 2 x for 30 minutes in PBT and 3 x 30 minutes in PBS. Then, brains were dehydrated in an ascending ethanol series (30 %, 50 %, 70 %, 90 %, 95 %, 100 %, 100 %) for 15 minutes each. Afterwards, they were transferred to a 1:1 mix of 100 % ethanol and methyl salicylate for 20 minutes and then cleared in pure methyl salicylate for 1h or overnight. After clearing, the brains were embedded between two coverslips in Permount (Fisher Scientific, Schwerte, Germany).

The A568-injected brains were fixated in 4 % paraformaldehyde-fixative for 30 minutes at room temperature. After fixation, the brains were dehydrated, cleared and embedded as described for the neurobiotin injections.

Visual stimulation and data analysis of contrast and luminance response properties

During recordings, the animals were placed in front of a screen at a distance of 20 cm, such that their left eye centered on the middle of the screen. Visual stimuli were generated by a custom written Matlab script (Matlab2015a, Mathworks) used with the psychtoolbox presented

at a frame rate of 120 Hz. The screen was linearly calibrated to ensure brightness consistency

across the whole visual field. If applicable to the stimulus, they were corrected for the animals'

viewing perspective to that all pixels of the screen subtended the same visual angles in the

field of view of the animal.

To examine the contrast and luminance response properties of the cells, we presented either

a brightness increment (ON step) or decrement (OFF step), changing the screen from mid grey

to white or to black, which results in a Weber contrast of 1 or -1, respectively. We presented

stimuli with two durations: a short impulse of 8 ms and an impulse of 500 ms duration. The

stimulus was preceded and followed by a 1 second adaptation phase presenting a mid-grey

screen. Each stimulus was repeated at least 10 times during a recording.

Custom-written code in Matlab (Matlab2022b, Mathworks) was used for all following analyses.

To compare the response characteristics of individual cells, we normalized the responses by

the maximum amplitude and averaged all stimulus repetitions. The 8 ms impulse responses

were fitted with a double lognormal function, including a positive and a negative deflection. The

fitting function included 6 parameters, which were optimized using the inbuilt Matlab function

'fminunc':

216
$$a_1 \times e^{-\frac{(\log(\frac{x}{\mu_1}))^2}{2\sigma_1{}^2}} - a_2 \times e^{-\frac{(\log(\frac{x}{\mu_2}))^2}{2\sigma_2{}^2}}$$

To investigate potential response clusters, which could result from different cell types with

idiosyncratic responses, we first reduced the parameter dimensions by performing a principal

component analysis (PCA). As an input to the PCA, all parameters were transformed into z-

scores to eliminate differences in units, with a mean 0 and standard deviation of 1. The results

provided a measure for the proportion of variance in the data explained by each principal

component (Fig S1 A, E), as well as the contribution of each parameter to each principal

component (Fig S1 B, F). To reduce dimensionality, we incorporated the principal components

that explained at least 80 % of the variability in the data. For both, ON and OFF impulse

responses, this required 3 principal components. With these 3 principal components we

performed k-means clustering to assess differences in the response properties between the

recorded cells. To estimate the optimal number of clusters, we calculated the silhouette score

(Fig. S1 C, G), which was used as a criterion for the probability that the data was correctly

assigned to the right number clusters. To further assess the separation between the clusters,

the silhouette value for each data point within the clusters were calculated (Fig S1 D, H):

numbers close to 1 indicate a clear separation between clusters, whereas low and negative

values indicate that the data is not certainly assigned correctly to its cluster.

The same analysis was applied to the 500 ms impulse responses with some adjustments. The
data was fitted by a single lognormal function, thus fitting only the cells' response to the onset
of the stimulus:

236
$$a \times e^{-\frac{(\log(\frac{x}{\mu}))^2}{2\sigma^2}}$$

237   This fit resulted in 3 fitting parameters, based on which the PCA was performed. For both,
238   ON and OFF data the first two principal components explained more than 90 % of the
239   variance of the data (Fig S1 I,M). Thus, the subsequent cluster analysis was performed on
240   the first two principal components and statistically analyzed by the silhouette method (Fig S1
241   K, L, O, P).

Visual stimulation and data analysis of spatial characteristics

We tested the spatial response properties of the cells by presenting a moving black bar on a
white background. The bar moved from left to right and vice versa, as well as up to down and
vice versa. Each bar orientation was presented 8 times, and the receptive field of the strongest
responses (either horizontal or vertical) was extracted. We normalized each trial by the
maximum response amplitude and averaged all normalized responses. Subsequently, we fitted
the averaged receptive fields using the inbuilt Matlab function 'isqcurvefit' combined with a
double Gaussian function, one of which had a positive and the other one a negative amplitude,
creating a difference-of-Gaussian shape:

251
$$a_c * e^{-\frac{(x-\mu_c)^2}{2\sigma_c^2}} - a_s * e^{-\frac{(x-\mu_s)^2}{2\sigma_s^2}}$$

This resulted in 6 parameters per receptive field. We reduced these parameters to 5 by
calculating the relation between the two amplitudes ($a = a_c/a_s$). We further reduced the
dimensionality of the data using a PCA (as described above). The first 4 principal components
explained more than 80% of the data (Fig S2 A). Based on these we then evaluated the optimal
number of clusters for k-means clustering, using the silhouette score and the gap statistic to
evaluate the separation of clusters (Fig S2 C, D).

Results

A cartridge in the hawkmoth lamina contained previously unknown cells

We used serial block-face scanning electron microscopy (SBF-SEM) (Denk & Horstmann, 2004) to reconstruct the fine structure of the neurons in one lamina cartridge of the hummingbird hawkmoth *Macroglossum stellatarum*.

The SBF-SEM data revealed a very structured retinotopic organization of lamina cells and photoreceptor axons, which formed tight cartridges that could readily be separated throughout the full extent of the imaged tissue (Fig 1B). Each cartridge contained nine photoreceptor axons (Fig 1C). Two of which were long visual fibers, extending into the medulla, whereas the remaining seven photoreceptors were short visual fibers that terminated 20 µm before the proximal end of the lamina. The 3D reconstruction of the photoreceptors revealed that they did not have any lateral processes. In addition to the photoreceptor axons, each cartridge contained five lamina monopolar cells (Fig 1C). This finding was surprising, as previous hawkmoth LMC classifications based on Golgi stainings described only four different types (Stöckl et al., 2016b). These results can be reconciled, since two of the cells we described matched closely in morphology with a previously described type.

The five LMCs showed extensive lateral branching (Fig 1D), which we used to define three lamina layers: a distal layer A, a middle layer B and proximal layer C. The axon terminals of the short visual fibers demarked the border between layers B and C, and the branching patterns of the LMCs defined the border between layers A and B. Of the five LMCs, the two cells with the largest axon diameters had very similar lateral branching patterns: they had long lateral processes in layers A and C, which left the home cartridge and extended up to three to four neighboring cartridges laterally. In the central layer B, they had mostly short lateral processes which remained in the home cartridge (Fig 1 C, D). Both cells had close contact to each other via their axon membranes throughout the full extent of the lamina (Fig. 1B right image). The LMC with the third biggest axon diameter had few lateral processes in layers A and C, and a regular spacing of short lateral processes in layer B, which were restricted to the home cartridge. While the axons of these three cell types remained in close contact to each other throughout the whole lamina, the two remaining cells were located further outside the cartridge boundaries (Fig 1 B right image). Their axons were considerably smaller in diameter. One of the cells had very regular, mid length, branched lateral processes in layers A and B, but no processes in layer C. The fifth cell had few lateral processes in layers A and C, which remained in the home cartridge, and were bulky and filled with vesicles. This cell type did not have processes in layer B. We found the same number and arrangement of cell types in all surrounding cartridges (35 in total) across the whole image stack.

Figure 1: **Anatomical reconstruction of a lamina cartridge in the hummingbird hawkmoth.**
A 3D volume rendering of the hummingbird hawkmoth brain; the lamina is highlighted in yellow. The position of the lamina section analyzed is indicated by the black box. **B** Serial block face electron microscopy. The image stack consisted of 3018 images of 10,000 x 10,000 pixels, with a pixel size of 8 nm. The intersection distance between the images was 70 nm. On the left: overview of the image stack, showing a cross-section of the lamina. On the right: the same image at higher magnification to highlight the focal cartridge that was analyzed (in color) **C** 3D reconstructions of all neurons within the reconstructed cartridge. From left to right: the cartridge neuron bundle, seven short visual fibers (svf), two long visual fibers (lvf) and five lamina monopolar cells. Shaded areas indicate the lamina layers A, B and C. These layers are based on the dendritic morphology as well as the terminals of the svfs. **D** Number and the extent of the LMC's lateral processes (measured as the straight-line distance from the axon center to the end point of each process). The shaded areas show the average cartridge diameter (=10 µm). **E** Based on vesicle clusters visible in the SBF-SEM data, we identified synaptic connection between the cells, and classified these into post- (left image) and presynaptic (right image) partners. **F** Pre - and post-synaptic connections of all LMC types of the focal cartridge, separated into synaptic partners (photoreceptors and other lamina neurons). The color code indicates the total number of synapses for the individual pairings.

A novel LMC classification for hawkmoths based on synaptic connectivity

To identify LMC types, we relied on the neuron's morphology and their synaptic connectivity, and compared these features with the existing literature on the two insect species with 3D morphology and connectome reconstructions; the fruit fly *Drosophila melanogaster*, and the swallowtail butterfly *Papilio xuthus*. To this aim, we identified pre - and post-synaptic connections of all five LMCs in the focal cartridge to both photoreceptors and other neurons (Fig. 1E). As in other insects, we termed the pair of neurons with the largest axon diameters and a central placement in the cartridge L1 and L2 (Boschek, 1971; Ribi, 1978, 1987; Shimohigashi & Tominaga, 1999). Their connectivity matches L1 and L2 cells across other insect species, as they have the highest number of synapses with photoreceptors out of all LMC types (Fig. 1F) (Rivera-Alba et al., 2011). While L1 and L2 are morphologically indistinguishable in the lamina of most insect species, their connectivity differs: L1 has more presynaptic connections compared to L2 in fruit flies (Rivera-Alba et al., 2011). We used this indicator to identify L1 and L2 in our 3D reconstructions (Fig. 1F).

Based on the brush-like structure of its lateral processes, its axon diameter and the location within the cartridge, we classified the third LMC type as L3. This cell received its main input from photoreceptors of the home cartridge, which aligns with findings in other insect species (Ribi, 1975a; Ribi, 1975b, 1987, Rivera-Alba et al., 2011). The final neuron, with processes in layers A and C only, has a morphological equivalent in *Drosophila* L5 (Fischbach & Dittrich, 1989, Rivera-Alba et al., 2011) and in desert ants (Ribi, 1975b). In both species, L5 possesses only few lateral processes in layer A and C. However, in *Drosophila*, its connectivity is only postsynaptic (Rivera-Alba et al., 2011), whereas we found mainly presynaptic connections.

The fourth identified LMC, by means of exclusion of the other cell types termed L4, differs distinctly from L4 LMCs in *Drosophila* (Fischbach & Dittrich, 1989), bees (Ribi, 1975a), ants (Ribi, 1975b) and *Papilio aegus* butterflies (Ribi, 1987), which possess long lateral processes in layer C only. It was, however, previously observed in several studies of hawkmoths (Stöckl et al., 2016b; Strausfeld & Blest, 1970) and does bear some resemblance to L4 in *Papilio xuthus* (Matsushita et al., 2022). While morphologically very different, its connectivity, with nearly no postsynaptic connections to photoreceptors, does align with L4 in the two species with known connectivity, *D. melanogaster* (Rivera-Alba et al., 2011) and *P. xuthus* (Matsushita et al., 2022).

Functional differences in L1 and L2 contrast and luminance coding

Having established a new classification of lamina cell types, and identified L1 and L2 neurons, we next investigated whether functional specializations existed between these two cell types,

which have previously been described in fruit flies. We therefore investigated imbalances in
their responses to light increments and decrements, as well as differences in their responses
to contrast stimuli, as the *Drosophila* L1, which initializes the ON pathway, responds with more
sustained responses to contrast steps than L2, which initializes the OFF pathway (Clark et al.,
2011; Joesch et al., 2010, Ketkar et al., 2023). To quantify the response characteristics of
hawkmoth L1 and L2, we recorded intracellularly (Fig 2A) and used neurobiotin injections to
characterize cell types.

The dye injections after recordings confirmed that we were recording from L1 and L2 (Fig 2B).
Interestingly, in all neurobiotin injections, we found at least 2 cells in the same cartridge to be
strongly labelled, suggesting that L1 and L2 were coupled via gap junctions, which neurobiotin
is known to pass. In these double-cell labelled brains, we found terminals in two distinct layers
of the medulla, the first one at the distal end of layer 1 and the second one at the distal end of
layer 2. This aligns the features of L1 and L2 in fruit flies, which are also coupled via gap
junctions (Joesch et al., 2010), and terminate in layers 1 and 5 (L1) and 2 (L2). To separate
the individual LMCs, we used injections of Alexa568 dye, which has a larger molecular
structure than neurobiotin and therefore passes gap junctions less frequently. Thereby, we
could isolate the two terminals in layers 1 and 2, confirming that only single cells were labelled
by registering only single cell bodies (Fig 2 B), which demonstrates that L1 and L2 terminate
in distinct layers in the medulla, as they do in other species.

Figure 2: **Contrast coding properties of L1 and L2.**
A We performed intracellular recordings in the lamina of the hummingbird hawkmoth, using an
LCD screen for visual stimulation. **B** Left: neurobiotin stainings of gap-junction coupled cells.

Staining shows two cell bodies as well as a double terminal in two different medulla layers.
Right: two separate endings in two different medulla layers, based on Alexa568 stainings. **C,**
D Example recordings for OFF (left) and ON (right) impulse stimuli of **C** 8 ms and **D** 500 ms,
respectively. Colored lines represent the raw traces, dashed line the fits. **E, F** Fits to all
individual cell responses for the **E** 8 ms and **F** 500 ms ON and OFF stimulation, respectively.
G, H Parameters of the fitted **G** 8 ms and **H** 500 ms ON and OFF responses: Left: time from
stimulus onset to first peak, time from the first peak to the second, inverse peak, and time from
the second peak to baseline. Right: Ratio between the amplitude of the first peak and the
second peak for ON and OFF. **H** Time to peak extracted from fit data: Left: stimulus onset to
peak, and time from peak to baseline. Right: half-width of the fits for ON and OFF responses.
I, J k-means clustering of the first three principal components for cell responses to **I** 8 ms and
J 500 ms stimuli, respectively, based on the fitting parameters of the cell responses.

Since the Alexas568 injections proofed to be distinctly less reliable than neurobiotin injections,

we could not use this method to screen for functional response differences between the two

cell types. We therefore analyzed whether the responses of all recorded cells, which were

identified as L1/L2, formed two distinct clusters that would delineate the two types.

We began our functional investigation with characteristics of the contrast and luminance coding

of light increments and decrements. L1 and L2 responded with a depolarization to OFF light

steps (Fig 2 C left) and a hyperpolarization to ON steps (Fig 2 C right). To brief, 8 ms impulses,

the cells responded with a quick response peak, followed by a membrane potential reversal

after the offset of the stimulus (which represented an inverse contrast step to the neurons). To

investigate potential differences in response amplitudes to ON and OFF steps between cell

types, as well as differences in the temporal response characteristics, the responses to ON

and OFF steps were fitted with a lognormal function (Fig 2D). From the fits, we extracted

information about the temporal dynamics of the cell such as the time from stimulus onset to

the peak of the cell, as well as time to membrane potential reversal after stimulus offset, the

time to return to baseline, and the ratio between amplitude at the onset and offset of the

stimulus (Fig 2G). We found that the responses to the brief impulse stimuli showed variation

between individual cells, but no consistent differences between the ON and OFF components

in response amplitude or temporal response characteristics (Fig 2G). Additionally, a cluster

analysis was applied to the resulting fitting parameters after dimensionality reduction.

Reviewing the data in the principal component space (Fig 2 I) showed that it was distributed

continuously rather than delineating into two clusters, as would be expected if L1 and L2 had

different response properties. While a statistical evaluation suggested the optimal number of

clusters was five for OFF responses (Fig S1 C), many data points were assigned to the clusters

with low certainty (Fig S1 D). Reviewing the parameters of the ON responses in principal

component space showed a continuous distribution with one outlier. The optimal number of

clusters was two (Fig S1 G), with the outlier being the only one data point assigned to the

second cluster. (Fig 2E right, Fig S1 H). Thus, we did not find that the neurons' responses to

ON and OFF steps clustered into two distinct groups that would represent L1 and L2.

We next tested 500 ms ON and OFF steps (Fig 2 D), as fruit fly L1 and L2 differ in their temporal
response characteristics to such contrast stimuli, with L1 responding with a combination of
transient and sustained responses, thus encoding both contrast and luminance, and L2
encoding primarily the contrast information in highly transient responses (Clark et al., 2011,
Ketkar et al., 2023). To these long contrast steps, most neurons responded with a rapid
decrease or increase in membrane potential (Fig 2 F), depending on the contrast polarity,
which went back to baseline about 10-15 ms later (Fig 2 H). Some neurons had more transient
responses, which did not return to baseline until stimulus offset (fig 2 H). To assess whether
these responses fell into two distinct categories, we performed a cluster analysis similar to the
8 ms contrast steps. As for those, assessing the response parameters in principal component
space showed a continuous distribution for both ON and OFF steps (Fig 2J). The optimal
number of clusters for OFF and ON was three and two (Fig S1 K,L & O,P), with only very few
cells represented in the second or third clusters, which did not fit the expectation of two equally
distributed response groups corresponding to L1 and L2, respectively.

Across short and long contrast steps with brightness increments and decrements, we did not
observe two distinct response clusters which we would have expected if L1 and L2 had clearly
distinguishable contrast response patterns.

Spatial properties of L1 and L2

We next investigated whether there were differences in the spatial morphology of L1 and L2,
and the corresponding spatial response properties. This question was initiated by the
morphological differences of the L1 and L2 neuron that were fully 3D reconstructed in the focal
cartridge (Fig. 1C, D): the focal L2 neuron had distinctly longer lateral processes than L1. A
skeleton reconstruction of the lateral processes of L1 and L2 in five additional cartridges
revealed, however, that these differences in lateral extent were not consistently aligned with
neuron type, but varied from cartridge to cartridge.

Correspondingly, we analyzed the spatial response properties of the neurons with intracellular
recordings. In the bright light intensities we used, these revealed receptive fields with a center-
surround structure (Fig 3A). We therefore fitted the receptive fields with a difference of
Gaussians (Fig 3B), which revealed an average center half-width of 2.4 degrees, and an
average half width of the inhibitory surround of 4.6 degrees (Fig C). Clustering of the receptive
field parameters similarly to the temporal properties revealed that the optimal number of
clusters was one (Fig S2 C,D), again suggesting no distinct differences in spatial responses
between L1 and L2.

Figure 3. **Spatial processing properties of L1 and L2.**
A Example recording of an L1 / L2 receptive field. Colored line shows the raw trace, dashed line shows the fit. **B** Difference-of-Gaussian fits of all individual receptive field measurements. **C** Half-width of the center of the receptive field, the width of the inhibitory field (measured from peak to peak), and on the right the half-width of the receptive field measured in dim light (data from Stöckl et al., 2020). **D** the amplitude of the left and the right inhibitory flank. **E** left: 2D projection of the synapse distribution in the different lamina layers. right: The synapse distribution across the radial extent of the neurons in the three lamina layers.

Synaptic distribution reveals spatial processing strategies

Since L1 and L2 in hawkmoths play a key role in spatial processing, both lateral summation in low light (Stöckl et al., 2020) and as demonstrated here, lateral inhibition in bright light

(Fig. 3B), we next investigated whether the distribution of pre - and postsynapses across layers
A, B and C would reveal how these processing strategies delineate across the morphology of
the neurons (Fig 3E). In layer B, the short lateral processes contained within the home
cartridge, carried only postsynapses, while presynapses were found on the long processes in
layers A and C. This suggests that lateral processing is confined to layers A and C. In these
layers, the postsynapses were present on processes that extended outside of the home
cartridge. This synaptic architecture provides some important insights into the mechanisms of
spatial processing in L1 and L2: the center of the receptive fields of the neurons matched the
spread of the postsynapses in layer B, suggesting that these were formed by the information
which the short processes of the neurons integrated. Lateral inhibition and spatial summation
could be performed both by the long processes in layers A and B, respectively. One noticeable
feature of the synaptic architecture is that the presynapses of the long processes were confined
almost entirely to the home cartridge. If these mitigate spatial summation or lateral inhibition,
the extent of this processing would therefore only be determined by the length the lateral
processes of other L1 and L2 neurons, as there would be no synaptic transmissions between
the long processes of L1 and L2 outside of the respective home cartridges of both pre- and
postsynaptic partners.

Discussion

479 Using serial block-face scanning electron microscopy (SBF-SEM), we provided a novel
480 classification of lamina monopolar cell types in a hawkmoth species, based on 3D-
481 morphological features and synaptic connectivity. Our analysis revealed five lamina monopolar
482 cells (LMCs) per cartridge, which we assigned to the established L1-L5 types homologous to
483 other insect species.

Reconciling lamina classifications across species

In order to extrapolate and compare functional characteristics of LMCs across insect species,
homology of these cell types is required. Establishing homology based on the neurons'
morphology in the lamina and medulla has proven to challenging. First off, different insect
species are reported with different numbers of LMCs per cartridge: while most species across
insect groups possess five LMC types (Diptera: fruit flies (Fischbach & Dittrich, 1989),
Hymenoptera: desert ants (Ribi, 1975b), Coleoptera: fireflies (Ohly, 1975), Lepidoptera:
skipper butterfly (Shimohigashi & Tominaga, 1999), hawkmoths, and in some cartridges of the
Asian swallowtail (Wakita et al., 2024). Other species, in the same insect groups, were only
reported to have four (Hymenoptera: honey bee (Ribi, 1975a), tropical sweat bee (Greiner et

al., 2004b), Lepidoptera: orchard swallowtail butterfly (Ribi, 1987)). It is possible that a fifth
LMC is present in all or some of the latter species, and was missed because two LMC types
share a similar morphology, as was the case for the hummingbird hawkmoth we investigated
here.

Second, some morphological features are highly consistent across species, while others can
differ drastically. Some neuron types share consistent morphological features, for example the
brush-like processes on one side of the axon only associated with L3 in ants (Ribi, 1975b),
honeybees (Ribi, 1975a), the fruit fly (Fischbach & Dittrich, 1989), firefly (Ohly, 1975), the
orchard swallowtail butterfly (Ribi, 1987) and the hummingbird hawkmoth (Stöckl et al., 2016b),
Fig. 1C)). However, neurons with this characteristic morphology are absent in other species,
such as sweat bees (Greiner et al., 2004b), and the Asian swallowtail (Wakita et al., 2024).
This morphological variation is striking, particularly since it is also present across relatively
closely related insect species. Another example of diverging morphology is L4, which possess
lateral processes that only extend within lamina layer C in fruit flies (Fischbach & Dittrich, 1989)
and in the orchard swallowtail butterfly (Ribi, 1987), but not in the hummingbird hawkmoth, or
the Asian swallowtail (Wakita et al., 2024). The axon diameter, and position within a cartridge
is also highly consistent for some neuron types, particularly L1 and L2, which are typically at
the center of a cartridge and have the largest axon diameters of all LMC types in house flies
(Pyza & Meinertzhagen, 1995), fruit flies (Fischbach & Dittrich, 1989), the honeybee (Ribi,
1975a), the hummingbird hawkmoth, but not in the Asian swallowtail butterfly (Wakita et al.,
2024).

Another commonly used approach to classify LMC types across insects is the termination layer
of the LMC axon terminals in the medulla, which provided the initial naming convention for fruit
fly LMCs (Fischbach & Dittrich, 1989). However, while different LMC types consistently project
to distinct medulla layers within a species, the termination layers themselves are not conserved
across insect groups: across the known lamina neuron types in Hymenoptera, L1 terminates
in layer 1 honeybees and desert ants, and in layer 2 in sweat bees, while this relation is
swapped for L2. L3 and L4 also do not terminate in the same layers across honeybees, sweat
bees, and desert ants (Ribi, 1975b), (Greiner et al., 2004b; Ribi, 1975a).

Thus, the morphology of the neurons in the lamina and medulla does not provide sufficient
information to establish homology. We suggest that combining neurons' synaptic connectivity
with their 3D morphology makes it possible to identify functional homologies, across
Lepidoptera (*Papilio xuthus* (Matsushita et al., 2022), *Macroglossum stellatarum* in this study)
and Diptera (*Drosophila melanogaster* (Rivera-Alba et al., 2011)). Despite their striking
morphological differences, L1, L2 and L3 in these species are the main postsynaptic partners

to photoreceptors. L4 has little to no postsynaptic connections to photoreceptors, but receives
most of its inputs from other (lamina) cells. L5 is mainly presynaptic in all 3 species.

The similarities in connectivity could be the result of an evolutionarily conserved wiring scheme
of the lamina across insect species, as is also known from other brain areas (Heinze, 2024),
while the species' specific differences in LMC morphology reflect adaptations to specific
processing requirements, in terms of luminance adaptation, or colour processing.

Functional homology of the main rely neurons of the motion pathway

Leveraging the proposed homology between LMCs in the hummingbird hawkmoth and fruit fly,
we investigated whether these LMCs share functional characteristics across species. We
focused on neurons with the best physiological characterization, L1 and L2, which are the main
contributors to the wide-field motion vision pathway in fruit flies (Borst, 2014; Borst et al., 2020;
Egelhaaf, 2023). They separate visual information in light increments (ON pathway), and
decrements (OFF pathway) at their terminals in the medulla (Joesch et al., 2010, Behnia et al.,
2014). Even though L1 and L2 are known to be interconnected via gap junctions in fruit flies
(Joesch et al., 2010), L2 in fruit flies shows distinct selectivity to OFF signals in its terminal
Calcium responses, suggesting that signal rectification happens in the cell. Similar to the fruit
fly, we demonstrated that in hummingbird hawkmoth L1 and L2 are linked via gap junctions
and their terminals also contact layers 1 and 2 in the medulla (Fig. 2B).

We attempted to identify which of the terminals was associated with L1 and L2, respectively,
as identified based on the SBF-SEM data. However, since the 3D morphology of L1 and L2
was highly similar, and the extend of the lateral processes could also not be used as an
indicator of neuron identity, as it varied more strongly within than across L1 and L2 cell types,
assigning identity of the recorded and subsequently labelled neurons via morphological means
was impossible. In our recordings of hawkmoths L1/L2, we also did not observe two distinct
response clusters to ON and OFF visual stimuli, which would represent distinct response
characteristics to light increments and decrements, suggesting that both morphologically, and
in terms of their response polarity to light, L1 and L2 are not distinguishable.

We therefore planned to establish neuron identity via their temporal response characteristics:
L1 and L2 are known to differ in their responses to changes in light in fruit flies: L1 exhibits
more sustained responses, encoding both contrast and luminance information, while L2
responds highly transiently, encoding only contrast information of a visual stimulus transient
versus sustained responses to luminance changes (Clark et al., 2011, Ketkar et al. 2023). In
hawkmoths, we did not find distinct response clusters that would indicate that L1 and L2 had
different response properties in terms of contrast and luminance. Both responded highly
transiently, as reported previously (Stöckl et al., 2020), indicating that they primarily encode

contrast information, but not luminance (Fig 2 F). Thus, if hawkmoths L1 and L2 also initialize
ON and OFF pathways in the motion circuitry, both pathways are provided with highly similar
response properties in terms of increment and decrement amplitude and contrast coding by
the two cell types.

Reconciling spatial and contrast processing in motion processing

Our identification of L1 and L2 reconciled two processing features in insect motion processing:
segregation of light information into increments and decrements (ON and OFF pathways), and
spatial integration. It has been suggested anatomically (Greiner et al., 2005; Stöckl et al.,
2016b) in various insect species, and demonstrated physiologically in the hummingbird
hawkmoth (Stöckl et al., 2020) that the lateral processes of LMCs that extend into neighboring
cartridges are capable of integrating visual information in dim light, to improve the signal to
noise ratio, and thus contrast sensitivity, of motion responses (Stöckl et al., 2017; Stöckl et al.,
2016a). How such spatial processing applies to the parallel ON and OFF arms of the motion
pathway was not clear, because the neurons previously classified as L1 and L2 in hawkmoths
had very different lateral extents, suggesting that spatial processing was applied
asymmetrically to the ON and OFF pathways.

With our novel LMC classification in hawkmoths we demonstrate that L1 and L2 supply the
same potential for spatial processing, in terms of lateral extent, to the ON and OFF arms of the
motion pathway. Our physiological investigation of the neuron's receptive fields supports this
notion, as we found no evidence for two distinct clusters in the spatial receptive fields (Fig. 3C,
Fig S2). Thus, assuming that L1 and L2 initiate On and OFF pathways in hawkmoths similar
to fruit flies, we demonstrate that spatial information is transmitted symmetrically into both
arms. Whether this is also the case in other insects with asymmetric spatial morphologies in
L1 and L2 (Greiner et al., 2004b; Ribi, 1975a; Ribi, 1987) remains to be tested.

Synaptic architecture for spatial processing

In addition to the previously described spatial integration of information in dim light, which
dynamically broadens receptive fields of L1/L2 in the hummingbird hawkmoth (Stöckl et al.,
2020), in our current recordings in bright light we observed another spatial processing
phenomenon, which is observed in many peripheral visual neurons in bright light: lateral
inhibition (Fig 2 B)(Hartline, 1969; Kramer & Davenport, 2015; Mimura, 1976; Zettler &
Jarvilehto, 1972), thought to enhance lateral contrast for improved spatial vision, particularly
in bright light (Priebe & Ferster, 2008; Strausfeld & Campos-Ortega, 1977).

With our 3D reconstructions and synapse tracking, we can provide mechanistic explanations
for these spatial processing features of lamina monopolar cells, based on the density and
distribution of pre- and postsynaptic connections of L1 and L2. Notably, we find that almost all

postsynaptic connections to photoreceptors in L1 and L2 are concentrated in layer B, with no

synaptic connections to other neurons in this layer. This suggests that layer B is the primary

site for sampling photoreceptor input.

Lateral processing in hummingbird hawkmoth L1 and L2 thus concentrates in layers A and C,

separated by the main input layer. Our connectivity data provides strong hypothesis for how

spatial processing functions segregate across these layers: we propose that the lateral

processes in layer A perform spatial summation of information in low light. The distribution of

postsynapses in layer A fits with the extent of the spatial receptive fields measured in low light

Fig. 4, when the neurons are performing spatial summation (Stöckl et al., 2020). Importantly,

our synaptic data answers an open question concerning the extent of spatial summation: it

was previously shown that the extent of the recorded receptive fields in low light matched the

extend of the long lateral processes of L1 / L2 in hummingbird hawkmoths. However, assuming

connections between these processes, the receptive fields should have been considerably

wider. Our data, however, demonstrates, that there are no (chemical) pre-synapses on the

lateral processes beyond the extent of the home cartridge, ruling out extra-cartridge

connections between L1/L2 neurons (Fig 3 D). Thus, the capacity for spatial summation

(assuming no involvement of additional neurons in the lamina) in hummingbird hawkmoths is

constrained by the extent of their lateral processes, which explains the close match between

their spatial receptive fields in dim light, and their process distributions observed previously

(Stöckl et al., 2020).

L1 / L2 processes in layer C contain considerably more presynapses than postsynapses than

in layer A. Together with short visual fibres terminating in layer B, this suggests that primary

input processing occurs before layer C. Instead, layer C appears to be well suited for lateral

inhibition. Correspondingly, the lateral extent and distribution of postsynapses in layer C

matches the visual angle of the lateral inhibitory flanks of the receptive fields in bright light

Fig. 4.

Based on these findings, we propose that the different lamina layers support distinct

connectivity and functional roles. Specifically, our findings support the existence of a layer-

based mechanism for spatial processing in the hawkmoth lamina, separating summation and

lateral inhibition in a similar, – albeit much simpler – architecture as the vertebrate cortex (Luo,

2021). Our results not only reinforce the notion that morphology and connectivity provide strong

indicators of cell function, but they also allow us to formulate new hypotheses about the

homology of functional circuitry across insect species.

Author Contributions Conceptualization: A.S.; Methodology: A.S., K.A., R.B.; Validation:
R.B.; Formal analysis: R.B.; Investigation: R.B.; Resources: A.S., K.A.; Data curation: R.B.;
Writing - original draft: R.B.; Writing - review & editing: R.B., K.A., A.S.; Visualization: R.B.;
Supervision: A.S.; Funding acquisition: A.S., K.A.

Acknowledgements We would like to thank Christian Stigloher for excellent suggestions
regarding 3D tracing tools, and Nadine Kaiser for supporting the 3D neuron reconstructions.

Funding We acknowledge funding to A.S. by the German Research Council (STO 1255 2-1),
the Young Scholar Fund of the University of Konstanz, and the European Research Council
(101116305 DynamicVision ERC-2023-STG), and to K.A. by the KAKENHI of the Japan
Society for the Promotion of Science (JSPS) (23K26914).

Competing Interest Statement: The authors declare no competing interests.

References

Behnia, R., Clark, D. A., Carter, A. G., Clandinin, T. R., & Desplan, C. (2014, Aug 28). Processing properties of ON and OFF pathways for Drosophila motion detection. *Nature, 512*(7515), 427-430.

Borst, A. (2014, Sep). Fly visual course control: behaviour, algorithms and circuits. *Nat Rev Neurosci, 15*(9), 590-599.

Borst, A., Haag, J., & Mauss, A. S. (2020, Mar). How fly neurons compute the direction of visual motion. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol, 206*(2), 109-124.

Boschek, C. B. (1971). On the fine structure of the peripheral retina and lamina ganglionaris of the fly, Musca domestica. *Z Zellforsch Mikrosk Anat, 118*(3), 369-409.

Braitenberg, V. (1967). Patterns of projection in the visual system of the fly. I. Retina-lamina projections. *Exp Brain Res, 3*(3), 271-298.

Cajal, S. R. y., & Sanchez, D. (1915). Contribución al conocimiento de los centros nerviosos de los insectos.

Cardona, A., Saalfeld, S., Schindelin, J., Arganda-Carreras, I., Preibisch, S., Longair, M., Tomancak, P., Hartenstein, V., & Douglas, R. J. (2012). TrakEM2 software for neural circuit reconstruction. *PLoS One, 7*(6), e38011.

Clark, D. A., Bursztyn, L., Horowitz, M. A., Schnitzer, M. J., & Clandinin, T. R. (2011, Jun 23). Defining the computational structure of the motion detector in Drosophila. *Neuron, 70*(6), 1165-1177.

Clark, D. A., & Demb, J. B. (2016, Oct 24). Parallel Computations in Insect and Mammalian Visual Motion Processing. *Current Biology, 26*(20), R1062-R1072.

Deerinck, T. J. (2010, 2010). NCMIR methods for 3D EM: a new protocol for preparation of biological specimens for serial block-face SEM. *Microscopy [Online]*, 6.

Denk, W., & Horstmann, H. (2004, Nov). Serial block-face scanning electron microscopy to reconstruct three-dimensional tissue nanostructure. *PLoS Biol, 2*(11), e329.

Egelhaaf, M. (2023, Jul). Optic flow based spatial vision in insects. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol, 209*(4), 541-561.

Fischbach, K.-F., & Dittrich, A. (1989). The optic lobe of Drosophila melanogaster. I. A Golgi analysis of wild-type structure. *Cell and tissue research, 258*(3), 441-475.

Golgi, C. (1873). Sulla struttura della sostanza grigia del cervello. *Gaz. Med. Ital. Lombardia, 6*, 244-246.

Greiner, B., Ribi, W. A., & Warrant, E. J. (2004a, Jun). Retinal and optical adaptations for nocturnal vision in the halictid bee Megalopta genalis. *Cell Tissue Res, 316*(3), 377-390.

Greiner, B., Ribi, W. A., & Warrant, E. J. (2005, Nov). A neural network to improve dim-light vision? Dendritic fields of first-order interneurons in the nocturnal bee Megalopta genalis. *Cell Tissue Res, 322*(2), 313-320.

Greiner, B., Ribi, W. A., Wcislo, W. T., & Warrant, E. J. (2004b, Nov). Neural organisation in the
first optic ganglion of the nocturnal bee Megalopta genalis. *Cell Tissue Res, 318*(2), 429-
437.

Hartline, H. K. (1969, Apr 18). Visual receptors and retinal interaction. *Science, 164*(3877),
270-278.

Heinze, S. (2024) Variations on an ancient theme — the central complex across insects.
Current Opinion in Behavioral Sciences. Volume 57, 101390.

Joesch, M., Schnell, B., Raghu, S. V., Reiff, D. F., & Borst, A. (2010, Nov 11). ON and OFF
pathways in Drosophila motion vision. *Nature, 468*(7321), 300-304.

Ketkar, M. D., Gur, B., Molina-Obando, S., Ioannidou, M., Martelli, C., & Silies, M. (2022, Mar
9). First-order visual interneurons distribute distinct contrast and luminance information
across ON and OFF pathways to achieve stable behavior. *Elife, 11*.

Ketkar, M. D., Shao, S., Gjorgjieva, J., & Silies, M. (2023, Jul 10). Multifaceted luminance gain
control beyond photoreceptors in Drosophila. *Current Biology, 33*(13), 2632-2645 e2636.

Kirschfeld, K. (1967). [The projection of the optical environment on the screen of the
rhabdomere in the compound eye of the Musca]. *Exp Brain Res, 3*(3), 248-270.

Kramer, R. H., & Davenport, C. M. (2015, Dec). Lateral Inhibition in the Vertebrate Retina: The
Case of the Missing Neurotransmitter. *PLoS Biol, 13*(12), e1002322.

Land, M. F., & Nilsson, D.-E. (2012). *Animal Eyes*. OUP Oxford.

Luo, L. (2021, Sep 3). Architectures of neuronal circuits. *Science, 373*(6559), eabg7285.

Macpherson, T., Matsumoto, M., Gomi, H., Morimoto, J., Uchibe, E., & Hikida, T. (2021, Dec).
Parallel and hierarchical neural mechanisms for adaptive and predictive behavioral
control. *Neural Netw, 144*, 507-521.

Matsushita, A., Stewart, F., Ilic, M., Chen, P. J., Wakita, D., Miyazaki, N., Murata, K., Kinoshita,
M., Belusic, G., & Arikawa, K. (2022, May 23). Connectome of the lamina reveals the
circuit for early color processing in the visual pathway of a butterfly. *Current Biology,
32*(10), 2291-2299 e2293.

Meinertzhagen, I. A., Menzel, R., & Kahle, G. (1983). The identification of spectral receptor
types in the retina and lamina of the dragonflySympetrum rubicundulum. *Journal of
Comparative Physiology ? A, 151*(3), 295-310.

Meinertzhagen, I. A., & O'Neil, S. D. (1991, Mar 8). Synaptic organization of columnar
elements in the lamina of the wild type in Drosophila melanogaster. *J Comp Neurol,
305*(2), 232-263.

Mimura, K. (1976). Some spatial properties in the first optic ganglion of the fly. *Journal of
Comparative Physiology ? A, 105*(1), 65-82.

Nassi, J. J., & Callaway, E. M. (2009, May). Parallel processing strategies of the primate visual
system. *Nat Rev Neurosci, 10*(5), 360-372.

Niven, J. E., & Laughlin, S. B. (2008, Jun). Energy limitation as a selective pressure on the
evolution of sensory systems. *J Exp Biol, 211*(Pt 11), 1792-1804.

Ohly, K. P. (1975). The neurons of the first synaptic region of the optic neuropil of the firefly,
Phausis splendidula l. (Coleoptera). *Cell Tissue Res, 158*(1), 89-109.

Priebe, N. J., & Ferster, D. (2008, Feb 28). Inhibition, spike threshold, and stimulus selectivity
in primary visual cortex. *Neuron, 57*(4), 482-497.

Pyza, E., & Meinertzhagen, I. A. (1995, Jan). Monopolar Cell Axons in the First Optic Neuropil
of the Housefly, Musca-Domestica L, Undergo Daily Fluctuations in Diameter That Have a
Circadian Basis. *Journal of Neuroscience, 15*(1), 407-418.

Ribi, W. A. (1975a). The first optic ganglion of the bee. *Cell and tissue research, 165*(1), 103-
111.

Ribi, W. A. (1975b, Jul 8). Golgi studies of the first optic ganglion of the ant, Cataglyphis
bicolor. *Cell Tissue Res, 160*(2), 207-217.

Ribi, W. A. (1977). Fine structure of the first optic ganglion (lamina) of the cockroach,
Periplaneta americana. *Tissue Cell, 9*(1), 57-72.

Ribi, W. A. (1978, Dec 28). Gap junctions coupling photoreceptor axons in the first optic
ganglion of the fly. *Cell Tissue Res, 195*(2), 299-308.

Ribi, W. A. (1987). Anatomical Identification of Spectral Receptor Types in the Retina and
Lamina of the Australian Orchard Butterfly, Papilio-Aegeus-Aegeus D. *Cell and tissue
research, 247*(2), 393-407.

Rivera-Alba, M., Vitaladevuni, S. N., Mishchenko, Y., Lu, Z., Takemura, S. Y., Scheffer, L.,
Meinertzhagen, I. A., Chklovskii, D. B., & de Polavieja, G. G. (2011, Dec 6). Wiring economy
and volume exclusion determine neuronal placement in the Drosophila brain. *Current
Biology, 21*(23), 2000-2005.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,
Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J. Y., White, D. J., Hartenstein, V., Eliceiri, K.,
Tomancak, P., & Cardona, A. (2012, Jun 28). Fiji: an open-source platform for biological-
image analysis. *Nat Methods, 9*(7), 676-682.

Shimohigashi, M., & Tominaga, Y. (1999, May 24). Synaptic organization in the lamina of the
superposition eye of a skipper butterfly, Parnara guttata. *J Comp Neurol, 408*(1), 107-124.

Stöckl, A. (2022, May 23). Connectomics: From form to function in butterfly colour vision.
Current Biology, 32(10), R459-R461.

Stöckl, A. L., O'Carroll, D., & Warrant, E. J. (2017, Jun 28). Higher-order neural processing
tunes motion neurons to visual ecology in three species of hawkmoths. *Proc Biol Sci,
284*(1857), 20170880.

Stöckl, A. L., O'Carroll, D. C., & Warrant, E. J. (2016a, Mar 21). Neural Summation in the
Hawkmoth Visual System Extends the Limits of Vision in Dim Light. *Current Biology, 26*(6),
821-826.

757
Stöckl, A. L., O'Carroll, D. C., & Warrant, E. J. (2020, Apr). Hawkmoth lamina monopolar cells
act as dynamic spatial filters to optimize vision at different light levels. *Sci Adv, 6*(16),
eaaz8645.

Stöckl, A. L., Ribi, W. A., & Warrant, E. J. (2016b, Jan 1). Adaptations for nocturnal and diurnal
vision in the hawkmoth lamina. *J Comp Neurol, 524*(1), 160-175.

Strausfeld, N. J. (1970). Golgi studies on insects Part II. The optic lobes of Diptera.
Philosophical Transactions of the Royal Society of London. B, Biological Sciences, 258(820),
135-223.

Strausfeld, N. J. (2012). *Arthropod brains: evolution, functional elegance, and historical
significance*. Belknap Press.

Strausfeld, N. J., & Blest, A. (1970). Golgi studies on insects Part I. The optic lobes of
Lepidoptera. *Philosophical Transactions of the Royal Society of London. B, Biological
Sciences, 258*(820), 81-134.

Strausfeld, N. J., & Braitenberg, V. (1970). The compound eye of the fly (Musca domestica):
connections between the cartridges of the lamina ganglionaris. *Zeitschrift für
vergleichende Physiologie, 70*(2), 95-104.

Strausfeld, N. J., & Campos-Ortega, J. A. (1977, Mar 4). Vision in insects: pathways possibly
underlying neural adaptation and lateral inhibition. *Science, 195*(4281), 894-897.

Wakita, D., Shibasaki, H., Kinoshita, M., & Arikawa, K. (2024, Feb). Morphology and spectral
sensitivity of long visual fibers and lamina monopolar cells in the butterfly Papilio xuthus. *J
Comp Neurol, 532*(2), e25579.

Warrant, E. J. (1999, May). Seeing better at night: life style, eye design and the optimum
strategy of spatial and temporal summation. *Vision Res, 39*(9), 1611-1630.

Watanabe, M., & Rodieck, R. W. (1989, Nov 15). Parasol and midget ganglion cells of the
primate retina. *J Comp Neurol, 289*(3), 434-454.

Wolburg-Buchholz, K. (1979, Mar 9). The organization of the lamina ganglionaris of the
hemipteran insects, Notonecta glauca, Corixa punctata and Gerris lacustris. *Cell Tissue
Res, 197*(1), 39-59.

Zettler, F., & Jarvilehto, M. (1972). Lateral Inhibition in an Insect Eye. *Zeitschrift für
vergleichende Physiologie, 76*(3), 233-+.

Supplement

Figure S1: **A** Cumulative explained variance of the individual principal components which are based on the fit data of 8 ms OFF impulse responses (Fig. 2). **B** The heatmap shows which fitting parameters contribute most to which of the principal components. **C** The silhouette score gives a measure for the optimal number of clusters. Values under 0.6 are considered very low and highlight a high uncertainty. **D** The silhouette plot for the suggested number of clusters. Negative values indicate a high chance of datapoints being assigned to the wrong cluster. Similarly, low positive values indicate a high uncertainty in cluster assignment. **E-H** same analysis as A-D for 8 ms ON responses, **I-L** for 500 ms OFF impulse responses, and **M-P** for 500 ms ON responses.

800 **Figure S2**: **A** Cumulative explained variance for the principal components corresponding to
801 the spatial receptive field fit (Fig. 3). **B** Correspondence of fitting parameters (see Methods)
802 and principal components. **C** Statistical analysis of the optimal number of clusters. **D** The
803 silhouette plot for five clusters. Negative and low values indicate a low certainty in assigning
804 the data points to the distinct clusters.

806