1

Reward impulsivity is linked to addictive-like behaviors towards sweet food and sugary drinks but not fatty food

Snigdha Mohan[1], Georgiy Bobashev[2], Victoria Moiseeva[1], Anna Shestakova[1], Ksenia Panidi[1*], Boris Gutkin[1,3]

[1]Centre for Cognition and Decision Making, Institute for Cognitive Neuroscience, HSE University, Moscow, Russia

[2]RTI International, 3040 E. Cornwallis Road, PO Box 12194, Research Triangle Park, NC, 27709, USA

[3]Group for Neural Theory, LNC2 INSERM U960, Département d'études cognitives, Ecole Normale Superieure, PSL Université Paris, 75005 Paris, France

* Corresponding author: k.a.panidi@gmail.com (KP)

Abstract

Overconsumption of hyperpalatable food is a worldwide problem. Although many previous studies have shown the link between monetary delay discounting and propensity to overconsume food, the studies did not focus on the link between delay discounting and various subcategories of palatable food. In the present study, we investigate the link between the individual propensity to choose smaller immediate rather than larger delayed reward and the degree of problematic eating behavior for five types of palatable food (sweets, sugary drinks, fatty food, salty snacks, and starchy food). The delay discounting task was used to measure propensity to choose immediate rather than delayed rewards, while the degree of addictive-like behavior was measured using the modified Yale food addiction scale (mYFAS) for each food category. Both model-free and model-based analysis showed that there is a significant correlation between the addiction score for sweet food and the tendency to choose smaller immediate rather than larger delayed rewards. A model-based analysis showed the same significant correlation for sugary drinks. However, no such correlation was observed for other food categories (fatty, salty, or starchy food), suggesting that

32 reward impulsivity may play different roles in addictive-like behavior towards various food

33 categories.

Introduction

Highly palatable food (i.e. food high in salt, fat and sugar) is known to be able to evoke addictive-

like behavior [1,2]. Studies on addictive eating behavior typically focus on energy-dense foods in

general, naming high-sugar and/or high-fat products as most common in binge-eating disorder and

bulimia nervosa. However, not many studies exist that explore differences in addictive-like

behavior towards different food categories containing various combinations of macronutrients,

although this issue may have important implications for public policy and clinical interventions.

Existing research suggests that both high-fat and high-sugar types of food may be associated with

addictive tendencies, but the underlying physiological and psychological mechanisms of this

association may differ. In the study of [3] among the set of 35 different foods products high in fat

content and glycemic load were most likely to be associated with problematic eating behavior. A

study with animal models demonstrated that food containing a combination of fat and sugar was

most likely to cause activity in the mesolimbic system compared to sugar or fat alone [4]. Significant

differences in demand curves for high-sugar/high-fat versus low-sugar/high-fat products were

found using economic methodology [5]. Individuals were also reported to be willing to pay more for

products containing a combination of fat and sugar rather than for fat or sugar alone, controlling

for caloric value, and this behavioral response correlated with greater activation of brain reward

circuits [6]. Some studies suggested that sugar and fat may differ in inducing additive-like behavior

because of the differences in how they affect dopaminergic system, which leads to withdrawal

symptoms for sugary but not fatty food [7]. Sweet taste was also proposed as a possible mediator of

addictive behavior and withdrawal symptoms, although the evidence to support this statement is

mixed [8].

The aim of the present study is to investigate the relationship between addictive tendencies for five

different categories of highly palatable food (sweets, salty snacks, fatty food, sugary drinks, and

starchy food) and individual impulsivity measured by a monetary delay discounting task. Delay

discounting is a concept that implies the decrease in the value of a reward when the reward is

delayed in time. Monetary delay discounting coefficient is currently used in many studies as an

indicator of reward impulsivity together with trait impulsivity measured by questionnaires like

Barratt Impulsiveness Scale [9], and motor impulsivity measured by the performance in a Go/NoGo

task [10] . The choice of smaller and immediate rewards is considered impulsive whereas choosing

larger delayed rewards is regarded as a self-controlled choice [11]. When an individual frequently

prefers an immediate smaller reward to a delayed larger reward this indicates a steeper discounting

curve and a greater degree of impulsivity [12].

Steep delay discounting in the monetary domain has been shown to be associated with various

types of addictive behaviors (see [13] for a review). The link between delay discounting and addictive

behavior towards food has been established as well [14,15]. A recent review provides evidence that

binge-eating disorder is positively associated with reward impulsivity measured by a delay

discounting task [16]. Delay discounting was also suggested to be a behavioral marker of obesity [17].

These findings might indicate that the common temporal discounting mechanism stands behind

addictive behavior in general. Therefore, if addictive behavior towards sugar stems from the same

mechanism we should observe a similar correlation between delay discounting in the monetary

domain and addictive behavior towards various foods high in sugar content. In order to separate

the effect of fat on sugar-containing food we specifically test for the link between delay

discounting and addictive tendencies towards sugary drinks, separately from other food categories,

like fatty, starchy, and salty food.

In our study, we asked participants to perform a multi-item monetary choice task to measure their

temporal discounting in the monetary domain. Participants also filled in the modified Yale Food

Addiction Scale (mYFAS) questionnaire for five food categories. Based on their answers we then

analyzed the link between the monetary delay discounting and the addiction score in the five food

categories. In the model-free analysis, we investigated the link between the probability to choose

immediate rather than delayed reward and the addiction scores for various food categories, without

any specific assumptions regarding the delay discounting model. In the model-based analysis, we

first use the participants' responses in the delay discounting task to fit exponential and hyperbolic

delay discounting functions, and select the model that fits the behavior best for each particular

participant. We then focus on two characteristics of delay discounting: the shape of the discounting

curve (exponential or hyperbolic) that best describes each participant's choices, and their

corresponding discounting coefficient. Exponential delay discounting describes consumers whose

preferences are time-consistent [18]. If a consumer prefers an apple over a chocolate bar a week from

now, she will still choose an apple over a chocolate bar when asked about what to eat today.

However, many people demonstrate hyperbolic discounting, which leads to preference reversals

typically observed when the behavior is impulsive. A participant offered an apple versus a

chocolate bar in a week from now will make a healthy choice for an apple; if asked to choose what

to eat right now they prefer a chocolate bar. Therefore, individuals may be first categorized based

on the discounting model (exponential or hyperbolic) that best describes their choices in a delay

discounting task. Additionally, individuals with the same discounting type may be further

compared based on their discounting coefficient. Those with higher discounting coefficient may

be considered more impulsive.

In the model-free analysis, we test the hypothesis that participants with higher addiction scores for

sugar-enriched food (such as sweets and sugary drinks) will have higher probability to choose an

immediate rather than delayed rewards in a monetary delay discounting task. We found that the

hypothesis is confirmed for sweet food, but not for sugary drinks or any other type of food.

Participants with higher addiction score for sweets showed higher sensitivity to the amount of

money received immediately, with higher immediate amount leading to higher probability to

choose immediate option compared to participants with lower addiction scores.

In the model-based analysis, we test two hypotheses. Our first hypothesis holds that individuals

who are hyperbolic rather than exponential discounters will be more likely to have higher mYFAS

scores for the food categories that contain high levels of sugar. Our second hypothesis suggests

that within the group of hyperbolic discounters steeper discounting will be associated with a higher

addiction score on the mYFAS scale for sugar-enriched food.

We found that indeed hyperbolic discounters are significantly more likely to have higher addiction

scores for sweet food and sugary drinks. Additionally, among hyperbolic discounters higher

discounting coefficient (higher impulsivity) was significantly associated with a higher addiction

score in the sweet food category. The same tendency was observed for sugary drinks, although it

did not reach a conventional statistical significance threshold, possibly because of a lower number

of participants showing high addiction scores in this particular category. No statistically significant

association was found for fatty, starchy, or salty food.

The present results suggest that reward impulsivity may play a specific role in addictive behaviors

towards high-sugar products even when fat is not present in them. The fact that no such relationship

was observed for fatty, starchy or salty food may reflect the differences in the addictive

mechanisms for sugar-enriched versus other types of food.

Materials and methods

Participants

Sixty-seven participants between the age of 18 and 40 years (mean 23y.o., min = 18y.o., max =

38y.o., 52 females) were recruited via a local social media platform and existing university

participation database. We excluded participants who self-reported having diabetes, any mental

disease, taking prescribed medication, or following any dietary restrictions. The upper age limit

145 was selected to be 40 because taste sensitivity declines with age and is significantly different after

146 the age of 50 [19].

148 The data collection started on 20/08/2020 and ended on 15/09/2020. At a preselection stage,

149 potential candidates filled in an online questionnaire (mYFAS for sweets category). We invited 35

150 participants who showed no addiction to sweets and 32 who showed at least some addiction to

151 sweets based on this questionnaire. The average age of the participants who are addicted to sweets

152 (PWAS) and non-addicted groups (non-PWAS) was 23.50 and 23.32, respectively.

154 For all participants, BMI (Body Mass Index) was calculated according to the classification of the

155 World Health Organization [20]. Body weight and height of the participants were self-reported and

156 used as the measurement to calculate their BMI score. The average BMI score for PWAS and non-

157 PWAS was 22.06 and 21.74, respectively. Our sample comprises individuals who mostly belong

158 within the normal BMI range of 18 to 24.9 with an average score of 22.01. In our sample 70% of

159 the participants had normal BMI, 16% of the participants were underweight, 6% were overweight,

160 and the other 7% were in the obese category.

162 All procedures were approved by the ethics committee of the HSE university. All experimental

163 procedures were performed in accordance with relevant guidelines and regulations.

165 *Experimental Protocol*

The laboratory experiment consisted of two tasks: completing the mYFAS questionnaire for all

five food categories and a computerized temporal discounting task. After completing both tasks

participants filled in the BIS/BAS personality scale.

Participants were asked not to eat 2 hours prior to the experiment to keep a similar level of hunger.

Because of COVID-19, to maintain safety participants and the experimenter wore masks during

the experiment. Before performing the task, all participants signed an informed consent form.

Participants were then given written instructions explaining the experimental procedures in detail.

Participants were informed that their choices during the delayed discounting task did not have any

real-life consequences; however, they were required to make choices as they would in reality. In

addition, they were informed that there is no correct or incorrect choice.

Participants then were asked to perform two tasks in a randomized order with instructions written

on the screen for each task again. The experimenter was not present to minimize the experimenter's

bias. After completing the experiment, participants were debriefed and paid 250 monetary units

(MU) in cash for their participation.

(a) Temporal Discounting Task

In our study, we used the multi-item temporal discounting choice task with a staircase procedure

[21]. Multi-item choice task has been used in addiction studies for alcohol consumption[22,23], smoking

[24,25], and gambling [26,27]. Similarly, several other studies used the staircase procedure to measure

discount rate for a multi-item temporal discounting task [28,29].

The procedure was similar to that of [30]. In each trial, two amounts of hypothetical reward with

respect to their time interval would appear on the screen. One reward was immediate and the other

was after a delay. The participants had to indicate the option they prefer in every trial by pressing

the right or left button. For every trial, a fixation cross was presented for 1 second, after which the

choice block between immediate and later rewards was presented. The chosen option was

highlighted by a box on the screen for 1 second after a 1.5-second interval. Participants made five

choices per block for six time delays (today vs. a delayed day): with 0 days always accounting for

present day monetary gain and 2, 14, 30, 60, and 90 days for delayed monetary reward.

The order of each block and time intervals shown on the screen were randomized. Every block

started with an option of 200 MU for the immediate option. In all trials, the delayed option was

fixed at 400 MU. When the individual chose an immediate reward, the amount of the immediate

reward was decreased for the next trial. On the other hand, if the participant chose the delayed

reward, the amount of the immediate reward was increased for the next trial. The adjustment

amount for the immediate reward decreased with successive choices was chosen following the

study by [31]. In their study, the first adjustment was half of the difference between the immediate

and the delayed reward and the next adjustment was half of the previous modification. In our study,

the subjects made five choices for each time interval in one block. The immediate reward in each

trial represents the subjective value of the delayed reward.

In our experiment for the temporal discounting task we used hypothetical monetary rewards

instead of real rewards. The main concern of using hypothetical monetary rewards is that

participants might not behave as they would in real life because of lack of motivation. However,

studies that have directly compared hypothetical and real monetary rewards did not find any

significant difference between the two [32,33].

(b) mYFAS

Our main instrument in measuring the degree of addiction to sugar enriched food is the mYFAS

2.0 based on seven symptoms of substance dependence using the Diagnostic and Statistical Manual

of Mental Disorders Fifth Edition criteria [34]. mYFAS 2.0 is a 13-item instrument questionnaire

that focuses on calculating an addictive score for food based on an individual's eating pattern of

highly palatable foods in the past year. This scale is a shorter version of the YFAS 2.0

questionnaire, which has been proved to have high convergent validity with respect to various

eating disorder measures and high discriminant validity with respect to alcohol addiction,

behavioral inhibition, and activation measures [35]. The mYFAS 2.0 questionnaire was shown to

have similar properties to YFAS 2.0 in terms of internal consistency and convergent validity

having been tested in various populations [36,37]. Therefore, mYFAS 2.0 is a valid and useful tool to

identify individuals with addiction to food and more specifically in our study with sugar

dependence.

Participants were asked to fill in the mYFAS 2.0 separately for five food categories: sweets

(doughnuts, cookies, ice cream, chocolate, candies), starch (white bread, pasta, rolls, rice), salty

snacks (chips, pretzels, crackers), fatty foods (steak, bacon, hamburger, cheeseburger, pizza,

french fries), and sugary drinks (soda pop). In the current study, a Russian version of mYFAS 2.0

was created. The validity of the questionnaire translation was tested through back-to-back

translation and cognitive interviews.

(c) BIS/BAS

The BAS/BIS questionnaire is 24-item self-report measure that identifies two motivational

systems: the behavioral inhibition system (BIS), and behavioral activation system (BAS) based on

a 4-point Likert scale [38]. BIS indicates the impulse an individual has to avoid aversive outcomes,

whereas BAS is related to the motivation an individual has to approach goal-oriented outcomes.

BAS is further divided into three categories—fun seeking, reward responsiveness, and drive. Each

category corresponds to measuring the motivation behind finding the novel rewards

spontaneously, sensitivity to pleasant reinforcers in the environment, and to follow one's goals,

respectively.

Data Analysis

(1) Model-free analysis

To analyze the behavior in the delay discounting task on a trial-by-trial level we used the logistic

regression model with subject-level random effects. We used each participants' responses on each

trial in a delay discounting task as the dependent variable (equal to 1 if delayed option was chosen,

and 0 otherwise) to analyze the probability to choose the delayed option. The explanatory variables

included the characteristics of the trial (the size of an immediate reward and the time delay to the

delayed reward), as well as the addiction scores for each of the five food categories. Additionally,

participants' age, gender, BMI and BIS and Reward Responsiveness scores from the BIS/BAS

questionnaire were used as control variables. As the delayed reward amount was kept constant in

the task, this variable was not included in the regression model. Importantly, we included the

interaction terms of the immediate reward amount and time delay with addiction scores, to test

whether the propensity towards addictive behavior is linked to the reward and delay sensitivity in

the monetary delay discounting task.

(2) Model-based analysis

The model-based data analysis was performed as follows. First, we used 25 choices each

participant made in a delay discounting task to fit two models of time discounting—exponential

and hyperbolic—with the Bayesian estimation procedure. Then, from these two models for each

individual we selected the one that explained their intertemporal choices best (according to the

Leave-One-Out information criterion). Therefore, after the estimation procedure, each participant

was characterized by two factors: (1) discounting model that fits her choices best (exponential or

hyperbolic), (2) estimated discounting coefficient that corresponds to this best-fitting model. We

further used regression analysis to test two hypotheses. We hypothesized that participants whose

intertemporal choice is best explained by an exponential discounting function will be less likely to

demonstrate high addiction scores compared to those with hyperbolic discounting. Second, we

expected that for participants in a hyperbolic discounting group higher discounting coefficient

(corresponding to steeper discounting) will be associated with higher addiction scores, while no

such association will be observed in the exponential discounting group.

Below we describe each step in detail.

(a) Discount factor

During the temporal discounting task participants chose between the sooner smaller and the larger

later reward on each trial. In accordance with economic theory, we assume that their choice is

based on the subjective present value of each option. We considered two possibilities for the delay

discounting function—exponential and hyperbolic discounting. In economic theory, if a

participant is characterized by the exponential discounting function then their preferences are time-

consistent and do not show preference reversals [39,40]. They are, hence, less likely to demonstrate

impulsive behavior such as addiction. By contrast, participants who are characterized by the

hyperbolic discounting function are time-inconsistent and are more likely to become addicted to

various substances [41].

Mathematically, the present subjective value of a reward is equal to the monetary amount

multiplied by a discount factor. Therefore, for each participant, we fitted two discounting models,

a hyperbolic and an exponential one [42,43]. Exponential discounting function was assumed to have

the following form:

$$V^{reward} = reward \times exp\left(-kD\right) \tag{1}$$

294 where D represents a time delay (in units of time), k is a discounting coefficient, and V^{reward} is the

295 present value of a delayed reward.

296 The hyperbolic discounting model was a 1-parameter hyperbolic discount function [44]:

$$V^{reward} = reward \times \frac{1}{1 + kD} = \frac{reward}{1 + kD} \tag{2}$$

where D and k represent time delay and a discounting coefficient respectively, and V^{reward} is the

present value of a delayed reward.

We further assumed that a participant's choice in every trial is determined by the comparison of

discounted utilities of each option and some random noise. For the participants' choice in each

trial we assume the random utility model with stochastic errors in *Luce* form [45]. According to this

model, a participant would choose the delayed reward if $log(V_B) - log(V_A) > \varepsilon$, where ε represents

a random noise variable, and an immediate reward otherwise. Here, V_B is the present value of a

delayed reward, V_A is the present value of an immediate reward, and ε represents random error

term. We used logs instead of the actual values for better convergence of estimation methods. In

accordance with a common modeling practice in economic studies, we assume that random errors

in each trial are distributed logistically [46,47]. Therefore, the probability of choosing the delayed

reward is as follows:

$$Pr(choose\ delayed) = Pr\ (log\ (V_B) - log\ (V_A) > \epsilon) = \frac{1}{1 + \left(\frac{V_A}{V_B}\right)^{\frac{1}{\mu}}} \qquad (3)$$

Here, μ represents the standard deviation of the noise. If μ tends toward zero, then participants

would always choose the reward with the highest present subjective value. As μ increases the

variance of a random error also increases. Hence, the choice between the two options becomes

almost random.

To obtain the point estimates of the individual discounting parameter we employed the Bayesian

approach. Estimation procedures followed the pipeline described in [48]. We chose uninformative

prior for k as a uniform distribution between 0 and 1 and partially informative prior for $\tau = \frac{1}{\mu}$ as

a uniform distribution between 0 and 10. These parameter constraints are wide enough to cover

the values typically found in other delay discounting studies [32,48]. The obtained posterior

distributions did not reach these boundaries, suggesting that the boundaries were not limiting.

After estimating the parameters of both exponential and hyperbolic discounting function for each

participant, we calculated the Leave-One-Out Cross-Validation information criterion (LOOIC).

For each participant the model with the best fit (lowest LOOIC value) was selected. The type of

discounting model that best describes each participant's preferences was then used as a binary

explanatory variable of interest (equal 1 for hyperbolic and 0 for exponential discounting) in the

regression analysis of addiction scores.

The estimation procedure was adapted from *hBayesDM* package in R software used to conduct

Markov chain Monte Carlo (MCMC) sampling-based inference. The sampling was performed

using 4 MCMC chains with 1,000 iterations for warmup and 4,000 iterations in total.

(b) Regression analysis

First, we tested the hypothesis that the delay discounting type is linked to addiction to sugar-

enriched food. For this purpose, we estimated an ordinal logistic regression model with an

addiction score on the mYFAS scale for each food category separately as a dependent variable.

Explanatory variables included delay discounting type (exponential or hyperbolic), gender, age,

and scores on the BIS and BAS Reward Responsiveness subscales. Because the subscales of the

BIS/BAS questionnaire are correlated with each other we used only BIS and Reward

Responsiveness scales as independent variables to avoid multicollinearity. These variables were

chosen because in many previous studies they demonstrated a significant relation with various

kinds of addictions, such as alcohol and marijuana, and gambling behavior [49–5149–51].

Second, we tested the hypothesis that for participants with hyperbolic discounting the higher value

of the discounting coefficient is associated with increased consumption of sugar-enriched food,

while no such relationship is observed for participants best characterized with the exponential

discounting model. For this purpose, we estimated the same ordered logit model with discounting

coefficient k as the main variable of interest. Since the discounting models between participants

differ, the values of k cannot be pulled together into one variable. Therefore, we included two

separate predictor variables. One predictor consists of k estimates for the exponential model (k_{exp})

multiplied by a dummy variable equal to 1 if the best model for a participant is exponential, and

zero otherwise (I_{exp}). The second predictor is constructed in a similar way with k values

corresponding to the hyperbolic model (k_{hyper}) multiplied by a dummy variable equal to 1 if the

best fitting model for a participant is hyperbolic, and zero otherwise (I_{hyper}). Since for every

participant there is only one best-fitting discounting model (exponential or hyperbolic), the

observations for hyperbolic and exponential k's are not included into the regression equation

simultaneously. Rather we estimate the effect of k on addiction scores separately for each

discounting type, while other control variable effects are estimated on the whole sample of

participants.

Results

In our dataset 30 participants showed no tendencies toward addiction to sweets (score 0-1), 20

participants demonstrated medium level of additive tendencies (score 2-5), and 12 participants had

severe form of addiction (score 6-13). Of 52 women in our sample, 45% belonged to the non-

PWAS category, 35% belonged to the group with moderate addiction to sweets, and 20% had

severe level of addiction to sweets. Of 16 men, 75% belonged to the non-PWAS group and both

the moderate and severe category of sweet addiction comprised 13% of males each. Discounting

model estimates revealed that 47.8% of participants were best characterized with the hyperbolic

discounting model and 52.2% with the exponential discounting model. Among the hyperbolic

discounters the average estimated discounting coefficient equaled 0.16 (s.e. = 0.033), which

corresponds to the devaluation of reward by approximately 14% for the first day of delay. Among

the exponential discounters the average estimated k equalled 0.093 (s.e. = 0.024), which indicates

that for these participants the reward value decreases by 9% per 1 day of delay.

Table 1 presents the estimation results of the regression model in the model-free analysis where

no specific assumptions about the shape of the discounting function are made. Here, the dependent

variable in each regression model is the probability to choose the larger delayed reward as opposed

to the smaller sooner reward in each trial in a delay discounting task. The main explanatory

variables of interest are the individual addiction score for various food categories and the

interaction term of the addiction score with the immediate reward amount and the time delay.

Estimation results showed that in the delay discounting task participants behaved in an expected

way with higher immediate amount and longer time delay leading to lower probability to choose

a delayed option in each trial. For each food category, addiction score did not show any significant

link with the probability to choose a delayed option, which indicates that high degree of addiction

is not associated with an a priori bias towards an immediate or a delayed option. However, the

interaction of addiction score with the immediate reward amount was significant for the Sweets

category, indicating that participants demonstrating high addiction to sweets had greater sensitivity

to the immediate reward amount compared to participants with lower addiction scores to sweets.

An increase in the immediate reward amount by 1 monetary unit would lead to greater decrease in

the probability to choose the delayed option for participants with higher addiction scores to sweets.

No significant association was observed for other food categories.

	Dependent variable: probability to choose delayed option				
Food category	Sweets	Fat	Drinks	Salt	Starch
Immediate amount	-0.007***	-0.009***	-0.009***	-0.009***	-0.009***
	(0.001)	(0.001)	(0.001)	(0.001)	(0.001)
Immediate amount x Addiction score	-0.001*	-0.0003	-0.0001	-0.001	-0.001
	(0.0003)	(0.0003)	(0.001)	(0.0005)	(0.001)
Time delay	-0.039***	-0.038***	-0.040***	-0.038***	-0.040***
	(0.004)	(0.003)	(0.003)	(0.003)	(0.003)
Time delay x Addiction score	-0.0002	-0.0003	0.003	-0.001	0.001
	(0.001)	(0.001)	(0.002)	(0.001)	(0.001)
Addiction score	0.111	-0.062	-0.205	0.082	0.023
	(0.130)	(0.122)	(0.260)	(0.170)	(0.203)
Gender	0.431	0.758	0.511	0.425	0.441
	(0.588)	(0.530)	(0.586)	(0.577)	(0.592)
Age	0.095	0.112*	0.087	0.093	0.093
	(0.060)	(0.054)	(0.060)	(0.059)	(0.060)
BAS Reward Responsiveness	-0.103	0.001	-0.108	-0.123	-0.095
	(0.118)	(0.108)	(0.117)	(0.116)	(0.119)
BIS	0.167	0.204*	0.147	0.164	0.150
	(0.097)	(0.087)	(0.095)	(0.095)	(0.095)
BMI $\geq$ 25	0.307	0.285	0.218	0.177	0.212
	(0.703)	(0.631)	(0.703)	(0.695)	(0.701)
Constant	-0.895	-3.749	0.160	-0.251	-0.411
	(3.046)	(2.828)	(2.988)	(2.932)	(3.017)
Observations	1,563	1,538	1,563	1,563	1,563
Log Likelihood	-851.478	-847.283	-854.476	-853.524	-854.070
Akaike Inf. Crit.	1,726.956	1,718.565	1,732.953	1,731.049	1,732.140
Bayesian Inf. Crit.	1,791.208	1,782.624	1,797.205	1,795.301	1,796.392

Table 1. Logistic regression model with subject-level random effects estimation results.

Dependent variable in each regression is the probability to choose a delayed reward in each trial

in a delay discounting task. Standard errors in parentheses. Stars indicate the following p-value

thresholds: *<0.05, **<0.01, ***<0.001.

Table 2 presents the estimation results of regression model in a model-based analysis where

participants were separated into exponential and hyperbolic discounters based on the discounting

model that best describes their intertemporal choices. Here, the type of discounting was the main

explanatory variable of interest.

	Dependent variable:				
	Sweets	Fat	Drinks	Salt	Starch
	(1)	(2)	(3)	(4)	(5)
Discounting type (hyperbolic=1, exponential=0)	1.144^{*}	0.868	1.245^{*}	0.390	0.649
	(0.484)	(0.521)	(0.632)	(0.595)	(0.514)
Gender	-0.277	0.192	0.793	0.004	-0.602
	(0.549)	(0.614)	(0.708)	(0.769)	(0.635)
Age	0.088	0.067	-0.037	0.016	0.056
	(0.056)	(0.062)	(0.082)	(0.079)	(0.065)
BAS Reward Responsiveness	0.128	0.133	-0.104	-0.033	0.248
	(0.118)	(0.130)	(0.143)	(0.140)	(0.135)
BIS	0.217^{*}	0.039	0.088	0.270^{*}	0.103
	(0.093)	(0.099)	(0.120)	(0.128)	(0.107)
BMI $\geq$ 25	0.394	0.168	-0.539	-0.302	-1.455
	(0.585)	(0.647)	(0.881)	(0.903)	(0.900)
Observations	67	67	67	67	67
R^2	0.254	0.072	0.110	0.109	0.181
chi^2 (df = 6)	19.214^{**}	4.722	6.373	6.661	12.547

Note: *p**p***p<0.001

Table 2. Ordered logit model estimation results. Dependent variable in each regression is the

individual mYFAS score on the corresponding food category. Standard errors in parentheses. Stars

indicate the following p-value thresholds: *<0.05, **<0.01, ***<0.001.

Estimation results suggest that the type of delay discounting is significantly positively associated

with an addiction score on the mYFAS scale for *Sweets* and *Drinks* categories. Namely,

participants who are hyperbolic rather than exponential delay discounters are more likely to have

higher scores on the mYFAS scale. Figure 1 provides further illustration of this result.

Additionally, we observe that a higher BIS score is also linked to increased mYFAS scores for

Sweets and *Salt* food categories.

Figure 1. Average addiction score on mYFAS scale in each food category for exponentially and

hyperbolically discounting participants. Participants exhibiting hyperbolic discounting

demonstrate higher average addiction scores.

421
422   Table 3 presents estimation results for the effects of discounting coefficient k on addiction scores.

	Dependent variable:				
	Sweets	Fat	Drinks	Salt	Starch
	(1)	(2)	(3)	(4)	(5)
k_{exp} x I_{exp}	-1.333	-3.301	-3.101	0.607	-2.911
	(2.530)	(3.715)	(4.642)	(3.424)	(3.016)
K_{hyper} x I_{hyper}	5.213**	2.488	3.426	2.852	3.404
	(1.772)	(1.868)	(2.051)	(2.285)	(2.106)
Gender	-0.210	0.191	0.518	0.036	-0.737
	(0.555)	(0.624)	(0.709)	(0.767)	(0.646)
Age	0.110	0.071	-0.033	0.030	0.057
	(0.058)	(0.065)	(0.085)	(0.081)	(0.067)
BAS Reward Responsiveness	0.153	0.144	-0.069	-0.032	0.282*
	(0.120)	(0.129)	(0.146)	(0.142)	(0.138)
BIS	0.342**	0.081	0.130	0.326*	0.123
	(0.110)	(0.109)	(0.125)	(0.144)	(0.108)
BMI $\geq$ 25	-0.089	-0.110	-0.777	-0.426	-1.835
	(0.598)	(0.699)	(0.933)	(0.894)	(0.944)
Observations	67	67	67	67	67
R^2	0.306	0.089	0.114	0.124	0.224

chi² (df = 7)	23.906**	5.842	6.602	7.682	15.872*

Table 3. Ordered logit model estimation results. Dependent variable in each regression is the individual mYFAS score on the corresponding food category. k_{exp} and k_{hyper} are discounting coefficients estimated assuming exponential and hyperbolic discounting model respectively. I_{exp} is a dummy variable equal to 1 if the participant's choices are best fitted with the exponential discounting model, and zero otherwise. I_{hyper} is a dummy variable equal to 1 if the participant's choices are best fitted with the hyperbolic discounting model, and zero otherwise. Standard errors in parentheses. Stars indicate the following p-value thresholds: *<0.05, **<0.01, ***<0.001.

As shown in Table 3, for exponential discounters the discounting coefficient is not associated with the addiction score in the *Sweets* category. However, for hyperbolic discounters a larger coefficient k (i.e., higher impulsivity in the monetary domain) is significantly linked to a higher addiction score on the mYFAS scale. Interestingly, BIS score is positively associated with the addiction score independently of delay discounting. The same tendency with respect to the delay discounting and mYFAS score is observed in the *Drinks* category; however, it failed to reach significance in our sample (p=0.095). Exponential discounters again do not show any significant correlation with mYFAS scores (p=0.50). No significant association is found in the *Fat*, *Salt,* and *Starch* categories.

Discussion

In the present study, we explore the association between the behavior in the monetary delay

discounting task and addiction to various types of food. Instead of measuring food addiction in

general, we use the mYFAS for five different food categories. Both model-free and model-based

analysis provides evidence that high degree of addiction to sweet food is associated with more

impulsive behavior and a greater sensitivity to monetary rewards in a delay discounting task. In

the model-free analysis no assumptions were made on the delay discounting functional form, and

participants' responses were analysed on a trial-by-trial basis. The analysis showed that higher

addiction score for sweets led to higher sensitivity to immediate monetary reward and higher

probability to make an impulsive monetary choice when this amount is increased. In the model-

based analysis, participants who showed hyperbolic instead of exponential discounting were more

likely to have high addiction scores for sweets. Additionally, for hyperbolic discounters, higher

delay discounting coefficient was also significantly positively associated with the addiction score

for sweet food. The same association was found for addiction to sweet drinks in a model-based

analysis. However, both, model-free and model-based analysis did not show significant

associations between monetary delay discounting and addiction score in other food categories,

such as fatty, starchy or salty food.

The fact that the association between delay discounting and addiction score was observed for

sugary drinks, which represent products that do not contain fat, suggests that sugar may have high

addictive potential independently of other substances like fats. The fact that no such association

was observed for fatty food, may suggest that addictive potential of fatty products may be

decreased in the absence of sugar, or that addictive mechanisms involved in addictive-like

behaviors towards fatty food are distinct from those involved in addictive behavior towards sugar-

enriched food.

Data availability

Complete dataset used in this study is freely available in the repository:

https://github.com/openaccessdata/sugar_addiction

Author contributions:

SM: conceptualization, data curation, formal analysis, drafting original writing;

GB: supervision, reviewing and editing;

VM: project administration, funding acquisition;

AS: supervision, project administration, funding acquisition;

KP: conceptualization, supervision, formal analysis, reviewing and editing;

BG: supervision, reviewing and editing.

References

1. Gordon, E., Ariel-Donges, A., Bauman, V. & Merlo, L. What Is the Evidence for "Food Addiction?" A Systematic Review. *Nutrients* **10**, 477 (2018).
2. Kalon, E., Hong, J. Y., Tobin, C. & Schulte, T. Psychological and Neurobiological Correlates of Food Addiction. *Int Rev Neurobiol* **129**, 85–110 (2016).
3. Schulte, E. M., Avena, N. M. & Gearhardt, A. N. Which foods may be addictive? The roles of processing, fat content, and glycemic load. *PLoS One* **10**, e0117959 (2015).
4. Hoch, T., Kreitz, S., Gaffling, S., Pischetsrieder, M. & Hess, A. Fat/carbohydrate ratio but not energy density determines snack food intake and activates brain reward areas. *Sci Rep* **5**, 10041 (2015).
5. Reslan, S., Saules, K. K. & Greenwald, M. K. Validating a behavioral economic approach to assess food demand: Effects of body mass index, dietary restraint, and impulsivity. *Appetite* **59**, 364–371 (2012).

6.   DiFeliceantonio, A. G. *et al.* Supra-Additive Effects of Combining Fat and Carbohydrate
on Food Reward. *Cell Metab* **28**, 33-44.e3 (2018).
7.   Avena, N. M., Rada, P. & Hoebel, B. G. Sugar and fat bingeing have notable differences
in addictive-like behavior. *J Nutr* **139**, 623–8 (2009).
8.   Greenberg, D. & St. Peter, J. V. Sugars and Sweet Taste: Addictive or Rewarding? *Int J*
Environ Res Public Health **18**, 9791 (2021).
9.   Patton, J. H., Stanford, M. S. & Barratt, E. S. Factor structure of the barratt impulsiveness
scale. *J Clin Psychol* **51**, 768–774 (1995).
10. Batterink, L., Yokum, S. & Stice, E. Body mass correlates inversely with inhibitory
control in response to food among adolescent girls: An fMRI study. *Neuroimage* **52**,
1696–1703 (2010).
11. Pursey, K. M., Davis, C. & Burrows, T. L. Nutritional Aspects of Food Addiction. *Curr*
Addict Rep **4**, 142–150 (2017).
12. Wallace, D. L. *et al.* The influence of ΔfosB in the nucleus accumbens on natural reward-
related behavior. *Journal of Neuroscience* **28**, 10272–10277 (2008).
13. Bickel, W. K., Koffarnus, M. N., Moody, L. & Wilson, A. G. The behavioral- and neuro-
economic process of temporal discounting: A candidate behavioral marker of addiction.
Neuropharmacology **76 Pt B**, 518–27 (2014).
14. Richards, T. J. & Hamilton, S. F. Obesity and Hyperbolic Discounting: An Experimental
Analysis on JSTOR. *Journal of Agrticultural and Resource Economics* **37**, 181–198
(2012).
15. Kekic, M. *et al.* Bad Things Come to Those Who Do Not Wait: Temporal Discounting Is
Associated With Compulsive Overeating, Eating Disorder Psychopathology and Food
Addiction. *Front Psychiatry* **10**, 978 (2019).
16. Carr, M. M., Wiedemann, A. A., Macdonald-Gagnon, G. & Potenza, M. N. Impulsivity
and compulsivity in binge eating disorder: A systematic review of behavioral studies.
Prog Neuropsychopharmacol Biol Psychiatry **110**, 110318 (2021).
17. Bickel, W. K. *et al.* Temporal discounting as a candidate behavioral marker of obesity.
Neuroscience and Biobehavioral Reviews vol. 129 307–329 Preprint at
https://doi.org/10.1016/j.neubiorev.2021.07.035 (2021).
18. Dasgupta, P. & Maskin, E. Uncertainty and Hyperbolic Discounting Uncertainty.
American Economic Review **95**, 1290–1299 (2013).
19. Schaie, K. W. *Intellectual Development in Adulthood*. (Academic Press, 1996).
20. World Health Organization. Obesity: preventing and managing the global epidemic.
Report of a WHO consultation. **894**, 1–253 (2000).
21. Du, W., Green, L. & Myerson, J. Cross-Cultural Comparisons of Discounting Delayed
and Probabilistic Rewards. *Psychol Rec* **52**, 479–492 (2002).
22. Boettiger, C. A. *et al.* Immediate reward bias in humans: fronto-parietal networks and a
role for the catechol-O-methyltransferase 158(Val/Val) genotype. *J Neurosci* **27**, 14383–
91 (2007).
23. Bobova, L., Finn, P. R., Rickert, M. E. & Lucas, J. Disinhibitory psychopathology and
delay discounting in alcohol dependence: personality and cognitive correlates. *Exp Clin*
Psychopharmacol **17**, 51–61 (2009).
24. Sweitzer, M. M., Donny, E. C., Dierker, L. C., Flory, J. D. & Manuck, S. B. Delay
discounting and smoking: Association with the Fagerström Test for Nicotine Dependence

but not cigarettes smoked per day. *Source: Nicotine & Tobacco Research* **10**, 1571–1575
(2008).

25.   Reynolds, B., Richards, J. B., Horn, K. & Karraker, K. Delay discounting and probability
discounting as related to cigarette smoking status in adults. *Behavioural processes* **65**, 35–
42 (2004).

26.   Dixon, M. R., Jacobs, E. A. & Sanders, S. Contextual Control of Delay Discounting by
Pathological Gamblers. *J Appl Behav Anal* **39**, 413–422 (2006).

27.   Ledgerwood, D. M., Alessi, S. M., Phoenix, N. & Petry, N. M. Behavioral assessment of
impulsivity in pathological gamblers with and without substance use disorder histories
versus healthy controls. *Drug Alcohol Depend* **105**, 89–96 (2009).

28.   Solway, A., Lohrenz, T. & Montague, P. R. Simulating future value in intertemporal
choice. *Sci Rep* **7**, 43119 (2017).

29.   Ciaramelli, E., Sellitto, M., Tosarelli, G. & di Pellegrino, G. Imagining Events Alternative
to the Present Can Attenuate Delay Discounting. *Front Behav Neurosci* **13**, 269 (2019).

30.   Terenzi, D., Mainetto, E., Barbato, M., Rumiati, R. I. & Aiello, M. Temporal and Effort
cost Decision-making in Healthy Individuals with Subclinical Psychotic Symptoms. *Sci
Rep* **9**, 2151 (2019).

31.   Myerson, J., Green, L., Scott Hanson, J., Holt, D. D. & Estle, S. J. Discounting delayed
and probabilistic rewards: Processes and traits. *J Econ Psychol* **24**, 619–635 (2003).

32.   Johnson, M. W. & Bickel, W. K. *Within-Subject Comparison of Reak and Hypothetical
Money Rewards in Delay Discounting. Journal of the Experimental Analysis of Behavior*
vol. 77 https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1284852/pdf/11936247.pdf
(2002).

33.   Coller, M. & Williams, M. B. Eliciting Individual Discount Rates. *Exp Econ* **2**, 107–127
(1999).

34.   American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders
(5th Ed.)*. (American Psychiatric Publishing, 2013).

35.   Gearhardt, A. N., Corbin, W. R. & Brownell, K. D. Development of the Yale Food
Addiction Scale Version 2.0. *Psychol Addict Behav* **30**, 113–21 (2016).

36.   Brunault, P. *et al.* The Modified Yale Food Addiction Scale 2.0: Validation Among Non-
Clinical and Clinical French-Speaking Samples and Comparison With the Full Yale Food
Addiction Scale 2.0. *Front Psychiatry* **11**, 892 (2020).

37.   Zhang, H. *et al.* Translation of the Chinese version of the modified Yale Food Addiction
Scale 2.0 and its validation among college students. *J Eat Disord* **9**, 116 (2021).

38.   Carver, C. S. & White, T. L. Behavioral Inhibition, Behavioral Activation, and Affective
Responses to Impending Reward and Punishment: The BIS/BAS Scales. *J Pers Soc
Psychol* **67**, 319–333 (1994).

39.   Frederick, S., Loewenstein, G. & O'donoghue, T. *Time Discounting and Time Preference:
A Critical Review. Source: Journal of Economic Literature* vol. 40
https://www.jstor.org/stable/pdf/2698382.pdf?refreqid=excelsior%3Aeac7c8e55d25f184b
f2437bb5215f0e7&ab_segments=&origin=&acceptTC=1 (2002).

40.   O'Donoghue, T. & Rabin, M. The economics of immediate gratification. *J Behav Decis
Mak* **13**, 233–250 (2000).

41.   MacKillop, J. *et al.* Delayed reward discounting and addictive behavior: a meta-analysis.
Psychopharmacology (Berl) **216**, 305–321 (2011).

42.   Green, L. & Myerson, J. Exponential Versus Hyperbolic Discounting of Delayed
588   Outcomes: Risk and Waiting Time 1. *American Zoology* **36**, 496–505 (1996).
589   43.   Green, L., Fristoe, N. & Myerson, J. *Temporal Discounting and Preference Reversals in*
590   *Choice between Delayed Outcomes. Psychonomic Bulletin & Review* vol. 1
591   https://link.springer.com/content/pdf/10.3758/BF03213979.pdf (1994).
592   44.   Mazur, J. E. An adjusting procedure for studying delayed reinforcement. *Quantitative*
593   *analyses of behavior: V. The effect of delay and of intervening events on reinforcement*
594   *value* 55–73 (1987).
595   45.   Andersen, S., Harrison, G. W., Lau, M. I. & Rutström, E. E. Eliciting risk and time
596   preferences. *Econometrica* **76**, 583–618 (2008).
597   46.   Harrison, G. W. & Elisabet Rutström, E. Risk aversion in the laboratory. *Research in*
598   *Experimental Economics* **12**, 41–196 (2008).
599   47.   Luce, R. D. *Individual Choice Behavior: A Theoretical Analysis.* (Dover Publications,
600   Mineola, 2005). doi:10.1037/14396-000.
601   48.   Ahn, W.-Y., Haines, N. & Zhang, L. Revealing Neurocomputational Mechanisms of
602   Reinforcement Learning and Decision-Making With the hBayesDM Package. *Comput*
603   *Psychiatr* **1**, 24–57 (2017).
604   49.   Kim, D.-Y. & Lee, J.-H. Effects of the BAS and BIS on decision-making in a gambling
605   task. *Pers Individ Dif* **50**, 1131–1135 (2011).
606   50.   Eitle, D. & Taylor, J. General Strain Theory, BIS/BAS Levels, and Gambling Behavior.
607   *Deviant Behav* **32**, 1–37 (2010).
608   51.   Simons, J. S. & Arens, A. M. Moderating effects of sensitivity to punishment and
609   sensitivity to reward on associations between marijuana effect expectancies and use.
610   *Psychol Addict Behav* **21**, 409–14 (2007).
611
612