Title

Coordinated Transcriptional Networks Program Organelle Expansion and Metabolic Flows for High Endothelial Morphology and Function

Yuhan Bi[1,2,3,6]*, Kevin Brulois[1,2,3], Aiman Ayesha[1,2,3], Menglan Xiang[1,2,3], Romain Ballet[1,2,3], Borja Ocón[1,2,3], Theresa Dinh[2,3], Nicole Lazarus[2,3], Manali Kun[,3], Griselda Ramos[2,3], Hiroto Kawashima[4,] Florea Lupu[5], Jonathan H Lin[1,2,3], Eugene C. Butcher[1,2,3]**# and Junliang Pan [2,3]***#.

1 Department of Pathology, Stanford University School of Medicine, Stanford, CA, USA. 2 Palo Alto Veterans Institute for Research, Palo Alto, CA, USA. 3 Veterans Affairs Palo Alto Health Care System, Palo Alto, CA, USA. 4 Graduate School of Pharmaceutical Sciences, Chiba University, Chiba, Japan 5 Oklahoma Medical Research Foundation, Oklahoma City, OK, USA.

6 Lead Contact

Co-senior Authors

*Correspondence: yub10@stanford.edu, ebutcher@stanford.edu, jpan@stanford.edu

Summary

High endothelial cells (HECs) are specialized vascular gatekeepers that control lymphocyte entry into lymph nodes, a process essential for immune surveillance and adaptive responses. However, how HECs coordinate their high biosynthetic demands with the secretory apparatus to support immune function remains unclear. Using multi-omic approaches, we identify IRE1α–XBP1–centered transcriptional networks that, together with CREB3L2–associated gene programs, coordinate inter-organelle metabolic and secretory pathways required for glycosylation and assembly of peripheral node addressin (PNAd), a sulfated sialomucin critical for lymphocyte homing. Genetic or pharmacological perturbation of these pathways disrupts HEC morphology and lymphocyte recruitment during homeostasis and impairs HEC induction during inflammation. Parallel transcriptional programs operate in mucin-producing intestinal goblet cells, suggesting common regulatory pathways linking metabolism and sulfated mucin specialization with the associated expansion of secretory organelles across immune and barrier tissues. Thus, our findings identify transcriptional programs that coordinately scale metabolic pathways and secretory organelles to support the biosynthetic infrastructure underlying the morphology and immune trafficking functions of HEVs.

Key points

XBP1-centered transcriptional networks coordinate metabolic and secretory programs for PNAd biosynthesis.

IRE1α or S1P inhibition flattens HEVs, prevents ectopic HEV formation and, reduces lymphocyte recruitment

Endothelial XBP1 deletion disrupts HEV morphology and lymphocyte trafficking.

Goblet cells share secretory transcriptional programs and regulatory logic with HEV.

Keywords

High endothelial cells, Goblet cells, Lineage specific secretory cells, Inter-organelle metabolic fluxes, Sulfated mucin type O-glycan specialization, Organellar architecture, Gene regulatory network, Lymphocyte trafficking, Mucosal immunity, Immunometabolism.

Introduction

All immune, vascular, and epithelial cells must dynamically integrate metabolic fluxes and organelle architecture to meet the biosynthetic demands of their specialized functions[1]. In the immune system, this integration is particularly critical, as cells must rapidly adapt to environmental cues during homeostasis, inflammation, and defense against pathogens. However, how metabolic pathways and organelle remodeling are logistically and transcriptionally coordinated to sustain specialized immune cell states remains incompletely understood[2-4].

High endothelial cells (HECs) exemplify this challenge. As specialized vascular gatekeepers lining high endothelial venules (HEVs) in lymph nodes (LNs), HECs control lymphocyte entry, a process essential for immune surveillance and adaptive immune responses[5,6]. Beyond lymph nodes, HEC-like cells can emerge *de novo* in chronically inflamed tissues and tumors, where they facilitate lymphocyte infiltration and tertiary lymphoid structure formation. This trafficking function is mediated by peripheral node addressin (PNAd), a set of sialomucins densely decorated with O-glycans bearing 6-sulfo sialyl Lewis X (6-sulfo sLe^X), the high-affinity ligand for L-selectin[7,8]. PNAd synthesis is metabolically intensive and depends on substantial secretory and glycosylation capacity, supported by expanded endoplasmic reticulum (ER), Golgi, and associated biosynthetic machinery that underlie the characteristic "high" morphology of HEVs and sustain the high-level, stepwise glycosylation and sulfation required for functional PNAd ligands[7,9-11]. These biosynthetic demands parallel those required for the production of sulfated mucins in intestinal goblet cells[12-14], suggesting that conserved principles link metabolism, glycosylation, secretory organelle architecture, and specialized secretory functions across immune and barrier tissues.

PNAd$^+$ HECs are induced in diverse inflammatory settings, including persistent infection, autoimmunity, graft rejection, and cancer, where their presence correlates with lymphocyte infiltration and local immune activation[15-17]. While lymphotoxin signaling and noncanonical NF-κB activation are required for HEV induction and maintenance[15,18-20], these pathways do not explain how HECs acquire their distinct cuboidal morphology and biosynthetic capacity for PNAd assembly. Thus, despite their central role in immune surveillance and inflammation, the transcriptional programs that integrate organelle and metabolic remodeling to enable HEC function remain poorly defined.

Unlike template-driven processes such as transcription or translation, PNAd biosynthesis depends on coordinated metabolic fluxes and spatial organization[7,21]. This includes sustained production of nucleotide-activated sugar donors and 3′-phosphoadenosine 5′-phosphosulfate (PAPS), their transport into the Golgi lumen, and precise deployment of glycosyltransferases and sulfotransferases across Golgi cisternae[22]. While individual enzymatic steps have been characterized, how these metabolic pathways are transcriptionally programmed and coupled to organelle expansion *in vivo* remains unknown.

These biosynthetic demands suggest that specialized transcriptional programs must coordinate metabolic pathways with secretory capacity to sustain PNAd production and the distinctive morphology of HEVs. Here, we identify conserved IRE1α–XBP1–centered gene regulatory networks (GRNs) as key components of the biosynthetic programs that sustain HEV identity and immune trafficking. Computational and biochemical analyses further implicate the bZIP transcription factor CREB3L2 as a component of the regulatory architecture associated with these programs. Using single-cell transcriptomics, genetic perturbation, and phylogenomic analyses, we show that these GRNs upregulate nutrient transporters and enzymes required for nucleotide sugar and PAPS synthesis, promote endomembrane transport of precursors into the Golgi,

coordinate spatial deployment of glycosyltransferases, and support the secretory capacity
underlying PNAd production. Disruption of these pathways impairs PNAd expression,
compromises lymphocyte homing, and prevents the ectopic emergence of functional HECs during
inflammation. Parallel transcriptional programs in goblet cells support sulfated mucin biosynthesis
and secretory specialization in barrier tissues[12 14,23-26]. Together, these findings identify IRE1α–
XBP1–centered transcriptional networks and implicate CREB3L2-associated pathways as key
components of the biosynthetic programs that sustain HEV identity and immune trafficking,
revealing a conserved regulatory logic underlying specialized immune and epithelial cell states.

Results

HECs Coordinately Express PNAd Biosynthesis, Secretory Organelle Machinery,
and UPR Programs to Support HEC Specialization

The biosynthesis of PNAd-bearing mucins, a defining feature of HECs, requires sustained
metabolic input and specialized organelle machinery, the expansion of which underlies the
characteristic 'high' endothelial morphology[27-29]. To define the metabolic programs supporting this
process, we analyzed single-cell transcriptomes of murine lymph node (LN) endothelial subsets[28]
(Suppl. Fig. 1A). Differential expression analysis revealed that HECs uniquely upregulate a suite
of genes required for PNAd biosynthesis, including nutrient transporters, metabolic enzymes for
precursor synthesis, Golgi-resident transporters, glycosyltransferases and sulfotransferases that
mediate stepwise glycan assembly (Fig. 1A, Suppl. Fig. 1B). Among the top 2,000 differentially
expressed genes (DEGs) enriched in HECs relative to other endothelial subsets, nearly all key
regulators of the inter-organelle metabolic pathways required for PNAd synthesis were
represented (Fig. 1A–C, DEG list provided in Table S1). These findings indicate that HECs
transcriptionally scale metabolic pathways to meet the biosynthetic demands of PNAd production.

These upregulated metabolic gene programs encompass every critical step of PNAd
biosynthesis, from precursor generation to glycan assembly (Fig. 1B). PNAd glycans, including
the 6-sulfo sialyl Lewis X (6-sulfo SLe^X) carbohydrate ligands for the lymphocyte homing
receptor L-selectin, are assembled within the Golgi apparatus from nucleotide sugar and sulfate
donors. HEC-enriched genes support the generation of these donors via multiple cytosolic
pathways. The hexosamine biosynthetic pathway (HBP) generates UDP-GlcNAc and UDP-
GalNAc from glucose, galactose, and glutamine through the actions of Gfpt1, Pgm3, Uap1,
and Gale, relying on nutrient uptake (via SLC2A1, SLC1A5) and acetyl-CoA derived from
mitochondrial fatty acid β-oxidation and the citrate-malate-pyruvate shuttle[21,30,31]. Additional
donors—including UDP-Gal, CMP-Neu5Ac, GDP-Fuc, and PAPS—are synthesized via the Leloir,
sialic acid, fucose and sulfate donor pathways[21,32,33], with specialized Golgi-resident transporters
(e.g., Slc35a1, Slc35b2/3, Slc35c1) directing these precursors into the Golgi lumen for glycan
assembly[34]. PNAd biosynthesis initiates at cis-Golgi with Galnt12-mediated transfer of GalNAc to
mucin core proteins such as Glycam1, CD34, Parm1, Podocalyxin, and CD24, followed by the
stepwise action of glycosyltransferases and sulfotransferases including C1galt1, Gcnt1, B3gnt3,
Fut7, Chst4, and St3gal4[7,27], along the Golgi stacks.

The 'high' endothelial morphology of HEC derives from the expansion of polyribosomes, ER,
Golgi, mitochondria, and autophagolysosomal systems, providing the architectural framework for
glyco-mucin synthesis and precursor transport between organelles. Alongside metabolic
programs, HECs upregulate gene programs required for the generation and function of organelles
that support high-output glycoprotein production (Fig. 1D–F). HEC-enriched genes include those

for ER-targeted protein translation, translocation and cotranslational N-glycosylation (Ssr1,
Sec61b, Spcs2, Rpn1, Dpagt1, Alg2, Stt3a, Stt3b, Ost4, Tmem258, Ddost)[35], ER chaperones
(Hspa5, Calr, Canx, Dnajb9, Pdia3, Pdia6)[36], and ER-associated degradation (Edem2, Derl1,
Os9, Erlec1, Vcp, Syvn1)[37]. Cargo trafficking is coordinated by COPII components (Mia3, Lman1,
Sar1b, Sec23b, Sec24d)[38] and COPI machinery (Kdelr1, Arf3, Copb2)[39], while Golgi organization
relies on scaffold proteins such as Golga2, Golgb1, and Golph3[40]. Additional HEC-enriched
programs support mitochondrial fatty acid oxidation and citrate shuttling (Mpc1, Cpt1a, Acad10,
Slc25a1, Acly)[41] as well as membrane maintenance and autophagy (Apoe, Agpat5, Lamp1,
Becn1)[42,43]. The inclusion of Slc35b1, an ATP/ADP exchanger that imports cytosolic ATP into the
ER lumen in exchange for ADP to support protein folding, further illustrates the tight coupling of
metabolic state with organelle function. Immunohistochemical datasets (Human Protein Atlas)
reveal protein-level enrichment in HECs of multiple proteins predicted by gene expression,
including proteins involved in metabolic flux (PASS2, PGM3, PMM2, CMAS), endomembrane
transport (SLC35B2, SLC35C1), PNAd biosynthesis (PARM1, GALNT1, CHST4), organelle
homeostasis (EDEM2, STX5A, KDELR2, GOLPH3), and the unfolded protein response
(SLC35B1, HSPA5) (Data S1). Together with prior studies[10,44-47], both transcriptional and
proteomic studies support a coordinated upscaling of metabolic and organellar machinery in HEC.
195
The expansive upregulation of these biosynthetic, metabolic, and organelle programs is
accompanied by selective activation of transcriptional regulators associated with the UPR. HECs
show marked enrichment of the IRE1α–XBP1 branch, including elevated expression of Hspa5,
Ern1 (IRE1α), spliced Xbp1(Fig. F). Expression of Atf6, but not Atf6b, is enriched. Notably, the
noncanonical UPR transcription factor Creb3l2, but not other Creb3 family members, is selectively
expressed in HECs, together with the S1P/S2P proteases (Mbtps1 and Mbtps2) required for
CREB3L2 activation. In contrast, the PERK-ATF4 branch is selectively suppressed in HECs, as
indicated by low Atf4 expression and elevated Dnajc3, an inhibitor of PERK signaling[48]. While
UPR signaling is classically associated with stress mitigation, these branches seem to exert
distinct metabolic outputs. IRE1α–XBP1 and CREB3L2 have been reported to drive adaptive
anabolic programs that couple nutrient sensing to ER vesicle trafficking[49,50], glycosylation[51] and
protein translation[52] while expanding ER and secretory capacity[53]. ATF6 promotes ER
proteostasis through induction of ER-associated degradation components[54,55], whereas PERK
broadly represses global translation to reduce ER load[56]. Thus, the engagement of IRE1α-XBP1
and CREB3L2 pathways in HEC, together with ATF6 activation and suppression of PERK
signaling, may establish a specialized UPR configuration for PNAd synthesis and lymphocyte
recruitment.
213
214 Taken together, these single-cell analyses reveal that PNAd biosynthesis is embedded within a
215 coordinated metabolic and regulatory network spanning multiple organelles in HECs. This
216 integrated program is predicted to coordinately scale metabolic pathways, secretory organelles,
217 and UPR signaling, thereby linking glycan biosynthesis to the distinctive morphology and
218 lymphocyte-recruiting function of the specialized "high" endothelial state.

Fig. 1 HECs Coordinately Express PNAd Biosynthesis, Secretory Organelle Machinery, and UPR Programs to Support HEC Specialization.

A. UMAP projection of scRNA-seq data from murine peripheral LN ECs[28], showing module scores that summarize the expression levels of genes controlling metabolic fluxes for PNAd synthesis in HECs compared to other EC subsets. Genes used for these modules are provided with expression values in C.

B. A schematic chart depicts the genes and flow of metabolic fluxes, starting with nutrient uptake and building block production, followed by their transport into the Golgi lumen, and culminating in the stepwise assembly of PNAd[7] on mucins within Golgi stacks. Transporters and enzymes involved in each step are color-coded in yellow and red, respectively, with those specifically involved in sulfation shown in blue.

C. Heat maps showing the enriched expression of genes in HECs that coordinate metabolic fluxes essential for PNAd biosynthesis.

D. UMAP projection of scRNA-seq data from murine peripheral LN ECs, showing module scores that summarize the expression levels of genes controlling organellar features. Mean expression profiles of the genes used for these module scores are provided in F.

E. Schematic chart illustrates the organellar protein machineries and their functions that shape the distinctive "high" cuboidal morphology of HECs.

F. Heat maps illustrate the HEC-enriched expression of genes associated with organellar structures and their functions. Values in heatmaps **C** and **F** are mean expression from three biological replicates and are shown on a color scale from min to max, with an asterisk (*) indicating the maximum value.

Goblet Cells Share Gene Programs for Sulfated Mucin Production and Organelle Expansion with HECs

Intestinal goblet cells (GCs) produce sulfated, complex carbohydrate–coated mucins that constitute a central component of the mucosal barrier, and defects in goblet cells or mucus integrity are associated with colitis[26]. Notably, Galnt12 and Chst4, highly enriched in HECs, are also selectively abundant in colonic GCs at both transcript and protein levels[26]. GALNT12 initiates mucin-type O-glycosylation and, through its lectin domain, preferentially produces heavily O-glycosylated structures like PNAd and mucus[57]. Sulfation of GC-derived mucins is catalyzed by CHST4[58,24], a sulfotransferase that is also required for PNAd biosynthesis in HEC[47]. We asked whether GCs employ similar molecular pathways to couple sulfated mucin synthesis with organelle function, by analyzing single-cell RNA sequencing data from mouse intestinal epithelial cells in the Tabula Muris atlas[59]. GCs exhibit a strikingly similar molecular profile, sharing nearly all gene sets involved in metabolic and secretory pathways supporting sulfated O-glycan production on mucins (Suppl. Fig. 2A and B). This convergence is particularly evident within precursor biosynthetic pathways and transporters. GCs also display enhanced expression of genes for ER- and Golgi-resident protein machineries that coordinate and streamline cargo trafficking along the secretory pathway (Suppl. Fig. 2C). Finally, GCs appear to engage analogous regulatory mechanisms, characterized by enriched expression of the Ern2-Xbp1 signaling axis[25,60,61], Creb3l1[24], and the intramembrane proteases Mbtps1[62]. Together, these findings suggest that GCs and HECs share conserved metabolic and secretory gene programs, secretory organelle machinery deployment, and regulatory logic to support high-output glycosylation.

XBP1 and CREB3L2 Are Predicted to Coordinate Conserved Transcriptional Networks that Couple PNAd Biosynthesis to Organelle Scaling

Having identified a shared metabolic and organellar gene program in HECs and GCs, we next sought to define the upstream regulatory networks that drive this convergence. Gene set overlap analysis across endothelial and intestinal epithelial subsets revealed that HECs and GCs share a highly correlated transcriptional signature (Fig. 2A). Among the top 2,000 DEGs enriched in HECs, 346 were co-upregulated in GCs, the majority of which encode components of metabolic pathways and organellar protein machineries (Suppl. 2B and C). This overlap suggests that a common set of *cis*- and *trans*-acting factors underlies their shared transcriptional architecture.

To identify candidate upstream transcription factors (TFs), we performed enrichment analysis on
these 346 shared genes using the Enrichr platform[63,64]. Across multiple datasets—including TF
perturbation signatures, co-expression networks, and curated TF-gene associations— the bZIP
family factors XBP1 and CREB3/CREB3L2 consistently emerged as the top regulatory
candidates (Fig. 2B, Suppl. Fig. 3A, Table S2). Independent analyses of HEC- and GC-specific
gene sets further supported a central role for XBP1 and CREB3L2 in coordinating the metabolic
and organellar programs that enable sulfated mucin biosynthesis (Suppl. Fig. 3B–C, Table S3-4).

To identify candidate cis-regulatory elements underlying this transcriptional control, we performed
de novo motif discovery using the HOMER platform on both HEC-enriched ATAC-seq peaks (Fig.
2C; Data S5) and HEC-enriched gene sets (Suppl. Fig. 3D)[65]. This analysis revealed significant
enrichment of three motif classes: a CRE-like motif ($TGAN_1TCA$) associated with CREB3L2
binding; canonical endoplasmic reticulum stress response elements (ERSE; ATTGG–N($\geq$1)–
CCACG), which mediate binding of NFYA/B and spliced XBP1 (XBP1s); and a bZIP-family
consensus motif containing a central ACGT core recognized by both XBP1s and CREB3L2[66,67],
among the most significantly enriched features. Comparative phylogenomics using Multiz
alignments from 58 eutherian mammals and PhyloP conservation scores from the UCSC Genome
Browser[68-70], further demonstrated that conserved noncoding genomic regions across multiple
gene sets harbor canonical sites associated with XBP1 and CREB3L2 activity. These include
shared "ACGT" core motifs capable of binding both TFs)[66,67], as well as factor-specific elements:
ERSE motifs that selectively bind XBP1s[71] but not CREB3L2, and CRE–like motifs ($TGAN_{1-2}TCA$)
preferentially recognized by CREB3L2[72] but not XBP1s. These elements were detected in multiple
genes required for PNAd biosynthesis in HECs, sulfated mucin production in GCs, and secretory
organelle function, as summarized in Fig. 2D.

Single-nucleotide–resolution maps of these conserved motifs, embedded within the relevant
noncoding regulatory modules, are provided in Data S2 based on aligned genomic sequences
from human, chimpanzee, monkey, rat, and mouse. Notably, XBP1- and CREB3L2-associated
motifs frequently co-occur with binding elements for additional regulatory factors, including E-box,
ETS, KLF/Sp1, and NF-κB, indicating integration with broader transcriptional architectures.
Conserved XBP1- and CREB3L2-associated motifs are also present within genes encoding cell
state–responsive regulators (Ltbr, Tnfrsf11a, Nfkb2, Relb, Nod1, Nod2, and Il13ra1) and lineage-
determining factors (Nr2f2[73] and Spdef[26]), suggesting that these bZIP-driven networks interface
with context-dependent programs to shape HEC and GC specialization.

Beyond HECs and GCs, analysis of Human Cell Atlas single-cell transcriptomic data indicate that
stomach plasma cells deploy a remarkably similar metabolic program for precursor nucleotide-
sugar biosynthesis and cell-enriched glycosyltransferases including MGAT1, MGAT2, MGAT3,
ST6GAL1 and FUT8, together with regulatory logic resembling that of HECs and GCs (Suppl.Fig.
4 and Table S6). This is consistent with prior studies showing that high glucose uptake and active
hexosamine biosynthetic pathway flux are essential for persistence of long-lived antibody-
secreting plasma cells[74,75]. A similar pattern was also evident in pituitary glandular cells, which
produce sulfoglycosylated peptide hormones through glycosyltransferases such as B4galnt3/4
and Chst8[76] (Table S6).
316
317　Together, these analyses identify key transcriptional regulators of the metabolic and secretory
318　programs of HECs. The conservation of XBP1 and CREB3L2 binding motifs across genes
319　embedded in these programs suggests that an integrated network links glycosylation capacity to
320　secretory pathway expansion through evolutionarily conserved cis-regulatory modules.
321　Consistent with this model, we identify conserved XBP1- and CREB3L2-binding sites in genes

governing specialized glycosylation in pituitary glandular cells (e.g., B4galnt3/4, Chst8) and long-lived plasma cells (Fut8, St6gal1) (Data S3). Supporting this framework, prior studies in goblet, plasma B and pituitary secretory cells demonstrate essential roles for IRE1β (ERN2)[77,78], XBP1[25,53], CREB3L1/2[24,52], and the S1P/S2P proteases (MBTPS1/2)[62,79] in establishing lineage-specific secretory pathways and high-level production of glycosylated proteins.

Fig. 2. **XBP1 and CREB3L2 Are Predicted to Link Conserved Networks Integrating PNAd Biosynthesis and Organelle Scaling.**

A. Gene set overlap of DEGs (top 2000 DEGs for individual cell subsets, computed independently for lymph node
endothelium and intestinal epithelium) revealed that HECs and intestinal GCs share a transcriptional signature enriched
for genes involved in integrating sulfated O-glycan synthesis with organellar machineries.
B. Transcription factor regulated gene set enrichment among the 346 co-upregulated genes shared between HECs
and GCs identified XBP1 and CREB3/CREB3L2 as the top candidate regulators. Enrichment was performed using the
Enrichr platform across the indicated datasets (the list of 346 shared genes is provided in Table S2). Statistical
significance is shown on the x-axis as $\sqrt{}$ ($-\log_{10} P$), compressing the dynamic range to facilitate comparisons. Dot size
represents Odds Ratio (scaled linearly from ~3.1 to 22.5), and dot color indicates $\log_2$ (Combined Score + 1). Grey
lines denote individual signatures, and the dashed vertical line indicates $P = 0.05$.
C. *De novo* motif discovery using the HOMER platform on HEC-enriched ATAC-seq peaks (compared with capillary
ECs) revealed significant enrichment of three major motif classes: a CRE-like motif (TGAN$_1$TCA) associated with
CREB3L2 binding; canonical endoplasmic reticulum stress response elements (ERSE; ATTGG–N($\geq$1)–CCACG), which
mediate binding of NFYA/B and spliced XBP1 (XBP1s); and a bZIP-family consensus motif containing a central ACGT
core recognized by both XBP1 and CREB3L2), among the most significantly enriched features. Shown is a lollipop plot
ranking the top motifs by significance ($-\log_{10}P$).
D. Genes harboring CRE–like motifs (TGAN$_{1-2}$TCA) specific for CREB3L2 are highlighted in red, those containing the
shared XBP1/CREB3L2 "ACGTG" core sequence in green, and genes with multiple conserved sites—including CRE–
like, ACGTG core, and/or ERSE elements—are shown in blue. These conserved motifs, situated within noncoding
regulatory modules, are provided at single-nucleotide resolution in Data S2.

XBP1 and CREB3L2 Activate Conserved Enhancer–Promoter Elements of PNAd
Biosynthetic Genes

Having identified conserved XBP1 and CREB3L2 binding motifs across genes involved in PNAd
biosynthesis and organelle function, we next tested whether these motifs and transcription factors
could directly regulate gene expression. We focused on glycosyltransferases and nucleotide
sugar biosynthetic enzymes localized to the Golgi and cytosol, including *Chst4* (sulfation), *Fut7*
(fucosylation), *C1galt1* (core 1 O-glycan synthesis), *Pgm3* (UDP-GalNAc production via the
HBP)[80], *Nans* (CMP-Neu5Ac synthesis)[81], *Slc35c1* (GDP-Fuc transporter)[34] and *Golph3* (a Golgi
Resident Protein that maintains the spatial organization of glycosyltransferases) [40] (Fig. 3A).

To assess direct transcriptional regulation, we generated luciferase reporter constructs driven by
enhancer and promoter regions from these genes, containing the evolutionarily conserved XBP1
and/or CREB3L2 binding sites identified in our motif analysis. Fig. 3B depicts the positions of
XBP1 and CREB3L2 binding motifs within the *Chst4* locus. These elements coincide with regions
of accessible chromatin in peripheral lymphoid, *Chst4*- and *Fut7*-expressing ECs, as indicated by
ATAC-seq, but are absent in non-expressing ITGA7$^+$ pericytes (IAPs). Electrophoretic mobility
shift assays (EMSAs) demonstrated site-specific binding of recombinant spliced XBP1 (XBP1s)
or active CREB3L2 to these motifs (Fig. 3C–D). Transient transfection–based reporter assays in
293T cells showed that overexpression of IRE1α, XBP1s, or an active form of CREB3L2
significantly transactivated the *Chst4* enhancer–promoter. In contrast, an IRE1α kinase mutant
(IRE1α K599A[82,83]), or overexpression of unspliced Xbp1 exhibited markedly reduced activity (Fig.
3E). The IRE1α K599A mutation prevents activation of the IRE1 RNase domain required for XBP1
mRNA splicing. Protein expression of the IRE1α, XBP1, and CREB3L2 constructs used in the
reporter assays was confirmed by western blot (Suppl. Fig. 5A, C–E). Wild-type IRE1α and IRE1α
K599A were expressed at comparable levels; however, the mutant induced substantially less
XBP1 mRNA splicing (Suppl. Fig. 5 B). Co-expression of XBP1s and CREB3L2 resulted in
enhanced activation, indicating cooperative control of *Chst4* transcription by these two regulatory
pathways (Fig. 3E).

The *Fut7* promoter also maps to regions of accessible chromatin in these lymphoid ECs (Fig. 3F)
and contains two conserved CREB3L2-binding sites but lacks Xbp1 motifs. This specificity was
supported by EMSA analyses showing selective CREB3L2 binding to the identified motifs (Fig.
G) and confirmed selective *Fut7* promoter activation by CREB3L2 but not XBP1s in transient

transfection-based reporter assays in 293T cells (Fig. 3H). Moreover, using bEnd.3 cells stably
integrated with a *Fut7* promoter-driven reporter (Fig. 3I), we found that CREB3L2 activation
by Brefeldin A (BFA)[84], which induces ER-Golgi fusion and CREB3L2 processing, markedly
upregulated reporter activity in bEnd3 cells. In contrast, the reporter activity was suppressed by
S1P (PF-429242) and S2P (nelfinavir) protease inhibitors[85,86], which block CREB3L2 activation;
while the IRE1α inhibitor STF-083010[87] had no effect.

Finally, XBP1s and CREB3L2 cooperatively enhanced transcription from reporter constructs
driven by the Pgm3, Nans, and C1galt1 enhancer–promoter elements, each of which contains
multiple conserved binding sites (Suppl. Fig. 6A–C). In contrast, overexpression of XBP1s and/or
CREB3L2 did not produce additive or cooperative activation of enhancer–promoter constructs
containing only a single binding site, such as the SLC35C1 and GOLPH3 reporters (Suppl. Fig.
6D–E). These findings suggest that XBP1s and CREB3L2 exert distinct, additive, or cooperative
transcriptional effects depending on motif composition.

Together, these results demonstrate that XBP1s and CREB3L2 can directly regulate transcription
of genes involved in precursor metabolism and glycan biosynthesis through conserved regulatory
motifs, extending their regulatory scope beyond canonical ER-localized targets.

Fig. 3. XBP1 and CREB3L2 Bind and Activate Conserved Regulatory Elements of PNAd Biosynthetic Genes
A. Diagram of the IRE1α–XBP1 and CREB3L2 activation pathways and their downstream transcriptional programs supporting PNAd biosynthesis.
B. Evolutionarily conserved XBP1- and CREB3L2-binding motifs within the *Chst4* locus colocalize with ATAC-accessible chromatin in Chst4[+] endothelial cells. The *Chst4* enhancer–promoter reporter construct positions the XBP1-dependent intronic enhancer upstream of the CREB3L2-binding promoter.
C–D. EMSA using fluorescently labeled Chst4 probes demonstrated that XBP1 and CREB3L2 bind to their respective motifs within the *Chst4* regulatory region. In (C), a probe containing the candidate XBP1-binding motif formed a specific DNA-protein complex with nuclear extracts from wild-type 293T cells, which was further enhanced by XBP1s overexpression. In (D), a probe encompassing the predicted CREB3L2-binding motif produced a specific complex with

nuclear extracts from CREB3L2-overexpressing cells. **Lane 1**, free probe; **Lane 2**, nuclear extracts from wild-type 293T
cells; **Lane 3**, nuclear extracts from XBP1s-overexpressing (C) or CREB3L2-overexpressing (D) cells. Binding
specificity was confirmed by competition with a 50-fold excess of unlabeled wild-type probe (Self, **Lane 4**), but not with
unlabeled mutant (**Lane 5**) or non-relevant (**Lane 6**) probes. Probe sequences are shown above each gel image.
Unbound labeled probe (**Lane 1**) migrated close to the dye front (cropped in image). Shifted protein–DNA complexes
are indicated by arrowheads.
E. IRE1, XBP1s, and CREB3L2 cooperatively activate the *Chst4* enhancer–promoter reporter in transiently transfected
293T cells. Results are shown as fold induction relative to CV (Control Vector).
F. Conserved CREB3L2-binding motifs in the *Fut7* locus map to regions of open chromatin. The positions of these
motifs are shown in the Fut7 promoter reporter construct.
G. A labeled *Fut7* probe encompassing the predicted CREB3L2-binding motif formed a specific DNA–protein complex
with nuclear extracts from CREB3L2-overexpressing 293T cells. EMSA conditions and lane assignments are as
described in (C–D); shifted protein–DNA complexes are indicated by arrowheads.
H. CREB3L2, but not IRE1α or XBP1s, selectively activates the *Fut7* promoter–driven luciferase reporter in transiently
transfected 293T cells. Results are shown as fold induction relative to CV (Control Vector).
I. CREB3L2 activation by brefeldin A (BFA) in endothelial cells increases *Fut7* reporter activity, whereas S1P/S2P
protease inhibitors PF-429242 and nelfinavir suppress CREB3L2-dependent activation. In contrast, IRE1α inhibition
(STF-083010) has no effect. Assays were performed in bEnd.3 cells stably harboring a Fut7 promoter reporter. Results
are shown as $\log_2$ (fold induction relative to control).
E, H and **I**. Data are from three experiments (independent transfections) and are shown with mean ± S.E.M. Statistical
significance was determined by unpaired two-tailed Student's *t*-test comparing each condition to control vector (CV) (E,
H) or vehicle control (I); Welch's correction was applied when variances were unequal. *: p value < 0.05; **: p value <
0.01; ***: p value < 0.001; ****: p value < 0.0001.

Pharmacological Inhibition of IRE1α- or the S1P Protease Flattens HEVs, Reduces PNAd Expression, and Impairs Lymphocyte Homing

Having established that XBP1 and CREB3L2 can directly activate PNAd biosynthetic genes
through conserved regulatory elements (Fig. 3), we next asked whether perturbation of these
pathways disrupts HEV morphology and function in vivo. To test this, we pharmacologically
perturbed IRE1α–XBP1 signaling and S1P-dependent transcription in wild-type C57BL/6 mice
using inhibitors validated in our cell culture studies. Mice were treated with STF-083010, a
selective inhibitor of IRE1α RNase activity that blocks generation of spliced Xbp1[87], or PF-429242,
an inhibitor of the Site-1 protease (S1P) required for proteolytic activation of several transcription
factors, including CREB3 family members such as CREB3L2[88].

Following treatment, peripheral lymph nodes (PLNs) were analyzed by whole-mount
immunofluorescence (IF) using anti-CD31 to label the vasculature and MECA-79 to detect the
extended core 1 epitope of PNAd[89,90]. PLNs from mice treated with STF-083010 every other day
for one week or PF-429242 daily for two weeks displayed marked structural abnormalities,
including reduced node size, flattened and elongated HEV walls, and diminished branching (Fig.
4A and D). PNAd expression was significantly reduced, as quantified by PNAd⁺ HEV surface area
relative to total vasculature (Fig. 4B and E).

To determine how rapidly HECs lose their specialized phenotype upon pathway inhibition, we
performed time-course analyses for each regimen. STF-083010 was administered every other
day for one week and lymph nodes were analyzed on Days 1, 4, and 8. Despite rapid inhibition
of Xbp1 mRNA splicing within ~24 hours *in vivo*[87], no overt changes were observed at day 1,
whereas modest reductions in PNAd expression and early morphological alterations appeared by
Day 4, followed by pronounced venular flattening and marked PNAd loss by Day 8 (Supplemental
Fig. 7A). PF-429242 was administered daily for two weeks, with analysis on Days 1, 4, 8, 12, and
15. Similarly, PF-429242 rapidly inhibits S1P protease activity within ~24 hours[85], yet minimal
changes were observed before Day 8, when flattening and loss of PNAd emerged and persisted
through Days 12–15 (Supplemental Fig. 7B). Together, inhibition of either pathway induced

progressive regression of the high endothelial phenotype, with reproducible loss of 'highness' and
PNAd loss within ~1 week, consistent with the timeframe reported following surgical occlusion of
afferent lymphatic vessels[11].
461
These defects in HEV were accompanied by a substantial decrease in total lymphocyte numbers
within the PLNs (Fig. 4C and F). To test whether lymphocyte homing was impaired, we next
assessed homing efficiency using adoptive transfer assays. Splenocytes were injected i.v. and
quantified in recipient LNs 2 hours later. Lymphocyte recruitment to PLNs (Fig.4 G, H) and
mesenteric lymph nodes (MLNs) (Suppl. Fig. 8A and C, left) were significantly reduced in both
inhibitor-treated groups, whereas lymphocyte homing to the spleen, and T cell homing to Peyer's
patches (PPs)—where HEVs are MAdCAM-1$^+$/PNAd^{-}[91]—remained unaffected (Fig.4 G, H, Suppl.
Fig. 8A and C, right). In contrast, B cell homing to PP was reduced by both inhibitors (Fig. 4G and
H, right): Although this reduction did not reach statistical significance, it may reflect effects on
IRE1α–XBP1 and S1P-dependent transcriptional programs involved in the synthesis of the B-cell
mucosal addressin (BMAd), a St6gal1-dependent glycotope that binds CD22 and supports
preferential B-cell homing to Peyer's patches [27,92]. Consistent with the role of HEV in determining
endogenous as well as injected lymphocyte accumulation, the donor-to-recipient lymphocyte
ratios in LN were comparable between treatment and control groups (Suppl. Fig. 8B, D). We also
examined the effects of IRE1α and S1P inhibition on short-term lymphocyte arrest and diapedesis
across HEVs[92]. CellTracker-labeled splenocytes were intravenously transferred into treated and
control mice, and their localization (in HEV and in the surrounding parenchyma) was assessed
30 minutes later by confocal microscopy. LNs from both STF-083010- and PF429242–treated
mice showed marked reductions in HEV-bound and extravasated lymphocytes (Fig. 4I for STF-
083010; Fig. 4J for PF429242), and a significant decrease in the ratio of extravasated to HEV
bound cells.
483
Because the S1P protease activates multiple transcription factors—including CREB3 family
members, ATF6, and SREBPs—PF-429242 inhibition may affect several regulatory pathways.
Among these, CREB3 family transcription factors have been implicated in secretory pathway
regulation, whereas ATF6 primarily promotes ER proteostasis through induction of ER-associated
degradation. Perturbation of these programs would be expected to impair secretory capacity and
glycoprotein biosynthesis, consistent with the reduced PNAd expression and HEV flattening
observed following PF-429242 treatment.
491
492 Taken together, these data indicate IRE1α–XBP1 signaling and S1P-dependent transcriptional
493 programs are required to maintain PNAd expression and function, and the characteristic "high"
494 endothelial morphology that defines HEV.
495
496

Fig. 4. Pharmacological Inhibition of IRE1α–XBP1 Signaling or S1P-Dependent Transcription Flattens HEVs and Impairs Lymphocyte Homing

A and D. Representative whole-mount immunofluorescence (IF) images of PLNs from mice treated with IRE1α inhibitor STF-083010 (**A**), Creb3l2 pathway inhibitor PF429242 (**D**) and control mice. Mice were injected intravenously (i.v.) with antibodies to label PNAd (MECA-79, cyan). Arrows denote thick HEV segments; arrowheads indicate thin, elongated HEVs. Scale bar: 500 μm.

B and E. Quantification of PNAd+ (MECA-79+) HEV surface area relative to total vasculature by *Imaris* analysis (See Star Methods). Data represent 1 field per PLN, 2–6 fields per mouse from 3 independent experiments (B: 1 control and 1 STF-083010–treated mouse per experiment; E: 1 control and 1 PF429242 treated mouse per experiment).

C and F. Lymphocytes count from 3 pooled PLNs (Inguinal, Axillary, and Branchial PLN) per STF-083010-treated (**C**), PF429242-(**F**) and control mouse.

G and H. Flow cytometry analysis of donor lymphocyte homing to PLNs and PPs 2 hours post-transfer in control vs. STF-083010– (**G**)and PF429242- (**H**) treated mice, expressed as percentage of control group mean localization ratio. **C and G:** Results are shown as mean ± SEM for n = 5 mice per group. **F and H:** Results are shown as mean ± SEM for n = 4 mice per group.

I and J. Left: Representative IF images of PLNs 30 minutes after lymphocyte injection for short-term diapedesis assay. Donor lymphocytes (CFSE-labeled, magenta), PNAd (cyan). Arrows highlight thick HEV segments with bound donor cells; arrowheads denote thin, elongated HEVs. Scale bar: 100 μm. **Right:** Percentage of donor cells localized inside MECA-79+ HEVs, quantified by *Imaris*. Data collected from 1 field/PLN, 2–3 fields/mouse across 3 independent experiments (1 control and 1 STF-083010– (**I**) or PF429242– (**J**) treated mouse per experiment).

All plotted data are shown as mean ± SEM. *: p value < 0.05; **: p value < 0.01; ***: p value < 0.001; ****: p value < 0.0001. Two-tailed unpaired t-test.

Endothelial Xbp1 Preserves HEV morphology, Supports Lymphocyte Recruitment, and Sustains Inflammatory Immune Responses

We next wished to determine whether the IRE1α–XBP1 pathway functions cell-intrinsically in endothelial cells to maintain HEV structure and function. We generated inducible, endothelial cell–specific *Xbp1* knockout mice (iEC-Xbp1-/-) by crossing *Cdh5-CreERT2* mice[93] with *Xbp1*[fl/fl] mice[94]. Cre-mediated recombination was induced by tamoxifen administration in adult mice. Tamoxifen-treated *Cdh5-CreERT2*–negative littermates served as wild-type controls. All groups were analyzed 4–10 weeks after tamoxifen treatment. Deletion of Xbp1 in ECs resulted in a significant reduction in LN size (Fig. 5A, left; Suppl. Fig. 9A). Whole-mount IF imaging revealed diminished PNAd expression (Fig. 5B), reduced HEV branching (Fig. 5A, right), and characteristic HEV flattening in iEC Xbp1-deficient mice (Fig. 5A, C; Suppl. Fig. 9B).

Lymphocyte recruitment was severely impaired in iEC_Xbp1-/- mice. Total lymphocyte numbers in peripheral LNs were reduced by ~65% compared to WT controls (Suppl. Fig. 9C), and short-term (2 hours) homing assays showed ~50% reduction in lymphocyte accumulation in both PLNs and MLNs, while control organs were unaffected (Fig. 5D; Suppl. Fig. 9D). Donor-to-recipient lymphocyte ratios remained comparable between groups (Suppl. Fig. 9E), confirming that reduced recruitment was not due to altered donor cell viability. *In vivo* diapedesis assays further revealed a ~60% decrease in lymphocyte adhesion to HEVs and an ~85% reduction in extravascular to HEV-associated lymphocyte ratios in iEC Xbp1-deficient LNs (Fig. 5E), confirming compromised lymphocyte–HEV interaction and transmigration.

To assess whether EC-specific Xbp1 depletion impacts immune function, we evaluated contact hypersensitivity responses (DTH) in iEC_Xbp1-/- and WT littermate mice following sensitization and challenge with the hapten DNFB (2,4-dinitrofluorobenzene)[47]. Xbp1 deletion significantly attenuated ear swelling and leukocyte infiltration (Fig. 5F, G). CD4+ T cells, including memory subsets (CD4+CD44^highCD45RB^low), were markedly reduced in the draining lymph nodes (auricular and inguinal) and inflamed ear tissue of iEC_Xbp1-/- mice compared to controls (Fig. 5H–I). MECA-79+ HEV-like vessels were absent from DNFB-inflamed ear skin in both genotypes (data not shown). These findings support a critical role for endothelial Xbp1 in induction of inflammatory immune responses within draining lymph nodes, although we cannot exclude the possibility that Xbp1 also influences inflamed venular structures in extra-lymphoid tissues. Together, these results demonstrate that endothelial Xbp1 is required to maintain the specialized "high" endothelial phenotype, promote PNAd expression, support HEV integrity, and enable effective lymphocyte recruitment during both homeostatic and inflammatory immune responses.

Fig.5. Endothelial Xbp1 Deletion Flattens HEVs and Reduces PNAd Expression, Lymphocyte Trafficking, and Inflammatory Immune Responses.

A. Representative vascular IF images of PLN from WT control and iEC_Xbp1-/-mice. Mice were intravenously injected with antibodies labeling PNAd (MECA-79, red) and capillaries (Ly6C, green). Representative images from five independent experiments, each with one mouse per group per experiment. Arrows show thick HEV segments, arrow heads indicate elongated thin HEV. Scale bars: 100 μm.

B. Reduced PNAd+ endothelium in EC-Xbp1 deficiency. PNAd$^+$ HEV surface area performed relative to the total vascular surface area quantified using *Imaris.*

C. Measurement of HEV diameter. Results are presented as the mean of 14-35 measurements per 10X field (~4 mm2/field). (**B-C**): Data are shown as mean ± SEM from 2-3 fields per PLN, 1-2 PLN per mouse, n=5 per group.

D. Donor lymphocyte localization in PLNs and Peyer's patches (PPs) 2 hours post-transfer in control and iEC_Xbp1-/- mice. Results are shown as a percentage of the mean localization ratio to the WT group (mean ± SEM of with 8 mice per group).

E. Left: Visualization of CFSE-labeled donor lymphocytes (white) within PLNs from control and iEC Xbp1-deficient mice, 30 minutes post transfer (i.v.). Vessels stained by i.v. injection of Mabs to PNAd (red) and capillaries (Ly6C, green). Scale bar: 100 µm. Right: Quantification of donor cell localization within PNAd$^+$ HEVs and calculation of intra- vs. extravascular lymphocyte ratios using *Imaris*. Data are presented as mean ± SEM from 1–2 lymph nodes per mouse, n=4 mice per group.

F-I: Attenuation of contact hypersensitivity responses in iEC_Xbp1-/-mice.

F. Ear swelling ratio (DNFB/vehicle) measured 24 hours after challenge. n = 8 mice per group.

G. H&E staining of ear tissue; stars indicate cartilage, arrowheads denote infiltrating leukocytes. Scale bar: 100 µm.

H-I. Flow cytometry analysis of total and memory CD4$^+$ T cells in auricular LN (ALN, H left), inguinal LN (ILN, H right) and inflamed ears (I). **H**: WT: n = 3; iEC_Xbp1-/-: n = 3. **I** WT: n = 7; iEC_Xbp1-/-: n = 8.

All plotted data are shown as mean ± SEM. Statistical analysis was performed using two-tailed unpaired t-tests. *: p value < 0.05; **: p value < 0.01; ***: p value < 0.001; ****: p value < 0.0001.

Xbp1 Cell-Intrinsically Coordinates Inter-Organelle Metabolic Programs, Organelle Integrity, and PNAd Biosynthesis in HECs

To define how Xbp1 sustains PNAd biosynthesis and organelle function *in vivo*, we performed single-cell RNA sequencing (scRNA-seq) of blood ECs isolated from lymph nodes of iEC Xbp1-deficient mice and littermate controls (Fig. 6A and B). UMAP analysis resolved nine EC populations consistent with our previously defined lymph node EC atlas[28], including capillary ECs (capECs), venous ECs (Vn), HECs, and intermediate states (CRP, TrEC) along the capillary-to-HEC differentiation trajectory[28] (Fig. 6A). As expected, *Xbp1* expression was reduced across all iEC-Xbp1-deficient EC subsets (Table S7). Compared with controls, fewer ECs were recovered from Xbp1-deficient mice, and the minor vnEC population was not represented (Fig. 6B).

To assess transcriptional impact of Xbp1 deletion, differentially expressed genes (DEGs) were identified within each EC subset and analyzed using Enrichr (Table S7-9). HECs exhibited the most pronounced transcriptional response to Xbp1 loss (Suppl. Fig. 10A). HEC-DEGs (downregulated in iEC_Xbp1−/− HECs, defining endothelial Xbp1-induced or dependent genes in HECs), were associated with XBP1-regulated gene sets and CREB3L2-linked signatures (Fig. 6C) and showed pronounced disruption of glycosylation and ER–Golgi secretory pathways, including processes related to IRE1α chaperone activity, protein processing in the ER, N- and O-linked glycosylation, oligosaccharyltransferase (OST) complex function, N-glycan trimming, the calnexin/calreticulin cycle, protein export, ER-associated degradation, and Golgi organization (Fig. 6D and E, Suppl. Fig. 11 A). Consistent with these pathway-level changes, many genes essential for PNAd biosynthesis and secretory organelle function were downregulated, including genes spanning nucleotide sugar production (*Gale, Gpi1, Pgm3, Uap1, Nans, Cmas*), metabolite transport (*Slc35a1, Slc35b3*), and glycan assembly enzymes (*Galnt12, C1galt1, C1galt1c1, Fut7, Chst4*). Key components of the secretory machinery were similarly downregulated (*Ssr1*, *Bet1*, *Derl1*, *Derl2*, *Mbtps1*, *Mbtps2* and *Golph3)*. (Fig. 6F). In contrast, expression of genes encoding core PNAd scaffolding mucins[7] was not decreased. Notably, Ern1 (IRE1α) expression was upregulated in iEC Xbp1-deficient HECs (Fig. 6F), a response previously associated with pro-apoptotic signaling and cell stress in other Xbp1 knockout models[95].

PNAd synthesis-associated pathways were generally much less affected by Xbp1 loss in other EC subsets, but Xbp1 deficiency impacted their gene expression as well (Suppl. Figs. 12-18,

Panels A; Table S8). DEG Genes downregulated by Xbp1 loss were enriched for immune and
inflammatory programs in CapIfn, arterial, and TrEC populations; lysosomal and autophagy-
associated processes in TrECs, and mitochondrial and energy homeostasis across capillary, pre-
arterial, arterial, and CRP populations. Genes upregulated in KO were enriched in adaptive
responses across EC populations, spanning stress, metabolic, and inflammatory associated
programs (Suppl. Figs. 11–18, Panels B; Table S9). Within this broader endothelial context, HECs
preferentially upregulated translational and ER stress–related pathways (Suppl. Fig.11B) and
CapIfn cells exhibited enhanced interferon signaling (Suppl. Fig.18B). Notably, canonical EC
subset identity marker genes were largely retained across all populations in iEC-Xbp1-deficient
mice (Suppl. Fig. 10B) — including pan-EC markers (*Cdh5*, *Pecam1, Erg, Ets1*), arterial and
capillary markers (*Bmx, Hey1, Dll4, Notch1, Notch4*), venous/HEC markers (*Nr2f2, Selp, Ackr1*),
and CRP markers (*Cxcr4, Apln, Pgf, Arhgap18*) — indicating that endothelial lineage identity is
preserved despite transcriptional perturbation, and that the observed gene expression changes
reflect stress and adaptive responses rather than a general collapse of EC identity.

These findings indicate that, while Xbp1 broadly supports endothelial homeostasis, HECs depend
on Xbp1-driven metabolic and secretory programs to sustain their functional specialization.

We next asked if the transcriptional changes in HEC translated into ultrastructural changes in
organelle architecture and cellular integrity. Xbp1-deficient HECs exhibited pronounced
morphological abnormalities compared to WT controls, including cellular flattening, distorted ER
membranes with ribosome dissociation, and prominent autophagic vacuoles containing
mitochondria (Fig. 6G and Suppl. 19A). Quantitative analysis confirmed significant increases in
multiple ultrastructural stress features in mutant HECs (Fig. 6H), including ER distortion,
autophagic vacuoles, mitophagy-associated structures, mitochondrial remnants, and osmiophilic
cytoplasmic regions. Adherent platelets were observed (Suppl. Fig. 19B). These morphological
abnormalities indicate severe organelle stress and compromised biosynthetic capacity, consistent
with the transcriptional changes identified by scRNA-seq.

Together, these results demonstrate that endothelial Xbp1 is essential for coordinating inter-
organelle metabolic and biosynthetic programs, preserving secretory organelle architecture, and
sustaining PNAd biosynthesis and HEC survival.

Fig. 6. Endothelial Xbp1 Coordinates Metabolic Programs and Organelle Integrity to Sustain PNAd Biosynthesis in High Endothelial Cells.
A–B. UMAP projections of lymph node blood endothelial cells colored by EC subset identity (A) and genotype (B; yellow, WT; red, iEC_Xbp1−/−), illustrating the transcriptional separation between conditions across all EC subsets.
C–E. Integrated enrichment analyses of DEGs between LN blood ECs from WT littermates and iEC_Xbp1−/− mice (the top 2,000 DEGs per EC subset). Bubble plots display enriched terms from transcription factor–related libraries (C; ARCHS4, TF perturbations), pathway databases (E; BioPlanet, CORUM, KEGG, MSigDB Hallmark, WikiPathways), and Gene Ontology (GO) terms (D; Biological Process and Cellular Component). The x-axis represents term significance (−log$_{10}$ (P value)), with the dashed line marking the nominal significance threshold (P = 0.05). Bubble size reflects the enrichment combined score (integrating effect size and significance), and bubble color indicates EC subset identity. Redundant terms within each database were collapsed to the most significant

representative. All panels share a unified x-axis and bubble size scale to allow direct comparison across datasets.
Raw Enrichr results are provided in Table S8.
F. scRNA-seq analysis of genes involved in PNAd biosynthesis and secretory organelle function in HECs from WT
and iEC-Xbp1-/- PLNs. Violin plots show normalized expression values; box plots indicate median and interquartile
range. n = 2 mice per group. Statistical testing was performed for exploratory purposes only due to limited sample
size. *p < 0.05; **p < 0.01; ***p < 0.001; ****p < 0.0001.
G. TEM of PLNs from WT and iEC-Xbp1-/- mice reveals pronounced ultrastructural abnormalities in iEC Xbp1-
deficient HECs, including cellular flattening, distorted endoplasmic reticulum (ER) membranes with ribosome
dissociation, and autophagic vacuoles containing mitochondria. N, nucleus; ER, endoplasmic reticulum. Scale bars: 2
μm, 10 μm, and 800 nm.
H. Quantification of ultrastructural abnormalities in WT and iEC-Xbp1−/− HECs from TEM micrographs, including ER
distortion, autophagic vacuoles, mitophagy-associated structures, degraded mitochondrial remnants, and osmiophilic
cytoplasmic regions. Bars represent mean ± SEM (n = 10 micrographs per genotype), with each dot corresponding to
an individual micrograph. For each genotype, 3 mice were analyzed; ≥2 tissue blocks per mouse were prepared; and
≥2 grids per block (10–12 sections per grid) were randomly sampled. Images were acquired from random fields on
each section. Statistical comparisons were performed using unpaired two-tailed t-tests with Welch's correction.

XBP1 Is Required for HEV Expansion, Ectopic PNAd[+] Vessel Induction, and Functional
Lymphocyte Homing during Inflammation.

During immune activation, HEVs undergo dynamic expansion through immune-driven
angiogenesis, facilitating enhanced lymphocyte recruitment and enlargement of draining LNs[96,97].
To determine whether Xbp1 activation is required for HEV expansion during immune stimulation,
we immunized mice locally using topical oxazolone treatment. In contrast to the expanded HEVs
observed in immunized wild-type lymph nodes (Fig. 7A-B and Suppl. Fig. 20A), iEC-specific
deletion of *Xbp1* or pharmacological inhibition with STF083010 resulted in markedly thinner and
sparsely branched PNAd[+] vessels with substantially reduced PNAd intensity. Consistent with
these changes, endogenous lymphocyte abundance and lymphocyte homing capacity were
profoundly diminished (Suppl. Fig. 20B-C). These results demonstrate a key role for Xbp1 in HEV
expansion and functional maturation during immune-driven angiogenesis. We next examined the
role of Xbp1 in the atypical lymphoid structures of omental milky spots (MSs), immune aggregates
within the peritoneal cavity that rely on induced PNAd[+] vessels for lymphocyte recruitment and
tissue organization[18,98]. Previous studies show that MS-associated HEVs are rapidly induced by
microbial stimuli such as lipopolysaccharide (LPS). Intraperitoneal injection of LPS (two injections
over 72 hours) triggered robust and rapid induction of PNAd[+] vessels in MSs, and this response
was significantly attenuated in iEC Xbp1-deficient mice (Fig. 7C–D).

HEV-like vessels also emerge *de novo* in sites of chronic inflammation[18,99] where they are
implicated in various human inflammatory diseases and cancer[100-103]. To model induction of
PNAd in immunologically driven inflammation, we assessed HEV-like vessel induction in an
oxazolone-induced model of atopic dermatitis[104]. We applied oxazolone to the ears of tamoxifen-
treated iEC_Xbp1-/-and WT mice and analyzed vessels ten days later. WT mice developed
prominent PNAd[+] HEV like vessels in inflamed ear skin, whereas iEC Xbp1-deficient mice showed
markedly reduced induction of these vessels (Fig. 7E–F).

Together, these findings establish Xbp1 as a critical regulator of PNAd[+] HEV expansion in LN and
and ectopic HEV formation in atypical lymphoid tissues and peripheral inflamed organs. By
enabling the molecular specialization of HEV, Xbp1 promotes efficient lymphocyte recruitment
and supports local tissue immune responses during inflammatory stimulation.

S1P Protease Inhibition Prevents Ectopic PNAd[+] Vessel Formation, HEV Immune
Angiogenesis, and Functional Lymphocyte Homing during Inflammation.

We next assessed the effects of the S1P protease inhibitor PF429242 in models of the MS-
induced HEV neogenesis and lymph node immune angiogenesis. S1P inhibition markedly
reduced the induction of PNAd⁺ HEVs in both MSs (Suppl. Fig. 21) and PLNs (Suppl. Fig. 22),
recapitulating key features of Xbp1 deficiency. The inhibitor reduced generation of the canonical
MECA79 glycotope (6-sulfo N-acetyllactosamine in O-linked branching structures), but even more
dramatically suppressed reactivity with antibody S2 recognizing alpha 2,3neuraminic acid
modified 6-sulfo LacNAc (6SO4 sialyl LacNAc), structures present on both O- and N-linked
glycans. Consistent with a broader disruption of HEV glycan elaboration, staining with antibody
F2, which recognizes sialyl Lewis X-containing N- and O-glycans, was also reduced in
homeostatic lymph nodes following PF429242 treatment (Suppl. Fig. 23). These suggest that
S1P-dependent transcription coordinates multiple glycosylation steps and is essential for ectopic
HEV formation[90,105].

To assess functional consequences, we performed homing assays in oxazolone-painted mice
treated with PF-429242. Both endogenous lymphocyte accumulation and recruitment of
intravenously transferred donor cells were significantly reduced in PF-treated mice (Suppl. Fig.
24), suggesting that SP1 - dependent HEV induction is essential for functional lymphocyte entry
into inflamed tissues. Although S1P protease regulates multiple transcription factors — including
CREB3 family members, ATF6, and SREBPs — and effects on other cells and pathways cannot
be excluded, these in vivo findings are consistent with a role for S1P-dependent transcriptional
programs in controlling HEV biosynthetic and functional gene networks. The specific contribution
of CREB3L2 among these substrates will require future endothelial-specific genetic studies.

730
731

Fig. 7. Endothelial Xbp1 is required for HEV expansion and Ectopic PNAd⁺ Vessel Induction during Inflammation.

A. Representative whole-mount IF images of PLNs from WT and iEC_Xbp1-/- mice, four days after OXA-induced cutaneous inflammation. Intravenous labeling with antibodies against PNAd (MECA-79, green), capillary EC (Ly6C, pink), and CD31 (red) highlights HEV morphology and surface markers. Dashed lines indicate LN boundaries; arrows denote PNAd extension along vessel segments. Scale bars, 100 µm.

B. Quantification of PNAd$^+$ HEV surface area relative to total vessel area, measured using *Imaris* analysis. Data
represent mean ± SEM from 2 fields per lymph node; 1–2 LNs per mouse (WT: n = 3; iEC_Xbp1-/-: n = 3).
C. Left: Whole-mount omental staining of WT and iEC_Xbp1-/- mice following two intraperitoneal LPS injections over
72 hours. Right: Representative milky spots showing induced PNAd$^+$ vessels (green) within CD31$^+$ vasculature (red).
Arrows highlight induction of PNAd+ HEVs. Scale bars, 100 µm.
D. Quantification of the PNAd$^+$/total vessel area ratio within milky spots, based on 10× fields (~4 mm²/field), analyzed
using *Imaris*. Data represent mean ± SEM from 3-4 fields per mouse: n = 4 mice per group.
E. Whole-mount IF of ear vasculature ten days after bilateral OXA application. Left: overview of entire ear; right: higher-
magnification views of inflamed skin showing PNAd$^+$ vessels (green) and CD31$^+$ endothelium (red). Dashed lines mark
ear boundaries; arrows indicate PNAd$^+$ HEV-like vessels. Scale bars, 100 µm.
F. Quantification of PNAd$^+$ vessel area in inflamed ear skin, analyzed using Imaris. Data are presented with mean ±
SEM from 1 field from each ear of WT (n = 5) or iEC_Xbp1-/- mice (n = 3).
B, D and F: Statistical analysis performed using unpaired two-tailed t-tests: *: p value < 0.05; **: p value < 0.01; ***: p
value < 0.001; ****: p value < 0.0001.
752

Discussion

Continuous lymphocyte recruitment into lymph nodes through HEVs puts exceptional biosynthetic demands on the vasculature, and how these demands are met at the transcriptional level has remained poorly understood. Here we show that IRE1α–XBP1–centered transcriptional networks, together with CREB3L2-associated gene programs, orchestrate the metabolic and organellar infrastructure required for PNAd biosynthesis and lymphocyte homing. These networks scale nucleotide sugar and sulfate donor synthesis, direct precursor transport across endomembrane compartments, and deploy glycosyltransferases and sulfotransferases within the Golgi — thereby coupling metabolic flux to ER and Golgi expansion and enabling the high-output glycoprotein production that underlies HEV identity.

The mechanistic centerpiece of this regulatory logic is the coupling of nucleotide sugar and PAPS biosynthesis to the spatial organization of the Golgi glycosylation machinery. Our data imply that XBP1 and CREB3L2 engage conserved cis-regulatory modules across a coordinated suite of genes spanning nutrient uptake, precursor metabolism, endomembrane transport, and glycan assembly. The frequent co-occurrence of XBP1 and CREB3L2 motifs within evolutionarily conserved regions suggests that these transcription factors may act cooperatively as well as independently, a model supported by reporter assays demonstrating both combinatorial and factor-specific regulation of PNAd biosynthetic gene promoters and enhancers. The concomitant upregulation of ER and Golgi biogenesis and function genes in HECs, and their disruption upon Xbp1 deletion or pharmacologic IRE1α or S1P inhibition, suggest that these transcriptional programs drive the scaling of secretory organelle capacity required to process the resulting glycoprotein load. Consistent with this framework, genes involved in precursor synthesis, glycosylation, and ER–Golgi trafficking are broadly represented among established direct transcriptional targets of XBP1 and CREB3L family factors[66,67,106]. This systems-level coupling is consistent with a mechanism by which the characteristic "high" morphology of HEVs and their glycosylation output are maintained as a coherent, integrated cell state rather than as independent features, though whether transcriptional coordination of these programs is sufficient to establish HEC morphology independently of other signals remains to be determined. Activation of XBP1 through IRE1α-mediated mRNA splicing and of CREB3L2 through S1P/S2P-mediated proteolysis enables these pathways to dynamically respond to increased secretory demand.

In vivo perturbation experiments at steady state further support this regulatory framework. Pharmacologic inhibition of IRE1α RNase activity and endothelial-specific deletion of Xbp1 demonstrate that the IRE1α–XBP1 axis functions as a principal regulator coordinating metabolic flux, PNAd biosynthesis, and the characteristic "high" morphology of HECs. Disruption of this pathway reduces expression of key PNAd biosynthetic genes, alters secretory organelle architecture, diminishes PNAd expression, flattens HEC, and impairs lymphocyte homing and diapedesis. Consistent with this model, pharmacologic inhibition of the site-1 protease using PF-429242 — which prevents activation of CREB3L2 — also collapses HEC and reduces PNAd expression and lymphocyte recruitment. Because S1P inhibition also blocks other S1P/S2P-dependent substrates, including ATF6, further studies will be required to define the specific contribution of CREB3L2. Nevertheless, CREB3L2 or its closely related homolog CREB3L1 has been associated with transcriptional programs engaged during increased secretory and glycosylation demand[50,52,66,107,108], and notably undergoes N-glycosylation itself, raising the possibility that its maturation and activation are coupled to the metabolic and structural requirements of glycan synthesis[109]. Through this dual control, these pathways can integrate ER adaptive capacity with metabolic and lineage-specific transcriptional programs, reinforcing glycosylation output and the scaling of secretory organelles that support PNAd production.

Beyond steady-state biosynthesis for HEC maintenance, these regulatory circuits also enable the
adaptive remodeling of secretory capacity required for HEV expansion or generation in response
to immune challenge. Pharmacologic inhibition of IRE1α or EC-specific deletion of Xbp1 impaired
HEC remodeling and lymphocyte recruitment during oxazolone-induced inflammation. EC-
specific deletion of Xbp1 also reduced ectopic induction of PNAd+ HECs in inflamed omental
lymphoid tissues, a phenotype similarly observed following treatment with the S1P inhibitor PF-
429242. These findings indicate that both the IRE1α–XBP1 and S1P-dependent transcriptional
pathways are required not only for maintenance of steady-state HEC identity but also for
inflammatory induction of specialized lymphocyte-recruiting endothelium. A parallel regulatory
framework appears to operate in goblet cells, where XBP1 and CREB3L1 support sulfated mucin
biosynthesis and expansion of secretory organelles, thereby sustaining mucosal barrier function
under basal conditions and during microbial challenge[25,50,77 26,62]. Together, these findings suggest
that coupling metabolic input to organelle architecture through shared transcriptional networks is
a conserved strategy for sustaining secretory output across distinct tissue contexts and
challenges.

More broadly, our findings in HECs support a generalizable regulatory principle in which IRE1–
XBP1-centered and CREB3 family networks coordinate metabolic and organellar programs to
enable glycosylation-dependent specialization across diverse secretory cell types. Consistent
with this framework, goblet cells and stomach-resident plasma cells engage parallel biosynthetic
programs to support high-level production of sulfated mucins and specialized N-glycosylation of
IgG, respectively (Suppl. Fig. 2 and 3), while pituitary glandular cells similarly enrich precursor-
synthesis pathways and lineage-restricted glycosyltransferases required for glycosylated peptide
hormone production[110,111] (Table S6). XBP1 is established as a key regulator in plasma[53] and
pituitary secretory cells[52], and recent studies further implicate CREB3L2 in both lineages[52,112],
suggesting a shared transcriptional architecture coupling secretory demand to lineage-specific
glycan output. This regulatory logic may extend to additional specialized secretory cells, including
chondrocytes and osteoblasts[50,113-116], further supporting its generality across diverse biosynthetic
contexts.

This regulatory architecture also appears to be evolutionarily conserved. A homologous
mechanism may operate in *Drosophila*, where CrebA — the sole CREB3 family member — and
Xbp1 together regulate secretory machinery in the salivary gland, suggesting deep evolutionary
conservation of this regulatory architecture[66,117,118]. This conserved network may represent a core
solution for scaling biosynthetic output of glycosylated macromolecules under both homeostatic
and stress-adaptive conditions. Further investigation into how these transcriptional circuits
interface with lineage-specific and signal-induced programs may yield insights into the modulation
of secretory capacity in health and disease. Additionally, these pathways may provide strategic
entry points for therapeutic glycoengineering or for manipulating vascular and epithelial
remodeling in inflammation-associated conditions.

Limitations of the study

Our study implicates IRE1α–XBP1–centered transcriptional networks in linking PNAd
biosynthesis to metabolic reprogramming and secretory organelle expansion in high endothelial
cells. Although downregulation of genes required for 6-sulfo sialyl Lewis X biosynthesis in iEC-
Xbp1–deficient HECs correlates with loss of PNAd glycotopes and "HEVness," our study does
not exclude additional effects of Xbp1 loss on mucin precursor processing, the spatial
organization of glycosyltransferases, or trafficking through the ER–Golgi system, which could also
contribute to reduced PNAd expression.

846
Despite the profound impact of endothelial Xbp1 deficiency on HEC biosynthetic capacity,
morphology, and lymphocyte recruitment, gene expression changes were also observed in other
endothelial subsets, including subsets with HEC precursor potential. Thus, contributions from
these subsets cannot be excluded. Definitive resolution of HEC-intrinsic requirements will require
HEC-restricted genetic models such as CHST4-CreERT2[119]. Similarly, we emphasize that the in
vivo functional requirement for CREB3L2 in HEV biology has not been formally demonstrated in
this study. A specific contribution of CREB3L2 is suggested by transcription factor binding site
motif enrichment in HEC-accessible chromatin and by transcriptional activation of PNAd
biosynthetic gene reporters in vitro; however, confirmation of its in vivo role will require future
endothelial-specific genetic studies.
857
858 In addition, how these transcriptional programs intersect with NF-κB signaling and are modulated
859 by cytokine cues, metabolic inputs, and organelle remodeling across distinct immunological
860 contexts remains incompletely understood. Addressing these questions will further refine our
861 understanding of how transcriptional coordination of metabolic and organellar programs shapes
862 immune cell trafficking and vascular specialization.
863

Resource availability

Lead contact
865
866 For more information and resource requests, please contact Yuhan Bi (yub10@stanford.edu),
867 who will collaborate with the Butcher Lab as needed.

Materials availability
868
869 Reagent resources are available upon request to the lead contact.

Code Availability
870
871 scRNA-seq analysis was performed using publicly available software. Code for computing
872 imputed expression values is available: https://github.com/kbrulois/magicBatch.
873
Data Availability
874
The authors declare that all data supporting the findings of this study are available within the
article and its supplementary information files. Raw and processed scRNA-seq data generated in
this study are available from the NCBI Gene Expression Omnibus repository under the accession
number GSE296739. Source data are provided with this paper. De novo motif discovery on HEC-
enriched genes is provided as a custom track in a UCSC Genome Browser session:
https://genome.ucsc.edu/s/m.xiang/HEV_genes conserved. Raw sequencing reads (FASTQ files)
for ATAC-seq performed on PLN and PP HECs, as well as other endothelial cell subsets, are
being deposited.

Acknowledgments

Supported by NIH grants R01 AI130471 and R01 AI047822, grant 1903-03787 from The Leona
M. & Harry B. Helmsley Charitable Trust to E.C.B. The Transmitted Electron Microcopy (TEM)
study was supported, in apart, by NIH S10 Award Number 1S10OD028536-01, titled "OneView
4kX4k sCMOS camera for transmission electron microscopy applications" from the Office of
Research Infrastructure Programs (ORIP). Y.B. and B.O. were supported by a Research Fellow

Award of the Crohn's and Colitis Foundation of America (835171 and 574148). M.X. was
supported by the Tobacco-Related Disease Research Program of University of California
(T31FT1867). We thank John J. Perrino for TEM sample preparation and imaging. We thank L.
Magalhaes for administrative support.

Author contributions

894 Y.B. conducted and oversaw experiments; K.B and Y.B. performed single-cell RNA-seq analysis.
895 M.X., Y.B, J.P. performed genomic analyses. Y.B. A.A., R.B., B.O., M.K., T.D. and N.L. performed
896 experiments. G.R. helped with mouse colonies. F.L. advised TEM imaging and analysis. H.K.
897 provided mAbs. J.H.L. provided Xbp1tm2Glm fl/fl mice and scientific advice. Y.B., J.P. and E.C.B.
898 conceived and supervised the study. Y.B., J.P. and E.C.B. wrote the manuscript.

Declaration of interests

900 The authors declare no competing interests.

Supplemental information

Document S1. Supplemental Figs. 1–24.

Data S1. Immunohistochemical profiling from the Human Protein Atlas validated these
transcriptomic predictions at the protein level, revealing selective enrichment in HECs of enzymes
supporting metabolic flux, nucleotide sugar transporters involved in endomembrane trafficking,
core components of the PNAd biosynthetic machinery regulators of secretory organelle
homeostasis and vesicular transport, and unfolded protein response factors, related to Fig. 1
908
909 Data S2. Base-resolution phylogenetic DNA footprinting of genes regulating metabolic flux for
910 sulfated mucin-type O-glycans and organelle machinery in HECs and GCs, related to Fig. 2D and
911 Discussion.
912
913 Data S3. Phylogenetic DNA footprinting at base resolution reveals regulatory elements associated
914 with specialized glycosylation in plasma cells and pituitary cells, related to Fig. 2D and
915 Discussion.
916
917 Table S1. Multiple excel sheets of 2000 upregulated DEGs of each subset of lymph node
918 endothelium and colon epithelium, related to Fig. 1 & 2.
919
920 Table S2-4. Integrated Enrichr TF enrichment analyses of HEC- and GC-enriched genes, related
921 to Fig. 2.
922
923 Table S5. Integrated HEC-enriched ATAC-seq peaks (compared with capECs), related to Fig. 2.
924
925 Table S6. Single-cell transcriptomic data of Stomach Resident Plasma Cells and Pituitary
926 Glandular Cells from Human Cell Atlas, related to Suppl. Fig. 4 and Discussion.
927
928 Table S7. Differential gene expression (DEG) analyses between wild-type (WT) and Xbp1-
929 deficient ECs across all EC subsets, related to Fig. 6 and Suppl. Figs. 10-18.

931 Table S8-10. Enrichr analyses performed for each EC population using DEGs from WT and Xbp1-
932 deficient ECs, related to Fig. 6 and Suppl. Figs. 10-18.

934 Table S11. Information for constructs, antibodies (Western Blot) and RT-PCR Primers.

Reference

1.   Palm, W., and Thompson, C.B. (2017). Nutrient acquisition strategies of mammalian cells.
Nature *546*, 234-242. 10.1038/nature22379.
2.   Buck, M.D., Sowell, R.T., Kaech, S.M., and Pearce, E.L. (2017). Metabolic Instruction of
Immunity. Cell *169*, 570-586. 10.1016/j.cell.2017.04.004.
3.   Chapman, N.M., and Chi, H. (2022). Metabolic adaptation of lymphocytes in immunity and
disease. Immunity *55*, 14-30. 10.1016/j.immuni.2021.12.012.
4.   Li, X., Sun, X., and Carmeliet, P. (2019). Hallmarks of Endothelial Cell Metabolism in
Health and Disease. Cell Metab *30*, 414-433. 10.1016/j.cmet.2019.08.011.
5.   von Andrian, U.H., and Mempel, T.R. (2003). Homing and cellular traffic in lymph nodes.
Nat Rev Immunol *3*, 867-878. 10.1038/nri1222.
6.   Butcher, E.C., and Picker, L.J. (1996). Lymphocyte homing and homeostasis. Science
272, 60-66. 10.1126/science.272.5258.60.
7.   Rosen, S.D. (2004). Ligands for L-selectin: homing, inflammation, and beyond. Annu Rev
Immunol *22*, 129-156. 10.1146/annurev.immunol.21.090501.080131.
8.   Streeter, P.R., Rouse, B.T., and Butcher, E.C. (1988). Immunohistologic and functional
characterization of a vascular addressin involved in lymphocyte homing into peripheral
lymph nodes. J Cell Biol *107*, 1853-1862. 10.1083/jcb.107.5.1853.
9.   Anderson, N.D., Anderson, A.O., and Wyllie, R.G. (1976). Specialized structure and
metabolic activities of high endothelial venules in rat lymphatic tissues. Immunology *31*,
455-473.
10.   Andrews, P., Milsom, D.W., and Stoddart, R.W. (1983). Glycoconjugates from high
endothelial cells. I. Partial characterization of a sulphated glycoconjugate from the high
endothelial cells of rat lymph nodes. J Cell Sci *59*, 231-244. 10.1242/jcs.59.1.231.
11.   Mebius, R.E., Streeter, P.R., Breve, J., Duijvestijn, A.M., and Kraal, G. (1991). The
influence of afferent lymphatic vessel interruption on vascular addressin expression. J Cell
Biol *115*, 85-95. 10.1083/jcb.115.1.85.
12.   Gustafsson, J.K., and Johansson, M.E.V. (2022). The role of goblet cells and mucus in
intestinal homeostasis. Nat Rev Gastroenterol Hepatol *19*, 785-803. 10.1038/s41575-022-
00675-x.
13.   Bergstrom, K.S., and Xia, L. (2013). Mucin-type O-glycans and their roles in intestinal
homeostasis. Glycobiology *23*, 1026-1037. 10.1093/glycob/cwt045.
14.   Gustafsson, J.K., and Hansson, G.C. (2025). Immune Regulation of Goblet Cell and
Mucus Functions in Health and Disease. Annu Rev Immunol *43*, 169-189.
10.1146/annurev-immunol-101721-065224.
15.   Meza-Perez, S., and Randall, T.D. (2017). Immunological Functions of the Omentum.
Trends Immunol *38*, 526-536. 10.1016/j.it.2017.03.002.
16.   Ruddle, N.H. (2023). Regulation, Maintenance, and Remodeling of High Endothelial
Venules in Homeostasis, Inflammation, and Cancer. Curr Opin Physiol *36*.
10.1016/j.cophys.2023.100705.
17.   Vella, G., Hua, Y., and Bergers, G. (2023). High endothelial venules in cancer: Regulation,
function, and therapeutic implication. Cancer Cell *41*, 527-545.
10.1016/j.ccell.2023.02.002.

18.   Blanchard, L., and Girard, J.P. (2021). High endothelial venules (HEVs) in immunity,
980   inflammation and cancer. Angiogenesis *24*, 719-753. 10.1007/s10456-021-09792-8.
981   19.   Browning, J.L., Allaire, N., Ngam-Ek, A., Notidis, E., Hunt, J., Perrin, S., and Fava, R.A.
982   (2005). Lymphotoxin-beta receptor signaling is required for the homeostatic control of HEV
983   differentiation and function. Immunity *23*, 539-550. 10.1016/j.immuni.2005.10.002.
984   20.   Sun, S.C. (2017). The non-canonical NF-kappaB pathway in immunity and inflammation.
985   Nat Rev Immunol *17*, 545-558. 10.1038/nri.2017.52.
986   21.   Scheper, A.F., Schofield, J., Bohara, R., Ritter, T., and Pandit, A. (2023). Understanding
987   glycosylation: Regulation through the metabolic flux of precursor pathways. Biotechnol
988   Adv *67*, 108184. 10.1016/j.biotechadv.2023.108184.
989   22.   Schjoldager, K.T., Narimatsu, Y., Joshi, H.J., and Clausen, H. (2020). Global view of
990   human protein glycosylation pathways and functions. Nat Rev Mol Cell Biol *21*, 729-749.
991   10.1038/s41580-020-00294-x.
992   23.   Luis, A.S., and Hansson, G.C. (2023). Intestinal mucus and their glycans: A habitat for
993   thriving microbiota. Cell Host Microbe *31*, 1087-1100. 10.1016/j.chom.2023.05.026.
994   24.   Asada, R., Saito, A., Kawasaki, N., Kanemoto, S., Iwamoto, H., Oki, M., Miyagi, H., Izumi,
995   S., and Imaizumi, K. (2012). The endoplasmic reticulum stress transducer OASIS is
996   involved in the terminal differentiation of goblet cells in the large intestine. J Biol Chem
997   *287*, 8144-8153. 10.1074/jbc.M111.332593.
998   25.   Kaser, A., Lee, A.H., Franke, A., Glickman, J.N., Zeissig, S., Tilg, H., Nieuwenhuis, E.E.,
999   Higgins, D.E., Schreiber, S., Glimcher, L.H., and Blumberg, R.S. (2008). XBP1 links ER
1000   stress to intestinal inflammation and confers genetic risk for human inflammatory bowel
1001   disease. Cell *134*, 743-756. 10.1016/j.cell.2008.07.021.
1002   26.   Nystrom, E.E.L., Martinez-Abad, B., Arike, L., Birchenough, G.M.H., Nonnecke, E.B.,
1003   Castillo, P.A., Svensson, F., Bevins, C.L., Hansson, G.C., and Johansson, M.E.V. (2021).
1004   An intercrypt subpopulation of goblet cells is essential for colonic mucus barrier function.
1005   Science *372*. 10.1126/science.abb1590.
1006   27.   Lee, M., Kiefel, H., LaJevic, M.D., Macauley, M.S., Kawashima, H., O'Hara, E., Pan, J.,
1007   Paulson, J.C., and Butcher, E.C. (2014). Transcriptional programs of lymphoid tissue
1008   capillary and high endothelium reveal control mechanisms for lymphocyte homing. Nat
1009   Immunol *15*, 982-995. 10.1038/ni.2983.
1010   28.   Brulois, K., Rajaraman, A., Szade, A., Nordling, S., Bogoslowski, A., Dermadi, D.,
1011   Rahman, M., Kiefel, H., O'Hara, E., Koning, J.J., et al. (2020). A molecular map of murine
1012   lymph node blood vascular endothelium at single cell resolution. Nat Commun *11*, 3798.
1013   10.1038/s41467-020-17291-5.
1014   29.   Veerman, K., Tardiveau, C., Martins, F., Coudert, J., and Girard, J.P. (2019). Single-Cell
1015   Analysis Reveals Heterogeneity of High Endothelial Venules and Different Regulation of
1016   Genes Controlling Lymphocyte Entry to Lymph Nodes. Cell Rep *26*, 3116-3131 e3115.
1017   10.1016/j.celrep.2019.02.042.
1018   30.   Paneque, A., Fortus, H., Zheng, J., Werlen, G., and Jacinto, E. (2023). The Hexosamine
1019   Biosynthesis Pathway: Regulation and Function. Genes (Basel) *14*.
1020   10.3390/genes14040933.
1021   31.   Pietrocola, F., Galluzzi, L., Bravo-San Pedro, J.M., Madeo, F., and Kroemer, G. (2015).
1022   Acetyl coenzyme A: a central metabolite and second messenger. Cell Metab *21*, 805-821.
1023   10.1016/j.cmet.2015.05.014.
1024   32.   Holden, H.M., Rayment, I., and Thoden, J.B. (2003). Structure and function of enzymes
1025   of the Leloir pathway for galactose metabolism. J Biol Chem *278*, 43885-43888.
1026   10.1074/jbc.R300025200.
1027   33.   Gunal, S., Hardman, R., Kopriva, S., and Mueller, J.W. (2019). Sulfation pathways from
1028   red to green. J Biol Chem *294*, 12293-12312. 10.1074/jbc.REV119.007422.

34. Hirschberg, C.B., Robbins, P.W., and Abeijon, C. (1998). Transporters of nucleotide
sugars, ATP, and nucleotide sulfate in the endoplasmic reticulum and Golgi apparatus.
Annu Rev Biochem *67*, 49-69. 10.1146/annurev.biochem.67.1.49.
35. Gemmer, M., and Forster, F. (2020). A clearer picture of the ER translocon complex. J
Cell Sci *133*. 10.1242/jcs.231340.
36. Wiseman, R.L., Mesgarzadeh, J.S., and Hendershot, L.M. (2022). Reshaping
endoplasmic reticulum quality control through the unfolded protein response. Mol Cell *82*,
1477-1491. 10.1016/j.molcel.2022.03.025.
37. Benyair, R., Ogen-Shtern, N., and Lederkremer, G.Z. (2015). Glycan regulation of ER-
associated degradation through compartmentalization. Semin Cell Dev Biol *41*, 99-109.
10.1016/j.semcdb.2014.11.006.
38. Robinson, C.M., Duggan, A., and Forrester, A. (2024). ER exit in physiology and disease.
Front Mol Biosci *11*, 1352970. 10.3389/fmolb.2024.1352970.
39. Dell'Angelica, E.C., and Bonifacino, J.S. (2019). Coatopathies: Genetic Disorders of
Protein Coats. Annu Rev Cell Dev Biol *35*, 131-168. 10.1146/annurev-cellbio-100818-
125234.
40. Muschalik, N., and Munro, S. (2018). Golgins. Curr Biol *28*, R374-R376.
10.1016/j.cub.2018.01.006.
41. Guertin, D.A., and Wellen, K.E. (2023). Acetyl-CoA metabolism in cancer. Nat Rev Cancer
23, 156-172. 10.1038/s41568-022-00543-5.
42. Debnath, J., Gammoh, N., and Ryan, K.M. (2023). Autophagy and autophagy-related
pathways in cancer. Nat Rev Mol Cell Biol *24*, 560-575. 10.1038/s41580-023-00585-z.
43. Kathryn A Jacobs, Y.H., Apostolis Panagiotis Nikolakopoulos, Swati Gupta, Gerlanda
Vella, Ana Filipa Domingues Laranjeiro, Julie Bonnereau, Mélanie Guyot, Larissa Mourao,
Feyza Cansiz, Gabriele Allies, Nicolas Van Baren, Sophie Lucas, Wiebke Sondermann,
Florian Rambow, Alpaslan Tasdogan, Colinda Scheele, Patrizia Agostinis, Gabriele
Bergers (2025). Autophagy maintains HEV identity and function during inflammation.
bioRxiv. https://doi.org/10.1101/2025.08.06.668862.
44. Freemont, A.J. (1988). Functional and biosynthetic changes in endothelial cells of vessels
in chronically inflamed tissues: evidence for endothelial control of lymphocyte entry into
diseased tissues. J Pathol *155*, 225-230. 10.1002/path.1711550308.
45. Kawashima, H., Hirakawa, J., Tobisawa, Y., Fukuda, M., and Saga, Y. (2009). Conditional
gene targeting in mouse high endothelial venules. J Immunol *182*, 5461-5468.
10.4049/jimmunol.0802327.
46. Uchimura, K., Gauguet, J.M., Singer, M.S., Tsay, D., Kannagi, R., Muramatsu, T., von
Andrian, U.H., and Rosen, S.D. (2005). A major class of L-selectin ligands is eliminated in
mice deficient in two sulfotransferases expressed in high endothelial venules. Nat
Immunol *6*, 1105-1113. 10.1038/ni1258.
47. Kawashima, H., Petryniak, B., Hiraoka, N., Mitoma, J., Huckaby, V., Nakayama, J.,
Uchimura, K., Kadomatsu, K., Muramatsu, T., Lowe, J.B., and Fukuda, M. (2005). N-
acetylglucosamine-6-O-sulfotransferases 1 and 2 cooperatively control lymphocyte
homing through L-selectin ligand biosynthesis in high endothelial venules. Nat Immunol *6*,
1096-1104. 10.1038/ni1259.
48. Tan, S.L., Gale, M.J., Jr., and Katze, M.G. (1998). Double-stranded RNA-independent
dimerization of interferon-induced protein kinase PKR and inhibition of dimerization by the
cellular P58IPK inhibitor. Mol Cell Biol *18*, 2431-2443. 10.1128/MCB.18.5.2431.
49. Liu, L., Cai, J., Wang, H., Liang, X., Zhou, Q., Ding, C., Zhu, Y., Fu, T., Guo, Q., Xu, Z., et
al. (2019). Coupling of COPII vesicle trafficking to nutrient availability by the IRE1alpha-
XBP1s axis. Proc Natl Acad Sci U S A *116*, 11776-11785. 10.1073/pnas.1814480116.
50. Melville, D.B., Montero-Balaguer, M., Levic, D.S., Bradley, K., Smith, J.R., Hatzopoulos,
A.K., and Knapik, E.W. (2011). The feelgood mutation in zebrafish dysregulates COPII-

1080   dependent secretion of select extracellular matrix proteins in skeletal morphogenesis. Dis
1081   Model Mech *4*, 763-776. 10.1242/dmm.007625.

1082   51.   Chen, K., and Shoulders, M.D. (2024). Protein Glycosylation Patterns Shaped By the
1083   IRE1-XBP1s Arm of the Unfolded Protein Response. Isr J Chem *64*.
1084   10.1002/ijch.202300162.

1085   52.   Khetchoumian, K., Balsalobre, A., Mayran, A., Christian, H., Chenard, V., St-Pierre, J.,
1086   and Drouin, J. (2019). Pituitary cell translation and secretory capacities are enhanced cell
1087   autonomously by the transcription factor Creb3l2. Nat Commun *10*, 3960.
1088   10.1038/s41467-019-11894-3.

1089   53.   Shaffer, A.L., Shapiro-Shelef, M., Iwakoshi, N.N., Lee, A.H., Qian, S.B., Zhao, H., Yu, X.,
1090   Yang, L., Tan, B.K., Rosenwald, A., et al. (2004). XBP1, downstream of Blimp-1, expands
1091   the secretory apparatus and other organelles, and increases protein synthesis in plasma
1092   cell differentiation. Immunity *21*, 81-93. 10.1016/j.immuni.2004.06.010.

1093   54.   Adachi, Y., Yamamoto, K., Okada, T., Yoshida, H., Harada, A., and Mori, K. (2008). ATF6
1094   is a transcription factor specializing in the regulation of quality control proteins in the
1095   endoplasmic reticulum. Cell Struct Funct *33*, 75-89. 10.1247/csf.07044.

1096   55.   Yamamoto, K., Sato, T., Matsui, T., Sato, M., Okada, T., Yoshida, H., Harada, A., and
1097   Mori, K. (2007). Transcriptional induction of mammalian ER quality control proteins is
1098   mediated by single or combined action of ATF6alpha and XBP1. Dev Cell *13*, 365-376.
1099   10.1016/j.devcel.2007.07.018.

1100   56.   Walter, P., and Ron, D. (2011). The unfolded protein response: from stress pathway to
1101   homeostatic regulation. Science *334*, 1081-1086. 10.1126/science.1209038.

1102   57.   Fernandez, A.J., Daniel, E.J.P., Mahajan, S.P., Gray, J.J., Gerken, T.A., Tabak, L.A., and
1103   Samara, N.L. (2019). The structure of the colorectal cancer-associated enzyme GalNAc-
1104   T12 reveals how nonconserved residues dictate its function. Proc Natl Acad Sci U S A
1105   *116*, 20404-20410. 10.1073/pnas.1902211116.

1106   58.   Abo, H., Muraki, A., Harusato, A., Imura, T., Suzuki, M., Takahashi, K., Denning, T.L., and
1107   Kawashima, H. (2023). N-acetylglucosamine-6-O sulfation on intestinal mucins prevents
1108   obesity and intestinal inflammation by regulating gut microbiota. JCI Insight *8*.
1109   10.1172/jci.insight.165944.

1110   59.   Tabula Muris, C., Overall, c., Logistical, c., Organ, c., processing, Library, p., sequencing,
1111   Computational data, a., Cell type, a., Writing, g., et al. (2018). Single-cell transcriptomics
1112   of 20 mouse organs creates a Tabula Muris. Nature *562*, 367-372. 10.1038/s41586-018-
1113   0590-4.

1114   60.   Neidhardt, L., Cloots, E., Friemel, N., Weiss, C.A.M., Harding, H.P., McLaughlin, S.H.,
1115   Janssens, S., and Ron, D. (2024). The IRE1beta-mediated unfolded protein response is
1116   repressed by the chaperone AGR2 in mucin producing cells. EMBO J *43*, 719-753.
1117   10.1038/s44318-023-00014-z.

1118   61.   Kaya, G.G., Schwarzer, R., Dannappel, M., Vlantis, K., Gobel, U., Kondylis, V.,
1119   Nedospasov, S.A., and Pasparakis, M. (2025). Unfolded protein response transcription
1120   factor XBP1 suppresses necroptosis-induced colitis by reinforcing the mucus barrier.
1121   Immunity *58*, 2208-2225 e2206. 10.1016/j.immuni.2025.07.023.

1122   62.   Brandl, K., Rutschmann, S., Li, X., Du, X., Xiao, N., Schnabl, B., Brenner, D.A., and
1123   Beutler, B. (2009). Enhanced sensitivity to DSS colitis caused by a hypomorphic Mbtps1
1124   mutation disrupting the ATF6-driven unfolded protein response. Proc Natl Acad Sci U S A
1125   *106*, 3300-3305. 10.1073/pnas.0813036106.

1126   63.   Kuleshov, M.V., Jones, M.R., Rouillard, A.D., Fernandez, N.F., Duan, Q., Wang, Z.,
1127   Koplev, S., Jenkins, S.L., Jagodnik, K.M., Lachmann, A., et al. (2016). Enrichr: a
1128   comprehensive gene set enrichment analysis web server 2016 update. Nucleic Acids Res
1129   *44*, W90-97. 10.1093/nar/gkw377.

64. Xie, Z., Bailey, A., Kuleshov, M.V., Clarke, D.J.B., Evangelista, J.E., Jenkins, S.L.,
Lachmann, A., Wojciechowicz, M.L., Kropiwnicki, E., Jagodnik, K.M., et al. (2021). Gene
Set Knowledge Discovery with Enrichr. Curr Protoc *1*, e90. 10.1002/cpz1.90.
65. Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X., Murre,
C., Singh, H., and Glass, C.K. (2010). Simple combinations of lineage-determining
transcription factors prime cis-regulatory elements required for macrophage and B cell
identities. Mol Cell *38*, 576-589. 10.1016/j.molcel.2010.05.004.
66. Fox, R.M., Hanlon, C.D., and Andrew, D.J. (2010). The CrebA/Creb3-like transcription
factors are major and direct regulators of secretory capacity. J Cell Biol *191*, 479-492.
10.1083/jcb.201004062.
67. Acosta-Alvear, D., Zhou, Y., Blais, A., Tsikitis, M., Lents, N.H., Arias, C., Lennon, C.J.,
Kluger, Y., and Dynlacht, B.D. (2007). XBP1 controls diverse cell type- and condition-
specific transcriptional regulatory networks. Mol Cell *27*, 53-66.
10.1016/j.molcel.2007.06.011.
68. Hinrichs, A.S., Karolchik, D., Baertsch, R., Barber, G.P., Bejerano, G., Clawson, H.,
Diekhans, M., Furey, T.S., Harte, R.A., Hsu, F., et al. (2006). The UCSC Genome Browser
Database: update 2006. Nucleic Acids Res *34*, D590-598. 10.1093/nar/gkj144.
69. Siepel, A., Bejerano, G., Pedersen, J.S., Hinrichs, A.S., Hou, M., Rosenbloom, K.,
Clawson, H., Spieth, J., Hillier, L.W., Richards, S., et al. (2005). Evolutionarily conserved
elements in vertebrate, insect, worm, and yeast genomes. Genome Res *15*, 1034-1050.
10.1101/gr.3715005.
70. Zoonomia, C. (2020). A comparative genomics multitool for scientific discovery and
conservation. Nature *587*, 240-245. 10.1038/s41586-020-2876-6.
71. Yamamoto, K., Suzuki, N., Wada, T., Okada, T., Yoshida, H., Kaufman, R.J., and Mori, K.
(2008). Human HRD1 promoter carries a functional unfolded protein response element to
which XBP1 but not ATF6 directly binds. J Biochem *144*, 477-486. 10.1093/jb/mvn091.
72. Vinson, C., Myakishev, M., Acharya, A., Mir, A.A., Moll, J.R., and Bonovich, M. (2002).
Classification of human B-ZIP proteins based on dimerization properties. Mol Cell Biol *22*,
6321-6335. 10.1128/MCB.22.18.6321-6335.2002.
73. You, L.R., Lin, F.J., Lee, C.T., DeMayo, F.J., Tsai, M.J., and Tsai, S.Y. (2005).
Suppression of Notch signalling by the COUP-TFII transcription factor regulates vein
identity. Nature *435*, 98-104. 10.1038/nature03511.
74. Corcoran, L.M., and Nutt, S.L. (2016). Long-Lived Plasma Cells Have a Sweet Tooth.
Immunity *45*, 3-5. 10.1016/j.immuni.2016.07.003.
75. Lam, W.Y., Becker, A.M., Kennerly, K.M., Wong, R., Curtis, J.D., Llufrio, E.M., McCommis,
K.S., Fahrmann, J., Pizzato, H.A., Nunley, R.M., et al. (2016). Mitochondrial Pyruvate
Import Promotes Long-Term Survival of Antibody-Secreting Plasma Cells. Immunity *45*,
60-73. 10.1016/j.immuni.2016.06.011.
76. Mi, Y., Fiete, D., and Baenziger, J.U. (2008). Ablation of GalNAc-4-sulfotransferase-1
enhances reproduction by altering the carbohydrate structures of luteinizing hormone in
mice. J Clin Invest *118*, 1815-1824. 10.1172/JCI32467.
77. Grey, M.J., De Luca, H., Ward, D.V., Kreulen, I.A., Bugda Gwilt, K., Foley, S.E.,
Thiagarajah, J.R., McCormick, B.A., Turner, J.R., and Lencer, W.I. (2022). The epithelial-
specific ER stress sensor ERN2/IRE1beta enables host-microbiota crosstalk to affect
colon goblet cell development. J Clin Invest *132*. 10.1172/JCI153519.
78. Tsuru, A., Fujimoto, N., Takahashi, S., Saito, M., Nakamura, D., Iwano, M., Iwawaki, T.,
Kadokura, H., Ron, D., and Kohno, K. (2013). Negative feedback by IRE1beta optimizes
mucin production in goblet cells. Proc Natl Acad Sci U S A *110*, 2864-2869.
10.1073/pnas.1212484110.

79. Caengprasath, N., Theerapanon, T., Porntaveetus, T., and Shotelersuk, V. (2021).
MBTPS2, a membrane bound protease, underlying several distinct skin and bone
disorders. J Transl Med *19*, 114. 10.1186/s12967-021-02779-5.
80. Stray-Pedersen, A., Backe, P.H., Sorte, H.S., Morkrid, L., Chokshi, N.Y., Erichsen, H.C.,
Gambin, T., Elgstoen, K.B., Bjoras, M., Wlodarski, M.W., et al. (2014). PGM3 mutations
cause a congenital disorder of glycosylation with severe immunodeficiency and skeletal
dysplasia. Am J Hum Genet *95*, 96-107. 10.1016/j.ajhg.2014.05.007.
81. van Karnebeek, C.D.M., Bonafe, L., Wen, X.Y., Tarailo-Graovac, M., Balzano, S., Royer-
Bertrand, B., Ashikov, A., Garavelli, L., Mammi, I., Turolla, L., et al. (2017). Corrigendum:
NANS-mediated synthesis of sialic acid is required for brain and skeletal development.
Nat Genet *49*, 969. 10.1038/ng0617-969a.
82. Han, D., Lerner, A.G., Vande Walle, L., Upton, J.P., Xu, W., Hagen, A., Backes, B.J.,
Oakes, S.A., and Papa, F.R. (2009). IRE1alpha kinase activation modes control alternate
endoribonuclease outputs to determine divergent cell fates. Cell *138*, 562-575.
10.1016/j.cell.2009.07.017.
83. Lipson, K.L., Ghosh, R., and Urano, F. (2008). The role of IRE1alpha in the degradation
of insulin mRNA in pancreatic beta-cells. PLoS One *3*, e1648.
10.1371/journal.pone.0001648.
84. Wood, S.A., Park, J.E., and Brown, W.J. (1991). Brefeldin A causes a microtubule-
mediated fusion of the trans-Golgi network and early endosomes. Cell *67*, 591-600.
10.1016/0092-8674(91)90533-5.
85. Hawkins, J.L., Robbins, M.D., Warren, L.C., Xia, D., Petras, S.F., Valentine, J.J.,
Varghese, A.H., Wang, I.K., Subashi, T.A., Shelly, L.D., et al. (2008). Pharmacologic
inhibition of site 1 protease activity inhibits sterol regulatory element-binding protein
processing and reduces lipogenic enzyme gene expression and lipid synthesis in cultured
cells and experimental animals. J Pharmacol Exp Ther *326*, 801-808.
10.1124/jpet.108.139626.
86. Cao, Z.J., You, J., Fan, Y.M., Yang, J.Y., Sun, J., Ma, X., Zhang, J., Li, Z., Wang, X., and
Feng, Y.X. (2025). Noncanonical UPR factor CREB3L2 drives immune evasion of triple-
negative breast cancer through Hedgehog pathway modulation in T cells. Sci Adv *11*,
eads5434. 10.1126/sciadv.ads5434.
87. Papandreou, I., Denko, N.C., Olson, M., Van Melckebeke, H., Lust, S., Tam, A., Solow-
Cordero, D.E., Bouley, D.M., Offner, F., Niwa, M., and Koong, A.C. (2011). Identification
of an Ire1alpha endonuclease specific inhibitor with cytotoxic activity against human
multiple myeloma. Blood *117*, 1311-1314. 10.1182/blood-2010-08-303099.
88. Lin, J.C., Liu, T.P., Chen, Y.B., and Yang, P.M. (2023). PF-429242 exhibits anticancer
activity in hepatocellular carcinoma cells via FOXO1-dependent autophagic cell death and
IGFBP1-dependent anti-survival signaling. Am J Cancer Res *13*, 4125-4144.
89. Yeh, J.C., Hiraoka, N., Petryniak, B., Nakayama, J., Ellies, L.G., Rabuka, D., Hindsgaul,
O., Marth, J.D., Lowe, J.B., and Fukuda, M. (2001). Novel sulfated lymphocyte homing
receptors and their control by a Core1 extension beta 1,3-N-
acetylglucosaminyltransferase. Cell *105*, 957-969. 10.1016/s0092-8674(01)00394-4.
90. Liu, W., Xiong, W., Liu, W., Hirakawa, J., and Kawashima, H. (2023). A novel monoclonal
antibody against 6-sulfo sialyl Lewis x glycans attenuates murine allergic rhinitis by
suppressing Th2 immune responses. Sci Rep *13*, 15740. 10.1038/s41598-023-43017-w.
91. Dinh, T.T., Xiang, M., Rajaraman, A., Wang, Y., Salazar, N., Zhu, Y., Roper, W., Rhee,
S., Brulois, K., O'Hara, E., et al. (2022). An NKX-COUP-TFII morphogenetic code directs
mucosal endothelial addressin expression. Nat Commun *13*, 7448. 10.1038/s41467-022-
34991-2.
92. Ballet, R., Brennan, M., Brandl, C., Feng, N., Berri, J., Cheng, J., Ocon, B., Alborzian Deh
Sheikh, A., Marki, A., Bi, Y., et al. (2021). A CD22-Shp1 phosphatase axis controls integrin

beta(7) display and B cell function in mucosal immunity. Nat Immunol *22*, 381-390.
10.1038/s41590-021-00862-z.

93.   Sorensen, I., Adams, R.H., and Gossler, A. (2009). DLL1-mediated Notch activation
regulates endothelial identity in mouse fetal arteries. Blood *113*, 5680-5688.
10.1182/blood-2008-08-174508.

94.   Hetz, C., Lee, A.H., Gonzalez-Romero, D., Thielen, P., Castilla, J., Soto, C., and Glimcher,
L.H. (2008). Unfolded protein response transcription factor XBP-1 does not influence prion
replication or pathogenesis. Proc Natl Acad Sci U S A *105*, 757-762.
10.1073/pnas.0711094105.

95.   Lee, A.H., Heidtman, K., Hotamisligil, G.S., and Glimcher, L.H. (2011). Dual and opposing
roles of the unfolded protein response regulated by IRE1alpha and XBP1 in proinsulin
processing and insulin secretion. Proc Natl Acad Sci U S A *108*, 8885-8890.
10.1073/pnas.1105564108.

96.   Kumar, V., Chyou, S., Stein, J.V., and Lu, T.T. (2012). Optical projection tomography
reveals dynamics of HEV growth after immunization with protein plus CFA and features
shared with HEVs in acute autoinflammatory lymphadenopathy. Front Immunol *3*, 282.
10.3389/fimmu.2012.00282.

97.   Tzeng, T.C., Chyou, S., Tian, S., Webster, B., Carpenter, A.C., Guaiquil, V.H., and Lu,
T.T. (2010). CD11c(hi) dendritic cells regulate the re-establishment of vascular
quiescence and stabilization after immune stimulation of lymph nodes. J Immunol *184*,
4247-4257. 10.4049/jimmunol.0902914.

98.   Rangel-Moreno, J., Moyron-Quiroz, J.E., Carragher, D.M., Kusser, K., Hartson, L.,
Moquin, A., and Randall, T.D. (2009). Omental milky spots develop in the absence of
lymphoid tissue-inducer cells and support B and T cell responses to peritoneal antigens.
Immunity *30*, 731-743. 10.1016/j.immuni.2009.03.014.

99.   Middleton, J., Americh, L., Gayon, R., Julien, D., Mansat, M., Mansat, P., Anract, P.,
Cantagrel, A., Cattan, P., Reimund, J.M., et al. (2005). A comparative study of endothelial
cell markers expressed in chronically inflamed human tissues: MECA-79, Duffy antigen
receptor for chemokines, von Willebrand factor, CD31, CD34, CD105 and CD146. J Pathol
206, 260-268. 10.1002/path.1788.

100.   Drayton, D.L., Liao, S., Mounzer, R.H., and Ruddle, N.H. (2006). Lymphoid organ
development: from ontogeny to neogenesis. Nat Immunol *7*, 344-353. 10.1038/ni1330.

101.   Habtezion, A., Nguyen, L.P., Hadeiba, H., and Butcher, E.C. (2016). Leukocyte Trafficking
to the Small Intestine and Colon. Gastroenterology *150*, 340-354.
10.1053/j.gastro.2015.10.046.

102.   Martinet, L., Garrido, I., Filleron, T., Le Guellec, S., Bellard, E., Fournie, J.J., Rochaix, P.,
and Girard, J.P. (2011). Human solid tumors contain high endothelial venules: association
with T- and B-lymphocyte infiltration and favorable prognosis in breast cancer. Cancer
Res *71*, 5678-5687. 10.1158/0008-5472.CAN-11-0431.

103.   Qian, C.N., Berghuis, B., Tsarfaty, G., Bruch, M., Kort, E.J., Ditlev, J., Tsarfaty, I., Hudson,
E., Jackson, D.G., Petillo, D., et al. (2006). Preparing the "soil": the primary tumor induces
vasculature reorganization in the sentinel lymph node before the arrival of metastatic
cancer cells. Cancer Res *66*, 10365-10376. 10.1158/0008-5472.CAN-06-2977.

104.   Choi, J., Sutaria, N., Roh, Y.S., Bordeaux, Z., Alphonse, M.P., Kwatra, S.G., and Kwatra,
M.M. (2021). Translational Relevance of Mouse Models of Atopic Dermatitis. J Clin Med
10. 10.3390/jcm10040613.

105.   Hirakawa, J., Tsuboi, K., Sato, K., Kobayashi, M., Watanabe, S., Takakura, A., Imai, Y.,
Ito, Y., Fukuda, M., and Kawashima, H. (2010). Novel anti-carbohydrate antibodies reveal
the cooperative function of sulfated N- and O-glycans in lymphocyte homing. J Biol Chem
285, 40864-40878. 10.1074/jbc.M110.167296.

106. Wang, Z.V., Deng, Y., Gao, N., Pedrozo, Z., Li, D.L., Morales, C.R., Criollo, A., Luo, X.,
Tan, W., Jiang, N., et al. (2014). Spliced X-box binding protein 1 couples the unfolded
protein response to hexosamine biosynthetic pathway. Cell *156*, 1179-1192.
10.1016/j.cell.2014.01.014.
107. Pittari, D., Dalla Torre, M., Borini, E., Hummel, B., Sawarkar, R., Semino, C., van Anken,
E., Panina-Bordignon, P., Sitia, R., and Anelli, T. (2022). CREB3L1 and CREB3L2 control
Golgi remodelling during decidualization of endometrial stromal cells. Front Cell Dev Biol
10, 986997. 10.3389/fcell.2022.986997.
108. Chan, C.P., Kok, K.H., and Jin, D.Y. (2011). CREB3 subfamily transcription factors are
not created equal: Recent insights from global analyses and animal models. Cell Biosci *1*,
6. 10.1186/2045-3701-1-6.
109. Sue, N., Thai, L.M., Saito, A., Boyer, C.K., Fordham, A.M., Yan, C., Davenport, A., Tao,
J., Bensellam, M., Cantley, J., et al. (2024). Independent activation of CREB3L2 by
glucose fills a regulatory gap in mouse beta-cells by co-ordinating insulin biosynthesis with
secretory granule formation. Mol Metab *79*, 101845. 10.1016/j.molmet.2023.101845.
110. Bousfield, G.R., and Harvey, D.J. (2019). Follicle-Stimulating Hormone Glycobiology.
Endocrinology *160*, 1515-1535. 10.1210/en.2019-00001.
111. Ulloa-Aguirre, A., Maldonado, A., Damian-Matsumura, P., and Timossi, C. (2001).
Endocrine regulation of gonadotropin glycosylation. Arch Med Res *32*, 520-532.
10.1016/s0188-4409(01)00319-8.
112. Lau, L.C., Vijay, G.K.M., Hall, D.R., and Singh, H. (2023). Non-canonical unfolded protein
response transcription factor CREB3L2 regulates survival of long-lived plasma cells.
Journal of Immunology *210*.
113. Saito, A., Hino, S., Murakami, T., Kanemoto, S., Kondo, S., Saitoh, M., Nishimura, R.,
Yoneda, T., Furuichi, T., Ikegawa, S., et al. (2009). Regulation of endoplasmic reticulum
stress response by a BBF2H7-mediated Sec23a pathway is essential for chondrogenesis.
Nat Cell Biol *11*, 1197-1204. 10.1038/ncb1962.
114. Cameron, T.L., Gresshoff, I.L., Bell, K.M., Pirog, K.A., Sampurno, L., Hartley, C.L.,
Sanford, E.M., Wilson, R., Ermann, J., Boot-Handford, R.P., et al. (2015). Cartilage-
specific ablation of XBP1 signaling in mouse results in a chondrodysplasia characterized
by reduced chondrocyte proliferation and delayed cartilage maturation and mineralization.
Osteoarthritis Cartilage *23*, 661-670. 10.1016/j.joca.2015.01.001.
115. Murakami, T., Saito, A., Hino, S., Kondo, S., Kanemoto, S., Chihara, K., Sekiya, H.,
Tsumagari, K., Ochiai, K., Yoshinaga, K., et al. (2009). Signalling mediated by the
endoplasmic reticulum stress transducer OASIS is involved in bone formation. Nat Cell
Biol *11*, 1205-1211. 10.1038/ncb1963.
116. Tvaroska, I. (2024). Glycosylation Modulates the Structure and Functions of Collagen: A
Review. Molecules *29*. 10.3390/molecules29071417.
117. Jackson, D.J., Peng, D., Shinde, S.A., Holenarasipura, A., Cahan, P., and Andrew, D.J.
(2025). CrebA regulation of secretory capacity: Genome-wide transcription profiling
coupled with in vivo DNA binding studies. bioRxiv. 10.1101/2025.06.12.659381.
118. Johnson, D.M., Wells, M.B., Fox, R., Lee, J.S., Loganathan, R., Levings, D., Bastien, A.,
Slattery, M., and Andrew, D.J. (2020). CrebA increases secretory capacity through direct
transcriptional regulation of the secretory machinery, a subset of secretory cargo, and
other key regulators. Traffic *21*, 560-577. 10.1111/tra.12753.
119. Che, Y., Yamamura, Y., Jung, S., Sabiu, G., Kondengadan, S.M., Lucia, S.E., Song, Y.,
Byun, H., Li, X., Seelam, A.J., et al. (2026). HEV-Targeted Antibody-Drug Conjugate
Promotes Long-Term Cardiac Allograft Acceptance. Circulation.
10.1161/CIRCULATIONAHA.125.076423.
1329
1330

Supplemental Figures

Supplemental Fig.1

 A

 B

Suppl. Fig. 1. (linked to Figure 1). **HECs Exhibit Enriched Expression of Metabolic and PNAd Biosynthetic Gene Programs. A.** UMAP projection of EC subsets from scRNA-seq profiling of murine peripheral lymph node ECs[26]. Cells are segregated by 11 subsets: arterial EC (pre-Art), high endothelial cells (HEC), non-HEC veins (Vn), and five capillary phenotype EC: CapEC1, CapEC2, capillary resident progenitors (CRP), transitional EC (TrEC), Interferon-stimulated gene-enriched CapEC (CapIfn). **B**. UMAP plots display module scores summarizing the expression of genes controlling building-block enzymes (left), mucins and assembly enzymes (middle), and nutrient-uptake/building-block transporters (right), all of which are significantly enriched in HECs compared with other EC subsets.

 Supplemental Fig.2

A

B <u>Metabolic Flux</u>

Supplemental Fig.2

C   Organellar Features

1349 **Suppl. Fig. 2.** (linked to Fig. 1). **Goblet Cells Share Metabolic and Secretory Gene Programs with HECs for**
1350 **Sulfated Mucin Production. A.** A schematic chart illustrates the progression of metabolic fluxes, beginning with
1351 nutrient uptake and the production of building blocks, followed by the endomembrane transport of building blocks, and
1352 culminating in the biosynthetic assembly of sulfated O-glycans within the Golgi apparatus in GCs. The fluxes are
1353 regulated by specific transporters and enzymes encoded by the genes highlighted in yellow and red, respectively, with
1354 those specifically involved in sulfation shown in blue. **B–C.** Heat maps display GC-enriched expression (bottom panels)
1355 of genes involved in metabolic pathways for sulfated mucin biosynthesis (B) and organelle-associated protein
1356 machinery (C). The expression profiles of the same genes in HECs are shown in the top panels. Mean expression
1357 values in panels B and C are derived from three independent samples and are shown on a color scale from min to max,
1358 with an asterisk (*) indicating the maximum value.
1359
1360
1361
1362
1363
1364
1365
1366
1367

Supplemental Fig.3

Suppl. Fig 3. (Linked to Figure 2). **Motif and Enrichment Analyses Identify XBP1 and CREB3L2 as Candidate**
Regulators of HEC and Goblet Cell Programs. A. Enrichr TF analysis of the up-regulated DEGs common to HEC
and GC (Table S2). **B.** Enrichr TF analysis of HEV enriched genes (Table S3). **C.** Enrichr TF analysis of intestinal GC
enriched genes (Table S4). **D.** De novo motif discovery in conserved regulatory regions flanking HEC-enriched genes
identified bZIP family binding sites, including the consensus sequence for XBP1 and CREB3L2**.** Motif analysis was
performed using HOMER on the mouse mm10 genome, with conserved regions defined by phastCons placental
mammal scores.
1423

Supplemental Fig.4

Suppl. Fig 4. (Linked to Figure 2). **Stomach-resident plasma cells are enriched for secretory and specialized IgG glycosylation gene programs.** Heatmap of selected genes across B memory cells (C0), plasma cells (C4), and germinal center (GC) B cells (C13) isolated from human stomach. Expression values were obtained from the Human Protein Atlas (HPA) single-cell RNA-seq dataset and are reported as nCPM (TMM-normalized counts per million). Values were log2-transformed (log2[nCPM + 1]) and displayed as row-wise z-scores across cell types for each gene. Genes are grouped into functional modules, including metabolic input and precursor synthesis, endomembrane transport, IgG N-glycosylation, ER protein production and quality control, vesicle trafficking and Golgi organization, and B cell identity markers. Plasma cells, in contrast to memory B cells and GC B cells, exhibit coordinated upregulation of secretory pathway regulators and glycosylation-associated genes.

Supplemental Fig.5

Suppl. Fig 5. (linked to Figure 3). **Over expression of different IRE1a, XBP1 and Creb3l2 constructs in HEK293T**
cells. A. Immunoblot analysis of IRE1 in HEK293T cells transfected with the indicated IRE1α, IRE1α K599A or control
vector (CV) (2.5 µg DNA per well, 6-well plate). Cells were harvested 48 h post-transfection. Two representative
samples are shown. **B.** RT-PCR analysis of cells transfected as in (A). Data are shown as mean ± SEM from four
independent transfections. Statistical significance was determined by unpaired two-tailed Student's t test versus control
vector (CV). *: p value < 0.05; **: p value < 0.01; ***: p value < 0.001; ****: p value < 0.0001. **C–E.** Immunoblot analysis
of XBP1s (C-D) and CREB3L2 (E) in HEK293T cells transfected with unspliced Xbp1 (C), spliced Xbp1 (D), or Creb3l2
(E). Cells were harvested 48 h post-transfection. Two representative samples are shown. RT-PCR primer sequences,
antibody and construct information are provided in Table S11.

Supplemental Fig.6

HEV enriched genes bearing multiple Xbp1 and Creb3l2 motifs

A Human PGM3-enhancer-promoter_LUC

Pgm3-enhancer-promoter-LUC

B Human NANS-enhancer-promoter_LUC

Nans-enhancer-promoter-LUC

C Human C1GALT1-enhancer-promoter_LUC

C1galt1-enhancer-promoter-LUC

HEV enriched genes bearing one Xbp1 or Creb3l2 motif

D Human SLC35C1-enhancer-promoter-LUC

Scl35c1-enhancer-promoter-LUC

E Human GOLPH3-enhancer-promoter-LUC

Golph3-enhancer-promoter-LUC

Suppl. Fig 6. (linked to Figure 3). **XBP1 and CREB3L2 Regulate PNAd Biosynthetic Gene Reporters Through Conserved Regulatory Elements. A–C**. Overexpression of Xbp1s and/or activated CREB3L2 cooperatively enhances luciferase activity driven by the Pgm3, Nans, and C1galt1 enhancer–promoter elements in 293T cells. The locations of the predicted XBP1- and CREB3L2-binding sites, together with ERSEII-like motifs, within each reporter construct are indicated in the left panels. Results are shown as fold induction relative to CV (Control Vector). **D–E.** In contrast, overexpression of Xbp1s and/or activated CREB3L2 does not produce additive or cooperative transactivation of enhancer–promoter constructs harboring a single XBP1 or CREB3L2 binding site (the SLC35C1 and GOLPH3 reporters). The locations of the predicted XBP1-/CREB3L2-binding site within each reporter construct are indicated in the left panels. Results are shown as fold induction relative to CV (Control Vector). All plotted data are from three experiments and are shown with mean ± S.E.M. Statistical significance was determined by unpaired two-tailed Student's t-test comparing each condition to control vector (CV); Welch's correction was applied when variances were unequal. *: p value < 0.05; **: p value < 0.01; ***: p value < 0.001; ****: p value < 0.0001.

Supplemental Fig.7

A

B

Suppl. Fig 7. (linked to Figure 4). **Time-course analysis of HEV regression following inhibition of IRE1α and S1P protease activity in vivo, respectively.** Representative whole-mount IF images of PLNs from mice treated with the IRE1α inhibitor STF-083010 (**A**: 3 days and 1 week) or the S1P protease inhibitor PF429242 (**B**: 3, 7, 11 days and 2 weeks), alongside untreated controls. Mice received i.v. injections of antibodies labeling PNAd (MECA-79, green). Arrowheads indicate thin, elongated HEVs. Scale bar: 100 μm.

Supplemental Fig.8

Suppl. Fig. 8. (Linked to Figure 4) **IRE1α or S1P Inhibition Reduces Lymphocyte Homing to Lymph Nodes. A, C.**
Localization of donor lymphocytes in MLN and spleen 2 hours after homing in mice treated with STF-083010 (**A**) or
PF429242 (**C**), compared to controls. Data represent the percentage of the mean localization ratio relative to control
mice. **B, D.** Donor-to-recipient cell count ratios in PLNs from mice treated with STF-083010 (B) or PF429242 (D). **A–D**:
Data are presented as mean ± SEM; n = 5 mice per group for A–B and n = 4 mice per group for C–D. Statistical
significance was assessed using unpaired two-tailed t-tests. *p < 0.05; **p < 0.01; ***p < 0.001; ****p < 0.0001.

Supplemental Fig.9

Suppl. Fig. 9. (Linked to Figure 5). **Endothelial Xbp1 Deletion Alters HEV Morphology and Lymphocyte Accumulation. A.** Representative images of LNs. Upper panel: from top to bottom—inguinal PLN, axillary PLN, and brachial PLN. Lower panel: MLN. **B**. Quantification of HEV diameter. Data represent 14–35 diameter measurements per 10× field (~4 mm²/field), averaged across 2–3 representative fields per PLN, with 1–2 PLNs analyzed per mouse (n = 5 mice per group). Each dot corresponds to an individual measurement. **C.** Total lymphocyte counts from three pooled peripheral lymph nodes (inguinal, axillary, and brachial) collected from each mouse. Each dot represents one mouse (WT: n = 6; iEC_Xbp1-/- n=5), and bars indicate the mean ± SEM. **D.** Localization of donor lymphocytes in the MLN and spleen of control and iEC Xbp1-deficient mice following a 2-hour short-term homing assay. Data are shown as the percentage of the mean localization ratio relative to the WT group. Each dot represents one mouse (n = 8 per group); bars indicate mean ± SEM. **E.** Quantification of donor lymphocytes as a percentage of total recipient lymphocytes recovered following the short-term homing assay. Each dot represents one mouse (n = 8 per group); bars indicate the mean ± SEM. **For panels C- E,** two-tailed unpaired t-tests were used for all comparisons. *p < 0.05; **p < 0.01; ***p < 0.001; ****p < 0.0001.

Supplemental Fig.10

Suppl. Fig. 10. (Link to Figure. 6). **scRNA-seq transcriptomic analysis of endothelial subsets in LNs of WT littermates and iEC_Xbp1−/− mice. A.** Transcriptional perturbation across EC subsets, calculated as Cohen's d × −log10(p value) using the top 100 DEGs per subset. **B.** Normalized mean expression of pan-EC markers (Cdh5, Pecam1, Erg, Ets1), vein markers (Nr2f2, Ackr1, Selp), artery and capillary markers (Notch1, Notch4, Dll4, Hey1, Bmx), and CRP markers (Cxcr4, Apln, Pgf, Arhgap18) across EC subsets from LNs. Each dot represents mean normalized expression. Error bars indicate ± SD. yellow, WT; red, iEC_Xbp1−/−. EC subsets: HEC, high endothelial cells; TrEC, transitional ECs; CRP, capillary-related population; CapIfn, interferon-responsive capillary ECs; CapEC1/2, capillary ECs 1 and 2; Pre-Art, pre-arterial ECs; Art, arterial ECs. n = 2 mice per group.

Supplemental Fig.11

Suppl. Fig. 11. (Link to Fig. 6). **HEC-enriched gene programs across WT and iEC_Xbp1-deficient EC populations.** Enrichr-based pathway and Gene Ontology enrichment analysis of DEGs (top 2,000) in HECs from WT and iEC_Xbp1−/− LNs, showing enriched terms derived from genes downregulated (A) and upregulated (B) in Xbp1-deficient (KO) cells relative to WT. The top 30 enriched terms ranked by specificity score (spec) are shown for visualization and are provided in Table S10. Complete Enrichr results are provided in Table S8 (downregulated gene set) and Table S9 (upregulated gene set). The x-axis represents −$\log_{10}$(P value). Dot size indicates the combined enrichment score, and dot color denotes EC subset identity.

Supplemental Fig.12

1601
1602
Suppl. Fig. 12. (Link to Fig. 6). **Art enriched gene programs across WT vs iEC Xbp1-deficient EC populations.**
Enrichr-based pathway and Gene Ontology enrichment analysis of DEGs (top 2,000) in arterial ECs from WT and
iEC_Xbp1−/− LNs, showing enriched terms derived from genes downregulated (A) and upregulated (B) in Xbp1-
deficient (KO) cells relative to WT. The top 30 enriched terms ranked by specificity score (spec) are shown for
visualization and are provided in Table S10. Complete Enrichr results are provided in Table S8 (downregulated gene
set) and Table S9 (upregulated gene set). The x-axis represents −log$_{10}$(P value). Dot size indicates the combined
enrichment score, and dot color denotes EC subset identity.
1610
1611

Supplemental Fig.13

Suppl. Fig. 13. (Link to Fig. 6). **Pre-Art enriched gene programs across WT vs iEC Xbp1-deficient EC populations.** Enrichr-based pathway and Gene Ontology enrichment analysis of DEGs (top 2,000) in pre-arterial ECs from WT and iEC_Xbp1−/− LNs, showing enriched terms derived from genes downregulated (A) and upregulated (B) in Xbp1-deficient (KO) cells relative to WT. The top 30 enriched terms ranked by specificity score (spec) are shown for visualization and are provided in Table S10. Complete Enrichr results are provided in Table S8 (downregulated gene set) and Table S9 (upregulated gene set). The x-axis represents −log$_{10}$(P value). Dot size indicates the combined enrichment score, and dot color denotes EC subset identity.

Supplemental Fig.14

1624
1625
Suppl. Fig. 14. A. (Link to Fig. 6)**. capEC1 enriched gene programs across WT vs iEC Xbp1-deficient EC**
populations. Enrichr-based pathway and Gene Ontology enrichment analysis of DEGs (top 2,000) in capEC1 from
WT and iEC_Xbp1−/− LNs, showing enriched terms derived from genes downregulated (A) and upregulated (B) in
Xbp1-deficient (KO) cells relative to WT. The top 30 enriched terms ranked by specificity score (spec) are shown for
visualization and are provided in Table S10. Complete Enrichr results are provided in Table S8 (downregulated gene
set) and Table S9 (upregulated gene set). The x-axis represents −log$_{10}$(P value). Dot size indicates the combined
enrichment score, and dot color denotes EC subset identity.
1633

Supplemental Fig.15

1635
1636
Suppl. Fig. 15. A. (Link to Fig. 6)**. capEC2 enriched gene programs across WT vs iEC Xbp1-deficient EC**
populations. Enrichr-based pathway and Gene Ontology enrichment analysis of DEGs (top 2,000) in capEC2 from
WT and iEC_Xbp1−/− LNs, showing enriched terms derived from genes downregulated (A) and upregulated (B) in
Xbp1-deficient (KO) cells relative to WT. The top 30 enriched terms ranked by specificity score (spec) are shown for
visualization and are provided in Table S10. Complete Enrichr results are provided in Table S8 (downregulated gene
set) and Table S9 (upregulated gene set). The x-axis represents −log$_{10}$(P value). Dot size indicates the combined
enrichment score, and dot color denotes EC subset identity.
1644

Supplemental Fig. 16

Suppl. Fig. 16. (Link to Fig. 6). **TrEC enriched gene programs across WT vs iEC Xbp1-deficient EC populations.** Enrichr-based pathway and Gene Ontology enrichment analysis of DEGs (top 2,000) in TrECs from WT and iEC_Xbp1−/− LNs, showing enriched terms derived from genes downregulated (A) and upregulated (B) in Xbp1-deficient (KO) cells relative to WT. The top 30 enriched terms ranked by specificity score (spec) are shown for visualization and are provided in Table S10. Complete Enrichr results are provided in Table S8 (downregulated gene set) and Table S9 (upregulated gene set). The x-axis represents −log$_{10}$(P value). Dot size indicates the combined enrichment score, and dot color denotes EC subset identity.

1656 **Supplemental Fig. 17**

1657
1658
Suppl. Fig. 17. (Link to Fig. 6). **CRP enriched gene programs across WT vs iEC Xbp1-deficient EC populations.**
Enrichr-based pathway and Gene Ontology enrichment analysis of DEGs (top 2,000) in CRPs from WT and
iEC_Xbp1−/− LNs, showing enriched terms derived from genes downregulated (A) and upregulated (B) in Xbp1-
deficient (KO) cells relative to WT. The top 30 enriched terms ranked by specificity score (spec) are shown for
visualization and are provided in Table S10. Complete Enrichr results are provided in Table S8 (downregulated gene
set) and Table S9 (upregulated gene set). The x-axis represents −log$_{10}$(P value). Dot size indicates the combined
enrichment score, and dot color denotes EC subset identity.
1666
1667
1668

Supplemental Fig. 18

Suppl. Fig. 18. (Link to Fig. 6). **Capfln enriched gene programs across WT vs iEC Xbp1-deficient EC**
populations. Enrichr-based pathway and Gene Ontology enrichment analysis of DEGs (top 2,000) in Capfln from
WT and iEC_Xbp1−/− LNs, showing enriched terms derived from genes downregulated (A) and upregulated (B) in
Xbp1-deficient (KO) cells relative to WT. The top 30 enriched terms ranked by specificity score (spec) are shown for
visualization and are provided in Table S10. Complete Enrichr results are provided in Table S8 (downregulated gene
set) and Table S9 (upregulated gene set). The x-axis represents −log$_{10}$(P value). Dot size indicates the combined
enrichment score, and dot color denotes EC subset identity.
1680

1681 **Supplemental Fig.19**

1687 **Suppl. Fig. 19.** (Linked to Fig. 6). **Ultrastructural defects in iEC Xbp1-deficient HECs. A.** iEC Xbp1-deficient HECs
1688 exhibit large degradative autophagic vacuoles and markedly distorted ER and Golgi membranes with dissociated
1689 ribosomes. **B.** Adherent platelets (arrows) are indicated. Scale bars, 5 μm.

Supplemental Fig.20

A STF-083010 i.p. 1 week; OXA 4 days

B OXA 4 days, Recipient PLN lymphocyte counts

OXA 4 days; Short-term (2 hours) homing assay

C

Suppl. Fig. 20. (Linked to Figure 7A). **Xbp1 Deficiency Reduces Lymphocyte Recruitment to Inflamed Lymph Nodes during Immune Activation. A.** Representative whole-mount staining of OXA-painted lymph nodes on Day 4 in control and STF-083010–treated mice. Mice received i.v. injections of MECA-79 (green) and CD31 (red) antibodies to visualize PNAd[+] HEVs and total vasculature, respectively. Data are representative of three independent experiments (each including one control and one STF-083010–treated mouse). **B–C.** Short-term homing of lymphocytes into OXA-painted lymph nodes on Day 4 in WT and iEC Xbp1-deficient mice. Mice were topically painted with OXA on Day 1, and homing assays were performed on Day 4. Fluorescently labeled lymphocytes were intravenously transferred, and homing was quantified after a 2-hour homing period. **B.** Total recipient lymphocyte counts from three pooled PLNs (inguinal, axillary, and brachial) collected from each mouse. Each dot represents one mouse (n = 3 per group); bars show mean ± SEM. **C.** Localization of donor lymphocytes in PLNs and spleen (control tissue) of WT and iEC Xbp1-deficient mice following the 2-hour homing assay. Data are presented as the percentage of the mean localization ratio relative to the WT group. Each dot represents one mouse (n = 3 per group); bars indicate mean ± SEM. **For panels B-C,** two-tailed unpaired t-tests were used for all comparisons. *p < 0.05; **p < 0.01.

Supplemental Fig.21

Suppl. Fig.21. (Linked to Fig. 7, S1P Protease inhibition part) **S1P Protease Inhibition Reduces Inflammation-Induced PNAd+ Vessel Formation in the Omentum. A.** Representative whole-mount staining of the omentum from control and PF429242-treated mice following two intraperitoneal injections of LPS over 72 hours. To visualize HEVs, mice were injected intravenously with antibodies labeling PNAd (MECA-79: Green; S2: Magenta) and CD31 (red). Dashed circles indicate regions containing induced HEVs. **B.** Representative images of omental milky spots showing induction of PNAd+ vessels within CD31+ vasculature (red) after PF429242 treatment. Arrows in A highlight the location of induced HEVs shown in B. Scale bar, 100 μm. Data are representative of three experiments (one control and one PF429242–treated mouse per experiment).

1757 **Supplemental Fig.22**

A

Inguinal Lymph Node PF429242 i.v. 2 weeks; LPS i.p. 72h

Suppl. Fig.22. (Linked to Fig. 7, S1P Protease inhibition part). **S1P Protease Inhibition Attenuates Remodeling of**
PNAd$^+$ Vessels in Peripheral Lymph Nodes during Inflammation. A. Representative whole-mount staining of the
PLNs from control and PF429242-treated mice following two intraperitoneal injections of LPS over 72 hours. To
visualize HEVs, mice were injected intravenously with antibodies labeling PNAd (MECA-79; green, S2; magenta) and
CD31 (red). Scale bar,100 µm. Data are representative of three experiments (one control and one PF429242–treated
mouse per experiment).

none

Supplemental Fig.23

A

Suppl. Fig. 23. (Linked to Fig. 7, S1P Protease inhibition part). **Differential PNAd glyco-epitope recognition following S1P inhibition. A.** Representative whole-mount IF images of PLNs from mice treated with the S1P inhibitor PF429242 (bottom; 2 weeks), alongside untreated controls (top). Mice were injected i.v. with antibodies labeling MECA-79 (green) and F2 (magenta). Scale bar: 1000 µm. Data are representative of three experiments (one control and one PF429242–treated mouse per experiment).

Supplemental Fig.24

 Suppl. Fig. 24. (Linked to Fig. 7, S1P Protease inhibition part). **S1P Protease Inhibition Reduces Lymphocyte**
Recruitment to Inflamed Lymph Nodes during Immune Activation. Mice were topically painted with OXA on Day 1
and draining lymph nodes were analyzed on Day 4. CFSE-labeled lymphocytes were transferred i.v., and homing was
assessed after 2 h. **A.** Total recipient lymphocyte counts from three pooled peripheral lymph nodes (inguinal, axillary,
brachial PLN) per mouse. Each dot represents one mouse (n = 4 per group); bars indicate mean ± SEM. **B.** Localization
of donor lymphocytes in PLNs and spleens of control and PF429242–treated mice after the 2-h homing assay, shown
as percentage of the mean localization ratio relative to the control group. Each dot represents one mouse (n = 3 per
group); bars indicate mean ± SEM. Two-tailed unpaired t-tests were used for all comparisons in panels A and B. *p <
0.05; **p < 0.01.

Methods

Mice

Xbp1[tm2Glm] mice[1] (provided by Dr. Jonathan Lin from Stanford University) were crossed with CDH5(PAC) Cre[ERT2] mice[2] to generate tamoxifen-inducible endothelial cell-specific Xbp1 knockout (KO) mice (Cre+ Xbp1[f/f]; iEC-Xbp1-/-). Littermates not harboring the Cre transgene were used as controls. Tamoxifen (75 mg/kg, Sigma-Aldrich, CAS # 10540-29-1) was administered via intraperitoneal (i.p.) injection four times, every other day, except for the omentum study, where tamoxifen was administered orally by gavage at the same dose. Wild-type (WT) C57BL/6J mice were obtained from the Jackson Laboratory (Bar Harbor, USA). Tamoxifen was prepared in corn oil at a concentration of 20 mg/ml by shaking overnight at 37°C in a light-protected vessel and stored at 4°C for the duration of the injections. Both male and female mice aged 6–20 weeks were used in experiments. Animals were housed under standard conditions (12 h/12 h light/dark cycle, 22 ± 2°C) with ad libitum access to food and tap water. All protocols discussed in the text and below have been approved or meet the guidelines of the accredited Department of Laboratory Animal Medicine and the Administrative Panel on Laboratory Animal Care at the VA Palo Alto Health Care System (VAPAHCS). Animals were sacrificed using approved procedures.

Preparation of lymphoid tissue BECs and flow cytometry

Axillary, inguinal, and brachial peripheral lymph nodes (PLNs) from 10–20 adult mice were dissociated as previously described[3,4]. To minimize technical variation, male and female PLN samples were combined prior to processing. Post-sequencing, the samples were computationally separated using the **AddModuleScore** function in the Seurat package (v3.1.1), leveraging the enrichment of male-specific Y-chromosomal genes and the female-specific gene *Xist*. Endothelial cells (ECs) were isolated following the procedure outlined in the **online protocol**[3]. Briefly, PLNs were pooled in HBSS buffer and minced with scissors. The tissue was washed 2–3 times with HBSS, then resuspended in HBSS containing 0.2 mg/ml collagenase P, 0.8 mg/ml Dispase II, and 0.01 mg/ml DNase. Samples were incubated at 37 °C for 10 minutes, followed by gentle disruption through repeated pipetting with P1000 pipette tips of successively smaller bore sizes. After allowing tissue fragments to settle, released cells were transferred into ice-cold FBS (final concentration, 30%). The remaining tissue was subjected to a second round of digestion with fresh buffer. Digested cells were filtered, and hematolymphoid cells were depleted using anti-CD45 MicroBeads (Miltenyi). For Figure 1, the gene set module scores were calculated using **AddModuleScore** function in the Seurat package (v3.1.1).

Isolated mouse ECs were surface-stained with monoclonal anti-mouse antibodies: APC-conjugated anti-CD31 (390; 102410; 1:4000), peridinin chlorophyll protein–cyanine 5.5–conjugated anti-CD45 (30-F11; 103131; 1:400), peridinin chlorophyll protein–cyanine 5.5–conjugated anti-Ter-119 (TER-119 TER-119; 116227; 1:400), peridinin chlorophyll protein–cyanine 5.5–conjugated anti-CD11a (H155-78; 141007; 1:400), peridinin chlorophyll protein–cyanine 5.5–conjugated anti-CD326 (G8.8; 118219; 1:400), phycoerythrin-cyanine 7-conjugated anti-Gp38 (8.1.1; 127411; 1:100), and APC/Cyanine7-conjugated anti-Ly6c (HK1.4; 128026, 1:200) (Biolegend). Anti-PNAd (MECA79) were produced in-house from hybridomas; labelled with DyLight Antibody Labeling Kits and used at a concentration of 2mg/ml (1:400).

For additional barcoding, ECs from the PLNs of iEC-Xbp1-/- and control mice (4-5 weeks post-tamoxifen treatment) were further stained with TotalSeq™-A anti-mouse Hashtag antibodies (1:200) (BioLegend). Dead cells were excluded by staining with 4',6-diamidino-2-phénylindole (DAPI) or the LIVE/DEAD Fixable Aqua dye (Invitrogen).

Approximately 5–10 × 10⁴ BECs (lin⁻Gp38⁻CD31⁺) were sorted into 100% fetal bovine serum using a FACS Aria (100 µm nozzle; ~2500 cells/second). Freshly sorted cell suspensions were diluted with PBS to a final FBS concentration of ~10%, then centrifuged at 400 × g for 5 minutes.

1864   The supernatant was carefully removed using micropipettes, and the cell pellets were
1865   resuspended in the residual volume (~30–50 µl). Cells were counted using a hemocytometer, and
1866   the concentration was adjusted to 500–1000 cells/µl by adding PBS with 10% FBS, if necessary.
1867

Single-cell RNA sequencing
Single-cell gene expression was measured using the 10x Chromium v3 platform with the
Chromium Single Cell 3' Library and Gel Bead Kit v2 (10X Genomics, PN-120237), following the
manufacturer's guidelines. Male and female cohorts were processed together and
computationally resolved post-sequencing. Libraries were sequenced on an Illumina NextSeq 500
using the 150-cycle High Output V2 Kit (Read 1: 26 bases, Read 2: 98 bases, Index 1: 8 bases).
The Cell Ranger package (v6.0.1) was used to align high-quality reads to the mm10
transcriptome. Quality control and data analysis were performed as described previously[7]. Briefly,
normalized log expression values were calculated using the scran package[8]. Imputed expression
values were generated using a customized implementation of the MAGIC algorithm[9] (Markov
Affinity-based Graph Imputation of Cells) with optimized parameters (t = 2, k = 9, ka = 3) available
at https://github.com/kbrulois/magicBatch.
Supervised cell selection was performed to exclude cells expressing non-blood endothelial gene
signatures, including lymphatic endothelial cells (Prox1, Lyve1, Pdpn), pericytes (Itga7, Pdgfrb),
fibroblastic reticular cells (Pdpn, Ccl19, Pdgfra), and lymphocytes (Ptprc, Cd52). The EC subsets
clusters were identified based on canonical marker expression and their position in tSpace
projections, yielding a total of 11 subsets[7]. Batch effects from technical replicates were corrected
using the MNN algorithm as implemented in the batchelor package (v1.0.1) via the fastMNN
function. Dimensionality reduction was performed using the UMAP algorithm. To account for cell-
cycle effects, the data were split into dividing and resting cells, and the fastMNN function was
used to align dividing cells with their resting counterparts. All subsets were represented by cells
from both male and female mice across each condition. Heatmaps were created using the
ComplexHeatmap package[15], Cell subsets were clustered based on their mean expression
profiles using ClustGeo and vegan packages. Violin plots were generated using the ggplot2
package, where y-axis units represent log-transformed normalized counts following imputation.
Differential gene expression (DEG) analyses were performed using two-sided Student's t-tests.
For comparisons between endothelial cell (EC) subsets (e.g., blood endothelial subset[4] vs
intestinal goblet cell subset[5]), as well as between wild-type (WT) and Xbp1-deficient ECs,
statistical analyses were conducted independently within each defined cell subset. Genes were
considered differentially expressed if they met the following criteria: average $\log_2$ fold change
($\log_2$FC) ≥ 0.0586 and false discovery rate (FDR) ≤ 9.94×10^{-44}. The least significant DEG
identified corresponded to a $-\log_{10}(P)$ value of 47.30. Complete DEG lists for EC subset
comparisons are provided in Table S1, and DEG results for WT versus Xbp1-deficient ECs across
individual EC subsets are provided in Table S7.
1902

1903   **Cell lines**
1904   bEnd.3 and HEK293T cells were obtained from ATCC (CRL-2299 and CRL-3216 respectively).
1905   HEK293T cells were grown in Dulbecco's modified Eagle's medium (DMEM) containing high
1906   glucose and L-pyruvate and supplemented with 10% heat-inactivated fetal bovine serum, at 37
1907   °C in 5%CO2. bEnd.3 and DN-MAML bEnd.3 cells were grown in the same media supplemented
1908   with penicillin-streptomycin[6]. Stable bEnd.3 cell transfectants overexpressing Fut7 promoter
1909   reporter were generated by transfection and selected with puromycin (1ug/ml).
1910

1911   **Plasmid constructs**
1912   *Chst4* enhancer-promoter and *Fut7* promoter fragments were PCR-amplified fragment from
1913   mouse genomic cDNA and cloned into pGL4.23[*luc2P*/Puro] and pGL4.22[*luc2P*/Puro]
1914   respectively to generate the luciferase (LUC) reporter constructs. pGL4.23[*luc2P*/Puro] and

1915   pGL4.22 [*luc2P*/Puro] vectors were obtained from Promega. *Pgm3*, *Nans*, *Galt, Scl35c1 and*
1916   *Golph3* enhancer-promoter_LUC reporter constructs were made by Vector Builder custom
1917   service. Control Renilla (Ren) Luciferase vector is from Promega.
1918   For co-transfection studies, mammalian expression vectors un-spliced Xbp1 (Xbp1), spliced Xbp1
1919   (Xbp1s), IRE1 dominant negative kinase mutant (IRE1-Mut), IRE1, Creb3l2 were commercially
1920   obtained. Constructs information is provided in Table S7.
1921

1922   **Transient transfection**
1923   For luciferase reporter assays, 1.5×10^5 HEK293T cells were plated in 24-well plates and
1924   transfected upon reaching ~70–80% confluency. Respective luciferase reporters, transcription
1925   factors, and a Renilla luciferase control vector (Promega) were co-transfected using the
1926   Lipofectamine 3000 system (Thermo Fisher Scientific) according to the manufacturer's
1927   instructions. Forty-eight hours post-transfection, cells were lysed using Lysis Reagent (Promega).
1928   Firefly and Renilla luminescence were measured using the Dual-Glo® Luciferase Assay System
1929   and read with a Turner Biosystems 20/20n luminometer. Luciferase activity was quantified as the
1930   ratio of Firefly to Renilla luminescence.
1931

1932   **Western blotting**
1933   Cells were lysed in RIPA buffer supplemented with protease inhibitors, and protein concentrations
1934   were determined by BCA assay. Equal amounts of protein were separated by SDS–PAGE and
1935   transferred onto PVDF membranes. Membranes were blocked and incubated with primary
1936   antibodies overnight at 4°C, followed by incubation with IRDye 680LT Goat anti-Rabbit IgG and
1937   IRDye 680LT Goat anti-Mouse IgG secondary antibodies (LI-COR Biosciences). Signals were
1938   detected using a LI-COR Odyssey imaging system.
1939

1940   **RT-PCR**
1941   Total RNA was isolated using the RNeasy Mini Kit (Qiagen) and reverse transcribed into cDNA
1942   using the High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems) according to the
1943   manufacturer's instructions. Quantitative PCR was performed using SYBR Green Master Mix on
1944   a real-time PCR system. Gene expression was normalized to Gapdh and calculated using the
1945 $2^{-\Delta\Delta Ct}$ method.
1946

1947   **Electrophoretic mobility shift assay (EMSA)**
1948   EMSAs were performed as previously described[6,7]. Double-stranded, fluorescence-labeled, and
1949   unlabeled probes were generated by annealing sense and antisense oligonucleotides
1950   (Sequences shown in Supplemental Fig. 4). Each binding reaction contained 1–2 pmol of labeled
1951   probe and 50 μg of nuclear extracts from 293T cells expressing recombinant spliced Xbp1 or
1952   CREB3L2. Gel shift reactions were conducted at 4 °C for 1 hour in 1.875 mg poly(d1-dC),100
1953   mM NaCl, 10 mM Tris, 1 mM DTT, 1 mM EDTA, Protease inhibitors (1:1000), 1% BSA. In
1954   reactions with cold competitors, 50-fold molar excess of ("cold") competitor unlabeled probes
1955   were added. Protein–probe complexes were separated on 5% Mini-PROTEAN® TBE gels (Bio-
1956   Rad) run in 4% TBE buffer (Bio-Rad). Gels were imaged using the Odyssey Imaging System (LI-
1957   COR).
1958

1959   **Pharmacological inhibition of IRE1α-Xbp1 and S1P Protease**
1960   For the pharmacological inhibition of IRE1α-Xbp1, WT C57BL/6J mice were injected with STF-
1961   083010 (30 mg/kg, i.p.) or DMSO, both given in 16% (vol/vol) Cremophor EL (Sigma-Aldrich)
1962   saline solution via i.p. injections every other day for four does[8]. For inhibiting active Creb3l2, WT
1963   C57BL/6J mice were administrated with PF-429242 (20 mg/kg, i.v.) or saline solution daily for two
1964   weeks[9,10].
1965

Imaging

Peripheral lymph nodes (PLNs), omentum, and ear skin were imaged following retroorbital injection of fluorescently labeled antibodies. Antibodies (10–25 µg) were administered 20–30 minutes prior to sacrifice and tissue collection. To visualize the overall vasculature, tissues were gently compressed to a thickness of approximately 35–50 µm on a glass slide. Imaging was performed using a Zeiss LSM 880 laser scanning microscope and a Nikon ECLIPSE Ti2 microscope.

Image quantification

Image analysis and parameter quantification were performed using *Imaris* 9.1. Parameters assessed included the total number of individual cell-sized particles expressing specific fluorescence signals and the volumetric surface area, representing the external surface of an isosurface. CFSE-stained cells were identified and counted using the **"Spots"** module. Volumetric measurements for **Ly6C⁺**, **MECA-79⁺**, and **CD31⁺** structures were obtained by generating 3D isosurfaces based on fluorescence detection with the **"Surfaces"** module. Total vascular surface area was measured from the **CD31⁺** isosurface. Surface statistics were generated automatically by the software and exported as CSV files for downstream quantification and statistical analysis. For analysis of cells within high endothelial venules (HEVs), a mask was applied to exclude voxels inside the region of interest (ROI), enabling quantification of cell distributions external to the surface objects using the **"Spots"** module.

Short-term homing and short-term diapedesis assays

Donor splenocytes were isolated and were labeled with 2.5 µM CFSE (Invitrogen) or CellTrace Yellow dye (Invitrogen) for 15 min at 37°C in RPMI without fetal bovine serum (FBS). A total of $50–100 \times 10^6$ donor cells were transferred into recipient mice (wild-type mice, wild-type mice treated with inhibitors, and iEC-Xbp1⁻/⁻ mice) via retroorbital injection.

For the homing assay, peripheral lymph nodes (PLNs; pooled axillary, inguinal, and brachial nodes; average of 3 PLNs per mouse), mesenteric lymph nodes (MLNs), Peyer's patches (PPs; average of 3 per mouse), and spleens were harvested 2 hours after donor cell injection. Single-cell suspensions were surface-stained for flow cytometry using the following monoclonal antibodies: PC-conjugated anti-CD4 (RM4-5; 100516; 1:200), BV421-conjugated anti-CD8β (YTS156.7.7; 126629; 1:200), PE-Cy7-conjugated anti-CD3 (145-2C11; 100320; 1:200), PE-conjugated anti-CD19 (6D5; 115508; 1:200), and FITC-conjugated anti-IgD (11.26c.2a; 11-5993-82; 1:100). Appropriate isotype controls were included. The Fc receptors were blocked using Rat Serum and anti-CD16/32 prior to the staining. Dead cells were excluded using Fixable Aqua Dead Cell Stain (Invitrogen), and counting beads were added to determine absolute cell numbers. T and B cell homing efficiencies were calculated as the ratio of recovered cells to input cells and normalized relative to the mean ratio of the wild-type control group.

For the diapedesis assay, tissues were collected 30 minutes after donor cell injection. Fluorescently labeled antibodies (10–25 µg) were administered retroorbitally 15 minutes prior to sacrifice. To visualize the overall vasculature, tissues were gently compressed to a thickness of approximately 35–50 µm on a glass slide. Imaging was performed using an LSM 880 laser scanning microscope (Zeiss) and an ECLIPSE Ti2 microscope (Nikon).

Delayed-Type Hypersensitivity (DTH) Assay

The DTH reaction in WT and iEC-Xbp1⁻/⁻ mice was induced as previously described[11]. Briefly, the DTH response was triggered through repeated epicutaneous application of 2,4-dinitrofluorobenzene (DNFB; Sigma-Aldrich). Primary immunization was achieved by applying

0.5% DNFB (50 µL in acetone: olive oil, 4:1) to bare abdominal skin on Day 1. On Day 7, mice were challenged by applying 0.5% DNFB (20 µL) to both sides of the left ear to elicit a secondary DTH response, while the right ear received 20 µL of solvent (acetone: olive oil, 4:1) as a control. 24 hours post-challenge, the thickness of both the left and right ears was measured with Ames calipers. The inflamed skin along with the draining lymph nodes (auricular and inguinal LNs) was harvested for FACS and H&E analysis.

Preparation of lymphocytes from skin and flow cytometry

Skin-infiltrating T cells were released from the connective tissue after separating the epidermal sheets as described[11-13]. T cells were surface-stained with monoclonal anti-mouse antibodies for flow cytometry, including BV421-conjugated anti-CD4 (RM4-5, 100563, 1:200); PE-Cy7-conjugated anti-CD3 (145-2C11; 100320; 1:200); BV650-conjugated anti-CD45Rb (C363-16A; 740449; 1:150); APC-conjugated anti-CD44 (IM7; 560567; 1:2000). The Fc receptors were blocked using Rat Serum and anti-CD16/32 prior to the staining. Fixable Aqua dye was applied for live/dead cell staining, and counting beads were included to calculate absolute cell numbers.

Transmission Electron Microscopy

One inguinal lymph node from each wild-type or iEC-Xbp1$^{-/-}$ mice was isolated and washed with cold PBS. The whole lymph nodes were fixed in Karnovsky's fixative: 2% Glutaraldehyde (EMS Cat# 16000) and 4% formaldehyde (EMS Cat# 15700) in 0.1M Sodium Cacodylate (EMS Cat# 12300) pH 7.4 for 1 hr. then placed on ice 24 hours. The fix was replaced with cold/aqueous 1% Osmium tetroxide (EMS Cat# 19100) and were then allowed to warm to Room Temperature (RT) for 2 hrs rotating in a hood, washed 3X with ultrafiltered water, then en bloc stained in 1% Uranyl Acetate at RT 2hrs while rotating. Samples were then dehydrated in a series of ethanol washes for 30 minutes each @ RT beginning at 50%, 70% EtOH then moved to 4°C overnight. They were place in cold 95% EtOH and allowed to warm to RT, changed to 100% 2X, then Propylene Oxide (PO) for 15 min. Samples are infiltrated with EMbed-812 resin (EMS Cat#14120) mixed 1:2, 1:1, and 2:1 with PO for 2 hrs each with leaving samples in 2:1 resin to PO overnight rotating at RT in the hood. The samples are then placed into EMbed-812 for 2 to 4 hours then placed into molds w/labels and fresh resin, orientated and placed into 65° C oven overnight.
Sections were taken around 80nm using an UC7 (Leica, Wetzlar, Germany) picked up on formvar/Carbon coated 100-mesh copper grids, stained for 40seconds in 3.5% Uranyl Acetate in 50% Acetone followed by staining in Sato's Lead Citrate for 2 minutes. Observed in the JEOL JEM-1400 120kV. Images were taken using a Gatan OneView 4k X 4k digital camera.

Electron microscopy sampling and quantification

For each genotype, 3 animals were analyzed. From each animal, ≥2 tissue blocks were prepared, and from each block, ≥2 grids (10–12 sections per grid) were randomly selected. Images were acquired from a random start field on each section at predefined magnifications. Ultrastructural features were quantified in ImageJ, including autophagic vacuoles, osmiophilic cytoplasm, and distorted ER, which were measured as percent area per micrograph, as well as mitophagy events and degraded mitochondria, which were quantified as the average number per micrograph. Data are presented as mean ± SEM, with n denoting the number of micrographs analyzed.

Lymph node and Ear immunization

Mice underwent a cutaneous immune challenge by applying 20 µL of 3–5% 4-Ethoxymethylene-2-phenyl-2-oxazolin-5-one (Sigma–Aldrich) dissolved in a 1:2 acetone: oil mixture. For lymph node analysis, the solution was applied to bare abdominal skin, while for skin analysis, it was applied to both sides of the ears. Peripheral lymph nodes (axillary, brachial, and inguinal) were collected on Day 4, and ear tissues were harvested on Day 10 following inflammation induction for imaging purposes.

HEV induction in Omentum

Six-week-old iEC-Xbp1⁻/⁻ and control mice were administered tamoxifen (75 mg/kg) orally via gavage every other day for a total of four doses. Four to five weeks after tamoxifen treatment, mice received LPS (5 mg/kg) intraperitoneally twice over a 72-hour period, followed by omentum vasculature imaging.

De novo TF motif discovery

ATAC-seq on purified HECs and Capillary ECs (CapECs) were performed essentially as previously described[14]. HEC-enriched ATAC-seq peaks were identified by differential accessibility analysis between HECs and CapECs. Peaks uniquely enriched in HECs (HEC>CapEC) were used for downstream motif analysis.

Evolutionarily conserved regulatory regions associated with HEV-enriched genes were defined as the gene body ±5 kb and intersected with conserved genomic elements scoring >300 in the UCSC Genome Browser phastCons Placental Elements track, which aligns 60 vertebrate genomes and reports nucleotide-level conservation[15,16]. All conserved regions of HEV genes are provided as a custom track in a UCSC Genome Browser session: https://genome.ucsc.edu/s/m.xiang/HEV_genes_conserved.

De novo transcription factor (TF) motif discovery was performed on HEC-enriched ATAC-seq peaks as well as on conserved regulatory regions of HEV-enriched genes in the mouse *mm10* genome using HOMER[17], applying default background models and standard significance thresholds.

TF regulation analysis

List of HEC and GC co-upregulated DEGs, HEC- and GC- enriched genes were analyzed to identify candidate upstream TF regulators using Enrichr[18-20]. Results of TF regulators from select databases are shown. Detailed analysis, including the gene lists used, is provided in Data 2-4.

Comparative Genomics and Phylogenomic Analysis

Comparative genomics and phylogenetic footprinting were performed using the UCSC Genome Browser to identify evolutionarily constrained regions across candidate genes, including promoter regions, 5′ and 3′ untranslated regions (UTRs), and intronic sequences. Analyses were based on the 58 eutherian mammals Multiz Alignment and PhyloP Basewise Conservation tracks to evaluate sequence conservation. To identify putative transcriptional regulatory elements, we utilized the TFBS Conserved track and the JASPAR TFBS database, focusing on conserved binding motifs for XBP1 and CREB3L2 within the identified regulatory modules.

Data and statistical analysis

Results are expressed as mean ± SEM and individual data points represent biological replicates. Statistics were calculated using GraphPad Prism, with differences among means tested for statistical significance as indicated in the corresponding figure legend. Statistical significance is indicated by the following: $*P \leq 0.05$, $**P \leq 0.01$, $***P \leq 0.001$ and $****P \leq 0.0001$. No statistical significance is denoted as 'na'.

Reference:

1.   Hetz, C., Lee, A.H., Gonzalez-Romero, D., Thielen, P., Castilla, J., Soto, C., and Glimcher, L.H. (2008). Unfolded protein response transcription factor XBP-1 does not influence prion replication or pathogenesis. Proc Natl Acad Sci U S A *105*, 757-762. 10.1073/pnas.0711094105.

2. Sorensen, I., Adams, R.H., and Gossler, A. (2009). DLL1-mediated Notch activation
regulates endothelial identity in mouse fetal arteries. Blood *113*, 5680-5688.
10.1182/blood-2008-08-174508.
3. Lee, M., Kiefel, H., LaJevic, M.D., Macauley, M.S., Kawashima, H., O'Hara, E., Pan, J.,
Paulson, J.C., and Butcher, E.C. (2014). Transcriptional programs of lymphoid tissue
capillary and high endothelium reveal control mechanisms for lymphocyte homing. Nat
Immunol *15*, 982-995. 10.1038/ni.2983.
4. Brulois, K., Rajaraman, A., Szade, A., Nordling, S., Bogoslowski, A., Dermadi, D.,
Rahman, M., Kiefel, H., O'Hara, E., Koning, J.J., et al. (2020). A molecular map of murine
lymph node blood vascular endothelium at single cell resolution. Nat Commun *11*, 3798.
10.1038/s41467-020-17291-5.
5. Tabula Muris, C., Overall, c., Logistical, c., Organ, c., processing, Library, p., sequencing,
Computational data, a., Cell type, a., Writing, g., et al. (2018). Single-cell transcriptomics
of 20 mouse organs creates a Tabula Muris. Nature *562*, 367-372. 10.1038/s41586-018-
0590-4.
6. Dinh, T.T., Xiang, M., Rajaraman, A., Wang, Y., Salazar, N., Zhu, Y., Roper, W., Rhee,
S., Brulois, K., O'Hara, E., et al. (2022). An NKX-COUP-TFII morphogenetic code directs
mucosal endothelial addressin expression. Nat Commun *13*, 7448. 10.1038/s41467-022-
34991-2.
7. Bi, Y., Shi, X., Zhu, J., Guan, X., Garbacz, W.G., Huang, Y., Gao, L., Yan, J., Xu, M., Ren,
S., et al. (2018). Regulation of Cholesterol Sulfotransferase SULT2B1b by Hepatocyte
Nuclear Factor 4alpha Constitutes a Negative Feedback Control of Hepatic
Gluconeogenesis. Mol Cell Biol *38*. 10.1128/MCB.00654-17.
8. Papandreou, I., Denko, N.C., Olson, M., Van Melckebeke, H., Lust, S., Tam, A., Solow-
Cordero, D.E., Bouley, D.M., Offner, F., Niwa, M., and Koong, A.C. (2011). Identification
of an Ire1alpha endonuclease specific inhibitor with cytotoxic activity against human
multiple myeloma. Blood *117*, 1311-1314. 10.1182/blood-2010-08-303099.
9. Lin, J.C., Liu, T.P., Chen, Y.B., and Yang, P.M. (2023). PF-429242 exhibits anticancer
activity in hepatocellular carcinoma cells via FOXO1-dependent autophagic cell death and
IGFBP1-dependent anti-survival signaling. Am J Cancer Res *13*, 4125-4144.
10. Wang, T.B., Geng, M., Jin, H., Tang, A.G., Sun, H., Zhou, L.Z., Chen, B.H., Shen, G., and
Sun, Q. (2021). SREBP1 site 1 protease inhibitor PF-429242 suppresses renal cell
carcinoma cell growth. Cell Death Dis *12*, 717. 10.1038/s41419-021-03999-9.
11. Reiss, Y., Proudfoot, A.E., Power, C.A., Campbell, J.J., and Butcher, E.C. (2001). CC
chemokine receptor (CCR)4 and the CCR10 ligand cutaneous T cell-attracting chemokine
(CTACK) in lymphocyte trafficking to inflamed skin. J Exp Med *194*, 1541-1547.
10.1084/jem.194.10.1541.
12. Benck, C.J., Martinov, T., Fife, B.T., and Chatterjea, D. (2016). Isolation of Infiltrating
Leukocytes from Mouse Skin Using Enzymatic Digest and Gradient Separation. J Vis Exp,
e53638. 10.3791/53638.
13. Kunkel, E.J., Campbell, J.J., Haraldsen, G., Pan, J., Boisvert, J., Roberts, A.I., Ebert, E.C.,
Vierra, M.A., Goodman, S.B., Genovese, M.C., et al. (2000). Lymphocyte CC chemokine
receptor 9 and epithelial thymus-expressed chemokine (TECK) expression distinguish the
small intestinal immune compartment: Epithelial expression of tissue-specific chemokines
as an organizing principle in regional immunity. J Exp Med *192*, 761-768.
10.1084/jem.192.5.761.
14. Buenrostro, J.D., Giresi, P.G., Zaba, L.C., Chang, H.Y., and Greenleaf, W.J. (2013).
Transposition of native chromatin for fast and sensitive epigenomic profiling of open
chromatin, DNA-binding proteins and nucleosome position. Nat Methods *10*, 1213-1218.
10.1038/nmeth.2688.

15.   Hinrichs, A.S., Karolchik, D., Baertsch, R., Barber, G.P., Bejerano, G., Clawson, H.,
Diekhans, M., Furey, T.S., Harte, R.A., Hsu, F., et al. (2006). The UCSC Genome Browser
Database: update 2006. Nucleic Acids Res *34*, D590-598. 10.1093/nar/gkj144.
16.   Siepel, A., Bejerano, G., Pedersen, J.S., Hinrichs, A.S., Hou, M., Rosenbloom, K.,
Clawson, H., Spieth, J., Hillier, L.W., Richards, S., et al. (2005). Evolutionarily conserved
elements in vertebrate, insect, worm, and yeast genomes. Genome Res *15*, 1034-1050.
10.1101/gr.3715005.
17.   Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X., Murre,
C., Singh, H., and Glass, C.K. (2010). Simple combinations of lineage-determining
transcription factors prime cis-regulatory elements required for macrophage and B cell
identities. Mol Cell *38*, 576-589. 10.1016/j.molcel.2010.05.004.
18.   Chen, E.Y., Tan, C.M., Kou, Y., Duan, Q., Wang, Z., Meirelles, G.V., Clark, N.R., and
Ma'ayan, A. (2013). Enrichr: interactive and collaborative HTML5 gene list enrichment
analysis tool. BMC Bioinformatics *14*, 128. 10.1186/1471-2105-14-128.
19.   Kuleshov, M.V., Jones, M.R., Rouillard, A.D., Fernandez, N.F., Duan, Q., Wang, Z.,
Koplev, S., Jenkins, S.L., Jagodnik, K.M., Lachmann, A., et al. (2016). Enrichr: a
comprehensive gene set enrichment analysis web server 2016 update. Nucleic Acids Res
44, W90-97. 10.1093/nar/gkw377.
20.   Xie, Z., Bailey, A., Kuleshov, M.V., Clarke, D.J.B., Evangelista, J.E., Jenkins, S.L.,
Lachmann, A., Wojciechowicz, M.L., Kropiwnicki, E., Jagodnik, K.M., et al. (2021). Gene
Set Knowledge Discovery with Enrichr. Curr Protoc *1*, e90. 10.1002/cpz1.90.
2185