A mouse model of PTEN Hamartoma Tumour Syndrome reveals that loss of the nuclear function of PTEN drives macrocephaly, lymphoid overgrowth, and late-onset cancer

Priyanka Tibarewal[1,*], Victoria Rathbone[1], Sarah E Conduit[1], Gala Anastasia Electra Classen[1], Fiona Black[1], Mohammad Amin Danesh[1], Georgia Constantinou[1], Zeinab Asgarian[2], Koujiro Tohyama[3], Nisha Kriplani[4], Virginia Alvarez Garcia[4], Elizabeth Foxall[1], Djenat Belarbi[5,6,7], Marie Leverve[5,6,7], Wayne Pearce[1], Mahreen Adil[1], Zofia Varyova[1], Lucia Conde[1], Adriana Alves[1], Glenn R Masson[8], Roger L Williams[8], Adrienne M Flanagan[9, 10], Javier Herrero[1], , Katerina Stroud[11], Marc Tischkowitz[11], Katherine Lachlan[12,13], Cheryl L Scudamore[14], Mark G H Scott[5,6,7], Nicholas R Leslie[4], Nicoletta Kessaris[2,*], Bart Vanhaesebroeck[1,*]

[1]Cancer Institute, University College London, UK
[2]Wolfson Institute of Biomedical Research, University College London, UK
[3]Department of Physiology, School of Dentistry, Iwate Medical University, Yahaba, Japan
[4]Institute of Biological Chemistry, Biophysics and Bioengineering, Heriot Watt University, Edinburgh, UK
[5]Université de Paris, 75014 Paris, France
[6]Institut Cochin, INSERM U1016, 75014 Paris, France
[7]CNRS, UMR8104, 75014 Paris, France
[8]MRC Laboratory of Molecular Biology, Cambridge, UK
[9]Research Department of Pathology, University College London, UK
[10]Cellular and Molecular Pathology, Royal National Orthopaedic Hospital, Stanmore, UK
[11]Department of Genomic Medicine, National Institute for Health Research Cambridge Biomedical Research Centre, University of Cambridge, Cambridge, UK
[12]Wessex Clinical Genetics Service, University Hospital Southampton, Princess Anne Hospital, Southampton, UK
[13]Human Development and Health, Faculty of Medicine, University of Southampton, Southampton, UK
[14]Exepathology, Exmouth, UK

*Corresponding authors

Author contributions: PT, NRL, NK and BV conceived the study, PT, VR, SEC, GAEC, FB, MAD, GC, ZA,KT, NK, VAG, EF, DB, ML,WP, MA, ZV, SH, LC, AA, GRM and MGHS performed and analysed the experiments, CLS performed the histological analysis, RLW, AMF and JH provided reagents/resources for the study, PT, NK and BV wrote the manuscript.

Competing interest statement: BV is a consultant for Pharming (Leiden, The Netherlands) and iOnctura (Geneva, Switzerland) and a shareholder of Open Orphan (Dublin, Ireland).

ABSTRACT

PTEN Hamartoma Tumour Syndrome (PHTS) is a rare disorder characterized by germline heterozygous mutations in the PTEN tumour suppressor gene, leading to multi-organ/tissue overgrowth, autism spectrum disorder and increased cancer risk. PHTS individuals display heterogeneity in phenotypes, which has been linked in part to the diverse genetic alterations in the *PTEN* gene and the multifaceted functions of this protein. Indeed, while PTEN primarily functions as a PIP_3 lipid phosphatase in the cytosol, regulating PI3K/AKT signalling, a pathway commonly

deregulated in cancer, it also plays crucial roles in maintaining chromosomal stability through
nuclear activities such as double strand (ds) DNA damage repair. Recent studies have identified a
subset of missense PHTS variants that cause nuclear exclusion of PTEN, impairing its nuclear
functions. Here, we present our findings from one such pathogenic variant, *PTEN-R173C*, frequently
found in PHTS and somatic cancers. Using cell biological and mouse modelling approaches, we show
that PTEN-R173C has higher PIP$_3$ phosphatase activity than wild-type PTEN, resulting in effective
regulation of canonical PI3K/AKT signalling. However, PTEN-R173C is unstable and excluded from
the nucleus. Aligning with their near normal PI3K/AKT signalling, *Pten$^{+/R173C}$* mice display a low
incidence of solid tumours compared to *Pten$^{+/-}$* mice. *Pten$^{+/R173C}$* mice also exhibit lymphoid
hyperplasia and macrocephaly which correlates with compromised nuclear functions of PTEN-
R173C. That nuclear functions are compromised is demonstrated by reduced dsDNA damage repair
in *Pten$^{+/R173C}$* mice. Integrating PHTS patient data with findings from our mouse model, our study
indicates that nuclear dysfunction of pathogenic *PTEN* variants is a key factor in predicting the onset
of the different PHTS-associated phenotypes. We speculate that late-onset cancer in individuals with
nuclear-excluded PTEN results from genetic alterations unrelated to PTEN itself, facilitated by
impaired PTEN-mediated dsDNA damage repair.

KEY WORDS: PTEN, PI3K, PHTS, overgrowth, mouse model, cancer, macrocephaly, germline, ASD

INTRODUCTION
PTEN is a ubiquitously expressed tumour suppressor (Alvarez-Garcia et al., 2019, Li et al., 1997, Steck
et al., 1997) that shows complete loss of expression in some cancers (e.g. prostate cancer and
glioblastoma) and overrepresentation of missense variants in other tumours (e.g. endometrial cancer)
(Alvarez-Garcia et al., 2019, Yehia et al., 2020). Heterozygous germline variants in PTEN cause PTEN
Hamartoma Tumour Syndrome (PHTS), a rare autosomal dominant disorder. A recent study puts the
prevalence of PHTS as 1:8000-13,000 (White et al., 2025) which is 10-20 fold higher than previous
estimates (Nelen et al., 1996). Clinical features of PHTS include macrocephaly, developmental delay
(DD), autism spectrum disorder (ASD), dermatological pathologies, gastrointestinal (GI) polyps,
immune dysfunction, vascular anomalies. Individuals with PHTS also face a significantly increased risk
of developing cancers particularly of the breast, endometrium and thyroid (Ngeow and Eng, 2019,
Yehia and Eng, 2018), with emerging evidence for ovarian cancer (Schei-Andersen et al., 2025).
PTEN acts in both the cytosol and nucleus. Cytosolic functions of PTEN predominantly include lipid
phosphatase   activity   on   phosphatidylinositol(3,4,5)trisphosphate   (PIP$_3$)   and
phosphatidylinositol(3,4)bisphosphate (PI(3,4)P$_2$), thereby antagonizing PI3K/AKT signalling (Myers et
al., 1998). PTEN also has protein-phosphatase activity, reducing phosphorylation of Ser/Thr/Tyr on
proteins such as PTK6, DREBRIN, DVL2, FAK and SHC (Leslie et al., 2009) , in addition to auto-
dephosphorylating its T366 site (Tibarewal et al., 2012). Nuclear PTEN has been implicated in cell cycle
control, dsDNA repair, maintenance of chromatin structure, genome integrity and transcriptional
control (Ho et al., 2019, Langdon, 2023).
The functional core of the PTEN protein consists of a catalytic (amino acids (AA) 16-185) and a C2
(AA 186-350) domain (Lee et al., 1999), with mutations in different regions differentially impacting
PTEN expression and function. 65% of PHTS patients express *PTEN* variants such as deletions, DNA
insertions or non-sense mutation leading to STOP codons (such as R130X and R233X) that result in
PTEN truncation and are therefore predicted to cause heterozygous loss-of-expression. The remaining
PHTS patients carry heterozygous missense mutations in the phosphatase or C2 domain, leading to
expression of PTEN proteins with altered function or subcellular distribution (Hendricks et al., 2022,

Spinelli et al., 2015). These include mutations in the catalytic pocket resulting in loss of all catalytic
activity [e.g. C124 (C124S/R), R130 (R130G/Q/L) and G129 (G129R)] (Lee et al., 1999) or selective loss
of PIP$_3$ phosphatase activity, while retaining protein phosphatase activity (e.g. G129E) (Myers et al.,
1998). Missense mutations leading to selective loss of protein phosphatase activity (Y138L/C) have
also been described, although these have not been reported in PHTS (Davidson et al., 2010, Tibarewal
et al., 2012). Recent reports have shown that a subset of PTEN variants in PHTS (including K289E,
D252G, F241S, I101T and others) are excluded from the nucleus and therefore lack all PTEN nuclear
functions (Langdon, 2023).

The PIP$_3$ phosphatase activity of PTEN has long been regarded as the main and critical factor in
PHTS pathogenesis, yet other lipid phosphatase-independent functions of PTEN are now also emerging
as important (Langdon, 2023). Various studies have attempted to establish genotype-phenotype
associations in PHTS. Limited patient data make it challenging to ascertain the impact of individual
variants in driving patient phenotypes. This challenge has been partially overcome by classifying the
variants based on their position in the *PTEN* gene and their impact on PTEN expression and function
(Lachlan et al., 2007, Leslie and Longy, 2016, Marsh et al., 1998, Marsh et al., 1999, Mighell et al.,
2018). Recently, a large paediatric and adult patient cohort study revealed that missense variants are
associated with early disease onset, with DD and macrocephaly being the reason for early diagnosis,
whereas *PTEN* truncation variants are associated with late onset disease, with cancer being the reason
for PHTS diagnosis at later stages in life (Hendricks et al., 2022). However, patient cohort studies have
been limited by the variability in age and sex of the individuals, one of the key challenges in rare disease
research.

Genetically modified mice have helped overcome some of the challenges in PTEN research by
allowing assessment of PTEN functions in an endogenous tissue context (Lee and Pandolfi, 2020). They
also serve as a platform for testing potential treatments, such as PI3K pathway inhibitors, aimed at
mitigating the effects of PTEN loss. An example are mice with heterozygous loss of PTEN expression
(*Pten$^{+/-}$*), representing *PTEN* variants that result in loss of expression and function. This mouse model
recapitulates many aspects of human PHTS remarkably well in terms of cancer development and
lymphoid and brain overgrowth (Podsypanina et al., 1999, Tibarewal et al., 2022).

Here we report the creation and characterisation of a novel mouse model for PHTS, carrying a
PTEN-R173C (NM_000314.8(PTEN):c.517C>T (p.Arg173Cys)) missense variant in the germline. R173 is
located in the TI (Thr-Ile) loop in the catalytic domain of PTEN important for PTEN catalytic activity (Lee
et al., 1999). Mutation of R173 to Cys or His in similar ratios are observed in PHTS and sporadic cancers,
with conflicting reports on catalytic activity impact, from having been reported active to inactive (Han
et al., 2000, Post et al., 2020, Smith and Briggs, 2016). R173 variants are frequently found in multiple
patient cohort studies with 34 PHTS patients been reported in literature to the best of our knowledge,
and an additional four new patients identified with this variant in the UK (**Supplementary Table 1**).
Most patients with R173 variants received PHTS diagnosis at young age due to presentation of
macrocephaly, DD and cutaneous pathology. Adult patients additionally presented with breast cancer,
benign and malignant tumours of the skin, thyroid and ovary, endometrial polyps and fibroids, GI
polyps, with malignant disease onset much later in life (over the age of 50 years). Vascular
malformations, mainly haemangiomas, were also seen in adult patients (**Table 1** and **Supplementary**
Table 1).

Here, we first revisited the impact of the R173 mutation on PTEN expression and function using
biochemical and cell-based models, with a focus on PTEN-R173C. We show that PTEN-R173C has
reduced protein stability but retains its ability to dephosphorylate PIP$_3$ and therefore regulate the
canonical PI3K/AKT pathway in the cytosol. However, we find that this PTEN mutant is largely excluded
from the nucleus.

Several mouse models have been developed to distinguish the relative importance of the cytosolic
versus nuclear functions of PTEN (reviewed in (Langdon, 2023)). For instance, *Pten*^{m3m4} mice,
engineered by Eng *et al.*, feature non-naturally occurring PTEN mutations that result in a
predominantly cytosolic PTEN distribution without affecting its PIP$_3$ phosphatase activity. These mice
have been extensively studied for the impact of the nuclear-cytoplasmic partitioning of PTEN on
neuronal and immune phenotypes, although their tumour spectrum has not been reported (Mester
et al., 2011, Tilot et al., 2014, Jaini et al., 2020). Other models, expressing PTEN missense variants
found in PHTS such as F341V (Caserta et al., 2015), and non-pathogenic mutations such as K13R, Y240F
and D384V (Igarashi et al., 2018, Ma et al., 2019), that do not affect the catalytic functions of PTEN but
impact its nuclear/sub-cellular localisation, have also been developed, but there is limited or no
patient data for these individual variants (Table 1 and Supplementary Table 2).
We created heterozygous germline *Pten*^{R173C} knock-in mice (*Pten*$^{+/R173C}$) as a unique model of a
frequent pathogenic PHTS PTEN variant representing nuclear-excluded PTEN missense variants. We
examined the consequences of this mutation on embryonic development, glucose metabolism,
development of PHTS-relevant cancers as well as immune, neurological and behavioural phenotypes.
By integrating our findings with phenotypic data from patients for this variant, and by comparison of
reported data on the *Pten*^{m3m4} mice, we delineate the role of the nuclear functions of PTEN in the
manifestation of PHTS phenotypes.
160
RESULTS
Increased PIP$_3$ phosphatase activity but reduced protein stability of PTEN-R173 mutant proteins
upon transient expression in mammalian cells
Conflicting data on the catalytic activity of PTEN-R173 variants have been published. PTEN-R173
mutant proteins, expressed in *E. coli*, have been reported to be catalytically inactive (Han et al., 2000)
however, mutant PTEN proteins expressed in non-mammalian systems often fail to fold properly,
resulting in lack of activity (Spinelli et al., 2015). Consistent with this, in our hands, PTEN-R173C protein
expressed in *E. coli* lacked *in vitro* PIP$_3$ lipid phosphatase activity (Fig. S1) and when expressed in Sf9
insect cells precipitated upon purification (Masson et al., 2016a) (data not shown), rendering it
unsuitable for use in *in vitro* assays.
A commonly used assay to determine the ability of PTEN and its mutants to regulate PI3K/AKT
signalling involves transfection into the PTEN-null U-87 MG glioblastoma cell line (further referred to
as U87) and assessment of basal AKT phosphorylation and cell proliferation (Davidson et al., 2010).
Transient lentiviral expression of untagged PTEN-R173C in U87 revealed consistently lower expression
levels compared to PTEN-WT (Fig. 1A, left panel), despite similar mRNA expression levels (Fig. 1A, right
panel). Similar results were seen with PTEN-R173H, suggesting that other PTEN-R173 mutant proteins
are less stable than PTEN-WT.
Expression of a dose range of PTEN proteins in U87 revealed that, compared to PTEN-WT, lower
levels of PTEN-R173C or PTEN-R173H effectively downregulated AKT P-S473 and S6 P-S240/244
phosphorylation (Fig. 1B) and suppressed U87 proliferation (Fig. 1C and S2A). We also assessed the
impact of PTEN-R173C expression in U87 cells on recruitment of a PIP$_3$-biosensor protein to the plasma
membrane. In both serum-starved or insulin-stimulated cells, the biosensor plasma membrane to
cytosol ratio (PM:cytosol) was significantly reduced upon treatment with the pan-class I PI3K inhibitor
GDC-0941 or upon expression of PTEN-WT or PTEN-R173C, with the strongest decrease upon PTEN-
R173C expression (Fig. 1D and S2B), despite its lower expression levels compared to PTEN-WT (Fig.
S2C).
Taken together, these data suggest that in mammalian cells, PTEN-R173C is less stable but
catalytically more active than PTEN-WT.
189
Evidence for a more open, proteolytically-sensitive protein conformation of PTEN-R173 mutants
We next investigated possible biochemical mechanisms underlying the reduced expression of PTEN-
R173C mutants in U87 cells. Cycloheximide chase experiments of U87 cells transiently transfected with

PTEN expression constructs indicated a reduced protein half-life of PTEN-R173C compared to PTEN-
WT (**Fig. 1E**, **S3A** and Ref.(Spinelli et al., 2015)). A recent study has shown similar findings with PTEN-
R173H (Jun et al., 2024).
PTEN protein stability and activity is regulated, in part, by phosphorylation on multiple C-terminal
sites (S380/T382/T383/S385) by kinases such as CK2 (Torres and Pulido, 2001). Such phosphorylation
results in a more closed, less active and more stable protein conformation (Torres and Pulido, 2001,
Vazquez et al., 2001, Birle et al., 2002, Odriozola et al., 2007, Dempsey et al., 2021). Upon transient
expression of similar levels of untagged proteins in U87 cells, PTEN-R173C showed a ~50% lower level
of S380/T382/T383 phosphorylation compared to PTEN-WT (**Fig. 1F**). Similar data were observed using
YFP-tagged Rluc-PTEN-R173C and PTEN-R173H in HEK-293T cells (**Fig. S3B**), suggesting that reduced
phosphorylation may contribute to the observed instability of PTEN-R173 mutants.
Dephosphorylation of S380/T382/T383/S385 on PTEN releases the inhibitory effect of these
phosphorylated residues on the C2 domain of PTEN, leading to a more open conformation with
increased membrane binding and therefore enhanced activity (Torres and Pulido, 2001, Vazquez et
al., 2001, Birle et al., 2002, Odriozola et al., 2007). It has also been shown that, once at the membrane,
the active form of PTEN becomes degraded (Maccario et al., 2010). A study using computational
modelling predicted increased protein flexibility of PTEN-R173C/H mutants, leading to impaired
protein stability (Smith and Briggs, 2016). In order to gain insight into these phenomena, we used an
intramolecular bioluminescent resonance energy transfer (BRET)-based biosensor Rluc-PTEN-YFP in
live HEK-293T and PTEN KO HeLa cells. This biosensor reveals dynamic changes in PTEN conformational
rearrangement and function due to shifts in energy transfer between the donor/acceptor couple
(Lima-Fernandes et al., 2014) (**Fig. S4A-B**). Using this approach, PTEN-R173C and PTEN-R173H were
found to be in a more open/flexible conformation compared to PTEN-WT (**Fig. 1G**).
We previously reported that PTEN protein dephosphorylated on S380/T382/T383/S385 is more
prone to ubiquitination and degradation, and that this involves the K66 ubiquitination site in PTEN
(Maccario et al., 2010, Gupta and Leslie, 2016). In line with this, when transiently expressed in U87
cells, the PTEN-R173C/K66R double mutant which cannot be ubiquitinylated, showed increased
protein expression levels compared to PTEN-R173C (**Fig. 1H**).
Altogether these data indicate that the PTEN-R173C protein has a more open conformation that
allows ubiquitination and degradation, leading to reduced protein levels.
223
PTEN-R173 mutants are predominantly cytoplasmic and largely excluded from the nucleus
The tumour suppressor functions of PTEN have been mostly attributed to its ability to regulate the
PI3K/AKT pathway, with the role of other PTEN functions such as protein phosphatase activity and
nuclear functions less well-studied in this context (Ho et al., 2019). Given that our results so far show
that PTEN-R173C/H mutants are effective at regulating the PI3K/AKT pathway, we turned our
attention to other PTEN functions that might explain the high mutation frequency of this site in
somatic cancers and PHTS.
Analysis of the subcellular distribution of PTEN-R173C by transiently expressing C-terminally GFP-
tagged PTEN-R173C or PTEN-WT in Lenti-X™ 293T cells revealed a significantly lower
nuclear:cytoplasmic ratio of the GFP signal in cells expressing PTEN-R173C-GFP than in cells with PTEN-
WT-GFP, suggesting that PTEN-R173C was largely excluded from the nucleus (**Fig. 1I**). Nuclear
exclusion of the R173 mutants was also revealed upon expression of YFP-tagged Rluc-PTEN-WT, -
R173C or -R173H in HEK-293T or PTEN KO HeLa cells (**Fig. S5**).
237
Homozygous germline PTEN-R173C expression is embryonically lethal in mice
In order to study PTEN-R173C in its endogenous tissue context and to better understand its role in
PHTS, we created heterozygous germline *Pten*[R173C] knock-in mice (*Pten*[+/R173C]) by CRISPR-Cas9
technology (**Fig. S6**). Intercrosses of *Pten*[+/R173C] mice did not yield live-born homozygous *Pten*[R173C/R173C]
mice (**Fig. 2A**). *Pten*[R173C/R173C] embryos were almost all reabsorbed at embryonic day (E)11.5, with
under-representation compared to Mendelian ratios at earlier time points (**Fig. 2A**). Intact

Pten$^{R173C/R173C}$ embryos were found up to E9.5 at which stage, 5 out of 9 embryos were smaller and had
not undergone turning, with the remaining 4 having been reabsorbed, suggesting that *Pten*$^{R173C/R173C}$
embryos die around E9.5 (Fig. 2A).

Given the embryonic lethality of homozygous *Pten* $^{R173C/R173C}$ mice, and the notion that PTEN
mutations in PHTS individuals are heterozygous, in the next series of experiments we used
heterozygous *Pten*$^{+/R173C}$ mice to further investigate the impact of the PTEN R173C mutation. For these
studies, we used littermate wild-type (*Pten*$^{+/+}$) mice as controls as well as *Pten*$^{+/-}$ mice, a well-
characterised model for PHTS (Knobbe et al., 2008, Di Cristofano et al., 1998, Suzuki et al., 1998,
Podsypanina et al., 1999, Freeman et al., 2006, Di Cristofano et al., 2001).

PTEN-R173C expressed from its endogenous promoter is unstable but retains PIP$_3$ phosphatase
activity
As a cell-based model to study endogenous PTEN-R173C protein expression and function, we derived
mouse embryonic fibroblasts (MEFs) of the following genotypes: *Pten*$^{+/R173C}$, *Pten*$^{R173C/R173C}$, *Pten*$^{-/R173C}$
and compared these to *Pten*$^{+/-}$, *Pten*$^{-/-}$ and *Pten*$^{+/+}$ MEFs.

PTEN protein expression levels in *Pten*$^{+/R173C}$ and *Pten*$^{+/-}$ MEFs were 60-70% of those in *Pten*$^{+/+}$ MEFs
(Fig. 2B). The PTEN levels observed in *Pten*$^{+/-}$ MEFs agree with previous studies where the *Pten*^{WT} allele
was shown to express higher than 50% level of PTEN protein (Alimonti et al., 2010). In *Pten*$^{R173C/R173C}$
and *Pten*$^{-/R173C}$ MEFs, PTEN protein levels were ~13% and ~2%, respectively, of those in *Pten*$^{+/+}$ MEFs
(Fig. 2C; Fig. S7), indicative of low protein stability of the PTEN-R173C protein, in line with our
exogenous expression studies above.

We next used immunoblotting to assess PI3K pathway activation, under standard cell culture
conditions and upon insulin-stimulation. *Pten*$^{+/R173C}$, *Pten*$^{+/-}$ and *Pten*$^{+/+}$ MEFs growing in complete
media showed similar basal levels of phosphorylation of AKT, S6 and other downstream effectors (Fig.
2B). Under these conditions, *Pten*$^{-/-}$ MEFs had increased levels of AKT P-S473, which was not observed
in *Pten*$^{R173C/R173C}$ MEFs (Fig. 2C). In *Pten*$^{-/R173C}$ MEFs, the levels of AKT P-S473 were elevated compared
to *Pten*$^{+/+}$ and *Pten*$^{+/-}$ MEFs, but still substantially lower compared to *Pten*$^{-/-}$ MEFs (Fig. S7).

Upon insulin stimulation, the increase in AKT P-S473 was dampened more effectively in
Pten$^{R173C/R173C}$ MEFs than in *Pten*$^{+/+}$ and *Pten*$^{-/-}$ MEFs, despite the very low levels of PTEN-R173C
expression (Fig. 2D).

Taken together, these data indicate that, even at low levels of expression, PTEN-R173C is still
capable of effectively downregulating unstimulated and insulin-stimulated AKT P-S473
phosphorylation and thus retains overall PIP$_3$ phosphatase activity.

PTEN-R173C regulates glucose metabolism *in vivo*
PTEN plays an important role in glucose metabolism by dampening insulin signalling, with *Pten*$^{+/-}$ mice
showing enhanced acute insulin signalling and increased glucose metabolism (Chen et al., 2018,
Garcia-Cao et al., 2012, Pal et al., 2012). PHTS patients also often show constitutive insulin sensitization
(Pal et al., 2012). In line with these reports, insulin tolerance tests on 3-month-old mice revealed
enhanced insulin sensitivity in *Pten*$^{+/-}$ mice, evidenced by lower levels of blood glucose post insulin
injection compared to WT littermates (Fig. 2E). In contrast, WT and *Pten*$^{+/R173C}$ mice showed a similar
insulin response up to 60 min after insulin injection (Fig. 2E). This indicates that, at the organismal
level, PTEN-R173C maintains the ability to downregulate acute insulin/PI3K/AKT signalling and
therefore glucose homeostasis (Fig. 2E).

PTEN-R173C expressed from an endogenous promoter is predominantly nuclear-excluded with
reduced ability to regulate dsDNA damage repair
To determine the subcellular distribution of PTEN-R173C upon expression from its endogenous locus,
we used immunocytochemistry and quantification of the PTEN nuclear/cytoplasmic ratio in
Pten$^{R173C/R173C}$ and *Pten*$^{-/R173C}$ MEFs. These data showed that PTEN-R173C expressed from its
endogenous locus was predominantly nuclear-excluded, in contrast to PTEN-WT protein which was

present in both the nucleus and cytoplasm (**Fig. 2F** and **S8A**). Given that R173 is not located in a known
PTEN nuclear localisation signal, one reason for the apparent nuclear exclusion of PTEN-R173C could
be sequestration at the plasma membrane, with little protein available for nuclear import. To
investigate this, we treated *Pten*$^{R173C/R173C}$ MEFs with the pan-class I PI3K inhibitor GDC-0941, with the
aim to reduce the PIP$_3$ substrate levels for PTEN at the plasma membrane and to potentially restore
the presence of PTEN-R173C in the nucleus. However, GDC-0941 treatment did not affect the
subcellular distribution of PTEN-R173C (**Fig. S8B**), suggesting that plasma membrane PIP$_3$ levels do not
significantly contribute to the nuclear exclusion of PTEN-R173C.
Nuclear PTEN has numerous functions including regulation of DNA repair, transcription, chromatin
structure, chromosomal stability and genome integrity by various mechanisms (Ho et al., 2019,
Langdon, 2023). We investigated one of these nuclear functions in *Pten*$^{+/R173C}$ mice, namely dsDNA
repair. Nuclear PTEN deficiency has been shown to cause a homologous recombination (HR) defect in
cells, leading to increased γH2AX expression and decreased Rad51 foci formation upon genotoxic
stress such as ionizing radiation (Bassi et al., 2013, Mendes-Pereira et al., 2009, Shen et al., 2007).
Ionizing radiation (IR) of *Pten*$^{+/R173C}$ and *Pten*$^{+/-}$ mice, followed by analysis 5 h post-IR treatment, led to
a similar increase in γH2AX levels in protein extracts from uteri, kidney and liver, compared to wild-
type mice (**Fig. 2G** and **S9**), indicative of a similarly reduced DNA repair capacity upon heterozygous
PTEN loss or heterozygous PTEN-R173C expression. This suggests that PTEN-R173C has a reduced
ability to regulate DNA damage repair compared to PTEN-WT.
314
Reduced cancer predisposition of *Pten*$^{+/R173C}$ mice compared to *Pten*$^{+/-}$ mice
Individuals with PHTS have an increased risk of cancers of multiple tissues including breast,
endometrium and thyroid (Yehia and Eng, 2018, Yehia et al., 2020, Yehia et al., 2019). To gain insight
into the cancer predisposition of germline *Pten*^{R173C} expression, we aged heterozygous *Pten*$^{+/R173C}$ mice
alongside *Pten*$^{+/-}$ and *Pten*$^{+/+}$ littermate controls. We used mice on a mixed C57Bl/6JxSv129
background, as on this background *Pten*$^{+/-}$ mice develop PHTS-relevant phenotypes, with an
overlapping tumour spectrum including hyperplasia/tumours of the thyroid, endometrium, lymphoid
tissue, small intestine and adenomyoepithelioma and malignant mammary tumours in the females
(Tibarewal et al., 2022). *Pten*$^{+/-}$ mice also develop lesions not seen in PHTS patients including,
pheochromocytoma and intraepithelial neoplasia (PIN) and adenocarcinoma of the prostate (Knobbe
et al., 2008, Di Cristofano et al., 1998, Suzuki et al., 1998, Podsypanina et al., 1999, Freeman et al.,
2006, Di Cristofano et al., 2001).
Gross morphology and development of *Pten*$^{+/R173C}$ mice was found to be similar to *Pten*$^{+/+}$
littermates, with a tendency for a slight increase in body weight at young age in both *Pten*$^{+/R173C}$ and
Pten$^{+/-}$ mice (**Fig. S10**).
Mice were aged and sacrificed at first appearance of signs of ill health or palpable masses. The
reasons for humane culling of mice are listed in **Supplementary Table 3**. Under these experimental
conditions, the median survival age of *Pten*$^{+/R173C}$ mice was 585 and 302 days for male and female mice,
respectively. This is significantly longer than the median survival age of 310 and 179 days for male and
female *Pten*$^{+/-}$ mice, respectively (**Fig.3A**). However, *Pten*$^{+/R173C}$ mice had a shorter lifespan compared
to wild-type mice (**Fig.3A**). This was especially the case for female *Pten*$^{+/R173C}$ mice, which were mainly
culled due to palpable masses or ill health.
Histopathological analysis of selected tissues from *Pten*$^{+/-}$ and *Pten*$^{+/R173C}$ mice revealed very similar
gross morphologies of all tissues analysed, with overall histopathology data summarized in **Fig. 3B** and
Supplementary Tables 4-7.
Pten$^{+/R173C}$ mice developed far fewer tumours than *Pten*$^{+/-}$ mice, with occasional hyperplasia of the
small intestine (2/14 mice) and low-grade PINs of prostate (4/14 mice), one case of mammary
adenocarcinoma (1/22 mice) and one case of endometrial hyperplasia (1/28) (**Fig. 3B; Fig. S11-S12;**
Supplementary Tables 4 and 5). Similar to *Pten*$^{+/-}$ mice, *Pten*$^{+/R173C}$ mice had a high incidence of
pheochromocytoma (**Fig. 3B, Fig. S11-S12; Supplementary Tables 4 and 5**).

One out of 28 *Pten*$^{+/R173C}$ mice displayed a single focus of endometrial hyperplasia, which correlated
with loss of immunostaining for PTEN and increased AKT P-S473 (**Fig. 3C**). Interestingly, loss of PTEN
immunoreactivity was also observed in microscopically *normal* endometrial glands from 5-month-old
and 10-month-old *Pten*$^{+/R173C}$ mice, but this was not accompanied by a concomitant increase in AKT P-
S473 immunoreactivity in these regions (**Fig. 3C**). In the absence of reagents which allow us to
specifically stain for WT or mutant PTEN protein and, based on the low expression level of endogenous
PTEN-R173C protein when expressed alone (**Fig. S7**), it is most likely that expression of the wild-type
Pten allele is lost in these morphologically normal glands. We speculate that, while the very low
expression level of PTEN-R173C from the endogenous allele may not be detectable by IHC, it is still
sufficient to partially downregulate PI3K signalling and AKT P-S473 and thereby prevent the
development of endometrial tumours. Although this observation is based on a single mouse, we
speculate that the hyperplastic gland found in the endometrium of this *Pten*$^{+/R173C}$ mouse may have
resulted either from loss-of-expression of the remaining *Pten*^{R173C} allele, or from acquisition of PI3K
pathway-activating mutations. These events likely led to PI3K pathway hyperactivation, ultimately
contributing to the development of endometrial hyperplasia. This hypothesis is supported by the fact
that somatic cancers with the PTEN-R173 mutation, along with mutation of the wild-type allele of
PTEN, also contain mutations in other cancer drivers such as *PIK3CA* (**Fig. 3D**).
362
Prominent but delayed lymphoid hyperplasia in *Pten*$^{+/R173C}$ mice compared to *Pten*$^{+/-}$ mice
Previous studies have reported a range of lymphoid and immune phenotypes in PHTS patients
including peripheral lymphoid hyperplasia, enlargement of tonsils and adenoids,
hypogammaglobulinemia, recurrent respiratory tract infection and autoimmune disorders. (Chen et
al., 2017, Drissen et al., 2024). Some of these phenotypes are found in a few PHTS patients with *PTEN-
R173* variants and other nuclear-excluded variants (**Table 1** and **Supplementary Tables 1&2**).
In *Pten* mutant mice*, lymphoid hyperplasia leads to enlarged cervical, brachial, inguinal and axillary
lymph nodes (Podsypanina et al., 1999, Alimonti et al., 2010). While this does not cause discomfort to
the mice, in our study it was a main reason for sacrificing *Pten*$^{+/-}$ mice (75% males and 80% females)
and many *Pten*$^{+/R173C}$ mice (21% males and 50% females), due to animal welfare regulations on the
allowed tissue overgrowth size (**Supplementary Table 3**).
Histological analysis revealed a similar severity grade of hyperplasia (assessed by semi-quantitative
histopathological scoring) in female *Pten*$^{+/R173C}$ and *Pten*$^{+/-}$ mice but of significantly lower grade in
Pten$^{+/R173C}$ males than *Pten*$^{+/-}$ males (**Fig. 3E**). Lymphoid hyperplasia has also been reported in
heterozygous *Pten*$^{+/m3m4}$ mice (Jaini et al., 2020).
Flow cytometry analysis of lymph nodes of 5-month-old female *Pten*$^{+/-}$ and 10-month-old female
Pten$^{+/R173C}$ mice, the age at which they have visibly enlarged lymph nodes, compared to littermate
Pten$^{+/+}$ mice, revealed a significant increase in the proportion of B-cells and a corresponding decrease
in T-cell proportions, a phenotype also reported in *Pten*$^{+/m3m4}$ mice (**Fig. 3F-G**).
Although we did not assess dsDNA damage repair in the lymphoid tissues of *Pten*$^{+/R173C}$ mice, the
lymphoid phenotypes correlate with loss of nuclear PTEN as this phenotype was also seen in *Pten*$^{+/m3m4}$
mice (Jaini et al., 2020).
385
Macrocephaly, increased neuronal complexity and white matter disturbances in *Pten*$^{+/R173C}$ mice
PHTS patients, including the majority of patients with PTEN-R173 mutations (**Supplementary Table 1**),
commonly present with macrocephaly, DD and ASD. We therefore explored neurological and
behavioural phenotypes in *Pten*$^{+/R173C}$ mice.
We examined the brains and neurons from *Pten*$^{+/+}$, *Pten*$^{+/R173C}$ and *Pten*$^{+/-}$ mice and, where possible,
conditional knock-out mice that selectively lack PTEN expression in cortical pyramidal neurons
(referred to here as cKO) (Kessaris et al., 2006, Lesche et al., 2002). These cKO mice were further
crossed with *Pten*$^{+/R173C}$ mice to obtain conditional knock-in mice (referred to here as cKI) which
express PTEN-R173C in cortical pyramidal neurons without a copy of the PTEN-WT allele and are
overall hemizygous for the *Pten*^{R173C} allele (*Pten*$^{-/R173C}$). The observed phenotypes are summarized in

Table 2, which also shows a comparison to phenotypes published for *Pten^{m3m4/m3m4}* mice. The latter model germline expression of non-naturally occurring PTEN mutations that leave PTEN PIP$_3$ phosphatase activity intact but lead to its nuclear exclusion, and whose neuronal phenotypes have been extensively characterised (Mester et al., 2011, Tilot et al., 2014, Jaini et al., 2020).

We first assessed the levels of PTEN protein and downstream pathways in protein lysates from cortical neurons. In the absence of a PTEN-WT allele, the level of PTEN protein in cKI was minimal and similar to that seen in protein extracts of cKO mice (Fig. S13A-B). However, the level of AKT P-S473 in cKI brains was significantly lower compared to the cKO mice (Fig. S13A and C). These observations mirror our data in MEFs (Fig. S7), indicating that PTEN-R173C is also highly active in the brain despite its low level of expression.

At 3 months of age, *Pten^{+/R173C}* and *Pten^{+/-}* mice of both sexes had enlarged brains with increased mass compared to *Pten^{+/+}* littermates (Fig. 4A-B, left panel). Whole-body mass was also increased in both *Pten* mutant genotypes (Fig. 4B, middle panel), however, the brain was disproportionally enlarged (Fig. 4B, left panel), indicating a macrocephalic phenotype. Also, at 3- and 6-months of age, and independent of the genetic background, *Pten^{+/R173C}* and *Pten^{+/-}* mice showed a similar increase in brain mass (Fig. S14). Given the apparently normal PI3K/AKT regulatory capacity of PTEN-R173C, and the neurological phenotypes and macrocephaly previously observed in nuclear-excluded PTEN in heterozygous *Pten^{+/m3m4}* and homozygous *Pten^{m3m4/m3m4}* mice (Mester et al., 2011, Tilot et al., 2014), these observations allow us to postulate that the nuclear function of PTEN is key to the development of macrocephaly.

We next quantified several brain parameters to characterize the macrocephalic phenotype in more detail. *Pten^{+/R173C}* mice had increased cortical thickness as measured at the level of the somatosensory barrel field (Fig. 4C-D). A significant increase was also observed in the thickness of the corpus callosum at four different Bregma levels (Fig. 4E) in both *Pten^{+/R173C}* and *Pten^{+/-}* mice. Consistent with macrocephaly, a lengthwise enlargement of the corpus callosum as well as an increase in the total area of this white matter tract at Bregma level were also observed (Fig. 4F). The increased thickness of the cortex was not caused by increased NEUN+ neuronal cell numbers or neuronal densities (Fig. 4G). In contrast, immunohistochemistry for OLIG2 revealed a small but significant increase in OLIG2-expressing oligodendrocyte lineage cell numbers in the cortex of *Pten^{+/R173C}* mice (Fig. 4H). Interestingly, this increase was not seen in our *Pten^{+/-}* mice and has not been reported in *Pten^{+/m3m4}* mice but was observed in *Pten^{m3m4/m3m4}* mice. Astrocyte overabundance and astrogliosis, as detected by GFAP immunostaining, were not observed (Fig. 4H). Interestingly, astrogliosis has been reported in *Pten^{m3m4/m3m4}* mice, suggesting that the remaining wild type *Pten* allele protects the brain from this immunoinflammatory phenotype.

To gain further insight into the causes of the macrocephaly, we quantified neuronal and glial parameters *in vitro*. Morphologies of cortical pyramidal neurons *in vitro* were assessed following transfection with a GFP expression vector to allow visualisation of the cells (Fig. 5A). *Pten^{+/R173C}* neurons had enlarged cell body size and increased process complexity compared to WT controls (Fig. 5B,C). While cKO neurons demonstrated a more severe phenotype, this phenotype was not observed in *Pten^{+/-}* neurons and not reported in *Pten^{+/m3m4}* mice but was observed in *Pten^{m3m4/m3m4}* mice (Fig. 5B,C).

To determine whether glial and/or axonal defects contribute to the enlargement of the white matter, we examined the corpus callosum of *Pten^{+/+}*, *Pten^{+/R173C}* and *Pten^{+/-}* mice using transmission electron microscopy (Fig. 5D). We found a small increase in average axonal diameter in *Pten^{+/R173C}* mice, with a reduction in the number of small diameter axons and a gain of larger diameter axons (Fig. 5E). This phenotype was not observed in *Pten^{+/-}* mice. Quantification of the G-ratio in the corpus callosum showed an increase in average myelin thickness in both *Pten^{+/R173C}* and *Pten^{+/-}* mice (Fig. 5F). However, scatter plot analysis of the g-ratio as a function of axonal diameter did not show significant changes in either *Pten^{+/R173C}* and *Pten^{+/-}* mice compared to controls (Fig. 5G).

Altogether, our experiments show that the macrocephalic *Pten^{+/R173C}* phenotype is caused by alterations in multiple parameters, including increases in neuronal cell body size, axonal thickness, axonal complexity and oligodendrocyte lineage cell numbers.

Pten^{+/R173C} mice have a similar degree of macrocephaly as *Pten^{+/m3m4}* heterozygous mice and have
multiple neuronal phenotypes overlapping with *Pten^{m3m4/m3m4}* homozygous mice and *Pten^{+/-}* mice
(Table 2). Importantly, the apparently intact or even increased PI3K/AKT regulatory capacity of PTEN-
R173C, and the phenotypic overlap with *Pten^{m3m4}* mice, suggests that the nuclear function of PTEN is
key to the development of macrocephaly and several neuronal phenotypes.

Defective dsDNA damage repair is increasingly being associated with macrocephaly and ASD with
studies showing increased dsDNA breaks in cells isolated from patients with these phenotypes (Wang
et al., 2020, Attia et al., 2020). To explore the possible involvement of impaired dsDNA damage repair
in neurons in the macrocephalic phenotype, we treated *Pten^{+/-}* and *Pten^{+/R173C}* mice with IR and
examined the levels of γH2AX in the brain 5 h post-IR treatment. While there was an increase in the
γH2AX levels in protein extracts from the brains of both *Pten^{+/-}* and *Pten^{+/R173C}* mice, compared to that
seen in littermate wild-type mice, the difference observed was not statistically significant (Fig. S15).
This might be due to the IR dosage and/or time of analysis, and warrants further investigation.

Pten^{+/R173C} mice display mild behavioural aberrations
ASD is common in PHTS (Yehia et al., 2020). Behavioural abnormalities are found in mice with germline
expression of heterozygous loss-of-function of *Pten* (Clipperton-Allen and Page, 2014) or homozygous
expression of the nuclear-excluded *Pten^{m3m4}* (Tilot et al., 2014). To examine such type of aberrations
in *Pten^{+/R173C}* mice, we used a range of tests associated with ASD-like mouse behaviours.

In Crawley's sociability test, *Pten^{+/R173C}* mice were indistinguishable from *Pten^{+/+}* littermate controls,
indicating normal social interactions (Fig. S16A-D). In the open field test, male mice were not
significantly different from their *Pten^{+/+}* littermates with regards to distance travelled in the open field
arena and time spent moving during the test (Fig. S17B,C). However, female *Pten^{+/R173C}* mice travelled
less and spent less time moving than WT controls (Fig. S17F,G), suggesting decreased exploratory
behaviour and/or decreased locomotion. This was not caused by increased anxiety since female (and
male) mice spent equal length of time exploring the different compartments of the open field test as
their WT controls (Fig. S17D,H), with defecation during the test being comparable between the two
groups (Fig. S17E,I). Reduced exploratory locomotion in females was also observed in a hole board
test (Fig. S18A). Changes in repetitive behaviours, as assayed in a marble burying test were absent in
both males and females (Fig. S18B).

DISCUSSION
Since the discovery of PTEN as a tumour suppressor in somatic cancer, multiple genetic mouse models
targeting *Pten* have been generated, highlighting the importance of PIP$_3$ phosphatase activity of PTEN
in tumour suppression, embryonic development, cellular homeostasis, neuronal development and the
development of autism spectrum traits (Lee and Pandolfi, 2020).

PTEN mouse models have also improved our understanding of PHTS, a rare genetic condition
caused due to heterozygous germline *PTEN* mutations. These include *Pten^{+/-}* mice which represent the
heterozygous genetic constellation of *PTEN* in PHTS and on a mixed C57BL6xSv129 background
develop multiple phenotypes overlapping with the clinical presentation of PHTS, including brain and
lymphoid overgrowth and the development of specific cancers (Tibarewal et al., 2022). While the PIP$_3$
phosphatase activity of PTEN has been the focus of research over the last three decades, there is an
increasing body of evidence for other PTEN functions in disease pathogenesis, in particular its nuclear
function (Langdon, 2023). Recently, the *Pten^{m3m4}* mouse model with germline expression of nuclear-
excluded, engineered non-naturally-occurring *Pten* mutations, has indicated a role for nuclear PTEN
defects in driving neuronal and immune phenotypes in PHTS (Tilot et al., 2014, Jaini et al., 2020).

In this study, we developed a new mouse model featuring the pathogenic PTEN-R173C variant
commonly found in PHTS and somatic cancer. We report that this PTEN variant retains its ability to
regulate PI3K/AKT signalling but is impaired in dsDNA damage repair through HR, likely due to nuclear
exclusion. Contrary to previous reports (Han et al., 2000, Smith and Briggs, 2016, Post et al., 2020), we
found that PTEN-R173 mutant proteins (R173C and R173H) were catalytically more active PIP$_3$ lipid

phosphatases than PTEN-WT, both in cells and tissues. However, these mutants displayed reduced
protein stability, possibly resulting from a more open protein configuration that is more susceptible to
ubiquitination-mediated degradation. Upon heterozygous germline expression in $Pten^{+/R173C}$ mice, this
biochemical feature results in PTEN protein expression levels comparable to those in $Pten^{+/-}$ mice but
an overall lipid phosphatase activity similar to wild-type mice, as demonstrated by effective
downregulation of insulin-stimulated PI3K/AKT signalling.

$Pten^{+/-}$ mice with heterozygous PTEN loss are predisposed to a range of tumours similar to those
observed in PHTS patients (Tibarewal et al., 2022). To assess the importance of nuclear functions of
PTEN in tumourigenesis, we aged $Pten^{+/R173C}$ mice and found only a modest predisposition to solid
tumour development compared to $Pten^{+/-}$ mice. This is reminiscent of mouse models of genes involved
in dsDNA damage repair via HR such as *Brca*, *Rad51* and *Palb2* (Matos-Rodrigues et al., 2021, Tsuzuki
et al., 1996, Roy et al., 2011, Rantakari et al., 2010, Yan et al., 2004, Hakem et al., 1996). Indeed, while
homozygous inactivation of these genes frequently leads to embryonic lethality, as also observed in
$Pten^{R173C/R173C}$ mice, their heterozygous inactivation only leads to a very low to absent tumour burden,
most likely because the lifespan of mice is insufficiently long to accumulate the additional mutations
required for tumourigenesis (Drost and Jonkers, 2009). That additional genetic defects may be
required to allow phenotypic output of PTEN-R173 defects is indicated by the observation that somatic
cancers with PTEN-R173 mutation have a range of co-occurring mutations, including but not limited
to oncogenic mutation in *PIK3CA* or loss of second allele of *PTEN* (**Fig. 3D**). Also in mouse models,
combination of *Pten* mutations with other oncogenic drivers such as mutant *Pik3ca* and *ErbB2* have
accelerated development of mammary tumours (Kato et al., 2020, Yip et al., 2020). Combining the
Pten-R173C mutation in mouse models carrying additional cancer drivers (such as *PIK3CA* mutation)
or defects in DNA repair genes (such as *Brca* or *Rad51*) could provide evidence for this speculation, but
such experiments are outside the scope of the current study.

Our data further show that, in a germline configuration, deficient nuclear localisation of PTEN, and
consequently its nuclear function, correlates with brain and lymphoid overgrowth. While our study
does not establish a direct correlation between these overgrowth phenotypes in our mice and dsDNA
damage repair, it is possible that such a link exists. Indeed, one study reported increased IR-induced
DNA strand breaks in lymphocytes from autistic children compared to non-autistic controls supporting
such a functional link (Attia et al., 2020). On the other hand, lymphoid cells isolated from PHTS patients
with PTEN-R173C mutation, who all present with ASD, did not exhibit increased DNA damage following
IR treatment compared to wild-type controls (Wei et al., 2024), suggesting that nuclear functions of
PTEN other than dsDNA repair may underlie the observed lymphoid overgrowth and macrocephaly in
$Pten^{+/R173C}$ mice. This is further supported by the absence of lymphoid overgrowth and macrocephaly
in patients and mouse models with germline mutations in other genes regulating dsDNA damage
repair by HR, such as the *Brca*, *Rad51* and *PalB2* genes.

We find that the macrocephalic $Pten^{+/R173C}$ phenotype correlates with alterations in multiple
parameters, including increases in neuronal cell body size, axonal thickness, axonal complexity and
oligodendrocyte lineage cell numbers. While this brain overgrowth phenotype in $Pten^{+/R173C}$ mice is
very clear, its impact on animal behaviour is less prominent. It has to be said that while mice with
heterozygous germline expression of PTEN accurately reflect the genetic makeup of patients with
PHTS, the behavioural abnormalities observed in heterozygous mutant mice are often subtle and
highly variable (Clipperton-Allen and Page, 2014). This necessitates large mouse numbers to achieve
quantifiable significant differences between genotypes. Consequently, the field of behavioural analysis
has mostly relied on the use homozygous mouse models. This approach is particularly true for PHTS,
with numerous studies conducted on homozygous brain-specific PTEN knockout models (Kwon et al.,
2006, Zhou et al., 2009) and homozygous $Pten^{m3m4}$ models (Tilot et al., 2014). Similar homozygous
models are also utilized in behavioural research on other rare diseases such as Tuberous Sclerosis
(Yuan et al., 2012, Bateup et al., 2013). Although these models do not share the exact genetic
composition of patients, they serve as platforms for behavioural and neurological studies and testing
therapeutics.

Our study suggests that the status of nuclear function of PTEN alongside its PIP_3 phosphatase
activity, allows refinement of genotype/phenotype analysis in PHTS. Based on our findings and
assessment of patient phenotypes (Table 1 and Supplementary Tables 1 and 2), our data indicate that
individuals carrying *PTEN* variants such as *$PTEN^{R173C}$* that retain their PIP_3 phosphatase function but are
nuclear-excluded tend to follow a characteristic progression of known PHTS disease phenotypes: early-
signs such as neurological abnormalities – including macrocephaly and global developmental delay -
alongside cutaneous features which together lead to a PHTS diagnosis during childhood. As they enter
early adulthood, they may be prone to developing vascular abnormalities and benign tumours in the
thyroid and GI tract, with an increased likelihood of developing malignant cancers later in life.
Our study also highlights the possibility of new therapeutic treatments for cancer in PHTS patients.
A previous study has shown that glioma cells expressing PTEN-Y204F, a mutation that leads to lack of
PTEN nuclear localisation and functions such as dsDNA damage repair are more sensitive to radiation
therapy (Ma et al., 2019). It is therefore tempting to speculate that radiation therapy and dsDNA
damage repair antagonist drugs, such as *PARP* inhibitors, might be additional therapeutic options
beyond PI3K pathway inhibitors for cancer in PHTS patients with R173 mutant-like PTEN variants.
564

Methods

Plasmids. DNA constructs are detailed in Supplementary Table 8. Point mutations were introduced in
PTEN WT cDNA in different vectors using primers listed in Supplementary Table 9 and KOD hot start
DNA polymerase (Sigma-Aldrich) using the manufacturer's instructions.
Mice. All mice were maintained at University College London in accordance with the UK Animals
(Scientific Procedures) Act 1986 and following UK Home Office guidance. All procedures were
authorized by a UK Home Office Project Licence subject to local ethical review. *Pten*[+/-] mice
(Podsypanina et al., 1999), *Pten*[flox/flox] mice (Lesche et al., 2002), *CAG CreER*[T2] mice (Schonhuber et al.,
2014) and *Emx1-Cre* mice (Kessaris et al., 2006) have been described elsewhere. *Pten*[+/R173C] mice were
generated by Taconic Biosciences using a CRISPR/Cas9 approach. gRNA was designed to contain the
R173C mutation (c.517C>T) and a silent mutation containing the *Eco*NI restriction site upstream of the
R173 site (to allow discrimination of wild-type and knock-in allele following PCR). Fig. S6 shows the
gRNA sequence and the targeting strategy. gRNA was microinjected into C57BL/6J mouse zygotes
along with recombinant Cas9. The zygotes were then transferred to pseudo-pregnant mothers. The
litters born (founder animals) were sequenced to check for the incorporation of the R173C mutation
and percentage of mosaicism. They were then used to as a breeder to generate F1 animals. F1 males
positive for the R173C mutation and negative for any off-target mutations were used to generate
further animals by IVF using C57BL/6J females. All mouse lines were maintained on a C57BL/6J
background. Sv129 mice were purchased from Charles River or Envigo. *Pten*[+/-] and *Pten*[+/R173C] mice
were crossed with Sv129 mice to obtain mixed background F1 mice that were used for some
experiments as described. For neuronal studies, conditional knock-out mice that selectively lack PTEN
expression in cortical pyramidal neurons *were generated by crossing Pten*[flox/flox] mice with *Emx1-Cre*
mice to give *Emx1-Cre*;*Pten*[fl/fl]; referred to here as cKO. The cKO mice were crossed with *Pten*[+/R173C]
mice to obtain *Emx1-Cre*;*Pten*[fl/R173C] ,referred to here as cKI, which express one only copy of *Pten*-R173C
in the cortical pyramidal neurons.
For genotyping, mouse tissue samples (from ear biopsy or from the embryonic yolk sac) were lysed in
25 mM NaOH/0.2 mM disodium EDTA for 1 h at 95°C. The samples were neutralized by adding an
equal volume of a buffer containing 40 mM Tris-HCl, pH 4.5. Two µl of the DNA extract was used for
PCR using and Titanium polymerase PCR kit (Takara) and primers listed in Supplementary Table 10.
For genotyping of *Pten*[+/R173C] mice, an additional restriction digestion step was performed on the PCR
products using *Eco*NI (New England Biolabs).
Mouse survival studies. For sample size calculations for long term survival study, we use ClinCalc
(www.clincalc.com). The mean survival age of *Pten*[+/-] on a mixed C57BL6J x Sv129 background has been
reported to be 12±1.8 months (Alimonti et al., 2010). The sample size required for *Pten*[+/R173C] mice
needed to see a 15% improvement in mean survival age compared to *Pten*[+/-] mice was calculated as
16 (probability of Type-I error was set to 0.05 and Power was set to 80%). We used at least 16 mice
for each genotype and each sex. At least 10 Littermate wild type mice were used for both *Pten*[+/-] and
Pten[+/R173C] mice. For survival analysis, mice were weighed twice a week. They were monitored and
euthanized upon showing a more than 20% reduction in body weight, any signs of ill health such as
hunched appearance, laboured breathing, sluggish behaviour, or palpable masses with a total surface
area of 1.4 cm^2 surface area. Animals showing no clinical signs were euthanized at the end of the study
(600 days for females and 730 days for males).
Cell lines. U-87 MG (U87), HEK-293T and Phoenix cells were purchased from ATCC, Lenti-X™ 293T cells
were purchased from Clontech. Human PTEN knockout HeLa cell line and control HeLa cells was
obtained from Abcam (ab255419 and ab255928); PTEN knockout was validated using western blotting
(Fig. S4C). All cell lines were maintained in DMEM (Sigma-Aldrich), supplemented with 10% Filtered
Bovine Serum (FBS) (Pan-Biotech) and Pen-Strep (Sigma-Aldrich) using standard cell culture methods.
Preparation of MEFs. For preparation of E13.5 MEFs, timed matings were set up between *Pten*[+/-] mice
and *Pten*[+/R173C] mice, *Pten*[flox/+]*CreER*[T2] and *Pten*[+/-] or *Pten*[+/R173C] mice. Pregnant females were euthanized
on day 13.5 and embryos harvested from the uterus. Following removal of the head and visceral
organs, the rest of the embryo was minced in 2x Trypsin/EDTA using a sterile scalpel and incubated at

37°C for 40 min. The cell suspension was plated on 10 cm dishes in DMEM (Sigma-Aldrich) medium
supplemented with 10% FBS (Pan-Biotech) and Pen-Strep (Sigma-Aldrich), and cells maintained using
standard cell culture procedures. For preparation of E9.5 MEFs, timed intercrosses were set up
between *Pten*[+/R173C] mice. Pregnant females were euthanized on day 9.5, embryos removed from the
uterus and gently dissociated by pipetting up and down using a p1000 pipette and 2 ml of media
containing DMEM (Sigma-Aldrich), supplemented with 10% FBS (Pan-Biotech) and Pen-Strep (Sigma-
Aldrich). The cell suspension was plated in 48-well plates and cells maintained using standard cell
culture procedures. To induce Cre-mediated recombination of the floxed *Pten* allele in *Pten*[flox/+]*CreER*[T2]
MEFs, 4-hydroxy tamoxifen (Sigma Aldrich) was added to the cells at a final concentration of 1 μM and
refreshed every 24 h for 3 days.

Primary neuron culture. Primary cortico-hippocampal cultures were prepared from P1 or P2 pups
after genotyping. Cortex and hippocampus were dissected, digested with Trypsin, dissociated using
glass pipettes and 40,000 cells were plated per 12 mm coverslip. Neurons were cultured in serum-free
medium (neurobasal, NBA) medium supplemented with B27 supplement, L-glutamine,
penicillin/streptomycin and 10 mM HEPES pH 7.4. Cells were cultured at 5% CO_2 at 37°C with
replacement of half of the medium every 2-3 days. Cells were transfected on DIV6 with a GFP
expression vector using Lipofectamine 3000 according to the manufacturer's instructions. On DIV
12/13 cells were fixed with 4% PFA/4% sucrose in PBS for 10 min at room temperature (RT) and washed
3x in PBS. Cells were blocked and permeabilized in immunofluorescence (IF) buffer (PBS supplemented
with 0.3% Triton X and 3% BSA) for 20 min and incubated with primary antibodies to NeuN (Millipore
MAB377 and GFP, Abcam 1:1000) for 2 h at RT in IF buffer in a humidity chamber. After 3 washes with
PBS, cells were incubated with cross-absorbed Alexa Fluor labelled secondary antibodies (Invitrogen)
in IF buffer for 1 h at RT in a humidity chamber in the dark, washed 3x with PBS and mounted with
Immunomount. Neurons were imaged using a LSM 880 microscope (Leica) acquiring Z-stacks at 40 x
magnification. Dendritic complexity was determined using the Sholl macro in FIJI. Interceptions were
analysed in 10 μm radii within a 200 μm distance from the centre of the soma. To measure cell body
size in FIJI, stacks were processed to z-projections, ROIS were drawn manually around the cell bodies
and the ROI area was measured in FIJI.

PTEN activity assays. For purification of bacterially-expressed GST-tagged PTEN proteins, pGEX6P1
vectors with cDNA for WT or mutant PTEN were introduced in *E. coli* and recombinant PTEN was
purified by glutathione-affinity chromatography followed by cleavage of the GST tag as described
(Davidson et al., 2010). PIP_3 assays using purified recombinant PTEN were performed with diC8 PIP_3
(Cell Signals Inc) and malachite green phosphate assay kit (Sigma Aldrich) and have been described
elsewhere (Maehama and Dixon, 1998). Purification of recombinant PTEN protein from insect cells
was as described in Ref.(Masson et al., 2016b).

Lentiviruses and retroviruses. Lentiviruses for PTEN and PIP_3 biosensor were generated by co-
transfecting Lenti-X™ 293T cells or HEK-293T cells with plasmids expressing the cDNA of interest along
with packaging vectors using TransIT-LT1 following the manufacturer's instructions. For preparation
of retroviruses, Phoenix cells were transfected with p53 shRNA expressing vector (Dickins et al., 2005)
using TransIT-LT1 following the manufacturer's protocol. 24 h post transfection, sodium butyrate was
added to the cells to a final concentration of 12.5 mM for 6 h, followed by washing the cells with warm
PBS and addition of fresh media. The supernatant containing lentiviral particles was collected after 20
h, passed through a 0.45 μm filter and stored at -80°C. Target-cells for transductions were plated at
40-50% confluency. Once the cells had attached, lentiviral/retroviral particles were added to the cells
along with polybrene (Sigma-Aldrich) at 20 μg/μl. The media was changed 24 h post transduction.

Immunoblotting analysis. Protein extracts from cell cultures were prepared by scraping cells into lysis
buffer containing 25 mM Tris-HCl pH 7.4, 150 mM NaCl, 1% Triton X-100, 10% glycerol, 1 mM EGTA, 1
mM EDTA, 5 mM sodium pyrophosphate, 10 mM β-glycerophosphate, 50 mM sodium fluoride, 1 mM
sodium orthovanadate, 1 mM DTT and Protease inhibitor cocktail (Millipore). Mouse tissues harvested
from mice upon euthanasia were homogenize using Lysing Matrix M tubes (MP Biomedicals) in double
the volume (v/w) of lysis buffer (25 mM Tris-HCl pH 7.4, 150 mM NaCl, 1% Triton X-100, 0.1% SDS, 10%

glycerol, 1 mM EGTA, 1 mM EDTA, 10 mM sodium pyrophosphate, 20 mM β-glycerophosphate, 100
mM sodium fluoride, 2 mM sodium orthovanadate, 1 mM DTT and protease inhibitors) on a FastPrep
24 homogeniser (MP Biomedicals) at 4 m/sec for 20 sec. Cell or tissue lysates were pre-cleared by
centrifugation at 20,000xg for 10 min and protein extracts analysed by immunoblotting. 293T cells
expressing Rluc-PTEN-YFP constructs were lysed in buffer containing 50 mM HEPES (pH 7.4), 250 mM
NaCl, 2mM EDTA, 0.5% NP-40, 10% glycerol supplemented with protease and phosphatase inhibitors
(Roche) and protein A/G PLUS-Agarose (30μl/IP) and GFP antibody (Roche, Cat #11814460001, Mouse,
Dilution 1:200) was used to immunoprecipitate YFP-tagged PTEN protein which was used for
immunoblotting as described below. Protein gel electrophoresis was conducted with 10 μg of total
soluble protein per lane on NuPage Bis-Tris 4-12% gradient polyacrylamide gels (Thermofisher
Scientific) following the manufacturer's protocols. Proteins were transferred onto PVDF membrane
(Millipore) and membranes blocked in 5% milk powder/TBST for 1 h at RT. Blocked membranes were
incubated overnight with primary antibodies (**Supplementary Table 11**). Antibody complexes were
detected by 1 h incubation at RT with HRP-conjugated secondary antibodies (GE healthcare). Blots
were developed with Immobilon Forte Western HRP substrate (Millipore) and chemiluminescence
imaged using an ImageQuant LAS4000 imaging system (GE healthcare).
qRT-PCR assays. Total RNA was extracted from U87cells non-transduced or transduced with
lentiviruses for PTEN-WT, PTEN-C124S, PTEN-R173C or PTEN-R173H, using RNAEasy kit (Qiagen)
following the manufacturer's instructions. 1 μg of RNA was converted to cDNA using iScript RT kit
(BioRad). 50 ng of cDNA was used for QRT-PCR reactions using primers in **Supplementary Table 12**.
The reactions were run on a QuantStudio™ 5 Real-Time PCR machine (Applied Biosystems). dCT was
calculated by deducting the CT for GAPDH from CT of PTEN. ddCT was calculated by deducting the dCT
of untransduced cells from the dCT of the sample of interest.
Cycloheximide chase studies. U87 cells were transduced with lentiviruses for PTEN-WT, PTEN-C124S,
PTEN-R173C or PTEN-D252G. 48 h post transduction, cells were treated with 200 μg/ml cycloheximide,
lysed at different time points and protein extracts analysed for PTEN expression by immunoblotting.
BRET assays. BRET assays using the Rluc-PTEN-YFP biosensor were performed as described in Refs
(Misticone et al., 2016, Lima-Fernandes et al., 2014). Briefly, 24 h post-transfection, cells were
detached from 12-well plates, using trypsin-EDTA, and resuspended in 1 ml of complete media. The
cells were subsequently distributed into poly-L-ornithine-coated (30 μg/ml) white 96-well optiplates
(Perkin Elmer). The next day media was replaced with Opti-MEM (Gibco) and Coelenterazine substrate
(Interchim) was added to a final concentration of 5 μM and incubated for 3 min at 25°C. BRET readings
were then collected using a Multilabel Reader Mithras2 LB 943 (Berthold Technologies). Sequential
integration of light output was measured for 1 sec using two filter settings (480+10nm for YFP and
540+20nm for Rluc). The BRET signal represents the ratio of the light emitted by YFP and the light
emitted by Rluc (YFP/Rluc). The ratio values were corrected by subtracting background BRET signals
obtained with Rluc-PTEN. MilliBRET (mBRET) values were calculated by multiplying these ratios by
1,000. Data are represented as specific mBRET values or ΔmBRET when compared to the signal
obtained with WT PTEN set to 0.
MTS assays. U87 cells were transduced with lentiviruses for GFP, PTEN-WT, PTEN-C124S, PTEN-R173C
or PTEN-R173H. 48 h post transduction, 5000 cells/well were plated in 96-well plates. A different plate
was prepared for each day. 20 μl of MTS assay reagent (Abcam) was added to the cells, incubated at
37°C for 1 h, followed by measurement of absorbance at 490 nm using a plate reader.
PIP$_3$ biosensor analysis and PTEN localization using fluorescence microscopy. For plasma membrane
PIP$_3$ quantification, U87-MG cells were transduced with lentiviral particles expressing the PIP$_3$
biosensor on its own or with lentiviral particles for PTEN-WT or PTEN-R173C, were pretreated $\pm$ GDC-
0941 (1 μM) for 1 h and stimulated $\pm$ insulin (100 nM) for 2 min. For PTEN localization studies, MEFs
were treated $\pm$ GDC-0941 (1 μM) 1 h prior to fixation. For both plasma membrane PIP$_3$ quantification
and localization of untagged or endogenous PTEN, cells were fixed with 4% PFA for 20 min, washed in
PBS, permeabilized with 0.1% Triton X-100 in PBS for 90 sec, washed 3 times with PBS and blocked
with PBS containing 1 or 3% (w/v) BSA for 30 min. Primary antibodies to PTEN were diluted in 1 or 3%

BSA block solution and incubated for 1 h at RT. Following 3 washes with PBS, secondary antibodies and
Phalloidin were diluted in 1 or 3% BSA block solution and incubated for 45 min at RT. Cells were then
washed 3 times and mounted using ProLong Gold antifade reagent containing DAPI (P36935, Life
Technologies). To examine GFP-PTEN localization during different stages of the cell cycle, Lenti-X™
293T cells were transfected with DNA for PTEN-WT-GFP or PTEN-R173C-GFP. 18 and 36 h post
transfection, culture medium was changed to fresh media containing 2 mM thymidine (Sigma). 18 h
post the second media change, the cells were released from double thymidine block by changing into
fresh, complete media. Cells were fixed with 4% PFA for 15 min, washed in 3 times in PBS and
permeabilized with 0.1% Triton X-100 in PBS for 5 min. Cells were then washed 3 times with PBS and
once with water before mounting using ProLong Gold antifade reagent containing DAPI. Confocal
microscopy was performed using a Zeiss LSM 880 microscope with AiryScan and a 63x PL APO objective
and Zen Black acquisition software or Zeiss LSM900 microscope and a 63x or 40x PL APO objective and
Zen Black acquisition software. Image processing was performed using Image J software (National
Institutes of Health, Rockville, MD, United States) and was limited to alterations of brightness,
subjected to the entire image. Quantification of plasma membrane PIP_3 intensity was performed using
Image J as described previously with minor modifications (Ooms et al., 2015). The average
fluorescence intensity was measured within a box of defined size (20 x 10 pixels) at three random
regions of the plasma membrane and three random regions of the cytosol (box size 20 x 20 pixels).
The plasma membrane and cytosol fluorescence intensity measurement were each averaged and the
ratio of plasma membrane to cytosolic fluorescence intensity determined. The PTEN nuclear to
cytoplasmic ratio was measured using Image J. First, the nuclear PTEN fluorescence intensity was
measured, followed by the cytoplasmic PTEN fluorescence intensity. The ratio of nuclear to
cytoplasmic PTEN fluorescence intensity was then calculated.

Insulin tolerance test (ITT). 3-month-old male $Pten^{+/-}$ and $Pten^{+/R173C}$ mice and their WT littermates
were starved for 6 h. The tip of the tail was punctured using a 26G needle and time-0 blood glucose
levels were recorded using a standard glucometer. The mice were given an IP injection of insulin to a
final concentration of 0.75 U/kg. Blood glucose readings were recorded at 15 min, 30 min, 45 min, 60
min, 90 min and 120 min post injection.

Irradiation of mice: 8-12-week-old mice were subjected to whole body irradiation of 7 Gy dose using
the Small Animal Radiation Therapy Platform (SARRP). 5 h post irradiation, the mice were euthanised
with a lethal injection of pentobarbital. Tissues of interest were harvested and snap frozen in liquid
nitrogen for biochemical analysis.

Flow cytometry. Mouse spleens and lymph nodes were mashed with a syringe plunger in RPMI
containing 10% FBS and filtered through a 70 µm filter. Cell suspensions from spleen were treated with
RBC lysis buffer (Biolegend). The cell suspensions were washed with PBS and stained with Fixable
Viability Dye eFluor™ 780 (ThermoFisher scientific) in PBS for 40 min followed by a few PBS washes.
For staining of surface antigens, antibodies were diluted in FACS staining buffer at 1:100 and added to
the cells for 30 min followed by two washes with FACS buffer. Details on antibodies and reagents used
is shown in **Supplementary Table 11**. The stained cells were analysed on BD FACSymphony™ analyser.

Tissue processing for histological analysis and immunohistochemical analysis. Mouse tissues were
fixed in 10% neutral buffered formalin for 24 h, followed by processing and embedding into paraffin
blocks. The tissues were orientated with the biggest surface down, to obtain complete sections. 3 µm
serial sections were cut and stained for Haematoxylin and Eosin, followed by immunohistochemistry
staining on the Leica BOND Rxm. Tissue sections for immunohistochemistry were incubated for 40 min
at 100°C in BOND epitope retrieval solution 2 (Leica, Germany), followed by blocking and staining using
the BOND polymer refine detection kit (Leica, DS9800), primary antibody (**Supplementary Table 11**)
and DAB enhancer (Leica, Germany).

Histopathological analysis. Non-neoplastic lesions, e.g. splenic extramedullary haematopoiesis and
lymph node hyperplasia, were graded using a standard semi-quantitative grading scheme from 0 to 5,
where 0 = lesion not present and 5 = the maximum lesion size/extent (Scudamore, 2014). In general,
where present, the presence of other proliferative lesions including hyperplasia and neoplasia was

recorded for each individual organ, allowing the overall incidence to be established for each lesion in
each group. Prostatic intraepithelial neoplasia (PIN) was graded as low or high grade according to the
consensus suggested by Ittmann *et al.* (Ittmann et al., 2013). Low-grade PIN lesions are focal, with one
to two layers of cells and mild nuclear atypia. High-grade lesions tend to be more extensive, filling the
prostatic lumen but without stromal invasion; have two or more cell layers often forming papillary or
cribriform patterns; increased nuclear atypia and mitoses.
Perfusion and preparation of brain sections for DAB immunostaining and immunofluorescence. Mice
were perfused with saline (0.9 % NaCl) followed by 4% PFA (w/v) in PBS through the left ventricle of
the heart. Brains were removed and post-fixed overnight at 4°C in 4% PFA, cryoprotected in 20%
sucrose in PBS for 24 h at 4°C and embedded in Tissue-Tek OCT compound (Sakura) in isopentane
cooled on dry ice. Frozen brains were stored at -80°C until use. Brains were processed on a Cryostat
(30 μm thickness) and sections were collected in PBS for immunohistochemistry. For DAB
immunostaining, floating sections were incubated with 0.3 % H_2O_2 in PBS for 30 min while shaking at
RT, and then blocked using blocking buffer (10% FBS in PBS supplemented with 0.1% Triton X-100) for
1 h at RT. After blocking, sections were incubated overnight at 4°C with primary antibodies diluted in
blocking buffer. Primary antibodies were detected with biotinylated secondary antibodies (Donkey
Anti-Mouse IgG Biotinylated Antibody, R&D systems) diluted 1:300 in blocking buffer for 1 h at RT. The
signal was amplified using the Vectastain ABC Kit HRP kit (Vector, PK-6100), according to the
manufacturer's instructions and developed using a DAB Peroxidase (HRP) Substrate Kit (Vector, SK-
4100) until the desired staining intensity was reached. Floating sections were mounted on Superfrost
plus slides in 0.2 % pork skin gelatine dissolved in 50 mM Tris-HCl pH 7.4, dried at RT overnight and
dehydrated in ethanol and xylene before mounting with DPX mounting medium (Merck). Images were
acquired using 20x magnification on a ZEISS Axio Scan.Z1 microscope and processed using ZEISS ZEN
lite software. For immunofluorescence, sectioned brains were blocked for 1 h before primary
antibodies were applied overnight. AlexaFluor 488 secondary antibody was used to detect OLIG2
(1:1000; Invitrogen, Carlsbad, CA) for 60 min at room temperature together with Hoechst 33258
(1:1000; Sigma) to detect cell nuclei. All secondary antibodies were diluted in block solution. Floating
sections were transferred onto Superfrost plus slides (BDH Laboratory Supplies) and air dried before
being coverslipped with Dako fluorescent mounting medium.
Analysis of corpus callosum and cortex thickness and cell counting. Matched sections at specified
Bregma levels were measured using the Zeiss Blue software measurement length tool. Counting boxes
were positioned onto matched slices in the barrel field of the primary somatosensory cortex
perpendicular to the corpus callosum. In each box, cells were counted from the border of the corpus
callosum to the pial surface. The length of the box was adjusted to fit the cortical thickness while the
width was not changed. All cells positive for NeuN within the box were counted and cell numbers
displayed as crude numbers in the box as well as density considering the area of the boxes analysed.
Four animals per genotype were used.
Electron microscopy (EM). EM was performed as described in Ref.(Kougioumtzidou et al., 2017).
Briefly, mice were prepared for EM by PBS perfusion followed by 2.5% (v/v) glutaraldehyde and 2%
(w/v) paraformaldehyde in 0.1 M sodium cacodylate buffer (pH 7.6). Brains were removed and kept
at 4°C for 48 h in the same fixative, before being washed with 0.1 M cacodylate buffer, rinsed briefly
with 0.1 M phosphate buffer and shipped to Japan at ambient temperature. Sagitally-sectioned brains
were processed for EM. After immersion in a 1% (w/v) osmium tetroxide solution for 2 h at 4°C, the
specimens were dehydrated through a graded alcohol series and embedded in Epon 812 (TAAB
Laboratories, UK). The required area was trimmed, and ultrathin sections were cut, collected on a
platinum-coated glass slide, stained with uranyl acetate and lead citrate and imaged in a scanning
electron microscope equipped with a back-scattered electron beam detector (Hitachi SU8010) at 1.5
kV accelerating voltage. EM images of sagittal sections of corpus callosum were taken at 15000x
magnification. To estimate g-ratios, the circumference of the axon and the external circumference of
the myelin sheath were measured using ImageJ, avoiding paranodal areas. The diameters of axon and

myelin sheath were calculated from these circumferential measurements, assuming a circular profile,
and the g-ratio was calculated.
Behavioural analysis. Behaviour testing was carried out as previously described (Magno et al., 2021).
12-week-old mice on mixed C57BL/6 x Sv129 background were used in this study. All animals were
kept on a 12 h light-dark cycle with food and water accessible ad libitum. The indicated numbers of
WT mice and their *Pten*[+/R173C] littermates were tested in the open field test, three chambers test, hole
board and marble burying. Mice tested in behavioural experiments were habituated to handling and
the experimenter for 9 days before the start of behavioural experiments to reduce stress and
habituated mice were generally handled by cupping. Mice were analysed in three cohorts and male
and female mice were tested on separate days. During experiments white noise was played at around
50 Db to mask external noises and all experiments were video recorded. Before every experiment,
mice were brought into the behavioural room 30 min before the first experiment for habituation and
animals that had finished a test were transferred into a new cage in order not to influence the
behaviour of the remaining experimental animals. Directly before the first experimental animal was
tested and after every mouse the set up (open field, sociability chamber, hole board) was cleaned
thoroughly with water and 70% ethanol in a consistent manner for every animal. Open field: the
experimental animal was placed in an open field box measuring 30 cm x 30 cm. The walls were white
and 40 cm high so that the animal could not get any visual clues. The mouse was allowed to roam
freely for 30 min to explore the arena without any prior habituation. Distance, time spent in different
areas and the time moving were analysed using Ethovision software while faecal boli were counted
manually. Sociability: The experimental animal was placed into the central compartment of Crawley's
chamber for 5 min with closed doors for habituation. After 5 min empty cages were placed into the
middle of the left and right compartments and a stranger mouse was gently put into one of the cages
(counterbalance left and right sides). The stranger mouse matched the subject mouse with regards to
the breeding background, age and gender and had no contact with the subject mouse ever before.
Both doors were opened at the same time and the experimental mouse was allowed to explore the
area for 10 min. A stopwatch was used to determine the time the subject mouse spent closely
interacting with either cage (direct contact/sniffing/cage climbing). The time spent with the empty
cage versus the first stranger was measured, referred to as sociability. Calculation of discrimination
indices: (time spent with mouse – time spent with empty cage)/(sum of the time spent with both).
Hole board: since the experimental animals were familiar with the open field arena from the open field
experiment, they were only briefly re-exposed to the empty arena for 10 min on day 1 of the hole
board test. On the next day a hole board with 16 holes evenly spaced was placed into the arena, the
subject mouse was put into the middle of the field and was recorded for 15 min. A manual cell counter
was used to determine the total number of head dips into the holes. Marble burying: a cage was filled
with 5cm of bedding that was tamped down lightly to produce an even surface. 12 glass marbles were
distributed evenly spaced out in the cage and the animal was allowed to roam freely for 15 min.
Marbles that were buried to 2/3 of their depth with bedding were counted.
Statistical analysis: GraphPad Prism (San Diego, CA, United States) was used for statistical analysis.
The method used for individual datasets is indicated in the figure legends.
859

860
ACKNOWLEDGEMENTS. Personal funding was from the Jean Shanks Pathological Society (Clinical PhD
fellowship; to V.R.), CRUK UCL Centre Non-Clinical Training Awards (C416/A29287 to G.C. and
CANTAC721\100022 to F.B.) and the European Commission (H2020-MSCA-IF-2018 GA: 838559; to
S.C.). Research in the B.V. laboratory was supported by PTEN Research (UCL-16-001, UCL-20-001),
Cancer Research UK (C23338/A25722) and the UK Biotechnology and Biological Sciences Research
Council (BB/W007460/1). Research in N.R.L. laboratory was supported by PTEN Research. Research in
the M.G.H.S. group was supported by Fondation ARC pour la recherche sur le cancer
(ARCPJA2022060005118) and Ligue Contre le Cancer (Comité de Paris). The UCL Cancer Institute
Translational Technology Platforms are supported by the CRUK UCL Centre Award (C416/A25145). The
N.K. lab is funded by grants from the UK Biotechnology and Biological Sciences Research Council
(BB/N009061/1) and the Wellcome Trust (108726/Z/15/Z). M.T. and K.S. are supported by PTEN
Research and the NIHR Cambridge Biomedical Research Centre (NIHR203312). The PHTS Patient
Registry UK is funded by PTEN Research (UOC-17-001). We thank Dr Isra Mohamed, Cambridge
University NHS Foundation Trust for providing information about patient P021. We thank staff at the
Technical Support Centre for Life Science Research in IMU for excellent technical support with EM
processing. We also thank the UCL Research Capital Infrastructure Fund (RCIF) and the National
Institute for Health Research University College London Hospitals Biomedical Research Centre for
upgrade of the UCL Cancer Institute Microscopy facility and Dr Maria Whitehead and Dr Paul Elvin and
members from the B.V. group for excellent feedback on the manuscript.
880

REFERENCES

ALIMONTI, A., CARRACEDO, A., CLOHESSY, J. G., TROTMAN, L. C., NARDELLA, C., EGIA, A., SALMENA, L., SAMPIERI, K., HAVEMAN, W. J., BROGI, E., RICHARDSON, A. L., ZHANG, J. & PANDOLFI, P. P. 2010. Subtle variations in Pten dose determine cancer susceptibility. *Nat Genet,* 42**,** 454-8.

ALVAREZ-GARCIA, V., TAWIL, Y., WISE, H. M. & LESLIE, N. R. 2019. Mechanisms of PTEN loss in cancer: It's all about diversity. *Semin Cancer Biol,* 59**,** 66-79.

ATTIA, S. M., AL-HAMAMAH, M. A., AHMAD, S. F., NADEEM, A., ATTIA, M. S. M., ANSARI, M. A., BAKHEET, S. A. & AL-AYADHI, L. Y. 2020. Evaluation of DNA repair efficiency in autistic children by molecular cytogenetic analysis and transcriptome profiling. *DNA Repair (Amst),* 85**,** 102750.

BASSI, C., HO, J., SRIKUMAR, T., DOWLING, R. J., GORRINI, C., MILLER, S. J., MAK, T. W., NEEL, B. G., RAUGHT, B. & STAMBOLIC, V. 2013. Nuclear PTEN controls DNA repair and sensitivity to genotoxic stress. *Science,* 341**,** 395-9.

BATEUP, H. S., JOHNSON, C. A., DENEFRIO, C. L., SAULNIER, J. L., KORNACKER, K. & SABATINI, B. L. 2013. Excitatory/inhibitory synaptic imbalance leads to hippocampal hyperexcitability in mouse models of tuberous sclerosis. *Neuron,* 78**,** 510-22.

BIRLE, D., BOTTINI, N., WILLIAMS, S., HUYNH, H., DEBELLE, I., ADAMSON, E. & MUSTELIN, T. 2002. Negative feedback regulation of the tumor suppressor PTEN by phosphoinositide-induced serine phosphorylation. *J Immunol,* 169**,** 286-91.

CASERTA, E., EGRIBOZ, O., WANG, H., MARTIN, C., KOIVISTO, C., PECOT, T., KLADNEY, R. D., SHEN, C., SHIM, K. S., PHAM, T., KARIKOMI, M. K., MAUNTEL, M. J., MAJUMDER, S., CUITINO, M. C., TANG, X., SRIVASTAVA, A., YU, L., WALLACE, J., MO, X., PARK, M., FERNANDEZ, S. A., PILARSKI, R., LA PERLE, K. M., ROSOL, T. J., COPPOLA, V., CASTRILLON, D. H., TIMMERS, C., COHN, D. E., O'MALLEY, D. M., BACKES, F., SUAREZ, A. A., GOODFELLOW, P., CHAMBERLIN, H. M., MACRAE, E. R., SHAPIRO, C. L., OSTROWSKI, M. C. & LEONE, G. 2015. Noncatalytic PTEN missense mutation predisposes to organ-selective cancer development in vivo. *Genes Dev,* 29**,** 1707-20.

CHEN, C. Y., CHEN, J., HE, L. & STILES, B. L. 2018. PTEN: Tumor Suppressor and Metabolic Regulator. *Front Endocrinol (Lausanne),* 9**,** 338.

CHEN, H. H., HANDEL, N., NGEOW, J., MULLER, J., HUHN, M., YANG, H. T., HEINDL, M., BERBERS, R. M., HEGAZY, A. N., KIONKE, J., YEHIA, L., SACK, U., BLASER, F., RENSING-EHL, A., REIFENBERGER, J., KEITH, J., TRAVIS, S., MERKENSCHLAGER, A., KIESS, W., WITTEKIND, C., WALKER, L., EHL, S., ARETZ, S., DUSTIN, M. L., ENG, C., POWRIE, F. & UHLIG, H. H. 2017. Immune dysregulation in patients with PTEN hamartoma tumor syndrome: Analysis of FOXP3 regulatory T cells. *J Allergy Clin Immunol,* 139**,** 607-620 e15.

CLIPPERTON-ALLEN, A. E. & PAGE, D. T. 2014. Pten haploinsufficient mice show broad brain overgrowth but selective impairments in autism-relevant behavioral tests. *Hum Mol Genet,* 23**,** 3490-505.

DAVIDSON, L., MACCARIO, H., PERERA, N. M., YANG, X., SPINELLI, L., TIBAREWAL, P., GLANCY, B., GRAY, A., WEIJER, C. J., DOWNES, C. P. & LESLIE, N. R. 2010. Suppression of cellular proliferation and invasion by the concerted lipid and protein phosphatase activities of PTEN. *Oncogene,* 29**,** 687-97.

DEMPSEY, D. R., VIENNET, T., IWASE, R., PARK, E., HENRIQUEZ, S., CHEN, Z., JELIAZKOV, J. R., PALANSKI, B. A., PHAN, K. L., COOTE, P., GRAY, J. J., ECK, M. J., GABELLI, S. B., ARTHANARI, H. & COLE, P. A. 2021. The structural basis of PTEN regulation by multi-site phosphorylation. *Nat Struct Mol Biol,* 28**,** 858-868.

DI CRISTOFANO, A., DE ACETIS, M., KOFF, A., CORDON-CARDO, C. & PANDOLFI, P. P. 2001. Pten and p27KIP1 cooperate in prostate cancer tumor suppression in the mouse. *Nat Genet,* 27**,** 222-4.

DI CRISTOFANO, A., PESCE, B., CORDON-CARDO, C. & PANDOLFI, P. P. 1998. Pten is essential for embryonic development and tumour suppression. *Nat Genet,* 19**,** 348-55.

DICKINS, R. A., HEMANN, M. T., ZILFOU, J. T., SIMPSON, D. R., IBARRA, I., HANNON, G. J. & LOWE, S.
W. 2005. Probing tumor phenotypes using stable and regulated synthetic microRNA
precursors. *Nat Genet,* 37**,** 1289-95.
DRISSEN, M., VOS, J. R., COLLADO CAMPS, E., SCHUURS-HOEIJMAKERS, J. H. M., SCHIEVING, J. H. &
HOOGERBRUGGE, N. 2024. Increased frequency of infections and autoimmune disease in
adults with PTEN Hamartoma Tumour Syndrome. *Eur J Med Genet,* 70**,** 104960.
DROST, R. M. & JONKERS, J. 2009. Preclinical mouse models for BRCA1-associated breast cancer. *Br J*
Cancer, 101**,** 1651-7.
FREEMAN, D., LESCHE, R., KERTESZ, N., WANG, S., LI, G., GAO, J., GROSZER, M., MARTINEZ-DIAZ, H.,
ROZENGURT, N., THOMAS, G., LIU, X. & WU, H. 2006. Genetic background controls tumor
development in PTEN-deficient mice. *Cancer Res,* 66**,** 6492-6.
GARCIA-CAO, I., SONG, M. S., HOBBS, R. M., LAURENT, G., GIORGI, C., DE BOER, V. C., ANASTASIOU, D.,
ITO, K., SASAKI, A. T., RAMEH, L., CARRACEDO, A., VANDER HEIDEN, M. G., CANTLEY, L. C.,
PINTON, P., HAIGIS, M. C. & PANDOLFI, P. P. 2012. Systemic elevation of PTEN induces a tumor-
suppressive metabolic state. *Cell,* 149**,** 49-62.
GUPTA, A. & LESLIE, N. R. 2016. Controlling PTEN (Phosphatase and Tensin Homolog) Stability: A
DOMINANT ROLE FOR LYSINE 66. *J Biol Chem,* 291**,** 18465-73.
HAKEM, R., DE LA POMPA, J. L., SIRARD, C., MO, R., WOO, M., HAKEM, A., WAKEHAM, A., POTTER, J.,
REITMAIR, A., BILLIA, F., FIRPO, E., HUI, C. C., ROBERTS, J., ROSSANT, J. & MAK, T. W. 1996. The
tumor suppressor gene Brca1 is required for embryonic cellular proliferation in the mouse.
Cell, 85**,** 1009-23.
HAN, S. Y., KATO, H., KATO, S., SUZUKI, T., SHIBATA, H., ISHII, S., SHIIBA, K., MATSUNO, S., KANAMARU,
R. & ISHIOKA, C. 2000. Functional evaluation of PTEN missense mutations using in vitro
phosphoinositide phosphatase assay. *Cancer Res,* 60**,** 3147-51.
HENDRICKS, L. A. J., HOOGERBRUGGE, N., VENSELAAR, H., ARETZ, S., SPIER, I., LEGIUS, E., BREMS, H.,
DE PUTTER, R., CLAES, K. B. M., EVANS, D. G., WOODWARD, E. R., GENUARDI, M.,
BRUGNOLETTI, F., VAN IERLAND, Y., DIJKE, K., THAM, E., TESI, B., SCHUURS-HOEIJMAKERS, J.
H. M., BRANCHAUD, M., SALVADOR, H., JAHN, A., SCHNAITER, S., ANASTASIADOU, V. C.,
BRUNET, J., OLIVEIRA, C., ROHT, L., BLATNIK, A., IRMEJS, A., GROUP, P. S., MENSENKAMP, A.
R. & VOS, J. R. 2022. Genotype-phenotype associations in a large PTEN Hamartoma Tumor
Syndrome (PHTS) patient cohort. *Eur J Med Genet,* 65**,** 104632.
HO, J., CRUISE, E. S., DOWLING, R. J. O. & STAMBOLIC, V. 2019. PTEN Nuclear Functions. *Cold Spring*
Harb Perspect Med.
IGARASHI, A., ITOH, K., YAMADA, T., ADACHI, Y., KATO, T., MURATA, D., SESAKI, H. & IIJIMA, M. 2018.
Nuclear PTEN deficiency causes microcephaly with decreased neuronal soma size and
increased seizure susceptibility. *J Biol Chem,* 293**,** 9292-9300.
ITTMANN, M., HUANG, J., RADAELLI, E., MARTIN, P., SIGNORETTI, S., SULLIVAN, R., SIMONS, B. W.,
WARD, J. M., ROBINSON, B. D., CHU, G. C., LODA, M., THOMAS, G., BOROWSKY, A. & CARDIFF,
R. D. 2013. Animal models of human prostate cancer: the consensus report of the New York
meeting of the Mouse Models of Human Cancers Consortium Prostate Pathology Committee.
Cancer Res, 73**,** 2718-36.
JAINI, R., LOYA, M. G., KING, A. T., THACKER, S., SARN, N. B., YU, Q., STARK, G. R. & ENG, C. 2020.
Germline PTEN mutations are associated with a skewed peripheral immune repertoire in
humans and mice. *Hum Mol Genet,* 29**,** 2353-2364.
JUN, H. J., PAULO, J. A., APPLEMAN, V. A., YARON-BARIR, T. M., JOHNSON, J. L., YEO, A. T., ROGERS, V.
A., KUANG, S., VARMA, H., GYGI, S. P., TROTMAN, L. C. & CHAREST, A. 2024. Pleiotropic tumor
suppressive functions of PTEN missense mutations during gliomagenesis. *iScience,* 27**,** 111278.
KATO, T., YAMADA, T., NAKAMURA, H., IGARASHI, A., ANDERS, R. A., SESAKI, H. & IIJIMA, M. 2020. The
Loss of Nuclear PTEN Increases Tumorigenesis in a Preclinical Mouse Model for Hepatocellular
Carcinoma. *iScience,* 23**,** 101548.

KESSARIS, N., FOGARTY, M., IANNARELLI, P., GRIST, M., WEGNER, M. & RICHARDSON, W. D. 2006. Competing waves of oligodendrocytes in the forebrain and postnatal elimination of an embryonic lineage. *Nat Neurosci,* 9**,** 173-9.

KNOBBE, C. B., LAPIN, V., SUZUKI, A. & MAK, T. W. 2008. The roles of PTEN in development, physiology and tumorigenesis in mouse models: a tissue-by-tissue survey. *Oncogene,* 27**,** 5398-415.

KOUGIOUMTZIDOU, E., SHIMIZU, T., HAMILTON, N. B., TOHYAMA, K., SPRENGEL, R., MONYER, H., ATTWELL, D. & RICHARDSON, W. D. 2017. Signalling through AMPA receptors on oligodendrocyte precursors promotes myelination by enhancing oligodendrocyte survival. *Elife,* 6.

KWON, C. H., LUIKART, B. W., POWELL, C. M., ZHOU, J., MATHENY, S. A., ZHANG, W., LI, Y., BAKER, S. J. & PARADA, L. F. 2006. Pten regulates neuronal arborization and social interaction in mice. *Neuron,* 50**,** 377-88.

LACHLAN, K. L., LUCASSEN, A. M., BUNYAN, D. & TEMPLE, I. K. 2007. Cowden syndrome and Bannayan Riley Ruvalcaba syndrome represent one condition with variable expression and age-related penetrance: results of a clinical study of PTEN mutation carriers. *J Med Genet,* 44**,** 579-85.

LANGDON, C. G. 2023. Nuclear PTEN's Functions in Suppressing Tumorigenesis: Implications for Rare Cancers. *Biomolecules,* 13.

LEE, J. O., YANG, H., GEORGESCU, M. M., DI CRISTOFANO, A., MAEHAMA, T., SHI, Y., DIXON, J. E., PANDOLFI, P. & PAVLETICH, N. P. 1999. Crystal structure of the PTEN tumor suppressor: implications for its phosphoinositide phosphatase activity and membrane association. *Cell,* 99**,** 323-34.

LEE, Y. R. & PANDOLFI, P. P. 2020. PTEN Mouse Models of Cancer Initiation and Progression. *Cold Spring Harb Perspect Med,* 10.

LESCHE, R., GROSZER, M., GAO, J., WANG, Y., MESSING, A., SUN, H., LIU, X. & WU, H. 2002. Cre/loxP-mediated inactivation of the murine Pten tumor suppressor gene. *Genesis,* 32**,** 148-9.

LESLIE, N. R. & LONGY, M. 2016. Inherited PTEN mutations and the prediction of phenotype. *Semin Cell Dev Biol,* 52**,** 30-8.

LESLIE, N. R., MACCARIO, H., SPINELLI, L. & DAVIDSON, L. 2009. The significance of PTEN's protein phosphatase activity. *Adv Enzyme Regul,* 49**,** 190-6.

LI, J., YEN, C., LIAW, D., PODSYPANINA, K., BOSE, S., WANG, S. I., PUC, J., MILIARESIS, C., RODGERS, L., MCCOMBIE, R., BIGNER, S. H., GIOVANELLA, B. C., ITTMANN, M., TYCKO, B., HIBSHOOSH, H., WIGLER, M. H. & PARSONS, R. 1997. PTEN, a putative protein tyrosine phosphatase gene mutated in human brain, breast, and prostate cancer. *Science,* 275**,** 1943-7.

LIMA-FERNANDES, E., MISTICONE, S., BOULARAN, C., PARADIS, J. S., ENSLEN, H., ROUX, P. P., BOUVIER, M., BAILLIE, G. S., MARULLO, S. & SCOTT, M. G. 2014. A biosensor to monitor dynamic regulation and function of tumour suppressor PTEN in living cells. *Nat Commun,* 5**,** 4431.

MA, J., BENITEZ, J. A., LI, J., MIKI, S., PONTE DE ALBUQUERQUE, C., GALATRO, T., ORELLANA, L., ZANCA, C., REED, R., BOYER, A., KOGA, T., VARKI, N. M., FENTON, T. R., NAGAHASHI MARIE, S. K., LINDAHL, E., GAHMAN, T. C., SHIAU, A. K., ZHOU, H., DEGROOT, J., SULMAN, E. P., CAVENEE, W. K., KOLODNER, R. D., CHEN, C. C. & FURNARI, F. B. 2019. Inhibition of Nuclear PTEN Tyrosine Phosphorylation Enhances Glioma Radiation Sensitivity through Attenuated DNA Repair. *Cancer Cell,* 35**,** 504-518 e7.

MACCARIO, H., PERERA, N. M., GRAY, A., DOWNES, C. P. & LESLIE, N. R. 2010. Ubiquitination of PTEN (phosphatase and tensin homolog) inhibits phosphatase activity and is enhanced by membrane targeting and hyperosmotic stress. *J Biol Chem,* 285**,** 12620-8.

MAEHAMA, T. & DIXON, J. E. 1998. The tumor suppressor, PTEN/MMAC1, dephosphorylates the lipid second messenger, phosphatidylinositol 3,4,5-trisphosphate. *J Biol Chem,* 273**,** 13375-8.

MAGNO, L., ASGARIAN, Z., PENDOLINO, V., VELONA, T., MACKINTOSH, A., LEE, F., STRYJEWSKA, A., ZIMMER, C., GUILLEMOT, F., FARRANT, M., CLARK, B. & KESSARIS, N. 2021. Transient developmental imbalance of cortical interneuron subtypes presages long-term changes in behavior. *Cell Rep,* 35**,** 109249.

MARSH, D. J., COULON, V., LUNETTA, K. L., ROCCA-SERRA, P., DAHIA, P. L., ZHENG, Z., LIAW, D., CARON,
S., DUBOUE, B., LIN, A. Y., RICHARDSON, A. L., BONNETBLANC, J. M., BRESSIEUX, J. M.,
CABARROT-MOREAU, A., CHOMPRET, A., DEMANGE, L., EELES, R. A., YAHANDA, A. M.,
FEARON, E. R., FRICKER, J. P., GORLIN, R. J., HODGSON, S. V., HUSON, S., LACOMBE, D., ENG,
C. & ET AL. 1998. Mutation spectrum and genotype-phenotype analyses in Cowden disease
and Bannayan-Zonana syndrome, two hamartoma syndromes with germline PTEN mutation.
Hum Mol Genet, 7**,** 507-15.
MARSH, D. J., KUM, J. B., LUNETTA, K. L., BENNETT, M. J., GORLIN, R. J., AHMED, S. F., BODURTHA, J.,
CROWE, C., CURTIS, M. A., DASOUKI, M., DUNN, T., FEIT, H., GERAGHTY, M. T., GRAHAM, J. M.,
JR., HODGSON, S. V., HUNTER, A., KORF, B. R., MANCHESTER, D., MIESFELDT, S., MURDAY, V.
A., NATHANSON, K. L., PARISI, M., POBER, B., ROMANO, C., ENG, C. & ET AL. 1999. PTEN
mutation spectrum and genotype-phenotype correlations in Bannayan-Riley-Ruvalcaba
syndrome suggest a single entity with Cowden syndrome. *Hum Mol Genet,* 8**,** 1461-72.
MASSON, G. R., BURKE, J. E. & WILLIAMS, R. L. 2016a. Methods in the Study of PTEN Structure: X-Ray
Crystallography and Hydrogen Deuterium Exchange Mass Spectrometry. *Methods Mol Biol,*
1388**,** 215-30.
MASSON, G. R., PERISIC, O., BURKE, J. E. & WILLIAMS, R. L. 2016b. The intrinsically disordered tails of
PTEN and PTEN-L have distinct roles in regulating substrate specificity and membrane activity.
Biochem J, 473**,** 135-44.
MATOS-RODRIGUES, G., MARTINI, E. & LOPEZ, B. S. 2021. Mouse Models for Deciphering the Impact
of Homologous Recombination on Tumorigenesis. *Cancers (Basel),* 13.
MENDES-PEREIRA, A. M., MARTIN, S. A., BROUGH, R., MCCARTHY, A., TAYLOR, J. R., KIM, J. S.,
WALDMAN, T., LORD, C. J. & ASHWORTH, A. 2009. Synthetic lethal targeting of PTEN mutant
cells with PARP inhibitors. *EMBO Mol Med,* 1**,** 315-22.
MESTER, J. L., TILOT, A. K., RYBICKI, L. A., FRAZIER, T. W., 2ND & ENG, C. 2011. Analysis of prevalence
and degree of macrocephaly in patients with germline PTEN mutations and of brain weight in
Pten knock-in murine model. *Eur J Hum Genet,* 19**,** 763-8.
MIGHELL, T. L., EVANS-DUTSON, S. & O'ROAK, B. J. 2018. A Saturation Mutagenesis Approach to
Understanding PTEN Lipid Phosphatase Activity and Genotype-Phenotype Relationships. *Am J*
Hum Genet, 102**,** 943-955.
MISTICONE, S., LIMA-FERNANDES, E. & SCOTT, M. G. 2016. Rapid Detection of Dynamic PTEN
Regulation in Living Cells Using Intramolecular BRET. *Methods Mol Biol,* 1388**,** 95-110.
MYERS, M. P., PASS, I., BATTY, I. H., VAN DER KAAY, J., STOLAROV, J. P., HEMMINGS, B. A., WIGLER, M.
H., DOWNES, C. P. & TONKS, N. K. 1998. The lipid phosphatase activity of PTEN is critical for its
tumor supressor function. *Proc Natl Acad Sci U S A,* 95**,** 13513-8.
NELEN, M. R., PADBERG, G. W., PEETERS, E. A., LIN, A. Y., VAN DEN HELM, B., FRANTS, R. R., COULON,
V., GOLDSTEIN, A. M., VAN REEN, M. M., EASTON, D. F., EELES, R. A., HODGSEN, S., MULVIHILL,
J. J., MURDAY, V. A., TUCKER, M. A., MARIMAN, E. C., STARINK, T. M., PONDER, B. A., ROPERS,
H. H., KREMER, H., LONGY, M. & ENG, C. 1996. Localization of the gene for Cowden disease to
chromosome 10q22-23. *Nat Genet,* 13**,** 114-6.
NGEOW, J. & ENG, C. 2019. PTEN in Hereditary and Sporadic Cancer. *Cold Spring Harb Perspect Med*.
ODRIOZOLA, L., SINGH, G., HOANG, T. & CHAN, A. M. 2007. Regulation of PTEN activity by its carboxyl-
terminal autoinhibitory domain. *J Biol Chem,* 282**,** 23306-15.
OOMS, L. M., BINGE, L. C., DAVIES, E. M., RAHMAN, P., CONWAY, J. R., GURUNG, R., FERGUSON, D. T.,
PAPA, A., FEDELE, C. G., VIEUSSEUX, J. L., CHAI, R. C., KOENTGEN, F., PRICE, J. T., TIGANIS, T.,
TIMPSON, P., MCLEAN, C. A. & MITCHELL, C. A. 2015. The Inositol Polyphosphate 5-
Phosphatase PIPP Regulates AKT1-Dependent Breast Cancer Growth and Metastasis. *Cancer*
Cell, 28**,** 155-69.
PAL, A., BARBER, T. M., VAN DE BUNT, M., RUDGE, S. A., ZHANG, Q., LACHLAN, K. L., COOPER, N. S.,
LINDEN, H., LEVY, J. C., WAKELAM, M. J., WALKER, L., KARPE, F. & GLOYN, A. L. 2012. PTEN

mutations as a cause of constitutive insulin sensitivity and obesity. *N Engl J Med,* 367, **1002-**
11.
PODSYPANINA, K., ELLENSON, L. H., NEMES, A., GU, J., TAMURA, M., YAMADA, K. M., CORDON-CARDO,
C., CATORETTI, G., FISHER, P. E. & PARSONS, R. 1999. Mutation of Pten/Mmac1 in mice causes
neoplasia in multiple organ systems. *Proc Natl Acad Sci U S A,* 96, **1563-8.**
POST, K. L., BELMADANI, M., GANGULY, P., MEILI, F., DINGWALL, R., MCDIARMID, T. A., MEYERS, W.
M., HERRINGTON, C., YOUNG, B. P., CALLAGHAN, D. B., ROGIC, S., EDWARDS, M.,
NICIFOROVIC, A., CAU, A., RANKIN, C. H., O'CONNOR, T. P., BAMJI, S. X., LOEWEN, C. J. R.,
ALLAN, D. W., PAVLIDIS, P. & HAAS, K. 2020. Multi-model functionalization of disease-
associated PTEN missense mutations identifies multiple molecular mechanisms underlying
protein dysfunction. *Nat Commun,* 11, **2073.**
RANTAKARI, P., NIKKILA, J., JOKELA, H., OLA, R., PYLKAS, K., LAGERBOHM, H., SAINIO, K., POUTANEN,
M. & WINQVIST, R. 2010. Inactivation of Palb2 gene leads to mesoderm differentiation defect
and early embryonic lethality in mice. *Hum Mol Genet,* 19, **3021-9.**
ROY, R., CHUN, J. & POWELL, S. N. 2011. BRCA1 and BRCA2: different roles in a common pathway of
genome protection. *Nat Rev Cancer,* 12, **68-78.**
SCHEI-ANDERSEN, A. J., WITJES, V. M., VOS, J. R., MENSENKAMP, A. R., VAN ALTENA, A., SCHIEVING,
J., SIMONS, M., SCHUURS-HOEIJMAKERS, J. H. M., GROUP, P. S. & HOOGERBRUGGE, N. 2025.
Non-serous ovarian cancer in PTEN Hamartoma Tumor Syndrome: additional evidence for
increased risk. *Fam Cancer,* 24, **28.**
SCHONHUBER, N., SEIDLER, B., SCHUCK, K., VELTKAMP, C., SCHACHTLER, C., ZUKOWSKA, M., ESER, S.,
FEYERABEND, T. B., PAUL, M. C., ESER, P., KLEIN, S., LOWY, A. M., BANERJEE, R., YANG, F., LEE,
C. L., MODING, E. J., KIRSCH, D. G., SCHEIDELER, A., ALESSI, D. R., VARELA, I., BRADLEY, A., KIND,
A., SCHNIEKE, A. E., RODEWALD, H. R., RAD, R., SCHMID, R. M., SCHNEIDER, G. & SAUR, D.
2014. A next-generation dual-recombinase system for time- and host-specific targeting of
pancreatic cancer. *Nat Med,* 20, **1340-1347.**
SCUDAMORE, C. 2014. Practical approaches to reviewing and recording pathology data. *In:* C.L., S. (ed.)
A Practical Guide to Histology of the Mouse Chichester, UK: Wiley-Blackwell.
SHEN, W. H., BALAJEE, A. S., WANG, J., WU, H., ENG, C., PANDOLFI, P. P. & YIN, Y. 2007. Essential role
for nuclear PTEN in maintaining chromosomal integrity. *Cell,* 128, **157-70.**
SMITH, I. N. & BRIGGS, J. M. 2016. Structural mutation analysis of PTEN and its genotype-phenotype
correlations in endometriosis and cancer. *Proteins,* 84, **1625-1643.**
SPINELLI, L., BLACK, F. M., BERG, J. N., EICKHOLT, B. J. & LESLIE, N. R. 2015. Functionally distinct groups
of inherited PTEN mutations in autism and tumour syndromes. *J Med Genet,* 52, **128-34.**
STECK, P. A., PERSHOUSE, M. A., JASSER, S. A., YUNG, W. K., LIN, H., LIGON, A. H., LANGFORD, L. A.,
BAUMGARD, M. L., HATTIER, T., DAVIS, T., FRYE, C., HU, R., SWEDLUND, B., TENG, D. H. &
TAVTIGIAN, S. V. 1997. Identification of a candidate tumour suppressor gene, MMAC1, at
chromosome 10q23.3 that is mutated in multiple advanced cancers. *Nat Genet,* 15, **356-62.**
SUZUKI, A., DE LA POMPA, J. L., STAMBOLIC, V., ELIA, A. J., SASAKI, T., DEL BARCO BARRANTES, I., HO,
A., WAKEHAM, A., ITIE, A., KHOO, W., FUKUMOTO, M. & MAK, T. W. 1998. High cancer
susceptibility and embryonic lethality associated with mutation of the PTEN tumor suppressor
gene in mice. *Curr Biol,* 8, **1169-78.**
TIBAREWAL, P., RATHBONE, V., CONSTANTINOU, G., PEARCE, W., ADIL, M., VARYOVA, Z., FOLKES, L.,
HAMPSON, A., CLASSEN, G. A. E., ALVES, A., CARVALHO, S., SCUDAMORE, C. L. &
VANHAESEBROECK, B. 2022. Long-term treatment of cancer-prone germline PTEN mutant
mice with low-dose rapamycin extends lifespan and delays tumour development. *J Pathol,*
258, **382-394.**
TIBAREWAL, P., ZILIDIS, G., SPINELLI, L., SCHURCH, N., MACCARIO, H., GRAY, A., PERERA, N. M.,
DAVIDSON, L., BARTON, G. J. & LESLIE, N. R. 2012. PTEN protein phosphatase activity
correlates with control of gene expression and invasion, a tumor-suppressing phenotype, but
not with AKT activity. *Sci Signal,* 5, **ra18.**

TILOT, A. K., GAUGLER, M. K., YU, Q., ROMIGH, T., YU, W., MILLER, R. H., FRAZIER, T. W., 2ND & ENG,
C. 2014. Germline disruption of Pten localization causes enhanced sex-dependent social
motivation and increased glial production. *Hum Mol Genet,* 23**,** 3212-27.
TORRES, J. & PULIDO, R. 2001. The tumor suppressor PTEN is phosphorylated by the protein kinase
CK2 at its C terminus. Implications for PTEN stability to proteasome-mediated degradation. *J
Biol Chem,* 276**,** 993-8.
TSUZUKI, T., FUJII, Y., SAKUMI, K., TOMINAGA, Y., NAKAO, K., SEKIGUCHI, M., MATSUSHIRO, A.,
YOSHIMURA, Y. & MORITAT 1996. Targeted disruption of the Rad51 gene leads to lethality in
embryonic mice. *Proc Natl Acad Sci U S A,* 93**,** 6236-40.
VAZQUEZ, F., GROSSMAN, S. R., TAKAHASHI, Y., ROKAS, M. V., NAKAMURA, N. & SELLERS, W. R. 2001.
Phosphorylation of the PTEN tail acts as an inhibitory switch by preventing its recruitment into
a protein complex. *J Biol Chem,* 276**,** 48627-30.
WANG, M., WEI, P. C., LIM, C. K., GALLINA, I. S., MARSHALL, S., MARCHETTO, M. C., ALT, F. W. & GAGE,
F. H. 2020. Increased Neural Progenitor Proliferation in a hiPSC Model of Autism Induces
Replication Stress-Associated Genome Instability. *Cell Stem Cell,* 26**,** 221-233 e6.
WEI, R., HITOMI, M., SADLER, T., YEHIA, L., CALVETTI, D., SCOTT, J. & ENG, C. 2024. Quantitative
evaluation of DNA damage repair dynamics to elucidate predictors of autism vs. cancer in
individuals with germline PTEN variants. *PLoS Comput Biol,* 20**,** e1012449.
WHITE, S. L., JAMIL, T., BELL, C., FISHBEIN, L., HAUGEN, B. R., GIGNOUX, C. R. & POZDEYEV, N. 2025.
Population Prevalence of the Major Thyroid Cancer–Associated Syndromes. *The Journal of
Clinical Endocrinology & Metabolism*.
YAN, D. H., WEN, Y., SU, L. K., XIA, W., WANG, S. C., ZHANG, S., GAN, L., LEE, D. F., SPOHN, B., FREY, J.
A., HORTOBAGYI, G. N. & HUNG, M. C. 2004. A delayed chemically induced tumorigenesis in
Brca2 mutant mice. *Oncogene,* 23**,** 1896-901.
YEHIA, L. & ENG, C. 2018. 65 YEARS OF THE DOUBLE HELIX: One gene, many endocrine and metabolic
syndromes: PTEN-opathies and precision medicine. *Endocr Relat Cancer,* 25**,** T121-T140.
YEHIA, L., KEEL, E. & ENG, C. 2020. The Clinical Spectrum of PTEN Mutations. *Annu Rev Med,* 71**,** 103-
116.
YEHIA, L., NGEOW, J. & ENG, C. 2019. PTEN-opathies: from biological insights to evidence-based
precision medicine. *J Clin Invest,* 129**,** 452-464.
YIP, H. Y. K., CHEE, A., ANG, C. S., SHIN, S. Y., OOMS, L. M., MOHAMMADI, Z., PHILLIPS, W. A., DALY, R.
J., COLE, T. J., BRONSON, R. T., NGUYEN, L. K., TIGANIS, T., HOBBS, R. M., MCLEAN, C. A.,
MITCHELL, C. A. & PAPA, A. 2020. Control of Glucocorticoid Receptor Levels by PTEN
Establishes a Failsafe Mechanism for Tumor Suppression. *Mol Cell,* 80**,** 279-295 e8.
YUAN, E., TSAI, P. T., GREENE-COLOZZI, E., SAHIN, M., KWIATKOWSKI, D. J. & MALINOWSKA, I. A. 2012.
Graded loss of tuberin in an allelic series of brain models of TSC correlates with survival, and
biochemical, histological and behavioral features. *Hum Mol Genet,* 21**,** 4286-300.
ZHOU, J., BLUNDELL, J., OGAWA, S., KWON, C. H., ZHANG, W., SINTON, C., POWELL, C. M. & PARADA,
L. F. 2009. Pharmacological inhibition of mTORC1 suppresses anatomical, cellular, and
behavioral abnormalities in neural-specific Pten knock-out mice. *J Neurosci,* 29**,** 1773-83.

1172
1173

1174

Table 1

		PHTS patients with *PTEN* R173 variants	PHTS patients with other nuclear excluded *PTEN* variants	Both groups combined
Patient demographics	Number of patients (N)	34	25	59
	Age range (median age)	7mo-63y (12.5y)	1y-40y (6y)	7mo-63y (10y)
	Gender distribution N (%)	m: 19 (56%) f: 14 (41%) NR: 1 (3%)	m: 12 (48%) m: 9 (36%) NR: 4 (16%)	m: 31(52%) f: 23 (39%) NR: 5 (8%)
Phenotypes (N (%))				
Neuro-developmental	Macrocephaly	32 (94%)	21 (84%)	53 (90%)
	Developmental disorders	18 (53%)	16 (64%)	34 (58%)
Cutaneous and vascular features	Cutaneous pathology	16 (47%)	12 (48%)	28 (47%)
	Vascular abnormalities	5 (15%)	3 (12%)	8 (14%)
	Lipomas	-	4 (16%)	4 (7%)
	Malignant cancer of skin	1 (3%) Age: 63y	-	1 (1.7 %)
Immune/Lymphoid features	Various	3 (9%)	1 (4%)	4 (7%)
Thyroid	Benign lesions	6 (18%)	4 (16%)	10 (17%)
	Malignant cancer	1 (3%) Age: 59y	1 (4%) 29y	2 (3%)
Gastro-intestinal	Benign polyps	4 (12%)	5 (20%)	9 (15%)
Breast	Benign lesions	2 (6%)	1 (4%)	3 (5%)
	Malignant cancer	2 (6%) Ages: 54y & 59y	-	2 (3%)
Ovary	Benign lesions	3 (9%)	-	3 (5%)
	Malignant cancer	1 (3%) Age: 63y	-	1 (1.7 %)
Uterus	Benign lesions	3 (9%)	-	3 (5%)
Other cancers	Testicular cancer	-	1 (4%) Age: 29y	1 (1.7%)
	Lung cancer	-	1 (4%) Age: NR	1 (1.7%)

Table 1. Clinical phenotypes of PHTS patients with *Pten-R173* variants and other nuclear excluded *Pten* variants

mo, age in months; y, age in years; m, males; f, females, NR, not recorded. Phenotypic categories include the following: Developmental disorders; global developmental delay, delay in speech development, mental retardation, autism spectrum disorder, special educational needs. Cutaneous pathology; hamartomas, tricholemmoma, mucocutaneous lesions, acral keratoses, palmoplantar keratoses, oral papillomas, penile freckling, facial papules, café-au-lait spots, skin tags. Vascular abnormalities; haemangiomas, venous malformations, arterio-venous malformations. Immune/

lymphoid features; recurrent upper respiratory tract infections, hypogammaglobulinemia, enlarged
tonsils and adenoids, lymphadenopathy, splenomegaly. Thyroid benign lesions; cyst, nodules,
adenomas. Thyroid malignant cancer; follicular cancer, papillary-follicular thyroid cancer. Breast
benign lesions; fibrocystic disease of the breast, fibroadenomas, benign breast disease. Breast
malignant cancer; ductal breast carcinoma, invasive ductal carcinoma and ductal carcinoma in-situ of
the breast. Ovary benign lesions; ovarian cysts, benign neoplasm. Ovary malignant cancer; malignant
neoplasm of the ovary. Uterus benign lesions; uterine fibroids, endometrial polyp. The phenotypes of
individual patients are listed in Supplementary Tables 1 and 2.
1193

1194   **Table 2**

1195

Phenotypes	Mouse genotype				
	Pten^{+R173C}	*Pten*$^{+/-}$	*Pten*$^{-/-}$	*Pten*$^{+/m3m4}$	*Pten*$^{m3m4/m3m4}$
Increased brain size	Yes	Yes	ND	Yes	Yes
Bigger corpus callosum	Yes	Yes	ND	ND	ND
Increased numbers of oligodendroglia cells	Yes	No	ND	No	Yes
Astrogliosis	No	No	ND	No	Yes
Increased cell body size	Yes	No	Yes	No	Yes
Increased neuronal complexity	Yes	No	Yes	No	No
Enlarged axons	Yes	No	ND	ND	ND
thinner myelin	Yes	Yes	ND	ND	ND
Sociability	No significant difference	ND	ND	No significant difference	Too social with revised criteria
Motor coordination	ND	ND	ND	ND	reduced
Locomotion	reduced	ND	ND	ND	ND

1196

1197   **Table 2. Neuronal and behavioural phenotypes in mice with *Pten* mutations**: Table shows data from
1198   *Pten*$^{+/R173C}$, *Pten*$^{+/-}$, and conditional *Pten*$^{-/-}$ mice from this study, and *Pten*$^{+/m3m4}$ *and* Pten$^{m3m4/m3m4}$ mice
1199   as reported in (Mester et al., 2011, Tilot et al., 2014). ND; not determined. The behavioural assays
1200   (sociability, motor coordination and locomotion) are relative to wild type littermate mice.

1201
1202

Figure 1

1203
1204
Figure 1. Characterisation of PTEN-R173C/H. (A-C) Transient lentiviral expression in U87 cells of PTEN-
WT or the indicated PTEN mutants at fixed (A and C) or increasing concentrations of lentiviral particles
(B). 48 h after transduction (A and B), cells were either lysed and PTEN expression and (B) AKT P-S473
and S6 P-S240/244 assessed by immunoblotting, or (A) qPCR analysis was performed to determine
PTEN mRNA levels (mean ± SEM, n=3, right panel). The PTEN-C124S mutant was used as a control. **(C)**
Cell proliferation was determined by MTS assay over 4 days. Graph shows the mean ± SEM of
absorbance at 490 nm relative to day 0 from 10 replicate wells of a representative experiment from
n=2. Immunoblots of PTEN protein and AKT P-S473 from the cells in (C) are shown in **Fig. S2A**. **(D)** U87
cells transduced with lentivirus expressing a PIP$_3$ biosensor, with or without PTEN-WT or PTEN-R173C,
were pre-treated ± GDC-0941 (1 µM) for 1 h, followed ± insulin treatment (100 nM) for 2 min. The cells
were stained with antibodies to PTEN and imaged by confocal microscopy (images in **Fig. S2B**). Graph
shows mean fluorescence intensity (MFI) of the PIP$_3$ biosensor at the plasma membrane normalized
to the MFI of the PIP$_3$ biosensor in the cytosol. Data shown as mean ± SEM from 3 independent
experiments, statistical analysis performed using 2-way ANOVA. Representative immunoblots for
PTEN protein and AKT P-S473 from the cells are shown in **Fig. S2C**. **(E-F)** Transient lentiviral expression

1220 of PTEN wild-type (WT) or the indicated PTEN mutants in U87 cells. (E) The cells were treated with
1221 cycloheximide and PTEN protein expression levels determined by immunoblotting at the indicated
1222 time points. Graph shows PTEN protein levels normalized to time 0, data shown as mean ± SEM, n=6.
1223 Representative blots shown in Fig. S3A. (F) Phosphorylation of PTEN on S380/T382/T383 was
1224 determined by immunoblotting. Quantification shows mean ± SEM, n=3, statistical analysis was
1225 performed using Unpaired Student's t-test. (**G**) mBRET and ΔmBRET values, obtained in live HEK-293T
1226 and HeLa PTEN KO cells, transfected with Rluc-PTEN-WT-YFP, Rluc-PTEN-R173C-YFP or Rluc-PTEN-
1227 R173H-YFP. Graphs represent mean ± SD of n=8 points from a representative experiment from 3
1228 independent experiments. (**H**) Transient lentiviral expression of PTEN-WT or the indicated PTEN
1229 mutants in U87 cells. Protein extracts from these cells were used for immunoblotting to assess PTEN
1230 protein levels. (**I**) Lenti-X™ 293T cells expressing GFP-tagged PTEN-WT or PTEN-R173C were arrested
1231 in the G1/S phase of cell cycle by double-thymidine block and imaged by confocal microscopy at the
1232 indicated time points after release from thymidine block, for expression of PTEN-GFP. Left panel shows
1233 representative images, arrows indicate nuclear PTEN-WT or reduced nuclear PTEN-R173C. Scale bar:
1234 10 μm. The graph shows the nuclear:cytoplasmic ratio of the mean fluorescence intensity (MFI) of the
1235 PTEN-GFP signal (mean ± SEM; n=3, statistical analysis performed using 2-way ANOVA). p-values for
1236 all statistical tests: *, $p < 0.05$; **, $p<0.01$; ***, $p<0.001$; ****, $p<0.0001$.

Figure 2. Characterisation of *Pten*[+/R173C] mice. (**A**) Homozygous *Pten*[R173C/R173C] mice die *in utero* after embryonic stage E9.5. Images show morphology of E9.5 *Pten*[+/+], *Pten*[+/R173C] and *Pten*[R173C/R173C] embryos. Scale bar: 1 mm. The table shows the number of embryos/pups of each genotype obtained at the indicated time points. ¥ indicates embryos smaller in size and * indicates embryos that had been reabsorbed. (**B, C**) Protein extracts from MEFs of the indicated genotype were used for immunoblotting with the antibodies shown. Data shown as mean ± SEM. At least 3 independent MEF lines of each were used for (B) and 2 independent MEF lines were used for (C). (**D**) MEFs of the

indicated genotype were treated with insulin (100 nM) for the indicated times and protein extracts
immunoblotted with the antibodies shown. Data shown as mean ± SEM from 2 independent MEF lines
for each genotype. (**E**) ITT assays on 3-month-old male mice of the indicated genotypes. XY curves
show blood glucose levels post insulin injection over time. Bar graphs show mean ± SEM of
measurements of Area Under the Curve (AUC). Statistical analysis performed using Unpaired Student's
t-test. (**F**) MEFs of the indicated genotype were immunostained with antibodies to PTEN (green) and
DAPI (blue) and imaged by confocal microscopy. Scale bar: 10 µm. Graphs show nuclear:cytoplasmic
ratio of the GFP MFI (mean ± SEM from 3 independent experiments). Statistical analysis was performed
using Unpaired Student's t-test. (**G**) 6-week-old mice of the indicated genotypes were treated with
7Gy γ-radiation and protein extracts from uterus were used for immunoblotting. Graph shows levels
of γH2AX relative to littermate *Pten*$^{+/+}$ controls. Data shown as mean ± SEM. Statistical analysis was
performed using Mann–Whitney U tests. Accompanying immunoblots are shown in **Fig. S9A**. p-values
for all statistical analysis: *, p< 0.05; **, p<0.01; ***, p<0.001; ****, p<0.0001. Mice used for
generating MEF lines in (A-D and F) and for experiments in (E and G) were on a C57BL/6J background.

Figure 3

Figure 3. Survival and histopathological analysis of *Pten*+/R173C mice. *Pten*+/R173C and *Pten*+/- mice on a mixed C57BL/6J x Sv129 background were allowed to age. Mice were euthanized for welfare reasons (ill health or palpable masses with a combined size of ≥ 1.4 cm^2 surface area) or at a specified age. (**A**) Kaplan–Meier survival curves for male and female mice. Statistical analysis was performed using Log-rank (Mantel–Cox) test and Gehan–Breslow–Wilcoxon test. Comparisons were made to *Pten*+/+ mice (p-values shown as black stars) or *Pten*+/- mice (p-values shown as red stars). Table show median survival age of the mice. (**B**) Incidence of specific tumour types in male and female mice, as assessed by histopathological analysis on all mice. Statistical analysis was performed using Fisher's exact test. (**C**) Representative IHC images of uteri from female *Pten*+/R173C and *Pten*+/- and littermate *Pten*+/+ controls at the indicated timepoints, black arrowheads indicate loss of PTEN immunoreactivity and the corresponding AKT P-S473 immunoreactivity. (**D**) Oncoplot showing co-occurrence of mutations in the

indicated genes in tumor samples harbouring PTEN-R173C mutation (cBioportal). (**E**) Grade of
lymphoid hyperplasia in mice at the end of study. Graph shows scatter plot of scores from individual
mice with line at median score. Statistical analysis was performed using Fisher's exact test. (**F-G**) Flow
cytometry analysis was performed on the lymph nodes of female *Pten*$^{+/R173C}$, *Pten*$^{+/-}$ and littermate
Pten$^{+/+}$ controls at 5 and 10 months of age. Graphs show the proportion of (F) B220+ B-cells and (G)
TCRβ+ T-cells. Statistical analysis was performed using Mann–Whitney U test. p-values for all statistical
analysis: *, p< 0.05; **, p<0.01; ***, p<0.001; ****, p<0.0001.

Figure 4

1280
1281

Figure 4. Enlarged brain phenotype with associated increases in grey and white matter areas in *Pten* mutant mice. (**A**) Representative NeuN immunostaining of coronal sections from *Pten*[+/+] and mutant animals at 3 months. The overlay of the contours of the depicted slices is also shown. Scale bar: 1000 μm. (**B**) Brain mass, body mass and brain mass as a % of body mass for male and female *Pten*[+/-], *Pten*[+/R173C] and WT littermate controls at 3 months. Data show mean ± SEM, statistical analysis performed using Two-tailed Mann-Whitney U test. (**C**) Representative NeuN immunostaining from coronal sections of WT and mutant animals at 3 months. Scale bar: 1000 μm. (**D**) Quantification of cortical thickness in the primary somatosensory cortex barrel field (area indicated with long arrows in C). n = 4 mice of each sex and genotype. Data show mean ± SEM, statistical analysis performed using Two-tailed Unpaired Student's t-test. (**E**) Quantification of the thickness of the corpus callosum at the indicated Bregma levels (area indicated with short arrows in C). n=4 mice of each sex and genotype.

Data show mean ± SEM, statistical analysis performed using Two-way RM ANOVA with uncorrected
Fisher's LSD test. (**F**) Representative sagittal sections from control and mutant mouse brains stained
with Toluidine Blue and quantification of the corpus callosum length, thickness, and area. Scale bar:
500 µm. n=3 mice per genotype. Data show mean ± SEM, statistical analysis performed using One-way
RM ANOVA with uncorrected Fisher's LSD test. (**G**) Top: Representative images of coronal sections
from the somatosensory cortex of *Pten*$^{+/+}$ and mutant animals at 3 months stained for NeuN. Scale bar:
250 µm. Bottom: Quantification of the number and density of neurons in the indicated boxed areas in
(G). n=4 mice of each sex each genotype. Data show mean ± SEM; statistical analysis performed using
Two-tailed Unpaired Student's t-test. (**H**) Top: Representative images of sections from *Pten*$^{+/+}$ and *Pten*
mutant animals at 3 months immunostained with OLIG2 and GFAP, Scale bar: 100 µm, and bottom:
quantification of OLIG2 positive cells in the motor cortex. n=4 mice per genotype. Data show mean ±
SEM and statistical analysis performed using Two-tailed Unpaired Student's t-test. p-values for all
statistical tests: *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$; ****, $p<0.0001$. Mice used were on a mixed
C57BL/6J x Sv129 background.

Figure 5

1307
Figure 5. Neuronal and glial deficits in *Pten*^+/R173C^ and *Pten*^+/-^ mice. (A) Representative images from
GFP-transfected cortico-hippocampal neurons prepared from the indicated *Pten* mutants and their
respective *Pten*^+/+^ littermate controls at DIV12/13. Neurons were stained for GFP. Scale bar: 20 μm. **(B)**
Quantification of cell body size of GFP-transfected cultured cortico-hippocampal neurons shown in (A).
from n>60 cells for each genotype from 3-6 independent experiments. Violin plots show mean ± SEM,
statistical analysis performed using Two-tailed Mann-Whitney U test. **(C)** Sholl analysis of GFP-
transfected cortico-hippocampal neurons shown in (A) from n=63–122 cells from 3–5 independent
experiments. Data show mean ± SEM, statistical analysis performed using Two-way RM ANOVA with
uncorrected Fisher's LSD test. **(D)** Representative electron micrographs of the corpus callosum in cross-
sections from 3-month-old *Pten*^+/+^, *Pten*^+/-^ and *Pten*^+/R173C^ mice. Scale bar: 1 μm. **(E)** Quantification of
axonal dimeters and their frequency difference in mutant mice compared to *Pten*^+/+^ littermate

controls. >4500 axons measured in each of three mice per genotype. Violin plots indicate mean ± SEM,
statistical analysis performed using Two-tailed Mann-Whitney U test. (**F**) Quantification of g-ratios for
corpus callosum axons. >4500 axons measured in each of three mice per genotype. Statistical analysis
performed using Two-tailed Mann-Whitney U test. (**G**) Scatter plot of g-ratios as a function of axon
diameter in 3-month-old *Pten*$^{+/+}$, *Pten*$^{+/-}$ and *Pten*$^{+/R173C}$ mice. >4500 axons measured in each of three
mice per genotype. p-values for all statistical tests: *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****,
$p < 0.0001$. Mice used in (A-C) were on a C57BL/6J background, and in (D-G) were on a mixed C57BL/6J
x Sv129 background.
1327