Phage-based microbiome manipulation elucidates interactions within community context

Tomas Hessler[1,2,3]*, Dawn Chiniquy[3], Benjamin A. Adler[1], Jordan Hoff[1], Eliana Tucker[3], Robert Huddy[4], Rohan Sachdeva[1], Shufei Lei[1], Susan T. L. Harrison[5,6,7], Jennifer Doudna[1], Rodolphe Barrangou[8], Spencer Diamond[1], Adam M. Deutschbauer[3,9], Jillian F. Banfield[1,2,3,10] *

[1]The Innovative Genomics Institute at the University of California, Berkeley, California, USA
[2]The Department of Earth and Planetary Science, University of California, Berkeley, California, USA
[3]Environmental Genomics and Systems Biology Division, Lawrence Berkeley National Laboratory, Berkeley, CA, USA
[4]Research Office, Faculty of Health Sciences, University of Cape Town, Cape Town 7700, South Africa
[5]The Center for Bioprocess Engineering Research, University of Cape Town, South Africa
[6]The Future Water Institute, University of Cape Town, South Africa
[7]Faculty of Engineering, Architecture & IT, University of Queensland, St Lucia, Australia
[8]North Carolina State University, Raleigh, USA
[9]Department of Plant and Microbial Biology, University of California, Berkeley, California, USA
[10] Department of Microbiology, Biomedicine Discovery Institute, Monash University, Melbourne, VIC, Australia

*Corresponding authors: tomas.hessler@berkeley.edu, jbanfield@berkeley.edu

Abstract

Testing microbial interaction hypotheses is a major challenge in microbiome research. Bacteriophages may be used to probe organism-organism interactions within complex microbiomes, yet are rarely used for this purpose. Here, we isolated nine narrow host range phages that replicate in several *Variovorax* species. As *Variovorax* CL14 degrades the plant root stunting hormone auxin, we used phages to eliminate *V.* CL14 from a rhizosphere consortium and reestablished the stunted root phenotype. We used three of the phages to deplete another *Variovorax*, SCN45, from complex communities to test correlation network-based hypotheses of thiamine interdependencies*.* Genome-resolved metagenomics revealed that three taxa decreased in relative abundance following *Variovorax* depletion and could be rescued by thiamine supplementation. Thus, we confirmed thiamine production as the mechanistic basis for interdependence. These experiments lay the foundation for research that employs wildtype and genetically modified phages to test interaction hypotheses and for targeted microbiome manipulation.

Introduction

Microorganisms form the foundation of food webs, and profoundly influence the health of plants,
humans, and other animals on a global scale. Yet microorganisms rarely function in isolation, and
instead form complex communities in which a multitude of interactions shape its structure
(Pacheco et al., 2019, Ma et al., 2020, Kost et al., 2023, Clegg & Gross, 2025).

Complex microbial communities have long been studied using metagenomics (Tyson et al., 2004),
metatranscriptomics (Gilbert et al., 2008, Frias-Lopez et al., 2008), metaproteomics (Ram et al.,
2005), and metabolomics (Segata et al., 2013), sometimes informed by stable isotope probing
(Starr et al., 2018). Although these methods may generate hypotheses about organism
interdependencies, our ability to test predicted interdependencies in a community context is
extremely limited and largely restricted to highly specialized techniques (Pierce and Dutton 2022).

Microbial community models can infer the sum of all interactions (Skwara et al., 2023, Quinn-
Bohmann et al., 2024), but lack of species-resolved mechanistic understanding limits our ability
to engineer these systems. Metabolic models can predict interactions by incorporating microbial
metabolic capacities (Dukovski et al., 2021), but this approach is currently limited to co-cultures
and simple synthetic communities. This motivates the search for approaches that can elucidate
organism-organism interactions in complex microbial ecosystems.

Bacteriophages offer a highly targeted way to probe microbial interactions due to their inherent
bacterial specificity and potent host-killing efficiency. Bacteriophages have been predominantly
used as therapeutics to remove or deplete pathogenic and antibiotic resistant microorganisms
(Buttimer et al., 2022; Thanki et al., 2022; Das et al., 2015, Strathdee et al., 2023; Liao et al.,
2025). Bacteriophage can also be further engineered for improved host killing and broadened
host range (Selle et al., 2020; Gencay et al., 2023).

Prior work has established that phage hold enormous potential for microbiome manipulation.
Carlson et al. (2023) demonstrated that the ammonium-generating capacity of fresh-water
enrichments could be altered through the use of applied bacteriophage. Li et al. (2022)
demonstrated the depletion of a specific *E. coli* using T7 phage can modulate host phenotypes,
specifically animal behaviour. While several studies have engineered phage for increased killing
efficiency (Selle et al., 2020), depletion of antibiotic resistance encoding plasmids (Bikard et al.,
2014) and modification of host range (Gencay et al., 2023).

Here, we isolated multiple lytic bacteriophages specific to a collection of *Variovorax* isolates, and
used them to modulate microbiomes to test several predicted microbial interactions. Our findings
demonstrate that phage can be used to probe microbial interactions in the context of complex
microbial communities.

Results

Variovorax phage isolation

To isolate *Variovorax* phages capable of selectively killing *Variovorax SCN45* we extracted phage viromes from six rhizosphere and compost samples (Oakland, CA) and tested for infectivity against a panel of *Variovorax* strains (Methods). We isolated nine phages that can replicate in one or more *Variovorax* species. These double-stranded DNA phages ranged in genome size from 43 to 159 Kb and with GC contents of 53% to 66%. Based on protein homology, seven of these phages (all with ~50 kbp genomes) were grouped with other RefSeq phages (Fig. 1a) classified predominantly with *Siphoviridae* (now *Siphovirus*) and share a high degree of average nucleotide diversity (ANI) with each other (Fig. 1b). The remaining two phages clustered with *Myoviridae* and *Autographiviridae* families. Two *Siphoviridae* phages (bapic4 and V.45_iii) and the large *Myoviridae* phage (V.45_v66) showed strong lytic activity against *V.* SCN45 (Fig. 1e). V.45_v66, bapic4 and V.45_iii were shown by TEM to have *Siphovirus*-like morphology, consisting of an icosahedral head and long filamentous tails, similar to the well-known lambda phage (Fig. 1d). All nine of the isolated phage are predicted to be restricted to a lytic lifestyle based on the absence of integrase genes. There was very limited cross-reactivity of the phages between the different *Variovorax* strains (Fig. 1c) based on spot assays. Only one phage, V.45_66, was able to form plaques on multiple hosts - *V.SCN45* and *V. paradoxus DSM 645* (Fig. S1).

Figure 1. Characterization of nine isolated *Variovorax*-infecting bacteriophages. (a) Protein similarity network of phages of phages deposited in RefSeq and those isolated in this study (black), with edges

indicating a significant shared protein content. (b) Pairwise average nucleotide identity (ANI) between the nine isolated *Variovorax* phages. (c) Plaque assays demonstrate limited cross-reactivity of the nine phages across four different *Variovorax* strains, with filled circles representing the ability to plaque on a given host. (d) Transmission electron microscopy (TEM) of the SCN45-infecting phage, showing icosahedral heads and filamentous tails characteristic of Siphoviridae. (e) High titre phage stocks visualized by 10-fold dilution 1uL spot assays from 10^{-6} to 10^{0}. (f) Genome-wide fitness profiling (RB-TnSeq) for *V. SCN45* genes, following exposure to phage bap4, identifies the Type IV pilus as the likely receptor used by the phage for infection. Phages V.45_66 and V.45_iii generated similar gene fitness results.

Variovorax SCN45 phages infect via chromosome-encoded type IV pilus

To identify loss-of-function gene disruptions that could confer *Variovorax* with increased fitness under phage infection we constructed an Rb-TnSeq mutant library of *Variovorax* SCN45 and employed methods similar to Mutalik et al. (2020). The disruption of genes of the type IV secretion system (T4SS) were found to substantially enhance the fitness of *Variovorax* when incubated with each of its three phages (Fig. 1f). None of these T4SS gene knockouts were enriched in the no-phage controls, indicating that the addition of phage selected for these gene mutants through the elimination of susceptible mutants.

Evaluating Phage-Mediated Depletion in the *Arabidopsis-Variovorax-Arthrobacter* System

Finkel et al. (2020) demonstrated that *Variovorax CL14* mitigates a root stunting phenotype in *Arabidopsis* by consuming excess auxin secreted by *Arthrobacter* that leads to root stunting. To test whether such well-characterized interactions could be tested through phage-based manipulation, rather than conventional strategies, we hypothesized that a co-culture of the two organisms would result in a normal root phenotype but phage-mediated depletion of *V.* CL14 would result in the accumulation of auxin and cause a stunted root phenotype. We employed a phage cocktail of the Siphoviridae phage V.cl_11 and V.cl_23 to deplete *V.* CL14 from both pure cultures and co-cultures with *Arthrobacter CL28* inoculated onto *Arabidopsis thaliana*. Phage treatment led to the reintroduction of root stunting phenotype (Fig. 2).

127
Figure 2. Phage-mediated depletion of Variovorax from an *Arabidopsis*-inoculated synthetic
community induces root stunting. (a) Model of root inhibition by the auxin-secreting *Arthrobacter* CL28,
which reverts to a normal root phenotype when the auxin-degrading *Variovorax* CL14 is present. If the
Variovorax-targeting phage cocktail (V_11, V_23) is effective in the rhizosphere, this would disrupt the
interaction and the root would revert to stunted growth. (b) *Arabidopsis* primary root elongation with
Arthrobacter CL28 causes a stunted root phenotype, which recovers with and *Variovorax* CL14 co-culture.
The addition of the *Variovorax*-targeting phage cocktail (V_11, V_23) to the co-culture of *Arthrobacter* CL28
and *Variovorax* CL14 recapitulates the root stunting phenotype observed in the presence of *Arthrobacter*
alone. Letters above bars indicate the results of a Tukey post hoc test (α=0.05).

Development of stable enrichment communities

To perform community manipulation experiments it is essential to generate stable, reproducible
consortia with reasonable complexity, ideally containing organisms that are predicted to be
interdependent on other community members. This study leveraged inoculum from lab-scale
bioreactors from which the *Variovorax SCN45* pure culture was previously isolated (Hessler et
al., 2023). *Variovorax SCN45* overproduces thiamine in pure culture and was predicted to supply
this essential vitamin to highly dependent thiamine auxotrophs in bioreactor communities based
on ecological network analyses (Hessler et al., 2023). The thiamine auxotrophs that were
correlated with *Variovorax* had very few correlations with other thiamine-producing organisms,
suggesting a strong dependence.

We supplemented additional *Variovorax* SCN45 cells to increase its abundance in inoculum
consortia based on previous findings that initial ratios of organisms can effect the final steady
state community composition (Gao et al., 2021). The inoculum was then grown in minimal media,
supplemented with various carbon sources, and passaged every 48 hours for a total of seven
passages (Fig. 3a) and community compositions assessed. Four carbon sources were chosen
for ongoing experiments because the communities were reproducible (Fig. 3, S2). The
communities included *Leifsonia*, multiple Xanthomondales and several *Microbacterium*, taxa that
were predicted to depend on *Variovorax* for thiamine (Hessler et al., 2023).

Figure 3. Development of stable enrichment communities. (a) Schematic of Bioreactor-derived
inoculum preparation and enrichment passaging. (b) Communities supplemented with guar gum showed
reproducible compositions across technical replicates and demonstrated stability across multiple passaging
cycles using 16S rRNA gene amplicon sequencing. PCA was performed on robust centered log-ratio (rclr)
transformed ASV relative abundances, demonstrating the compositional (c) reproducibility and (d)
longitudinal stability were consistent across the cultures supplemented with arabinogalactan with guar gum,
guar gum alone, kojac gum, and locust bean gum.

Phage-mediated *Variovorax* depletion leads to broad community impacts

Following the passage of the day 8 communities, either the phage V.45_v66 or a cocktail of three
Variovorax phages (V.45_v66, V.45_iii, and V.45_bapic4) was added to the communities (Fig. 4
a). The relative abundance of *Variovorax* in the guar gum enrichment was reduced from
approximately 22% in the day 8 communities to 2.4% in the day 10 community but further
recovered to 11% by day 12 (Fig. 3b). This pattern was consistent across all four communities
(Fig. S2). There was minimal difference in the abundance of *Variovorax* between the single phage
V.45_v66 and three-phage cocktail, except in the kojac gum- community, where the three-phage
cocktail resulted in a lower *Variovorax* abundance (mean 4%) compared to the single V.45_v66
phage (mean 10%) following passage six (day 12).

The overall community structure was impacted by the addition of phages, as evidenced by a
notable shift in the principal component two of the PCA (Fig. 4c-e) performed on log-normalized
abundance data from amplicon sequencing data. Despite phage addition, community stability was
re-established, both in terms of species composition (Fig. 3c-f) and optical density over
subsequent passages (Fig. S3). Bacteria other than *Variovorax* that can produce thiamine did
not proliferate following *Variovorax* depletion (Fig. 4f). There was no evidence that *Variovorax*
phage interacted with non-target organisms that encode a Type IV secretion system (Fig. 4g).

181
Figure 4. Phage-mediated *Variovorax* depletion across enrichment cultures. (a) Schematic adapted
from Figure 3, illustrating the introduction of phage to 2 x10^4 PFU/ml on day 8 following passaging. (b)
Genome resolved metagenomics revealed a marked reduction of *Variovorax* (blue) on day 10 post-phage.
Principal component analysis (PCA) of log-normalized abundance data from amplicon sequencing data (c)
reveals a separation along PC1 driven by carbon source, while (d) PC2 captures the divergence from pre-
and post-phage conditions, and (e) day of the experiment. The summed relative abundance of all organisms
encoding (f) a complete thiamine biosynthesis pathway and (g) a type IV secretion system (excluding
Variovorax from 100% relative abundance). Shown across three stages: pre-phage addition (day 6 and day

8 communities), post-phage addition (day 10, 12, and 14 communities) and post-phage communities
supplemented with 10 mg/L thiamine.
192

Variovorax depletion confirms dependence of specific thiamine auxotrophs

We assessed which organisms exhibited statistically significant abundance changes between the
'pre-phage communities' (day six and day eight communities) and the communities following the
addition of phage (day 10, 12, 14 communities). After phage administration, we added thiamine
to the control enrichment communities to identify the organisms that were specifically reliant on
Variovorax for thiamine. Amplicon sequencing of the four enrichment cultures revealed that some
organisms that decreased in abundance following *Variovorax* depletion, and then increased
significantly in abundance following thiamine supplementation (Fig. 5a). These included a
Lysobacter (Xanthomonadales*)*, a *Leifsonia,* and microorganisms belonging to the family of
Microbacteriaceae (Fig. 5b). Genome-resolved metagenomics of the samples revealed that the
Leifsonia (Fig. S5), a *Microbacterium* and a *Stenotrophomonas* (Xanthomonadales) all lack a
complete thiamine biosynthesis pathway (Supplementary data).
205
206   To determine if the association between *Leifsonia* species and thiamine-producing bacteria may
207   be more general, we assessed 32 *Leifsonia* genomes present in GTDB for the ability to synthesize
208   thiamine. We found only a single *Leifsonia* genome to have a complete pathway (Supplementary
209   data).

Figure 5. Previously identified organisms experimentally confirmed to rely on *Variovorax* for thiamine. (a) Amplicon sequence variants (ASVs) exhibiting significant changes in relative abundance between the pre- and post phage communities. Many ASVs increased in abundance following *Variovorax* depletion while others showed reduced abundance, and with a subset of these demonstrate recovery following the addition of thiamine. (b) Six ASVs corresponding to multiple *Microbacteria*, a *Leifsonia* and a *Lysobacter* show a reduced abundance following the depletion of *Variovorax* but returned to near original levels with the supplementation of thiamine. (c) Closely related strains of these organisms appear in the correlation network that was initially used to predict *Variovorax* as a thiamine-producer and show correlations with *Variovorax*.

SNPs enabled post-infection recovery of *Variovorax* The *Variovorax* SCN45 genome is comprised of a 6.8 Mb chromosome and 1.2 Mb circular genetic element that is related to a *Variovorax paradoxus S110* sequence described as a "second chromosome" by Han et al. (2010). *Variovorax* SCN45 lacks a CRISPR-Cas systems but encodes 24 PADLOC(v4.3.0)-predicted phage defence systems. We were unable to detect any deletions of these genes, or of any genes (including those

232 of the T4SS), in   post-phage *Variovorax* populations based on read mapping and gene
233 abundances.
234
To seek evidence for mutations that could confer phage immunity in surviving *Variovorax*
populations, we monitored *Variovorax* single-nucleotide polymorphisms (SNPs) across the four
experiments.Before the introduction of phage, the majority of reads map to the reference genome
without SNPs. However, following phage addition, there is a marked increase in the
representation of SNPs and an increase in the number of SNP sites. In prephage communities
the majority of wildtype haplotypes are present at approximately 90% (indicating that at a given
SNP site, 90% of mapped reads support the wildtype base). However, following phage infection
the proportion of SNP sites with the wildtype haplotype reduces, demonstrating selection of new
or previously less abundant SNPs (Fig. 6a, S6).
244
Although most SNPs post-phage infection were synonymous, a subset were not. To identify new
gene variants under strong selection, we identified regions of the genome with SNPs that were
closely spaced enough that linkage could be established. Some regions with high densities of
linked SNPs occurred in both the no-phage controls and post-phage populations (Fig. 6b) and
thus were likely beneficial under all conditions (Fig. 6c). These included genes related to aerobic
benzoyl-CoA catabolism, a type V secretion system, enoyl-CoA hydratase and cell envelope
structural proteins. However, there was a single region of the *Variovorax* genome that showed
high SNP linkage density only in the post-phage populations. This locus encoded multiple
conjugal transfer proteins (Fig. 6c). Most importantly, we identified two non-synonymous SNPs in
the *trbE* conjugal transfer protein (homologous to *virB4*) within the P-loop domain which alters the
charge and hydrophobicity of this region of the protein.
256
To identify SNPs that were not close enough to other mutations to provide linkage, we recovered
all non-synonymous SNPs that increased in abundance by at least 30% in the post-phage
populations relative to the pre-phage and no-phage control populations. We identified 275 gene
variants found exclusively in post-phage populations (Supplementary data). Several were
associated with known phage defence systems including the retron XIII, Lamassu family and
putative PDC-S02 defence systems. Most interesting were three SNPs detected in three or more
of the four communities. This group of genes included a filamentous haemagglutinin family outer
membrane protein, a redoxin/thioredoxin-like sensor protein associated with a fatty acid
biosynthesis operon, and a protein with domain of unknown function (DUF932). This DUF932
gene occurs within an integrated plasmid that encodes *parA*, *parB*, an integrase, a plasmid
transfer protein (*traI*) and a replication initiation protein (*repA*). Notably, multiple non-synonymous
SNPs occurred in the DUF932 protein, and these variants were exceptionally abundant,
exceeding 70% in the post-phage populations. In contrast, the trbE gene displayed non-
synonymous SNPs in 21-33% of reads from post-phage populations. Interestingly, the start codon
of this DUF932 gene was mutated to AAG (lysine) in 50-60% of each of the four populations. This
likely impacts translation initiation and results in truncation of the protein as the next in-frame AUG
codon occurs in position 66.
274

Figure 6. Phage infection induces shifts in *Variovorax* populations and selects for consistent gene variants. (a) Haplotype frequency distributions of *Variovorax* populations in the locust bean gum communities shows the impact by phage application, with expansion of low abundance SNPs, followed by return to pre-phage distributions. (b) Post-phage *Variovorax* populations show increased linked-SNP density in a conjugative transfer gene operon, visualised by a heatmap. Linked-SNPs associated with three 16S rRNA genes have been removed for clarity. (c) A schematic illustrating different categories for selection in day 12 communities. (d) Gene operons and AlphaFold-predicted protein structures showing consistent mutation sites within the conjugal transfer protein TrbE and domain of unknown function within an integrated genetic element.

Variovorax relies on it's own thiamine biosynthesis in all enrichment cultures

To investigate genes important for *Variovorax* fitness in enrichment cultures (in the absence of phage), we introduced *Variovorax* SCN45 mutant libraries into each of the four enrichment cultures (Fig. 7a). In all four of the communities, knockouts of genes involved in the biosynthesis thiamine (*thiC*, *thiG*, *thiE, purE* and *purK)* had a strong negative fitness impact on *Variovorax* (See Fig. S6 for an overview of the pathway). These results indicate that in the four enrichments, *Variovorax* was reliant on its own biosynthesis for thiamine and little extracellular thiamine was likely available.

Rate of recovery of *Variovorax* populations correlates with reduced dependence on Type IV secretion system

Phage treatment reduced the abundance of *Variovorax* to approximately 3% in all day 10 post-phage communities, but *Variovorax* showed the fastest recovery in the guar and locust-bean gum communities over the following days (Fig. 7a). The *Variovorax* populations in these two communities (Fig. 7b, S5) had SNP concentrations close to pre-phage levels by day 14.

We amended enrichments with the Rb-TnSeq mutant library to investigate the consequences of specific gene knockouts for *Variovorax* fitness in the four communities (in the absence of phages; Fig. 7c). In the guar and locust-bean gum communities the knockout of the T4SS was advantageous, based on selection for these genotypes. In contrast, these knockouts were deleterious in the arabino-guar and kojac gum substrate-amended communities.

Figure 7. The rate recovery of *Variovorax* populations from phage infection corresponds to the fitness conferred by the type IV secretion system (T4SS). (a) Relative abundance of *Variovorax* is reduced to ~3% in all post-phage communities, with a gradual recovery over the following passages, again fastest in the guar and locust bean gum communities. (b) Genome-wide SNP sites in Variovorax populations increase immediately following phage infection (day 10) but decline to pre-phage levels, with the guar and locust bean gum populations recovering the fastest. Read mapping was normalized to 20x coverage for all *Variovorax* populations. (c) Genome-wide fitness profiling (Rb-TnSeq) for *V. SCN45* showing the fitness of gene knockouts in each of the four communities (y-axis) versus in pure culture in R2A, rich media (x-axis), highlighting the type IV secretion system and thiamine biosynthesis genes.(b)

Discussion

Previously, networks constructed using organism abundance data allowed us to identify thiamine auxotrophs that were strongly correlated with *Variovorax* (SCN45) in the bioreactor communities studied here (Hessler et al., 2023). Subsequently, we showed that *Variovorax* produces high

concentrations of extracellular thiamine in pure culture, however we did not verify that this thiamine production was sustained within natural communities. To better explore possible interdependence related to vitamin provision required experiments conducted in communities from which *Variovorax* could be selectively removed, without directly impacting the rest of the community. We hypothesized that highly targeted community manipulation could be achieved via use of *Variovorax*-specific phages.

To demonstrate the feasibility of testing organism-organism interactions using phage-mediated *Variovorax* depletion, we leveraged the *Arabidopsis*-*Variovorax* CL14-*Arthrobacter* system first characterized by Finkel et al. (2020) in which *V. CL14* mitigates root stunting in *Arabidopsis* by degrading auxin produced by *Arthrobacter*. Following the addition of isolated phages V.cl_11 and V.cl_23 to this co-culture, we observed a pronounced root-stunting phenotype, indicating that the phages successfully depleted *Variovorax* CL14 (Fig. 2). These findings highlight the potential of extending phage-mediated testing of microbe-microbe and microorganism-plant interactions to rhizosphere microbiome manipulations for agricultural applications.

Next, we used three isolated *Variovorax* SCN45 phages to selectively deplete this *Variovorax* from four enrichment communities and monitored the effects on the remaining community (Fig. 4, 5). The subsequent addition of thiamine facilitated the recovery of only a *Leifsonia*, a Xanthomondales and a *Microbacterium*, each lacking complete thiamine biosynthesis pathway. This provides clear evidence that *Variovorax*-produced thiamine was supporting the growth of these specific organisms in the communities and demonstrates the effectiveness of phage-mediated knock-down experiments for capturing these metabolic interactions. Related, but non-identical, strains of *Leifsonia*, *Microbacterium,* and the Xanthomondales were among a larger group of organisms associated with *Variovorax* in the correlation network from which the thiamine-exchange hypothesis was developed (Hessler et al., 2023).

Using a *Variovorax* SCN45 RB-TnSeq mutant library we found that disruptions in the *Variovorax* type IV secretion system genes conferred phage immunity, indicating that the three phages tested all infect via this complex. The T4SS is a common target for phage attachment and genome injection (Russell 1995; Berry et al., 2011). One potential concern is that depletion of *Variovorax* might occur following phage infection, yet decreased abundance of correlated organisms could be due to toxicity following uptake of viral DNA via T4SSs. Countering this, we find that encoding of a type IV secretion system was not a predictor of whether an organism decreased in abundance following phage addition. Thus, we concluded that these phages could be used to directly identify the organisms that are dependent on *Variovorax* SCN45 in the bioreactor consortia and the basis of the dependence.

Variovorax populations demonstrated partial to complete recovery from phage infection by the end of the experiment. We were unable to identify any gene deletions, frameshift mutations or non-synonymous SNPs in T4SS genes. However, by investigating genes with enriched non-synonymous SNPs, we identified gene variants of the *TrbE* conjugal transfer protein in *Variovorax* that arose following phage treatment. *TrbE* physically interacts with the T4SS and is a member of the *VirB4* family of proteins responsible for substrate recruitment and transfer to the secretion

364 channel of T4SS (Li et al., 2018; Redzej et al., 2022). We observed two non-synonymous SNPs
365 within the P-loop domain of this protein which is responsible for ATP binding and hydrolysis. We
366 speculate that the *TrbE* mutations interfere with the functioning of the T4SS required for phage
367 infection.
368
In addition to *trbE* variants, we observed non-synonymous SNPs in a small-protein-coding
(DUF932) gene within a putative integrated plasmid in all phage-treated *Variovorax* populations.
Although the precise role of this gene is unclear, its consistent selection after phage application
strongly suggests involvement in phage defence. The blocking of phage entry by resident mobile
genetic elements, including phage, is well-documented (Kliem & Dreiseikelmann, 1989; Berry and
Christie, 2011; Kamruzzaman et al., 2022; Rivard et al., 2024; Johnson et al., 2022; Taylor et al.,
2024). For example, the *pVII* minor capsid protein in *Pseudomonas* phage *Pf4*, interacts directly
with the host *PilC* protein, responsible for T4SS assembly in *Pseudomonas*, rendering it inactive
and preventing phage superinfection (Wang et al., 2022). We suspect that *Variovorax* developed
phage resistance through both chromosomal and mobile element-mediated modulation of the
T4SS. These mutations are likely to have existed within the *Variovorax* populations prior to phage
application and highlight the role of population variation in surviving phage infection.
381
The addition of a single or multiple phages showed little difference in *Variovorax* depletion or its
rate of recovery. This absence of additive effects has been observed in other studies (Pyenson
et al. 2024), including efforts to remove *Ralstonia* from tomato plants (Franco Ortega et al., 2024).
In our study, this may have occurred to all three *Variovorax*-targeting phages infecting via the
same receptor. In future studies, the use of multiple phages targeting different receptors would
likely be beneficial, as has been shown in phage-therapy research treating bacterial infections in
humans (Yoo et al., 2024). However, complete bacterial removal is rare and, in clinical cases, has
required the simultaneous use of antibiotics (Pirnay et al., 2024). Notably, in the *Ralstonia*-
infected tomato plant studied by Wang et al. (2019), the addition of phages resulted in the
enrichment of taxa that naturally suppressed the growth of *Ralstonia* through antibiotic production.
These studies suggest that phage application alone is unlikely to remove target bacterium long-
term, and that additional ecological or therapeutic pressures are required to sustain suppression
of the target organism. However, as we show here, even partial and temporary depletion of
populations is useful to test interaction hypotheses.
396
Variovorax recovered at variable rates following phage treatment, with communities in which
T4SS knockouts conferred positive fitness exhibiting the fastest recovery. Prior work has shown
that community composition can substantially impact phage-host dynamics (Blazanin and Turner,
2021; Alseth et al. 2019). Phage susceptibility can also be strongly influenced by the natural
variation in physiology of genetically identical cells, such as cell size (Schenk & Sieber, 2019) and
receptor protein copy numbers (Chapman-McQuiston & Wu, 2008; Igler et al., 2022). This leads
us to hypothesize that community- and environmental selection against T4SS expression in
Variovorax in the guar and locust bean gum communities could have led to reduced presentation
of this complex by *Variovorax*, thereby limiting infection by our phages targeting this system. The
observed variation in *Variovorax* recovery highlights how community context could impact phage

infection, and may be an important area of research for the application of phage-based
microbiome manipulations.
409
Here, we establish the basis of specific microbe-microbe interactions in natural enrichment-based
communities by isolating and using phages to selectively deplete a member of interest. This
approach addresses a critical challenge in microbial ecology by easily enabling the direct testing
of interaction hypotheses within natural consortia, including interactions involving taxa that are
otherwise difficult to culture or manipulate. This method is scalable, cost-effective and uses
standard microbiology techniques without requiring specialized equipment. Although suitable
phage isolation does require the isolation of the microorganism of interest, future integration of
phage engineering with advances in protein design (Pacesa et al., 2025) may circumvent this
current limitation. Overall, phage-based approaches represent valuable tools for understanding
the structure of microbial communities and offer promising avenues for microbiome engineering.

Methodology

Bacterial inoculum preparation

Variovorax was grown overnight from a single colony in R2A media. 100 ml of planktonic phase
cells were recovered from two bench-scale thiocyanate-degrading bioreactor systems, from which
Variovorax was initially isolated a year prior. Flocculated particles were allowed to settle to the
bottom of the container for 30 min and the top 50 ml were carefully removed. 10 ml of bioreactor
planktonic phase were added into 50 ml falcon tubes. 5 ml of *Variovorax* overnight culture was
added to this falcon tube. 15 ml of 50% glycerol was then added and mixed by swirling and
inverting each tube. 200 uL of each culture were distributed into cryo tubes, frozen in liquid
nitrogen, and stored at -80C. This approach was based on previous inoculum ratio testing
experiments (Gao et al., 2021), which demonstrated that a higher abundance of a member in the
inoculum improves its chances of establishing within the community.

Community passaging

Inoculum glycerol stocks were thawed on ice and gently mixed by pipetting. 50 uL of glycerol
stock was added to 10 ml of M9 minimal media in quadruplicate supplemented with 50 ug/L
cycloheximide, 10 mg/L pantothenic acid, and a trace metal solution described by Watts et al.
(2019). Culturing was performed using four independent technical replicates to provide
robustness for statistical tests. The electron donors used were 0.2 g/L of kojac gum, locust bean
gum, guar gum, and combined 0.1 g/L guar gum and 0.1 g/L arabinogalactan. Under each
electron donor condition, Following inoculation, cultures were incubated at 30C for 48 hours with
shaking at 140 rpm. Cultures were passaged using 200 uL of spent medium into 10ml of fresh
media. This was repeated every 48 hours for a total of seven passages over 14 days. This
culturing protocol was heavily informed by Goldford et al. (2018). The glycerol stocks were then
used to inoculate fresh tubes followed by two passages as described above. At each passaging
step, 8 ml of culture were centrifuged at 10 000 X g for 10 minutes and cell pellets were frozen at

-80C for subsequent DNA extraction. 200 uL of spent medium was used to perform OD
measurements using a 96-well plate reader at 600 nm.

Virome preparations

Three sources of viromes were collected: (i) soils, (ii) composts and (iii) and thiocyanate-
degrading bioreactors. Briefly, approximately 10 grams of soil or compost were collected in falcon
tubes and incubated with 25 ml of SM buffer with shaking for one hour. In the case of bioreactor
viromes, biofilm material was removed from the bioreactors using serological pipettes and
subjected to metal bead beating on a vortex at maximum speed for ten minutes to disrupt the
biofilm. For all three types of viromes, the particulate matter was then allowed to settle for 30
minutes before the supernatant was removed and centrifuged at 10 000 X g for 10 minutes and
the supernatant then passed through a 0.22 um filter.

Phage liquid enrichment

Bacteria were grown overnight from single colonies in 10 ml of R2A media and incubated at 30C
at 160 rpm. Cultures were back diluted into fresh R2A containing 2 mM MgCl2 and 2 mM CaCl2
to an OD600 of 0.03-0.05. Cultures were allowed to grow until reaching an OD600 of 0.15-0.3.
Subsequently, 100 uL of viromes were then added to the tubes and the cultures were incubated
for a further 24 hours. The cultures were then sampled and cells were removed by centrifugation
at 10 000 X g for 10 minutes. This supernatant served as the virome for the next round of
enrichment. This process was repeated for a total of five enrichments.

Phage isolation: Plaque assays

Plaque assays were performed on the *Variovorax* strains *V.* SCN45, *V.* CL14, *V. boronicumulans*
(DSM 21722), *V. ginsengisoli* DSM 25185 and *V. paradoxus* DSM 645. These strains were grown
overnight from single colonies in 10 ml of R2A media and incubated at 30C at 160 rpm. Cultures
were back diluted into fresh R2A to an OD600 of 0.03-0.05. Cultures were allowed to grow until
reaching an OD600 of 0.15 - 0.3. Subsequently, 100uL of log-phase cells were added to 20 uL of
0.2 um filter-sterilized viromes and incubated at RT for 10 minutes. These were mixed with 50C
R2A (2 mM MgCl2 and 2 mM CaCl2) 0.5% top agar and plated onto R2A 1.5% bottom agar plates.
Plates were incubated at 30C for 48 hours and inspected for plaques. Plaques were picked using
sterile 100 uL tips into 100 uL of SM buffer.

Phage genome sequencing, assembly and annotation

DNA from phage suspensions were extracted using Zymo Quick-DNA viral kits using
manufacturers instructions. Extracted DNA was then illumina shotgun sequenced on a Novaseq
using 150PE reads with a 300 bp insert size and yielding 1 Gbp per sample. Viral genomes were
assembled using Metaviral-Spades (Antipov et al., 2020) and extended using COBRA (Chen et
al., 2024). The genomes were then annotated using vogDB (Trgovec-Greif et al., 2024), Pfam
(Bateman et al., 2004) and KEGG (Kanehisa & Goto, 2000) hidden markov models using
hmmscan. Protein sharing of the nine phage and Refseq and was performed using Vcontact2

(Bin-Jang et al., 2019). Phage genome ANI was determined using Vclust (Zielezinski et al., 2024;
https://github.com/refresh-bio/vclust).

Transmission electron microscopy (TEM)

485 Virus particles were visualized using a FEI Tecnai T12 Transmission Electron Microscope at an
486 accelerating voltage of 80 kV. Carbon-coated copper grids were charged in a vacuum chamber
487 followed by application of 5uL of virus suspension and allowed to dry. The grids were washed
488 with deionised water three times before negative staining in 1.0 % ammonium acetate for 60s.
489 Grids were allowed to air dry before being visualized.
490

Arabidopsis inoculation and co-cultures with Variovorax CL14, Arthrobacter CL28, and V.cl_11+V.cl_23 phage for root length phenotyping

Arabidopsis thaliana Col-0 seeds were surface-sterilized in 50% bleach for 20 min with 0.05%
Tween 20 for 20 min, removing all bleach and rinsing five times in sterile water. Sterile seeds
were stratified in the dark for 2 to 3 days at 4°C. A nylon mesh filter with 100 micron pore size
was cut to fit on the 100 mm x 100 mm square petri dish as previously described (Cole et al.
2017). After stratification, ~7 seeds were placed on the mesh filter on 1.0MS, 0.5% sucrose, 0.6%
phytagel plates and grown under long day conditions for 7d (16-h light/8-h dark regime at 22-°C).
Bacteria strains preparation: *Variovorax* CL14 and the root growth inhibiting strain Arthrobacter
CL28 were streaked on agar plates from glycerol stocks, a single colony was transferred to 3 ml
liquid R2A media and grown overnight. Cultures were gently pelleted then washed twice with 10
mM MgCl2, normalizing the volume of each culture to OD600 = 1, then adjust to OD600=0.04
before spreading 50 µl on each plate (Full MS, 0.6% phytagel without sucrose. Note: Full MS was
used instead of 1/2X as previously described to maintain the required Mg and Ca concentrations
for optimal phage lysis (2mM)). To prepare the phage inoculation, 5ul of a phage cocktail
(Siphoviridae phage V.cl_11+V.cl_23) was diluted in 500 ul 20 mM MgCl2 and 100 ul of this
solution was spread onto each plate. After the plants grew 7d, the mesh was transferred to each
of the bacteria and phage inoculated plates and returned to the growth chamber vertically grown
in short day conditions (16-h dark/8-h light regime at 22-°C). Plates were imaged 10 days after
transfer. Primary root length was measured using ImageJ software, first setting the scale to the
grid on the plate (13 mm) before measuring primary root elongation for each plant. There were 3
replicate plates (containing ~ plants each) for each control and treatment: uninoculated, no
bacteria + V11/V23 phage, *Variovorax* CL14, *Arthrobacter* CL28, *Variovorax* CL14 + *Arthrobacter*
CL28, *Variovorax* CL14 + *Arthrobacter* CL28 + V11/V23 phage.

Phage-perturbed community experiments

Stable communities grown on selected carbon sources and were exposed to *Variovorax*-specific
phages (V45_v66, or a three-phage cocktail of V45_v66, V.45_iii and V.45_bapic4) 8 hours after
the passaging of the day 8 communities (ie. following passage 5), using 50 uL of 4×10^4 PFU/uL
each per 10 ml culture. Experiments were conducted in quadruplicate with a no-phage control
included for comparison. The final varied condition involved the presence or absence of 10 mg/L

thiamine, administered at the time of phage addition and maintained during subsequent
passages.

DNA extraction, amplicon and shotgun sequencing

Genomic DNA was extracted from cell pellets using the Qiagen DNeasy PowerSoil Pro Kit
according to the manufacturer's instructions. V4-V5 16S rRNA gene amplicons were generated
by PCR using primers 515F (GTGNCAGCMGCCGCGGTAA) and 926R
(CCGYCAATTYMTTTRAGTTT) and cycling conditions 95C 10 min, 72C 90s, 50C 60s, 72C 90s
for 30 cycles and a final extension at 72C for 10 min. PCR products were cleaned using Ampure
XP magnetic beads. PCR products were sequenced by Innovative Genomic Institute's Next
Generation Sequencing Core on an Illumina MiSeq instrument using a read length of 2 x 150 bp
with approximately 50 000 reads per sample. Generated amplicon sequencing data was
processed yielding ASVs as performed by Kust et al. (2023). Whole genome community library
preparation and shotgun sequencing was performed by Novogene. Paired-end Illumina libraries
were prepared from genomic DNA with an insert size of 350 bp and sequenced on an Illumina
Novaseq producing approximately 6 Gbp of raw sequencing data per sample. The *Variovorax*
isolate was subsequently sequenced using PacBio long-read sequencing at 60x coverage, to
generate two circular chromosomes.
538

Metagenomic Read Processing, Assembly and binning

Illumina 150 PE reads from community shotgun sequencing were processed as documented in
the ggKbase data preparation pipeline (https://ggkbase-help.berkeley.edu/overview/data-
preparation-metagenome/). Reads were processed using BBtools (Bushnell 2020) and sickle
(Joshi and Fass, 2011), and were then assembled using IDBA-UD (Peng et al., 2012). Open
reading frames (ORFs) were predicted using Prodigal V2.6.3 (Hyatt et al., 2010). Contigs were
binned using CONCOCT (Alneberg et al., 2013), Maxbin2 (Wu et al., 2016), MetaBat2 (Kang et
al., 2019), and VAMB (Nissen et al., 2021). The best bins per sampled were selected using DAS
tools (Sieber et al., 2018) and a final non-redundant genome set across all enrichment
communities and the previously published Thiocyanate bioreactor genome set found at
https://ggkbase.berkeley.edu/SCN_92/organisms was selected using dRep (Olm et al., 2017) at
95% ANI.

Microbial genome annotation, classification and strain analysis

Microbial genomes were then annotated using hmmscan against Pfam (Bateman et al., 2004)
and KEGG (Kanehisa & Goto, 2000) databases and were taxonomically classified using GTDB-
tk (Chaumeil et al., 2020). All read mappings were performed with bowtie2 (Langmead et al.,
2012). Genome coverage and relative abundance across samples were determined using coverm
(Aroney et al., 2025). Genome-genome correlation was determined using Fastspar (Watts et al.,
2019) which implements the Sparcc algorithm (Friedman et al., 2012). *Variovorax* strain analysis
was performed using inStrain (Olm et al., 2021) and the PacBio assembled *Variovorax* genome.
Phage defence systems were predicted in the *Variovorax* genome using padlock (Payne et al.,

2022, https://padloc.otago.ac.nz/padloc/). Predicted Protein structure predictions were performed
using Alphafold3 (Abramson et al., 2024. https://alphafoldserver.com/)

Statistical analyses

Distributions in organism abundances were compared by two-sample t-test assuming unequal
variance was performed using the ttest_ind function of the scipy.stats Python package. The
abundance of ASVs was calculated excluding the abundance of *Variovorax* contributing to 100%
abundance. Relative abundance data were normalized by robust centered log-ratio (rclr)
transformation using the decostand function from the vegan R package (Oksanen et al., 2001).
Principal component analysis (PCA) was performed on the normalized abundance data using the
prcomp function in R to visualize variation in community composition across samples.

RB-TnSeq library construction

A DNA-barcoded Tn5 transposon mutant library of *Variovorax* SCN45 was generated as
described previously (Wetmore et al., 2015). Briefly, an *E. coli* strain (WM3064) carrying the
transposon vector library pKMW3 was conjugated with wild-type *Variovorax* SCN45 at a 1:1 ratio
on R2A agar plates supplemented with diaminopimelic acid (50 ug/ml). Following 6 hours of
incubations, the transformants were plated onto R2A agar plates containing 100 μg/ml kanamycin.
Several thousand *Variovorax* colonies were scraped and collected into a tube before being
resuspended in 100 ml of R2A with Kanamycin and incubated at 30 C for 6 hours. Glycerol stocks
were made at a 1:1 ratio of mid-log phase Variovorax cells and 25% glycerol, flash frozen in liquid
nitrogen, and stored at -80C. Simultaneously 10 ml of mid-log phase cells were pelleted through
centrifugation and their DNA extracted and sequenced by TnSeq-like Illumina sequencing
protocol.

RB-TnSeq library incubations

A 2 ml mutant library glycerol stock was thawed and inoculated into 25 ml of R2A with 50 ug/ml
Kanamycin and incubated at 30°C with shaking until cells entered mid log phage (approximately
8 hours). Four tubes of 2 ml of culture, termed time zero controls, were then recovered and cells
pelleted by centrifugation and frozen at -80°C. The remaining cultured cells were then washed
twice in relevant media (M9 media without carbon source or R2A) with spins at 3000g for 3
minutes. The cells were resuspended in media and inoculated into relevant or culture to a final
OD600 of 0.02. The cultures were grown in 10ml glass tubes and incubated at 30 °C with shaking.
(i) For phage-challenge experiments 10 uL of phage (4 x10^4 PFU/ul) were added to the libraries
after 4 hours of subsequent incubation and allowed to incubate for a total of 24 h. (ii) For
community incubations, cultures were passaged every 48 h for four passages. Cells were
recovered from four replicate cultures by centrifugation and stored at -80 °C.Genomic DNA was
then extracted from the stored cell pellets and PCR and sequencing by BarSeq performed as
described previously (Wetmore et al., 2015).A 2 ml mutant library glycerol stock was thawed and
inoculated into 25 ml of R2A with 50 ug/ml Kanamycin and incubated at 30°C with shaking until
cells entered mid log phage (approximately 8 hours). Four tubes of 2 ml of culture, termed time
zero controls, were then recovered and cells pelleted by centrifugation and frozen at -80°C. The

remaining cultured cells were then washed twice in relevant media (M9 media without carbon source or R2A) with spins at 3000g for 3 minutes. The cells were resuspended in media and inoculated into relevant or culture to a final OD600 of 0.02. The cultures were grown in 10ml glass tubes and incubated at 30 °C with shaking. (i) For phage-challenge experiments 10 uL of phage (4 x10^4 PFU/ul) were added to the libraries after 4 hours of subsequent incubation and allowed to incubate for a total of 24 h. (ii) For community incubations, cultures were passaged every 48 h for four passages. Cells were recovered from four replicate cultures by centrifugation and stored at - 80 °C.Genomic DNA was then extracted from the stored cell pellets and PCR and sequencing by BarSeq performed as described previously (Wetmore et al., 2015).

Data availability

The contigs representing the Variovorax genome assembled by PacBio can be found under bioproject PRJNA1212206 genome accessions CP179718-CP179720. The assembled genomes of the isolated phage have been deposited in GenBank under accession numbers PV558788 - PV558796. The reads associated with these genomes and assemblies can be found under accession numbers SRR33449703 - SRR33449695. The 16S rRNA amplicon sequencing data can be found under Bioproject PRJNA1254325, under accession numbers SRR33264975 - SRR33264785. Shotgun sequencing can be found under bioproject PRJNA1254232, including reads (SRR33436150-SRR33436076), assemblies (SAMN48138949 - SAMN48139025), and novel MAGs (SAMN48122594 - SAMN48122609). These MAGs can also be found on ggkbase https://ggkbase.berkeley.edu/Phage_treated_Variovorax_enrichments_dereplicated_MAGs/ and MAGs detected in this study that had been previously published can be found on ggkbase https://ggkbase.berkeley.edu/SCN_92/.

Acknowledgments

This material by m-CAFEs Microbial Community Analysis and Functional Evaluation in Soils, (m-CAFEs@lbl.gov) a Science Focus Area led by Lawrence Berkeley National Laboratory is based upon work supported by the U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research under contract number DE-AC02-05CH11231. Thank you to the staff at the UC Berkeley Electron Microscope Laboratory for advice and assistance in electron microscopy sample preparation and data collection. Thanks to Leylen Miloslavich, Marta Ortega and Kaitlin Creamer for assistance with lab operations, Henrik V. Scheller and Benjamin J Cole for *Arabidopsis* seeds, and to Jeffery L. Dangl for the *Variovorax CL14* strain and Rebecca S Bart for the *Arthrobacter CL28* isolates.

References

Alneberg, Johannes, Brynjar Smári Bjarnason, Ino de Bruijn, Melanie Schirmer, Joshua Quick, Umer Z. Ijaz, Nicholas J. Loman, Anders F. Andersson, and Christopher Quince. "CONCOCT: clustering contigs on coverage and composition." *arXiv preprint arXiv:1312.4038* (2013).

Alseth, E. O., Pursey, E., Luján, A. M., McLeod, I., Rollie, C., & Westra, E. R. (2019). Bacterial biodiversity drives the evolution of CRISPR-based phage resistance. Nature, 574(7779), 549-552.

Antipov, Dmitry, Mikhail Raiko, Alla Lapidus, and Pavel A. Pevzner. "Metaviral SPAdes: assembly of viruses from metagenomic data." Bioinformatics 36, no. 14 (2020): 4126-4129.

Bateman, Alex, Lachlan Coin, Richard Durbin, Robert D. Finn, Volker Hollich, Sam Griffiths-Jones, Ajay Khanna et al. "The Pfam protein families database." *Nucleic acids research* 32, no. suppl_1 (2004): D138-D141.

Bikard, David, Chad W. Euler, Wenyan Jiang, Philip M. Nussenzweig, Gregory W. Goldberg, Xavier Duportet, Vincent A. Fischetti, and Luciano A. Marraffini. "Exploiting CRISPR-Cas nucleases to produce sequence-specific antimicrobials." Nature biotechnology 32, no. 11 (2014): 1146-1150.

Bin Jang, Ho, Benjamin Bolduc, Olivier Zablocki, Jens H. Kuhn, Simon Roux, Evelien M. Adriaenssens, J. Rodney Brister et al. "Taxonomic assignment of uncultivated prokaryotic virus genomes is enabled by gene-sharing networks." *Nature biotechnology* 37, no. 6 (2019): 632-639.

Blazanin, M., & Turner, P. E. (2021). Community context matters for bacteria-phage ecology and evolution. *The ISME Journal*, *15*(11), 3119-3128.

Abramson, Josh, Jonas Adler, Jack Dunger, Richard Evans, Tim Green, Alexander Pritzel, Olaf Ronneberger et al. "Accurate structure prediction of biomolecular interactions with AlphaFold 3." Nature (2024): 1-3.

Berry, Trista M., and Peter J. Christie. "Caught in the act: the dialogue between bacteriophage R17 and the type IV secretion machine of plasmid R1." *Molecular microbiology* 82, no. 5 (2011): 1039-1043.

Bushnell, B. (2020). BBMap. sourceforge. net/projects/bbmap.https://cir.nii.ac.jp/crid/1370294643771707027

Buttimer, Colin, Tom Sutton, Joan Colom, Ellen Murray, Pedro H. Bettio, Linda Smith, Andrei S. Bolocan et al. "Impact of a phage cocktail targeting Escherichia coli and Enterococcus faecalis as members of a gut bacterial consortium in vitro and in vivo." Frontiers in Microbiology 13 (2022): 936083.

Carlson, H.K., Piya, D., Moore, M.L., Magar, R.T., Elisabeth, N.H., Deutschbauer, A.M., Arkin, A.P. and Mutalik, V.K., 2023. Geochemical constraints on bacteriophage infectivity in terrestrial environments. *ISME communications*, *3*(1), p.78.

Chapman-McQuiston, E., and X. L. Wu. "Stochastic receptor expression allows sensitive bacteria to evade phage attack. Part I: experiments." Biophysical journal 94, no. 11 (2008): 4525-4536.

682   Chaumeil, Pierre-Alain, Aaron J. Mussig, Philip Hugenholtz, and Donovan H. Parks. "GTDB-Tk: a toolkit
683   to classify genomes with the Genome Taxonomy Database." (2020): 1925-1927.
684
685   Chen, LinXing, and Jillian F. Banfield. "COBRA improves the completeness and contiguity of viral genomes
686   assembled from metagenomes." Nature Microbiology 9, no. 3 (2024): 737-750.
687
688   Clegg, Tom, and Thilo Gross. "Cross-feeding creates tipping points in microbiome diversity." Proceedings
689   of the National Academy of Sciences 122, no. 19 (2025): e2425603122.
690
691   Cole, Benjamin J., Meghan E. Feltcher, Robert J. Waters, Kelly M. Wetmore, Tatiana S. Mucyn, Elizabeth
692   M. Ryan, Gaoyan Wang et al. "Genome-wide identification of bacterial plant colonization genes." PLoS
693   Biology 15, no. 9 (2017): e2002860.
694
695   Das, Mayukh, Tushar Suvra Bhowmick, Stephen J. Ahern, Ry Young, and Carlos F. Gonzalez. "Control of
696   Pierce's disease by phage." *PLoS One* 10, no. 6 (2015): e0128902.
697
698   Dukovski, Ilija, Djordje Bajić, Jeremy M. Chacón, Michael Quintin, Jean CC Vila, Snorre Sulheim, Alan R.
699   Pacheco et al. "A metabolic modeling platform for the computation of microbial ecosystems in time and
700   space (COMETS)." *Nature protocols* 16, no. 11 (2021): 5030-5082.
701
702   Franco Ortega, S., Fields, B., Narino Rojas, D., Mikonranta, L., Holmes, M., Harper, A. L., & Friman, V. P.
703   (2024). Phage biocontrol success of bacterial wilt depends on synergistic interactions with resident
704   rhizosphere microbiota. Microbial Biotechnology, 17(11), e70049.
705
706   Finkel, Omri M., Isai Salas-González, Gabriel Castrillo, Jonathan M. Conway, Theresa F. Law, Paulo José
707   Pereira Lima Teixeira, Ellie D. Wilson, Connor R. Fitzpatrick, Corbin D. Jones, and Jeffery L. Dangl. "A
708   single bacterial genus maintains root growth in a complex microbiome." *Nature* 587, no. 7832 (2020): 103-
709   108.
710
711   Frias-Lopez, Jorge, Yanmei Shi, Gene W. Tyson, Maureen L. Coleman, Stephan C. Schuster, Sallie W.
712   Chisholm, and Edward F. DeLong. "Microbial community gene expression in ocean surface waters."
713   Proceedings of the National Academy of Sciences 105, no. 10 (2008): 3805-3810.
714
715   Friedman, Jonathan, and Eric J. Alm. "Inferring correlation networks from genomic survey data." (2012):
716   e1002687.
717
718   Gao, Chun-Hui, Hui Cao, Peng Cai, and Søren J. Sørensen. "The initial inoculation ratio regulates bacterial
719   coculture interactions and metabolic capacity." *The ISME journal* 15, no. 1 (2021): 29-40.
720
721   Gilbert, Jack A., Dawn Field, Ying Huang, Rob Edwards, Weizhong Li, Paul Gilna, and Ian Joint. "Detection
722   of large numbers of novel sequences in the metatranscriptomes of complex marine microbial communities."
723   *PloS one* 3, no. 8 (2008): e3042.
724
725   Goldford, Joshua E., Nanxi Lu, Djordje Bajić, Sylvie Estrela, Mikhail Tikhonov, Alicia Sanchez-Gorostiaga,
726   Daniel Segrè, Pankaj Mehta, and Alvaro Sanchez. "Emergent simplicity in microbial community assembly."
727   Science 361, no. 6401 (2018): 469-474.
728

Gencay, Y.E., Jasinskytė, D., Robert, C., Semsey, S., Martínez, V., Petersen, A.Ø., Brunner, K., de
Santiago Torio, A., Salazar, A., Turcu, I.C. and Eriksen, M.K., 2024. Engineered phage with antibacterial
CRISPR–Cas selectively reduce E. coli burden in mice. Nature Biotechnology, 42(2), pp.265-274.

Han, Jong-In, Hong-Kyu Choi, Seung-Won Lee, Paul M. Orwin, Jina Kim, Sarah L. LaRoe, Tae-gyu Kim et
al. "Complete genome sequence of the metabolically versatile plant growth-promoting endophyte
Variovorax paradoxus S110." *Journal of bacteriology* 193, no. 5 (2011): 1183-1190.

Hessler, Tomas, Robert J. Huddy, Rohan Sachdeva, Shufei Lei, Susan TL Harrison, Spencer Diamond,
and Jillian F. Banfield. "Vitamin interdependencies predicted by metagenomics-informed network analyses
and validated in microbial community microcosms." Nature Communications 14, no. 1 (2023): 4768.

Hyatt, Doug, Gwo-Liang Chen, Philip F. LoCascio, Miriam L. Land, Frank W. Larimer, and Loren J.
Hauser. "Prodigal: prokaryotic gene recognition and translation initiation site identification." *BMC*
bioinformatics 11 (2010): 1-11.

Igler, C., 2022. Phenotypic flux: The role of physiology in explaining the conundrum of bacterial
persistence amid phage attack. *Virus evolution*, *8*(2), p.veac086.

Johnson, Christopher M., M. Michael Harden, and Alan D. Grossman. "Interactions between mobile genetic
elements: An anti-phage gene in an integrative and conjugative element protects host cells from predation
by a temperate bacteriophage." PLoS Genetics 18, no. 2 (2022): e1010065.

Joshi, N. A.; Fass, J. N. (**2011**). *Sickle: A sliding-window, adaptive, quality-based trimming tool for FastQ*
files. (Version 1.33) [Software], https://github.com/najoshi/sickle.

Kamruzzaman, Muhammad, Amy J. Mathers, and Jonathan R. Iredell. "A Novel Plasmid Entry Exclusion
System in pKPC_UVA01, a Promiscuous Conjugative Plasmid Carrying the bla KPC Carbapenemase
Gene." Antimicrobial Agents and Chemotherapy 66, no. 3 (2022): e02322-21.

Kanehisa, M., & Goto, S. (2000). KEGG: kyoto encyclopedia of genes and genomes. *Nucleic acids*
research, *28*(1), 27-30.

Kang, F. Li, E. Kirton, A. Thomas, R. Egan, H. An, Z. Wang, MetaBAT 2: an adaptive binning algorithm for
robust and efficient genome reconstruction from metagenome assemblies. PeerJ 7, e7359 (2019).

Kost, Christian, Kiran Raosaheb Patil, Jonathan Friedman, Sarahi L. Garcia, and Markus Ralser. "Metabolic
exchanges are ubiquitous in natural microbial communities." Nature Microbiology 8, no. 12 (2023): 2244-
2252.

Kliem, Manfred, and Brigitte Dreiseikelmann. "The superimmunity gene sim of bacteriophage P1 causes
superinfection exclusion." *Virology* 171, no. 2 (1989): 350-355.

Kust, Andreja, Jackie Zorz, Catalina Cruañas Paniker, Keith Bouma-Gregson, Netravathi Krishnappa,
Jillian F. Banfield, and Spencer Diamond. "Model cyanobacterial consortia reveal a consistent core
microbiome independent of inoculation source or cyanobacterial host species." bioRxiv (2023): 2023-12.

Langmead, Ben, and Steven L. Salzberg. "Fast gapped-read alignment with Bowtie 2." *Nature methods* 9,
no. 4 (2012): 357-359.

Li, Y., Zhu, F., Li, Y., Pan, S., Wang, H., Yang, Z., ... & Liu, B. (2022). Bacteriophages allow selective
depletion of gut bacteria to produce a targeted-bacterium-depleted mouse model. *Cell Reports Methods*,
2(11).
782

783   Li, Yang Grace, and Peter J. Christie. "The Agrobacterium VirB/VirD4 T4SS: mechanism and architecture
784   defined through in vivo mutagenesis and chimeric systems." *Agrobacterium Biology: From Basic Science*
785   *to Biotechnology* (2018): 233-260.
786

787   Liao, H., Wen, C., Huang, D., Liu, C., Gao, T., Du, Q., ... & Friman, V. P. (2025). Harnessing phage
788   consortia to mitigate the soil antibiotic resistome by targeting keystone taxa Streptomyces. *Microbiome*,
789   *13*(1), 1-20.
790

791   Ma, Bin, Yiling Wang, Shudi Ye, Shan Liu, Erinne Stirling, Jack A. Gilbert, Karoline Faust et al. "Earth
792   microbial co-occurrence network reveals interconnection pattern across microbiomes." *Microbiome* 8, no.
793   1 (2020): 1-12.
794

795   Mutalik, V. K., Adler, B. A., Rishi, H. S., Piya, D., Zhong, C., Koskella, B., ... & Arkin, A. P. (2020). High-
796   throughput mapping of the phage resistance landscape in E. coli. *PLoS biology*, *18*(10), e3000877.
797

798   Nissen, Jakob Nybo, Joachim Johansen, Rosa Lundbye Allesøe, Casper Kaae Sønderby, Jose Juan
799   Almagro Armenteros, Christopher Heje Grønbech, Lars Juhl Jensen et al. "Improved metagenome
800   binning and assembly using deep variational autoencoders." *Nature biotechnology* 39, no. 5 (2021): 555-
801   560.
802

Oksanen, G. L. Simpson, F. G. Blanchet, R. Kindt, P. Legendre, P. R. Minchin, R. B. O'Hara, P. Solymos,
M. H. H. Stevens, E. Szoecs, H. Wagner, M. Barbour, M. Bedward, B. Bolker, D. Borcard, G. Carvalho,
M. Chirico, M. De Caceres, S. Durand, H. B. A. Evangelista, R. FitzJohn, M. Friendly, B. Furneaux, G.
Hannigan, M. O. Hill, L. Lahti, D. McGlinn, M.-H. Ouellette, E. Ribeiro Cunha, T. Smith, A. Stier, C. J. F.
Ter Braak, J. Weedon, vegan: Community Ecology Package, (2001);
https://doi.org/10.32614/CRAN.package.vegan.
809

810   Olm, Matthew R., Christopher T. Brown, Brandon Brooks, and Jillian F. Banfield. "dRep: a tool for fast
811   and accurate genomic comparisons that enables improved genome recovery from metagenomes through
812   de-replication." *The ISME journal* 11, no. 12 (2017): 2864-2868.
813

814   Olm, Matthew R., Alexander Crits-Christoph, Keith Bouma-Gregson, Brian A. Firek, Michael J. Morowitz,
815   and Jillian F. Banfield. "inStrain profiles population microdiversity from metagenomic data and sensitively
816   detects shared microbial strains." *Nature Biotechnology* 39, no. 6 (2021): 727-736.
817

818   Pacheco, Alan R., and Daniel Segrè. "A multidimensional perspective on microbial interactions." FEMS
819   microbiology letters 366, no. 11 (2019): fnz125.
820

821   Pacesa, M., Nickel, L., Schellhaas, C., Schmidt, J., Pyatova, E., Kissling, L., Barendse, P., Choudhury, J.,
822   Kapoor, S., Alcaraz-Serna, A. and Cho, Y., 2024. BindCraft: one-shot design of functional protein binders.
823   bioRxiv, pp.2024-09.
824

Payne L.J., Meaden S., Mestre M.R., Palmer C., Toro N., Fineran P.C. and Jackson S.A.* (2022)
PADLOC: a web server for the identification of antiviral defence systems in microbial genomes.

Nucleic Acids Research 50, W541-W550. doi: 10.1093/nar/gkac400

Peng, Yu, Henry CM Leung, Siu-Ming Yiu, and Francis YL Chin. "IDBA-UD: a de novo assembler for
single-cell and metagenomic sequencing data with highly uneven depth." Bioinformatics 28, no. 11
(2012): 1420-1428.

Pierce, Emily C., and Rachel J. Dutton. "Putting microbial interactions back into community contexts."
Current Opinion in Microbiology 65 (2022): 56-63.

Pirnay, Jean-Paul, Sarah Djebara, Griet Steurs, Johann Griselain, Christel Cochez, Steven De Soir, Tea
Glonti et al. "Personalized bacteriophage therapy outcomes for 100 consecutive cases: a multicentre,
multinational, retrospective observational study." Nature Microbiology (2024): 1-20.

Pyenson, N.C., Leeks, A., Nweke, O., Goldford, J.E., Schluter, J., Turner, P.E., Foster, K.R. and Sanchez,
A., 2024. Diverse phage communities are maintained stably on a clonal bacterial host. *Science*, *386*(6727),
pp.1294-1300.

Quinn-Bohmann, Nick, Tomasz Wilmanski, Katherine Ramos Sarmiento, Lisa Levy, Johanna W. Lampe,
Thomas Gurry, Noa Rappaport et al. "Microbial community-scale metabolic modelling predicts personalized
short-chain fatty acid production profiles in the human gut." *Nature Microbiology* 9, no. 7 (2024): 1700-1712.

Quinones-Olvera, Natalia, Siân V. Owen, Lucy M. McCully, Maximillian G. Marin, Eleanor A. Rand, Alice
C. Fan, Oluremi J. Martins Dosumu et al. "Diverse and abundant phages exploit conjugative plasmids."
Nature Communications 15, no. 1 (2024): 3197.

Ram, Rachna J., Nathan C. VerBerkmoes, Michael P. Thelen, Gene W. Tyson, Brett J. Baker, Robert C.
Blake, Manesh Shah, Robert L. Hettich, and Jillian F. Banfield. "Community proteomics of a natural
microbial biofilm." *Science* 308, no. 5730 (2005): 1915-1920.

Ramirez, Jaime, Francisco guarner, Luis Bustos Fernandez, Aldo Maruy, Vera Lucia Sdepanian, and Henry
Cohen. "Antibiotics as major disruptors of gut microbiota." Frontiers in cellular and infection microbiology
10 (2020): 572912.

Redzej, Adam, Marta Ukleja, Sarah Connery, Martina Trokter, Catarina Felisberto-Rodrigues, Adam Cryar,
Konstantinos Thalassinos, Richard D. Hayward, Elena V. Orlova, and Gabriel Waksman. "Structure of a
VirD4 coupling protein bound to a VirB type IV secretion machinery." *The EMBO journal* 36, no. 20 (2017):
3080-3095.

Rivard, Nicolas, Malika Humbert, Kévin T. Huguet, Aurélien Fauconnier, César Pérez Bucio, Eve Quirion,
and Vincent Burrus. "Surface exclusion of IncC conjugative plasmids and their relatives." Plos Genetics 20,
no. 10 (2024): e1011442.

Russell, Marjorie. "Moving through the membrane with filamentous phages." *Trends in microbiology* 3, no.
6 (1995): 223-228.

Aroney S.T.N.(1), Rhys J. P. Newell (1), Jakob N. Nissen (2), Antonio Pedro Camargo (3 and 4), Gene W.
Tyson (1), Ben J. Woodcroft. (2025). CoverM: Read alignment statistics for metagenomics.
https://doi.org/10.48550/arXiv.2501.11217

Segata, Nicola, Daniela Boernigen, Timothy L. Tickle, Xochitl C. Morgan, Wendy S. Garrett, and Curtis
Huttenhower. "Computational meta'omics for microbial community studies." *Molecular systems biology* 9,
no. 1 (2013): 666.

Schenk, Hanna, and Michael Sieber. "Bacteriophage can promote the emergence of physiologically
sub-optimal host phenotypes." bioRxiv (2019): 621524.

Selle, K., Fletcher, J.R., Tuson, H., Schmitt, D.S., McMillan, L., Vridhambal, G.S., Rivera, A.J.,
Montgomery, S.A., Fortier, L.C., Barrangou, R. and Theriot, C.M., 2020. In vivo targeting of
Clostridioides difficile using phage-delivered CRISPR-Cas3 antimicrobials. MBio, 11(2), pp.10-1128.

Skwara, Abigail, Karna Gowda, Mahmoud Yousef, Juan Diaz-Colunga, Arjun S. Raman, Alvaro Sanchez,
Mikhail Tikhonov, and Seppe Kuehn. "Statistically learning the functional landscape of microbial
communities." Nature ecology & evolution 7, no. 11 (2023): 1823-1833.

Starr, Evan P., Shengjing Shi, Steven J. Blazewicz, Alexander J. Probst, Donald J. Herman, Mary K.
Firestone, and Jillian F. Banfield. "Stable isotope informed genome-resolved metagenomics reveals that
Saccharibacteria utilize microbially-processed plant-derived carbon." *Microbiome* 6 (2018): 1-12

Strathdee, Steffanie A., Graham F. Hatfull, Vivek K. Mutalik, and Robert T. Schooley. "Phage therapy:
From biological mechanisms to future directions." Cell 186, no. 1 (2023): 17-31.

Sieber, Christian MK, Alexander J. Probst, Allison Sharrar, Brian C. Thomas, Matthias Hess, Susannah
G. Tringe, and Jillian F. Banfield. "Recovery of genomes from metagenomes via a dereplication,
aggregation and scoring strategy." *Nature microbiology* 3, no. 7 (2018): 836-843.

Taylor, Véronique L., Megha Shah, Pramalkumar H. Patel, Ahmed Yusuf, Cayla M. Burk, Zemer Gitai,
Alan R. Davidson, Matthias D. Koch, and Karen L. Maxwell. "Prophages block cell surface receptors to
ensure survival of their viral progeny." bioRxiv (2024): 2024-03.

Thanki, Anisha M., Guillaume Mignard, Robert J. Atterbury, Paul Barrow, Andrew D. Millard, and
Martha RJ Clokie. "Prophylactic delivery of a bacteriophage cocktail in feed significantly reduces
Salmonella colonization in pigs." Microbiology spectrum 10, no. 3 (2022): e00422-22.

Trgovec-Greif, Lovro, Hans-Jörg Hellinger, Jean Mainguy, Alexander Pfundner, Dmitrij Frishman, Michael
Kiening, Nicole Suzanne Webster, Patrick William Laffy, Michael Feichtinger, and Thomas Rattei.
"VOGDB—database of virus orthologous groups." *Viruses* 16, no. 8 (2024): 1191.

Tyson, Gene W., Jarrod Chapman, Philip Hugenholtz, Eric E. Allen, Rachna J. Ram, Paul M. Richardson,
Victor V. Solovyev, Edward M. Rubin, Daniel S. Rokhsar, and Jillian F. Banfield. "Community structure and
metabolism through reconstruction of microbial genomes from the environment." *Nature* 428, no. 6978
(2004): 37-43.

Watts, Mathew P., Liam P. Spurr, Kim-Anh Lê Cao, Ryan Wick, Jillian F. Banfield, and John W. Moreau.
"Genome-resolved metagenomics of an autotrophic thiocyanate-remediating microbial bioreactor
consortium." Water research 158 (2019): 106-117.

Watts, Stephen C., Scott C. Ritchie, Michael Inouye, and Kathryn E. Holt. "FastSpar: rapid and scalable
correlation estimation for compositional data." *Bioinformatics* 35, no. 6 (2019): 1064-1066.

Wang, Xiaofang, Shuo Wang, Mingcong Huang, Yilin He, Saisai Guo, Keming Yang, Ningqi Wang et al.
"Phages enhance both phytopathogen density control and rhizosphere microbiome suppressiveness." Mbio
15, no. 6 (2024): e03016-23.

Wang, Xiaofang, Zhong Wei, Keming Yang, Jianing Wang, Alexandre Jousset, Yangchun Xu, Qirong Shen,
and Ville-Petri Friman. "Phage combination therapies for bacterial wilt disease in tomato." *Nature
Biotechnology* 37, no. 12 (2019): 1513-1520.

Wetmore, Kelly M., Morgan N. Price, Robert J. Waters, Jacob S. Lamson, Jennifer He, Cindi A. Hoover,
Matthew J. Blow et al. "Rapid quantification of mutant fitness in diverse bacteria by sequencing randomly
bar-coded transposons." *MBio* 6, no. 3 (2015): 10-1128.

Wu Y.-W. , B. A. Simmons, S. W. Singer, MaxBin 2.0: an automated binning algorithm to recover
genomes from multiple metagenomic datasets. Bioinformatics 32, 605–607 (2016).

Yoo, Seongjun, Kang-Mu Lee, Nayoung Kim, Thao Nguyen Vu, Ricardo Abadie, and Dongeun Yong.
"Designing phage cocktails to combat the emergence of bacteriophage-resistant mutants in multidrug-
resistant Klebsiella pneumoniae." Microbiology spectrum 12, no. 1 (2024): e01258-23.

Zielezinski, Andrzej, Adam Gudys, Jakub Barylski, Krzysztof Siminski, Piotr Rozwalak, Bas E. Dutilh, and
Sebastian Deorowicz. "Ultrafast and accurate sequence alignment and clustering of viral genomes."
bioRxiv (2024): 2024-06.

Supplementary materials

Figure S1. Plaque morphology from spot assays performed on *Variovorax* species with the nine isolated *Variovorax* phages. Only phage V45_66 was able to plaque on more than one host.

978
979 **Figure S2**. Genome-resolved metagenomics reveals consistent reduction in the relative abundance in
980 Variovorax across all four enrichment cultures following phage application. The 16 most abundant
981 organisms per enrichment culture recovered are shown.

982
983
984 **Figure S3.** The optical density (600nm) of the 48 hour old communities following inoculation and each
985 subsequent passage. All communities show a sharp reduction between day 2 and day 4 communities but
986 then reach a stable state from day eight onwards.

993
994 **Figure S4.** *Leifsonia* MAG relative abundance is consistent with the ASV data. The abundance of this MAG
995 is shown across the four carbon source enrichment cultures before phage addition (day 6 and day 8
996 communities), following phage infection (day 10, day 12, day 14 communities), and following phage
997 infection with supplementation of thiamine.
998
999
1000

1001
1002   **Figure S5.** Haplotype frequencies associated with phage exposure to each of the four enrichment
1003   communities and their no-phage controls.

1004
1005 **Figure S6.** Thiamine biosynthesis pathway illustrating the multiple routes for the production of the thiazole
1006 and pyrimidine moieties of thiamine. Adapted from https://www.kegg.jp/ (Kanehisa & Goto, 2000).
1007