1

Context-dependent oviposition reveals strong association between acceptance and

preference in the Mediterranean fruit fly

Benoît Facon[1], Madeline Chauve[1], Virginie Ravigné[2], Bruno Serrate[1], Myriam

Robejean[1], Antoine Fraimout[1], Julien Foucaud[1]

1 CBGP, INRAE, IRD, CIRAD, Institut Agro, University of Montpellier, 755 avenue du

Campus Agropolis, CS 34988, Montferrier sur Lez cedex 30016, France

2 CIRAD, UMR PHIM, – PHIM, University of Montpellier, CIRAD, INRAE, Institut

Agro, IRD, TA A-120/K, Campus international de Baillarguet, avenue du Campus

d'Agropolis, Montpellier Cedex 5 34398, France

 * To whom correspondence should be addressed. E-mail: benoit.facon@inrae.fr

Summary

Oviposition behavior in phytophagous insects is influenced by different stimuli and plays a

key role in pest dynamics and crop loss. This study used 3D-printed artificial fruits varying in

colour (yellow, blue, white) and odour (cherry, orange, banana) to test how visual and

olfactory cues affect oviposition acceptance (no-choice) and preference (choice). In no-choice

assays, the nine artificial fruits displayed sufficiently different visual and olfactory cues to

trigger different egg-laying outputs (by a factor 1:3 between the least attractive fruit, white

fruit with banana scent and the most attractive fruit, yellow fruit with cherry scent). While

cues acted independently in no-choice settings, significant interactions were observed in

choice conditions, highlighting multimodal sensory integration. In choice assays, the number

of eggs laid and female preference depended on both the characteristics of fruits and their

context. However a strong correlation was found between acceptance and preference. The

relationship found between acceptance and preference implied that when a fruit seemed

preferred in no-choice assays, it was even more preferred in one-choice assays. We finally

discussed the practical implications for behavior-based pest management strategies.

Introduction

Phytophagous pest insects are of particular importance for food safety as they are responsible

for between 20 and 40% of crop production losses worldwide (Culliney 2014, Deutsch et al.

2018). For a large number of these species, larvae are unable to move from one plant to

another to find the best conditions for their successful growth and development. Therefore,

the choice of the host plant in which larvae eventually develop relies on gravid females and

their oviposition behavior. Consequently, phenotypic variation in females' oviposition

behavior is tightly linked to offspring growth and survival, and constitutes an important

fitness component on which selection can act (Jaenike 1978, Cunningham 2012). Aside from

its relevance in the field of evolutionary ecology (see for instance Futuyma & Moreno 1988,

Bernays & Chapmann 1994), the study of insect oviposition behavior is particularly important

for the management of pest species. In an effort to reduce the use of synthetic chemical

pesticides, studying oviposition behavior is needed for the development of behavior-based

pest control techniques such as masking of host plant compounds to disrupt recognition,

selecting crop varieties emitting less attractive volatiles, using trap crops or combining

repellent and attractant elements in push–pull systems (Anton & Cortesero 2022). Integrating

evolutionary theory to studying oviposition behavior seems therefore to be a promising

approach to determine the efficiency of alternative pest control techniques, and the

evolutionary response of insect populations to this new form of selection (Sentis et al. 2022).

Oviposition behavior involves a complex suite of traits. Insects locate and discriminate among

plants by a variety of sensory means that ultimately lead to plant acceptance or rejection

(Jaenike 1978). Even though olfaction seems sufficient to locate and identify a suitable host

for some phytophagous species, other senses, i.e., vision and contact chemoreception may

also play roles in this process (Tasin et al. 2011, Bernays & Chapman 1994, Balkenius et al.

2009), with cross-stimuli interactions being the rule rather than the exception. The whole fruit

display, involving scent, colour, pattern, shape, touch, and taste, is perceived by sensory

systems and integrated in higher brain centers to trigger the correct behavioral response

(Riffell 2020, Saitta et al. 2023).

Recent studies have highlighted non-trivial interactions between stimuli (mainly visual and

olfactory). For instance, some insects such as the melon ladybird, *Chnootriba elaterii*, cannot

discriminate plants based solely on visual or olfactory signals (Saitta et al. 2023). They need

to perceive both cues simultaneously to select their host plant. When sensory cues come into

conflict, i.e., visual stimuli do not match olfactory stimuli, then insect response can be altered

in a highly variable manner depending on the odours and colours involved (Yv et al. 2015,

Riffell 2020). Just as the suite of traits involved in oviposition behavior, the stimuli produced

by the plants are fundamentally multivariate, with each stimulus potentially varying in its

mode of action. Accounting for these multimodal interactions is therefore important to

understand oviposition choice of phytophagous insects (Balkenius & Balkenius 2016).

Finally, in other insects, the olfactory input modulates the visual system, whereas the visual

system does not modulate the olfactory responses (Riffell 2020).

The response to stimuli is also highly context-dependent, depending in particular on the

current availability of different host species (Schäpers et al. 2017). For instance, *Drosophila*

melanogaster females prefer to oviposit on ethanol-containing substrates, but only as long as

they are very close to an ethanol-free substrate (Sumethasorn & Turner 2016). The

importance of the environmental context has led to the definition of two quantitative traits:

oviposition acceptance and oviposition preference (Fanara et al. 2023). Acceptance is defined

as the propensity of the female to lay eggs on a particular host when it is the only available

possibility, as typically measured by no-choice experiments. In the field, this would

correspond to monocultures where all plants produce the same stimuli. Preference is the

propensity of the female to lay eggs on a host when other hosts are available. Preference is

best assessed through one- or multiple-choice assays. In the field, such situations would be

found in mixed cultures or in natural plots harboring a diversity of host plants. Resolving the

relationship between oviposition acceptance and preference is particularly important for the

development of evolutionary-informed pest control strategies. Several studies have tested this

relationship using different insect species with contrasting results (Yang et al. 2015, Charlery

de la Masselière et al. 2017, Olazcuaga et al. 2019, Fanna et al. 2023). Charlery de la

Masselière et al. (2017) found that acceptance and preference showed a significant positive

correlation for four Tephritidae species out of six studied species. On the contrary, Fanara et

al. (2023) found that preference is not correlated to acceptance in *Drosophila melanogaster*,

suggesting that these behaviors are independent.

To our knowledge, no study to date has simultaneously quantified acceptance and preference

while experimentally disentangling the relative contributions of both visual and olfactory

stimuli. By integrating these four components within a unified framework, our study provides

a comprehensive assessment of how multimodal cues shape oviposition decisions and how

acceptance and preference interact to determine final host-use outcomes using the

Mediterranean fruit fly *Ceratitis capitata* as a study model. Belonging to the Tephritidae

family (true fruit flies), *C. capitata* is a multivoltine generalist species with more than 300

described host plants (Dias et al. 2018). Originating from tropical sub-Saharan Africa, it now

has a global distribution and is considered as one of the most important insect pests for fresh

fruit and vegetables production worldwide because of both direct damage to fruits and

quarantine regulations (USDA 2006). Females lay their eggs in ripening or ripe fruits, leading

to accelerated maturation and premature fruit drop, rendering them unfit for consumption and

unsuitable for marketing (Dias et al. 2018). To study oviposition choice in *C. capitata*, we

produced 3D-printed oviposition supports to mimic fruits that can vary in colour and odour,

two essential cues for host choice in *C. capitata*. In this study, we measured the behavior of

female flies in response to three colours (blue, yellow, and white) and three odours (cherry,

banana, and orange). We used a full-factorial design to thoroughly determine the impact of

visual and olfactory cues on the oviposition behavior of *C. capitata* in two types of

environments: no-choice and one-choice assays, allowing to respectively evaluate oviposition

acceptance and preference. By delving into the intricacies of the oviposition behavior, we aim

at providing valuable insight into the determinants of a major trait of insect pest biology and

hope to contribute to guide efforts in agroecological crop protection.

Materials and Methods

Study system and experimental conditions

The individuals used in this study were derived from a population reared in the arthropod

rearing and phenotyping facilities of the Biology Centre for Population Management

(Montpellier, France) for two years following crosses between individuals collected in the

wild between 2021 and 2022 from three different southern France localities: San-Giuliano

(Corsica), Torreilles (Pyrénées-Orientales), and Millas (Pyrénées-Orientales). Both rearing

and experiments were conducted in climate chambers under controlled conditions

(temperature: $25 \pm 1°C$, humidity: $60 \pm 10\%$, and photoperiod: 12L:12D). For the mass

rearing of this population, white mesh cages of $30 \times 30 \times 30$ cm containing approximately

1,800 individuals with a sex ratio of 1:1 were used. Larvae were reared on an artificial diet

composed of dehydrated carrot powder, brewer's yeast, dehydrated potato, water, and

Nipagin/Sodium benzoate. In laboratory conditions, the developmental cycle from egg to

adult emergence lasted approximately 17 days. These cages allowed the creation of cohorts of

gravid females every week for the experiments. After adult emergence, females had the

opportunity to mate for approximately five days before being isolated in experimental cages.

The experiments were conducted in smaller white mesh cages of $15 \times 15 \times 15$ cm (BugDorm-

4M1515 insect cage that has nylon netting (44 x 32 mesh) with 700 µm aperture). Each cage,

representing a replicate, contained eight reproductive females of the same age (five days).

Artificial fruits were chosen over real fruits for three main reasons. Firstly, fruit availability

varies seasonally. Additionally, the inter-fruit variability within a species may be high. Lastly,

previous studies have shown that visual cues such as fruit shape and size can strongly

influence oviposition behavior in fruit flies (Prokopy et al. 1984, Prokopy & Roitberg 1984).

Accordingly, we used 3D-printed fruits to control for shape and size, allowing us to

specifically test the effects of independent variation of colour and odour.. These artificial

fruits consisted of a black disk of 4 cm in diameter and 1.3 cm in height to hold an odourous

liquid, topped by a coloured hemisphere with 32 small perforations for the female to insert her

ovipositor. The hemisphere measured 4 cm in diameter and 2 cm in height (Appendix 1).

Odourous liquids consisted in 2 mL of 1% water-diluted aroma prepared on the day of the

behavioral assays (Appendix 1). Odour release rates were not quantified in the present study.

As such, the results should be interpreted in a comparative framework and not directly

extrapolated to field conditions. We selected three visual stimuli (blue, yellow, and white) and

three natural fruit aromas (cherry, banana, and orange - Culinaide, France) based on literature

and preliminary experiments. Although there are some contradictory data (Diaz-Fleischer et

al. 1999), we would expect yellow to be the most attractive visual stimuli, white the least

attractive and blue intermediate (Katsoyannos et al. 1986). Although "colour" has a specific

definition in the context of insect vision (van der Kooi et al. 2021), this term will be used

throughout the rest of the manuscript for the sake of simplicity. No study has yet been

conducted on natural fruit aromas, but in the field orange is a good, i.e., heavily infested, host.

Banana is a bad (rarely infested) host, and cherry is an intermediate host

(https://edis.ifas.ufl.edu/publication/IN371,

https://www.agric.wa.gov.au/medfly/mediterranean-fruit-fly-host-plants).

Oviposition behavior assays

The nine colour-odour combinations (Appendix 2), hereafter fruit types, were all tested in

four different choice types: no-choice assays and three different one-choice assays. In no-

choice assays (homogeneous environment), two identical artificial fruits were placed in each

cage. In one-choice assays (heterogeneous environment), the two fruits could differ only in

colour (visual choice), only in odour (olfactory choice), or in both colour and odour (visual

and olfactory choice). It has to be noted that the two artificial fruits were placed 3 cm apart

within the experimental cage. Before being reused, artificial fruits were thoroughly washed

with detergent and tap water to remove any residue from *C. capitata* pheromones or fruit

aroma.

To estimate oviposition preferences while minimizing both initial behavioural noise and

experience-dependent effects, females were tested for the same type of artificial fruits in two

standardized 4-hour sessions separated by a two-day interval. The number of eggs laid were

summed for each female group over the two testing sessions. Assessing innate oviposition

preferences involves a trade-off between short assays, which may be dominated by

exploratory or stochastic behaviour, and longer or repeated assays, which may induce learning

or physiological constraints such as egg-load depletion. The present design reflects a

compromise between these opposing sources of bias. To avoid potential bias related to the

placement of artificial fruits in the cages and the positioning of cages within the climate

chamber, each fruit type was randomly assigned across all cages, and the position of the fruits

within the cage (left or right) was also randomly determined. For the no-choice assays, we

conducted 14 replicates for each of the nine fruit types. For the one-choice assays, we

performed seven replicates per fruit pair in the visual choice treatment, seven replicates in the

odour choice treatment, and eight replicates in the combined visual and odour choice

treatment. Because all assays could not be conducted simultaneously, experiments were

organized into temporal blocks corresponding to the weeks during which measurements were

performed.

The day before the first measurement, females were isolated in experimental cages using a

mouth aspirator. In each cage, eight females were placed with water and food (white sugar

and inactive dry yeast). On the day of the first measurement, water and food were removed

from the cages. The artificial fruits were prepared according to the tested colours and odours

and placed in the cages for four hours (from 10 a.m. to 2 p.m). After four hours of oviposition,

artificial fruits were removed from the cages and replaced with water and food sources. Then

the bases of the artificial fruits containing the laid eggs were photographed and counted

automatically with an ad-hoc artificial intelligence model. This model made use of the

detectron2 platform (https://github.com/facebookresearch/detectron2) and was trained from a

Mask-RCNN ResNet 101 model, using approximately 400 annotated 1000x1000px vignettes

(80% used for training and 20% for validation). Annotation consisted of individually masking

each egg in the vignette using CVAT v1.1.0. We verified this model on an independent

dataset of 49 human-counted photographs, and found to attain satisfying metrics (Kendall's τ

$= 0.933$; Appendix 3). Two days after the first measurement, the second measurement was

conducted following the same protocol.

Statistical analysis

Statistical analyses were performed using R v.4.3.2 (R core team 2024). All data and R code

are available at: https://doi.org/10.57745/8VMCMQ.

Two preliminary checks were first addressed. Firstly, the mortality rate was estimated to

check whether experimental manipulations disturbed the individuals. Across all assays, the

average female mortality rate between the two measurements of the same block was 1.66%,

which is very low, indicating that the females were not particularly stressed during the

experiments. Despite low mortality, the number of living females was still accounted as an

offset covariate in subsequent statistical analyses. Secondly, a potential left-right bias was

also tested for in the no-choice assays. In no-choice assays, there was no significant difference

in the numbers of eggs laid in the left and right artificial fruits (paired t-test, $t = 0.63$, df $=$

125, $p = 0.53$), indicating that only colour and odour can affect the number of eggs laid in

each of the two fruits per cage.

Then, three response variables were studied: the total number of eggs of the pair of artificial

fruits per cage, the number of eggs laid in the focal fruit, and the preference estimated as the

number of eggs laid in the focal fruit divided by the total number of eggs of the pair of

artificial fruits per cage. Effects of the tested odours and colours were estimated using

Generalized Linear Mixed Models (GLMM) with the R package glmmTMB (Brooks et al.,

2017), in which blocks were treated as a random effect and the number of females as an offset

covariate. The selection of the most appropriate error distribution and link function was based

on residual checks provided by the R package DHARMa (Hartig 2024).

To understand the effect of colours and odours on the total number of eggs laid, we first

performed a GLMM (family: gaussian) with the fruit colour, odour and the colour-odour

interaction in no-choice assays. We then tested for the effect of colours and odours of both

artificial fruits on the total number of eggs on the global dataset (no-choice and one-choice

assays). Not having any strong *a priori* on the relevant interactions to account for, we applied

a systematic model selection procedure on the most complete model (i.e. colour and odour of

the focal fruit, colour and odour of the complementary fruit, and all potential interactions)

using the package MuMIn (Bartoń, 2024).

Next, to assess the context-dependence of oviposition behavior, the number of eggs laid on

the focal fruit was modelled using GLMM (family: quasi-poisson) as resulting from the focal

fruit, the choice type and the fruit x choice type interaction. The focal fruit has nine types (3

colours $\times$ 3 odours) and the variable "choice type" has four categories (a- no-choice assay, b-

visual heterogeneity, c- olfactory heterogeneity, d- visual and olfactory heterogeneity).

Based on the data obtained from one-choice assays, observed preferences were calculated for

each pair of artificial fruits using the formula:

$$OP = \frac{NF}{NF + NC}$$

with NF = number of eggs from the focal fruit, and NC = number of eggs from the

complementary fruit.

For each pair of fruits presented in one-choice assays, expected preferences, i.e., preferences

that should be observed in absence of interaction between fruits, were estimated from the data

obtained from no-choice assays. The expected preferences (EP) were calculated using the

following formula:

$$EP = \frac{MF}{MF + MC}$$

with MF = mean number of eggs from the focal fruit (estimated in no-choice), and MC =

mean number of eggs from the complementary fruit (estimated in no-choice).

Preference was modelled as resulting from the effect of the focal fruit, the choice type and

their interaction using GLMM (family: ordbeta). We finally analyzed the correlation between

expected and observed preferences using a GLMM with observed preference as the response

variable, expected preference, choice type and the interaction as explanatory variables using

all the heterogeneous assays.

Results

Total number of eggs

In no-choice experiments, visual stimuli ($\chi^2 = 12.30$, df $= 2$, $p = 0.002$) and olfactory stimuli

($\chi^2 = 16.98$, df $= 2$, $p = 0.0002$) significantly influenced the total number of eggs laid. The

interaction between visual and olfactory stimuli was not significant ($\chi^2 = 2.93$, df $= 4$, $p = $

0.57, Appendix 4). It was thus possible to rank the artificial fruits based on the total number

of eggs (Figure 1). Overall, yellow was the colour that most stimulated egg-laying, while

white was the least stimulating and blue was intermediate. Cherry and orange odours were not

significantly different, and both stimulated egg-laying more than the banana odour (Appendix

5).

Figure 1: Total number of eggs (with their 95% CI) laid by the eight females during the four-hour assays. The

nine colour-odour combinations are ranked in an increasing order. The significance underlying the compact

letter display is Bonferroni adjusted.

In the global dataset (no-choice and one-choice assays), the most parsimonious model

explaining the total number of eggs laid included a significant effect of the colour and odour

of the focal fruit ($\chi^2 = 82.09$, df = 2, $p < 2.2.10^{-16}$ and $\chi^2 = 98.81$, df = 2, $p < 2.2.10^{-16}$,

respectively), a significant effect of the colour and odour of the complementary fruit ($\chi^2 =$

40.82, df = 2, $p = 1.37.10^{-9}$ and $\chi^2 = 54.97$, df = 2, $p = 1.16.10^{-12}$, respectively), an interaction

between the colour and odour of the focal fruit ($\chi^2 = 20.47$, df = 4, $p = 0.0004$), an interaction

between the colour and odour of the complementary fruit ($\chi^2 = 14.68$, df = 4, $p = 0.005$), an

interaction between the odours of the two fruits ($\chi^2 = 80.08$, df = 4, $p < 2.2.10^{-16}$), and an

interaction between the colour of the focal fruit and the odour of the complementary fruit ($\chi^2 =$

9.57, df = 4, $p = 0.05$, Appendix 6).

Figure 2: Number of eggs (with their 95% CI) laid by the eight females in the focal during the four-hour assays in
the no-choice or acceptance assays (A), in the colour-choice assays (B), in the odour-choice assays (C) and in
odour and colour choice assays (D).

Number of eggs laid in the focal fruit and preference

The number of eggs laid in the focal fruit significantly depended on fruit characteristics ($\chi^2 =$ 21.47, df = 8, $p = 0.006$), on the choice type ($\chi^2 = 14.76$, df = 3, $p = 0.002$) and their interaction ($\chi^2 = 115.15$, df = 24, $p = 7.02.10^{-14}$, Figure 2, Appendices 7 and 8).

The preference was strongly affected by the choice type ($\chi^2 = 56.91$, df = 3, $p = 2.7.10^{-12}$), the interaction between choice type and focal fruit ($\chi^2 = 288.30$, df = 24, $p < 2.2.10^{-16}$) but not by the focal fruit itself ($\chi^2 = 3.54$, df = 8, $p = 0.90$, Figure 3, Appendix 7).

Figure 3: Preference for the focal fruit (with their 95% CI) during the four-hour assays in the no-choice or acceptance assays (A), in the colour-choice assays (B), in the odour-choice assays (C) and in odour and colour choice assays (D).

Observed preference was strongly correlated with the expected preference ($\chi^2 = 18.49$, df = 1,

$p = 1.7.10^{-5}$), but not by the choice type ($\chi^2 = 0.11$, df = 2, $p = 0.95$) or the interaction between

expected preference and choice type ($\chi^2 = 0.13$, df = 2, $p = 0.94$, Appendix 9). The observed

preferences can be largely explained by the expected preferences, as indicated by the linear

regression: with a high

coefficient of determination ($R^2 = 0.647$). This relationship implies that when a fruit is

preferred in no-choice assays, it is even more preferred in one-choice assays.

Figure 4: Mean observed preferences versus expected preferences for the three types of choice type. The solid
line represents the linear regression fit between observed and expected data and the dotted line represents
y=x.

Discussion
304
Our study adds to the list of works supporting that females of phytophagous insects can alter

their oviposition behavior depending on the current availability of different oviposition sites

(Jaenike 1978, Schäppers et al. 2017). Here we demonstrate that, in the Mediterranean fruit

fly, this alteration stems the high context-dependence of the action of visual and olfactory

stimuli on oviposition behavior (Light & Jang, 1996). In no-choice assays, all the nine

artificial fruits (three colours × three odours) exceeded the threshold required to stimulate

egg-laying, but they displayed sufficiently different visual and olfactory cues to trigger

significant different egg-laying outputs (by a factor 1:3 between the two extreme fruits). Our

results are in agreement with recent research demonstrating that generalist phytophagous

insects have the capacity to rank hosts when given a choice, in contradiction to the "neural

limitations" hypothesis postulating that generalist species should not have the neural capacity

to identify cues specific to every possible host (Silva & Clarke 2020). In addition to the

classification of different colours and odours, it is important to note that the two types of

stimuli acted independently of each other in no-choice assays. On the contrary, in the full

dataset, including one-choice assays, different types of interactions between these stimuli

(visual and olfactory) needed to be accounted for to properly model the data in addition to the

simple effect of each stimulus. First, the two types of stimulus (visual and olfactory)

significantly interacted within each of the two artificial fruits to explain the number of eggs

laid by the female flies. The interaction between the odours of the two fruits and between the

colour of the focal fruit and the odour of the complementary fruit also had to be accounted for

to best explain the total number of eggs.

A complementary way of analyzing our data is to look specifically at the number of eggs laid

in the focal fruit on the overall dataset. As with no-choice assays, the number of eggs laid in

the focal fruit in one-choice assays depends on its characteristics in terms of colour and odour,

but this time in interaction with the characteristics of the complementary fruit. In other words,

the number of eggs laid in a given fruit will be different if the fruit next to it is different from

it in odour, colour or both. As expected, we found the same pattern when studying

preferences: the effect of stimuli from the focal fruit depends of the female's evaluation of

both stimuli (visual and olfactory) from the complementary fruit. We found that visual and

olfactory stimuli do not act independently but interact significantly, influencing oviposition

site selection. This finding aligns with studies on sensory integration in insects, particularly

flower-visiting species like moths suggesting that while each stimulus (visual or olfactory)

can have a direct effect on oviposition behavior, both stimuli are processed together in the

brain, contributing to a unified perception of the external environment (Balkenius et al.,

2009). This multimodal integration of visual and olfactory stimuli enhances behavioral

precision and reflects the adaptive benefits of integrating sensory inputs in a dynamic

environment (Hebets & Papaj 2005). The underlying neural mechanisms of this multimodal

integration are still not fully understood, but recent studies have shown that responses to

combined visual and olfactory stimuli are modulated in the mushroom bodies and at antennal

lobe levels of insects (Rennou & Anton 2020). However a study by Vinauger et al. (2019)

showed that the modulation of peripheral visual and olfactory sensory stimuli is asymmetric

in the mosquito *Aedes aegypti*: CO_2 modulates lobula neuropil responses to discrete visual

stimuli but visual stimuli do not modulate responses to CO_2 in olfactory glomeruli. Our

experimental approach allowed us to preserve the integrity of the different sensory channels,

which better reflects the natural conditions which insects are confronted to, with multiple

simultaneous sensory stimuli. This approach provides a complementary understanding to

previous studies that have often isolated sensory stimuli by removing certain modalities (e.g.,

by ablation of antennae or eye painting; see Riffell, 2020).

Although the female's response to a given fruit is modulated not only by the fruit individual

characteristics (colour and odour) but also by the characteristics of the other available fruit,

our study shows that the observed preferences in one-choice are strongly correlated with

expected preferences calculated with data from no-choice assays. This result challenges the

study by Fanara et al. (2023) that showed a clear decoupling between observed preferences

and expected preferences in *Drosophila melanogaster*. Among other existing studies, either

the number of fruit pairs tested is too limited to allow statistically robust analyses, or

preference is assessed using multi-choice (cafeteria-type) assays. Both approaches limit

meaningful quantitative comparisons between acceptance and preference. (Karageorgi et al.

2017, Olazcuaga et al. 2019, Yang et al. 2008, 2015). It would be of great interest to replicate

this experimental protocol on a variety of species to better understand the determinants

(phylogenetic, ecological, etc.) of the observed patterns. It is also important to note that the

relationship between observed preferences and expected preferences is not influenced by the

type of heterogeneity between the two fruits (visual, olfactory or both). It is not the nature of

the stimulus (visual vs olfactory) that matters, but the relative motivation to lay that the

stimulus provides for females after multimodal integration (Renou & Anton 2020). In *C.*

capitata, our results show that if the outcome of bimodal interaction between olfactory and

visual cues of a fruit is higher than another fruit in no-choice assays, then it will be preferred

in one-choice assays by a bias that can be estimated by the coefficients of the linear regression

between observed preferences and expected ones. This result suggests that, to be used as a

trap plant or in a push-pull, a host plant must elicit more egg-laying than the crop to be

protected in no-choice assays. We hope that our results will stimulate new studies that will

seek to better understand how the female weigh the various signals to make an appropriate

decision for oviposition. Identifying optimal cues in different ecological settings will be key

to developing sustainable strategies based on manipulation of insect behaviour.

Data accessibility

380   All data and R code are available online at:

381   https://doi.org/10.57745/8VMCMQ

Supplementary material

384   Supplementary materials is available online at:

385   https://www.biorxiv.org/content/10.1101/2025.06.20.660717v1

Acknowledgements

388   JF and BF acknowledge financial support by the INRAE scientific department SPE and the

389   UMR CBGP from its internal grant support.

Conflict of interest disclosure

392   The authors of this preprint declare that they have no financial conflict of interest with the

393   content of this article. JF and BF are PCI Zool recommenders.

References

Anton S. & Cortesero A.-M. (2022) Plasticity in chemical host plant recognition in
herbivorous insects and its implication for pest control. Biology, 11, 1842,
doi:10.3390/biology11121842

Balkenius A., Bisch-Knaden S. & Hansson B. (2009) Interaction of visual and odour cues in
the mushroom body of the hawkmoth *Manduca sexta*. The Journal of Experimental Biology,
212, 535-541, doi:10.1242/jeb.021220

Balkenius A. & Balkenius C. (2016) Multimodal interaction in the insect brain. BMC
Neuroscience, 17, 29, doi:10.1186/s12868-016-0258-7

Bartoń K (2024) MuMIn: Multi-Model Inference. R package version 1.48.4, https://CRAN.R-
project.org/package=MuMIn

Bernays E.A. & Chapman R.F. (1994) Host-Plant Selection by Phytophagous Insects.
Chapman & Hall, New York, 95-165. doi:10.1007/b102508

Brooks M.E., Kristensen K., van Benthem K.J., Magnusson A., Berg C.W., Nielsen A., Skaug
H.J., Maechler M. & Bolker B.M. (2017) glmmTMB Balances Speed and Flexibility Among
Packages for Zero-inflated Generalized Linear Mixed Modeling. The R Journal, 9, 378–400,
doi:10.32614/RJ-2017-066

Charlery de la Masselière M., Facon B., Hafsi A. & Duyck P.-F. (2017) Diet breadth
modulates preference - performance relationships in a phytophagous insect community.
Scientific Reports, 7, 1-9, doi:10.1038/s41598-017-17231-2

Cook S. M., Khan Z. R. & Pickett J. A. (2007) The Use of Push-Pull Strategies in Integrated
Pest Management. Annual Review of Entomology, 52, 375□400,
doi:10.1146/annurev.ento.52.110405.091407

Culliney T. W. (2014) Crop losses to arthropods. In Integrated Pest Management, eds D.
Pimentel and R. Peshin (Cham: Springer), 201–225. doi: 10.1007/978-94-007-7796-5_8

Cunningham J. P. (2012) Can mechanism help explain insect host choice? Journal of
Evolutionary Biology, 25, 244□251, doi:10.1111/j.1420-9101.2011.02435.x

Deutsch C.A., Tewksbury J.J., Tigchelaar M., Battisti D.S., Merrill S.C., Huey R.B. & Naylor

R.L. (2018) Increase in crop losses to insect pests in a warming climate. Science, 361, 916–

919, doi: 10.1126/science.aat3466

Dias N.P., Zotti M.J., Montoya P., Carvalho I.R. & Nava D.E. (2018) Fruit fly management

research: A systematic review of monitoring and control tactics in the world. Crop Protection,

112, 187-200, doi:10.1016/j.cropro.2018.05.019

Díaz-Fleischer F., Papaj D.R., Prokopy R.J., Norrbom A.L. & Aluja, M. (1999) Evolution of

fruit fly oviposition behavior. In Fruit Flies (Tephritidae), pp. 829–860. CRC Press

Fanara J.J., Beti M.I.L., Gandini L. & Hasson, E. (2023). Oviposition behaviour in

Drosophila melanogaster: Genetic and behavioural decoupling between oviposition

acceptance and preference for natural fruits. Journal of Evolutionary Biology, 36, 251263,

doi:10.1111/jeb.14109

Foster S. & Harris M. O. (1997) Behavioral manipulation methods for insect pest-

management. Annual Review of Entomology, 42, 123-146,

doi:10.1146/annurev.ento.42.1.123

Futuyma D.J. & Moreno G. (1988) The evolution of ecological specialization. Annual Review

of Ecology, Evolution, and Systematics, 19, 207-233,

doi:10.1146/annurev.es.19.110188.001231

Hartig F. (2024). DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)

Regression Models. R package version 0.4.7, https://CRAN.R-project.org/package=DHARMa

Hebets E.A. & Papaj D.R. (2005) Complex signal function: developing a framework of

testable hypotheses. Behavioral Ecology and Sociobiology, 57, 197-214, doi:10.1007/s00265-

004-0865-7

Jaenike J. (1978) On optimal oviposition behavior in phytophagous insects. Theoretical

Population Biology, 14, 350356. Doi:10.1016/0040-5809(78)90012-6

Karageorgi M., Bräcker L.B., Lebreton S., Minervino C., Cavey M., Siju K.P., Grunwald

Kadow I.C., Gompel N., Prud'homme B. (2017) Evolution of multiple sensory systems drives

novel egg-laying behavior in the fruit pest Drosophila suzukii. Current Biology, 27, 847-853,

doi:10.1016/j.cub.2017.01.055

Katsoyannos B. I., Panagiotidou K. & Kechagia, I. (1986) Effect of color properties on the
selection of oviposition site by *Ceratitis capitata*. Entomologia Experimentalis et Applicata,
42, 187□193. doi:10.1111/j.1570-7458.1986.tb01020.x

Light D.M. & Jang E.B. (1996) Plant volatiles evoke and modulate tephritid behavior. In Fruit
Fly Pests: A World Assessment of Their Biology and Management (eds. B.A.McPheron &
G.J. Steck), pp. 123–133. Delray Beach, FL: St. Lucie Press.

Olazcuaga L., Rode N.O., Foucaud J., Facon B., Ravigné V., Ausset A., Leménager N.,
Loiseau A., Gautier M., Estoup A. & Hufbauer R.A. (2019) Oviposition Preference and
Larval Performance of *Drosphila suzukii* (Diptera: Drosophilidae), Spotted-Wing Drosophila:
Effects of Fruit Identity and Composition. Environmental Entomology, 48, 867-881,
doi:10.1093/ee/nvz062

R Core Team (2024). R: A Language and Environment for Statistical Computing. R
Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org

Rennou M. & Anton S. (2020) Insect olfactory communication in a complex and changing
world. Current Opinion in Insect Science, 42, 1-7, doi:10.1016/j.cois.2020.04.004

Riffell J.A. (2020) The neuroecology of insect-plant interactions: The importance of
physiological state and sensory integration. Current Opinion in Insect Science, 42, 118-124,
doi:10.1016/j.cois.2020.10.007

Saitta V., Rebora M., Piersanti S. & Salerno G. (2023). Visual and chemical cues in the host
plant selection of the melon ladybird *Chnootriba elaterii* (Coleoptera□: Coccinellidae).
Arthorpod-Plant Interactions. doi:10.1007/s11829-023-10018-5

Schäpers A., Petrén H., Wheat C.W. Wiklund C. & Friberg M. (2017). Female fecundity
variation affects reproducibility of experiments on host plant preference and acceptance in a
phytophagous insect. Proceedings of the Royal Society B: Biological Sciences, 284,
20162643. doi:10.1098/rspb.2016.2643

Sentis A., Hemptinne J.-L., Magro A. & Outreman Y. (2022) Biological control needs
evolutionary perspectives of ecological interactions. Evolutionary Applications, 15, 1537-
1554, doi:10.1111/eva.13457

Silva R. & Clarke A.R. (2020) The "sequential cues hypothesis": a conceptual model to

explain host location and ranking by polyphagous herbivores. Insect Science, 27, 1136-1147,

doi:10.1111/1744-7917.12719

Sumethasorn M. & Turner T.L. (2016) Oviposition preferences for ethanol depend on spatial

arrangement and differ dramatically among closely related Drosophila species. Biology Open,

5, 1642-1647, doi:10.1242/bio.019380

Tasin M., Lucchi A., Ioriatti C., Mraihi M., De Cristofaro A., Boger Z. & Anfora G. (2011)

Oviposition Response of the Moth *Lobesia botrana* to Sensory Cues from a Host Plant.

Chemical Senses, 36, 633-639, doi:10.1093/chemse/bjr027

USDA (2006) California county agricultural commissioners' data 2005, p. 80. Sacramento,

CA: United States Department of Agriculture, National Agricultural Statistics Service.

Vinauger C., Van Breugel F., Locke L.T., Tobin K.K.S., Dickinson M.H., Fairhall A.L.,

Akbari O.S. & Riffell J.A. (2019) Visual-olfactory integration in the human disease vector

mosquito *Aedes aegypti*. Current Biology, 29, 2509-2516.e5, doi:10.1016/j.cub.2019.06.043

Yang C.H., Belawat P., Hafen E., Jan L.Y. & Jan Y.N. (2008) Drosophila egg-laying site

selection as a system to study simple decision-making processes. Science, 319, 1679-1683,

doi:10.1126/science.1151842

Yang C.H., He R. & Stern U. (2015). Behavioral and circuit basis of sucrose rejection by

Drosophila females in a simple decision-making task. Journal of Neuroscience, 35, 1396–

1410, doi:10.1523/JNEUROSCI.0992-14.2015

Yv F.L., Hai X., Wang Z., Yan A., Liu B. & Bi Y. (2015) Integration of Visual and Olfactory

Cues in Host Plant Identification by the Asian Longhorned Beetle, *Anoplophora glabripennis*

(Motschulsky) (Coleoptera: Cerambycidae). PloS ONE, 10, e0142752,

doi:10.1371/journal.pone.0142752

505