Thriving Across Depths: How Blue Light Shapes a Large PSI Supercomplex and Specic

Photosynthetic Traits in the seagrass *Posidonia oceanica*.

Quentin Charras Ferroussier[1,2], Charlie Mathiot[3], Dmitry A. Semchonok[4,5], Eduard Elias[6],

Ahmad Farhan Bhatti[6], Régine Lebrun[7], Dorian Guillemain[8], Marina I. Siponen[9], Roberta

Croce[6] and Colette Jungas[1✉].

[1]Aix Marseille Univ, CEA, CNRS, BIAM, Team LGBP, Saint Paul-Lez-Durance, France F-1311. [2]Science for
Life Laboratory, School of Engineering Sciences in Chemistry, Biotechnology and Health, KTH-Royal Institute
of Technology, KTH University, Stockholm, Sweden.
[3]Aix Marseille Univ, CEA, CNRS, BIAM, Team Photosynthesis and Environment, Saint Paul-Lez-Durance,
France F-1311.
[4]Interdisciplinary Research Center HALOmem, Charles Tanford Protein Center, Martin Luther University
Halle-Wittenberg, Kurt-Mothes-Straße 3a, Halle/Saale, Germany.
[5]Instituto de Tecnologia Química e Biológica, Universidade Nova de Lisboa, Av. da República, 2780-157
Oeiras, Portugal.
[6]Biophysics of Photosynthesis, Department of Physics and Astronomy, Faculty of Science, Vrije Universiteit
Amsterdam, the Netherlands.
[7]Aix Marseille Univ, CNRS, Plate-forme Protéomique, Marseille Protéomique (MaP), IMM FR 3479, 31
Chemin Joseph Aiguier, 13009 Marseille, France.
[8]CNRS, IRD, IRSTEA, OSU Institut Pythéas, Aix-Marseille University, Marseille, France.
[9]Aix Marseille Univ, CEA, CNRS, BIAM, Team Proteintec, Saint Paul-Lez-Durance, France F-13115.

Abstract

Photosynthetic organisms rely on finely tuned mechanisms to optimize photosynthesis under

different light conditions. While these processes are well-characterized in land plants, the

adaptive strategies of marine plants remain largely unexplored. The Mediterranean seagrass

Posidonia oceanica (Alismatales), a key ecosystem engineer thriving from the surface up to

40m depth and one of the largest long-term blue carbon sinks in coastal environments. Here,

we investigate how *P. oceanica* adjusts its photosynthetic apparatus in response to varying

light spectra encountered at different seawater depths. Contrary to land plants, *P. oceanica*

maintains a relatively high PSI/PSII ratio and a high content of the major light-harvesting

complex II (LHCII), regardless of depth. Notably, the antenna size of the photosystems

remains stable across depths, although we document significant depth-dependent

reorganization of the thylakoid membrane ultrastructure. Moreover, we identify a novel large

PSI-LHCII supercomplex (L-PSI-LHCII) in *P. oceanica*, characterized by additional Lhca

proteins, reduced red-shifted absorption, and increased chlorophyll *b* content. Ultrafast

spectroscopy reveals the distinct energy transfer dynamics within this complex. The presence

of a similar supercomplex in other marine plants, such as *Zostera marina*, suggests a
conserved adaptive strategy among seagrasses.

Introduction

Blue carbon refers to carbon sequestered by vegetated coastal ecosystems, including
salt marshes, **mangroves**, and **seagrass meadows** over millennia. Although seagrasses
occupy only 0.01% of the Earth's surface, they account for approximately 4% of global
carbon stocks (Duarte et al., 2013). Among these, *Posidonia oceanica*, a Mediterranean
endemic, forms extensive meadows from shallow waters down to 40 m depth (Pasqualini et
al., 1998), covering roughly 25% of the region's coastal seabeds (Telesca et al., 2015). Its
slow horizontal expansion (1–5 cm/year) enables the formation of persistent structures, with
some meadows estimated to be up to 80,000 years old (Arnaud-Haond et al., 2012).

P. oceanica is a key ecosystem engineer supporting forming ecological niches for a
multitude of taxa (Díaz-Almela and Duarte, 2008; Boudouresque et al., 2012) and is
recognized as a highly efficient carbon sink. The reported long-term sequestration of ~270 t
C/ha/year (Pergent-Martini et al., 2021), significantly surpassing the ~4 t C/ha/year
sequestration typical of tropical forests (Yadav et al., 2022). This sequestration capacity
corresponds to 11–42% of historical CO_2 emissions from Mediterranean countries since the
Industrial Revolution (Pergent et al., 2014), primarily through the matte, the dense, decay-
resistant layer of rhizomes, roots and sediment (Boudouresque et al., 2016).

Despite its ecological and biogeochemical significance, *P. oceanica* meadows have
experienced a decline of approximately 34% over the past five decades (Pergent et al., 2014)
with current annual losses exceeding 1.5%. Anthropogenic pressures including coastal
development, increased turbidity, mechanical damage from anchoring, and rising sea surface
temperatures are the primary drivers of decline (Marbà et al., 1996; Boudouresque et al., 2012).
While restoration initiatives, such as transplantation, have yielded variable outcomes
(Molenaar, 1992; BOUDOURESQUE et al., 2013), a more comprehensive understanding of
the physiological and biochemical mechanisms underlying *P. oceanica*'s adaptation and
resilience is urgently needed for conservation and management strategies under accelerating
global change.

In marine environments, both light intensity and quality rapidly decline with depth,
with red light disappearing entirely beyond ~10 m and overall intensity reduced 20 to 50

times at deeper meadows (see below). *P. oceanica* thrives across these varying conditions and understanding its adaptive mechanisms provides valuable insights into photosynthetic processes. Photosystems I (PSI) and photosystem II (PSII) and their antenna systems (LHC) drive linear electron flow to produce ATP and NADPH, used for downstream CO_2 fixation. Optimal electron transfer requires balanced excitation between PSI and PSII, maintained through acclimation strategies such as adjusting PSI/PSII ratios (Chow et al., 1990; Wientjes et al., 2017), LHCII content (Wagner et al., 2008; Kouřil et al., 2013; Albanese et al., 2016), antenna size, and state transitions (Goldschmidt-Clermont and Bassi, 2015; Grieco et al., 2015; Rantala et al., 2020). Underwater, the dominant blue-green wavelengths are primarily absorbed by chlorophyll *b*, abundant in LHCII linked mostly to PSII (Su et al., 2017), enhancing its absorption capacity compared to PSI-LHCI (Bos et al., 2023). Thus, PSI excitation may become limited with increasing depth.

We examined the depth-dependent photosynthetic adaptations of *P. oceanica*, focusing on LHCII levels, PSI/PSII stoichiometry, and antenna sizes at depths of 2, 15, and 26 m. we unexpectedly found stable, elevated LHCII content and PSI/PSII ratios across all depths compared to terrestrial plants. Using a new protocol developed to isolate thylakoids from polyphenol-rich marine plants (Charras et al., 2024), we identified a Chl *b*-enriched PSI supercomplex, consisting of canonical PSI-LHCI linked to phosphorylated LHCII trimers and an additional Lhca1-Lhca4 heterodimer, uniformly expressed at all tested depths. Notably, the far-red absorbing Chls typical of land plants (Morosinotto et al., 2003; Romero et al., 2009) were absent in PSI of *P. oceanica* and related Mediterranean marine species, highlighting a common blue-shifted adaptation in all tested seagrasses, and the expression of the L-PSI-LHCII in deep-growing marine plants.

Results

The light environment of the *P. oceanica* meadows selected for the study

The light spectrum at each depth tested was determined (**Suplemental Figure 1**). From the surface to a depth of 2 m, light intensity decreases by a factor of ~2.5 (from 2000 to 800 $\mu mol.m^{-2}.s^{-1}$) largely in the 600-750 nm region. However, at depths of 15 and 26 m, the irradiance is between 20 and 50 times lower than at the surface (120 and 40 $\mu mol.m^{-2}.s^{-1}$, respectively) and completely depleted of red light; only green and blue-green light remain, peaking around 450-500 nm.

Depth-dependent changes in chloroplast ultrastructure

Long-term acclimation of plants to varying light conditons occurs through the composition
and structural organization of the thylakoid membrane (Buschmann et al., 1978; Ferroni et
al., 2016; Schumann et al., 2017). The chloroplast acclimation of *P. oceanica* was examined
at the structural level in leaves collected at different depths (2, 15, and 26 m) using electron
microscopy (**Figure 1**). The micrographs and the grana size quantification (**Suplemental**
Figure 2) revealed significant alterations in membrane organization: At 15 m, the grana
thickness increased **(Figure 1B and 1E)** compared to 2 m (200 nm vs. 140 nm) (**Figure 1A**
and 1D), while the grana diameter remained unchanged (around 420 nm). Surprisingly, at 26
m (**Figure 1C and 1F**), grana thickness decreased (80 nm vs. 140 nm at 15 m), but their
diameter was larger (510 nm vs. 420 nm at 15 m). Similar observations have been made in
Arabidopsis thaliana exposed to light favoring PSII excitation (Wagner et al., 2008),
consistent with the predominance of blue-green light in deeper water.

115
116
Figure 1: Transmission electron microscopy (TEM) analysis of the *P. oceanica*
chloroplasts from three growing depths. Electron micrographs of negative-stained
chloroplasts cross section (up) and thylakoid membrane (down) from *P. oceanic*a leaves that
grew at (**A-B**) 2 m depth, (**C-D**) 15 m depth, and (**E-F**) 26 m depth. **cp**: chloroplast, **cw**: cell

wall, **mt**: mitochondria, **m**: margins, **pg**: plastoglobules, **sl**: stroma lamellae. The
corresponding scale bars are specified below each figure panel.
123
P. oceanica shows a low Chl a/b ratio at all depths
Changes in the Chl*a/b* ratio is a proxy of changes in the number of LHCII. Surprisingly, in *P.*
oceanica, we found that the Chl*a/b* ratio varied little at the depths tested (2 m: 2.14±0.004, 15
m: 2.04±0.007, 26 m: 2.06±0.02) (**Figure 2A**). This ratio is very low compared with that of
A. thaliana, which is around 3.4-3.7 in high light and 2.7-2.9 in low light (Kouřil et al., 2013;
Wientjes et al., 2013). Over a 4-year record of Chl *a/b* data, however, the values show some
seasonal variations; in winter, the values at 2am are slightly higher (2.13-2.21) than at 15 and
26 m (2-2.06), while in summer are even lower (Chl *a/b* around 1.7-1.8) and stable at all
depths (**Supplemental Figure 3A**). These data suggest that LHCII is present in large amounts
in this plant at all depths.
134
The PSI/PSII ratio shows slight variation throughout depth
Since the difference in light spectra at different depths can result in different excitation of the
two photosystems, we investigated if changes occur in PSI/PSII ratio with depths.
We first assessed the PSI and PSII stoichiometry via immunodetection of PSII (CP43, CP47)
and PSI (PsaG, Lhca2) subunits. The analysis of the blots (**Figure 2B and suplemental**
Figure 4A) could not highlight significant variations, irrespective of the growing season of
the collected sample (**Supplemental Figure 4B).** We then quantified the functional PSI/PSII
stoichiometry using flash-induced electrochromic shift (ECS) at 520 nm on fresh untreated
and DCMU infiltrated leaves (Bailleul et al., 2010). The stoichiometry variation was difficult
to assess due to the deviation between experiments and replicates (**Supplemental Figure 4D**
and 4E), yet, at all depths tested, *P. oceanica* exhibits a PSI/PSII stoichiometry around 1.15 ±
0.06 (**Figure 2C**), higher than most land plants such as *A. thaliana* (0.6 to 0.9 grown under
white light and sunlight, respectively (Hogewoning et al., 2012; Wientjes et al., 2017).

Figure 2: Chlorophyll content and photosystems stoichiometry in *P. oceanica* thylakoids at different growing depths. **(A)** Chl *a/b* of the thylakoid membranes of *P. oceanica* collected at 2, 15, and 26 m depth in winter. Data for each depth are expressed as the mean ± SD of 2 technical replicates from a thylakoid preparation obtained with 30 shoots (4 leaves/shoot). **(B)** Immunoblot of thylakoid membranes from *P. oceanica* collected at 2, 15, and 26 m depth. 0,5 μg of Chl was loaded on the gel. The primary antibodies used are indicated. Quantification of the band's intensity is reported in Suplemental Figure 4a. **(C)** PSI/PSII stoichiometry ± SD in fresh young winter leaves estimated by difference of ECS signal between untreated (PSII+PSI) and DCMU-HA infiltrated (PSI only) leaves. Data are expressed as the mean from two independent experiments (**Suplemental Figure 4D and 4E**) at the 3 tested depths (n=3 different leaves from each depth and for each experiment).

Analysis of PSI and PSII supercomplexes and characterization of a Large PSI-LHCII supercomplex.

Insights into the composition of PSI and PSII supercomplexes were obtained through the BN-PAGE analysis of *P. oceanica*'s α-DDM-solubilized thylakoids from different depths (**Figure 3A).** A distinct and intense band in between the PSII-LHCII bands and the PSI-LHCI band was observed. 2D urea-PAGE analysis shows that this band contains the main protein components of PSI-LHCI, with in addition LHCII subunits (Lhcb1 and Lhcb2) like the PSI-LHCII state transition complex in *A. thaliana* (Wu et al., 2023) and *Z. mays* (Pan et al., 2018). This Larger PSI-LHCII was designated as L-PSI-LHCII and its composition was analyzed by mass spectrometry.

The LC-MS-MS data showed that L-PSI-LHCII has a higher relative content of Lhca1 (around 1.8) and Lhca4 (around 1.5) peptides in comparison with PSI-LHCI (**Figure 3C),** indicating the presence of an additional Lhca1-Lhca4 dimer (Lhca1²-Lhca4²) bound to PSI-LHCII.

As for the composition of the LHCII trimer, we found a 2:1 stoichiometry of Lhcb1 and Lhcb2 (**Supplemental Figure 5 A-C**), similar to *A. thaliana* and maize, as well as threonine

phosphorylation of the Lhcb2 subunit. L-PSI-LHCII and PSI-LHCI bands exhibit consistent intensity across all depths, regardless of the growing season (**Figure 3A**, **Suplemental Figure 6 and 7**). Quantification of the Native-PAGE bands reveals a L-PSI-LHCII/PSI-LHCI stoichiometry of 0.4, so the large complex represents at least 30% of the total PSI pool at all depths. L-PSI-LHCII seemed stable as a very weak amount of phosphorylated Lhcb2 protein was found in the free LHCII bands in the Native-PAGE (**Suplemental Figure 5D**). Moreover, phosphorylation levels of Lhcb2 in thylakoids show no significant variations between depths, consistent with the constant accumulation of the L-PSI-LHCII across depths when samples are collected during the day, regardless of the season of collection (**Suplemental Figure 5E**). In contrast, plants harvested before sunrise have a lower concentration of this complex (**Suplemental Figure 8**).

Figure 3: Analysis of thylakoid composition and the Large PSI-LHCII supercomplex.
(A) BN-PAGE of solubilized thylakoid collected at different growing depths in winter. For
each tested depth, 5μg of was loaded onto the gel. **(B)** Identification of the peptide content of
the BN-PAGE bands by a second dimension on Urea-PAGE. Gel stained using SyproRuby
(C) Proteomic analysis (LC-MSMS) of L-PSI-LHCII. Results were compared to PSI–LHCI.
A number of peptide spectra matches (PSMs) are represented as fold change of the unique
peptides from the 4 LHCAs normalized by the PSM of some of the PSI-core unique-peptides
(indicated on the x-axis) between the L-PSI-LHCII and the PSI-LHCI. **(D)** Absorption

spectrum of LHCII trimers purified from *P. oceanica* and *A. thaliana*. **(E)** Chl content of *P.*
oceanica LHCII **(F)** Normalized (680 nm peak) absorption spectrum of PSI-LHCII compared
to PSI-LHCI from *P. oceanica* and *A. thaliana*. The absorption difference spectrum between
PSI from *P. oceanica* and *A. thaliana* was multiplied by 50 for better visualization. **(G)** 77K
fluorescence emission spectra of PSI-LHCI and L-PSI-LHCII compared with the PSI-LHCI
of *A. thaliana*, normalized at the 720 nm band maxima.
205

Chlorophyll *b* enrichment in the LHCII, PSI-LHCI and L-PSI-LHCII complexes

Next we isolated the main pigment-protein complexes using sucrose density gradient
ultracentrifugation. The absorption spectrum of *P. oceanica* trimeric LHCII showed higher
amplitudes in the 450-500 nm and 650 nm regions compared to that of *A. thaliana* (**Figure**
3D), indicating a higher Chl *b* content. This was confirmed by pigment analysis, which
showed a Chl *a/b* ratio of 0.95 $\pm$0.01, (**Figure 3E**). In the model organisms
Chlamydomonas reinhardtii, *Physcomitrella patens, A. thaliana*, and *Zea mays*, the LHCII
trimer contains 42 Chls with a Chl *a/b* ratio of 1.33 and therefore binds 24 Chl *a* and 18 Chl *b*
(Huang et al., 2021; Zhang et al., 2023; Wu et al., 2023; Pan et al., 2018). Assuming the same
total number of pigments, *P. oceanica* LHCII is expected to bind 20.5 Chl *a* and 21.5 Chl *b*
(**Figure 3E**), suggesting that 3 Chl *a* have been replaced by Chl *b*.
The comparison of *P. oceanica* PSI-LHCI and the L-PSI-LHCII absorption spectra with that
of *A. thaliana* PSI-LHCI (**Figure 3F**) shows a Chl *b* enrichment in the seagrass. This was
confirmed by measuring the Chl *a/b* ratio of 6.84 $\pm$ 0.05, and 4.33 $\pm$0.03 in the PSI-LHCI
and L-PSI-LHCII respectively, which are clearly lower than those of *A. thaliana*
(**Suplemental Figure 9**).
In addition, the absorption in the far red of the PSIs from *P. oceanica* is less intense than in *A.*
thaliana (**Figure 3F and Suplemental Figure 10**), indicating that these complexes contain
less low energy pigments. This is confirmed by the 77K emission fluorescence spectra that
for both PSI-LHCI and L-PSI-LHCII show a peak at 718-719 nm, significantly blue-shifted
compared to PSI-LHCI of *A. thaliana* (735 nm) (**Figure 3G**).
227

Structural organization of L-PSI-LHCII.

To gain further insight into the structural organization of the L-PSI-LHCII, we performed
single-particle analysis using negative staining electron microscopy (**Figure 4**). The 2D class
average projections show a density corresponding to PSI-LHCI, one to trimeric LHCII and an
extra density corresponding to the additional Lhca1-Lhca4 dimer. The Lhca1-Lhca4 dimer

interacts PSI-core on the PsaB side (**Figure 4D-F**), while the LHCII trimer is connected to

the PsaA, side, as observed in land plants (Wu et al., 2023).

Figure 4: Architecture of the L-PSI-LHCII.

(**A-F**) Negative staining single particles analysis of the L-PSI-LHCII purified from *P.*

oceanica. (**a**) 2D EM projection class average size = 1441 particles (**b**) 2D class average size

= 2787, (**c**) 2D class average size = 2629 particles. (**D-F**) The atomic models of the existing

structure (5ZJI) were superposed on the L-PSI-LHCII projection 2D class averages: PSI core

(green), Lhca1 (red), Lhca2 (magenta), Lhca3 (yellow), Lhca4 (blue) and LHCII trimer

(cyan).

Energy transfer dynamics in the PSI-LHCI and L-PSI-LHCII complexes

To investigate excitation energy transfer (EET) and trapping in *P. oceanica* PSI-LHCI and L-

PSI-LHCII complexes, we performed time-resolved fluorescence measurements with 400 nm

and 475 nm excitation, which excite the core and the outer antennae differently (see

Suplemental Figure 11 for streak camera images). Global analyses were performed on these

datasets and four components were sufficient to fit each dataset satisfactorily (**Figure 5A-B**).

The fast component (<10 ps) represents EET from high to low-energy Chls, while the ns

component, with small amplitude and a blue-shifted spectrum corresponds to disconnected

Chls. The dominant ~20 ps (in PSI-LHCI) and ~28 ps (in L-PSI-LHCII) component, has a

similar spectrum in all datasets and is associated with trapping excitations in the RC from

bulk Chls, mainly in the PSI core. A red-shifted 58 ps (in PSI-LHCI) and 94 ps (in L-PSI-LHCII) component, with larger amplitude upon 475 nm excitation, describes trapping in the RC from excitations mainly originating in the antenna. The longer lifetime in L-PSI-LHCII reflects its larger antenna. Notably, the second component also shows a slight increase in lifetime in L-PSI-LHCII (from ~20 ps to ~28 ps) and relative amplitude, suggesting the presence of an antenna complex without red forms that transfers energy rapidly to the PSI core.

Figure 5: Excitation energy transfer and trapping of PSI-LHCI and L-PSI-LHCII. Decay-associated spectra (DAS) for the measurement of the PSI-LHCI sample (solid lines) and L-PSI-LHCII sample (short-dots) upon 400 nm **(A)** and 475 nm **(B)** excitation. The * indicates a fixed lifetime component. **(C)** Simplified kinetic model for the target analysis, see **Suplemental Figure 12** for the full scheme and an accompanying descriptive text. **(D)**

Species-associated Spectra (SAS) as extracted from the target analysis. Bulk Chls are in black
and the red forms in red.

To better understand the EET dynamics, we simultaneously fitted the four datasets using a
target model (summarized in **Figure 5C**). The full kinetic scheme and fitting quality are
presented in **Suplemental Figure 12**. Target analysis reveals that excitations in the bulk Chls
are trapped in 23 ps, consistently with previous studies on different organisms (Engelmann et
al., 2006; Wientjes et al., 2011; Le Quiniou et al., 2015) as expected as the PSI core is highly
conserved among plants, algae and cyanobacteria (Alboresi et al., 2017). EET from red forms
to bulk Chls occurs in 47 ps, while the reverse transfer has a lifetime of 139 ps. This is a
consequence of the significantly higher amount of bulk Chls than red forms in the system,
providing an entropic advantage to this compartment, although it is higher in energy. The
Species-Associated Spectra (SAS) (**Figure 5D)** confirm the attribution of the compartments
to bulk Chls and red forms. The extra LHCII trimer is energetically reasonably well
connected to the bulk Chls (forward time constant of 35 ps and a reverse time constant of 686
ps). The extra Lhca14 dimer is instead less well connected to the bulk: it exchanges energy
with the bulk at a forward rate of $(175 \text{ ps})^{-1}$ and a reverse rate of $(819 \text{ ps})^{-1}$.

The antenna size of both Photosystems remains constant with depth.
The low Chl *a/b* ratio of *P. oceanica* suggests that more LHCII than in e.g *A.thaliana* are
present in the membrane. Using the method from (Ünlü et al., 2014) we could estimate that
there should be between 6 and 8 LHCII trimers per PSII, with only minor changes between
depths (**Suplemental Figure 3B**).
The functional antenna size of PSII was determined following the fluorescence rise kinetics
in DCMU-treated leaves (Erickson et al., 1984) (**Figure 6A**). The Antenna size was slightly
lower at 2 m compared to 15/26 m but, overall, it appeared highly similar at all depths
(**Figure 6A**), indicating that a change in PSII antenna size is not a major adaptation strategy
to depth in *P. oceanica*. Next, we tested the the distribution of LHCII between the two
photosystems. The 77K fluorescence emission spectra of the isolated thylakoids showed a
depth-dependent increase of the 715 nm emission relative to 680 nm (**Figure 6B**). These
differences can have various origins as the signal depends not only on the number of
pigments associated with the two photosystems but also on their lifetimes (in this respect is
important to consider that PSI lifetimes becomes long and multiexponential at 77K) and the
connections between the complexes. Thus, both the composition and organization of the

thylakoids could explain the changes in fluorescence emission with depth (Rantala et al.,

2017).

For a more quantitative approach, we performed time-resolved fluorescence measurements on

winter (**Figure 6C**) and summer thylakoids (**Suplemental Figure 13A**). This technique

allows us to disentangle the contributions of the two photosystems. The lifetime of PSI will

inform about the antenna size of the complex and the area under the components of PSI and

PSII proximates   the relative number of pigments associated with the two photosystems.

Exiting at 400 nm and 475 nm provide information on the migration time from the antenna to

the PSI core.

Three components were sufficient to fit the datasets (**Figure 6C and Suplemental Figure**

13A). The first component (<10 ps, black in **Figure 6C**) represents EET from blue to red

chlorophylls in PSI corresponding to the fast component in the isolated PSI complexes

(**Figure 6C**). The second component (red in **Figure 6C**) exhibits the far-red signature of PSI

and has similar lifetimes and spectra (**Suplemental Figure 13B** for PSI DAS comparison)

across depths, indicating no differences in PSI antenna size. However, both spectra and

lifetimes are excitation wavelength-dependent. The difference in PSI lifetime between

excitations (~100 ps at 475 nm and ~80 ps at 400 nm) indicates that the LHCs (which contain

Chl *b*) significantly contribute to the PSI antenna with some complexes transferring energy

relatively slowly to the RC. This agrees with the measurements on the isolated L-PSI-LHCII

complex. Moreover, the PSI spectrum upon 475 nm excitation shows a relatively larger

amplitude at 680 nm than above 700 nm, (**Suplemental Figure 13**), indicating that the outer

antenna complexes mainly contribute to the emission at 680 nm, while the far-red emission

(>700 nm) originates from the core. The third component (blue in figure 6) also has similar

lifetimes and spectra in all datasets and is associated with PSII. While lifetimes and spectra

remain similar at different depths for all components, depth-dependent changes are observed

in the relative areas of the components associated with PSI and PSII, increasing slightly with

depth with the largest variation (from 0.9 to 1.1) in summer leaves. This result indicates a

change in the excitation between PSI and PSII, resulting from a difference in the number of

Chls associated with the two photosystems at different depths. Considering that the PSII

antenna size does not change (**Figure 6A**), there are still two remaining explanations for this

result: (i) a change in the PSI antenna size and (ii) a change in PSI/PSII ratio. Since the PSI

lifetime and spectra remain unchanged with depths, indicating no changes in PSI antenna

size, we conclude that these differences are due to a small increase in the PSI/PSII ratio from

2 m to 26 m depths.

Figure 6: PSI and PSII light harvesting in *P. oceanica* at different growing depths.

(**A**) Functional antenna size of PSII. PSII turnover rates (in $e^-.s^{-1}$), derived from the
fluorescence rise kinetics of DCMU-blocked leaves collected in winter, were measured at
three moderate (100, 125, and 170 $\mu mol.m^{-2}.s^{-1}$) intensities of red light. Slope values measure
PSII functional antenna size. (**B**) Normalized 77K emission fluorescence of thylakoids from
leaves collected at depths of 2, 15, and 26 m in winter**. (C)** Decay-associated spectra obtained
from global fitting of time-resolved fluorescence spectra. Thylakoids from leaves collected at
depths of 2, 15, and 26 m in winter were excited at 400 (top) and 475 nm (bottom). (**D**)
Kinetics of NPQ induction. NPQ is probed on low light-adapted *P. oceanica* leaves exposed
to moderate actinic illumination (200 $\mu mol.m^{-2}.s^{-1}$) n=3. (**E**) Immunodetection of PsbS in *P.*
oceanica leaf exctract. Large rubisco subunit was used as a standard for protein loading.
351

Non-photochemical quenching and Psbs expression at different growth depths.
Balancing the energy distribution between photosystems may also occur through modulation
of their photochemical efficiency (ultimately affecting their functional antenna size). Shallow
P. oceanica leaves showed a ~3 times higher non-photochemical quenching (NPQ) than 26 m
depth leaves, while 15 m depth leaves displayed an intermediate NPQ level (**Figure 6D**).
PsbS expression followed this pattern (**Figure 6E**), showing an increase in shallow/high-light
plants, and conversely a strong decrease in deep/low-light plants. Like *A. thaliana*, PsbS
expression is light-depending (Albanese et al., 2016) and NPQ depends on the amount of the
PsbS protein (Li et al., 2002).
361

L-PSI-LHCII complex in other Mediterranean seagrasses
The presence of the L-PSI-LHCII was investigated across four mediterranean seagrass
species with different depth growth limits: *Cymodocea nodosa* (1-40 m), *Zostera marina* (up
to 15 m) and *Ruppia maritima*, and *Zostera noltii* (up to 2 m). All species were harvested at
0.5-1 m depth. 77k fluorescence spectra of isolated PSI revealed a maximum emission peak at
720 nm for all species (**Figure 7A**). Comparing Chl *a/b* ratios in purified thylakoids showed
that the shallow plants *R. maritima* (3.01) and *Z. noltii* (2.77) have a ratio higher than *P.*
oceanica grown at similar irradiance, whereas *C. nodosa* (2.14) is closer to *P. oceanica*
(1.95), suggesting a higher LHCII abundance in species with wider depth distribution. Z.
marina (0.5-15 m) had an intermediate Chl *a/b* ratio (2.27) between shallow and deep
seagrasses (**Figure 7B**).
BN-PAGE was performed to analyze thylakoid complexes in the selected plant species. *Z.*
marina and *C. nodosa* showed a strong green band, matching the molecular weight of L-PSI-
LHCII (**Figure 7C**). In contrast, *Z. noltii* and *R. maritima* exhibited migration profiles similar
to *A. thaliana*, as the position of the L-PSI-LHCII is occupied by PSII supercomplexes as

seen by the 2D-PAGE analysis (**Suplemental Figure 14**) suggesting the absence of L-PSI-

LHCII in plants specifically thriving in shallow waters. To confirm the absence of L-PSI-

LHCII in *A. thaliana* and shallow seagrasses, the Lhca4 protein signal was compared to that

of Lhca2 in *P. oceanica*, *A. thaliana*, *R. maritima*, and *Z. noltii* using immunoblotting

(**Suplemental Figure 15**). The Lhca4/Lhca2 ratio from the blots was normalized to the

Lhca4/Lhca2 ratio in the purified PSI-LHCI, where the stoichiometry between both proteins

was assumed to be 1. In *P. oceanica*, the ratio was 1.43 ± 0.06, while in *R. maritima*, *Z.*

noltii, and *A. thaliana*, it was 1.02 ± 0.06, 0.99 ± 0.07, and 1.07 ± 0.07, respectively. This

indicates that seagrasses with a shallow distribution, similar to the land angiosperm *A.*

thaliana, have only a few/no PSI complexes with an additional Lhca1-Lhca4 antenna. In

contrast, the strong accumulation of L-PSI-LHCII is a characteristic of seagrasses able to

colonize a wider bathymetric area.

Figure 7: Thylakoid composition of different marine plant species. (A) 77K fluorescence emission spectra of the isolated PSI-LHCI from the selected plant species (indicated) normalized at 720 nm. The complexes were isolated by CN-PAGE. **(B)** Chl *a/b* of the thylakoid membranes from the selected plants. Data are expressed as the mean ± SD, n = 3. **(C)** BN-PAGE of solubilized thylakoids extracted in different seagrass species.

Discussion

Chl *b* within LHCII, mostly associated with PSII, enhance its absorption of blue-green wavelengths, whereas red and far-red light preferentially excite PSI. Consistent with this spectral specialization, *P. oceanica* exhibits an elevated PSI to PSII ratio relative to *A. thaliana*. Furthermore, *P. oceanica* demonstrates an increased expression of LHCII (**Figure 2**), which, alongside the PSI-LHCI, are enriched in Chl *b* and PSI supercomplexes were characterized by the absence of the red forms (**Figure 3**). Consequently, the antenna size of PSI and its blue light absorption are enhanced in *P. oceanica* compared to land angiosperms, with at least 30% of PSI existing as L-PSI-LHCII supercomplexes, characterized by the inclusion of a Chl *b*–rich LHCII trimer and an additional Lhca1-Lhca4 heterodimer.

Surprisingly, LHCII levels, PSI/PSII ratio and PS's antenna size in *P. oceanica* remain high and rather stable even at 2 m depth despite the substantial differences in both the quality and intensity of light throughout depth. In contrast, as seen in other organisms (Zuo, 2025) , we showed a greater NPQ activation in shallow plants, with this capacity varying by depth through PsbS protein expression level (**Figure 7**). More importantly, microscopy observations showed that thylakoid's organisation changed drastically (**Figure 1**) as a consequence of the depth-specific light constraints. Specifically, the reduced grana size observed at 2 meters depth may enhance PSII repair by facilitating the mobility of light-damaged PSII complexes toward the grana margins to facilitate repair mechanisms as suggested by (Procaccini et al., 2017). Conversely, at 26 meters, a wider yet thinner grana combined may facilitate light penetration into the grana's internal stacks. A comparable structure has been already observed in *A. thaliana* where it would allow greater efficiency for linear electron transfer under low light (Garty et al., 2024). Nevertheless, (Procaccini et al., 2017) showed that the linear electron transfer (LET) measured *in situ* in *P. oceanica* decreases in deeper meadows as a consequence of reduced irradiance. All together, these observations suggest that photosynthetic efficiency in *P. oceanica* is primarily influenced by

light intensity and quality, as well as the organization of thylakoid membranes, rather than
solely by antenna size or photosystem stoichiometry. We propose that *P. oceanica* maintains
its photosynthetic apparatus in a configuration maximizing light harvesting at all depth. Yet, a
dynamic modulation of antenna connectivity notably through NPQ, particularly in shallow
waters, would be essential for coping with fluctuating light intensities throughout the day
(Procaccini et al., 2017). A complementary explanation is that as depth increases, canopy
density decreases, partially compensating for the reduced light availability. Conversely, in
shallower meadows, leaf density results in self-shading (Dalla Via et al., 1998) and they are
more prone to colonization by epiphytic organisms (Tsirika et al., 2007), meaning that the
actual light received by the leaves is lower than the incident irradiance measured
(**Suplemental Figure 1**).

In land plants, red forms located in Lhca3 and Lhca4 absorb light below the P700
reaction center energy. Although they are essential for harvesting far-red light, they slow
PSI's energy trapping (Engelmann et al., 2006; Jennings et al., 2013; Chukhutsina et al.,
2020). In marine environments depleted of far-red light, *P. oceanica* lost those red-forms,
allowing it to increase its PSI antenna size, through the connection of the extra Lhca1-Lhca4
dimer, while preserving high energy-trapping efficiency (**Figure 5**). Additionally, the longer
trapping time of *P. oceanica*'s PSI in the membrane compared to L-PSI-LHCII (100 ps vs. 55
ps at 475 nm excitation) suggests that PSI *in vivo* includes at least one more LHCII trimer,
possibly lost during purification as seen in other studies (Croce, 2020) (**Figure 6,**
Suplemental Figure 12 and 13). It can be hypothesized that the L-PSI-LHCII could serve as
an anchor point for the binding of one or multiple LHCII, notably via the canonical and/or
extra Lhca4 subunits as suggested in *A.thaliana* (Benson et al., 2015).

The presence of L-PSI-LHCII in *P. oceanica*, absent or only marginally expressed in
land plants (Crepin et al., 2020), suggests an evolutionary adaptation to low and blue-shifted
underwater light. We propose that this complex, which may play a pivotal functional and
ecological role in euryphotic seagrasses with an extensive bathymetric distribution and
persists as a vestigial element in land angiosperms, potentially representing an evolutionary
relic from the aquatic origins of angiosperms (Gomez et al., 2015). Furthermore, similar
strategies such as a high PSI/PSII ratio, high LHCII content together with an enrichment of
Chl *b* in the LHCII*,* are also expressed in green marine macroalgae *Bryopsis maxima* and
Ulva pertusa (Yamazaki et al., 2005)[64], highlighting an example of convergent evolution

between green algae and seagrasses. Based on our analysis of *P. oceanica*'s adaptation
strategies to different light conditions at various depths, it is clear that this plant exhibits
unique photosynthetic characteristics distinct from those of land angiosperms but shared with
green marine algae.

Since the half of the 20th century, coastal urbanization has significantly degraded *P.*
oceanica habitats by causing direct destruction and increasing water turbidity, thereby
limiting the growth of *Posidonia oceanica* to 30 m depth in urbanized areas. *P. oceanica* is
now facing new challenges, such as rising sea levels, which will reduce the light balance at
the lower limit of the meadow in the future, and especially at the current time the rapid-
growing marine angiosperm *Halophila stipulacea*, which is capable of acclimating to depths
of up to 60 meters. Although it also expresses the L-PSI-LHCII complex (**Supplemental**
Figure 16), *H. stipulacea* does not form the dense, persistent matte characteristic of *P.*
oceanica, which are critical for carbon sequestration. Effective conservation of *P. oceanica*
must prioritize protecting existing meadows and limiting coastal urbanization. Enhancing
water transparency could allow *P. oceanica* to recolonize greater depths and improve its
resilience in the face of rising sea levels while controlling the spread of invasive *H. stipulacea*
would preserve *P. oceanica* potential niche. Collectively, these efforts will help sustain and
possibly expand this essential marine carbon sink.

METHODS

Sample collection

P. oceanica shoots were collected from three healthy meadow patches at 2, 15, and 26 m
depth near Frioul Island, Marseille, France (43.1611 N, 5.1732 E), avoiding the fragile
upper (0.5 m) and lower (29 m) limits. Sampling occurred in January and June of 2019–
2022 between 9–11 am. Plants were kept at 4 °C in the dark and processed in the lab
within 3 hours: leaves were separated at 3 °C under green light, and epiphytes removed by
seawater washing. *C. nodosa* and shallow *P. oceanica* were collected in Agay (43.4299 N,
6.8680 E) in June 2021 (0.5 m depth), and *Z. marina* and *Z. noltii* in Port de Bouc
(41.2158 N, 4.5152 E) in July 2021 (0.5 m depth). A. thaliana WT Col-0 plants were
grown for 4 weeks under 120 µmol photons m^{-2} s^{-1} white light.

Isolation and separation of photosystem supercomplexes

Thylakoid membranes were prepared as described in (Charras et al., 2024).

Thylakoids membrane preparation for separation of photosystem supercomplexes by Blue-Native-PAGE.

Thylakoid membranes were washed 2 times in a solubilization buffer (50 mM BisTris pH=7, 25 mM EDTA, 20% glycerol, 10 mM NaF). The chlorophyll *a/b* ratio and chlorophyll concentration were determined according (Croce et al., 2002; Chazaux et al., 2022). At 4°C, purified thylakoids (1 mg/ml) were solubilized with a final concentration of 2% (v/w) n-Dodecyl-□-D-Maltopyranoside solution (α-DDM, Anatrace, Ref: D310HA 10 GM) for 20 min, then centrifuged at 21,000 g. The solubilizate was then used for Blue native-PAGE sample preparation.

Separation of photosystem supercomplexes by Blue native-PAGE

As in Järvi et al., 2011[37], the solubilizate was supplemented with 1/10 volume of Serva Blue loading buffer (100 mM BisTris pH 7.0, 0.5 M amino-caproic acid, 30% (w/v) sucrose and 50 mg/mL Serva Blue G). About 5-10 µg of chlorophyll was then loaded on the commercial gel (Invitrogen™ NativePAGE™ 4 -16□%, Bis-Tris, 1.0□mm; ref BN1001BOX) set in Minicellule XCell *SureLock*™ Invitrogen™. The anode buffer contained 25 mM BisTris, 25 mM tricine pH=7, and the cathode buffer contained 50 mM BisTris pH=7, 0.015% □DDM, and 0.01% Serva Blue G250. Electrophoresis was performed at 4°C for 7 hours. The intensity was fixed and set so that at the start of the run, the voltage should be around 40V.

Second Dimension urea-PAGE identification of polypeptides content from supercomplexes isolated by BN-PAGE

The lanes of interest were cut from the 1D BN-PAGE and incubated for 30 min in Laemmli denaturing buffer (125 mM Tris pH 6.8, 20% glycerol, 2% SDS, 100 mM DTT, 0.01% bromophenol blue). The lane was then deposited on the top of 13% acrylamide/bisacrylamide (37.5/1) resolving gel containing 6 M urea and fixed with agarose (1% agarose, 0. M Tris pH 6.8, 0.001% bromophenol blue). The anode and cathode chambers were filled with a Laemmli running buffer (25 mM Tris pH 8.3, 0.192 M glycine, 0.1% SDS). Migration was performed for approximately 2h30 at 90 V.

Urea-PAGE and immunoblot

Thylakoid preparations were denatured at 70 °C for 30 min in denaturing Laemmli buffer, then loaded on a 14% acrylamide gel (37.5:1 acrylamide/bisacrylamide) containing 6 M urea (Tris-Glycine Laemmli system). For Phos-tag™ analysis, 12% gels with 50 μM Phos-Tag (Wako Pure Chemicals Industries) and 100 μM MnCl were used. After electrophoresis, gels were stained with Sypro™ Ruby or used for immunoblotting. Phos-tag™ gels were equilibrated in transfer buffer with 50 mM EDTA before transfer using the BioRad Trans-Blot Turbo system. Immunoblots were developed using Agrisera antibodies and chemiluminescence detection (Fusion FX7, Vilber). Leaf extracts were prepared as described in (Charras et al., 2024).

L-PSI-LHCII and LHCII purification

Purified thylakoids (1 mg/ml) were solubilized (same buffer as Native-PAGE without glycerol) with a final concentration of 2% (v/w) α-DDM for 20 min, then centrifuged at 21,000 g. The supernatant was loaded onto a sucrose gradient (0.75 M sucrose, 40 mM Hepes pH 7.4, 0.015% α-DDM), formed by freeze-thawing. After centrifugation, target bands were collected, dialyzed against 40 mM Hepes pH 7.4 concentrated. For LHCII, the sample underwent a second sucrose gradient, followed by dialysis and concentration. For L-PSI-LHCII, PSI-LHCI contaminants were partially removed after the first gradient using anion exchange chromatography (MonoQ, 0–500 mM NaCl in 40 mM Hepes pH 7.4, 0.015% α-DDM).

Electron microscopy analysis of chloroplast ultrastructure

Samples were prepared as in (Nja et al., 2018). Grids were observed under the Tecnai G2 electron microscope.

Sample preparation for negative staining single particle electron microscopy analysis

5 μL of the protein sample (3 mg.mL^{-1}) were applied to the glow-discharged (PELCO easyGlow, 15mA, 25 s) Quantifoil 100 Carbon Support Films grid: Cu 300 and left on the grid for 45 sec. The excess sample was blotted away (Whatman #1). 5 μl of 2% uranyl acetate was applied to the grid. Excess stain excess was blotted away (Whatman #1) for 30s. Grids were washed by applying a 5% uranyl acetate droplet (5 μl), removed immediately with blotting paper (Whatman #1). After drying out, the grids were loaded in the Thermo Fisher Scientific Glacios using an autoloader under cryo-EM conditions.

Electron microscopy data acquisition

Data acquisition was performed on the Thermo Fisher Scientific Glacios equipped with a field emission gun and operated at 200 kV in bright field imaging mode. Movies were recorded using a Thermo Scientific Falcon 3EC Direct electron Detector in linear mode at a nominal magnification of 92,000, corresponding to a pixel size of 1.567 Å/pix with 30 frames at a dose of 30 e-/Å2 per frame and an exposure time of 30 s per movie.

Image processing

The dataset of PSI protein (fraction p1q_15) was derived from 1326 movies. The raw movies were imported into SCIPION 3.0 (de la Rosa-Trevín et al., 2013) for the following processing steps. The movies were motion-corrected using the MotionCor2-1.4.7 protocol (Zheng et al., 2017) and the CTF estimation was done by the Gctf – 1.06 protocol (Zhang, 2016).

A total of 337227 particles were extracted after xmipp manual/auto particle picking[55]. After multiple rounds of 2D classification using cryoSPARC v3.0.1 protocol (Punjani et al., 2017), the main protein complexes of interest: PSI and PSI-LHCII, with apparent features, were sorted. Based on the 2D classification, the particles exhibiting the clear signatures of PSI-LHCI and L-PSI-LHCII were manually sorted into two groups: 1) PSI-LHCI, containing 168432 particles and 2) L-PSI-LHCII with 14255 particles.

77K fluorescence spectra

For the L-PSI–LHCII and PSI-LHCI from *P. oceanica* the supercomplexes were isolated as described above (sucrose gradient and AEC) and for the other plant species, the supercomplexes were isolated by CN-PAGE. The band was sliced and deposited in an aluminum holder and snapped frozen in liquid nitrogen at 77 K. By means of an optic fiber, chlorophyll is excited by a blue (~455 nm) LED, and the resulting fluorescence signal is detected through a red-colored Kodak Wratten filter. Excitation and detection are mediated with an Ocean Optics USB2000 CCB spectrometer, piloted with an in-lab-developed PC acquisition software.

Determination of PSI/PSII stoichiometry via ECS spectroscopy

In the green lineage, absorbance changes at 515–520 nm directly reflect thylakoid trasmembrane potential (Joliot and Joliot, 1989). A Joliot-type spectrophotometer (Continuum Nd:YAG Laser + OPO) was used with laser flashes (680 nm) for excitation,

and BG39 filters to block actinic light. In *P. oceanica* leaves, a saturating flash induces a

ΔA520 nm peak, indicating membrane potential from one-electron transfer per photosystem

(PSII + PSI). When PSII is inhibited (DCMU 10 µM + hydroxylamine 10 mM), a reduced

peak reveals PSI's sole contribution. The PSI/PSII ratio is then calculated from peak

amplitudes (Bailleul et al., 2010). This method reflects only effective charge separation,

independent of antenna size variations.

Determination of PSII antenna sizes via DCMU-fluorescence rise

PSII functional antenna sizes have been determined using a Dual-PAM 100 fluorimeter

(Melis and Homann, 1975; Lavergne and Trissl, 1995). In brief, the fluorescence rise (FR)

kinetics was measured on *P. oceanica* leaves upon pulses (500 ms) of red light. Leaves were

pre-treated with DCMU (to block all electron transfers downstream PSII) and kept in

darkness at room temperature in fresh seawater. The area above the sigmoid curve

(normalized from 0 to 1) is proportional to the still-oxidized primary acceptor Q_A fraction.

The evolution of the Q_A reduction rate can be obtained by integrating the FR curve: the initial

rate at light onset (slope at origin) gives the PSII functional antenna size.

Non-photochemical quenching of fluorescence

Fluorescence kinetics were followed on fresh, low light-adapted *P. oceanica* leaves with a

DualPAM-100 fluorometer. The maximal fluorescence yield *Fm* is first probed by a

preliminary saturating light pulse on a dark background. After light onset, successive

saturating pulses probe the evolution of the maximal fluorescence yield during

photochemistry *Fm'*. The NPQ parameter is computed as (*Fm-Fm'*)/*Fm'* for each

measurement.

Time-resolved fluorescence spectroscopy

Time-resolved fluorescence measurements on *P. oceanica* samples were performed at room

temperature using a synchroscan streak camera setup as previously described in Hu et al.

(2023, New Phytol.). Femtosecond pulses at either 400 nm or 475 nm with a repetition rate of

250 kHz were used. The excitation spot size was ~150 µm. An excitation power study was

performed for each sample. The thylakoids were treated with 20 µM DCMU to close the

reaction centers. The samples were magnetically stirred in 1 cm × 1 cm cuvette, and the OD

at Q_y maximum was ~0.5 for the thylakoids and ~0.3 cm^{-1} for the PSI samples. The

fluorescence was detected at a right angle using a spectrograph (Chromex 250IS, 50 l/mm)

and a synchroscan streak camera (Hamamatsu C5680). Two-dimensional streak images were

corrected for the detector's background and wavelength-dependent sensitivity. The resulting

averaged streak images were sliced up into time traces spanning ~3 nm. Processed streak

images were then globally analyzed using the Glotaran (Mullen and Stokkum, 2007;

Snellenburg et al., 2012) and TIMP package for R to determine the fluorescence lifetimes and

decay-associated spectra (DAS). Briefly, the streak images were fitted with a sum of four

exponentials. The amplitudes of these exponentials across emission wavelengths resulted in

decay-associated spectra.

Target analysis

The streak images of the PSI-LHCI and L-PSI-LHCII samples excited at both 400 and 475

nm were simultaneously analysed in a target analysis using the pyglotaran Python package

(Weißenborn et al., 2021; van Stokkum et al., 2023). The full kinetic scheme is presented in

Suplemental Figure 12. To allow for the estimation of the initial excitation densities for the

different compartments the areas under the SAS were constrained to be equal (Maiuri et al.,

2013). The initial excitation densities are placed in the precursor compartments (precursor

compartment names start with ET and they are grouped in the grey bar in **Suplemental**

Figure 12. They were free fitting parameters for each measurement. The precursor

compartments transfer rapidly (2.0-3.3 ps) and within the instrument response time (modeled

as a Gaussian function with FWHM ranging from 5.3-6.7 ps) to the principal compartments,

which are the 'Bulk', 'Red', 'LHCII' and 'xLhca14'. The precursor compartments represent

the fast energy equilibration within the subcomplexes and can be compared to the first (black)

DAS of the global analyses. For each dataset/measurement the SAS of the precursor

compartments were constrained to be equal. For each dataset/measurement there is also the

'Disc.' compartment to model the disconnected Chls. The measurements conducted on the

PSI-LHCI samples were fitted with a model that for the principal compartments only

contained the 'Bulk' and the 'Red' compartments, whereas for the measurements on the L-

PSI-LHCII sample the model additionally contained the 'LHCII' and the extra Lhca1-Lhca4

dimer (Lhca1²-Lhca4²) compartment. The SAS of the principal compartments were

constrained to be equal across all datasets. The backsweep of the synchroscan streak camera

was also modeled and its frequency was fixed to 76 MHz. The fit quality of the target model

is excellent and shown for a range of wavelengths in **Suplemental Figure 17.**

Absorption spectrum deconvolution

For the spectral deconvolution the measured absorption spectra of the *P. oceanica* LHCII

trimer, PSI-LHCI and L-PSI-LHCII complexes were first normalised to their integrated

absorption in the Q_y region (630-750 nm) and then scaled to their relative Chl content as

presented in **Suplemental Figure 9A**, taking into account the oscillator strength ratio of 1:0.7

for Chl *a:b*. The extra Lhca14 spectrum was obtained by subtracting the LHCII trimer and the

PSI-LHCI absorption spectrum from the L-PSI-LHCII absorption spectrum.

Mass spectrometry

Native-PAGE bands from the L-PSI-LHCII and *p.o* PSI-LHCI complexes were analyzed in

triplicate by LC-MS/MS (Q-Exactive Plus, ThermoFisher), following (Proietti et al., 2023)

with minor modifications. Samples were digested with Trypsin/Lys-C (Promega), and data

processed using Proteome Discoverer 2.4.1.15 with the *A. thaliana* database (SwissProt,

TxID 3702, version 2017-10-25) supplemented with *P. oceanica* reaction center and Lhca

sequences. Peptides were validated via Sequest, using best PSM score (ΔCn $\leq$ 0.05) and 1%

FDR. Spectral counting of unique peptide PSMs for Lhca proteins, normalized to core

subunit PSMs, allowed estimation of relative Lhca abundance and L-PSI-LHCII/PSI-LHCI

ratio.

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this work the author(s) used ChatGPT in order to improve language

and readability. After using this tool/service, the author(s) reviewed and edited the content as

needed and take(s) full responsibility for the content of the publication.

References:

Albanese, P., Manfredi, M., Meneghesso, A., Marengo, E., Saracco, G., Barber, J.,
Morosinotto, T., and Pagliano, C. (2016). Dynamic reorganization of photosystem II
supercomplexes in response to variations in light intensities. *Biochimica et Biophysica*
Acta (BBA) - Bioenergetics **1857**:1651–1660.

Alboresi, A., Le Quiniou, C., Yadav, S. K. N., Scholz, M., Meneghesso, A., Gerotto, C.,
Simionato, D., Hippler, M., Boekema, E. J., Croce, R., et al. (2017). Conservation
of core complex subunits shaped the structure and function of photosystem I in the
secondary endosymbiont alga Nannochloropsis gaditana. *New Phytol* **213**:714–726.

Arnaud-Haond, S., Duarte, C. M., Diaz-Almela, E., Marbà, N., Sintes, T., and Serrão, E.
A. (2012). Implications of Extreme Life Span in Clonal Organisms: Millenary Clones
in Meadows of the Threatened Seagrass Posidonia oceanica. *PLOS ONE* **7**:e30454.

Bailleul, B., Cardol, P., Breyton, C., and Finazzi, G. (2010). Electrochromism: a useful
probe to study algal photosynthesis. *Photosynth Res* **106**:179–189.

Benson, S. L., Maheswaran, P., Ware, M. A., Hunter, C. N., Horton, P., Jansson, S., Ruban, A. V., and Johnson, M. P. (2015). An intact light harvesting complex I antenna system is required for complete state transitions in Arabidopsis. *Nature Plants* **1**:1–9.

Bos, P. R., Schiphorst, C., Kercher, I., Buis, S., de Jong, D., Vunderink, I., and Wientjes, E. (2023). Spectral diversity of photosystem I from flowering plants. *Photosynth Res* **155**:35–47.

Boudouresque, C.-F., Bernard, G., Bonhomme, P., Charbonnel, E., Diviacco, G., Meinesz, A., Pergent, G., Pergent-Martini, C., Ruitton, S., and Tunesi, L. (2012). *Protection and conservation of Posidonia oceanica meadows*. RAMOGE and RAC/SPA.

BOUDOURESQUE, C. F., GRAVEZ, V., MEINESZ, A., MOLENAAR, H., PERGENT, G., and VITIELLO, P. (2013). Posidonie□: réimplantations et transplantations Advance Access published 2013.

Boudouresque, C. F., Pergent, G., Pergent-Martini, C., Ruitton, S., Thibaut, T., and Verlaque, M. (2016). The necromass of the Posidonia oceanica seagrass meadow: fate, role, ecosystem services and vulnerability. *Hydrobiologia* **781**:25–42.

Buschmann, C., Meier, D., Kleudgen, H. K., and Lichtenthaler, H. K. (1978). REGULATION OF CHLOROPLAST DEVELOPMENT BY RED AND BLUE LIGHT. In *Annual European Symposium on Photomorphogenesis* (ed. Song, P.-S.), pp. 195–198. Pergamon.

Charras, Q., Rey, P., Guillemain, D., Dourguin, F., Laganier, H., Peschoux, S., Molinié, R., Ismaël, M., Caffarri, S., Rayon, C., et al. (2024). An efficient protocol for extracting thylakoid membranes and total leaf proteins from Posidonia oceanica and other polyphenol-rich plants. *Plant Methods* **20**:38.

Chazaux, M., Schiphorst, C., Lazzari, G., and Caffarri, S. (2022). Precise estimation of chlorophyll a, b and carotenoid content by deconvolution of the absorption spectrum and new simultaneous equations for Chl determination. *Plant J* **109**:1630–1648.

Chow, W. S., Melis, A., and Anderson, J. M. (1990). Adjustments of photosystem stoichiometry in chloroplasts improve the quantum efficiency of photosynthesis. *Proc Natl Acad Sci U S A* **87**:7502–7506.

Chukhutsina, V. U., Liu, X., Xu, P., and Croce, R. (2020). Light-harvesting complex II is an antenna of photosystem I in dark-adapted plants. *Nat. Plants* **6**:860–868.

Crepin, A., Kučerová, Z., Kosta, A., Durand, E., and Caffarri, S. (2020). Isolation and characterization of a large photosystem I-light-harvesting complex II supercomplex with an additional Lhca1-a4 dimer in Arabidopsis. *Plant J* **102**:398–409.

Croce, R. (2020). Beyond "seeing is believing": the antenna size of the photosystems in vivo. *The New Phytologist* Advance Access published 006 2020, doi:10.1111/nph.16758.

Croce, R., Canino, G., Ros, F., and Bassi, R. (2002). Chromophore organization in the higher-plant photosystem II antenna protein CP26. *Biochemistry* **41**:7334–7343.

Dalla Via, J., Sturmbauer, C., Schönweger, G., Sötz, E., Mathekowitsch, S., Stifter, M.,
and Rieger, R. (1998). Light gradients and meadow structure in Posidonia
oceanica:ecomorphological and functional correlates. *Mar. Ecol. Prog. Ser.* **163**:267–
278.

de la Rosa-Trevín, J. M., Otón, J., Marabini, R., Zaldívar, A., Vargas, J., Carazo, J. M.,
and Sorzano, C. O. S. (2013). Xmipp 3.0: an improved software suite for image
processing in electron microscopy. *J Struct Biol* **184**:321–328.

Díaz-Almela, E., and Duarte, C. (2008). Management of Natura 2000 habitats Advance
Access published 2008.

Duarte, C. M., Losada, I. J., Hendriks, I. E., Mazarrasa, I., and Marbà, N. (2013). The
role of coastal plant communities for climate change mitigation and adaptation.
Nature Clim Change **3**:961–968.

Engelmann, E., Zucchelli, G., Casazza, A. P., Brogioli, D., Garlaschi, F. M., and
Jennings, R. C. (2006). Influence of the Photosystem I–Light Harvesting Complex I
Antenna Domains on Fluorescence Decay. *Biochemistry* **45**:6947–6955.

Erickson, J. M., Rahire, M., Bennoun, P., Delepelaire, P., Diner, B., and Rochaix, J.-D.
(1984). Herbicide resistance in *Chlamydomonas reinhardtii* results from a mutation in
the chloroplast gene for the 32-kilodalton protein of photosystem II. *Proc. Natl. Acad.*
Sci. U.S.A. **81**:3617–3621.

Ferroni, L., Suorsa, M., Aro, E., Baldisserotto, C., and Pancaldi, S. (2016). Light
acclimation in the lycophyte *S elaginella martensii* depends on changes in the
amount of photosystems and on the flexibility of the light☐harvesting complex II
antenna association with both photosystems. *New Phytol* **211**:554–568.

Garty, Y., Bussi, Y., Levin-Zaidman, S., Shimoni, E., Kirchhoff, H., Charuvi, D., Nevo,
R., and Reich, Z. (2024). Thylakoid membrane stacking controls electron transport
mode during the dark-to-light transition by adjusting the distances between PSI and
PSII. *Nat. Plants* **10**:512–524.

Goldschmidt-Clermont, M., and Bassi, R. (2015). Sharing light between two photosystems:
mechanism of state transitions. *Current Opinion in Plant Biology* **25**:71–78.

Gomez, B., Daviero-Gomez, V., Coiffard, C., Martín-Closas, C., and Dilcher, D. L.
(2015). *Montsechia* , an ancient aquatic angiosperm. *Proc. Natl. Acad. Sci. U.S.A.*
112:10985–10988.

Grieco, M., Suorsa, M., Jajoo, A., Tikkanen, M., and Aro, E.-M. (2015). Light-harvesting
II antenna trimers connect energetically the entire photosynthetic machinery —
including both photosystems II and I. *Biochimica et Biophysica Acta (BBA) -*
Bioenergetics **1847**:607–619.

Hogewoning, S. W., Wientjes, E., Douwstra, P., Trouwborst, G., van Ieperen, W., Croce,
R., and Harbinson, J. (2012). Photosynthetic Quantum Yield Dynamics: From
Photosystems to Leaves. *Plant Cell* **24**:1921–1935.

Huang, Z., Shen, L., Wang, W., Mao, Z., Yi, X., Kuang, T., Shen, J.-R., Zhang, X., and
Han, G. (2021). Structure of photosystem I-LHCI-LHCII from the green alga
Chlamydomonas reinhardtii in State 2. *Nat Commun* **12**:1100.

Jennings, R. C., Zucchelli, G., and Santabarbara, S. (2013). Photochemical trapping
heterogeneity as a function of wavelength, in plant photosystem I (PSI–LHCI).
Biochimica et Biophysica Acta (BBA) - Bioenergetics **1827**:779–785.

Joliot, P., and Joliot, A. (1989). Characterization of linear and quadratic electrochromic
probes in Chlorella sorokiniana and Chlamydomonas reinhardtii. *Biochimica et*
Biophysica Acta (BBA) - Bioenergetics **975**:355–360.

Kouřil, R., Wientjes, E., Bultema, J. B., Croce, R., and Boekema, E. J. (2013). High-light
vs. low-light: Effect of light acclimation on photosystem II composition and
organization in Arabidopsis thaliana. *Biochimica et Biophysica Acta (BBA) -*
Bioenergetics **1827**:411–419.

Lavergne, J., and Trissl, H. W. (1995). Theory of fluorescence induction in photosystem II:
derivation of analytical expressions in a model including exciton-radical-pair
equilibrium and restricted energy transfer between photosynthetic units. *Biophys J*
68:2474–2492.

Le Quiniou, C., Tian, L., Drop, B., Wientjes, E., van Stokkum, I. H. M., van Oort, B.,
and Croce, R. (2015). PSI–LHCI of Chlamydomonas reinhardtii: Increasing the
absorption cross section without losing efficiency. *Biochimica et Biophysica Acta*
(BBA) - Bioenergetics **1847**:458–467.

Li, X.-P., Müller-Moulé, P., Gilmore, A. M., and Niyogi, K. K. (2002). PsbS-dependent
enhancement of feedback de-excitation protects photosystem II from photoinhibition.
Proc. Natl. Acad. Sci. U.S.A. **99**:15222–15227.

Maiuri, M., Snellenburg, J. J., van Stokkum, I. H. M., Pillai, S., Wongcarter, K., Gust,
D., Moore, T. A., Moore, A. L., van Grondelle, R., Cerullo, G., et al. (2013).
Ultrafast energy transfer and excited state coupling in an artificial photosynthetic
antenna. *J Phys Chem B* **117**:14183–14190.

Marbà, N., Duarte, C. M., Cebrián, J., Gallegos, M. E., Olesen, B., and Sand-Jensen, K.
(1996). Growth and population dynamics of Posidonia oceanica on the Spanish
Mediterranean coast: elucidating seagrass decline. *Marine Ecology Progress Series*
137:203–213.

Melis, A., and Homann, P. H. (1975). KINETIC ANALYSIS OF THE FLUORESCENCE
INDUCTION IN 3-(3,4-DICHLOROPHENYL)-1,1-DIMETHYLUREA POISONED
CHLOROPLASTS. *Photochem Photobiol* **21**:431–437.

Molenaar, H. (1992). Investigation of transplantation in cuttings of the marine phanerogam
posidonia oceanica (l.) delile. Modelling of its architecture and growth Advance
Access published 1992.

Morosinotto, T., Breton, J., Bassi, R., and Croce, R. (2003). The Nature of a Chlorophyll
Ligand in Lhca Proteins Determines the Far Red Fluorescence Emission Typical of
Photosystem I *. *Journal of Biological Chemistry* **278**:49223–49229.

Mullen, K. M., and Stokkum, I. H. M. van (2007). TIMP: An R Package for Modeling
813   Multi-way Spectroscopic Measurements. *Journal of Statistical Software* **18**:1–46.

814   **Nja, R. B., Merceron, B., Faucher, M., Fleurat-Lessard, P., and Béré, E.** (2018). NaCl –
815   Changes stem morphology, anatomy and phloem structure in Lucerne (Medicago
816   sativa cv. Gabès): Comparison of upper and lower internodes. *Micron* **105**:70–81.

817   **Pan, X., Ma, J., Su, X., Cao, P., Chang, W., Liu, Z., Zhang, X., and Li, M.** (2018).
818   Structure of the maize photosystem I supercomplex with light-harvesting complexes I
819   and II. *Science* **360**:1109–1113.

820   **Pasqualini, V., Pergent-Martini, C., Clabaut, P., and Pergent, G.** (1998). Mapping
821   ofPosidonia oceanicausing Aerial Photographs and Side Scan Sonar: Application off
822   the Island of Corsica (France). *Estuarine, Coastal and Shelf Science* **47**:359–367.

823   **Pergent, G., Bazairi, H., Bianchi, C. N., Boudouresque, C. F., Buia, M. C., Calvo, S.,**
824   **Clabaut, P., Harmelin-Vivien, M., Mateo, M. A., Montefalcone, M., et al.** (2014).
825   Climate change and Mediterranean seagrass meadows: a synopsis for environmental
826   managers. *Mediterranean Marine Science* **15**:462–473.

827   **Pergent-Martini, C., Pergent, G., Monnier, B., Boudouresque, C.-F., Mori, C., and**
828   **Valette-Sansevin, A.** (2021). Contribution of *Posidonia oceanica* meadows in the
829   context of climate change mitigation in the Mediterranean Sea. *Marine Environmental*
830   *Research* **165**:105236.

831   **Procaccini, G., Ruocco, M., Marín-Guirao, L., Dattolo, E., Brunet, C., D'Esposito, D.,**
832   **Lauritano, C., Mazzuca, S., Serra, I. A., Bernardo, L., et al.** (2017). Depth-specific
833   fluctuations of gene expression and protein abundance modulate the photophysiology
834   in the seagrass Posidonia oceanica. *Sci Rep* **7**:42890.

835   **Proietti, S., Falconieri, G. S., Bertini, L., Pascale, A., Bizzarri, E., Morales-Sanfrutos, J.,**
836   **Sabidó, E., Ruocco, M., Monti, M. M., Russo, A., et al.** (2023). *Beauveria bassiana*
837   rewires molecular mechanisms related to growth and defense in tomato. *Journal of*
838   *Experimental Botany* **74**:4225–4243.

839   **Punjani, A., Rubinstein, J. L., Fleet, D. J., and Brubaker, M. A.** (2017). cryoSPARC:
840   algorithms for rapid unsupervised cryo-EM structure determination. *Nat Methods*
841   **14**:290–296.

842   **Rantala, M., Tikkanen, M., and Aro, E.-M.** (2017). Proteomic characterization of
843   hierarchical megacomplex formation in Arabidopsis thylakoid membrane. *The Plant*
844   *Journal* **92**:951–962.

845   **Rantala, M., Rantala, S., and Aro, E.-M.** (2020). Composition, phosphorylation and
846   dynamic organization of photosynthetic protein complexes in plant thylakoid
847   membrane. *Photochem. Photobiol. Sci.* **19**:604–619.

848   **Romero, E., Mozzo, M., van Stokkum, I. H. M., Dekker, J. P., van Grondelle, R., and**
849   **Croce, R.** (2009). The Origin of the Low-Energy Form of Photosystem I Light-
850   Harvesting Complex Lhca4: Mixing of the Lowest Exciton with a Charge-Transfer
851   State. *Biophysical Journal* **96**:L35–L37.

Schumann, T., Paul, S., Melzer, M., Dörmann, P., and Jahns, P. (2017). Plant Growth
under Natural Light Conditions Provides Highly Flexible Short-Term Acclimation
Properties toward High Light Stress. *Frontiers in Plant Science* **8**.

Snellenburg, J. J., Laptenok, S., Seger, R., Mullen, K. M., and Stokkum, I. H. M. van
(2012). Glotaran: A Java-Based Graphical User Interface for the R Package TIMP.
Journal of Statistical Software **49**:1–22.

Su, X., Ma, J., Wei, X., Cao, P., Zhu, D., Chang, W., Liu, Z., Zhang, X., and Li, M.
(2017). Structure and assembly mechanism of plant C2S2M2-type PSII-LHCII
supercomplex. *Science* Advance Access published August 25, 2017,
doi:10.1126/science.aan0327.

Telesca, L., Belluscio, A., Criscoli, A., Ardizzone, G., Apostolaki, E. T., Fraschetti, S.,
Gristina, M., Knittweis, L., Martin, C. S., Pergent, G., et al. (2015). Seagrass
meadows (Posidonia oceanica) distribution and trajectories of change. *Sci Rep*
5:12505.

Tsirika, A., Skoufas, G., and Haritonidis, S. (2007). Seasonal and bathymetric variations of
epiphytic macroflora on Posidonia oceanica (L.) Delile leaves in the National Marine
Park of Zakynthos (Greece): Epiphytic macroflora on Posidonia oceanica leaves.
Marine Ecology **28**:146–153.

Ünlü, C., Drop, B., Croce, R., and van Amerongen, H. (2014). State transitions in
Chlamydomonas reinhardtii strongly modulate the functional size of photosystem II
but not of photosystem I. *Proc. Natl. Acad. Sci. U.S.A.* **111**:3460–3465.

van Stokkum, I. H. M., Müller, M. G., Weißenborn, J., Weigand, S., Snellenburg, J. J.,
and Holzwarth, A. R. (2023). Energy transfer and trapping in photosystem I with and
without chlorophyll-*f*. *iScience* **26**:107650.

Wagner, R., Dietzel, L., Bräutigam, K., Fischer, W., and Pfannschmidt, T. (2008). The
long-term response to fluctuating light quality is an important and distinct light
acclimation mechanism that supports survival of Arabidopsis thaliana under low light
conditions. *Planta* **228**:573–587.

Weißenborn, J., Snellenburg, J., Weigand, S., and Stokkum, I. H. M. V. (2021).
pyglotaran: a Python library for global and target analysis Advance Access published
March 18, 2021, doi:10.5281/ZENODO.4619635.

Wientjes, E., van Stokkum, I. H. M., van Amerongen, H., and Croce, R. (2011). The Role
of the Individual Lhcas in Photosystem I Excitation Energy Trapping. *Biophysical*
Journal **101**:745–754.

Wientjes, E., van Amerongen, H., and Croce, R. (2013). LHCII is an antenna of both
photosystems after long-term acclimation. *Biochimica et Biophysica Acta (BBA) -*
Bioenergetics **1827**:420–426.

Wientjes, E., Philippi, J., Borst, J. W., and van Amerongen, H. (2017). Imaging the
Photosystem I/Photosystem II chlorophyll ratio inside the leaf. *Biochimica et*
Biophysica Acta (BBA) - Bioenergetics **1858**:259–265.

Wu, J., Chen, S., Wang, C., Lin, W., Huang, C., Fan, C., Han, D., Lu, D., Xu, X., Sui, S.,
et al. (2023). Regulatory dynamics of the higher-plant PSI–LHCI supercomplex
during state transitions. *Molecular Plant* **16**:1937–1950.

Yadav, V. S., Yadav, S. S., Gupta, S. R., Meena, R. S., Lal, R., Sheoran, N. S., and
Jhariya, M. K. (2022). Carbon sequestration potential and CO2 fluxes in a tropical
forest ecosystem. *Ecological Engineering* **176**:106541.

Yamazaki, J., Suzuki, T., Maruta, E., and Kamimura, Y. (2005). The stoichiometry and
antenna size of the two photosystems in marine green algae, Bryopsis maxima and
Ulva pertusa, in relation to the light environment of their natural habitat. *Journal of*
Experimental Botany **56**:1517–1523.

Zhang, K. (2016). Gctf: Real-time CTF determination and correction. *J Struct Biol* **193**:1–12.

Zhang, S., Tang, K., Yan, Q., Li, X., Shen, L., Wang, W., He, Y.-K., Kuang, T., Han, G.,
Shen, J.-R., et al. (2023). Structural insights into a unique PSI–LHCI–LHCII–Lhcb9
supercomplex from moss Physcomitrium patens. *Nat. Plants* **9**:832–846.

Zheng, S. Q., Palovcak, E., Armache, J.-P., Verba, K. A., Cheng, Y., and Agard, D. A.
(2017). MotionCor2: anisotropic correction of beam-induced motion for improved
cryo-electron microscopy. *Nat Methods* **14**:331–332.

Zuo, G. (2025). Non☐photochemical quenching (NPQ) in photoprotection: insights into
NPQ levels required to avoid photoinactivation and photoinhibition. *New Phytologist*
246:1967–1974.

Author contribution

CJ initiated and developed the research project. CJ, and QCF planned and designed the

experiments.

QCF performed sample preparation, Electron microscopy, biochemistry and pigment analysis

and part of the biophysical analysis.

CM collected and analysed the data from DUAL-PAM,and Joliot spectroscopy, and measured

the 77K fluorescence spectra of thylakoids). DAS performed single-particle negative staining

electron microscopy (EM) experiments, including data collection, image analysis, and

interpretation, and contributed to result discussions. AFB measured the time-resolved

fluorescence kinetics on thylakoids and performed preliminary data analysis. EE measured

time-resolved kinetics on the purified PSI complexes, and performed the global and target

analysis of all time-resolved data sets. RC supervised the work in Amsterdam and contributed

to the analysis and interpretation of the data. MS trained QCF on protein purification and

purified the PSI and the L-PSI-LHCII supercomplex. RL performed Mass spectroscopy

experiments (data collection and analysis). DG and colleagues organized and performed all

sample collections.

QCF prepared most figures and additional files with the contribution of CM and EE.

CJ and QCF wrote the manuscript, with contributions from EE and RC. DAS, MS, DG, RC,

RL edited and reviewed the manuscript. All authors read and approved the manuscript.

Acknowledgements

We are grateful to Sandrine Ruitton, Pascal Mirleau, Deny Malengros, Fabrice Garcia,

Michel Lafont, Sandrine Chenesseau, Frédérique Legendre and Bruno Belloni from the OSU

Institut Pytheas (Aix Marseille University, CNRS, IRD, IRSTEA, OSU Institut Pythéas,

Marseille, France) for the *Posidonia oceanica* samples collection. *Z. marina* and *Z. Noltii*

were gifted by Pascal Bazile (8 Vie Pour la Planète, Projet Zorro) and *C. nodosa* by Heike

Molinaar (Université Côte d'Azur).

The electron microscopy experiments were performed on the PiCSL-FBI core facility

(Nicolas BROUILLY, Fabrice RICHARD and Aïcha AOUANE, IBDM, AMU-Marseille),

member of the France-BioImaging national research infrastructure (ANR-10-INBS-04). A

special thanks to Aïcha Aouane for her expertise and help in the plant samples preparation for

electron microscopy. Thanks to Cécile Lecampion for the statistics analysis on the grana size

quantification. We thank Dr. Kastritis and Dr. Hamdi for their assistance with transmission

electron microscopy at Martin Luther University Halle-Wittenberg, Germany, and appreciate

their support in providing access to the microscopy facility. Thanks to Rainer Hienerwadel

for his expertise with the Joliot spectroscopy for ETR measurements and to Alexis Riche for

his help and advice for 77 K fluorescence experiments. Thanks to Stefano Caffarri for the

correction of an early version of the manuscript and general advice on biochemistry

techniques. We are grateful to EMBO who granted QCF with a Scientific Exchange Grant

(number 9348) and thanks to Alexey Amunts. Huge thanks to Laurie Casalot, Lea Sylvi,

Manon Bartoli, Corinne Valette and Sophie Guasco for providing the equipment necessary

for sample preparation. We are deeply grateful to Paul Hudson for providing a scientific

environment of excellence.

Funding

This research was supported by the Swedish Foundation for Strategic Research SSF (Grant

Number ARC19-0051).

The work at VU Amsterdam was supported by the EU project Capitalise and by the Dutch

Organization for Scientific Research (NWO) via a TOP grant (to RC). Additionally, this work

was partly supported by FCT - Fundação para a Ciência e a Tecnologia, I.P., through

MOSTMICRO-ITQB R&D Unit (DOI: 10.54499/UIDB/04612/2020; DOI:

10.54499/UIDP/04612/2020) and LS4FUTURE Associated Laboratory (DOI:

10.54499/LA/P/0087/2020) (Hold by DAS during the editing and revision of the MS).