Tissue-specific I-Smad mechanisms revealed by structure-function analysis in *Drosophila*

Ania M. Simoncek[1], Steven J. Sviridoff[1], Joshua N. Hays[1], Noah J. Graichen[1], Mikolaj J.

Sulkowski[1*]

[1]Biology Department, Southern Connecticut State University, New Haven, USA

*Corresponding author

Abstract

Inhibitory Smads (I-Smads) regulate TGF-β/BMP signaling through multiple distinct

mechanisms, but whether different tissues preferentially employ specific mechanisms remains

unknown. To address this question, we performed structure-function analyses of the *Drosophila*

I-Smad Dad and its vertebrate orthologs Smad6 and Smad7 in neural and wing tissues,

measuring in vivo outputs of BMP signaling. We identified a critical 24-amino acid putative

DNA-binding domain (DNABD) within the MH1 domain of the Drosophila I-Smad, Dad, that is

essential for inhibitory function in wing tissue but unessential in neural tissue. Structural

analyses revealed that ΔDNABD disrupts a beta hairpin structure homologous to R-Smad DNA-

binding regions. We also found that Dad requires an intact MH1 domain to disrupt wing

development, whereas either MH1 or MH2 can independently disrupt BMP signaling in motor

neurons. These findings support a model where Dad functions through MH1-mediated

transcriptional regulation in wing primordium, but through multiple mechanisms in neurons.

Comparative analysis revealed that vertebrate I-Smad orthologs also show tissue-specific activity

patterns, with structural predictions suggesting that Smad6 retains ancestral DNA-binding

capacity while Smad7 has evolved enhanced MH2-mediated functions. These results reveal

context-dependent mechanisms of I-Smads that further the understanding of TGF-β/BMP

pathway regulation.

Introduction

Transforming growth factor β (TGF-β) signaling is an ancient and evolutionarily conserved

pathway for intercellular communication that controls diverse life processes such as axis

formation, cell differentiation, and tissue homeostasis. Its pleiotropy necessitates precise intra-

and extracellular modulation to elicit the appropriate effects in target cells (Massagué 2012).

Dysregulation of TGF-β signaling underlies the pathophysiology of numerous disease states and

developmental disorders. For instance, as TGF-β inhibits mitosis and promotes apoptosis in

many tissues, loss of TGF-β signaling often engenders tumorigenesis. In later oncogenic stages,

however, cells become resistant to the apoptosis-inducing effects, and TGF-β promotes tumor

growth and invasiveness via dedifferentiation, angiogenesis, and immune suppression (Derynck

et al. 2001; Syed 2016; Chan et al. 2022). Thus, the ability of cells to receive and interpret

physiologically appropriate levels of TGF-β signaling is vital across metazoan life stages.

Inhibitory Smads (I-Smads) are intracellular proteins that ensure an appropriate response

to TGF-β signaling. They are transcriptional targets of the signaling pathway, functioning in a

negative-feedback mechanism (Miyazawa and Miyazono 2017). Dysfunction of I-Smads is
implicated in the pathogenesis of human diseases such as colorectal cancer, atrial fibrosis, and
inflammatory bowel disease (Broderick et al. 2007; Fortini et al. 2014; He et al. 2011; Tang et al.
2019). Although mammalian genomes encode two I-Smads (*Smad6* and *Smad7*), the *Drosophila*
genome contains only a single I-Smad encoding gene, *Daughters against Decapentaplegic* (*Dad*)
(Tsuneizumi et al. 1997; Massagué et al. 2005). This lack of genetic redundancy, combined with
powerful genetic tools and well-characterized phenotypes, make *Drosophila* an ideal system in
which to study the mechanisms of I-Smad function *in vivo*.
I-Smads belong to the Smad family of proteins, which transduce signals from TGF-β
receptors. The Smad family also includes R-Smads, which are the receptor-regulated Smads that
become activated by type I receptors, and Co-Smads (common-mediator Smads), which form
complexes with activated R-Smads to enable their nuclear translocation and transcriptional
activity (Schmierer and Hill 2007). Vertebrate R-Smads include Smad2 and Smad3, which
respond to TGF-β and activin signals, and Smad1, Smad5, and Smad8/9, which respond to BMP
signals (Massagué et al. 2005). Smads are characterized by the presence of two main functional
domains: the MH1 domain, which mediates DNA binding and nuclear import, and the MH2
domain, which facilitates protein-protein interactions, including receptor binding, Smad
oligomerization, and transcriptional activation. These two domains are connected by a variable
linker region that contains regulatory sites for post-translational modifications.
I-Smads share a similar domain organization with other Smads. The crystal structure of
SMAD7 reveals that the MH2 domain shares a high degree of structural similarity with the MH2
domains of other Smads (Murayama et al. 2020). However, there is currently no crystal structure
available for the I-Smad N-terminal region, and no consensus on whether it constitutes a true
MH1 domain. The two mammalian I-Smad paralogs, Smad6 and Smad7, are divergent both in
their structure and function. Smad7 inhibits both TGF-β and bone morphogenetic protein (BMP)
signaling, whereas Smad6 specifically inhibits BMP signaling (Hanyu et al. 2001; Goto et al.
2007). Smad7 is also more potent than Smad6 when repression is quantified via luciferase assay
(Hanyu et al. 2001). Previous studies, largely performed in vitro, indicate that Dad specifically
inhibits the BMP branch of TGF-β signaling, and this inhibition has been attributed to Dad
preventing Thickveins, a BMP type I receptor, from phosphorylating the R-Smad Mothers
against decapentaplegic (Mad) (Inoue et al. 1998; Kamiya et al. 2008; Li et al. 2017).
Intriguingly, I-Smads have been shown to antagonize TGF-β signaling by several distinct
mechanisms in addition to the abovementioned inhibition of type I receptors (Miyazawa and
Miyazono 2017). I-Smads can form complexes with R-Smads, thus preventing their association
with co-Smads (Hata and Chen 2016; Yan et al. 2016). I-Smads cooperate with E3 ubiquitin
ligases to target R-Smads and other TGF- β components for destruction (Zhu et al. 1999; Kavsak
et al. 2000; Ebisawa et al. 2001; Murakami et al. 2003; Morén et al. 2005). I-Smads can also
accumulate in the nucleus and directly regulate transcription to counter the effects of R-Smads
(Bai et al. 2000; Hanyu et al. 2001; Zhang et al. 2007). Given that I-Smads function through such
diverse mechanisms and considering that many of the previous studies were performed in vitro,

we sought to further explore the mechanisms of Dad and I-Smad orthologs by conducting
structure/function studies in different in vivo cellular contexts.
Using a series of transgenic animals, here we provide evidence that the human orthologs
of Dad, SMAD6 and SMAD7, have diverging functions in *Drosophila* wing and neural tissue.
We further demonstrate that Dad lacking a specific, putative DNA binding region of its MH1
domain is not sufficient to disrupt wing development yet is sufficient to inhibit BMP signaling in
neural tissue. By comparing protein structure predictions, we extend these results to hypothesize
that Smad6/Dad inhibit TGF-β signaling by a divergent mechanism from Smad7, and that this
mechanism necessitates the MH1 domain, possibly for DNA binding. We also find that in both
brain and wing tissue, Dad does not require palmitoylation at a C-terminal cysteine (C556)
previously shown to be required for inhibition in ovarian tissue (Li et al. 2017). Together these
findings shed light on the molecular mechanisms of TGF-β regulation and provide compelling
evidence for the evolution of distinct, tissue-specific functionality among I-Smad orthologs.
94
Results
Creation of a series of transgenic animals to investigate tissue-specific I-Smad functionality
To investigate the structural basis for alternative I-Smad mechanisms, we created a series of
transgenic *Drosophila* to express vertebrate I-Smad orthologs, Smad6 and Smad7, and targeted
alterations of the *Dad* gene (Figure 1A). In addition to full-length *Smad6* and *Smad7*, a
transgenic line expressing a shorter isoform of *Smad7* lacking the MH1 domain (*Smad7S*) was
developed. Dad truncations covering the MH1 and MH2 domains were constructed, along with
truncations deleting specific residues implicated in DNA binding and trimerization (Δ*DNABD*
and Δ*Trimer*). We also created a substitution replacing a C-terminal cysteine of Dad that is a
reported target of palmitoylation (Dad^{C556A}).
105
Phenotypic analysis of wing tissue following construct expression
To assess the potency of our gene products to inhibit BMP signaling in developing wing tissue,
constructs were expressed with *Ap-Gal4*, which is expressed primarily in the dorsal compartment
of the wing imaginal disc (O'Keefe et al., 1998, Butterworth & King, 1965). We found that
phenotypes clustered into three groups based on severity. Group 1, which includes genetic
background control, ΔMH1, and ΔDNABD, had wing phenotypes indistinguishable from
controls. The mean wing areas (in mm^2) for males were 1.39±0.146, 1.50±0.089 and 1.53±0.124,
and for females were 1.96±0.163, 1.90±0.091, and 1.88±0.147, respectively. Group 2 includes
ΔMH2, Smad7S, Smad6, and ΔTrimer. This group had a moderately severe phenotypes, with
mean areas for males of 1.26±0.085, 1.05±0.143, 1.23±0.183, and 0.749±0.107. Group 2 females
wings were 1.60± 0.076, 1.20±0.198, 1.26±0.167, and 0.710±0.163, respectively. Finally, group
3, which includes Dad, Smad7L, and Dad^{C556A}, had the most severe phenotypes with mean areas
for males of 0.214±0.045, 0.219±0.079, and 0.262±0.047, and for females of 0.304±0.076,
0.258±0.112, and 0.280±0.063, respectively (Figure 1B).

Based upon these results, we conclude that Dad lacking its MH1 domain is insufficient to inhibit wing tissue growth. Dad with a 24 aa deletion in a putative DNA binding site (ΔDNA-BD) within the MH1 domain is also insufficient. Conversely, Smad7 lacking the MH1 domain is sufficient to inhibit wing growth. Even with a substitution converting the cysteine at position 556 to alanine (C556A), Dad remained effective in inhibiting wing growth. These results indicate that Dad must include MH1 domain to disrupt wing growth, while the vertebrate counterpart Smad7 can disrupt growth without the MH1 domain. Furthermore, Dad lacking the MH2 domain is still capable of producing some disruption in wing development.

Figure 1: Disruption of *Drosophila* wing development by I-Smad structure/function constructs and vertebrate orthologs. (A) Protein domain organization of I-Smad constructs. Scale indicates length in amino acids. Locations of putative DNA binding sites (blue) and putative trimer interface (red) are indicated. (B) Quantification of wing area (mm^2) in adult flies expressing I-Smad constructs in wing primordium with *Ap-Gal4* and genetic background controls (*Ap-Gal4/+*). Data points for individual female or male wings are represented by ♀ or ♂ symbols, respectively. Bars represent mean ± SD. Statistical comparisons performed using Dunnett's test comparing individual genotypes with controls. **** $p < 0.0001$, ** $p < 0.01$

Structural analysis of MH1 domains
As the MH1 domain of Dad was sufficient to disrupt wing development, we sought to identify
potential differences in protein structures among the MH1 domains of I-Smad orthologs. We first
generated a model of the MH1 domain of the R-Smad, Smad3 using AlphaFold 3 (Figure 2A).
The model matched the previously published crystal structure (Shi et al. 1998). Notably, the β-
hairpin structure and its interaction with DNA were correctly replicated. When we compared
predictions of the MH1 structures of Dad, Smad6, and Smad7 we found that the β-hairpin
structure is longer and more disorganized in Smad7 compared with Dad or Smad6, which had β-
hairpin structures with antiparallel stretches that more closely resemble the DNA binding
structure in Smad3 (Figure 2B).
150
Effects of human I-Smad orthologs in neural tissue
To gauge the ability of specific I-Smad structural regions to inhibit BMP signaling in motor
neurons, we expressed constructs using *OK371-Gal4*, a Gal4 enhancer trap in the proximity of
the *DVGLUT* gene, which drives expression specifically in glutamatergic motor neurons (Mahr
and Aberle 2006; Sanyal 2009). We began by determining the ability of the vertebrate I-Smads,
Smad6 and Smad7, to inhibit BMP signaling in this neural tissue. BMP signaling was quantified
by visualizing levels of phosphorylated Mad (pMad) in motor nuclei. Smad6, Smad7L, and
Smad7S decreased nuclear pMad relative to genetic background control (mean±s.e.m.:
0.650±0.0.098, 0.315±0.040, and 0.495±0.061, respectively) (Figure 2). To evaluate a
downstream effector of BMP signaling, we utilized a GFP protein trap in the BMP target gene
twit (Sulkowski et al., 2016; Kim & Marquez, 2012; Venken et al., 2011). Smad7L decreased
Twit-GFP levels to 0.519±0.0.048 relative to control. Interestingly, Smad6 and Smad7S did not
cause a significant reduction in Twit-GFP levels (0.897±0.070 and 0.792±0.072 relative to
control, respectively). Thus, vertebrate I-Smad orthologs were sufficient in inhibiting BMP
signaling in neural as well as wing disc tissues.

166
Figure 2: Human I-Smads inhibit BMP signaling in *Drosophila* tissues. Transgenic control (*TwitGFP,*
OK371-Gal4/+) compared with *OK371-Gal4*-driven overexpression of *UAS-Dad*, *UAS-Smad6*, *UAS-*
Smad7L, and *UAS-Smad7S*. (A) Motor neurons in ventral nerve cord triple-labeled for Elav (blue), Twit-
GFP (green) and pMad (red). Scale bar represents 20 microns. (B) Quantification of pMad and Twit-GFP
intensity normalized to genetic background control. Bars represent mean ± SEM. Each dot represents
mean data from motor neurons of one CNS. Statistical comparisons performed using Dunn's test with
Bonferroni correction. **** p<0.0001, ** p<0.01.
174
Regulation of non-canonical BMP signaling at neuromuscular synapses by I-Smads
BMP signaling at the neuromuscular junction (NMJ) encompasses both the canonical retrograde
pathway, which culminates with pMad accumulation in the nucleus (Aberle et al. 2002; Marqués
et al. 2002; McCabe et al. 2003), and a non-canonical pathway, characterized by pMad
accumulation at synaptic active zones (Dudu et al. 2006; Higashi-Kovtun et al. 2010; O'Connor-
Giles et al. 2008; Smith et al. 2012; Sulkowski et al. 2014, 2016). To test whether Dad affects the
noncanonical BMP pathway, we examined synaptic pMad levels in animals overexpressing Dad
in motor neurons. We saw a significant reduction in pMad accumulation at NMJ synapses in
animals overexpressing Dad in motor neurons relative to controls (0.313±0.043; p<0.005). We
also observed defects in NMJ morphology (bouton number and size) reminiscent of BMP
pathway perturbations (data not shown). Therefore, I-Smads can inhibit both canonical and
noncanonical BMP signaling at the NMJ.

Figure 3: Presynaptic Dad expression inhibits local BMP signaling at NMJ synapses. Transgenic control (*TwitGFP, OK371-Gal4/+*) compared with *OK371-Gal4*-driven overexpression of *UAS-Dad.* (A) Muscle 4 NMJ triple-labeled for the presynaptic membrane marker HRP (blue), active zone marker Brp (green), and synaptic pMad (red). Scale bar represents 10 microns. (B) Quantification of synaptic pMad intensity. Each dot represents one data point, which is the mean of one animal's NMJ 4s from A2 and A3 normalized to control (4 NMJs per data point). Bars represent mean ± SD. Statistical comparison performed with Student's t test. *** $p<0.005$.

I-Smad MH1 and MH2 domain truncations in neural tissue

To evaluate the necessity of I-Smad structural domains to inhibit BMP signaling in motor neurons, expression of *Dad* constructs with truncations of the MH1 and MH2 domains (ΔMH1 and ΔMH2) were driven with *OK371-Gal4* (Figure 4). Both ΔMH1 and ΔMH2 were sufficient to suppress nuclear pMad accumulation (0.201±0.015 and 0.396±0.058 relative to control, respectively). ΔMH1 expression significantly reduced Twit-GFP reporter gene expression to 0.525±0.047 relative to control. ΔMH2 reduced Twit-GFP to 0.734±0.083, but this reduction was not statistically significant. This analysis reveals that the functional domains of *Dad* can independently inhibit BMP signaling in neural tissue, with the mechanisms utilizing the MH2 domain appearing to be slightly more effective in motor neurons. This contrasts with the results of expression in wing primordium tissue, where there was no difference observed when ΔMH1 was expressed and only a small reduction in area when ΔMH2 was expressed (Figure 4C).

Figure 4: Dad can repress BMP signaling in Brains through multiple mechanisms. Transgenic
control (*TwitGFP, OK371-Gal4/+*) compared with OK371-Gal4 driven overexpression of UAS-ΔMH1
and UAS-ΔMH2 (A) Motor neurons in ventral nerve cord triple-labeled for Elav (blue), Twit-GFP (green)
and pMad (red). Scale bar represents 20 microns. (B) Quantification of pMad and Twit-GFP intensity.
Bars represent mean ± SEM. (C) Representative images of wing phenotypes for indicated genotypes.
Scale bar represents 1mm. Statistical comparisons performed using Dunn's test with Bonferroni
correction. **** $p<0.0001$, * $p<0.05$.

Deletion of putative DNA binding sites and trimerization interfaces
To determine whether specific regions of the I-Smad MH1 and MH2 domains are sufficient to
inhibit BMP signaling in motor neurons, we drove the expression of *Dad* constructs with
deletions in putative DNA binding sites (ΔDNABD) or trimer interfaces (ΔTrimer) with OK371-
Gal4 (Figure 5). Structural predictions show that the ΔDNABD mutation removes the beta-
hairpin structure implicated in DNA binding (Figure 5A). Both ΔDNABD and ΔTrimer
significantly reduced the accumulation of nuclear pMad relative to control (0.292±0.0276 and
0.341±0.0357, respectively). While ΔTrimer caused a significant decrease in Twit-GFP reporter
gene levels (0.656±0.062), ΔDNABD expression did not significantly change Twit-GFP
expression (0.849±0.0457). Therefore, Dad lacking putative DNA binding sites within the MH1
domain was not sufficient for repressing this target gene's expression. However, this same
structure is sufficient for decreasing pMad accumulation, indicating that these processes are
separable and can be attributed to different structural motifs. As the ΔDNABD construct was not

sufficient for altering wing morphology (Figures 1B and 5D), it appears that different tissue
types preferentially employ different I-Smad mechanisms.
233

234
Figure 5: Dad lacking putative DNA binding region is sufficient for repressing pMad accumulation
and Twit-GFP levels in brains, but not for repressing wing development. (A) AlphaFold3 structure
prediction of Dad and Dad$^{\Delta DNABD}$ MH1 domains. Putative Dad DNA binding domain is depicted in yellow
(absent in Dad$^{\Delta DNABD}$). (B-C) Transgenic control (*TwitGFP, OK371-Gal4/+*) compared with *OK371-*
Gal4-driven overexpression of *UAS-Dad$^{\Delta Trimer}$* and *UAS-Dad$^{\Delta DNABD}$*. (B) Motor neurons in ventral nerve
cord triple-labeled for Elav (blue), Twit-GFP (green) and pMad (red). Scale bar represents 20 microns.
(C) Quantification of pMad and Twit-GFP intensity normalized to genetic background control. Bars
represent mean ± SEM. Statistical comparisons performed using Dunn's test with Bonferroni correction

(pMad) or Tukey HSD test (TwitGFP). *** p<0.001, ** p<0.01, * p<0.05. (D) Representative images of

wing phenotypes for indicated genotypes. Scale bar represents 1 mm.

Removal of C-terminal palmitoylation site

Previous studies identified a conserved C-terminal cysteine (C556) of Dad that serves as a

palmitoylation site required for membrane association and BMP signaling inhibition (Li et al.

2017). We generated a *Dad* construct containing a substitution of C556 with alanine (Dad^{C556A}).

This variant was expressed using *Ap-Gal4* in the wing and *OK371-Gal4* in the brain to assess

tissue-specific effects on BMP signaling. In wing tissue, expression of Dad^{C556A} produced a

strong phenotype that placed it into the most severe category along with Dad (Figure 1B). This

finding suggests that C556 is not essential for I-Smad function in wing tissue, contrary to its

function based on ovarian studies, where this same modification reduced effectiveness to levels

similar to genetic background control (Li et al. 2017). Expression of Dad^{C556A} in motor neurons

resulted in a dramatic reduction in both pMad and TwitGFP levels (0.275±0.0319 and

0.395±0.0469 relative to control, respectively). These results indicate that C556-dependent

palmitoylation is not universally required for Dad function, and that this process may be tissue

specific, rather than universally applicable to I-Smad functionality.

Figure 6: Dad functions normally without cysteine 556 in wing and brain. Transgenic control
(*TwitGFP, OK371-Gal4/+*) compared with *OK371-Gal4*-driven overexpression of *UAS-Dad^{C556A}*. (A)
Motor neurons in ventral nerve cord triple-labeled for Elav (blue), Twit-GFP (green) and pMad (red).
Scale bar represents 20 microns. (B) Quantification of pMad and Twit-GFP intensity normalized to
genetic background control. Bars represent mean ± SEM. Statistical comparisons performed using Dunn's
test with Bonferroni correction. **** p<0.001. (C) Representative images of wing phenotypes for
indicated genotypes. Scale bar represents 1 mm.

Structural predictions of physical interactions between Dad and target molecules

Our results support a model where I-Smads inhibit TGF-β /BMP signaling through several
distinct mechanisms, and different tissues preferentially employ these different mechanisms. As
we used structural prediction to compare a putative DNA binding motifs of R-Smads and I-
Smads (Figures 2A and 5A), we wanted to test whether structural predictions can also visualize
the physical interactions between I-Smads and the other target proteins, R-Smads and Type-I
receptors. Indeed, AlphaFold3 models support a strong, complementary conformation between
Dad and the R-Smad Mad and between Dad and the Type-I receptor Thickveins (Tkv) (Figure
7). The I-Smad/R-Smad conformation with the closest fit was between two I-Smads and one R-
Smad (Figure 7A). There were five C-terminal residues of Dad that formed hydrogen bonds with

the MH2 domain of Mad: M520, H523, G524, M528, and R568. Structural predictions most strongly support an association between a single I-Smad and Type-I receptor (Figure 7B). We identified nine C-terminal residues of Dad that formed hydrogen bonds with the cytoplasmic region of Tkv: T406, W406, D487, L495, M528, P530, T544, I553, E561. In agreement with prior knowledge, these residues are all found within the C-terminal MH2 domain of the I-Smad.

Figure 7: Structural predictions support multiple modes of I-Smad function. (A) AlphaFold3 predicted protein interaction between two Dad molecules (I-Smad) and one Mad (R-Smad). Predicted hydrogen bonding residues are labeled. (B) AlphaFold3 predicted protein interaction between one Dad (I-Smad) and one Thickveins (Type I receptor). Predicted hydrogen bonding residues are labeled. (C) I-Smads function through at least three distinct mechanisms: 1. Preventing receptors from phosphorylating

292   R-Smads, 2. Binding to R-Smads and preventing formation of R-Smad/Co-Smad complex, and 3.
293   Interacting with DNA to counteract TGF-β/BMP transcriptional effects. In neural tissue, I-Smads
294   function preferentially through either the first or second mechanisms. In wing tissue, I-Smads rely on the
295   third mechanism.
296
297

Discussion
Structural basis of I-Smad tissue-specific mechanisms
The modes by which I-Smads have been shown to inhibit TGF-β/BMP signaling can be
classified into three distinct mechanisms (Figure 7C): 1. Interfering with Type I receptors, either
by direct physical interaction, or by recruitment of effector molecules such as BAMBI, E3
ubiquitin ligases (Smurf1 and Smurf2), or GADD34; 2. Preventing R-Smad/Co-Smad complex
formation both by direct physical interaction and promoting ubiquitination; 3. Directly regulating
transcription to counteract expression changes of TGF-β/BMP target genes (Miyazawa and
Miyazono 2017). Our current study supports the hypothesis that certain tissue types utilize
specific I-Smad mechanisms preferentially over others. Confirming previous results, we found
that expression of the *Drosophila* I-Smad, Dad, in wing primordium results in abnormally small,
shriveled wings (Figure 1B) (Inoue et al. 1998). Seeking to determine which parts of Dad are
sufficient for this phenotype, we identified a 24 aa deletion in a putative DNA binding region
within the MH1 domain (ΔDNABD) (Figure 1A & 1B). Protein structure prediction reveals a
complete disruption of the beta hairpin structure that resembles the R-Smad structure which
closely contacts the DNA molecule (Figures 2A & 4A).
ΔDNABD expressed in motor neurons disrupted BMP signaling in this neural tissue
(Figure 5), indicating that an alternative mechanism is used in this tissue compared to wing disc.
Indeed, Dad with only an MH1 or an MH2 domain (ΔMH1 and ΔMH2) disrupted BMP signaling
in neural tissue (Figure 4). However, Dad with only MH2 domain (ΔMH1), was not sufficient to
disrupt wing growth (Figures 1B and 4C). Based on these results, we propose that Dad functions
primarily through MH1-mediated direct transcriptional regulation in wing tissue, but through
multiple mechanisms in neural tissue (Figure 7C). To support this model, we analyzed predicted
protein structures and observed three characteristics: 1. The DNA binding loop of Dad shares a
similar structure with that of R-Smads (Figure 2A); 2. The structure of I-Smad associating with
the Type-I receptor reveals binding between specific residues; and 3. The structure of I-Smads
associating with R-Smads also reveals interaction sites (Figure 7A,B).
These alternative mechanisms may hold the key to maintaining appropriate levels of
TGF-β/BMP signaling in different tissue types. For example, we found that expression of Dad
with a cysteine to alanine substitution at position 556 (C556A) was fully effective at disrupting
both wing development and BMP signaling outputs in motor neurons (Figures 1B & 6). Previous
work shows that this mutation abolishes the ability of Dad to cause loss of germ cells in ovarian
tissue (Li et al. 2017). As this residue is reported to serve as a palmitoylation site that promotes
association with receptors on the plasma membrane, it appears that Dad functions independently

of this membrane association in wing and brain tissue. Further experiments such as genetic
epistasis will be required to determine the exact mechanism(s) used in these tissue types.
334

Multiple functions performed by distinct structural regions in other proteins
The fact that a single protein can utilize different structural regions to work through such
disparate interactions is remarkable but is not without precedent. Trio, a large, multi-domain
protein, exemplifies this versatility. The N-terminal GEF1, central GEF2, and C-terminal protein
kinase domains each catalyze distinct reactions that affect different aspects of cellular physiology
(Bircher and Koleske 2021). While GEF1 activates Rac1 to promote membrane ruffling and cell
migration, GEF2 stimulates RhoA to induce stress fiber formation, with the protein kinase
domain providing additional signaling capabilities. Another example is the extracellular protein
Cerberus, which uses distinct mechanisms to inhibit three different pathways: directly binding to
Wnt proteins in the extracellular space, preventing Nodal from interacting with its receptors, and
antagonizing BMP signaling through direct protein interaction (Piccolo et al. 1999; Kawano and
Kypta 2003; Tavares et al. 2007; Aykul et al. 2015). Thus, proteins can utilize different structural
motifs to carry out distinct interactions, although further research is necessary to determine
whether these different mechanisms are used preferentially depending on the cellular context.
349

Switching I-Smad mechanisms based on intracellular environment
It is likely that the complement of molecules within a cell determines which mechanism an I-
Smad will work through in that cell. Recent works have begun to elucidate the identities of I-
Smad partner molecules. For example, Liu et al. recently identified a gene encoding part of the
Integrator complex in a screen for genes required for female germline maintenance (Liu et al.
2023). In germline tissue and S2 cells, the Integrator complex is required for Dad to degrade
Tkv. It is possible that certain Integrator complex components are absent from wing discs, and
thus Dad functions through an alternative mechanism. Testing this hypothesis may prove
challenging as there are eight genes encoding Integrator complex components (Hao et al. 2008).
Under the assumption that I-Smads evolved from an ancestral Smad gene that encoded a
protein with DNA binding properties, we favor a model wherein *Smad6* and *Smad7* began
differentiating after their duplication in the vertebrate lineage. Considering the higher percent
sequence identity between Dad and Smad6, as well as the specificity of both molecules for the
BMP branch of TGF- β signaling (Goto et al. 2007), it can be inferred that Smad6 retains more
of the ancestral function shared with Dad. Therefore, Smad7 may have evolved additional
mechanisms, for example an increased effectiveness of the MH2-mediated mechanisms (Figure
1B)(Hanyu et al. 2001), and a reduction of the MH1-mediated mechanisms. The structural
similarity of the DNA binding loop between R-Smads, Smad6, and Dad, but not Smad7 (Figure
2A) supports this hypothesis.
369
370
371

Applications and future directions
These findings indicate that any potential diagnosis or treatment involving I-Smads must
consider the mechanism(s) employed by the target tissue type. We demonstrate that to inhibit
growth of wing tissue, the *Drosophila* I-Smad functions through an MH1-mediated modality,
most likely involving DNA binding. However, future work is necessary to test the ramifications
and limits of this phenomenon. It is vital to determine which mechanisms work in additional
tissue types, such as intestinal epithelium, and whether the mechanisms change in pathological
states like colorectal cancer. It will also be interesting to further explore the evolutionary
relationships between Dad, Smad6, and Smad7 to learn more about their common and divergent
functionality. Our versatile in vivo platform can be used to conduct further structure/function
analyses of I-Smad orthologs, such as deletion of the homologous Smad6 DNA binding region
and domain swap experiments. Ultimately, extension of these findings to vertebrate systems will
be required to test their broader applicability to modulating TGF-β/BMP signaling in diverse
cellular contexts.
386
Methods
Molecular biology and transgenesis
Plasmids were created using standard molecular biology methods. All plasmid sequences were
verified using whole plasmid nanopore sequencing (Eurofins Genomics; Louisville, KY, USA).
Injections were performed by Rainbow Transgenics, Inc. (Camarillo, CA, USA). All constructs
were inserted into the VK00033 locus on the third chromosome. Insertions were verified by
sequencing. Complete lists of plasmids and primers used in this study are included below:

Plasmid	Creation	Source	Notes
pUAST-attB	N/A	DGRC Stock 1419 ; https://dgrc.bio.indiana.edu//stock/1419; RRID:DGRC_1419	Destination vector for transformation
pBSPPC	N/A	pBSPPc was a gift from Cuiqing Ma (Addgene plasmid # 116918 ; https://n2t.net/addgene:116918; RRID:Addgene_116918) (Xu et al. 2013)	Intermediate vector
CS2 Smad6	N/A	CS2 Smad6 was a gift from Joan Massague (Addgene plasmid # 14960 ; https://n2t.net/addgene:14960; RRID:Addgene_14960) (Hata et al. 1998)	Source of Smad6 CDS
TFORF0111	N/A	TFORF0111 was a gift from Feng Zhang (Addgene plasmid # 141518 ; https://n2t.net/addgene:141518; RRID:Addgene_141518) (Joung et al. 2023)	Source of Smad7S CDS
pBSPPc-Smad6	Insert from CS2 Smad6 cloned into pBSPPc using BamHI/XbaI sites	Current study	

pUAST-attB-Smad6	Insert from pBSPPc-Smad6 cloned into pUAST-attB using XhoI/XbaI sites	Current study	Transformation vector for Smad6
pBSPPc-NMC	Additional cloning sites added to pBSPPc by inserting oligonucleotide into PstI/BamHI sites	Current study	To accommodate Smad7S insert
pBSPPc-NMC-Smad7S	Insert from TFORF0111 cloned into pBSPPc-NMC using BmtI/SpeI sites	Current study	
pUAST-attB-Smad7S	Insert from pBSPPc-NMC-Smad7S (BamHI/XbaI) cloned into pUAST-attB using XhoI/XbaI sites	Current study	Transformation vector for Smad7S
LD47465	N/A	LD47465 (DGRC Stock 1333567 ; https://dgrc.bio.indiana.edu//stock/1333567 ; RRID:DGRC_1333567)	Source for Dad CDS
pUAST-attB-Dad	PCR product from LD47465 with XhoI/XbaI sites added cloned into pUAST-attB		Transformation vector for UAS-Dad
PCMV5-Smad7-HA	N/A	pCMV5-Smad7-HA was a gift from Jeff Wrana (Addgene plasmid # 11733 ; https://n2t.net/addgene:11733; RRID:Addgene_11733)(Hayashi et al. 1997)	Source for Smad7L CDS
PCMV5-Smad7	Deleted HA tag from pCMV5-Smad7-HA with PCR	Current study	
pUAST-attB-Smad7L	Insert from pCMV5-Smad7 cloned into pUAST-attB using EcoRI/XbaI sites	Current study	Transformation vector for UAS-Smad7L
pUAST-attB-Dad-ΔMH1	Deleted	Current study	Transformation vector for ΔMH1
pUAST-attB-Dad-ΔDNA-BD		Current study	Transformation vector for ΔDNABD
pUAST-attB-Dad-ΔMH2		Current study	Transformation vector for ΔMH2
pUAST-attB-Dad-ΔTrimer		Current study	Transformation vector for ΔTrimer

Primer/Oligonucleotide	Sequence (5'-3')	Notes
pBSPPc-NMC-sense	GCATGCTAGCTTAAGATCTG	Hybridized to create PstI/BamHI overhangs to clone into pBSPPc
pBSPPc-NMC-antisense	GATCCAGATCTTAAGCTAGCATGCTGCA	

pOT2-Dad-XhoI-F	GCTCGAGATGATATTCCCAAG AGAAAAGAAGGT	To obtain Dad CDS from LD47465 and add XhoI/XbaI sites
pOT2-Dad-XbaI-R	CTCTAGATCACCGCAGATGAC TAAAGTGA	
Smad7-HA-Del-F	TAATCTAGAGGATCCCG	To delete HA tag from pCMV5-Smad7-HA
Smad7-HA-Del-R	CCGGCTGTTGAAGATG	
MH1-Del-F	GAAACAGAATCCCCAAC	To delete MH1 domain from pUAST-attB-Dad
MH1-Del-R	TGGAATCGTAAGTGTGG	
DNA-BD-F	AAGGAACTGAAGCGAC	To delete putative DNA binding region from pUAST-attB-Dad
DNA-BD-R	GCACGGTATCAGGATG	
MH2-Del-F	CATCTGCGGTGATCTAG	To delete MH2 domain from pUAST-attB-Dad
MH2-Del-R	CACCTGACTGTTGATG	
Trimer-Del-F	GTGGCCAGTGAGGTG	To delete putative trimer interface region from pUAST-attB-Dad
Trimer-Del-R	GATATTGACTGCATTCGTTTT G	
C556A-F1	TATCATGGGCGCGCCCTGCTG GC	To create pUAST-attB-DadC556A two PCR products were combined with Gibson assembly.
C556A-R1	GTTGACCAGCTGCCGCCATC	
C556A-F2	GATGGCGGCAGCTGGTCAAC	
C556A-R2	GCCAGCAGGGCGCGCCCATG ATA	

Drosophila stocks and husbandry

Animals were reared at 25°C on standard cornmeal molasses medium. To control for crowding, crosses were set up with eight females crossed with six males, and adults were passed to a new vial every 3-4 days. For expression in motor neurons, *OK371-Gal4* (Mahr and Aberle 2006) was used. For expression in wing primordium, *Ap-Gal4* was used. A complete list of the strains used for this study is included below:

Abbreviation	**Genotype**	**Source**
w1118	*w[1118]*	BIN: 2376 (BDSC)
OK371	*w[1118]; P{w[+mW.hs]=GawB}VGlut1[OK371]*	BIN 26160 (BDSC)
Ap-Gal4	*w[1118]; P{y[+t7.7] w[+mC]=GMR42A06-GAL4}attP2*	BIN 41425 (BDSC)
Dad^{RNAi}	*y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.JF02133}attP2*	BIN 26235 (BDSC)
TwitGFP	*w[1118]; Mi{y[+mDint2]=MIC}twit[MI06552]*	BIN: 41449 (BDSC)
UAS-Dad	*w[1118]; +; P{UAS-Dad}VK00037*	Current study
UAS-Smad6	*w[1118]; +; P{UAS-SMAD6}VK00037*	Current study
UAS-Smad7	*w[1118]; +; P{UAS-SMAD7.1}VK00037*	Current study
UAS-Smad7S	*w[1118]; +; P{UAS-SMAD7.3}VK00037*	Current study
ΔMH1	*w[1118]; +; P{UAS-DadΔMH1}VK00037*	Current study
ΔMH2	*w[1118]; +; P{UAS-DadΔMH2}VK00037*	Current study
ΔDNABD	*w[1118]; +; P{UAS-DadΔDNABD}VK00037*	Current study
ΔTrimer	*w[1118]; +; P{UAS-DadΔTrimer}VK00037*	Current study

Dad^{C556A}	$w[1118]; +; P\{UAS\text{-}Dad^{C556A}\}VK00037$	Current study

Immunostaining and microscopy

Nervous system imaging and immunostaining were performed as described (Brown et al. 2018). Briefly, wandering third instar larvae were dissected in hemolymph-like-3.1 (HL3.1) saline (Feng et al. 2004), fixed in 4% paraformaldehyde diluted with 1XPBS for 20 minutes, and washed several times in wash buffer (1XPBS with 0.3% TX-100). Primary antibodies were used at the following concentrations: rabbit anti-pSmad1/5 (Cell Signaling Technology Cat# 9516, RRID:AB_491015) at 1:500, rabbit anti-pMad PS1 (Carl-Henrik Heldin Göteborg University, Sweden Cat# pMad, RRID:AB_2617125) at 1:500, chicken anti-GFP (Abcam Cat#ab13960, RRID: AB_300798) at 1:1000, and rat anti-Elav (DSHB Cat# 7E8A10, RRID: AB_528218) at 1: 500. Secondary antibodies were used at the following concentrations: goat anti-chicken Alexa Fluor 488 (Thermo Fisher Scientific Cat# A-11039, RRID:AB_2534096) at 1:500, goat anti-rabbit Alexa Fluor 568 (ThermoFisher Scientific Cat# A-11011, RRID: AB_143157) at 1:500, goat-anti-rat Alexa Fluor 647 (Thermo Fisher Scientific Cat #A-21247, RRID: AB_141778), and goat anti-HRP Alexa Fluor 488 (Jackson ImmunoResearch Labs Cat# 123-545-021, RRID:AB_2338965). Samples were processed in parallel such that all incubation times were identical. Brains were mounted in SlowFade Diamond (ThermoFisher Cat#S36972) and filets were mounted in Prolong Diamond (ThermoFisher Cat# P36961). Images were acquired with a Zeiss LSM 880 or Nikon Ti-2 confocal microscope using a 20X or 63X objective. Images were acquired and analyzed while blinded to genotype.

Wing dissection, imaging, and analysis

Newly eclosed (between 8-24 hrs.) adults were immediately fixed in 100% ethanol. Wings were dissected and transferred to 70% ethanol to rehydrate. Wings were then arranged on a microscope slide and covered with ProLong Diamond mounting medium (ThermoFisher Scientific Cat# P36965). A coverslip was placed on top, and the slide was allowed to cure overnight covered with a small weight to flatten the preparation. Edges were sealed with nail polish. Slides were imaged with compound microscope using 20X objective. Area was determined by manually tracing the border of the wing image using the polygon tool in ImageJ and quantifying area of the resulting polygon. Images were acquired and analyzed while blinded to genotype.

Structural prediction and analyses

Full-length protein sequences were obtained from the National Center for Biotechnology Information (NCBI) protein database (https://www.ncbi.nlm.nih.gov/protein/) and verified for completeness and accuracy. Complete amino acid sequences were submitted to AlphaFold3 server (https://alphafoldserver.com/) using default parameters. Predicted structures were downloaded from AlphaFold3 in PBD format. Structural visualization was performed using UCSF ChimeraX (Meng et al. 2023). Confidence scores provided by AlphaFold3 were used to

441 assess the reliability of predicted structural regions, with high-confidence regions (pLDDT > 70)
442 considered reliable for structural interpretation.
443
Data analysis and statistics
Pixel intensity was measured from maximum projection confocal micrographs using a custom
ImageJ script (AIABS) as previously described (Brown et al. 2019). Intensity measurements
were performed while blinded to genotype. All data were subjected to outlier analysis with the Z-
test method with threshold ≥ 2 STDEV from the mean. Normality of data points for each
genotype was tested using a Shapiro-Wilk test. Similarity of variances was tested using Levene's
test. For data sets with multiple genotypes where measurements from each genotype were
normally distributed and variances were similar, differences across genotypes were tested by
one-way ANOVA. As a post-hoc test, when samples were compared only to control, Dunnett's
test was performed. When all samples were compared to each other, Tukey's test was conducted.
For data that were not normally distributed or had unequal variances, differences across
genotypes were tested by the Kruskal-Wallis test. Pairwise differences were tested by post-hoc
Dunn's test with Bonferroni correction. For comparing two groups with normal distributions,
Student's t-test was used. Data were analyzed with IBM SPSS Statistics (Version 27) and R
(Version 4.3.2; R Core Team 2023). Graphs were created with the *ggplot2* R package (Wickham
et al. 2016). Scripts used for statistical analysis and plot creation available upon request.
460

Acknowledgements
This work was supported by The National Institute of Neurological Disorders and Stroke of The
National Institutes of Health under Award Number R15NS116630 to M.J.S. We thank the
Drosophila Genomics Resource Center (NIH Grant 2P40OD010949) for plasmids. We thank the
Developmental Studies Hybridoma Bank, created by the NICHD of the NIH and maintained at
The University of Iowa, Department of Biology, Iowa City, IA 52242, for antibodies. Molecular
graphics and analyses were performed with UCSF ChimeraX, developed by the Resource for
Biocomputing, Visualization, and Informatics at the University of California, San Francisco,
with support from National Institutes of Health R01-GM129325 and the Office of Cyber
Infrastructure and Computational Biology, National Institute of Allergy and Infectious Diseases.
We thank Stuart Newfeld and Gregory Reeves for sharing *Drosophila* stocks. Thanks to Daniel
Cox, Sean Grace, Nicholas Edgington, and Rachel Jeffrey for feedback and discussions. Thanks
to members of the Sulkowski lab for proofreading and editing the manuscript.
474

475 **Author contributions**
476 MJS and AMS conceived of the study. MJS, SJS, AMS, JNH, and NJG created transformation
477 vectors. AMS, SJS and NJG conducted experiments and analyzed data. NJG conducted structural
478 analysis. JNH isolated and sequenced genomic DNA from transformants. All authors contributed
479 to writing the manuscript.
480

Literature Cited

481
Aberle H, Haghighi AP, Fetter RD, McCabe BD, Magalhães TR, Goodman CS. 2002. wishful
thinking encodes a BMP type II receptor that regulates synaptic growth in
Drosophila. *Neuron* **33**: 545–558.

Aykul S, Ni W, Mutatu W, Martinez-Hackert E. 2015. Human Cerberus Prevents Nodal-
Receptor Binding, Inhibits Nodal Signaling, and Suppresses Nodal-Mediated
Phenotypes. *PLOS ONE* **10**: e0114954.

Bai S, Shi X, Yang X, Cao X. 2000. Smad6 as a transcriptional corepressor. *J Biol Chem* **275**:
8267–8270.

Bircher JE, Koleske AJ. 2021. Trio family proteins as regulators of cell migration and
morphogenesis in development and disease - mechanisms and cellular contexts. *J*
Cell Sci **134**: jcs248393.

Broderick P, Carvajal-Carmona L, Pittman AM, Webb E, Howarth K, Rowan A, Lubbe S,
Spain S, Sullivan K, Fielding S, et al. 2007. A genome-wide association study shows
that common alleles of SMAD7 influence colorectal cancer risk. *Nat Genet* **39**:
1315–1317.

Brown JR, Phongthachit C, Sulkowski MJ. 2019. Immunofluorescence and image analysis
pipeline for Drosophila motor neurons. *Biol Methods Protoc* **4**: bpz010.

Chan MK-K, Chung JY-F, Tang PC-T, Chan AS-W, Ho JY-Y, Lin TP-T, Chen J, Leung K-T, To K-
F, Lan H-Y, et al. 2022. TGF-β signaling networks in the tumor microenvironment.
Cancer Lett **550**: 215925.

Derynck R, Akhurst RJ, Balmain A. 2001. TGF-beta signaling in tumor suppression and
cancer progression. *Nat Genet* **29**: 117–129.

Dudu V, Bittig T, Entchev E, Kicheva A, Jülicher F, González-Gaitán M. 2006. Postsynaptic
Mad Signaling at the Drosophila Neuromuscular Junction. *Curr Biol* **16**: 625–635.

Ebisawa T, Fukuchi M, Murakami G, Chiba T, Tanaka K, Imamura T, Miyazono K. 2001.
Smurf1 interacts with transforming growth factor-beta type I receptor through
Smad7 and induces receptor degradation. *J Biol Chem* **276**: 12477–12480.

Fortini BK, Tring S, Plummer SJ, Edlund CK, Moreno V, Bresalier RS, Barry EL, Church TR,
Figueiredo JC, Casey G. 2014. Multiple Functional Risk Variants in a SMAD7
Enhancer Implicate a Colorectal Cancer Risk Haplotype. *PLoS ONE* **9**: e111914.

Goto K, Kamiya Y, Imamura T, Miyazono K, Miyazawa K. 2007. Selective inhibitory effects of
Smad6 on bone morphogenetic protein type I receptors. *J Biol Chem* **282**: 20603–
20611.

Hanyu A, Ishidou Y, Ebisawa T, Shimanuki T, Imamura T, Miyazono K. 2001. The N domain
of Smad7 is essential for specific inhibition of transforming growth factor-β
signaling. *J Cell Biol* **155**: 1017–1028.

Hao R, Chen L, Wu J-W, Wang Z-X. 2008. Structure of Drosophila Mad MH2 domain. *Acta*
Crystallograph Sect F Struct Biol Cryst Commun **64**: 986–990.

Hata A, Chen Y-G. 2016. TGF-β Signaling from Receptors to Smads. *Cold Spring Harb*
Perspect Biol **8**. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5008074/
(Accessed March 10, 2023).

Hata A, Lagna G, Massagué J, Hemmati-Brivanlou A. 1998. Smad6 inhibits BMP/Smad1
signaling by specifically competing with the Smad4 tumor suppressor. *Genes Dev*
12: 186–197.

Hayashi H, Abdollah S, Qiu Y, Cai J, Xu YY, Grinnell BW, Richardson MA, Topper JN,
Gimbrone MA, Wrana JL, et al. 1997. The MAD-related protein Smad7 associates
with the TGFbeta receptor and functions as an antagonist of TGFbeta signaling. *Cell*
89: 1165–1173.

He X, Gao X, Peng L, Wang S, Zhu Y, Ma H, Lin J, Duan DD. 2011. Atrial fibrillation induces
myocardial fibrosis through angiotensin II type 1 receptor-specific Arkadia-
mediated downregulation of Smad7. *Circ Res* **108**: 164–175.

Higashi-Kovtun ME, Mosca TJ, Dickman DK, Meinertzhagen IA, Schwarz TL. 2010. Importin-
β11 Regulates Synaptic Phosphorylated Mothers Against Decapentaplegic, and
Thereby Influences Synaptic Development and Function at the Drosophila
Neuromuscular Junction. *J Neurosci* **30**: 5253–5268.

Inoue H, Imamura T, Ishidou Y, Takase M, Udagawa Y, Oka Y, Tsuneizumi K, Tabata T,
Miyazono K, Kawabata M. 1998. Interplay of Signal Mediators of Decapentaplegic
(Dpp): Molecular Characterization of Mothers against dpp, Medea, and Daughters
against dpp. *Mol Biol Cell* **9**: 2145–2156.

Joung J, Ma S, Tay T, Geiger-Schuller KR, Kirchgatterer PC, Verdine VK, Guo B, Arias-Garcia
MA, Allen WE, Singh A, et al. 2023. A transcription factor atlas of directed
differentiation. *Cell* **186**: 209-229.e26.

Kamiya Y, Miyazono K, Miyazawa K. 2008. Specificity of the inhibitory effects of Dad on
TGF-beta family type I receptors, Thickveins, Saxophone, and Baboon in
Drosophila. *FEBS Lett* **582**: 2496–2500.

Kavsak P, Rasmussen RK, Causing CG, Bonni S, Zhu H, Thomsen GH, Wrana JL. 2000.
Smad7 binds to Smurf2 to form an E3 ubiquitin ligase that targets the TGF beta
receptor for degradation. *Mol Cell* **6**: 1365–1375.

Kawano Y, Kypta R. 2003. Secreted antagonists of the Wnt signalling pathway. *J Cell Sci*
116: 2627–2634.

Li W, Li W, Zou L, Ji S, Li C, Liu K, Zhang G, Sun Q, Xiao F, Chen D. 2017. Membrane
targeting of inhibitory Smads through palmitoylation controls TGF-β/BMP signaling.
Proc Natl Acad Sci U S A **114**: 13206–13211.

Liu S, Baeg GH, Yang Y, Goh FG, Bao H, Wagner EJ, Yang X, Cai Y. 2023. The Integrator
complex desensitizes cellular response to TGF-β/BMP signaling. *Cell Rep* **42**:
112007.

Mahr A, Aberle H. 2006. The expression pattern of the Drosophila vesicular glutamate
transporter: A marker protein for motoneurons and glutamatergic centers in the
brain. *Gene Expr Patterns* **6**: 299–309.

Marqués G, Bao H, Haerry TE, Shimell MJ, Duchek P, Zhang B, O'Connor MB. 2002. The
Drosophila BMP type II receptor Wishful Thinking regulates neuromuscular synapse
morphology and function. *Neuron* **33**: 529–543.

Massagué J. 2012. TGFβ signalling in context. *Nat Rev Mol Cell Biol* **13**: 616–630.

Massagué J, Seoane J, Wotton D. 2005. Smad transcription factors. *Genes Dev* **19**: 2783–
2810.

McCabe BD, Marqués G, Haghighi AP, Fetter RD, Crotty ML, Haerry TE, Goodman CS,
O'Connor MB. 2003. The BMP homolog Gbb provides a retrograde signal that
regulates synaptic growth at the Drosophila neuromuscular junction. *Neuron* **39**:
241–254.

Meng EC, Goddard TD, Pettersen EF, Couch GS, Pearson ZJ, Morris JH, Ferrin TE. 2023.
UCSF ChimeraX: Tools for structure building and analysis. *Protein Sci* **32**: e4792.

Miyazawa K, Miyazono K. 2017. Regulation of TGF-β Family Signaling by Inhibitory Smads.
Cold Spring Harb Perspect Biol **9**: a022095.

Morén A, Imamura T, Miyazono K, Heldin C-H, Moustakas A. 2005. Degradation of the
tumor suppressor Smad4 by WW and HECT domain ubiquitin ligases. *J Biol Chem*
280: 22115–22123.

Murakami G, Watabe T, Takaoka K, Miyazono K, Imamura T. 2003. Cooperative Inhibition of
Bone Morphogenetic Protein Signaling by Smurf1 and Inhibitory Smads. *Mol Biol*
Cell **14**: 2809–2817.

O'Connor-Giles KM, Ho LL, Ganetzky B. 2008. Nervous Wreck Interacts with Thickveins
and the Endocytic Machinery to Attenuate Retrograde BMP Signaling during
Synaptic Growth. *Neuron* **58**: 507–518.

Piccolo S, Agius E, Leyns L, Bhattacharyya S, Grunz H, Bouwmeester T, De Robertis EM.
1999. The head inducer Cerberus is a multifunctional antagonist of Nodal, BMP and
Wnt signals. *Nature* **397**: 707–710.

Sanyal S. 2009. Genomic mapping and expression patterns of C380, OK6 and D42
enhancer trap lines in the larval nervous system of Drosophila. *Gene Expr Patterns
GEP* **9**: 371–380.

Schmierer B, Hill CS. 2007. TGFβ–SMAD signal transduction: molecular specificity and
functional flexibility. *Nat Rev Mol Cell Biol* **8**: 970–982.

Shi Y, Wang Y-F, Jayaraman L, Yang H, Massagué J, Pavletich NP. 1998. Crystal Structure of
a Smad MH1 Domain Bound to DNA: Insights on DNA Binding in TGF-β Signaling.
Cell **94**: 585–594.

Smith RB, Machamer JB, Kim NC, Hays TS, Marqués G. 2012. Relay of retrograde
synaptogenic signals through axonal transport of BMP receptors. *J Cell Sci* **125**:
3752–3764.

Sulkowski M, Kim Y-J, Serpe M. 2014. Postsynaptic glutamate receptors regulate local BMP
signaling at the Drosophila neuromuscular junction. *Dev Camb Engl* **141**: 436–447.

Sulkowski MJ, Han TH, Ott C, Wang Q, Verheyen EM, Lippincott-Schwartz J, Serpe M. 2016.
A Novel, Noncanonical BMP Pathway Modulates Synapse Maturation at the
Drosophila Neuromuscular Junction. *PLOS Genet* **12**: e1005810.

Syed V. 2016. TGF-β Signaling in Cancer. *J Cell Biochem* **117**: 1279–1287.

Tang Y, Reissig S, Glasmacher E, Regen T, Wanke F, Nikolaev A, Gerlach K, Popp V, Karram
K, Fantini MC, et al. 2019. Alternative Splice Forms of CYLD Mediate Ubiquitination
of SMAD7 to Prevent TGFB Signaling and Promote Colitis. *Gastroenterology* **156**:
692-707.e7.

Tavares AT, Andrade S, Silva AC, Belo JA. 2007. Cerberus is a feedback inhibitor of Nodal
asymmetric signaling in the chick embryo. *Development* **134**: 2051–2060.

Tsuneizumi K, Nakayama T, Kamoshida Y, Kornberg TB, Christian JL, Tabata T. 1997.
Daughters against dpp modulates dpp organizing activity in Drosophila wing
development. *Nature* **389**: 627–631.

Xu Y, Tao F, Ma C, Xu P. 2013. New constitutive vectors: useful genetic engineering tools
for biocatalysis. *Appl Environ Microbiol* **79**: 2836–2840.

Yan X, Liao H, Cheng M, Shi X, Lin X, Feng X-H, Chen Y-G. 2016. Smad7 Protein Interacts
with Receptor-regulated Smads (R-Smads) to Inhibit Transforming Growth Factor-β
(TGF-β)/Smad Signaling. *J Biol Chem* **291**: 382–392.

Zhang S, Fei T, Zhang L, Zhang R, Chen F, Ning Y, Han Y, Feng X-H, Meng A, Chen Y-G. 2007.
Smad7 Antagonizes Transforming Growth Factor β Signaling in the Nucleus by
Interfering with Functional Smad-DNA Complex Formation. *Mol Cell Biol* **27**: 4488–
4499.

Zhu H, Kavsak P, Abdollah S, Wrana JL, Thomsen GH. 1999. A SMAD ubiquitin ligase targets
the BMP pathway and affects embryonic pattern formation. *Nature* **400**: 687–693.

624