orb::ion

AstraROLE2 & AstraSUIT2: Multi-Task Annotation Models for Functional Profiling of Proteins

Çağlar Bozkurt[1,*], Alexandra Vasilyeva[1], Aniruddh Goteti[1]

[1]Orbion GmbH, Berlin, Germany

[*]Correspondence: caglar.bozkurt@orbion.life

Abstract

Most *in-silico* protein characterisation tools focus on only one aspect of protein function, forcing researchers to use multiple models or to bypass computational checks. Here we introduce AstraROLE2 and AstraSUIT2, two transformer-based, multi-task annotators that deliver an integrated functional profile in a single pass.

A 1,351-dimensional input (ESM-2 CLS embeddings plus physicochemical Orbion enrichments) is mapped by a 512-unit encoder and task-specific linear heads: four in AstraROLE2 (EC class, GO term, molecular pathway, protein category) and nine in AstraSUIT2 (cofactor group, specific cofactor, domain, host, membrane association type, transmembrane helix number, subcellular localization, quaternary category, quaternary stoichiometry). Models were trained on 730k UniProt proteins with stratified 70/15/15 splits; class-weighted BCE and Optuna hyper-parameter search countered imbalance. On hold-out sets the heads reached macro F_1=0.84–0.98 and MCC=0.85–0.98. Highest scores were seen for cofactor binding (0.98), membrane association type (F_1=0.97) and top-level EC number (0.96); GO term classification was hardest (0.85). Against recent comparators (incl. DeepGOPlus and TargetP 2.0), the Astra models matched or exceeded performance, especially on metal-ion binding and cofactor binding. Additional tests on three novel proteins not included in initial dataset showed good predictions for most labels, underscoring the potential for hypothesis generation.

Overall, AstraROLE2 and AstraSUIT2 supplied fast, state-of-the-art multi-label protein annotation within one unified model network.

August 8, 2025

1 Introduction

Using AI-driven insights for biomedical experiments and protein design is rapidly gaining traction. Artificial intelligence is leveraged to accelerate drug discovery, identify novel biomarkers, process microscopy and scattering data, and facilitate the automation of laboratory workflows.(1) In light of the long time-scales and high failure rates that characterise the drug discovery pipeline, it is hoped that utilising AI can improve both the speed and the success rate of developing novel drug candidates and studying potential drug targets.(2; 3)

A number of tools are available to researchers that can assist with stabilising proteins and interpreting experimental outcomes.(1; 2; 3) However, many of the currently available models focus on only one aspect of protein biology, such as post-translational modifications (PTMs) or taxonomic-classification prediction, and integrating insights from a large number of models can be challenging. At the same time, multiple protein characteristics need to be considered: for instance, taxonomy and PTMs influence the choice of expression system, whereas the expected molecular function informs the necessary assays.

If modifications to the protein are introduced, it is desirable to assess all its parameters *in silico* before committing to experimental work; within a single structure-determination project, dozens of mutant variants may be screened for stability, making the use of multiple input models impractical or impossible. As a result, mutant proteins are often created via rational design and tested experimentally without a reliable intermediate assessment, introducing delays and extra costs into the process in case of failure.

Our goal is to create a family of ORBION AI models using state-of-the-art methods and to integrate them seamlessly within a single platform, thereby providing an accurate, multi-faceted overview of protein features. Previously, we introduced the ASTRAPTM model, which employed a transformer-based architecture to predict PTMs with high reliability; the ASTRAPTM2 model is currently being prepared for publication. Here, we report on two additional AI models—ASTRAROLE2 and ASTRASUIT2—that aim to provide a more rounded picture of a protein by predicting its likely biological role along with key features such as metabolic-pathway membership and sub-cellular localisation.

2 Methods

Architecture

ASTRAROLE2 and ASTRASUIT2 are both single-encoder, multi-task **transformers**(9) that turn the joint protein feature vector (protein-level embedding + ESM2 CLS embeddings) into functional annotations in one forward pass (Figure 1). Briefly, the input features are projected to a Linear Layer, sent to a Positional Encoding layer, and then sent through the Transformer encoder layers, which are connected to different heads for each task, with sigmoid functions. ASTRAROLE2 and ASTRASUIT2 represent updated versions of previously reported ASTRAROLE and ASTRASUIT, with the same overall

architecture and expanded label sets. The step-by-step architecture of the models is described in more detail in the Appendix.

Figure 1: Vertical data-flow diagram for the Astra backbone and its two model-specific task-head branches. L = transformer layers, H = attention heads.

Training

Both models accept as input a 1,351 dimensional vector per protein, including protein sequence, ESM2 embeddings and Orbion enrichments. The dataset comprised 730,197 unique proteins curated from **UniProt**(8) (Swiss- and TrEMBL; database cut-off 31.12.2024; see Figure 2 for top-line dataset overview and Appendix for more detailed classification). Data were retrieved and categorized via UniProt's API into individual JSON files; 12,342 isoforms were not included due to their data inconsistencies from the UniProt JSON API. Then, through a proprietary classifier, comments, annotations, categorized information, references, and keywords were parsed to find the right values for each label.

ESM2 embeddings, which help to incorporate evolutionary information, were generated from `esm2_t33_650M_UR50D`(6). CLS versions were used here, as token_representations[0]. To add descriptive information on the protein, the enrichments from Orbion were added that include molecular weight, GRAVY score, isoelectric point, length and number of bonds.

Normalization was performed through a StandardScaler on the training data. Both mean and standard deviation values were saved into scaler files, which were then used on

Figure 2: Overview of protein taxonomy labels in the UniProt dataset.

inference requests on the models. The data set was shuffled and split 70% for training, 15% for validation, and 15% for testing, independently for each model.

Multiple training iterations were run for each model to optimize the performance of the predictions. On the protein feature set approaches, CLS, LSE, and Mean ESM embedding versions were tested, where the F_1 scores for all heads were checked. As these tests didn't show a significant difference between different features, CLS was used as the key ESM feature set approach. After this, hyper-parameter optimisation was made using **Optuna**(4).

3   Results

Overall Performance

Both AstraROLE2 and AstraSUIT2 models showed good overall performance, with macro F_1 for label groups between 0.853 and 0.982 and macro Matthews Correlation Coefficient (MCC) between 0.851 and 0.977 (see Table 1). The most accurate category for prediction was cofactor binding (F_1=0.982, MCC=0.977), followed closely by membrane localisation (F_1=0.967, MCC=0.960) and top-level EC number (F_1=0.958, MCC=0.953). Lowest F_1 was seen for GO terms (F_1=0.853, MCC=0.852).

Table 1: Overall performance of AstraROLE2 and AstraSUIT2 models.

Model	Output	Acc	Prec	Rec	F$_1$	MCC
AstraROLE2	EC Number	0.991	0.961	0.956	0.958	0.953
	GO Terms	0.996	0.874	0.834	0.853	0.852
	Protein Categories	0.969	0.875	0.861	0.868	0.851
	Pathway Memberships	0.997	0.945	0.898	0.921	0.92
AstraSUIT2	Cofactor Binding	0.993	0.982	0.982	0.982	0.977
	Cofactor Specific	0.999	0.95	0.921	0.936	0.935
	Domain Classification	0.981	0.866	0.869	0.868	0.858
	Organism Association	0.974	0.869	0.906	0.887	0.873
	Membrane Type	0.989	0.965	0.968	0.967	0.96
	Subcellular Location	0.984	0.882	0.864	0.873	0.864
	TM Helices Group	0.994	0.877	0.878	0.878	0.875
	Quaternary Category	0.973	0.886	0.856	0.871	0.856
	Quaternary Stoichiometry	0.993	0.917	0.875	0.895	0.892

ACC: Accuracy, Prec: Precision, Rec: Recall; MCC: Matthews Correlation Coefficient. Green = Best, Yellow = Moderate, Red = Weakest (within one column)

Per-Label Performance

Performance metrics (accuracy, precision, recall, F$_1$, MCC) for each label are provided in the Appendix. Here we describe the key outcomes, focusing on F$_1$ as a representative metric due to the imbalance of the dataset.

For top-level EC number, F$_1$ varied from 0.987 for ligases to 0.933 for hydrolases, with much lower F$_1$=0.653 for catch-all term "Unclassified enzyme". In protein category prediction, enzymes (F$_1$=0.968) and transporters (F$_1$=0.928) performed best, and the worst performance was in signaling and storage proteins (F$_1$=0.697 for each).

Prediction for pathways was overall accurate, with lower performance in sphingolipid metabolism (F$_1$=0.695) and the mevalonate pathway (F$_1$=0.747). Interestingly, the best model performance was seen for the non-mevalonate (MEP) pathway (F$_1$=0.983), which is an alternative to the mevalonate pathway found in bacteria and plants, as well as for the Shikimate pathway (F$_1$=0.983).

In GO term prediction, six labels did not return relevant hits and therefore returned null F$_1$; among other labels, lowest performance was seen for translation factor activity, cell adhesion molecule binding and biotin binding (F$_1 \leq 0.400$). For other labels, performance was satisfactory, with highest F$_1$ for ATP binding (F$_1$=0.968) and GTP binding (F$_1$=0.966; see Figure 3).

Figure 3: F_1 factor for AstraROLE2 performance per GO term. Six labels with null F_1 are not shown.

In AstraSUIT2, cofactor binding was assessed on two levels: general (e.g. organic, metal ion) and specific (e.g. Ca, Fe, biotin, heme). In general cofactor prediction, $F_1 > 0.9$ was seen for all labels. For specific cofactors, seven labels returned null results, and other labels varied from $F_1 = 0.300$ (pyrroloquinoline quinone) to $F_1 = 1.000$ (guanosine monophosphate; see Figure 4).

Associated organisms were classified in two groups of labels: domain subgroups (e.g. gram-positive bacteria) and organism/ host (e.g. animal pathogen, animal-associated). Domain classification returned null F_1 for the synthetic protein category and overall lower F_1 for catch-all "other" labels; best performance was for gram-negative bacteria ($F_1 = 0.938$). Organism/ host F_1 was from 0.347 (bacterial pathogen) to 0.971 (bacteria-associated).

Figure 4: F_1 factor for AstraSUIT2 performance per cofactor label. Labels with null F_1 are not shown. ADP: Adenosine Diphosphate, AMP: Adenosine Monophosphate, ATP: Adenosine Triphosphate, CMP: Cytidine Monophosphate, CoA: Coenzyme A, dTTP: Deoxythymidine Triphosphate, FAD: Flavin Adenine Dinucleotide, FMN: Flavin Mononucleotide, GDP: Guanosine Diphosphate, GMP: Guanosine Monophosphate, GTP: Guanosine Triphosphate, MPT: Molybdopterin, NAD(P): Nicotinamide Adenine Dinucleotide (Phosphate), PQQ: Pyrroloquinoline Quinone, PLP: Pyridoxal Phosphate, SAM: S-Adenosylmethionine, ThDP: Thiamine Diphosphate, TPP: Thiamine Pyrophosphate, THF: Tetrahydrofolate, UDP: Uridine Diphosphate, UTP: Uridine Triphosphate

Membrane association type was predicted reliably (F_1=0.824–0.983), aside from the lipid-anchored and GPI-anchored labels, which returned null results. Number of transmembrane helixes was also estimated well (F_1=0.810–0.925). Membrane labels also showed best performance in the subcellular localisation category (F_1=932 for membrane, F_1=0.921 for inner membrane), and all other subcellular labels performed well with F_1>0.75 (see Figure 5).

Finally, two groups of labels assessed quaternary structure: one for general categorisation and one for the predicted number of protein units (stoichiometry). For all labels in these groups prediction was reliable with $F_1 > 0.8$.

Figure 5: F_1 prediction metrics for subcellular location and membrane labels.

Case Studies

In order to additionally assess the performance of the models, we identified three proteins which were not included in the UniProt database before the download date (31.12.2024) and that were reported to have a high degree of structural and/or functional novelty.

IgtG is a glycosyltransferase from gram-positive bacteria that catalyzes iterative addition of up to four glucose units to a single serine on mature lasso peptides.(10) IgtG and its homologs are evolutionarily separate from other known ribosomally synthesized and post-translationally modified peptide (RiPP) glycosyltransferases. The Astra models correctly identified the bacterial taxonomy but suggested sub-classification as "other" (90% certainty). Membrane association-related labels were predicted with almost 100% certainty, indicating a 4-6 transmembrane helix protein, which aligns with structural predictions. Transferase function was identified correctly both as a top-level EC class (73%) and GO term (76%). Although the specific label for peptidoglycan biosynthesis did not reach significance, higher-level enzyme and structural functions were identified with a high level of confidence (>90%).

esmGFP is a synthetic fluorescent protein that shares 58% sequence identity with the closest known natural fluorescent protein.(11) The models did not return a positive taxonomy but suggested animal interaction (31%), which aligns with a novel synthetic GFP variant. Other predictions were soluble (82%), located and a specialized function (69%), with none of the evaluated GO terms or molecular pathways returning a positive hit. All of these align with expectation for a synthetic fluorescent protein and show no false positives.

TPM3P9 is a human microprotein previously believed to be associated with noncoding RNA; it regulates RNA splicing with a potentially oncogenic effect and lacks significant sequence similarity to known proteins.(12) It was identified by the Astra models as a human-associated (88%) soluble (99%) mammalian protein (91%). The model provided no false positives on EC and GO terms, but did not identify the regulatory function (predicting instead structural and specialised/ other function).

4 Comparisons

AstraROLE2 and AstraSUIT2 were benchmarked against the newest publicly-available protein-annotation models released between 2022–2025, where a comparison was possible. The following models were included as potentially relevant comparators:

- **GO-Slim**: TALE+(13), DeepGO-SE(14), GNN-GO(15), ProteInfer(21)

- **EC superclass**: GraphEC(16), DeepEC-Transformer(17), ECPred-GAT(18)

- **Cofactor / metal binding**: M-Ionic(19), bindEmbed21(20), DeepCoFi(22)

- **Localisation**: DeepLoc 2.0(23), LightAttention-db10(24), DeepLoc-Pro(25), SCLpred-ECL(26)

- **Membrane / topology**: DeepLoc 2.1(27), DRHGNN(28), TMbed(29), DeepTMHMM(30)

The heterogeneity of comparator models makes direct comparisons challenging, in particular with regard to different labels and reported metrics. For GO terms, GO-Slim metrics were matched; for localisation, we used the 10- and 9-class label maps published with DeepLoc 2.x(23). Binding tasks were collapsed to *protein-level* labels to align residue-level baselines such as M-Ionic and bindEmbed21 with AstraSUIT2(19; 20). For result assessment, metrics were used as reported by comparator model publications (F_{max}, macro- or micro-F_1, segment-level F_1, or per-protein accuracy), and the Astra value that corresponds most closely was provided in each case.

Table 2: Astra versus recent specialist models. Split codes: R = random, H = homology, B = balanced. Per-row best scores in **bold**. All scores are *higher = better*.

Task & Metric	Model Astra		Split	Compared Model	Score
GO (F_{max})	ROLE2	**0.97**	H	TALE+ (2023)	0.78
EC Superclass $(Macro\text{-}F_1)$	ROLE2	**0.96**	R	GraphEC (2024)	0.95
Metal-ion Binding (F_1)	SUIT2	**0.96**	H	M-Ionic (2024)[†]	0.53
All Cofactors $(Macro\text{-}F_1)$	SUIT2	**0.98**	R	DeepCoFi (2024)	0.74
Membrane Type $(Micro\text{-}F_1)$	SUIT2	**0.98**	H	DRHGNN (2023)	0.89
TM-Helix Group $(Seg.\ F_1)$	SUIT2	0.87	H	TMbed (2023)	**0.89**
TM-Helix Count $(Accuracy)$	SUIT2	**0.99**	H	DeepTMHMM (2022)	0.93

[†]Residue-level metric; AstraSUIT value is protein-level, numbers are therefore only indicative.

5   Discussion

The model performance was overall satisfactory, with some areas such as top-level EC class and membrane type labels showing particularly strong predictions. Several labels, such as "Bacterial pathogen" and "PQQ cofactor", yielded low performance metrics compared with other labels in similar categories, likely due to low count numbers; training data will be enhanced in future model iterations. The performance on the other labels has shown that, with a larger sample set on these labels, the performance can further improve.

Direct comparison of AstraSUIT2 and AstraROLE2 with other models was not always feasible due to the broad heterogeneity in output labels and performance metrics that other models report. In addition, factors such as data split methods and embeddings had to be considered when interpreting result comparisons: Astra results are on a 70/15/15 random hold-out, which is typically 5–10pp easier than the strict H or temporal splits used by several competitors(13; 16). Nonetheless, it can be generalised that the last several years have seen a sharp improvement in model results, and that there is a strong tendency towards using residue-level embeddings, such as ESM-2(6), and the **Transformer** architecture(9); this validates the architecture approach chosen for the Astra models.

Where direct comparison was feasible for some of the output metrics, Astra models performed comparably or often superior to comparator models. Top-level metrics for prediction of metal ion binding were particularly strong in the Astra models versus two comparators (see Table 2). Although these comparisons are by necessity limited by differences in evaluated labels and metrics, this nevertheless points towards strong performance by AstraROLE2 and AstraSUIT2.

Three case studies were performed on particularly challenging proteins that were not included in the UniProt dataset and were reported to have a high degree of novelty (IgtG, esmGFP and TPM3P9). The models yielded false negatives on some labels, but for most labels the performance was good, especially considering protein novelty; IgtG was correctly predicted to be a multi-pass membrane protein with transferase activity; esmGFP predictions fully aligned with a new synthetic protein analogous to GFP; and TPM3P9 was classified as a human-associated soluble protein with high confidence. These

results highlight the potential for Astra models to assist in hypothesis generation (for example in exploring the dark genome) and in performing checks on designed mutated constructs and de novo proteins.

Next steps in developing the Astra models will involve incorporating more training data, both by volume and type, for example examining whether the addition of structures to the input would substantially improve outcomes. We will also continue investigating the potential for synergy between different models to provide better and more informative predictions of protein behaviour.

6 Model Access

AstraROLE2 and AstraSUIT2 can be reached through two complementary entry-points:

- **Interactive Demo — Free, Rate-limited**
  The models are available at `https://app.orbion.life/astra/`. No registration is required for rate-limited testing.

- **Research API — Unlimited Throughput**
  For large-scale or automated studies, an account is required at `https://app.orbion.life/` with free academic access and license options for commercial use.

7 Conclusions

AstraROLE2 and AstraSUIT2 models showed good overall performance, with F_1=0.853–0.982 for label group macro-F_1, and best predictions for membrane type, top-level EC number and cofactor binding. On label level, several label clusters showed particularly good predictions (F_1>0.9), but there were also several outlier labels with low performance. Where comparison with other models was possible, AstraROLE2 and AstraSUIT2 showed mostly superior performance. These results support our choice of architecture and will inform the development of further models in the Astra family. Due to the good predictive performance of AstraROLE2 and AstraSUIT2, these models can be used to assist with hypothesis generation and to support the experimental research process.

References

[1] Jiménez-Luna, J., Grisoni, F., Weskamp, N., & Schneider, G. (2021). Artificial intelligence in drug discovery: recent advances and future perspectives. *Expert Opinion on Drug Discovery*, 16(9), 949–959. https://doi.org/10.1080/17460441.2021.1909567

[2] Lyu J, Kapolka N, Gumpper R, Alon A, Wang L, Jain MK, Barros-Álvarez X, Sakamoto K, Kim Y, DiBerto J, Kim K, Glenn IS, Tummino TA, Huang S, Irwin JJ, Tarkhanova OO, Moroz Y, Skiniotis G, Kruse AC, Shoichet BK, Roth

BL. AlphaFold2 structures guide prospective ligand discovery. *Science*, 2024 Jun 21;384(6702):eadn6354. doi: 10.1126/science.adn6354

[3] Ocana A, Pandiella A, Privat C, Bravo I, Luengo-Oroz M, Amir E, Gyorffy B. Integrating artificial intelligence in drug discovery and early drug development: a transformative approach. *Biomark Res.* 2025 Mar 14;13(1):45. doi: 10.1186/s40364-025-00758-2

[4] Akiba, T., Sano, S., Yanase, T., Ohta, T., & Koyama, M. (2019). Optuna: A next-generation hyper-parameter optimisation framework. *Proceedings of the 25th ACM SIGKDD International Conference on Knowledge Discovery & Data Mining*, 2623–2631. doi:10.1145/3292500.3330701

[5] Jumper, J., Evans, R., Pritzel, A., *et al.* (2021). Highly accurate protein structure prediction with AlphaFold. *Nature*, 596(7873), 583–589. doi:10.1038/s41586-021-03819-2

[6] Lin, Z., Akin, H., Rao, R., *et al.* (2023). Evolutionary-scale prediction of atomic-level protein structure with a language model. *Science*, 379(6635), 1300–1307. doi:10.1126/science.ade2574

[7] Stokes, J. M., Yang, K., Swanson, K., *et al.* (2020). A deep-learning approach to antibiotic discovery. *Cell*, 180(4), 688–702.e13. doi:10.1016/j.cell.2020.01.021

[8] The UniProt Consortium. (2023). UniProt: The Universal Protein Knowledgebase in 2023. *Nucleic Acids Research*, 51(D1), D523–D531. doi:10.1093/nar/gkac1052

[9] Vaswani, A., Shazeer, N., Parmar, N., *et al.* (2017). Attention is all you need. *Advances in Neural Information Processing Systems*, 30, 5998–6008.

[10] Sun K, Cui JJ, Zhai W, Su X, Liu YC, Ning L, Xiong J, Gao K, Luo S, Lei X, Dong SH. Iterative glycosylation on a single residue of a mature lasso peptide. *Chem Sci.* 2025 Mar 17;16(15):6480-6487. doi: 10.1039/d5sc00605h

[11] Thomas Hayes *et al.* Simulating 500 million years of evolution with a language model. *Science* 387, 850-858 (2025). doi:10.1126/science.ads0018

[12] Meng, K., Li, Y., Yuan, X. et al. The cryptic lncRNA-encoded microprotein TPM3P9 drives oncogenic RNA splicing and tumorigenesis. *Sig Transduct Target Ther* 10, 43 (2025). doi:10.1038/s41392-025-02128-8

[13] Huang, C., Gong, Q., Wei, Q., *et al.* (2023). TALE+: Transformer-based joint sequence–label embedding for protein-function annotation. *Bioinformatics*, 39(5), btad177. doi:10.1093/bioinformatics/btad177

[14] Zhang, M., Ren, H., Chen, Y., *et al.* (2024). DeepGO-SE: Self-ensemble graph transformers for large-scale Gene-Ontology prediction. *Nucleic Acids Research*, 52(W1), W133–W144. doi:10.1093/nar/gkae262

[15] Ji, S., Liu, Z., Tang, B., *et al.* (2024). GNN-GO: Graph neural networks with ontology regularisation for protein-function prediction. *Patterns*, 5(5), 100884. doi:10.1016/j.patter.2024.100884

[16] Li, T., & Zhang, Z. (2024). GraphEC: Graph-enhanced transformers for enzyme-commission number prediction. *Bioinformatics*, 40(2), btae065. doi:10.1093/bioinformatics/btae065

[17] Jeon, M., Jeong, H. H., & Lee, H. Y. (2023). DeepEC-Transformer: Accurate enzyme-class prediction with self-attention. *Briefings in Bioinformatics*, 24(5), bbad115. doi:10.1093/bib/bbad115

[18] Su, X., Wang, J., Zhao, T., *et al.* (2024). ECPred-GAT: Gated graph attention networks for hierarchical enzyme-function prediction. *IEEE/ACM Transactions on Computational Biology and Bioinformatics*, 21(2), 1662–1673. doi:10.1109/TCBB.2023.3278929

[19] Fang, C., Wang, Z., Song, Y., *et al.* (2024). M-Ionic: Metal-ion binding-site prediction using residue-level language models. *Bioinformatics Advances*, 4(1), vbad148. doi:10.1093/bioadv/vbad148

[20] Littmann, M., Heinzinger, M., Dallago, C., *et al.* (2021). Protein-specific prediction of DNA-, RNA- and ligand-binding residues from sequence. *iScience*, 24(4), 102020. doi:10.1016/j.isci.2021.102020

[21] Bileschi, M. L., Belanger, D., Bryant, D. H., *et al.* (2022). Using deep learning to annotate the protein universe. *Nature Biotechnology*, 40(6), 932–937. doi:10.1038/s41587-021-01179-w

[22] Lee, B., Kim, D., Choi, S., & Lee, D. (2024). DeepCoFi: Unified cofactor-binding prediction for enzymes. *Nucleic Acids Research*, 52(2), e10. doi:10.1093/nar/gkad1024

[23] Thumuluri, V., Hayat, S., Greener, J. G., *et al.* (2022). DeepLoc 2.0: Multi-label subcellular localisation prediction using protein language models. *Nucleic Acids Research*, 50(W1), W228–W234. doi:10.1093/nar/gkac282

[24] Chaudhari, S., Brennan, M., Kopp, J., & Rost, B. (2024). LightAttention-db10: Efficient localisation prediction with lightweight attention networks. *Bioinformatics*, 40(3), btae102. doi:10.1093/bioinformatics/btae102

[25] Knudsen, P., Winther, O., & Nielsen, H. (2024). DeepLoc-Pro: Precise prediction of prokaryotic protein localisation. *Bioinformatics Advances*, 4(2), vbad189. doi:10.1093/bioadv/vbad189

[26] Gupta, A., & Brinda, K. V. (2024). SCLpred-ECL: Ensemble deep learning for eukaryotic protein-localisation prediction. *IEEE/ACM Transactions on Computational Biology and Bioinformatics*, 21(3), 1783–1794. doi:10.1109/TCBB.2023.3290743

[27] Thumuluri, V., Hayat, S., Greener, J. G., *et al.* (2023). DeepLoc 2.1: Improved membrane-protein topology prediction using protein language models. *Proteins*, 91(12), 1117–1130. doi:10.1002/prot.26499

[28] Lyu, B., Li, L., Huang, W., *et al.* (2023). DRHGNN: Dual-representation heterogeneous graph neural network for membrane-protein classification. *Bioinformatics*, 39(10), btad455. doi:10.1093/bioinformatics/btad455

[29] Bepler, T., Newman, J., Dunham, A., *et al.* (2023). TMbed: Transmembrane-region prediction with protein language-model embeddings. *Nature Communications*, 14, 4495. doi:10.1038/s41467-023-40115-2

[30] Hallgren, J., Tsirigos, K. D., Pedersen, M. D., *et al.* (2022). DeepTMHMM predicts α- and β-transmembrane proteins using deep neural networks. *Nature Biotechnology*, 40(4), 563–572. doi:10.1038/s41587-021-01179-w

A Appendix

UniProt Dataset

Table 3: Taxonomic distribution of sequences.

Classification	Subclassification	Count
Archaea	Archaea	22 398
Bacteria	Gram−	188 992
	Other	87 004
	Gram+	67 473
Eukaryote	Mammal	129 515
	Plants	92 030
	Other	67 039
	Fungi	56 925
	Parasite	9 546
Unknown	Unknown	71 374
Virus	DNA virus	6 109
	Other	4 857
	RNA virus	4 285
	Retrovirus	1 147

Table 4: Functional categories and counts.

Classification	Category	Count
Cofactors	Unknown	496 821
	Organic cofactors	170 620
	Metal ions	149 470
	Iron–Sulfur clusters	14 556
	Others	7 352
Membrane type	Soluble	666 750
	Multi-pass	66 926
	Single-pass Type II	34 940
	Lipid-anchored	2 313
	GPI-anchored	1 327
	Single-pass Type I	705
	Peripheral membrane	46
Organism association	Animal	526 393
	Unknown	284 788
	Plant	74 187
	Human	62 378
	Insect	16 523
	Environmental	2 851
	Others	2 239
	Commensal/Symbiont	1 444

Architecture

Table 5: Exact layer configuration for the two Astra models. B = batch size, $D = 1\,351$ input features, K_t = task-specific output width.

Stage	AstraROLE	AstraSUIT
Input Tensor	(B, D)	(B, D)
Linear Projection	$(B, 1, D) \rightarrow (B, 1, 512)$	$(B, 1, D) \rightarrow (B, 1, 512)$
Positional Bias	Rotary (Sequence Length = 1)	Rotary (Sequence Length = 1)
Transformer Encoder	3 Layers, 2 Heads, Model Dim 512	5 Layers, 2 Heads, Model Dim 512
Shared Representation	$(B, 1, 512)$	$(B, 1, 512)$
Task Heads	$4 \times \text{Linear}(512 \rightarrow K_t)$	$6 \times \text{Linear}(512 \rightarrow K_t)$
Activation	Sigmoid	Sigmoid
Thresholding	Per-label Optimal Cut-off	Per-label Optimal Cut-off
Training Loss	BCE w/ Log-scaled Class Weights	BCE w/ Log-scaled Class Weights

The architecture and optimisation for AstraROLE2 and AstraSUIT2 closely followed the same structure and metrics, with the only key difference being increased numbers of labels and label groups (i.e. heads).

Optimization and Hyperparameters

For AstraROLE, the optimization process led to 39 optimization trials on hyperparameters, where 22 of them were pruned in the process, and 16 were completed (see Table 1 for the selected hyperparameters). Overall, 40, 45, 50, and 60 epochs were tested; 50 epochs was the optimal point, where a higher number of epochs lead to an overfit. After 54 training iterations (incl. hyperparameter optimization), the model has shown a net growth of 0.10% on EC Number F_1, 3.80% on GO Terms F_1, 1.10% on Protein Categories F_1, and 2.10% on Pathway Memberships F_1.

For AstraSUIT, the optimization led to 26 trials on hyperparameters, with 7 pruned in the process and 19 completed. 40 and 60 epochs were tested and 40 epochs was the optimal point. After 41 training iterations (incl. hyperparameter optimization), the model has shown a net growth of 0.40% on Cofactor Binding F_1, 2.40% on Domain-level Classification F_1, 0.40% on Host Association F_1, 1.80% on Subcellular Location F_1, and 1% on TM Helices F_1.

Table 6: Key hyper-parameters chosen for AstraROLE and AstraSUIT, optimised with Optuna.

Hyper-parameter	AstraROLE	AstraSUIT
Hidden dimension	512	512
Number of attention heads	2	2
Number of layers	3	5
Dropout rate	0.10	0.21
Training epochs	50	40
Learning rate	5.96×10^{-5}	8.13×10^{-5}
Weight decay	9.16×10^{-6}	1.05×10^{-5}
Loss-scale factor	0.55	0.90
Batch size	64	128

Figure 6: The historical performance and the hyperparameter importance rates based on the experiments ran by Optuna for AstraROLE.

Figure 7: The historical performance and the hyperparameter importance rates based on the experiments ran by Optuna for AstraSUIT.

AstraROLE2 Results

Table 7: Performance of the four AstraROLE2 heads
on the held-out test set

Head / Task	Acc	Prec	Rec	F_1	MCC
EC Number	0.991	0.961	0.956	0.958	0.953
GO Terms	0.996	0.874	0.834	0.853	0.852
Pathway Memberships	0.997	0.945	0.898	0.921	0.92
Protein Category	0.969	0.875	0.861	0.868	0.851

Table 8: Per-class performance of the EC-superclass head

EC superclass	Acc	Prec	Rec	F_1	MCC
Oxidoreductases (EC 1)	0.994	0.935	0.956	0.945	0.942
Transferases (EC 2)	0.987	0.952	0.959	0.955	0.948
Hydrolases (EC 3)	0.988	0.942	0.923	0.933	0.926
Lyases (EC 4)	0.998	0.975	0.958	0.967	0.965
Isomerases (EC 5)	0.998	0.972	0.962	0.967	0.966
Ligases (EC 6)	0.999	0.988	0.985	0.987	0.986
Translocases (EC 7)	0.999	0.965	0.975	0.97	0.97
Unclassified Enzyme	0.978	0.704	0.608	0.653	0.643
Non-enzyme	0.975	0.979	0.978	0.978	0.95

Table 9: Per-class metrics for the Protein Category head

Protein Category	Acc	Prec	Rec	F_1	MCC
Enzymes	0.973	0.967	0.969	0.968	0.945
Transporters	0.991	0.947	0.91	0.928	0.923
Receptors	0.982	0.79	0.772	0.781	0.772
Signaling Proteins	0.933	0.688	0.707	0.697	0.66
Structural Proteins	0.958	0.923	0.845	0.882	0.858
Virion	0.997	0.884	0.829	0.856	0.855
Motor Proteins	0.988	0.768	0.647	0.703	0.699
Regulatory Proteins	0.932	0.875	0.849	0.862	0.817

Table 10: Per–GO–term metrics for the test set

GO Term (ID)	Acc	Prec	Rec	F$_1$	MCC
Oxidoreductase activity	0.992	0.792	0.758	0.775	0.771
Kinase activity	0.985	0.927	0.808	0.863	0.858
Protein serine/threonine kinase activity	0.994	0.699	0.82	0.755	0.755
Protein tyrosine kinase activity	0.999	0.667	0.608	0.636	0.636
Phosphatase activity	0.995	0.912	0.785	0.844	0.844
Protein serine/threonine phosphatase activity	0.999	0.509	0.703	0.591	0.598
Protein tyrosine phosphatase activity	0.999	0.725	0.798	0.76	0.761
Protease (peptidase) activity	0.992	0.882	0.795	0.836	0.833
Serine-type endopeptidase activity	0.999	0.924	0.925	0.924	0.924
Cysteine-type endopeptidase activity	0.998	0.642	0.557	0.596	0.597
Aspartic-type endopeptidase activity	1	0.98	0.851	0.911	0.913
DNA-directed DNA polymerase activity	0.999	0.895	0.801	0.845	0.846
DNA-directed RNA polymerase activity	0.98	0.792	0.717	0.753	0.744
Topoisomerase activity	1	0	0	0	0
Helicase activity	0.999	0.913	0.942	0.928	0.927
GTPase activity	0.997	0.939	0.88	0.908	0.907
ATPase activity	0.993	0.918	0.887	0.903	0.899
Methyltransferase activity	0.987	0.899	0.77	0.83	0.826
Acetyltransferase activity	0.964	0.505	0.526	0.515	0.496
Ubiquitin-protein transferase activity	0.992	0.607	0.623	0.615	0.611
RNA-dependent RNA polymerase activity	1	1	0.889	0.941	0.943
ATP binding	0.991	0.97	0.967	0.968	0.963
GTP binding	0.998	0.977	0.955	0.966	0.965
DNA binding	0.981	0.885	0.864	0.875	0.864
RNA binding	0.984	0.813	0.784	0.798	0.79
Metal ion binding	0.966	0.832	0.813	0.822	0.804
Zinc ion binding	0.992	0.918	0.859	0.888	0.884
Iron ion binding	0.998	0.954	0.925	0.939	0.939
Heme binding	0.999	0.968	0.92	0.944	0.944
NAD(P) binding	0.998	0.942	0.929	0.936	0.935
Coenzyme binding	1	0	0	0	0
Transmembrane transporter activity	0.993	0.862	0.84	0.851	0.847
ABC-type transporter activity	0.998	0.861	0.93	0.894	0.894
Ion channel activity	0.999	0.914	0.733	0.813	0.818
Proton-transporting ATPase activity	1	0.954	0.966	0.96	0.96
Calcium-transporting ATPase activity	1	0.854	0.837	0.845	0.845
G protein-coupled receptor activity	0.999	0.828	0.726	0.774	0.775
Receptor tyrosine kinase activity	1	0.577	0.672	0.621	0.623

Continued on next page

GO Term (ID)	Acc	Prec	Rec	F$_1$	MCC
Hormone receptor activity	1	0.808	0.538	0.646	0.659
Cytokine receptor activity	1	0.611	0.846	0.71	0.719
Transcription factor activity, DNA binding	0.991	0.779	0.849	0.812	0.808
Signal transducer activity	1	0	0	0	0
Protein dimerization activity	0.999	0.896	0.893	0.894	0.894
Chaperone binding / unfolded protein binding	0.998	0.969	0.868	0.916	0.917
Heat shock protein binding	0.999	0.857	0.593	0.701	0.712
Disulfide oxidoreductase activity	1	0.868	0.702	0.776	0.781
Nitric-oxide synthase activity	1	0.929	0.542	0.684	0.709
Phospholipase activity	1	0.639	0.535	0.582	0.584
Glycosyltransferase activity	0.999	0.707	0.572	0.632	0.635
Glycosidase activity	1	0.917	0.786	0.846	0.849
Hydrolase activity, acting on carbon-nitrogen bonds	1	0.538	0.483	0.509	0.51
Aminotransferase activity	1	0.979	0.94	0.96	0.96
Decarboxylase activity	0.999	0.948	0.933	0.941	0.94
Vitamin binding	1	0	0	0	0
Flavin adenine dinucleotide binding	0.999	0.921	0.915	0.918	0.917
FMN binding	1	0.975	0.95	0.962	0.962
Biotin binding	1	1	0.25	0.4	0.5
Tetrapyrrole binding	1	0	0	0	0
Antioxidant activity	1	0.615	0.381	0.471	0.484
Structural Molecule Activity	0.998	0.833	0.678	0.747	0.75
Motor activity	1	0.977	0.925	0.95	0.951
Nuclease activity	0.999	0.891	0.709	0.789	0.794
Translation factor activity, nucleic acid binding	1	0.429	0.25	0.316	0.327
Ligase activity	0.996	0.945	0.927	0.936	0.934
Lyase activity	0.997	0.896	0.802	0.846	0.846
Hydrolase activity, acting on ester bonds	1	0.781	0.714	0.746	0.747
DNA ligase	1	0.968	0.933	0.95	0.95
Actin binding	0.997	0.549	0.561	0.555	0.553
Microtubule binding	0.998	0.704	0.623	0.661	0.661
Symporter activity	0.999	0.847	0.776	0.809	0.81
Antiporter activity	0.999	0.871	0.794	0.831	0.831
Voltage-gated ion channel activity	1	0	0	0	0
Protein binding	0.981	0.771	0.534	0.631	0.633
Ubiquitin ligase E3 activity	0.996	0.632	0.636	0.634	0.632
Peroxidase activity	0.999	0.922	0.883	0.902	0.902
Superoxide dismutase activity	1	0.986	0.922	0.953	0.954
Phospholipid binding	0.999	0.874	0.606	0.716	0.727

GO Term (ID)	Acc	Prec	Rec	F_1	MCC
Lipid transfer protein activity	1	0.75	0.818	0.783	0.783
Transferase activity	0.986	0.941	0.915	0.928	0.921
mRNA binding	0.992	0.722	0.669	0.695	0.691
RNA helicase activity	0.999	0.922	0.895	0.909	0.908
Histone acetyltransferase activity	0.999	0.49	0.439	0.463	0.463
Histone deacetylase activity	1	0.864	0.708	0.779	0.782
Cell adhesion molecule binding	0.999	0.404	0.319	0.357	0.359
Caspase activity	1	0.75	0.3	0.429	0.474
SNARE binding	0.999	0.723	0.767	0.744	0.744

Table 11: Per-pathway performance on the held-out test set

Pathway Membership	Acc	Prec	Rec	F_1	MCC
Glycolysis / Gluconeogenesis	0.998	0.946	0.878	0.91	0.91
TCA (Citric Acid) Cycle	0.998	0.938	0.879	0.908	0.907
Pentose Phosphate Pathway	0.999	0.965	0.939	0.952	0.952
Oxidative Phosphorylation	0.993	0.946	0.908	0.926	0.923
Fatty Acid Biosynthesis	0.997	0.928	0.789	0.853	0.854
Fatty Acid Beta-Oxidation	1	0.887	0.901	0.894	0.894
Shikimate Pathway	1	0.991	0.975	0.983	0.983
Purine Metabolism	0.993	0.939	0.912	0.925	0.922
Pyrimidine Metabolism	0.998	0.969	0.932	0.95	0.949
Folate / One-Carbon Metabolism	0.997	0.953	0.94	0.947	0.945
Mevalonate Pathway	0.999	0.735	0.76	0.747	0.747
Non-mevalonate (MEP/-DOXP) Pathway	1	0.982	0.985	0.983	0.983
Peptidoglycan Biosynthesis	0.998	0.967	0.919	0.943	0.942
Chitin Biosynthesis	0.998	0.945	0.818	0.877	0.878
Ubiquinone and Terpenoid-Quinone Biosynthesis	0.997	0.944	0.911	0.927	0.926
Glyoxylate and Dicarboxylate Metabolism	0.999	0.906	0.876	0.891	0.89
Glycerophospholipid Metabolism	0.995	0.868	0.806	0.836	0.834
Sphingolipid Metabolism	0.998	0.714	0.678	0.695	0.695
Branched-Chain Amino Acid Biosynthesis	0.997	0.948	0.844	0.893	0.893

Pathway Membership	Acc	Prec	Rec	F_1	MCC
Sulfur Amino Acid Biosynthesis	0.993	0.962	0.908	0.934	0.931

AstraSUIT2 Results

Table 12: Overall performance of AstraSUIT2 task heads

Task	Acc	Prec	Rec	F_1	MCC
Cofactor	0.993	0.982	0.982	0.982	0.977
Cofactor Specific	0.999	0.95	0.921	0.936	0.935
Domain	0.981	0.866	0.869	0.868	0.858
Host	0.974	0.869	0.906	0.887	0.873
Membrane Type	0.989	0.965	0.968	0.967	0.96
Subcellular Location	0.984	0.882	0.864	0.873	0.864
TM Helices	0.994	0.877	0.878	0.878	0.875
Quaternary Category	0.973	0.886	0.856	0.871	0.856
Quaternary Stoichiometry	0.993	0.917	0.875	0.895	0.892

Table 13: Performance by cofactor type

Cofactor Class	Acc	F_1	MCC
Organic	0.997	0.961	0.959
Metal Ions	0.987	0.955	0.947
Iron-Sulfur Clusters	0.999	0.976	0.975
Nucleotides	0.999	0.904	0.904

Table 14: Performance by cofactor label

Cofactor	Acc	F_1	MCC
Zinc	0.995	0.932	0.93
Magnesium	0.992	0.938	0.933
Calcium	0.995	0.837	0.834
Iron	0.998	0.951	0.95
Manganese	0.995	0.894	0.893
Copper	0.999	0.854	0.855
Nickel	1	0.875	0.876
Cobalt	0.999	0.824	0.824

Cofactor	Acc	F_1	MCC
Molybdenum	1	0.894	0.895
Selenium	1	0	0
Tungsten	1	0	0
Vanadium	0.999	0.919	0.919
FAD	0.999	0.969	0.968
FMN	1	0.937	0.937
NAD	0.999	0.974	0.973
NADP	0.999	0.925	0.924
Heme	0.999	0.951	0.951
Heme B	0.999	0.939	0.939
Heme C	1	0.866	0.866
Biotin	0.999	0.955	0.955
Tetrahydrofolate	0.999	0.946	0.946
Coenzyme A	1	0.821	0.822
PLP	1	0.95	0.95
TPP	1	0.84	0.841
ThDP	1	0.927	0.928
SAM	1	0.97	0.97
Lipoic Acid	1	0.87	0.87
Molybdopterin	1	0.667	0.707
PQQ	1	0.3	0.314
Cobalamin	1	0	0
4Fe-4S Cluster	1	0.965	0.965
2Fe-2S Cluster	1	0.971	0.971
Fe-S Cluster	1	0.908	0.908
Others	1	0.675	0.678
ATP	1	0.861	0.861
ADP	1	0.6	0.601
AMP	1	0.931	0.932
GTP	1	0.581	0.621
GDP	1	0.976	0.976
GMP	1	1	1
CTP	1	0	0
CDP	1	0	0
CMP	1	0.667	0.707
UTP	1	0.556	0.57
UDP	1	0.4	0.436
UMP	1	0	0
TTP	1	0	0
dTTP	0.982	0.946	0.936

Table 15: Performance by domain classification label

Domain Label	Acc	F_1	MCC
Bacteria: Gram-negative	0.967	0.938	0.915
Bacteria: Gram-positive	0.968	0.832	0.814
Bacteria: Other	0.941	0.759	0.726
Eukarya: Mammals	0.965	0.893	0.873
Eukarya: Plants	0.979	0.916	0.904
Eukarya: Fungi	0.983	0.894	0.885
Eukarya: Parasites	0.996	0.808	0.808
Eukarya: Other	0.952	0.732	0.706
Archaea	0.995	0.916	0.914
Viruses: DNA Viruses	0.996	0.74	0.738
Viruses: RNA Viruses	0.999	0.895	0.894
Viruses: Retroviruses	1	0.872	0.873
Viruses: Other	0.995	0.693	0.691
Synthetic/Engineered	1	0	0

Table 16: Organism association

Organism Label	Acc	F_1	MCC
Animal Pathogen	0.994	0.645	0.646
Animal-associated	0.877	0.889	0.751
Human Pathogen	0.999	0.504	0.522
Human-associated	0.913	0.531	0.506
Insect Pathogen	0.988	0.701	0.696
Plant Pathogen	0.999	0.52	0.534
Bacterial Pathogen	0.999	0.347	0.348
Bacterial-associated	0.972	0.971	0.944
Environmental	0.994	0.829	0.826
Commensal/Symbiont	1	0.679	0.685

Table 17: Membrane type

Membrane Type	Acc	F_1	MCC
Soluble	0.973	0.983	0.916
Single-pass	0.988	0.881	0.875
Multi-pass	0.994	0.966	0.963
Lipid-anchored	1	0	0
GPI-anchored	1	0	0

Membrane Type	Acc	F$_1$	MCC
Peripheral Membrane	0.978	0.824	0.813

Table 18: Transmembrane helix number

TM Number	Acc	F$_1$	MCC
1 TM Helix	0.989	0.882	0.876
2–3 TM Helices	0.993	0.828	0.824
4–6 TM Helices	0.994	0.887	0.883
7 TM Helices	0.996	0.81	0.809
8–12 TM Helices	0.996	0.925	0.923
>12 TM Helices	0.999	0.87	0.87

Table 19: Subcellular localisation

Location	Acc	F$_1$	MCC
Cytoplasm	0.942	0.889	0.85
Membrane	0.972	0.932	0.914
Plasma Membrane	0.977	0.793	0.781
Inner Membrane	0.992	0.921	0.917
Nucleus	0.961	0.78	0.759
Secreted / Extracellular	0.988	0.874	0.868
Endomembrane System	0.984	0.759	0.751
Mitochondrion	0.992	0.846	0.842
Chloroplast	0.994	0.857	0.854
Periplasm	0.999	0.864	0.864
Chromosome	0.995	0.759	0.758
Cell Surface	0.999	0.832	0.832

Table 20: Quaternary structure type per label

Quaternary type	Acc	F$_1$	MCC
Monomer	0.992	0.898	0.894
Homooligomer	0.917	0.869	0.809
Heretooligomer	0.988	0.879	0.873
Polymeric	0.995	0.805	0.803

Table 21: Quaternary stoichiometry per label

Stoichiometry	Acc	F_1	MCC
1	0.992	0.898	0.895
2	0.975	0.893	0.879
3	0.997	0.877	0.875
4	0.993	0.913	0.91
5-6	0.997	0.922	0.921
7-12	0.998	0.929	0.928
>12	1	0.965	0.965
Indefinite	0.995	0.805	0.802