Glycolipid recognition and binding by Siglec-6 hinges on interactions with the cell membrane

Silvia D'Andrea[1], Edward N. Schmidt[2], Duong Bui[2,], Ojas Singh[1], Ling Han[2], Lara K. Mahal[2], John S. Klassen[2], Matthew S. Macauley[2] and Elisa Fadda[3*]

1.Department of Chemistry, Maynooth University, Maynooth, Ireland; 2. Department of Chemistry, University of Alberta, Edmonton, Canada, 3. School of Biological Sciences, University of Southampton, Southampton, United Kingdom.

Abstract

Sialic acid-binding immunoglobulin-type lectins (Siglecs) regulate immune response through interactions with sialylated glycan epitopes on glycoproteins and glycolipids. Human Siglecs count 14 unique proteins allocated to two subsets; group 1 includes four evolutionarily conserved proteins with clear orthologues in all mammals with low sequence similarity and group 2 includes ten rapidly evolving proteins with high sequence similarity and species specificity. In all Siglecs, the recognition and binding of the sialic acid on the glycan target involves a conserved, or canonical, Arg residue. Yet, for a subset of human Siglecs, namely MAG, Siglec-6, and Siglec-11, this Arg appears not to be essential, suggesting that a different binding mechanism may be at play. In this work, we used all-atom molecular dynamics (MD) simulations, binding assays, and mutagenesis to investigate the structural, mechanistic and energetic details of the binding of Siglec-6 to monosialylated gangliosides. Our results show that Siglec-6 relies only partially on its conserved Arg (Arg122) for recognition of membrane-bound gangliosides and that it supplements its binding free energy through interactions with the phospholipids in the membrane surrounding the target epitope. We identified residues Lys126 and Trp127 as key players in this unique membrane interaction, and we confirmed by mutagenesis assays that the loss of these residues abrogates binding. These results provide a step-change in our understanding of the diversification of human Siglecs across group 2, where different members likely evolved as molecular precision tools, to bind specific sialosides by adapting their structure to recognize not only the epitope, but also the biological environment in which it is found.

---

[*] For correspondence: Elisa Fadda, elisa.fadda@soton.ac.uk

Introduction

Sialic acid-binding immunoglobulin (Ig)-like lectins (Siglecs) are immunoregulatory transmembrane receptors that recognise sialylated glycoconjugates. Siglecs modulate immune responses by triggering inhibitory or activating signals, maintaining immune homeostasis[1–4]. The ability to distinguish self from non-self through sialic acid recognition makes Siglecs critical regulators of immune function and attractive therapeutic targets for autoimmune diseases, cancer, and infections[5], as well as for the development of chimeric antigen receptor (CAR) T-cell immunotherapeutic strategies[6–8]. Humans Siglecs count 14 distinct members that are commonly classified into two groups, see **Figure 1**. Evolutionarily conserved Siglecs[9], here labelled as Group 1, are present in all mammalian species[10] and have a low degree of homology, i.e. 25–30% sequence identity, namely sialoadhesin (Siglec-1), CD22 (Siglec-2), myelin-associated glycoprotein (MAG or Siglec-4), and Siglec-15. Group 2 comprises CD33-related Siglecs (CD33rSiglecs), which are not conserved across species, but share a higher sequence similarity (~50–99%)[1], suggesting that diversification could have occurred in response to species-specific selective pressures[10].

Figure 1. Schematic representation of human Siglecs. Evolutionarily conserved Siglecs (Group 1) are shown on the left-hand side panel (pink box). The CD33-related Siglecs (Group 2) are shown on the right-hand side panel, (light-green box). The Siglecs extracellular domains are represented by a composition of Ig domains (C2-set) rendered in grey as a solvent accessible surface, highlighted within green circles. The terminal Ig V-set binding domain is

rendered in orange and highlighted within an orange circle with a bound sialic acid in magenta (PDB 1OD9). Cytoplasmic domains are indicated with boxes and labelled according to the legend. Siglec-6 is highlighted within a yellow box as it is the focus of this work. All structural elements in this image were rendered with pymol (www.pymol.org) and incorporated in an original design based on Figure 1 in ref. [2]. In the insert on the top right-hand side, the conserved Arg is shown in the 3D structure of the Ig domain V-set of Siglec-6 (orange cartoons) bound to the NeuNAc-a(2-3)-Gal fragment (sticks with purple C atoms, red O atoms and blue N atoms) as an example.

In terms of 3D structure, human Siglecs are characterized by a different number of Ig C2-set domains terminating with an Ig V-set domain that contains the canonical binding site, where a conserved Arg binds sialosides[1], see **Figure 1**. The V-set domain includes a structurally flexible C-C' loop, characterised by a high sequence variability across Siglecs. Both of these features allow the C-C' loop to play a critical role in modulating recognition[1] and contribute to the Siglecs' binding specificity[5,11].

Target sialosides can be presented to Siglecs within a wide variety of different structural contexts, namely functionalising glycans linked to membrane bound or soluble glycoproteins, terminating the chains of membrane-embedded glycolipids or on secreted oligosaccharides[12,13]. In most Siglecs, the loss of the conserved Arg in the V-set domain results in loss of binding, except in Siglec-6, where binding is only weakened by the loss of the canonical Arg, as demonstrated in recent work by some of us[14]. Earlier work indicates that MAG and Siglec-11[15–18] can also operate independently of the canonical Arg, suggesting an evolution of alternative binding mechanisms in selected members of the Siglec family. To this end, previous work[14] by some of us suggested the involvement of a non-canonical Arg in the V-set domain in the binding mechanism of Siglec-6, spearheading further investigation.

In this work, we used computational and molecular biology approaches to investigate the recognition and binding specificity of Siglec-6 for monosialylated gangliosides, namely GM1, GM2 and GM3, which are known Siglecs ligands[12,14]. We first gathered a detailed understanding of the binding mechanism of Siglec-6 at the atomistic-level of detail through extensive molecular dynamics (MD) simulations. Based on structural analysis and the results of MD simulations, we were able to exclude the role of any other Arg residues in the V-set domain of Siglec-6. Instead, we found that the interaction between the Siglec-6 and the gangliosides still involves the canonical binding site, with the conserved Arg122 coordinating the carboxylic group of the sialic acid, but not continuously. This occurs because the interaction with Arg122 is structurally stabilised and energetically supplemented by an additional contact that Siglec-6 establishes with

the lipid bilayer through residues Trp127 and Lys126, see **Figure 2.a**. Mutagenesis and cell binding assays, both on liposomes and on nanodisks, support the atomistic-scale insight, also demonstrating a unique interaction between Siglec-6 and the bilayer devoid of GM1. Additionally, we show through COIN-nMS assays that the binding of free (not membrane-linked) oligosaccharides hinges entirely on the canonical Arg122. In the following sections we present the results and discuss the biological implications of the unique specialization of Siglec-6 in the recognition and binding of membrane-bound glycan epitopes with a direct comparison to other members of the Siglecs family.

Figure 2. a. 3D structure of Siglec-6 in complex with GM1 embedded in a lipid bilayer with composition distearoylphosphatidylcholine (DSPC) 60% and cholesterol 40%. In this representative snapshot from the MD trajectory, collected at 0.715 µs from the equilibrated MD ensemble, Arg 122 is engaged in a salt bridge with the Neu5Ac of GM1. The protein is represented with cyan cartoon rendering, the GM1 with sticks and C atoms in purple, O in red and N in blue. DSPC and cholesterol are rendered with semi-transparent sticks, with C atoms in grey, O in

red, N in blue and P in yellow. Key residues are labelled with numbering corresponding to the human Siglec-6 (UniProtID O43699). **b.** Structures of GM1, GM2 and GM3 represented with the SNFG nomenclature. Labels below each structure include the oligosaccharides GlyTouCan IDs. **c.** Kernel Density Estimates (KDE) analysis of the tilt angle values measured though the 1.0 µs MD trajectories ran for isolated GM1 (light green) and GM3 (dark purple). KDE maxima are 31.49º and 40.69º measured for GM1 and GM3, respectively. **d.** 3D structure of an isolated GM1 molecule (sticks with C atoms in purple, O in red and N in blue) embedded in one of the bilayer leaflets (sticks with all atoms in grey) used to represent the axes used to measure the tilt angle (θ) indicating the orientation relative to the bilayer of the Neu5Ac and thus its accessibility. **e.** Close-up view on the Siglec-6 binding site obtained through a counterclockwise rotation of approximately 120º relative to the structure in panel a). Key residues are labelled, while the embedding of the Trp127 sidechain in the bilayer is highlighted by a surface rendering of the lipids. The terminal Gal of GM1 is highlighted with C atoms in yellow for his role in orienting the C-C' loop in the bound complex, also shown in yellow. **f.** Time evolution along the MD trajectory of the distance (Å) between the Arg122 and the Neu5Ac carboxylate group. Data points correspond to the largest distance value calculated between four pairs of atoms R122-NH1(2) and O11(12)-Neu5Ac. A distance of 5 Å was chosen as a threshold for the formation of the salt bridge. **g.** 3D structure of Siglec-6 in complex with GM2 (sticks with C atoms in green) embedded in a lipid bilayer with composition distearoylphosphatidylcholine (DSPC) 60% and cholesterol 40%. In this representative snapshot from the MD trajectory, collected at 1.0 µs from the equilibrated MD ensemble, Arg 122 is still engaged in a salt bridge with the Neu5Ac of GM2, but the Siglec-6 is not embedded in the bilayer through the Trp 127 and Lys 126 due to a conformational change of the C-C' loop (yellow cartoon section). Colouring described in the legend and rendering style as in panel a). **h.** Comparison of the canonical binding site of Siglec-6 in complex with GM1, the image on the left shows a snapshot from the MD simulation when Arg122 is engaged in the salt bridge, collected at 0.715 µs, while the image on the right shows a snapshot when the salt bridge is broken, collected at 1.028 µs. **i)** Close-up view on the Siglec-6 binding site with key residues labelled. Molecular rendering done with Visual Molecular Dynamics[19] (VMD; https://www.ks.uiuc.edu/Research/vmd/). Tilt angle analysis and other graphs done with *matplotlib* (https://matplotlib.org/) and *seaborn* libraries in python (https://seaborn.pydata.org/).

Results

Orientation and accessibility of the ganglioside epitopes in the bilayer. We used all-atom classical MD simulations to investigate the mechanism that regulates ganglioside recognition and binding in Siglec-6 and to determine its ligand preference. Earlier work by some of us[14] shows that while Siglec-6 binds GM1 in an Arg-independent manner, it does not bind GM2 or GM3, which differ from GM1 by the lack of the terminal Gal and Gal-β(1-3)-GalNAc, respectively, see **Figure 2.b**. To verify that this effect was not determined by a different degree of exposure of the sialic acid through the membrane in the different gangliosides, we ran 1 µs MD simulations of isolated GM1 and GM3, as the two extremes in the monosialylated ganglioside series, in duplicates, with one ganglioside embedded in the upper and one in the lower leaflet of the bilayer, see **Figure 2.d**. The analysis of the tilt angle (θ) describing the orientation of the Neu5Ac along the trajectory relative to an axis perpendicular to the plane of

the bilayer, see **Figure 2.c**, shows that the Neu5Ac is equally exposed in GM1 and GM3, and thus equally accessible for recognition and binding. These results are in excellent agreement with previous work[20].

Structure of the Siglec-6/GM1 complex and binding mechanism. From a representative conformer of the GM1 embedded in the membrane obtained from the MD simulations described above, we built the 3D structure of the complex with Siglec-6, with the canonical Arg122 involved in a salt bridge with the Neu5Ac of the ligand. To date, no experimental 3D structure of Siglec-6 is available in public repositories, however, the common architecture and high degree of sequence and structure homology across Siglecs support confident structure predictions by machine learning (ML) leveraging on existing structural data in the RCSB PDB. In this work we used the AlphaFold (AF)[21] model AF-O43699-F1 of Siglec-6 deposited in the EBI-EMBL AF protein structure database[22]. Structural alignment of the AF model with the high resolution crystal structure of Siglec-3 (PDB 7AW6 at 1.95 Å), which shares 60% sequence identity with Siglec-6, results in an RMSD value of 0.704 Å calculated over backbone atoms. The Siglec-6/GM1 complex was built by structural alignment of the membrane bound GM1 from the MD simulations to the Neu5Ac in the sialoside analogue bound to Siglec-3 in the crystal structure (PDB 7AW6), to which the Siglec-6 was aligned. Comparison between the resulting Siglec-6/GM1 complex and the Siglec-3/sialoside analogue complexe shows that the glycan in the sialoside analogue is rotated by approximately 180° relative to the membrane bound GM1. This suggests that free and bound sialosides could potentially bind Siglecs in different orientations, due to the C2V symmetry of the carboxylate group and the broad accessibility of the Siglecs canonical binding site, discussed further in the following sections.

The Siglec-6 sequence (453 aa) carries seven *N*-glycosylation sequons, with only one in the V-set domain at N103. Although we have no information on the glycosylation of Siglec-6 from mast cells or memory B-cells, earlier work[23] indicates that occupancy of these sites is low or partial in normal (non aberrant) syncytiotrophoblasts, while it appears to be higher in recombinant constructs (317 aa). Molecular weights of 50 kDa are reported for Siglec-6 from normal pregnancy and preeclampsia placenta cells lysates[23], which corresponds to a low level of *N*-glycosylation. In preeclampsia placental lysates, molecular weights of up to 70 kDa are found, which suggests *N*-glycosylation at potentially all seven sites of Siglec-6. Molecular weights from 57 up to 90 kDa are reported on commercial sites for recombinant products. In the absence of more detailed insight on the nature and occupancy of the sites, we ran simulations

on two 3D models of the Siglec-6/GM1 complex, one with a non-glycosylated Siglec-6 and the other with a fully glycosylated Siglec-6 as a control, with all sites occupied with a biantennary mono galactosylated complex *N*-glycan (GlyTouCan ID G99129GB; 1478.5 Da). Below we describe the results we obtained for the non-glycosylated system as they correspond more closely to the Siglec-6 expression in normal placental syncytiotrophoblasts. The results obtained for the complex with the fully *N*-glycosylated Siglec-6 are included in the Supplementary Material.

After a multi-stage structure equilibration, see Methods section in the Supplementary Material for details, the conformational dynamics of the non-glycosylated complex was analyzed for 2.4 µs MD simulation ran with a deterministic sampling scheme to explore the stability of the interactions and to monitor functionally relevant conformational changes, energetically accessible at room temperature. The results indicate that, 1) the Neu5Ac of GM1 forms a salt bridge with the canonical Arg122, and 2) this interaction is structurally and energetically supplemented by the insertion in the membrane of the Siglec-6 V-set domain through the Trp127 indole sidechain and through ancillary binding of the phospholipid heads by the adjacent Lys126, see **Figure 2.a**. Furthermore, the terminal Gal of GM1 interacts with the C-C' loop (aa 70 to 75) through hydrogen bonds with Asp70 and Glu73. In the context of this interaction network, the salt bridge between the canonical Arg122 and the Neu5Ac of GM1 appears to be energetically dispensable at room temperature. Indeed, along the MD trajectory we observe that Arg122 interrupts its contact with Neu5Ac for significant amounts of sampling time, see **Figures 2.f** and **2.h**. After these intervals, the salt bridge gets restored with no disruption of the Siglec-6 bound conformation, which is primarily stabilised by its interactions with the membrane, supported by contacts between the C-C' loop and the terminal Gal of GM1.

The structure of the ganglioside determines Siglec-6 binding selectivity. Earlier work by some of us[14] shows that Siglec-6 binds GM1 selectively when embedded in a membrane, but not GM2 or GM3. Our results show that the terminal Gal of GM1 likely contributes to the stability of the complex through interactions with the C-C' loop. To test if our binding model supports the earlier experimental data and if the interaction between the C-C' loop and the terminal Gal is indeed critical, we built the 3D structure of the Siglec-6/GM2 complex from a conformation selected from the equilibrated ensemble of the Siglec-6/GM1 complex, by removing the terminal Gal of GM1, and thus biasing the new model system towards a stable conformation. We ran two independent MD trajectories with production runs of 1.0 µs and 0.5 µs each from uncorrelated

starting structures. Both simulations show that the absence of the terminal Gal triggers a conformational change in the flexible C-C' loop, which can extend into the space that was occupied by the Gal in the complex with GM1, see **Figures 2.g** and **2.i**. This shift contributes to pulling the Ig V-set domain away from the membrane, lifting out of the bilayer the embedded Trp127 and Lys126 sidechains, ultimately compromising the stability of the complex. Because GM3 is even shorter, missing the terminal Gal-β(1-3)-GalNAc relative to GM1, we expect the same outcome.

Siglec-6 binding of ganglioside-enriched liposomes. As the MD simulations predicted Lys126 and Trp127 in mediating Siglec-6-ganglioside binding when the ganglioside is presented from a lipid bilayer, we created point K126A and W127A mutations and stably expressed these in CHO cells. Results are shown in **Figure 3**. Additionally, as we had previously demonstrated that Arg122 was not required for Siglec-6 to engage with sialosides presented from a bilayer[14], we posited that another positively charged residue in the V-set may compensate in the absence of Arg122. We analyzed the binding pocket of Siglec-6 and found that Lys129 is in relatively close spatial proximity to Arg122 and hypothesized that Lys129 may be able to compensate in the absences of Arg122. Expression of the Siglec-6 mutants were validated by flow cytometry to be expressed at similar levels as WT Siglec-6, see **Figure 3.a**. Therefore, we tested these Siglec-6-expressing cells for their ability to engage fluorescent liposomes bearing a neoglycolipid that we previously optimized as a Siglec-6 ligand[14,24], see **Figure 3.b**. In this assay, two controls were used: untransfected CHO cells and liposomes lacking the neoglycolipid. We observed a modest 25% reduction, which was statistically significant, in liposome binding to R122A, K126A, and K129A Siglec-6 cells. In line with the results from our previous study, we observed that liposome binding was only modestly decreased (~75% of WT) upon the mutation of the canonical arginine residue (Arg122), a feat unique to Siglec-6. The double mutant of R122A and K129A Siglec-6 showed a greater reduction in binding compared to each single mutant (~50% of WT) but was still above background signal, suggesting that additional interactions contribute to recognition of GM1. However, in the W127A mutant, nearly all binding to the glycolipid liposomes was lost (~5% of WT). Moreover, a near complete loss in binding was observed to the K126A W127A double mutant. These results demonstrate that Trp127 is the major driving force for the recognition of glycolipids presented from a bilayer by Siglec-6.

Figure 3. a. Expression levels of the Siglec-6 WT and mutants obtained by flow cytometry. **b.** Depiction and results of the cell assay used to assess the ability of Siglec-6 mutants to bind glycolipid liposomes using flow cytometry. **c.** Depiction and results of the ELISA approach used to assess the function of the W127A Siglec-6 Mutant. **d.** 3D structures of the Ig V-set domains of MAG (blue cartoons; PDB 2ZG3), Siglec-6 (cyan cartoons; this work) and Siglec-11 (purple cartoons; PDB AF_AFQ96RL6F1). The residues in the membrane-facing loops that could potentially interact with the bilayer are labelled and highlighted with sticks. **e.** Sequence alignment of all human Siglecs performed with Clustal Omega[25] (https://www.ebi.ac.uk/jdispatcher/msa/clustalo) shows that only Siglec-6 has a KW (or similar) combination of residues on the same loop, yet a number of other Siglecs, namely 5, 8, 10, 11,

14 and 16, have a combination of one aromatic and one positively charged residues across the two loops facing the cell membrane when binding gangliosides.

To assess whether W127A Siglec-6 perturbs the structure of Siglec-6, we tested the ability of W127A Siglec-6 to bind to glycolipids outside a bilayer in an ELISA, using conditions that we have previously optimized[26,27], see **Figure 3.c**. We observed no significant difference in binding between WT and W127A Siglec-6 to a neoglycolipid ligand developed previously[14] to engage Siglec-6. Moreover, none of the full length Siglec-6 mutants showed significantly perturbed expression levels. These results strongly suggest that W127A Siglec-6 is functional, and that the loss of binding for W127A Siglec-6 observed in the cell assay is not due to structural changes in the protein due to the mutation.

Siglec-6 binding to GM1-3 as free oligosaccharides. The characterisation of the recognition and binding mechanism of gangliosides by Siglec-6 was extended to include native MS (nMS) binding assays with GM1, GM2, and GM3 as free oligosaccharides, i.e. GM1os, GM2os and GM2os. Results from the Concentration Independent (COIN)-Catch-and-Release (CaR)-nMS assay[28] are shown in **Figure 4.a** for Siglec-6 WT and for the W127A mutant, and in **Figure 4.b** for the R122A mutant. Unlike the case of the membrane-linked gangliosides, Siglec-6 binds the monosialylated gangliosides GM1-3os with similarly low affinity. Binding of the free oligosaccharides is largely unaffected by the mutation of Trp127 (W127A), further suggesting that this mutation does not significantly perturb the structure of Siglec-6 unlike binding to glycolipids in a membrane. Binding is significantly weakened by the loss of the canonical Arg122. Taken together these results suggest that, unlike the case of membrane-linked gangliosides, binding of GM1-3os occurs primarily through the canonical Arg122. As an interesting point, the complex of CD33 with a sialoside analogue (PDB 7AW6), shows an alternative conformational arrangement of the glycan epitope relative to the conformation required for a membrane-bound ganglioside. Indeed, in this complex the reducing end (GM1-equivalent) Gal-β(1-4)-[2-aminoethyl]-Glc moiety is directed towards the C-C' loop of CD33, rather than in the opposite direction, see **Figure 4.h**.

Figure 4. **a.** Binding affinities (*K*d, mM) of the ganglioside oligosaccharides GM1-3os for Siglec-6 (Fc) WT (light blue bars) and W127A mutant (orange bars) measured in aqueous ammonium acetate (200 mM, pH 6.8, 25 ºC) by COIN-CaR-nMS. Bar plots with *seaborn* (https://seaborn.pydata.org/). **b.** Binding affinities (*K*d, mM) of the ganglioside oligosaccharides GM1-3os for Siglec-6 (Fc) R122A mutant (green bars) measured in aqueous ammonium acetate (200 mM, pH 6.8, 25 ºC) by COIN-nMS. **c – g** IMS-CaR-nMS measurements performed in negative ion mode for 0.8 µM Siglec 6 (Fc) WT with 5 µM 10% GM1/DMPC ND in 200 mM aqueous ammonium acetate solution (pH 7.4). **c.** IMS heat map (m/z versus IMS drift time *t*D) and **d.** corresponding full mass spectrum extracted from IMS heat map using Driftscope. **e.** IMS heat map and **f.** corresponding extracted CID mass spectrum showing (glyco)lipid ions

released from Siglec-6; ions with m/z 7,000±100 were isolated by quadrupole (isolation window highlighted in yellow in **e**, followed by IMS and CID; a collision energy of 100 V was applied in the Transfer region. **g.** IMS-CaR-nMS measurements for Siglec-6 (Fc) mutants (each at 0.8 μM) and 10% GM1/DMPC ND (5 μM) in 200 mM aqueous ammonium acetate solutions (pH 7.4); CID mass spectra (extracted form IMS heat maps) showing (glyco)lipid ions released from (top) Siglec-6 R122A and (bottom) Siglec-6 W127A. **h.** 3D structure of the V-set domain of CD33 in complex with a sialoside analogue (PDB 7AW6). The protein is rendered in cyan as a surface, while the sialoside is rendered with sticks with C atoms in cyan, O in red and N in blue. SNFG symbols of the sialoside (left) and of the GM1 (right) are shown to indicate the position of the glucose at the reducing end.

Siglec-6 binding to GM1-enriched lipid nanodiscs. We used Catch-and-Release nMS performed with Ion Mobility Separation (IMS-CaR-nMS) to assess the ability of Siglec-6 to bind phospholipids in addition to gangliosides. To this end, we performed experiments with Siglec-6-FcWT, and R122A and W127A mutants with a GM1 nanodisc (ND) prepared with 10% GM1 and DMPC. Shown in **Figures 4.c** and **4.d** are: representative IMS heat maps (m/z versus IMS drift time t_D) and mass spectrum measured in negative ion mode for 0.8 μM Siglec 6 WT and 5 μM 10% GM1/DMPC ND (corresponding to 100 μM GM1 and ~900 μM DMPC), respectively. Both Siglec-6 and ND appeared as broad peaks, and centred at m/z 7,000 (t_D = 12.5 to 19.0 ms) and 9,500 (t_D = 22 to 32 ms), respectively. Due to the broad m/z distributions of both Siglec-6 and ND, the free and lipid-bound Siglec-6 ions and those of the ND presumably overlapped in the mass spectrum. To address this potential issue, we used quadrupole isolation (m/z 7,000 ± 100) along with IMS to separate the Siglec-6 (and Siglec-6−lipid complexes) ions from the ND ions. Collision-induced dissociation (CID) was performed in the Transfer region (100 V collision energy) to release lipid ions exclusively bound to the protein. The IMS heat map of the IMS-CaR-nMS assay and the CID mass spectrum are shown in **Figures 4.e** and **4.f**, respectively. The CID shows the release of deprotonated GM1 ions, with deprotonated DMPC ions also detected at lower abundance. Taken together the IMS-CaR-nMS results support the binding of Siglec-6 WT to both GM1 and to phospholipids. We performed analogous binding assays on the Siglec-6 R122A and W127A mutants with GM1/DMPC ND. The CID mass spectra extracted from IMS are shown in **Figure 4.g**. The results indicate that the Siglec-6 R122A mutant binds phospholipids analogously as the WT, meanwhile Siglec-6 W127A produces no detectable lipid ions, which confirms that the loss of Trp 127 compromises lipid binding.

Siglec-6 binds liposomes and nanodisks devoid of GM1. As the MD simulations and the mutagenesis studies demonstrated, Trp127 interacts with the bilayer and is critical for the binding of membrane-bound glycolipids by Siglec-6. Therefore, we hypothesized that there is direct binding between the naked liposomes and Siglec-6 and used the cell-based assay to assess if this was the case. Indeed, WT Siglec-6+ cells showed significantly higher binding to naked liposomes compared to Siglec-6- CHO cells, see **Figure 5.a**. Moreover, the binding of naked liposomes was greatly decreased (~5-fold) to W127A Siglec-6 CHO cells. In line with the results from **Figure 3.b**, naked liposome binding was not reduced to background levels with W127A Siglec-6, suggesting that while Trp127 is the major contributor to the bilayer binding, other residues such as K126A also support the interaction. In support of the results obtained on naked liposomes, we also performed IMS-CaR-nMS measurements to test lipid binding of Siglec-6 WT and mutants on 'empty' DMPC ND (5 µM), see **Figure 5.b**. The CID mass spectra show that Siglec-6 WT binds phospholipids and that the R122A mutant retains lipid-binding. Results obtained for the Siglec-6 W127A mutant demonstrate that the loss of Trp127 compromises the interaction with the bilayer.

Figure 5 a. Binding of naked liposomes (liposomes that lack a ligand) to WT, W127A and Siglec-6- CHO cells using the cell assay. **b)** IMS-CaR-nMS measurements for Siglec 6 (Fc) WT and mutants (each at 0.8 µM) with DMPC ND (5 µM) in 200 mM aqueous ammonium acetate solutions (pH 7.4); **b.** CID mass spectra (extracted form IMS heat maps) showing lipid ions released from (top) Siglec 6 WT, (middle) Siglec 6 R122A and (bottom) Siglec 6 W127A.

Discussion

Characterising glycan-binding specificity across human Siglecs is a difficult task, hindered by their weak binding affinity, potential promiscuity, and structural plasticity facilitating both *cis* and *trans* ligand interactions. Identifying the molecular determinants regulating their glycan preference is even more complex, as it requires gathering statistically reproducible, atomistic-level information on the structure and dynamics of the Siglecs, integrated within a sufficiently-detailed 3D model of their own biological environment. In this study, we show how all-atom classical MD simulations were used to guide experiments towards the characterisation of the unique binding mechanism and glycan-specificity of human Siglec-6, an inhibitory receptor expressed in mast cells, memory B-cells and, unique to humans[29], placental syncytiotrophoblasts.

Our results show that Siglec-6 evolved the ability of recognising and binding monosialylated gangliosides not only by engaging the glycan epitope poking out of the bilayer, but also by supplementing its binding affinity through interactions with the surrounding phospholipids. As suggested by MD simulations and supported by liposome and ND binding assays, the interactions between the Siglec-6 and the cell membrane hinge on the insertion of Trp127 into the bilayer, assisted by the adjacent Lys126, which interacts with the phospholipid heads. Different examples of proteins anchoring phospholipid bilayers through exposed aromatic and charged amino acids have been described within various biological contexts, e.g. where such interactions are instrumental for the orientation of transmembrane proteins[30–33] within the bilayer, as well as for the interaction of proteins and peptides with the membrane[34–36] and for the binding of membrane-embedded ligands[35,37–39]. To our knowledge, this is the first instance in which binding to the cell membrane has been characterised as a key step in the ligand-recognition of a carbohydrate-binding protein.

Further to this, we find that the structure of the ganglioside epitope is a determinant for Siglec-6 recognition and binding, with GM1 being the only monosialylated lipid-linked ganglioside recognised within a phospholipid bilayer. As in other Siglecs[1], Siglec-6 binding specificity is controlled by the C-C' loop, which interacts with the terminal Gal in GM1. GM2 and GM3 lack this terminal Gal and Gal-β(1-3)-GalNAc motif, respectively. In a model of the Siglec-6/GM2 complex that we studied by two independent MD simulations, we see that the C-C' loop is conformationally dynamic, enabled by the absence of the terminal Gal, triggers the disruption of the contact with the membrane and ultimately promoting the dissociation of the complex.

Mutagenesis and binding assays on GM1-enriched liposomes and ND confirm that bilayer interactions mediated by Siglec-6 requires Trp127, with the support of the adjacent Lys126. As demonstrated through binding assays on 'empty' liposomes and ND, the interaction with the bilayer persists even in the absence of gangliosides. These results inform an interesting hypothesis about the ganglioside recognition strategy, whereby Siglec-6 scans the cell membrane surface through low affinity interactions with lipids, in search for higher affinity and specificity binding epitopes, such as GM1. The MD simulations of isolated GM1 and GM3 gangliosides in a model bilayer suggest that such a strategy may be the most efficient process. Indeed, while we measured that the sialic acid in both GM1 and GM3 remains accessible through the lipids, the size of the exposed epitope is rather small, making it unlikely to be able to engage the Siglec-6 V-set unless in very close proximity.

Siglec-6 binding has been shown to be independent of the canonical Arg[14], while MAG and Siglec-11 have been also reported to share such feature[15]. In agreement, our MD simulations show that the interaction between Siglec-6 and GM1 is only partially supported by the salt bridge between Arg122 and the Neu5Ac, with binding assays and mutagenesis in this and in earlier work[14] showing a reduction of the binding affinity in the R122A mutant. Sequence alignment shows that the KW motif implicated in the membrane interaction is a unique feature of Siglec-6, see **Figure 3.e**. Yet, structural analysis supported by the insight we gathered from the MD simulations, show that contacts with the membrane could potentially involve two loops, see **Figure 3.d.** Other Siglecs, namely 5, 8, 10, 11, 14, and 16, carry at least one aromatic and one positively charged residue (at physiological pH) on either loop. Although we cannot exclude that membrane interactions may occur in other Siglecs through this alternative 'two-loop' strategy, some of us showed through MD simulations that Siglec-10, which has a particularly extensive combinations of 'ideal' membrane-binding residues across the two loops, namely Arg47 and Trp50 on one, and Arg122 and Tyr125 on the other, see **Figure 3.e,** does not engage with the lipids in the bilayer[40]. This is not an entirely surprising result, as Siglec-10 does not need to bind the cell membrane because its binding specificity is for polysialylated epitopes in glycoproteins or on secreted glycans[40–43]. Siglec-11 does not need to bind the membrane either, as it does not bind gangliosides[14] but α(2-8) polysialic acids, which have a rigid architecture and terminate the arms of *N*-glycans[44]. Earlier work[14] indicates that MAG binds GM1 only weakly, but not GM2/3, and that it has a higher affinity of polysialylated gangliosides; this interaction appears to be dependent on its canonical Arg118. Ultimately, membrane binding appears, thus far, to be a feature unique to Siglec-6. Yet, the broad accessibility of the binding site in the terminal V-set

domain and the conservation of the salt-bridging canonical Arg, indicates that Siglecs are able to bind free sialosides within a varying range of low binding affinities. Indeed, contrary to the case of membrane-bound gangliosides, here we have shown through nMS binding assays that Siglec-6 binds GM1-3 as free oligosaccharides (GM1-3os) with similarly low binding affinities with a mechanism that is entirely dependent on the canonical Arg122. Based on the crystal structure of the highly homologous Siglec-3 in complex with a sialoside analogue (PDB 7AW6), the relative orientation of the bound glycan epitope could also change to complement different (less constrained) environmental conditions.

Conclusions

In this work we show through a combination of computational and experimental methods how the recognition and binding of membrane-bound gangliosides by Siglec-6 hinges on its interaction with the cell membrane. The insertion of a Trp127 indole sidechain within the bilayer, assisted by the electrostatic interaction between Lys126 and the phospholipid heads, complements the ligand binding energetics, making the canonical Arg122 dispensable. Furthermore, the interaction with the membrane persists in the absence of the ligands, suggesting its involvement in recognition. These results corroborate a unique and distinctive binding mechanism adopted by Siglec-6, supporting the evolution of many different human Siglecs as molecular precision tools for the recognition of specific sialosides in their distinct biological environment. Further studies in this area are currently in progress.

Acknowledgements

IT Solutions at the University of Southampton is gratefully acknowledged for the generous allocation of computational resources on the HPC cluster Iridis. The Science Foundation of Ireland (SFI) Frontiers for the Future Programme is gratefully acknowledged for financial support of SDA and OS postgraduate training (20/FFP-P/8809). The opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the Science Foundation of Ireland. MSM thanks NSERC (RGPIN-2018-03815) and GlycoNet (CR-03) for funding. JSK acknowledges GlycoNet Integrated Services (CFI-MSI) for generous support and LKM is grateful for financial support through the Canada Excellence Research Chairs program.

References

(1) Crocker, P. R.; Paulson, J. C.; Varki, A. Siglecs and Their Roles in the Immune System. *Nat Rev Immunol* **2007**, *7* (4), 255–266.

(2) Läubli, H.; Varki, A. Sialic Acid-Binding Immunoglobulin-like Lectins (Siglecs) Detect Self-Associated Molecular Patterns to Regulate Immune Responses. *Cell. Mol. Life Sci.* **2020**, *77* (4), 593–605.

(3) Macauley, M. S.; Crocker, P. R.; Paulson, J. C. Siglec-Mediated Regulation of Immune Cell Function in Disease. *Nat. Rev. Immunol.* **2014**, *14* (10), 653–666.

(4) Pillai, S.; Netravali, I. A.; Cariappa, A.; Mattoo, H. Siglecs and Immune Regulation. *Annu. Rev. Immunol.* **2012**, *30* (1), 357–392.

(5) Duan, S.; Paulson, J. C. Siglecs as Immune Cell Checkpoints in Disease. *Annu. Rev. Immunol.* **2020**, *38* (1), 365–395.

(6) Jetani, H.; Navarro-Bailón, A.; Maucher, M.; Frenz, S.; Verbruggen, C.; Yeguas, A.; Vidriales, M. B.; González, M.; Rial Saborido, J.; Kraus, S.; Mestermann, K.; Thomas, S.; Bonig, H.; Luu, M.; Monjezi, R.; Mougiakakos, D.; Sauer, M.; Einsele, H.; Hudecek, M. Siglec-6 Is a Novel Target for CAR T-Cell Therapy in Acute Myeloid Leukemia. *Blood* **2021**, *138* (19), 1830–1842.

(7) Fry, T. J.; Shah, N. N.; Orentas, R. J.; Stetler-Stevenson, M.; Yuan, C. M.; Ramakrishna, S.; Wolters, P.; Martin, S.; Delbrook, C.; Yates, B.; Shalabi, H.; Fountaine, T. J.; Shern, J. F.; Majzner, R. G.; Stroncek, D. F.; Sabatino, M.; Feng, Y.; Dimitrov, D. S.; Zhang, L.; Nguyen, S.; Qin, H.; Dropulic, B.; Lee, D. W.; Mackall, C. L. CD22-Targeted CAR T Cells Induce Remission in B-ALL That Is Naive or Resistant to CD19-Targeted CAR Immunotherapy. *Nat. Med.* **2018**, *24* (1), 20–28.

(8) Kenderian, S. S.; Ruella, M.; Shestova, O.; Klichinsky, M.; Aikawa, V.; Morrissette, J. J. D.; Scholler, J.; Song, D.; Porter, D. L.; Carroll, M.; June, C. H.; Gill, S. CD33-Specific Chimeric Antigen Receptor T Cells Exhibit Potent Preclinical Activity against Human Acute Myeloid Leukemia. *Leukemia* **2015**, *29* (8), 1637–1647.

(9) Angata, T.; Varki, A. Discovery, Classification, Evolution and Diversity of Siglecs. *Mol. Aspects Med.* **2023**, *90* (101117), 101117.

(10) Bornhöfft, K. F.; Goldammer, T.; Rebl, A.; Galuska, S. P. Siglecs: A Journey through the Evolution of Sialic Acid-Binding Immunoglobulin-Type Lectins. *Dev. Comp. Immunol.* **2018**, *86*, 219–231.

(11) Yamaji, T.; Teranishi, T.; Alphey, M. S.; Crocker, P. R.; Hashimoto, Y. A Small Region of the Natural Killer Cell Receptor, Siglec-7, Is Responsible for Its Preferred Binding to Alpha 2,8-Disialyl and Branched Alpha 2,6-Sialyl Residues. A Comparison with Siglec-9. *J. Biol. Chem.* **2002**, *277* (8), 6324–6332.

(12) Schnaar, R. L. Gangliosides as Siglec Ligands. *Glycoconj. J.* **2023**, *40* (2), 159–167.

(13) Gonzalez-Gil, A.; Schnaar, R. L. Siglec Ligands. *Cells* **2021**, *10* (5), 1260.

(14) Schmidt, E. N.; Lamprinaki, D.; McCord, K. A.; Joe, M.; Sojitra, M.; Waldow, A.; Nguyen, J.; Monyror, J.; Kitova, E. N.; Mozaneh, F.; Guo, X. Y.; Jung, J.; Enterina, J. R.; Daskhan, G. C.; Han, L.; Krysler, A. R.; Cromwell, C. R.; Hubbard, B. P.; West, L. J.; Kulka, M.; Sipione, S.; Klassen, J. S.; Derda, R.; Lowary, T. L.; Mahal, L. K.; Riddell, M. R.; Macauley, M. S. Siglec-6 Mediates the Uptake of Extracellular Vesicles through a Noncanonical Glycolipid Binding Pocket. *Nat. Commun.* **2023**, *14* (1), 2327.

(15) Varki, A.; Angata, T. Siglecs--the Major Subfamily of I-Type Lectins. *Glycobiology* **2006**, *16* (1), 1R – 27R.

(16) Vinson, M.; Strijbos, P. J.; Rowles, A.; Facci, L.; Moore, S. E.; Simmons, D. L.; Walsh, F. S. Myelin-Associated Glycoprotein Interacts with Ganglioside GT1b. A Mechanism for Neurite Outgrowth Inhibition. *J. Biol. Chem.* **2001**, *276* (23), 20280–20285.

(17) Angata, T.; Kerr, S. C.; Greaves, D. R.; Varki, N. M.; Crocker, P. R.; Varki, A. Cloning and Characterization of Human Siglec-11. *J. Biol. Chem.* **2002**, *277* (27), 24466–24474.

(18) Tang, S.; Shen, Y. J.; DeBellard, M. E.; Mukhopadhyay, G.; Salzer, J. L.; Crocker, P. R.; Filbin, M. T. Myelin-Associated Glycoprotein Interacts with Neurons via a Sialic Acid Binding Site at ARG118 and a Distinct Neurite Inhibition Site. *J. Cell Biol.* **1997**, *138* (6), 1355–1366.

(19) Humphrey, W.; Dalke, A.; Schulten, K. VMD: Visual Molecular Dynamics. *J. Mol. Graph.* **1996**, *14* (1), 33–38, 27–28.

(20) Patel, D. S.; Park, S.; Wu, E. L.; Yeom, M. S.; Widmalm, G.; Klauda, J. B.; Im, W. Influence of Ganglioside GM1 Concentration on Lipid Clustering and Membrane Properties and Curvature. *Biophys. J.* **2016**, *111* (9), 1987–1999.

(21) Jumper, J.; Evans, R.; Pritzel, A.; Green, T.; Figurnov, M.; Ronneberger, O.; Tunyasuvunakool, K.; Bates, R.; Žídek, A.; Potapenko, A.; Bridgland, A.; Meyer, C.; Kohl, S. A. A.; Ballard, A. J.; Cowie, A.; Romera-Paredes, B.; Nikolov, S.; Jain, R.; Adler, J.; Back, T.; Petersen, S.; Reiman, D.; Clancy, E.; Zielinski, M.; Steinegger, M.; Pacholska, M.; Berghammer, T.; Bodenstein, S.; Silver, D.; Vinyals, O.; Senior, A. W.; Kavukcuoglu, K.; Kohli, P.; Hassabis, D. Highly Accurate Protein Structure Prediction with AlphaFold. *Nature* **2021**. https://doi.org/10.1038/s41586-021-03819-2.

(22) Varadi, M.; Anyango, S.; Deshpande, M.; Nair, S.; Natassia, C.; Yordanova, G.; Yuan, D.; Stroe, O.; Wood, G.; Laydon, A.; Žídek, A.; Green, T.; Tunyasuvunakool, K.; Petersen, S.; Jumper, J.; Clancy, E.; Green, R.; Vora, A.; Lutfi, M.; Figurnov, M.; Cowie, A.; Hobbs, N.; Kohli, P.; Kleywegt, G.; Birney, E.; Hassabis, D.; Velankar, S. AlphaFold Protein Structure Database: Massively Expanding the Structural Coverage of Protein-Sequence Space with High-Accuracy Models. *Nucleic Acids Res.* **2022**, *50* (D1), D439–D444.

(23) Awoyemi, T.; Tannetta, D.; Zhang, W.; Kandzija, N.; Motta-Mejia, C.; Fischer, R.; Heilig, R.; Raiss, S.; Redman, C.; Vatish, M. Glycosylated Siglec-6 Expression in Syncytiotrophoblast-Derived Extracellular Vesicles from Preeclampsia Placentas. *Biochem. Biophys. Res. Commun.* **2020**, *533* (4), 838–844.

(24) Schmidt, E. N.; Guo, X. Y.; Bui, D. T.; Jung, J.; Klassen, J. S.; Macauley, M. S. Dissecting the Abilities of Murine Siglecs to Interact with Gangliosides. *J. Biol. Chem.* **2024**, *300* (7), 107482.

(25) Sievers, F.; Wilm, A.; Dineen, D.; Gibson, T. J.; Karplus, K.; Li, W.; Lopez, R.; McWilliam, H.; Remmert, M.; Söding, J.; Thompson, J. D.; Higgins, D. G. Fast, Scalable Generation of High-Quality Protein Multiple Sequence Alignments Using Clustal Omega. *Mol. Syst. Biol.* **2011**, *7*, 539.

(26) Rodrigues, E.; Jung, J.; Park, H.; Loo, C.; Soukhtehzari, S.; Kitova, E. N.; Mozaneh, F.; Daskhan, G.; Aghanya, V.; Sarkar, S.; Streith, L.; Laurent, C. S.; Julien, J.-P.; West, L. J.; Williams, K. C.; Klassen, J. S.; Macauley, M. S. A Versatile Soluble Siglec Scaffold for Sensitive and Quantitative Detection of Glycan Ligands. *bioRxiv*, 2020. https://doi.org/10.1101/2020.07.28.223164.

(27) Schmidt, E. N.; Jung, J.; Macauley, M. S. Flow Cytometry-Based Detection of Siglec Ligands. *Methods Mol. Biol.* **2023**, *2657*, 181–193.

(28) Bui, D. T.; Favell, J.; Kitova, E. N.; Li, Z.; McCord, K. A.; Schmidt, E. N.; Mozaneh, F.; Elaish, M.; El-Hawiet, A.; St-Pierre, Y.; Hobman, T. C.; Macauley, M. S.; Mahal, L. K.; Flynn, M. R.; Klassen, J. S. Absolute Affinities from Quantitative Shotgun Glycomics Using Concentration-Independent (COIN) Native Mass Spectrometry. *ACS Cent. Sci.* **2023**, *9* (7), 1374–1387.

(29) Brinkman-Van der Linden, E. C. M.; Hurtado-Ziola, N.; Hayakawa, T.; Wiggleton, L.; Benirschke, K.; Varki, A.; Varki, N. Human-Specific Expression of Siglec-6 in the Placenta. *Glycobiology* **2007**, *17* (9), 922–931.

(30) Vostrikov, V. V.; Daily, A. E.; Greathouse, D. V.; Koeppe, R. E., 2nd. Charged or Aromatic

Anchor Residue Dependence of Transmembrane Peptide Tilt. *J. Biol. Chem.* **2010**, *285* (41), 31723–31730.

(31) Sustich, S. J.; Afrose, F.; Greathouse, D. V.; Koeppe, R. E., 2nd. Influence of Interfacial Tryptophan Residues on an Arginine-Flanked Transmembrane Helix. *Biochim. Biophys. Acta Biomembr.* **2020**, *1862* (2), 183134.

(32) Wimley, W. C.; White, S. H. Experimentally Determined Hydrophobicity Scale for Proteins at Membrane Interfaces. *Nat. Struct. Biol.* **1996**, *3* (10), 842–848.

(33) Frazier, A. A.; Wisner, M. A.; Malmberg, N. J.; Victor, K. G.; Fanucci, G. E.; Nalefski, E. A.; Falke, J. J.; Cafiso, D. S. Membrane Orientation and Position of the C2 Domain from cPLA2 by Site-Directed Spin Labeling. *Biochemistry* **2002**, *41* (20), 6282–6292.

(34) Christiaens, B.; Symoens, S.; Verheyden, S.; Engelborghs, Y.; Joliot, A.; Prochiantz, A.; Vandekerckhove, J.; Rosseneu, M.; Vanloo, B. Tryptophan Fluorescence Study of the Interaction of Penetratin Peptides with Model Membranes: Penetratin: Interaction with Model Membranes. *Eur. J. Biochem.* **2002**, *269* (12), 2918–2926.

(35) Barkdull, A. P.; Holcomb, M.; Forli, S. A Quantitative Analysis of Ligand Binding at the Protein-Lipid Bilayer Interface. *Commun. Chem.* **2025**, *8* (1), 89.

(36) Suárez, T.; Gallaher, W. R.; Agirre, A.; Goñi, F. M.; Nieva, J. L. Membrane Interface-Interacting Sequences within the Ectodomain of the Human Immunodeficiency Virus Type 1 Envelope Glycoprotein: Putative Role during Viral Fusion. *J. Virol.* **2000**, *74* (17), 8038–8047.

(37) Zhang, W.; Crocker, E.; McLaughlin, S.; Smith, S. O. Binding of Peptides with Basic and Aromatic Residues to Bilayer Membranes: Phenylalanine in the Myristoylated Alanine-Rich C Kinase Substrate Effector Domain Penetrates into the Hydrophobic Core of the Bilayer. *J. Biol. Chem.* **2003**, *278* (24), 21459–21466.

(38) Stahelin, R. V.; Long, F.; Diraviyam, K.; Bruzik, K. S.; Murray, D.; Cho, W. Phosphatidylinositol 3-Phosphate Induces the Membrane Penetration of the FYVE Domains of Vps27p and Hrs. *J. Biol. Chem.* **2002**, *277* (29), 26379–26388.

(39) Bazzi, M. D.; Nelsestuen, G. L. Association of Protein Kinase C with Phospholipid Monolayers: Two-Stage Irreversible Binding. *Biochemistry* **1988**, *27* (18), 6776–6783.

(40) Sobczak, K.; Antoñana-Vildosola, A.; Valverde, P.; Travecedo, M. A.; Jame-Chernaboo, Z.; Schmidt, E. N.; D'Andrea, S.; Valdaliso-Díez, E.; Oyenarte, I.; Laugieri, M. E.; Joe, M.; Mozaneh, F.; Lin, S.-Y.; Bosch, A.; Moure, M. J.; Franconetti, A.; Lee, S. Y.; de Durana, J. E.-D.; Pérez-Gutierrez, L.; Palazón, A.; Marcelo, F.; Fadda, E.; Corzana, F.; Gimeno, A.; Macauley, M. S.; Jiménez-Barbero, J.; Ereño-Orbea, J. The Unique Molecular Recognition Features of Siglec-10: Structural Insights into Sialoglycan and Antibody Interactions. *bioRxiv*, 2025. https://doi.org/10.1101/2025.06.10.658867.

(41) Boelaars, K.; van Kooyk, Y. Targeting Myeloid Cells for Cancer Immunotherapy: Siglec-7/9/10/15 and Their Ligands. *Trends Cancer* **2024**, *10* (3), 230–241.

(42) Bandala-Sanchez, E.; Zhang, Y.; Reinwald, S.; Dromey, J. A.; Lee, B.-H.; Qian, J.; Böhmer, R. M.; Harrison, L. C. T Cell Regulation Mediated by Interaction of Soluble CD52 with the Inhibitory Receptor Siglec-10. *Nat. Immunol.* **2013**, *14* (7), 741–748.

(43) DeBono, N. J.; D'Andrea, S.; Bandala-Sanchez, E.; Goddard-Borger, E.; Zenaidee, M. A.; Moh, E. S. X.; Fadda, E.; Harrison, L. C.; Packer, N. H. The Molecular Basis of Immunosuppression by Soluble CD52 Is Defined by Interactions of N-Linked and O-Linked Glycans with HMGB1 Box B. *J. Biol. Chem.* **2025**, *301* (4), 108350.

(44) Schnaar, R. L.; Gerardy-Schahn, R.; Hildebrandt, H. Sialic Acids in the Brain: Gangliosides and Polysialic Acid in Nervous System Development, Stability, Disease, and Regeneration. *Physiol. Rev.* **2014**, *94* (2), 461–518.