Vocal convergence during formation of social relationships in vampire bats

Julia Vrtilek*[1], Grace Smith-Vidaurre*[2,3,4,5], Eric Fosler-Lussier[6], Rachel A. Page[7], Gerald G. Carter[7,8,9]

Affiliations
1. Department of Evolution, Ecology and Organismal Biology; The Ohio State University, Columbus, Ohio, USA
2. Department of Biology, New Mexico State University, Las Cruces, NM, USA
3. Department of Integrative Biology; Michigan State University, East Lansing, MI, USA
4. Department of Computational Mathematics, Science, and Engineering; Michigan State University, East Lansing, MI, USA
5. Ecology, Evolution, and Behavior Program; Michigan State University, East Lansing, MI, USA
6. Department of Computer Science and Engineering; The Ohio State University, Columbus, Ohio, USA
7. Smithsonian Tropical Research Institute, Balboa, Ancón, Panamá
8. Department of Ecology and Evolutionary Biology; Princeton University, Princeton, New Jersey, USA
9. Howard Hughes Medical Institute, Chevy Chase, Maryland, USA

*These authors contributed equally to this work

Corresponding authors: Julia Vrtilek: jkvrtilek@gmail.com; Gerald Carter: gc1511@princeton.edu

Keywords: contact calls, social bonds, social relationships, vampire bats, vocal convergence, vocal learning

Abstract

In many group-living birds and mammals, the formation of affiliative relationships is hypothesized to cause vocal convergence (an increase in call similarity between individuals). However, testing this causal effect can be difficult, because it requires experimentally forming new relationships. Here, we demonstrate convergence in the contact calls of common vampire bats (*Desmodus rotundus*) that we introduced and experimentally housed together. To estimate and disentangle the roles of kinship, co-housing (familiarity), allogrooming, and food sharing in predicting call similarity, we first measured call similarity using 35 features of 693,494 contact calls from 95 bats, then fit a series of Bayesian generalized multi-membership models. We also measured changes in call similarity for a subset of individuals that were recorded before and after co-housing. We found that co-housing caused vocal convergence. Furthermore, food-sharing rates among familiar nonkin within the same co-housed group predicted contact call similarity. This finding suggests that the development of cooperative relationships causes further vocal convergence beyond the initial convergence caused by co-housing. Our findings have implications for the development of cooperative relationships and vocal learning.

Introduction

Vocal behaviour provides an important window into the social lives of many birds and mammals. Exchanges of vocalizations called "contact calls" seem to facilitate recognition, reunions, and collective movements among affiliated individuals across several species (e.g. African elephants (1), common marmosets (2), and white-winged vampire bats (3)). In many group-living birds and mammals, the contact calls of unrelated members of the same group sound more similar than those of strangers (e.g. black-capped chickadees (4), orange-fronted conures (5), budgerigars (6), yellow-naped amazons (7), pygmy and common marmosets (8–10), Guinea baboons (11), grey mouse lemurs (12), gelada (13), African elephants (14), bottlenose dolphins (15,16), orcas (17), greater spear-nosed bats (18) and Taiwanese leaf-nosed bats (19)). Based on this observation, a longstanding hypothesis is that the formation of affiliative relationships ("social bonds") can increase similarity in the acoustic structure of calls between individuals ("vocal convergence") (20).

Although vocal similarity or convergence is often associated with affiliative relationships (10,18,19,21–25), there are multiple possible mechanisms for how and why vocal convergence occurs. First, familiar animals might have the ability to modify the structure of vocalizations based on auditory inputs (vocal production learning), causing similarity between familiar individuals (e.g. 4). However, vocal convergence exists in many mammalian lineages where vocal production learning has not yet been demonstrated (26). In these cases, it is possible that animals might learn from conspecifics to produce existing sounds in their repertoire in new contexts (usage learning). Finally, familiar animals might use the same vocalizations more often simply because their physiological or motivational states are often matching (26).

Regardless of the underlying mechanism, demonstrating a causal link from affiliation to vocal convergence can be difficult. For example, correlations between affiliation and vocal similarity can result from kinship, either through heritable variation or shared environment. The same correlation can result from pre-existing vocal similarity causing higher affiliation (assortment based on acoustic similarity). To demonstrate causation, an ideal study would induce new relationships to form among nonkin, then observe changes in call similarity (e.g. 4).

Here, we applied this approach to demonstrate convergence in the contact calls of common vampire bats (*Desmodus rotundus*). Vampire bats produce contact calls when they are socially isolated (27) and use the calls to vocally recognize affiliated individuals (28). By analysing 693,494 contact calls from 95 bats, some of which were experimentally co-housed together, we found that co-housing causes vocal convergence. Next, to tease apart how pairwise call similarity was shaped by relationships within each captive group, we estimated effects of genetic relatedness, allogrooming, and food sharing (29–32). Both kin and nonkin females can develop reciprocal allogrooming relationships that can become reciprocal food-sharing relationships, in which a fed bat regurgitates potentially life-saving ingested blood to a starving partner (29–31). We found evidence that these food-sharing relationships cause further vocal convergence between nonkin after the initial convergence caused by co-housing.

Methods

Study subjects

The 95 common vampire bats (*Desmodus rotundus*) used in this study came from 8 different origins (e.g. captive colonies or field sites; for details, see Supplementary Information (SI), Table S1). To identify individual bats, all bats were marked using 1-4 unique bat bands. We labelled bats as "adults" if they were older than 2 years old at time of recording or, if their exact age was unknown, by the presence or absence of an epiphyseal gap between their metacarpals. We refer to volant non-adults as "subadult". Non-volant pups that were still nursing from their mother were not included in the study.

Kinship, social familiarity, and cooperative relationships

Bats captured from different sites hundreds of kilometres apart were considered both unrelated (non-kin) and unfamiliar. Kinship of bats from the same colony was estimated through a combination of microsatellite markers and maternal pedigrees, as described in previous studies (29,31). Pairs of bats with unknown kinship were excluded from analyses where kinship was a predictor.

Most pairs (n = 2605) were unfamiliar because they had never met; 582 pairs were captured from different field sites hundreds of km apart, then introduced in captivity and caged together (co-housed) for at least 4 months; 612 pairs were captured from the same field site then caged together for at least 4 months; and 435 pairs were living in captivity for years. Another 231 pairs had unknown levels of kinship and familiarity, because they were captured from the same wild roost about four years apart. Within each co-housed captive group, pairs had variable rates of cooperative behaviour. We had measures of food-sharing rates for 307 pairs, allogrooming rates for 190 pairs, and both measures for 406 pairs. These data were collected from a series of past studies where experimenters repeatedly induced food sharing by removing a bat, fasting it for over 24 hours, and reintroducing it to its home cage, then used focal follow sampling to observe allogrooming, regurgitated food sharing, or both (29–31).

Recording contact calls from individually isolated bats

Contact calls are frequency modulated syllables that are produced by socially isolated bats (27). To record contact calls, we isolated an individual bat inside a plastic storage bin (about 75 to 200 litres in volume) lined with acoustic foam and/or soft fabric to dampen echoes. To keep the bat within 10 to 30 cm of the microphone, we placed it inside a small soft-mesh butterfly cage or a tube of plastic mesh. Calls were recorded with an Avisoft CM16 ultrasound condenser microphone (frequency range 10-200 kHz, purchased prior to 2017 from Avisoft Bioacoustics, Berlin, Germany). Sound was digitised with 16-bit resolution at sampling rates ranging between 250-500 kHz through an Avisoft UltraSoundGate (one-channel or four-channel) to a laptop running the program Avisoft Recorder. To remove any effects of digitisation rate on subsequent call measures, recordings digitised at a rate above 250 kHz were downsampled to 250 kHz using the *seewave* R package (33). To filter out sounds that were not contact calls, we removed calls with durations less than 3 ms or longer than 50 ms and with peak

frequencies below 10 kHz. For further explanation of call selection, see SI,
Supplementary Methods.
We analysed 693,494 contact calls from 95 of the 115 common vampire bats that
were individually isolated and recorded from 2011 to 2019. We only analysed the 95
bats that produced at least 100 contact calls of sufficient quality to measure acoustic
features (described below). The number of recording sessions per bat varied from 1 to
22 (mean = 7.36). The mean number of calls recorded per bat was 7300 (median =
2500, minimum = 118, maximum = 98161).
142
Measuring acoustic features of calls
We wrote a custom R script to identify start and end times of individual calls
using both amplitude and spectral density (the distribution of amplitude across
frequencies). This automated method was tested and validated with 1535 manually
labelled recordings containing ~7000 call selections. Using these start and end times,
we then measured 35 features of each call (Figure 1, Table S2). We used the R
package *warbleR* (34) to measure 27 spectral and temporal parameters and the R
package *soundgen* (35) to estimate 8 additional measurements from the fundamental:
maximum slope, minimum slope, and absolute minimum slope; the number of positive
slopes and of negative slopes; the number of turns; the mean slope; and the number of
segments. We eliminated 'clipped' calls, which we identified as calls that reached a
relative amplitude of more than 0.99 using the *normalize* function from the R package
tuneR (36). All scripts were run using the Ohio Supercomputer (37).
156

157
Figure 1. Vampire bat contact call. Call measures shown here include: 1 duration, 2
time quantile 25, 3 time median, 4 time quantile 75, 5 mean frequency, 6 frequency
quantile 25, 7 frequency median, 8 frequency quantile 75, 9 maximum dominant
frequency, 10 mean dominant frequency, 11 minimum dominant frequency, 12 start
dominant frequency, 13 end dominant frequency, 14 peak frequency. Table S2 defines
the above measures and 21 additional measures not shown here.
164

Assessing vocal individuality
To assess and confirm baseline individuality in contact call structure, we
estimated the classification rate for assigning calls to the correct caller using a linear
discriminant function analysis (DFA) with leave-one-out cross-validation, including all
variables measured (via the MASS R package (38)). To choose a minimum number of
calls per bat, we compared these classification rates to the same rates from a single
DFA without cross-validation. We excluded bats with fewer than 100 sampled calls
because within-bat classification rates were virtually identical for bats with greater than
100 calls. To confirm that calls could be assigned to the caller with greater-than-chance
accuracy, we ran a permutation test: we repeated a DFA without cross-validation 1000
times with randomized bat names to generate an expected distribution of accuracy rates
under the null hypothesis that calls contain no information about caller identity.
177
178 *Estimating contact call similarity*
179   We defined contact call similarity for each pair of bats as:
180
181 $$S = 1 - \left(\frac{D}{maximum\ (D)}\right) \tag{1}$$
182
where D is the Mahalanobis distance between the two group centroids (bats) from a
DFA classifying calls (n = 693,494) to bat (n = 95). These Mahalanobis distances
represent the average multivariate distance (or dissimilarity) between every two bats.
We normalized our measure of call similarity to range from 0 to 1; however, we replaced
the single similarity score of 0 with 0.001 (the next lowest call similarity was 0.03),
because similarity cannot be precisely zero and beta regression cannot model an
outcome of zero.
190
Estimating effects of kinship and social relationship on call similarity
To assess predictors of contact call similarity while controlling for the effect of
both bat identities, we fit five Bayesian generalized (beta distribution) multi-membership
models (Table 1) using the R package *brms* (39). Multi-membership models are used to
model dyadic data by fitting group-structured random effects, which capture the
dependence of each pair on the identities of both individuals. This model structure
allows us to estimate a random intercept for each bat and its influence on its pairings
with other bats (40). To assess causal hypotheses, we estimated multiple predictors
across seven models based on known and hypothesized causal paths (Figure 2a).
Our first goal was to disentangle the effects of kinship from social familiarity. In
model 1, we estimated how well kinship predicted contact call similarity across the 2179
pairs of adult bats with known kinship estimates. Kinship is confounded with group-level
familiarity and within-group affiliation rates, and any effects of kinship in model 1 could
therefore be driven by social interaction history rather than kinship. In model 2, we
therefore estimated how well kinship predicted contact call similarity when conditioning
on (i.e. statistically controlling for) affiliation rates in the same 2179 pairs. Predictors
were kinship, whether the bats were co-housed (true/false), and within-group affiliation
lograte, defined as:
209

$$\frac{log(observed + 1)}{log(sampling)}$$ (2)

where "observed" was the seconds of either allogrooming or regurgitated food sharing
between the two bats and "sampling" was the total seconds in which allogrooming or
food sharing could have been observed between the two bats. Both allogrooming and
food sharing are directed behaviours that indicate an affiliative social preference. We
used lograyes rather than rates because seconds of allogrooming and food sharing are
lognormal. We scaled (i.e. z-transformed) both kinship and affiliation lograte to convert
units to standard deviations.
Another way to control for kinship is to only test unrelated pairs. In model 3, we
measured how well being caged together (true/false) predicted contact call similarity in
the 1159 pairs of unrelated female adults caught from different sites. All pairs had
kinship estimates of 0, allowing us to test for an effect of nonkin being caged together
on their contact call similarity. We focused on female pairs because the few males in
these colonies were sub-adults or young born into the group.
Our next goal was to separate the effects of group-level familiarity and dyad-level
affiliation rate among nonkin females. In model 4, we measured how well within-group
affiliation lograte (scaled lograte) predicted contact call similarity in 272 unrelated pairs
of female adults that were caged together. We removed pairs with kinship estimates
greater than 0.05 to test for an effect of nonkin social interactions on contact call
similarity. We also removed pairs that could never have been in contact. We again
focused on female pairs because social affiliation data were focused on females and
undersampled or missing for males (29–32). We did not include allogrooming and
allofeeding together as covariates in model 4 because they are highly correlated proxies
for social affiliation, rather than different effects. The affiliation rate therefore includes
either allogrooming, regurgitated food sharing, or both. The results of model 4 must
therefore be interpreted carefully, because affiliation rate is based more on allogrooming
in some pairs than others. For example, in one colony from 2010-2014, food-sharing
rates were sampled more than allogrooming, but in another colony in 2019, food-
sharing rates were unknown (missing) for all pairs. Moreover, food sharing represents a
much higher level of affiliation than allogrooming. For example, when bats are
introduced in captivity (i.e. become familiar), most pairs form allogrooming relationships
and only a few begin food-sharing relationships (29). In other words, co-housing predicts
allogrooming better than food sharing.
To remedy these issues, we used our last model to focus on food-sharing rates,
which is the best indication of a within-group social bond (29). In model 5, we measured
how well food-sharing rate (scaled lograte) predicted contact call similarity in the 139
unrelated pairs of female adults that were caged together. We again removed pairs with
kinship estimates greater than 0.05, pairs that could never have shared food and pairs
with unknown food-sharing rates.
Across all models, we used default priors and scaled continuous predictors. We
used posterior predictive checks to assess model fit (Figure S2). To ensure Markov
Chain Monte Carlo (MCMC) convergence, we checked the R-hat values across 4
chains, which ranged from 0.9999 to 1.0004, showing that all 4 chains reached the
same estimates for coefficients. We used chain and warmup lengths of 6000 and 2000

samples, respectively. The lowest Bulk and Tail effective sample sizes across all
models were 8747 and 10064 respectively.
257

Model	Question addressed	Covariates included as predictors of contact call similarity	Bat pairs in sample
1	How well does kinship predict contact call similarity?	kinship	pairs with estimated kinship = 2179
2	How well does kinship predict contact call similarity after controlling for co-housing and affiliation?	kinship, co-housing (true/false), affiliation rate (including zeros for bats that never met)	pairs with estimated kinship = 2179
3	In nonkin females, how well does co-housing predict contact call similarity?	co-housing (true/false)	pairs of nonkin adult females = 1159
4	In nonkin females, how well does affiliation predict contact call similarity?	within-group affiliation rate (excluding bats that never met)	pairs of nonkin adult females that were caged together = 272
5	In nonkin females, how well does food sharing predict contact call similarity?	within-group food-sharing rate (excluding bats that never met)	pairs of nonkin adult females with estimated food-sharing rates = 139

258 **Table 1. Purpose of each statistical model.** We selected this set of models based on
259 causal models shown in Figure 2a.
260
Comparing call similarity pre-introduction and post-introduction
We could determine how call similarity changed over time using a subset of
cases in 2016 where 7 bats from Tolé were recorded both before and after being
introduced to 5 bats from Las Pavas (Figure S3). We predicted that contact call
convergence (increases in call similarity) should be greater for introduced pairs (from
different sites) than for previously familiar pairs (from the same site). To compare call
similarity before and after introduction, we first performed two discriminant function
analyses (DFAs) classifying calls to bats. Both DFAs included 5 Las Pavas bats and 7
Tolé bats, but the first DFA included recordings from the Tolé bats *before introduction* to
the Las Pavas bats, and the second DFA included the same Tolé bats recorded *after*
the introduction. From each DFA, we extracted Mahalanobis distances and call
similarities as described above. We then estimated the change in similarity as the
difference between post-introduction and pre-introduction call similarities.
To test the hypothesis that increases in similarity were greater for introduced
pairs than for familiar pairs, we used two approaches. First, we used a nonparametric

permutation test (Mantel test with Spearman correlation) of the correlation between the
change in contact call similarity and a binary familiarity matrix indicating familiar vs.
introduced pairs. Second, we used a Bayesian multi-membership model (Gaussian
distribution, default priors in brms, 4 chains with length of 6000 interactions, 2000
warmup samples) to test whether familiarity predicted the change in contact call
similarity.
Finally, we tested the prediction that call classification to the correct capture site
should be more accurate prior to the merging of the two groups. To do this, we used two
permuted DFAs, which are designed to test for correct classification of calls to social
group (41). In the first permuted DFA, we classified calls to either the Las Pavas group
or the *pre-introduction* Tolé group. In the second permuted DFA, we classified calls to
either the Las Pavas group or the *post-introduction* Tolé group.
288

Results

Contact call variation within and between individuals
Contact calls had clear individual signatures but were also highly variable within
bat (Figure S1). Discriminant function analysis (DFA) correctly assigned the 693,494
calls to the 95 callers with 36% probability, which is more than twice the expected
chance level of 14% (95% quantiles of expected values: 14.14% to 14.15%). The call
assignment accuracy of assigning calls back to individual bats from the cross-validated
DFA and from the non-cross-validated DFAs were almost perfectly correlated across
the 95 bats (r = 0.9996), indicating robustness of the DFA used for estimating call
similarity.
299

Kinship-biased contact call similarity can be explained by social familiarity
Model 1 revealed that kinship predicts contact call similarity (Figure 2b, Figure
S4). However, kinship is confounded with social affiliation in vampire bats (30,31,42). We
therefore used Model 2 to test for an effect of kinship while conditioning on (i.e.
"controlling for") affiliation rate and co-housing. We no longer found an effect of kinship
while conditioning on these measures of social affiliation, but we did find that call
similarity was predicted by both affiliation rate and co-housing while conditioning on
kinship and each other (Figure 2b). This result does not mean that kinship did not
influence call similarity; only that the correlations between kinship and contact call
similarity can also be explained by social familiarity. In other words, both affiliation rate
and familiarity predicted contact call similarity, regardless of kinship.
311

Co-housing causes contact call similarity among nonkin
To further test whether co-housing causes contact call convergence among
nonkin, we used Model 3 to focus on 1159 pairs of unrelated female adults that were
captured from different sites. Compared to model 2, which included kin, we found that
co-housing was associated with an even larger increase in call similarity among pairs of
unrelated adult females (Figure 2b). As expected, call similarity for nonkin from distant
sites that were housed together in captivity was dramatically higher than nonkin from
different sites that never met, and the highest mean similarity in nonkin contact calls

occurred within nonkin pairs from the same wild roost or in the same long-term captive
colony (Figure S5).
322
323   *Within co-housed groups, food-sharing rates predict further vocal convergence*
324   Although Model 2 revealed an effect of affiliation (allogrooming and food sharing,
Figure S6), further analyses showed that this effect was driven by food sharing rather
than allogrooming (Figure 2b, Figure S7). Specifically, Model 4 estimated the effect of
within-group dyadic affiliation rate on call similarity among unrelated female adults, and
this model found no clear effect of affiliation rate beyond the effect of previous
familiarity. However, Model 5 showed that contact call similarity was predicted by within-
group rates of regurgitated food sharing. This effect of food sharing on call similarity
was largely driven by the 2014 long-term captive colony (Figure S7), which had the
highest rates of food sharing and the longest periods of familiarity.
333

334
Figure 2. Contact call similarity in vampire bats is predicted by co-housing,
affiliation rate, and food-sharing. Panel a: causal model for vocal convergence, with
black arrows indicating causal paths of interest, and grey arrows indicating causal paths
that cause potential confounding. Path A: Greater kinship causes greater affiliation. We
estimate affiliation through allogrooming and food-sharing. Path B: Kinship might cause
greater similarity in contact calls if call features are heritable. Path C: Affiliation might
cause vocal convergence. Path D: Affiliation is caused by co-housing. Path E: Co-
housing leads to acoustic exposure, which might also cause vocal convergence
independent of variation in affiliation. **Panel b:** seven regression coefficients from five
models (colours) selected to disentangle causal effects of kinship, co-housing, affiliation
rate, and food-sharing. Dots and error bars show means and 95% credible intervals

from Bayesian posterior probability distributions. See Table S3 for coefficients and
credible intervals.
348
Vocal convergence evident from call similarity before and after co-housing
We saw evidence for vocal convergence over time (increased contact call
similarity after introduction) in 25 of 35 pairs from different sites (71%), compared to 11
of 21 familiar pairs from Tolé (52%). We found that a greater increase in call similarity
(i.e. greater vocal convergence) was positively associated with being an introduced pair
of bats from different sites (Mantel test: Spearman's correlation = 0.28, n = 12 bats, p =
0.011; Bayesian multi-membership model: coefficient for being introduced = 0.06,
Bayesian 95% credible interval = 0.02 - 0.11). Consistent with vocal convergence, the
ability to correctly classify calls to capture site (Las Pavas or Tolé) decreased from 72%
to 65% after bats from both capture sites were merged into one colony.
359

Discussion
Variation in social affiliation drives convergence of contact calls in vampire bats
For this study, we analysed a nine-year collection of contact calls recorded from
a large sample of vampire bats for which we had known kinship and social interaction
histories. We observed multiple cases of contact call convergence within both kin and
non-kin pairs with different levels of familiarity (from completely unfamiliar to long-term
co-housing) and relationship development (mere familiarity to reciprocal food-sharing
relationship). Using these data, we demonstrated that familiarity, which was induced by
co-housing bats in captivity, causes vocal convergence. We showed that the correlation
between kinship and call similarity can be explained by social affiliation among kin.
Interestingly, food-sharing rates among familiar nonkin within the same co-housed
group predicted contact call similarity, even beyond the effect of familiarity caused by
co-housing. These findings are the first clear evidence of vocal convergence in vampire
bats, and the first evidence of vocal convergence in bats based on within-group
variation in cooperative relationships.
The mean levels of call similarity among nonkin were exactly what one would
predict if familiarity causes vocal convergence (Figure S5). We observed the lowest
contact call similarity among nonkin that never met and clearly higher call similarity
among nonkin that were captured from different wild roosts and then co-housed in
captivity. We observed the highest levels of call similarity equally among wild-caught
nonkin from the same roost that were co-housed together, and among captive-born
nonkin co-housed in the same long-term captive colony for years (Figure S5).
Food-sharing was associated with vocal convergence, but unexpectedly, we
found that call similarity among co-housed bats was not predicted by allogrooming. This
finding was clear from the difference between the effect of overall affiliative behaviour
(allogrooming and food sharing) and the effect of only food-sharing rate (Figures S7 and
S8). One interpretation of this result is that introduced (co-housed) bats vocally
converge to some degree, and then additional convergence occurs for the subset of bat
pairs with even stronger bonds (i.e. indicated by frequent food sharing). Compared to
food-sharing relationships, allogrooming relationships are less stable and less selective.

All food-sharing relationships include allogrooming but not all allogrooming relationships
involve food sharing, and when we co-housed previously unfamiliar adult female
vampire bats for almost a year, 78% of those pairs began allogrooming while only 16%
of them began food sharing (29). Moreover, the allogrooming relationships in this study
were changing more than in a typical colony due to sickness behaviour from infections
(43) and experimental manipulations such as forced proximity (29,32). Our analyses
averaged these complex dynamics over the course of an entire study into a single mean
pairwise allogrooming rate, which might not necessarily reflect the observed call
similarity for the two bats that were each recorded at different times. In contrast, food-
sharing rates in a long-term captive colony were reliable, strong, and stable indicators of
a close social bond.
401
Variation in affiliation and vocal convergence in other species
We observed vocal convergence in vampire bats at two levels: group-level
familiarity (co-housing) and within-group close social bonds (food-sharing relationships).
We believe this to be the first evidence of partner-specific vocal convergence in bats, in
contrast with convergence on a group-level signature, which has been observed in the
harsh broadband "screech" contact calls of greater spear-nosed bats (*Phyllostomus*
hastatus) (18) and the tonal constant-frequency echolocation calls of Taiwanese leaf-
nosed bats (*Hipposideros terasensis*) (19). Outside bats, partner-specific vocal
convergence has been reported in multiple birds and mammals, including in bottlenose
dolphins (15,44), orange-fronted conures (45), African elephants (46), and Campbell's
monkeys (23).
Vocal convergence at multiple levels of association (e.g. relationship and group)
has been observed in several lineages of avian and mammalian vocal learners. In
budgerigars, males entering a new flock adapt their social call to match the group (47),
and mated pairs develop a shared call (48). Monk parakeets show acoustic
convergence at the pair, flock, and site social scales (49). In crossbills, ecologically
diverged but sympatric groups have discrete call variants, and some bonded pairs
produce nearly identical call structures (50). In mammals, evidence of multi-level vocal
convergence has come from cetaceans (e.g. 15,16,44) and nonhuman primates (e.g.
8,11,24). Bottlenose dolphins have individual signature whistles (15,44), but one group
has also been shown to share a stereotyped whistle contour (16). Male Guinea baboons
live in a multilevel society and individuals have more similar grunts within gangs (the
largest group size) than between gangs, and within a party (several parties to a gang)
than between parties (11). In the pygmy marmoset *Cebuella pygmaea*, there is strong
evidence for convergence at the pair level and weak evidence at the group level:
individuals converge the duration and structure of their trill calls in response to pairing
with a new mate, but only converge duration when introduced to a group of strangers
(8,24). In general, vocal convergence at multiple levels of social familiarity may be
widespread, but there is not yet a clear pattern in how or why it is used by callers and
receivers.
432
Is vocal convergence an adaptive social behaviour or a byproduct of vocal plasticity?
Many authors argue that vocal learning requires complex and therefore costly
neurological adaptations, and is therefore unlikely to evolve unless there is a significant

fitness advantage (51). One possible explanation for vocal convergence in vampire bats
and other echolocating bats is that vocal plasticity evolved for echolocation, and that the
calls of echolocating bats might be altered by their auditory inputs as a byproduct of
audio-vocal feedback mechanisms that function for echolocation. According to this
hypothesis, vocal convergence does not serve a social function, and instead only
results from variation in an individual's exposure to the calls of other individuals.
However, this hypothesis is contradicted by our findings: captive vampire bats that could
always hear all colony members in their cages converged with food-sharing partners
more than with other cagemates. By contrast, a simple exposure effect would lead only
to group-level convergence on calls made by the most vocal bats. Instead, we observed
biased vocal convergence towards specific group members depending on the
relationships between them. Future work could further assess this hypothesis by
comparing how social contact (clustering, allogrooming, food sharing) – as distinguished
from acoustic contact alone – influences call similarity.
Vocal convergence might have social causes but not social consequences. An
alternative hypothesis is that vocal convergence in vampire bats has social
consequences that influence the development of cooperative relationships. For
example, it may be a form of behavioural coordination – a key feature of the formation
of many social bonds (e.g. coordinated courtship behaviours, vocal exchanges and turn-
taking, and reciprocal allogrooming). If partner-specific vocal convergence requires
time, it might act as an honest signal of affiliation and investment. On the other hand, if
vampire bats can rapidly imitate the calls they hear, then call matching may be used to
address specific receivers. For now, the functions of vocal convergence remain unclear.
459
Are vampire bats true "vocal learners"?
Definitions of vocal production learning differ among authors. Some authors
include modifying extant call types (which is often difficult to distinguish from vocal
usage learning), sometimes called "limited vocal learning". Others emphasize the
acquisition of entirely new call types (which is the best possible evidence for vocal
learning), sometimes called "complex vocal learning" (26,52–57). Complex vocal
production learning is rare among non-human mammals, but several nonhuman species
– including cetaceans, pinnipeds, elephants and several species of bats – can copy
artificial or novel sounds, including human speech (14,26,52,58–63).
Our findings show that vampire bats are at least 'limited' vocal learners, in the
sense that social interactions cause increased similarity in either call structure, usage,
or both. Although our data are consistent with the hypothesis that adult vampire bats are
capable of vocal production learning, further study on variation in call syllables and
responses to playbacks would be necessary to assess whether vampire bats are
modifying the acoustic features of existing calls in response to auditory input
(20,52,53,55). Interestingly, the line between convergence in syllable structure versus
syllable usage becomes blurred if contact call variation in vampire bats is not discrete
but rather continuously variable. Future work could further delineate contact calls into
different syllable types and assess the amount of convergence and individuality within
these subjectively defined "call types".
Evidence consistent with vocal learning has been found in at least 11 species of
bats, distributed across eight families (63). Only six of these species have shown *direct*

evidence of vocal learning (18,19,59–61,64,65). In the pale spear-nosed bat
(*Phyllostomus discolor*) adults can shift the fundamental frequency of their tonal social
calls towards a downward-pitch-shifted version of their calls (60), and pups change their
isolation calls to resemble their mother's individual frequency modulation over time (64).
In the greater spear-nosed bat (*Phyllostomus hastatus*) individuals changed frequency
and temporal characteristics of noisy, broadband "screech calls" to converge on a new
group signature as their group composition changed (18). In Taiwanese leaf-nosed bats
(*Hipposideros terasensis*), adult bats in the same colony shifted their echolocation calls
in the same direction gradually throughout the year, and individuals that joined the
colony shifted to match the colony's mean call frequency (19). In the greater horseshoe
bat (*Rhinolophus ferrumequinum*), pups shifted towards the frequency of their mother's
echolocation calls more than to other group members (65). In Egyptian fruit bats
(*Rousettus aegyptiacus*), pups raised in normal conditions narrowed their vocal
repertoire to fewer adult vocalizations, while experimentally isolated pups lagged
dramatically in call development or never developed adult calls (61). In the greater sac-
winged bat (*Saccopteryx bilineata*), pups of both sexes learned territorial song syllables
from adult males (22,59,66). In general, echolocating bats are remarkably specialized for
flexibility in vocal production (52,60) and have evolved sophisticated audio-vocal
systems that allow them to adapt their echolocation calls in response to acoustic inputs
(67–70). Given the exceptional vocal flexibility of bats, the evidence for vocal learning
across the bat phylogeny, and the fact that bat vocal repertoires are large and not well
understood (68), it is possible that all bats are capable of vocal convergence or even
vocal production learning.
505
Conclusion
Manipulations of co-housing in captivity showed that social familiarity causes
convergence in contact call structure in vampire bats. Further vocal convergence occurs
among familiar female bats that form close social bonds, reflected by dyadic rates of
food sharing. Although captive vampire bats in our studies were acoustically exposed to
all colony members in their cages, they appeared to converge with food-sharing
partners more than other familiar cagemates, suggesting that vocal convergence is not
completely explained by mere exposure. The social causes and consequences of this
newly discovered vocal convergence have yet to be fully explored.
515
516 **Ethics**
517 Procedures with animals were approved by the Institutional Animal Care and Use
518 Committees at the University of Maryland (protocol R-10–63) and the Smithsonian
519 Tropical Research Institute (2015-0915-2018- A9; 2017-0102-2020; 2015-0501-2022),
520 and by the Panamanian Ministry of the Environment (SE/A-76-16 and SEX/A-67-2019).
521
522 **Data accessibility**
523 Data and R code for reproducing analyses is available on GitHub (71) and figshare (72):
524 https://github.com/jkvrtilek/call-convergence-2025
525 https://doi.org/10.6084/m9.figshare.29191334
526
527 **Declaration of AI use**

528   We have not used AI-assisted technologies in creating this article.
529
Authors' contributions
Julia Vrtilek: data curation, formal analysis, investigation, visualization, writing—original
draft, and writing—review and editing; Grace Smith-Vidaurre: data curation, software,
formal analysis, investigation, validation, visualization, writing—review and editing; Eric
Fosler-Lussier: software, writing—review and editing; Rachel Page: resources,
supervision, writing—review and editing; Gerald Carter: conceptualization, data
curation, formal analysis, funding acquisition, investigation, methodology, project
administration, resources, supervision, visualization, writing—original draft, and
writing—review and editing. All authors gave final approval for publication and agreed to
be held accountable for the work performed therein.
540

541   **Competing interests**
542   We declare we have no competing interests.
543

544   **Funding**
545   This publication is based upon work supported by the National Science Foundation
546   under grant no. IOS-2015928.
547

548   **Acknowledgements**
549   We thank Gregg Cohen, Imran Razik, and Jerry Wilkinson for help with data collection.
550   We thank May Dixon, Raven Hartman, Virginia Heinen, Tobias Nguyen, Jacob
551   Chisausky, and two anonymous reviewers for feedback that greatly improved the
552   manuscript.
553

References

1. Soltis J, Leong K, Savage A. African elephant vocal communication I: antiphonal calling
behaviour among affiliated females. Anim Behav. 2005 Sept 1;70(3):579–87.

2. Miller CT, Beck K, Meade B, Wang X. Antiphonal call timing in marmosets is behaviorally
significant: Interactive playback experiments. J Comp Physiol A. 2009 Aug;195(8):783–9.

3. Carter GG, Fenton MB, Faure PA. White-winged vampire bats (*Diaemus youngi*) exchange
contact calls. Can J Zool. 2009 July;87(7):604–8.

4. Mammen DL, Nowicki S. Individual Differences and Within-Flock Convergence in Chickadee
Calls. Behavioral Ecology and Sociobiology. 1981;9(3):179–86.

5. Balsby TJS, Bradbury JW. Vocal matching by orange-fronted conures (*Aratinga canicularis*).
Behav Processes. 2009 Oct;82(2):133–9.

6. Dahlin CR, Young AM, Cordier B, Mundry R, Wright TF. A test of multiple hypotheses for
the function of call sharing in female budgerigars, *Melopsittacus undulatus*. Behav Ecol
Sociobiol. 2014 Jan 1;68(1):145–61.

7. Wright TF. Regional dialects in the contact call of a parrot. Proc R Soc B. 1997
Jan;263(1372):867–72.

8. Elowson AM, Snowdon CT. Pygmy marmosets, *Cebuella pygmaea*, modify vocal structure
in response to changed social environment. Anim Behav. 1994 June 1;47(6):1267–77.

9. Oren G, Shapira A, Lifshitz R, Vinepinsky E, Cohen R, Fried T, et al. Vocal labeling of
others by nonhuman primates. Science. 2024 Aug 30;385(6712):996–1003.

10. Zürcher Y, Willems EP, Burkart JM. Are dialects socially learned in marmoset monkeys?
Evidence from translocation experiments. PLoS One. 2019 Oct 23;14(10):e0222486.

11. Fischer J, Wegdell F, Trede F, Dal Pesco F, Hammerschmidt K. Vocal convergence in a
multi-level primate society: insights into the evolution of vocal learning. Proc Biol Sci. 2020
Dec 23;287(1941):20202531.

12. Langehennig-Peristenidou A, Scheumann M. Sex differences in the impact of social
relationships on individual vocal signatures in grey mouse lemurs (*Microcebus murinus*).
Philos Trans R Soc B. 2024 May 20;379(1905):20230193.

13. Painter MC, Gustison ML, Snyder-Mackler N, Tinsley Johnson E, le Roux A, Bergman TJ.
Acoustic variation and group level convergence of gelada, *Theropithecus gelada*, contact
calls. Animal Behaviour. 2024 Jan 1;207:235–46.

14. Poole JH, Tyack PL, Stoeger-Horwath AS, Watwood S. Elephants are capable of vocal
learning. Nature. 2005 Mar;434(7032):455–6.

15. King SL, Sayigh LS, Wells RS, Fellner W, Janik VM. Vocal copying of individually distinctive
signature whistles in bottlenose dolphins. Proc R Soc B. 2013 Apr 22;280(1757):20130053.

16. Jones BL, Daniels R, Tufano S, Ridgway S. Five members of a mixed-sex group of
bottlenose dolphins share a stereotyped whistle contour in addition to maintaining their
individually distinctive signature whistles. PLoS One. 2020 May 22;15(5):e0233658.

17. Deecke VB, Barrett-Lennard LG, Spong P, Ford JKB. The structure of stereotyped calls
reflects kinship and social affiliation in resident killer whales (*Orcinus orca*).
Naturwissenschaften. 2010 May 1;97(5):513–8.

18. Boughman JW. Vocal learning by greater spear-nosed bats. Proc R Soc B.
1998;265(1392):227–33.

19. Hiryu S, Katsura K, Nagato T, Yamazaki H, Lin LK, Watanabe Y, et al. Intra-individual
variation in the vocalized frequency of the Taiwanese leaf-nosed bat, *Hipposideros*
terasensis, influenced by conspecific colony members. J Comp Physiol A. 2006 Aug
1;192(8):807–15.

20. Tyack PL. Convergence of calls as animals form social bonds, active compensation for
noisy communication channels, and the evolution of vocal learning in mammals. J Comp
Psychol. 2008;122(3):319–31.

21. Barker AJ, Veviurko G, Bennett NC, Hart DW, Mograby L, Lewin GR. Cultural transmission
of vocal dialect in the naked mole-rat. Science. 2021 Jan 29;371(6528):503–7.

22. Knörnschild M, Nagy M, Metz M, Mayer F, von Helversen O. Learned vocal group
signatures in the polygynous bat *Saccopteryx bilineata*. Anim Behav. 2012 Oct 1;84(4):761–
9.

23. Lemasson A, Ouattara K, Petit EJ, Zuberbühler K. Social learning of vocal structure in a
nonhuman primate? BMC Evol Biol. 2011 Dec 16;11(1):362.

24. Snowdon CT, Elowson AM. Pygmy Marmosets Modify Call Structure When Paired.
Ethology. 1999;105(10):893–908.

25. Briefer EF, McElligott AG. Social effects on vocal ontogeny in an ungulate, the goat, *Capra
hircus*. Anim Behav. 2012 Apr 1;83(4):991–1000.

26. Janik VM, Knörnschild M. Vocal production learning in mammals revisited. Philos Trans R
Soc B. 2021 Oct 25;376(1836):20200244.

27. Carter GG, Logsdon R, Arnold BD, Menchaca A, Medellin RA. Adult Vampire Bats Produce
Contact Calls When Isolated: Acoustic Variation by Species, Population, Colony, and
Individual. PLoS One. 2012 June 14;7(6):e38791.

28. Carter GG, Wilkinson GS. Common vampire bat contact calls attract past food-sharing
partners. Anim Behav. 2016 June 1;116:45–51.

29. Carter GG, Farine DR, Crisp RJ, Vrtilek JK, Ripperger SP, Page RA. Development of New
Food-Sharing Relationships in Vampire Bats. Curr Biol. 2020 Apr 6;30(7):1275-1279.e3.

30. Carter GG, Wilkinson GS. Food sharing in vampire bats: reciprocal help predicts donations
more than relatedness or harassment. Proc R Soc B. 2013 Feb 22;280(1753):20122573.

31. Carter GG, Wilkinson GS. Social benefits of non-kin food sharing by female vampire bats.
Proc R Soc B. 2015 Nov 22;282(1819):20152524.

32. Razik I, Brown BKG, Carter GG. Forced proximity promotes the formation of enduring
cooperative relationships in vampire bats. Biol Lett. 2022 Apr 6;18(4):20220056.

33. Sueur J, Aubin T, Simonis C, Lellouch L, Aumond P, Baudouin A de, et al. seewave: Sound
Analysis and Synthesis [Internet]. 2023 [cited 2025 Feb 22]. Available from: https://cran.r-
project.org/web/packages/seewave/index.html

34. Araya-Salas M, Smith-Vidaurre G. warbleR: an r package to streamline analysis of animal
acoustic signals. Methods Ecol Evol. 2017;8(2):184–91.

35. Anikin A. soundgen: Sound Synthesis and Acoustic Analysis [Internet]. 2025 [cited 2025
Feb 22]. Available from: https://cran.r-project.org/web/packages/soundgen/index.html

36. Ligges U, Krey S, Mersmann O, Schnackenberg S, Guénard G, Ellis DPW. tuneR: Analysis
of Music and Speech [Internet]. 2024 [cited 2025 Aug 25]. Available from:
https://cran.csiro.au/web/packages/tuneR/index.html

37. Ohio Supercomputer Center [Internet]. Columbus OH: Ohio Supercomputer Center; 1987.
Available from: http://osc.edu/ark:/19495/f5s1ph73

38. Venables WN, Ripley BD. Modern Applied Statistics with S [Internet]. Fourth edition. New
York: Springer; 2002. Available from: https://www.stats.ox.ac.uk/pub/MASS4/

39. Bürkner PC. Advanced Bayesian Multilevel Modeling with the R Package brms. The R J.
2018;10(1):395–411.

40. Hart JDA, Weiss MN, Brent LJN, Franks DW. Common permutation methods in animal
social network analysis do not control for non-independence. Behav Ecol Sociobiol. 2022
Oct 29;76(11):151.

41. Mundry R, Sommer C. Discriminant function analysis with nonindependent data:
consequences and an alternative. Anim Behav. 2007 Oct 1;74(4):965–76.

42. Wilkinson GS. Reciprocal food sharing in the vampire bat. Nature. 1984
Mar;308(5955):181–4.

43. Razik I, Stockmaier S, Abou-Elias M, Carter GG. Opportunistic evidence of the impact of
bacterial infections on social integration in vampire bats [Internet]. bioRxiv; 2024 [cited 2025
Apr 29]. p. 2023.04.17.537180. Available from:
https://www.biorxiv.org/content/10.1101/2023.04.17.537180v5

44. King SL, Janik VM. Bottlenose dolphins can use learned vocal labels to address each other.
PNAS. 2013 Aug 6;110(32):13216–21.

45. Balsby TJS, Momberg JV, Dabelsteen T. Vocal Imitation in Parrots Allows Addressing of
Specific Individuals in a Dynamic Communication Network. PLoS One. 2012 Nov
21;7(11):e49747.

46. Pardo MA, Fristrup K, Lolchuragi DS, Poole JH, Granli P, Moss C, et al. African elephants
address one another with individually specific name-like calls. Nat Ecol Evol. 2024 June
10;1–12.

47. Hile AG, Plummer TK, Striedter GF. Male vocal imitation produces call convergence during
pair bonding in budgerigars, *Melopsittacus undulatus*. Anim Behav. 2000 June
1;59(6):1209–18.

48. Bartlett P, and Slater PJB. The effect of new recruits on the flock specific call of budgerigars
(*Melopsittacus undulatus*). Ethol Ecol Evol. 1999 Apr 1;11(2):139–47.

49. Smith-Vidaurre G, Araya-Salas M, Wright TF. Individual signatures outweigh social group
identity in contact calls of a communally nesting parrot. Behav Ecol. 2020 Mar
20;31(2):448–58.

50. Sewall KB. Limited adult vocal learning maintains call dialects but permits pair-distinctive
calls in red crossbills. Anim Behav. 2009 May 1;77(5):1303–11.

51. Nowicki S, Searcy WA. The evolution of vocal learning. Curr Opin Neurobiol. 2014 Oct
1;28:48–53.

52. Knörnschild M. Vocal production learning in bats. Curr Opin Neurobiol. 2014 Oct 1;28:80–5.

53. Tyack PL. A taxonomy for vocal learning. Philos Trans R Soc B. 2019;375(1789):20180406.

54. Jarvis ED. Evolution of vocal learning and spoken language. Science. 2019 Oct
4;366(6461):50–4.

55. Lattenkamp EZ, Vernes SC. Vocal learning: a language-relevant trait in need of a broad
cross-species approach. Curr Opin Behav Sci. 2018 June 1;21:209–15.

56. Martins PT, Boeckx C. Vocal learning: Beyond the continuum. PLOS Biol. 2020 Mar
30;18(3):e3000672.

57. Vernes SC, Kriengwatana BP, Beeck VC, Fischer J, Tyack PL, ten Cate C, et al. The multi-
dimensional nature of vocal learning. Philos Trans R Soc B. 2021 Sept
6;376(1836):20200236.

58. Janik VM. Cetacean vocal learning and communication. Curr Opin Neurobiol. 2014 Oct
1;28:60–5.

59. Knörnschild M, Nagy M, Metz M, Mayer F, von Helversen O. Complex vocal imitation during
ontogeny in a bat. Biol Lett. 2009 Oct 7;6(2):156–9.

60. Lattenkamp EZ, Vernes SC, Wiegrebe L. Vocal production learning in the pale spear-nosed
bat, *Phyllostomus discolor*. Biol Lett. 2020 Apr 15;16(4):20190928.

61. Prat Y, Taub M, Yovel Y. Vocal learning in a social mammal: Demonstrated by isolation and
playback experiments in bats. Sci Adv. 2015 Mar 27;1(2):e1500019.

62. Reichmuth C, Casey C. Vocal learning in seals, sea lions, and walruses. Curr Opin
Neurobiol. 2014 Oct 1;28:66–71.

63. Vernes SC, Wilkinson GS. Behaviour, biology and evolution of vocal learning in bats. Philos
Trans R Soc B. 2019;375(1789):20190061.

64. Esser KH, Schmidt U. Mother-Infant Communication in the Lesser Spear-nosed Bat
Phyllostomus discolor (Chiroptera, Phyllostomidae) — Evidence for Acoustic Learning.
Ethology. 1989;82(2):156–68.

65. Jones G, Ransome RD. Echolocation calls of bats are influenced by maternal effects and
change over a lifetime. Proc R Soc B. 1993;252(1334):125–8.

66. Knörnschild M, Behr O, von Helversen O. Babbling behavior in the sac-winged bat
(*Saccopteryx bilineata*). Naturwissenschaften. 2006 Sept 1;93(9):451–4.

67. Fenzl T, Schuller G. Dissimilarities in the vocal control over communication and
echolocation calls in bats. Behav Brain Res. 2007 Sept 4;182(2):173–9.

68. Salles A, Bohn KM, Moss CF. Auditory communication processing in bats: What we know
and where to go. Behav Neurosci. 2019;133:305–19.

69. Jakobsen L, Surlykke A. Vespertilionid bats control the width of their biosonar sound beam
dynamically during prey pursuit. PNAS. 2010 Aug 3;107(31):13930–5.

70. Gillam E, Fenton MB. Roles of Acoustic Social Communication in the Lives of Bats. In:
Fenton MB, Grinnell AD, Popper AN, Fay RR, editors. Bat Bioacoustics. New York, NY:
Springer; 2016. p. 117–39. (Springer Handbook of Auditory Research).

71. Vrtilek JK, Carter G. call-convergence-2025 [Internet]. 2025 [cited 2025 Mar 30]. Available
from: https://github.com/jkvrtilek/call-convergence-2025

72. Vrtilek JK, Smith-Vidaurre G, Carter GG. Data for "Vocal convergence during formation of
cooperative relationships in vampire bats" [Internet]. figshare; 2025 [cited 2025 Aug 7].
Available from: https://doi.org/10.6084/m9.figshare.29191334

73. Ripperger SP, Carter GG, Duda N, Koelpin A, Cassens B, Kapitza R, et al. Vampire Bats
that Cooperate in the Lab Maintain Their Social Networks in the Wild. Curr Biol. 2019 Dec
2;29(23):4139-4144.e4.

Supplementary Information (SI)

Supplementary Methods

Variation in contact call structure

We defined contact calls based on a combination of behavioural context and call
features. Specifically, we only analysed calls that were produced by a bat that was
isolated from group members, had durations of 3-50 ms, and had peak frequencies of at
least 10 kHz. Calls were excluded by our analysis pipeline whenever we failed to
measure frequencies from the fundamental, indicating low signal-to-noise ratio. We
found that removing these calls increased the overall call classification rate for
assigning calls to bat.
Within these call parameters, we failed to find any clear evidence for discrete call
types despite considerable variation both among and within callers (Figure S1). Given
that contact call variation appeared to be graded (continuous) rather than discrete, we
decided to take the conservative first step of assessing similarity for all contact calls
rather than subjectively defining subtypes and measuring similarity within each type.
The latter approach would also elevate the false discovery rate due to multiple testing
and the many possible ways to label types. Future work can assess the degree to which
contact call variation is continuous rather than discrete and describe how vocal
individuality differs between structurally defined types of syllables.
746

747
Figure S1. Representative variation in contact calls within and among individual
vampire bats. Spectrograms of example contact calls from four female vampire bats
(row) across 4 different nights of being individually isolated. Each row represents an
individual; the top row is from 2011-2013, the second row is from 2016-2017, and the
bottom two rows are from 2019. Recordings in a row are in order from earliest to latest.
The variable structure of contact calls suggest that they might convey other information
beyond the caller's individual identity, such as motivational state.
755

Model 1 posterior predictive check

Model 2 posterior predictive check

Model 3 posterior predictive check

Model 4 posterior predictive check

Model 5 posterior predictive check

756
757 **Figure S2. Posterior predictive checks for 5 Bayesian models.** Comparison of
758 distribution of observed contact call similarity values (dark black line) to the predicted
759 values from 100 datasets simulated from the fitted model (light blue lines) suggests that
760 the models capture the essential features of the data-generating process.
761

762
763 **Figure S3. Timeline of call recordings for analysis of call convergence within-**
764 **pairs over time.** Top panel: recording dates (dots) for 5 bats (lines) captured from Las
765 Pavas, Panama. Bottom panel: recording dates for 7 bats from Tolé, Panama. Dotted
766 red line: date of introduction between Tolé and Las Pavas bats in small groups (2-5
767 bats). Solid red line: date of introduction between all captive bats in large flight cage.
768

769 **Table S1. Bats recorded for this study.** Of the 115 bats recorded for this study, 95
770 produced more than 100 contact calls. The number of bats below adds to 97 because 2
771 bats from the 2016-2017 colony were recaptured and included in the 2019 colony.
772

Years recorded	# bats	Familiarity	Kinship	Origin	Age and Sex
2011-2014	30	Long-term captive colony (28,30,31)	Previously estimated using maternal pedigree and microsatellite markers (31)	Multiple zoos	12 adult F 4 subadult F 5 adult M 9 subadult M
2016	4	Never caged with or introduced to any other bats in the study, but potentially familiar	Unrelated to other bats, but unknown relatedness among each other	Cattle pasture, Chilibre, Panama	1 adult F 3 adult M
2016	1	Never caged with or introduced to any other bat in the study	Unrelated to other bats	Forest, Gamboa, Panama	1 adult M
2016-2017	40	Co-housed (caged together) for over 10 months (29,73)	Previously estimated using microsatellite markers (73); bats from different sites are unrelated	Cattle pasture, Las Pavas, Panama	5 adult F
				Captive-born to Las Pavas bats	1 subadult F 1 subadult M
				Tree hollow, Tolé, Panama	19 adult F 7 adult M
				Captive-born to Tolé bats	6 subadult F 1 subadult M
2019	22	Co-housed for 4 months (32)	Bats from different sites are unrelated; kinship unknown for bats from the same site	Same tree, Tolé, Panama	2 recaptured adult F 7 new adult F
				Tree hollow, La Chorrera, Panama	7 adult F
				Cave, Lake Bayano, Panama	6 adult F

774 **Table S2. Spectral and temporal measurements used in linear discriminant**
775 **function analyses.** *Duration* through *meanpeakf* descriptions are edited from the
776 *warbleR* (34) and *seewave* (33) documentation. *Maxslope* through *segments* are
777 summary measures of the fundamental from the *soundgen* package (35).
778

Measure	Explanation
duration	Length of signal (in ms)
meanfreq	Mean of frequency spectrum (i.e. weighted average of frequency by amplitude within supplied band pass) (in kHz)
sd	Standard deviation of frequency (in kHz).
freq.median	The frequency at which the signal is divided in two frequency intervals of equal energy (in kHz)
freq.Q25	The frequency at which the signal is divided in two frequency intervals of 25% and 75% energy respectively (in kHz)
freq.Q75	The frequency at which the signal is divided in two frequency intervals of 75% and 25% energy respectively (in kHz)
freq.IQR	Frequency range between 'freq.Q25' and 'freq.Q75' (in kHz)
time.median	The time at which the signal is divided in two time intervals of equal energy (in s)
time.Q25	The time at which the signal is divided in two time intervals of 25% and 75% energy respectively (in s).
time.Q75	The time at which the signal is divided in two time intervals of 75% and 25% energy respectively (in s).
time.IQR	Time range between 'time.Q25' and 'time.Q75' (in s).
skew	Asymmetry of the frequency spectrum
kurt	Peakedness of the frequency spectrum
sp.ent	Energy distribution of the frequency spectrum
time.ent	Energy distribution on the time envelope
entropy	Product of sp.ent and time.ent
sfm	Spectral flatness
meandom	Average of dominant frequency measured across the acoustic signal
mindom	Minimum of dominant frequency measured across the acoustic signal
maxdom	Maximum of dominant frequency measured across the acoustic signal
dfrange	Range of dominant frequency measured across the acoustic signal
modindx	Modulation index, calculated as the cumulative absolute difference between adjacent measurements of dominant frequencies divided by the dominant frequency range
startdom	Dominant frequency measurement at the start of the signal
enddom	Dominant frequency measurement at the end of the signal
dfslope	Slope of the change in dominant frequency through time (kHz/s)
peakf	Frequency with the highest energy from the frequency spectrum
meanpeakf	Frequency with highest energy from the mean frequency spectrum (the mean relative amplitude of the frequency distribution of a time wave)
maxslope	Maximum slope of fundamental frequency contour
minslope	Minimum slope of fundamental frequency contour
absminslope	Absolute value of the minimum slope of fundamental frequency contour
pos.slopes	Number of positive slopes in the fundamental frequency contour
neg.slopes	Number of negative slopes in the fundamental frequency contour
turns	Number of turns (changes between positive and negative slopes) in the fundamental frequency contour
meanslope	Mean slope of the fundamental frequency contour
segments	Number of separate segments in the fundamental frequency contour

Supplementary Results

782
Figure S4. Kinship predicts contact call similarity across pairs with known
kinship. Left panel shows how call similarity increases with kinship. Blue line shows a
general additive model fit with the ggplot2 R package. Right panel shows mean contact
call similarity with bootstrapped 95% confidence intervals (CIs) for three bins of kinship.
The 95% CIs treat pairs as independent observations, so these plots are only
descriptive; inferences should be drawn from the multi-membership model results
(Figure 2b).
790

791
Figure S5. Contact call similarity by levels of familiarity. Top panel shows the
distributions and boxplots of the contact call similarity; bottom panel shows means with
bootstrapped 95% CIs for contact call similarity. Both are shown at four levels of
familiarity (x-axis): (1) unfamiliar nonkin that never met (baseline), (2) previously
unfamiliar nonkin captured from different wild roosts then co-housed in captivity, (3)
nonkin captured from the same roost then co-housed together in captivity, and (4)
familiar nonkin co-housed in the same long-term captive colony. Vocal convergence is
demonstrated by the difference from the pairs that never met (horizontal line). The 95%
CIs treat pairs as independent observations, so these plots are only descriptive;
inferences should be drawn from the multi-membership model results (Figure 2b).
802

Figure S6. The association between nonkin affiliation rate and nonkin contact call similarity across colonies. Only pairs that were nonkin (r <0.05) are plotted. The 2014 dataset included only food donations to fasted bats in a long-term captive colony (left panel). The 2017 and 2019 colonies were shorter-term colonies that merged samples of bats from different wild populations. The 2017 colony (orange) included mostly allogrooming and some food donations. The 2019 dataset (purple) included only allogrooming in pairs from different sites, because kinship was unknown in pairs from the same site (although including kin pairs does not change the conclusions from these analyses). The 95% CIs on slopes treat pairs as independent observations, so these plots are only descriptive; inferences should be drawn from the multi-membership model results (Figure 2b).

816
Figure S7. Nonkin pairs that engage in more food sharing tend to have more
similar contact calls. Greater food-sharing predicted greater contact call similarity
among nonkin females (A). The absence and presence of food sharing is possibly
linked to greater contact call similarity in nonkin females (B) and in all nonkin pairs (C).
The 95% CIs treat pairs as independent observations so these plots are only
descriptive; inferences should be drawn from the multi-membership model results
(Figure 2b).
824

825 **Table S3. Model coefficients.** Other columns are the estimate of the average
826 (expected) posterior probability, the estimated error (the posterior standard deviation),
827 the 95% credible interval, the R-hat convergence diagnostic (convergence = 1.00), and
828 the Bulk and Tail effect sample sizes.

Model	Term	Estimate	Est. Error	Lower 95% CI	Upper 95% CI	R-hat	Bulk ESS	Tail ESS
1	kinship (scaled)	0.037	0.005	0.028	0.047	1.000	12111	11709
2	affiliation log rate* (scaled)	0.025	0.006	0.013	0.037	1.001	6270	9051
2	kinship (scaled)	0.003	0.005	-0.006	0.013	1.000	7686	9742
2	co-housed (T/F)	0.148	0.013	0.123	0.173	1.001	6031	8581
3	co-housed (T/F)	0.207	0.015	0.178	0.238	1.001	7717	9579
4	affiliation log rate* (scaled)	-0.002	0.020	-0.040	0.036	1.000	8090	10802
5	food sharing log rate* (scaled)	0.068	0.026	0.017	0.119	1.000	10274	11916

829 *Log rate is (natural log (seconds of observation + 1)) / natural log (sampled hours).
830
831