Vocal convergence during formation of social relationships in vampire bats

Julia Vrtilek*[1], Grace Smith-Vidaurre*[2,3,4,5], Eric Fosler-Lussier[6], Rachel A. Page[7],
Gerald G. Carter[7,8,9]

Affiliations
1. Department of Evolution, Ecology and Organismal Biology; The Ohio State University, Columbus, Ohio, USA
2. Department of Biology, New Mexico State University, Las Cruces, NM, USA
3. Department of Integrative Biology; Michigan State University, East Lansing, MI, USA
4. Department of Computational Mathematics, Science, and Engineering; Michigan State University, East Lansing, MI, USA
5. Ecology, Evolution, and Behavior Program; Michigan State University, East Lansing, MI, USA
6. Department of Computer Science and Engineering; The Ohio State University, Columbus, Ohio, USA
7. Smithsonian Tropical Research Institute, Balboa, Ancón, Panamá
8. Department of Ecology and Evolutionary Biology; Princeton University, Princeton, New Jersey, USA
9. Howard Hughes Medical Institute, Chevy Chase, Maryland, USA

*These authors contributed equally to this work

Corresponding authors: Julia Vrtilek: jkvrtilek@gmail.com; Gerald Carter: gc1511@princeton.edu

Keywords: contact calls, social bonds, social relationships, vampire bats, vocal convergence, vocal learning

Abstract

Vocal behavior provides an important window into animal social lives. A longstanding hypothesis for many group-living birds and mammals is that the formation of affiliative relationships causes an increase in call similarity between individuals (vocal convergence). Testing this causal effect can be difficult, however, because it requires experimentally forming new relationships. Here, we demonstrate convergence in the contact calls of common vampire bats (*Desmodus rotundus*) that we introduced and experimentally housed together. To estimate and disentangle the roles of kinship, co-housing (familiarity), allogrooming, and food sharing in predicting call similarity, we first measured call similarity using 35 features of 565,557 contact calls from 94 bats, then fit a series of Bayesian generalized multi-membership models. We also measured changes in call similarity for a subset of individuals that were recorded before and after co-housing. We found that co-housing caused vocal convergence. Furthermore, food-sharing rates among familiar nonkin within the same co-housed group predicted contact call similarity. This finding suggests that the development of cooperative relationships causes further vocal convergence beyond the initial convergence caused by co-housing. Our findings have implications for the development of cooperative relationships and vocal learning.

Introduction

Vocal behavior provides an important window into the social lives of many birds and mammals. Exchanges of vocalizations called "contact calls" seem to facilitate recognition, reunions, and collective movements among affiliated individuals across several species (e.g. African elephants (1), common marmosets (2), and white-winged vampire bats (3)). In other group-living birds and mammals, the contact calls of unrelated members of the same group sound more similar than those of strangers (e.g. orange-fronted conures (4), budgerigars (5), yellow-naped amazons (6), pygmy and common marmosets (7–9), Guinea baboons (10), grey mouse lemurs (11), gelada (12), African elephants (13), bottlenose dolphins (14,15), orcas (16), greater spear-nosed bats (17) and Taiwanese leaf-nosed bats (18)). Based on this observation, a longstanding hypothesis is that the formation of affiliative relationships ("social bonds") can increase similarity in the acoustic structure of calls between individuals ("vocal convergence") (19).

Although vocal similarity or convergence is often associated with affiliative relationships (9,17,18,20–24), there are multiple possible mechanisms for vocal convergence. First, familiar animals might have the ability to modify the structure of vocalizations based on auditory inputs (vocal production learning), causing similarity between familiar individuals. However, vocal convergence exists in many mammalian lineages where vocal production learning has not yet been demonstrated (25). In these cases, it is possible that animals might learn from conspecifics to produce existing sounds in their repertoire in new contexts (usage learning). Finally, familiar animals might use the same vocalizations more often because their physiological or motivational states are often matching (25).

Regardless of the underlying mechanism, demonstrating a causal link from
affiliation to vocal convergence can be difficult. For example, correlations between
affiliation and vocal similarity can result from kinship, either through heritable variation
or shared environment. The same correlation can result from pre-existing vocal
similarity causing higher affiliation (assortment based on acoustic similarity). To
demonstrate causation, an ideal study would induce new relationships to form among
nonkin, then observe changes in call similarity.
Here, we applied this approach to demonstrate convergence in the contact calls
of common vampire bats (*Desmodus rotundus*). Vampire bats produce contact calls
when they are socially isolated (26) and use the calls to vocally recognize affiliated
individuals (27). We analyzed 565,557 contact calls from 94 bats, some of which were
experimentally co-housed together. We found that co-housing causes vocal
convergence. Next, to tease apart how pairwise call similarity was shaped by
relationships within each captive group, we estimated effects of genetic relatedness,
allogrooming, and food sharing (28–31). Both kin and nonkin females can develop
reciprocal allogrooming relationships that can become reciprocal food-sharing
relationships, in which a fed bat regurgitates potentially life-saving ingested blood to a
starving partner (28–30). We found evidence that these food-sharing relationships
cause further vocal convergence between nonkin after the initial convergence caused
by co-housing.
94

Methods
Recording contact calls from individually isolated bats
Contact calls are frequency modulated syllables between 3-50 ms in duration
with peak frequencies of 10-30 kHz that are produced by socially isolated bats (26). To
record contact calls with high signal-to-noise ratios, we isolated an individual bat inside
a plastic storage bin (about 75 to 200 liters in volume) lined with acoustic foam and/or
soft fabric to dampen echoes. To keep the bat within 10 to 30 cm of the microphone, we
placed it inside a small soft-mesh butterfly cage or a tube of plastic mesh. Calls were
recorded with an Avisoft CM16 ultrasound condenser microphone (frequency range 10-
200kHz, Avisoft Bioacoustics, Berlin, Germany). Sound was digitized with 16-bit
resolution at sampling rates ranging between 250-500 kHz through an Avisoft
UltraSoundGate (one-channel or four-channel) to a laptop running the program Avisoft
Recorder. To remove any effects of digitization rate on subsequent call measures,
recordings digitized at a rate above 250 kHz were downsampled to 250 kHz using the
seewave R package (32). To filter out sounds that were not contact calls, we removed
calls with durations less than 3 ms and longer than 50 ms and with peak frequencies
outside the range of 10-30 kHz. For further explanation of call selection, see
Supplementary Information (SI).
From 2011 to 2019, we recorded 565,557 contact calls from 113 individually
isolated common vampire bats, but we only analyzed 565,192 calls from 94 of those
bats (4371 pairs of bats). We only included bats from which we sampled over 80 contact
calls of sufficient quality to allow for automated measurements of the call fundamental

(mean calls per bat = 6013, median = 1945, minimum = 84, maximum = 86,573). The
number of recording sessions per bat varied from 1-22 (mean = 7.33).
119
Study subjects
The 94 bats used in this study came from 8 different origins (e.g. captive colonies
or field sites; for details, see Supplementary Information (SI), Table S1). To identify
individual bats, all bats were marked using 1-4 unique bat bands. We labeled bats as
"adults" if they were older than 2 years old at time of recording, or if their exact age was
unknown, by the presence or absence of an epiphyseal gap between their metacarpals.
We refer to volant non-adults as "subadult". Non-volant pups that were still nursing from
their mother were not included in the study.
128
Kinship, social familiarity, and cooperative relationships
Bats captured from different sites hundreds of kilometers apart were considered
both unrelated (non-kin) and unfamiliar. Kinship of bats from the same colony was
estimated through a combination of microsatellite markers and maternal pedigrees, as
described in previous studies (28,30). Pairs of bats with unknown kinship were excluded
from analyses where kinship was a predictor.
Most pairs (n = 2540) were *unfamiliar* because they had never met, 582 pairs
were captured from different field sites hundreds of km apart, then *introduced in*
captivity and caged together (co-housed) for at least 4 months; *612 pairs* were captured
from the same field site then caged together for at least 4 months (n = 612); and 406
pairs *were* living in captivity for years (n = 406). Other pairs (n = 231) had unknown
levels of kinship and familiarity, because they were captured from the same wild roost
about four years apart. Within each co-housed captive group, pairs had variable rates of
cooperative behavior. We had measures of food-sharing rates for 284 pairs,
allogrooming rates for 190 pairs, and both for 406 pairs. These data were collected from
a series of past studies where experimenters repeatedly induced food sharing by
removing a bat, fasting it for over 24 hours, and reintroducing it to its home cage, then
used focal follow sampling to observe allogrooming and regurgitated food sharing (28–
30).
148
Measuring features of calls
We identified start and end times of individual calls with a custom R script that
uses both amplitude and spectral density to segment audio recordings into calls. The
script excludes calls which exceed the maximum amplitude of the microphone. This
automated method was tested and validated with 1535 manually labeled recordings
containing ~7000 call selections. Using these start and end times, we then measured 35
features of each call (Figure 1). We used the R package *warbleR* (33) to measure 27
spectral and temporal parameters and the R package *soundgen* (34) to estimate 8
additional measurements from the fundamental: maximum slope, minimum slope, and
absolute minimum slope; the number of positive slopes and of negative slopes; the
number of turns; the mean slope; and the number of segments. All scripts were run
using the Ohio Supercomputer (35).
161
162

163
Figure 1. Measures from spectrogram of contact call. Call measures (defined in
Table S1) include: 1 duration, 2 time quantile 25, 3 time median, 4 time quantile 75, 5
mean frequency, 6 frequency quantile 25, 7 frequency median, 8 frequency quantile 75,
9 maximum dominant frequency, 10 mean dominant frequency, 11 minimum dominant
frequency, 12 start dominant frequency, 13 end dominant frequency, 14 mean peak
frequency, 15 peak frequency. The following variables are not shown: time interquartile
range, frequency interquartile range, standard deviation of frequency, dominant
frequency range, dominant frequency slope, skew, kurt, spectral entropy, time entropy,
entropy, spectral flatness, modulation index, and fundamental frequency summary
measures (maximum, minimum, and absolute minimum slope; the number of positive
slopes and of negative slopes; the number of turns; the mean slope; and the number of
segments).
176
Assessing vocal individuality
To assess and confirm baseline individuality in contact call structure, we
estimated the classification rate for assigning calls to the correct caller using a linear
discriminant function analysis (DFA) with leave-one-out cross-validation, including all
variables measured. To choose a minimum number of calls per bat for later analyses,
we compared these classification rates to the same rates from a single DFA without
cross-validation. We excluded bats with fewer than 80 sampled calls because within-bat
classification rates were virtually identical for bats with greater than 80 calls. To confirm
that calls could be assigned to the caller with greater-than-chance accuracy, we ran a
permutation test: DFA without cross-validation was repeated 1000 times with
randomized bat names to generate an expected distribution of accuracy rates under the
null hypothesis that calls contain no information about caller identity.
189
Estimating contact call similarity

We defined contact call similarity for each pair of bats as:

$$S = 1 - \left(\frac{D}{maximum\ (D)}\right)$$ (1)

where D is the Mahalanobis distance between the two group centroids (bats)
from a DFA classifying calls (n = 565,192) to bat (n = 94). These Mahalanobis distances
represent the average multivariate distance (or dissimilarity) between every two bats.
We normalized our measure of call similarity to range from 0 to 1; however, we replaced
the single similarity score of 0 with 0.001 (the next lowest call similarity is 0.03),
because similarity cannot be zero and beta regression cannot model an outcome of
zero.

Estimating effects of kinship and social relationship on call similarity
To assess predictors of contact call similarity while controlling for the effect of
both bat identities, we fit five Bayesian generalized (beta distribution) multi-membership
models (Table 1) using the R package *brms* (36). Multi-membership models are used to
model dyadic data by fitting group-structured random effects, which capture the
dependence of each pair on the identities of both individuals. This model structure
allows us to estimate a random intercept for each bat and its influence on its pairings
with other bats (37). To assess causal hypotheses, we estimated multiple predictors
across seven models based on known and hypothesized causal paths (Figure S2).
Our first goal was to disentangle the effects of kinship from social familiarity. In
model 1, we estimated how well kinship predicted contact call similarity across the 2068
pairs of adult bats with known kinship estimates. Kinship is confounded with group-level
familiarity and within-group affiliation rates, and any effects of kinship in model 1 could
therefore be driven by social interaction history rather than kinship. In model 2, we
therefore estimated how well kinship predicted contact call similarity when conditioning
on (i.e. statistically controlling for) affiliation rates in the same 2068 pairs. Predictors
were kinship, whether the bats were co-housed (true/false), and within-group affiliation
lograte, defined as:

$$\frac{log(observed + 1)}{log(sampling)}$$ (2)

where "observed" is the seconds of either allogrooming or regurgitated food
sharing between the two bats and "sampling" is the total seconds in which allogrooming
or food sharing could have been observed between the two bats. Both allogrooming and
food sharing are directed behaviors that indicate an affiliative social preference. We
used logrates rather than rates because seconds of allogrooming and food sharing are
lognormal. We scaled (i.e. z-transformed) both kinship and affiliation lograte to convert
units to standard deviations.
Another way to control for kinship is to only test unrelated pairs. In model 3, we
measured how well being caged together (true/false) predicted contact call similarity in
the 1114 pairs of unrelated female adults caught from different sites. All pairs had
kinship estimates of 0, allowing us to test for an effect of nonkin being caged together

on their contact call similarity. We focused on female pairs because the few males in
these colonies were sub-adults or young born into the group.
Our next goal was to separate the effects of group-level familiarity and dyad-level
affiliation rate among nonkin females. In model 4, we measured how well within-group
affiliation lograte (scaled lograte) predicted contact call similarity in 265 unrelated pairs
of female adults that were caged together. We removed pairs with kinship estimates
greater than 0.05 to test for an effect of nonkin social interactions on contact call
similarity. We also removed pairs that could never have been in contact. We again
focused on female pairs because social affiliation data were focused on females and
undersampled or missing for males (28–31).
We did not include allogrooming and allofeeding together as covariates in model
4 because they are highly correlated proxies for social affiliation, rather than different
effects. The affiliation rate therefore includes either allogrooming, regurgitated food
sharing, or both. The results of model 4 must therefore be interpreted carefully, because
affiliation rate is based more on allogrooming in some pairs than others. For example, in
one colony from 2010-2014, food-sharing rates were sampled more than allogrooming,
but in another colony in 2019, food-sharing rates were unknown (missing) for all pairs.
Moreover, food sharing represents a much higher level of affiliation than allogrooming.
For example, when bats are introduced in captivity (i.e. become familiar), most pairs
form allogrooming relationships and only a few begin food-sharing relationships (28). In
other words, co-housing predicts allogrooming better than food sharing.
To remedy these issues, we next focused on food-sharing rates, which is the
best indication of a within-group social bond (28). In model 5, we measured how well
food-sharing rate (scaled lograte) predicted contact call similarity in the 132 unrelated
pairs of female adults that were caged together. We again removed pairs with kinship
estimates greater than 0.05, pairs that could never have shared food and pairs with
unknown food-sharing rates.
Across all models, we used default priors and scaled continuous predictors. We
used posterior predictive checks to assess model fit (Figure S3). To ensure Markov
Chain Monte Carlo (MCMC) convergence, we checked the Rhat values across 4 chains,
which ranged from 0.9999 to 1.0004, showing that all 4 chains reached the same
estimates for coefficients. We used chain and warmup lengths of 6000 and 2000
samples, respectively. The lowest Bulk and Tail effective sample sizes across all
models were 8747 and 10064 respectively.
269

Model	Question addressed	Covariates included as predictors of contact call similarity	Bat pairs in sample
1	How well does kinship predict contact call similarity?	kinship	pairs with estimated kinship = 2068
2	How well does kinship predict contact call similarity after controlling for co-housing and affiliation?	kinship, co-housing (true/false), affiliation rate (including zeros for bats that never met)	pairs with estimated kinship = 2068
3	In nonkin females, how well does co-housing predict contact call similarity?	co-housing (true/false)	pairs of nonkin adult females = 1114
4	In nonkin females, how well does affiliation predict contact call similarity?	within-group affiliation rate (excluding bats that never met)	pairs of nonkin adult females that were caged together = 265
5	In nonkin females, how well does food sharing predict contact call similarity?	within-group food-sharing rate (excluding bats that never met)	pairs of nonkin adult females with estimated food-sharing rates = 132

270 **Table 1. Purpose of each statistical model.** We selected this set of models based on
271 causal models shown in Figure S2.
272
Comparing call similarity pre- and post-introduction
We could determine how call similarity changed over time using a subset of
cases in 2016 where 7 bats from Tolé were recorded both before and after being
introduced to 5 bats from Las Pavas (Figure S4). We predicted that contact call
convergence (increases in call similarity) should be greater for introduced pairs (from
different sites) than for previously familiar pairs (from the same site). To compare call
similarity before and after introduction, we first performed two discriminant function
analyses (DFAs) classifying calls to bats. Both DFAs included 5 Las Pavas bats and 7
Tolé bats, but the first DFA included recordings from the Tolé bats *before introduction* to
the Las Pavas bats, and the second DFA the same Tolé bats recorded *after the*
introduction. From each DFA, we extracted Mahalanobis distances and call similarities
as described above, then we subtracted call similarities post-introduction from call
similarities pre-introduction to estimate the change in similarity.
To test the hypothesis that increases in similarity were greater for introduced
pairs than for familiar pairs, we used two approaches. First, we used a nonparametric
permutation test (Mantel test with Spearman correlation) of the correlation between the
change in contact call similarity and a binary familiarity matrix indicating familiar vs.
introduced pairs. Second, we used a Bayesian multi-membership model (Gaussian
distribution, default priors in brms, 4 chains with length of 6000 interactions, 2000

warmup samples) to test whether familiarity predicted the change in contact call
similarity.
Finally, we tested the prediction that call classification to the correct capture site
should be more accurate prior to the merging of the two groups. To do this, we used two
permuted DFAs, which are designed to test for correct classification of calls to social
group (38). In the first permuted DFA, we classified calls to either the Las Pavas group
or the *pre-introduction* Tolé group. In the second permuted DFA, we classified calls to
either the Las Pavas group or the *post-introduction* Tolé group.
300

Results

Contact call variation within and between individuals
Contact calls had clear individual signatures, but were also highly variable within
bat (SI, Figure S1). Discriminant function analysis (DFA) correctly assigned the 565,192
calls to the 94 callers with 38% probability, which is more than twice the expected
chance level of 15.3% (95% quantiles = 15.30% to 15.32%). The call assignment
accuracy of assigning calls back to individual bats from the cross-validated DFA and
from the non-cross-validated DFAs were almost perfectly correlated across the 94 bats
(r = 0.9998), indicating robustness of the DFA used for estimating call similarity.
310
Kinship-biased contact call similarity can be explained by social familiarity
Model 1 revealed that kinship predicts contact call similarity (Figure 2, Figure
S5). However, kinship is confounded with social affiliation in vampire bats (29,30,39).
We therefore used Model 2 to test for an effect of kinship while conditioning on (i.e.
"controlling for") affiliation rate and co-housing. We no longer found an effect of kinship
while conditioning on these measures of social affiliation, but we did find that call
similarity was predicted by both affiliation rate and co-housing while conditioning on
kinship and each other (Figure 2). This result does not mean that kinship did not
influence call similarity; only that the correlations between kinship and contact call
similarity can be explained by social familiarity. In other words, both affiliation rate and
familiarity predicted contact call similarity, regardless of kinship.
322
Co-housing causes contact call similarity among nonkin
To further test whether co-housing causes contact call convergence among
nonkin, we used Model 3 to focus on 1114 pairs of unrelated female adults that were
captured from different sites. Compared to model 2, which included kin, we found that
co-housing was associated with an even larger increase in call similarity among pairs of
unrelated adult females (Figure 2). As expected, call similarity for nonkin from distant
sites that were housed together in captivity was dramatically higher than nonkin from
different sites that never met, and the highest mean similarity in nonkin contact calls
occurred within nonkin pairs from the same wild roost or in the same long-term captive
colony (Figure S6).
333
Within co-housed groups, food-sharing rates predict further vocal convergence

Although Model 2 revealed an effect of affiliation (allogrooming and food sharing,
Figure S7), further analyses showed that this effect was driven by food sharing rather
than allogrooming (Figure 2, Figure S8). Specifically, Model 4 estimated the effect of
within-group dyadic affiliation rate on call similarity among unrelated female adults, and
this model found no clear effect of affiliation rate beyond the effect of previous
familiarity. However, Model 5 showed that contact call similarity was predicted by within-
group rates of regurgitated food sharing. This effect of food sharing on call similarity
was driven by the 2014 long-term captive colony (Figure S8), which had the highest
rates of food sharing and the longest periods of familiarity.
344
Vocal convergence evident from call similarity before and after co-housing
We saw evidence for vocal convergence over time (increased contact call
similarity after introduction) in 21 of 35 pairs from different sites (60%), compared to 9 of
21 familiar pairs from Tolé (43%). We found that a greater increase in call similarity (i.e.
greater vocal convergence) was positively associated with being an introduced pair of
bats from different sites (Mantel test: Spearman's correlation = 0.28, n = 12 bats, p =
0.007; Bayesian multi-membership model: coefficient for being introduced = 0.06,
Bayesian 95% credible interval = 0.02 - 0.11). Consistent with vocal convergence, the
ability to correctly classify calls to capture site (Las Pavas or Tolé) decreased from 77%
to 67% after bats from both capture sites were merged into one colony.
355

356
Figure 2. Contact call similarity is predicted by co-housing, affiliation rate, and
food-sharing. Posterior probability distributions with means (circles) and their Bayesian
95% credible intervals (circles error bars) for regression coefficients (y-axis) from one of
five Bayesian models (each a different color). Models were selected to disentangle
effects of kinship, co-housing, affiliation rate, and food-sharing. See Figure S2 for
diagram of causal relationships and Table S3 for coefficients and credible intervals.
363

Discussion

Variation in social affiliation drives convergence of contact calls in vampire bats

For this study, we analysed a nine-year collection of contact calls recorded from over a hundred vampire bats for which we had known kinship and social interaction histories. We observed multiple cases of contact call convergence within both kin and non-kin pairs with different levels of familiarity (from completely unfamiliar to long-term co-housing) and relationship development (mere familiarity to reciprocal food-sharing relationship). Using these data, we demonstrated that familiarity, which was induced by co-housing bats in captivity, causes vocal convergence. We showed that the correlation between kinship and call similarity can be explained by social affiliation among kin. Interestingly, food-sharing rates among familiar nonkin within the same co-housed group predicted contact call similarity, even beyond the effect of familiarity caused by co-housing. These findings are the first clear evidence of vocal convergence in vampire bats, and the first evidence of vocal convergence in bats based on within-group variation in cooperative relationships.

The mean levels of call similarity among nonkin were exactly what one would predict if familiarity causes vocal convergence (Figure S6). We observed the lowest contact call similarity among nonkin that never met and clearly higher call similarity among nonkin that were captured from different wild roosts and then co-housed in captivity. We observed the highest levels of call similarity equally among wild-caught nonkin from the same roost that were co-housed together, and among captive-born nonkin co-housed in the same long-term captive colony for years (Figure S6).

Food-sharing was associated with vocal convergence, but unexpectedly, we found that call similarity among co-housed bats was not predicted by allogrooming. This finding was clear from the difference between the effect of all affiliative behavior (allogrooming and food sharing) and the effect of only food-sharing rate (Figures S7 and S8). One interpretation of this result is that introduced (co-housed) bats vocally converge to some degree, and then additional convergence occurs for the subset of bat pairs with even stronger bonds (i.e. indicated by frequent food sharing). Compared to food-sharing relationships, allogrooming relationships are less stable and less selective. All food-sharing relationships include allogrooming but not all allogrooming relationships involve food sharing, and when we co-housed previously unfamiliar adult female vampire bats for almost a year, 78% of pairs began allogrooming while only 16% began food sharing (28). Moreover, the allogrooming relationships in this study were changing more than in a typical colony due to experimental manipulations such as forced proximity (28,31) and sickness behavior from infections (40). Our analyses averaged these complex dynamics over the course of an entire study into a single mean pairwise allogrooming rate, which might not necessarily reflect the observed call similarity for the two bats that were each recorded at different times. In contrast, food-sharing rates in a long-term captive colony were reliable, strong, and stable indicators of a close social bond.

Variation in affiliation and vocal convergence in other species

We observed vocal convergence in vampire bats at two levels: group-level familiarity (co-housing) and within-group close social bonds (food-sharing relationships).

We believe this to be the first evidence of partner-specific vocal convergence in bats, in
contrast with convergence on a group-level signature, which has been observed in the
harsh broadband "screech" contact calls of greater spear-nosed bats (*Phyllostomus*
hastatus) (17) and the tonal constant-frequency echolocation calls of Taiwanese leaf-
nosed bats (*Hipposideros terasensis*) (18). Outside bats, partner-specific vocal
convergence has been reported in multiple birds and mammals, including in bottlenose
dolphins (14,41), orange-fronted conures (42), African elephants (43), and Campbell's
monkeys (22).
Vocal convergence at multiple levels of association (e.g. relationship and group)
has been observed in several lineages of avian and mammalian vocal learners. In
budgerigars, males entering a new flock adapt their social call to match the group (44),
and mated pairs develop a shared call (45). Monk parakeets show acoustic
convergence at the pair, flock, and site social scales (46). In crossbills, ecologically
diverged but sympatric groups have discrete call variants, and some bonded pairs
produce nearly identical call structures (47). In mammals, evidence of multi-level vocal
convergence has come from cetaceans (e.g. 14,15,41) and nonhuman primates (e.g.
7,10,23). Bottlenose dolphins have individual signature whistles (41,14), but one group
has also been shown to share a stereotyped whistle contour (15). Male Guinea baboons
live in a multilevel society and individuals have more similar grunts within gangs (the
largest group size) than between gangs, and within a party (several parties to a gang)
than between parties (10). In the pygmy marmoset *Cebuella pygmaea*, there is weak
evidence for convergence at the group level and strong evidence for the pair level:
individuals introduced to strangers converge on a trill call duration but not a shared
structure, but they do converge their trill calls in response to pairing with a new mate
(7,23). In general, although vocal convergence at multiple levels of social familiarity may
be widespread, there is not yet a clear pattern in how or why it is used by callers and
receivers.
436
Is vocal convergence an adaptive social behavior or a byproduct of vocal plasticity?
Many authors argue that vocal learning requires complex and therefore costly
neurological adaptations, and is therefore unlikely to evolve unless there is a significant
fitness advantage (48). One possible explanation for vocal convergence in vampire bats
and other echolocating bats is that vocal plasticity evolved for echolocation, and that the
calls of echolocating bats might be altered by their auditory inputs as a byproduct of
audio-vocal feedback mechanisms that function for echolocation. According to this
hypothesis, vocal convergence does not serve a social function, and instead only
results from variation in an individual's exposure to the calls of other individuals.
However, this hypothesis is contradicted by our findings: captive vampire bats that could
always hear all colony members in their cages converged with food-sharing partners
more than other cagemates. By contrast, a simple exposure effect would lead only to
group-level convergence on calls made by the most vocal bats. Instead, we observed
biased vocal convergence towards specific group members depending on the
relationships between them. Future work could further assess this hypothesis by
comparing the influence of social contact (clustering, allogrooming, food sharing) to
acoustic contact alone. Vocal convergence might also have social causes, but not have
social consequences.

An alternative hypothesis is that vocal convergence in vampire bats has social
consequences that might influence the development of cooperative relationships.
Although the functions of vocal convergence remain unclear, it may be a form of
behavioral coordination – a key feature of the formation of many social bonds (e.g.
coordinated courtship behaviors, vocal exchanges and turn-taking, and reciprocal
allogrooming). If partner-specific vocal convergence requires time, it might act as an
honest signal of affiliation and investment. On the other hand, if vampire bats can
rapidly imitate the calls they hear, then call matching may be used to address specific
receivers.
464
Are vampire bats true "vocal learners"?
Definitions of vocal production learning differ among authors. Some authors
include modifying extant call types (often difficult to distinguish from vocal usage
learning, sometimes called "limited vocal learning"), while others emphasize the
acquisition of entirely new call types (the best possible evidence for vocal learning,
sometimes called "complex vocal learning") (25,49–54). Complex vocal production
learning is rare among non-human mammals, but several nonhuman species (including
cetaceans, pinnipeds, elephants and several species of bats) can copy artificial or novel
sounds, including human speech (13,25,49,55–60).
Our findings show that vampire bats are at least 'limited' vocal learners, in the
sense that social interactions cause increased similarity in either call structure, usage,
or both. Although our data are consistent with the hypothesis that adult vampire bats are
capable of vocal production learning, further study on variation in call syllables and
responses to playbacks can assess whether vampire bats are modifying the acoustic
features of existing calls in response to auditory input (19,49,50,52). Interestingly, the
line between convergence in syllable structure versus syllable usage becomes blurred if
contact call variation in vampire bats is not discrete but rather continuously variable.
Future work could further delineate contact calls into different syllable types and assess
the amount of convergence and individuality within these subjectively defined "call
types".
Evidence consistent with vocal learning has been found in at least 11 species of
bats, distributed across eight families (60). Only six of these species have shown *direct*
evidence of vocal learning (17,18,56–58,61,62). In the pale spear-nosed bat
(*Phyllostomus discolor*) adults can shift the fundamental frequency of their tonal social
calls towards a downward-pitch-shifted version of their calls (57), and pups change their
isolation calls to resemble their mother's individual frequency modulation over time (61).
In the greater spear-nosed bat (*Phyllostomus hastatus*) individuals changed frequency
and temporal characteristics of noisy, broadband "screech calls" to converge on a new
group signature as their group composition changed (17). In Taiwanese leaf-nosed bats
(*Hipposideros terasensis*), adult bats in the same colony shifted their echolocation calls
in the same direction gradually throughout the year, and individuals that joined the
colony shifted to match the colony's mean call frequency (18). In the greater horseshoe
bat (*Rhinolophus ferrumequinum*), pups shift towards the frequency of their mother's
echolocation calls more than to other group members (62). In Egyptian fruit bats
(*Rousettus aegyptiacus*), pups raised in normal conditions narrowed their vocal
repertoire to fewer adult vocalizations, while experimentally isolated pups lagged

dramatically in call development or never developed adult calls (58). In the greater sac-winged bat (*Saccopteryx bilineata*), pups of both sexes learn territorial song syllables from adult males (21,56,63). Echolocating bats are specialized for flexibility in vocal production (49,57) and have evolved sophisticated audio-vocal systems that allow them to adapt their echolocation calls in response to acoustic inputs (64–67). Given the exceptional vocal flexibility of bats, the evidence for vocal learning across the bat phylogeny, and the fact that bat vocal repertoires are large and not well understood (65), it is possible that all bats are capable of vocal convergence or even vocal production learning.

Conclusion

Manipulations of co-housing in captivity showed that social familiarity causes convergence in contact call structure in vampire bats. Further vocal convergence occurs among familiar female bats that form close social bonds, reflected by dyadic rates of food sharing. Although captive vampire bats in our studies were acoustically exposed to all colony members in their cages, they appeared to converge with food-sharing partners more than other familiar cagemates, suggesting that vocal convergence is not completely explained by mere exposure. The social causes and consequences of this newly discovered vocal convergence have yet to be fully explored.

Ethics

Procedures with animals were approved by the Institutional Animal Care and Use Committees at the University of Maryland (protocol R-10–63), and the Smithsonian Tropical Research Institute (2015-0915-2018- A9; 2017-0102-2020; 2015-0501-2022), and by the Panamanian Ministry of the Environment (SE/A-76-16 and SEX/A-67-2019).

Data accessibility

Data and R code for reproducing analyses is available on github (68): https://github.com/jkvrtilek/call-convergence-2025

Declaration of AI use

We have not used AI-assisted technologies in creating this article.

Authors' contributions

Julia Vrtilek: data curation, formal analysis, investigation, visualization, writing—original draft, and writing—review and editing; Grace Smith-Vidaurre: data curation, software, formal analysis, investigation, validation, visualization, writing—review and editing; Eric Fosler-Lussier: software, writing—review and editing; Rachel Page: resources, supervision, writing—review and editing; Gerald Carter: conceptualization, data curation, formal analysis, funding acquisition, investigation, methodology, project administration, resources, supervision, visualization, writing—original draft, and writing—review and editing. All authors gave final approval for publication and agreed to be held accountable for the work performed therein.

Competing interests

We declare we have no competing interests.

Funding

This publication is based upon work supported by the National Science Foundation under grant no. IOS-2015928.

Acknowledgements

We thank Gregg Cohen and Imran Razik for help with data collection. We thank May Dixon, Raven Hartman, Virginia Heinen, and Tobias Nguyen for comments that improved the manuscript.

References

1. Soltis J, Leong K, Savage A. African elephant vocal communication I: antiphonal calling behaviour among affiliated females. Anim Behav. 2005 Sep 1;70(3):579–87.

2. Miller CT, Beck K, Meade B, Wang X. Antiphonal call timing in marmosets is behaviorally significant: Interactive playback experiments. J Comp Physiol A. 2009 Aug;195(8):783–9.

3. Carter GG, Fenton MB, Faure PA. White-winged vampire bats (Diaemus youngi) exchange contact calls. Can J Zool. 2009 Jul;87(7):604–8.

4. Balsby TJS, Bradbury JW. Vocal matching by orange-fronted conures (Aratinga canicularis). Behav Processes. 2009 Oct;82(2):133–9.

5. Dahlin CR, Young AM, Cordier B, Mundry R, Wright TF. A test of multiple hypotheses for the function of call sharing in female budgerigars, *Melopsittacus undulatus*. Behav Ecol Sociobiol. 2014 Jan 1;68(1):145–61.

6. Wright TF. Regional dialects in the contact call of a parrot. Proc R Soc B. 1997 Jan;263(1372):867–72.

7. Elowson AM, Snowdon CT. Pygmy marmosets, *Cebuella pygmaea*, modify vocal structure in response to changed social environment. Anim Behav. 1994 Jun 1;47(6):1267–77.

8. Oren G, Shapira A, Lifshitz R, Vinepinsky E, Cohen R, Fried T, et al. Vocal labeling of others by nonhuman primates. Science. 2024 Aug 30;385(6712):996–1003.

9. Zürcher Y, Willems EP, Burkart JM. Are dialects socially learned in marmoset monkeys? Evidence from translocation experiments. PLoS One. 2019 Oct 23;14(10):e0222486.

10. Fischer J, Wegdell F, Trede F, Dal Pesco F, Hammerschmidt K. Vocal convergence in a multi-level primate society: insights into the evolution of vocal learning. Proc Biol Sci. 2020 Dec 23;287(1941):20202531.

11. Langehennig-Peristenidou A, Scheumann M. Sex differences in the impact of social
relationships on individual vocal signatures in grey mouse lemurs (*Microcebus*
murinus). Philos Trans R Soc B. 2024 May 20;379(1905):20230193.

12. Painter MC, Gustison ML, Snyder-Mackler N, Tinsley Johnson E, le Roux A,
Bergman TJ. Acoustic variation and group level convergence of gelada,
Theropithecus gelada, contact calls. Animal Behaviour. 2024 Jan 1;207:235–46.

13. Poole JH, Tyack PL, Stoeger-Horwath AS, Watwood S. Elephants are capable of
vocal learning. Nature. 2005 Mar;434(7032):455–6.

14. King SL, Sayigh LS, Wells RS, Fellner W, Janik VM. Vocal copying of individually
distinctive signature whistles in bottlenose dolphins. Proc R Soc B. 2013 Apr
22;280(1757):20130053.

15. Jones BL, Daniels R, Tufano S, Ridgway S. Five members of a mixed-sex group of
bottlenose dolphins share a stereotyped whistle contour in addition to maintaining
their individually distinctive signature whistles. PLoS One. 2020 May
22;15(5):e0233658.

16. Deecke VB, Barrett-Lennard LG, Spong P, Ford JKB. The structure of stereotyped
calls reflects kinship and social affiliation in resident killer whales (*Orcinus orca*).
Naturwissenschaften. 2010 May 1;97(5):513–8.

17. Boughman JW. Vocal learning by greater spear-nosed bats. Proc R Soc B.
1998;265(1392):227–33.

18. Hiryu S, Katsura K, Nagato T, Yamazaki H, Lin LK, Watanabe Y, et al. Intra-
individual variation in the vocalized frequency of the Taiwanese leaf-nosed bat,
Hipposideros terasensis, influenced by conspecific colony members. J Comp
Physiol A. 2006 Aug 1;192(8):807–15.

19. Tyack PL. Convergence of calls as animals form social bonds, active compensation
for noisy communication channels, and the evolution of vocal learning in mammals.
J Comp Psychol. 2008;122(3):319–31.

20. Barker AJ, Veviurko G, Bennett NC, Hart DW, Mograby L, Lewin GR. Cultural
transmission of vocal dialect in the naked mole-rat. Science. 2021 Jan
29;371(6528):503–7.

21. Knörnschild M, Nagy M, Metz M, Mayer F, von Helversen O. Learned vocal group
signatures in the polygynous bat *Saccopteryx bilineata*. Anim Behav. 2012 Oct
1;84(4):761–9.

22. Lemasson A, Ouattara K, Petit EJ, Zuberbühler K. Social learning of vocal structure
in a nonhuman primate? BMC Evol Biol. 2011 Dec 16;11(1):362.

23. Snowdon CT, Elowson AM. Pygmy Marmosets Modify Call Structure When Paired. Ethology. 1999;105(10):893–908.

24. Briefer EF, McElligott AG. Social effects on vocal ontogeny in an ungulate, the goat, *Capra hircus*. Anim Behav. 2012 Apr 1;83(4):991–1000.

25. Janik VM, Knörnschild M. Vocal production learning in mammals revisited. Philos Trans R Soc B. 2021 Oct 25;376(1836):20200244.

26. Carter GG, Logsdon R, Arnold BD, Menchaca A, Medellin RA. Adult Vampire Bats Produce Contact Calls When Isolated: Acoustic Variation by Species, Population, Colony, and Individual. PLoS One. 2012 Jun 14;7(6):e38791.

27. Carter GG, Wilkinson GS. Common vampire bat contact calls attract past food-sharing partners. Anim Behav. 2016 Jun 1;116:45–51.

28. Carter GG, Farine DR, Crisp RJ, Vrtilek JK, Ripperger SP, Page RA. Development of New Food-Sharing Relationships in Vampire Bats. Curr Biol. 2020 Apr 6;30(7):1275-1279.e3.

29. Carter GG, Wilkinson GS. Food sharing in vampire bats: reciprocal help predicts donations more than relatedness or harassment. Proc R Soc B. 2013 Feb 22;280(1753):20122573.

30. Carter GG, Wilkinson GS. Social benefits of non-kin food sharing by female vampire bats. Proc R Soc B. 2015 Nov 22;282(1819):20152524.

31. Razik I, Brown BKG, Carter GG. Forced proximity promotes the formation of enduring cooperative relationships in vampire bats. Biol Lett. 2022 Apr 6;18(4):20220056.

32. Sueur J, Aubin T, Simonis C, Lellouch L, Aumond P, Baudouin A de, et al. seewave: Sound Analysis and Synthesis [Internet]. 2023 [cited 2025 Feb 22]. Available from: https://cran.r-project.org/web/packages/seewave/index.html

33. Araya-Salas M, Smith-Vidaurre G. warbleR: an r package to streamline analysis of animal acoustic signals. Methods Ecol Evol. 2017;8(2):184–91.

34. Anikin A. soundgen: Sound Synthesis and Acoustic Analysis [Internet]. 2025 [cited 2025 Feb 22]. Available from: https://cran.r-project.org/web/packages/soundgen/index.html

35. Ohio Supercomputer Center [Internet]. Columbus OH: Ohio Supercomputer Center; 1987. Available from: http://osc.edu/ark:/19495/f5s1ph73

36. Bürkner PC. Advanced Bayesian Multilevel Modeling with the R Package brms. The R J. 2018;10(1):395–411.

37. Hart JDA, Weiss MN, Brent LJN, Franks DW. Common permutation methods in
animal social network analysis do not control for non-independence. Behav Ecol
Sociobiol. 2022 Oct 29;76(11):151.

38. Mundry R, Sommer C. Discriminant function analysis with nonindependent data:
consequences and an alternative. Anim Behav. 2007 Oct 1;74(4):965–76.

39. Wilkinson GS. Reciprocal food sharing in the vampire bat. Nature. 1984
Mar;308(5955):181–4.

40. Razik I, Stockmaier S, Abou-Elias M, Carter GG. Opportunistic evidence of the
impact of bacterial infections on social integration in vampire bats. bioRxiv; 2024
https://www.biorxiv.org/content/10.1101/2023.04.17.537180v5

41. King SL, Janik VM. Bottlenose dolphins can use learned vocal labels to address
each other. PNAS. 2013 Aug 6;110(32):13216–21.

42. Balsby TJS, Momberg JV, Dabelsteen T. Vocal Imitation in Parrots Allows
Addressing of Specific Individuals in a Dynamic Communication Network. PLoS
One. 2012 Nov 21;7(11):e49747.

43. Pardo MA, Fristrup K, Lolchuragi DS, Poole JH, Granli P, Moss C, et al. African
elephants address one another with individually specific name-like calls. Nat Ecol
Evol. 2024 Jun 10;1–12.

44. Hile AG, Plummer TK, Striedter GF. Male vocal imitation produces call convergence
during pair bonding in budgerigars, *Melopsittacus undulatus*. Anim Behav. 2000 Jun
1;59(6):1209–18.

45. Bartlett P, and Slater PJB. The effect of new recruits on the flock specific call of
budgerigars (*Melopsittacus undulatus*). Ethol Ecol Evol. 1999 Apr 1;11(2):139–47.

46. Smith-Vidaurre G, Araya-Salas M, Wright TF. Individual signatures outweigh social
group identity in contact calls of a communally nesting parrot. Behav Ecol. 2020 Mar
20;31(2):448–58.

47. Sewall KB. Limited adult vocal learning maintains call dialects but permits pair-
distinctive calls in red crossbills. Anim Behav. 2009 May 1;77(5):1303–11.

48. Nowicki S, Searcy WA. The evolution of vocal learning. Curr Opin Neurobiol. 2014
Oct 1;28:48–53.

49. Knörnschild M. Vocal production learning in bats. Curr Opin Neurobiol. 2014 Oct
1;28:80–5.

50. Tyack PL. A taxonomy for vocal learning. Philos Trans R Soc B.
2019;375(1789):20180406.

51. Jarvis ED. Evolution of vocal learning and spoken language. Science. 2019 Oct
4;366(6461):50–4.

52. Lattenkamp EZ, Vernes SC. Vocal learning: a language-relevant trait in need of a
broad cross-species approach. Curr Opin Behav Sci. 2018 Jun 1;21:209–15.

53. Martins PT, Boeckx C. Vocal learning: Beyond the continuum. PLOS Biol. 2020 Mar
30;18(3):e3000672.

54. Vernes SC, Kriengwatana BP, Beeck VC, Fischer J, Tyack PL, ten Cate C, et al.
The multi-dimensional nature of vocal learning. Philos Trans R Soc B. 2021 Sep
6;376(1836):20200236.

55. Janik VM. Cetacean vocal learning and communication. Curr Opin Neurobiol. 2014
Oct 1;28:60–5.

56. Knörnschild M, Nagy M, Metz M, Mayer F, von Helversen O. Complex vocal
imitation during ontogeny in a bat. Biol Lett. 2009 Oct 7;6(2):156–9.

57. Lattenkamp EZ, Vernes SC, Wiegrebe L. Vocal production learning in the pale
spear-nosed bat, *Phyllostomus discolor*. Biol Lett. 2020 Apr 15;16(4):20190928.

58. Prat Y, Taub M, Yovel Y. Vocal learning in a social mammal: Demonstrated by
isolation and playback experiments in bats. Sci Adv. 2015 Mar 27;1(2):e1500019.

59. Reichmuth C, Casey C. Vocal learning in seals, sea lions, and walruses. Curr Opin
Neurobiol. 2014 Oct 1;28:66–71.

60. Vernes SC, Wilkinson GS. Behaviour, biology and evolution of vocal learning in
bats. Philos Trans R Soc B. 2019;375(1789):20190061.

61. Esser KH, Schmidt U. Mother-Infant Communication in the Lesser Spear-nosed Bat
Phyllostomus discolor (Chiroptera, Phyllostomidae) — Evidence for Acoustic
Learning. Ethology. 1989;82(2):156–68.

62. Jones G, Ransome RD. Echolocation calls of bats are influenced by maternal
effects and change over a lifetime. Proc R Soc B. 1993;252(1334):125–8.

63. Knörnschild M, Behr O, von Helversen O. Babbling behavior in the sac-winged bat
(*Saccopteryx bilineata*). Naturwissenschaften. 2006 Sep 1;93(9):451–4.

64. Fenzl T, Schuller G. Dissimilarities in the vocal control over communication and
echolocation calls in bats. Behav Brain Res. 2007 Sep 4;182(2):173–9.

65. Salles A, Bohn KM, Moss CF. Auditory communication processing in bats: What we
know and where to go. Behav Neurosci. 2019;133:305–19.

66. Jakobsen L, Surlykke A. Vespertilionid bats control the width of their biosonar sound

beam dynamically during prey pursuit. PNAS. 2010 Aug 3;107(31):13930–5.

67. Gillam E, Fenton MB. Roles of Acoustic Social Communication in the Lives of Bats.
In: Fenton MB, Grinnell AD, Popper AN, Fay RR, editors. Bat Bioacoustics. New
York, NY: Springer; 2016. p. 117–39. (Springer Handbook of Auditory Research).

68. Vrtilek JK, Carter G. call-convergence-2025 [Internet]. 2025 [cited 2025 Mar 30].
Available from: https://github.com/jkvrtilek/call-convergence-2025

69. Ripperger SP, Carter GG, Duda N, Koelpin A, Cassens B, Kapitza R, et al. Vampire
Bats that Cooperate in the Lab Maintain Their Social Networks in the Wild. Curr Biol.
2019 Dec 2;29(23):4139-4144.e4.

70. Araya-Salas M, Smith-Vidaurre G. warbleR: Streamline Bioacoustic Analysis
[Internet]. 2025 [cited 2025 Feb 22]. Available from: https://cran.r-
project.org/web/packages/warbleR/index.html

731
732

Supplementary Information (SI)

Supplementary Methods

Variation in contact call structure
We defined contact calls based on a combination of behavioral context and call
features. Specifically, all contact calls that we analyzed were produced by a bat that
was isolated from group members, and the calls had durations of 3-50 ms and had peak
frequencies of 10-30 kHz. To remove ambiguous cases, we also removed calls for
which measurements of the fundamental frequency failed (which decreased the number
of sampled bats by 3 males and 1 female), because these calls either had poor signal-
to-noise ratio or they lacked clear tonal structure. Removing them increased the overall
call classification rate for assigning calls to bat.
Within these call parameters, we failed to find any clear evidence for discrete call
types despite considerable variation both among and within callers (Figure S1). Given
that contact call variation appeared to be graded (continuous) rather than discrete, we
decided to take the conservative first step of assessing similarity for all contact calls
rather than subjectively defining subtypes and measuring similarity within each type.
The latter approach would also elevate the false discovery rate due to multiple testing
and the many possible ways to label types. Future work can assess the degree to which
contact call variation is actually continuous rather than discrete, and describe how vocal
individuality differs between structurally defined types of syllables.
755
756

Figure S1. Representative variation in contact calls within and among individual vampire bats. Spectrograms of example contact calls from four female vampire bats (row) across 4 different nights of being individually isolated. Each row represents an individual; the top row is from 2011-2013, the second row is from 2016-2017, and the bottom two rows are from 2019. Recordings in a row are in order from earliest to latest. The variable structure of contact calls suggest that they might convey other information beyond the caller's individual identity, such as motivational state.

766
Figure S2. Causal model for vocal convergence among all pairs. Arrows are causal
paths. Black arrows are causal effect of interest. Grey arrows are causal paths that
cause potential confounding. Greater kinship causes greater affiliation in vampire bats
(A, "affiliation" = allogrooming lograte + food-sharing lograte) and it might also cause
greater similarity in contact calls (B), if calls are heritable. Affiliation might cause vocal
convergence (C), but affiliation is also dependent on co-housing (D), and the mere
acoustic exposure due to co-housing might also cause vocal convergence (E).
774

Model 1 posterior predictive check

Model 2 posterior predictive check

Model 3 posterior predictive check

Model 4 posterior predictive check

Model 5 posterior predictive check

Figure S3. Posterior predictive checks for 5 Bayesian models. Comparison of distribution of observed contact call similarity values (dark black line) to the predicted values from 100 datasets simulated from the fitted model (light blue lines) suggests that the models capture the essential features of the data-generating process.

781
782 **Figure S4. Timeline of call recordings for analysis of call convergence within-**
783 **pairs over time.** Top panel: recording dates (dots) for 5 bats (lines) captured from Las
784 Pavas, Panama. Bottom panel: recording dates for 7 bats from Tolé, Panama. Dotted
785 red line: date of introduction between Tolé and Las Pavas bats in small groups (2-5
786 bats). Solid red line: date of introduction between all captive bats into large flight cage.
787
788

789 **Table S1. Bats recorded for this study.** Of the 116 bats recorded for this study, 94
790 produced more than 80 contact calls. The number of bats below adds to 96 because 2
791 bats from the 2016-2017 colony were recaptured and included in the 2019 colony.

Years recorded	# bats	Familiarity	Kinship	Origin	Age and Sex
2011-2014	29	Long-term captive colony (27,29,30)	Previously estimated using maternal pedigree and microsatellite markers (30)	Multiple zoos	11 adult F 4 subadult F 5 adult M 9 subadult M
2016	4	Never caged with or introduced to any other bats in the study, but potentially familiar	Unrelated to other bats, but unknown relatedness among each other	Cattle pasture, Chilibre, Panama	1 adult F 3 adult M
2016	1	Never caged with or introduced to any other bat in the study	Unrelated to other bats	Forest, Gamboa, Panama	1 adult M
2016-2017	40	Co-housed (caged together) for over 10 months (28,69)	Previously estimated using microsatellite markers (69); bats from different sites are unrelated	Cattle pasture, Las Pavas, Panama	5 adult F
				Captive-born to Las Pavas bats	1 subadult F 1 subadult M
				Tree hollow, Tolé, Panama	19 adult F 7 adult M
				Captive-born to Tolé bats	6 subadult F 1 subadult M
2019	22	Co-housed for 4 months (31)	Bats from different sites are unrelated; kinship unknown for bats from the same site	Same tree, Tolé, Panama	2 recaptured adult F 7 new adult F
				Tree hollow, La Chorrera, Panama	7 adult F
				Cave, Lake Bayano, Panama	6 adult F

792
793

Table S2. Spectral and temporal measurements used in linear discriminant
function analyses. Duration through meanpeakf are slightly edited from the warbleR
(70) and seewave (32) documentation. Maxslope through segments are summary
measures of fundamental frequency measurements from the *soundgen* package (34).
Time interquartile range is calculated by subtracting time quantile 25 from time quantile
75. Frequency interquartile range is calculated by subtracting frequency quantile 25
from frequency quantile 75. Standard deviation of frequency is calculated from
frequency range. Dominant frequency range is calculated from maximum and minimum
dominant frequency. Dominant frequency slope is calculated from start dominant
frequency, end dominant frequency and duration.
804

Measure	Explanation
duration	Length of signal (in ms)
meanfreq	Mean of frequency spectrum (i.e. weighted average of frequency by amplitude within supplied band pass) (in kHz)
sd	Standard deviation of frequency (in kHz).
freq.median	The frequency at which the signal is divided in two frequency intervals of equal energy (in kHz)
freq.Q25	The frequency at which the signal is divided in two frequency intervals of 25% and 75% energy respectively (in kHz)
freq.Q75	The frequency at which the signal is divided in two frequency intervals of 75% and 25% energy respectively (in kHz)
freq.IQR	Frequency range between 'freq.Q25' and 'freq.Q75' (in kHz)
time.median	The time at which the signal is divided in two time intervals of equal energy (in s)
time.Q25	The time at which the signal is divided in two time intervals of 25% and 75% energy respectively (in s).
time.Q75	The time at which the signal is divided in two time intervals of 75% and 25% energy respectively (in s).
time.IQR	Time range between 'time.Q25' and 'time.Q75' (in s).
skew	Asymmetry of the frequency spectrum
kurt	Peakedness of the frequency spectrum
sp.ent	Energy distribution of the frequency spectrum
time.ent	Energy distribution on the time envelope
entropy	Product of sp.ent and time.ent
sfm	Spectral flatness
meandom	Average of dominant frequency measured across the acoustic signal
mindom	Minimum of dominant frequency measured across the acoustic signal
maxdom	Maximum of dominant frequency measured across the acoustic signal
dfrange	Range of dominant frequency measured across the acoustic signal
modindx	Modulation index, calculated as the cumulative absolute difference between adjacent measurements of dominant frequencies divided by the dominant frequency range
startdom	Dominant frequency measurement at the start of the signal
enddom	Dominant frequency measurement at the end of the signal
dfslope	Slope of the change in dominant frequency through time (kHz/s)
peakf	Frequency with the highest energy from the frequency spectrum
meanpeakf	Frequency with highest energy from the mean frequency spectrum (the mean relative amplitude of the frequency distribution of a time wave)
maxslope	Maximum slope of fundamental frequency contour
minslope	Minimum slope of fundamental frequency contour
absminslope	Absolute value of the minimum slope of fundamental frequency contour

pos.slopes	Number of positive slopes in the fundamental frequency contour
neg.slopes	Number of negative slopes in the fundamental frequency contour
turns	Number of turns (changes between positive and negative slopes) in the fundamental frequency contour
meanslope	Mean slope of the fundamental frequency contour
segments	Number of separate segments in the fundamental frequency contour

Supplementary Results

808
Figure S5. Kinship predicts contact call similarity across all pairs with known
kinship. Left panel shows how call similarity increases with kinship. Blue line shows a
general additive model in the ggplot2 R package. Right panel shows mean contact call
similarity with bootstrapped 95% confidence intervals (CIs) for three bins of kinship. The
95% CIs treat pairs as independent observations, so inferences should be drawn from
the multi-membership model results (Figure 2).
815

816
Figure S6. Contact call similarity varies by levels of familiarity. Top panel shows
the distributions and boxplots of the contact call similarity, and bottom panel shows
means with bootstrapped 95% CIs of the contact call similarity, at five conditions
(degrees) of familiarity (x-axis) including (1) unfamiliar nonkin that never met (baseline
level of call similarity among strangers), (2) previously unfamiliar nonkin captured from
different wild roosts then co-housed in captivity, (3) nonkin captured from the same
roost then co-housed together in captivity, and (4) familiar nonkin co-housed in the
same long-term captive colony. Vocal convergence is demonstrated by the difference
from the pairs that never met (horizontal line). The 95% CIs treat pairs as independent
observations, so inferences should be drawn from the multi-membership model results
(Figure 2).
828

829
Figure S7. The association between nonkin affiliation rate and nonkin contact call similarity differed across colonies. All pairs were nonkin (r <0.05). The 2014 dataset included only food donations to fast bats in a long-term captive colony (left panel). The 2017 and 2019 colonies were shorter term colonies that merged samples of bats from different wild populations. The 2017 colony included mostly allogrooming and some food donations (orange). The 2019 dataset included only allogrooming in pairs from different sites (purple), because kinship was unknown in pairs from the same site (but including kin pairs does not change the conclusions from these analyses). The 95% CIs on slopes treat pairs as independent observations, so inferences should be drawn from the multi-membership model results (Figure 2).

841
Figure S8. Nonkin pairs that engage in more food sharing have more similar
contact calls. Greater food-sharing predicted greater contact call similarity among
nonkin females in the 2014 long-term stable colony and possibly the 2017 merged
colony (A). The absence and presence of food sharing is linked to greater contact call
similarity in nonkin females (B) and in all nonkin pairs (C). The 95% CIs treat pairs as
independent observations, so inferences should be drawn from the multi-membership
model results (Figure 2).
849
850

851 **Table S3. Model coefficients.** Other columns are the estimate of the average
852 (expected) posterior probability, the estimated error (the posterior standard deviation),
853 the 95% credible interval. The Rhat convergence diagnostic (convergence = 1.00) and
854 the Bulk and Tail effect sample sizes.

Model	Term	Estimate	Est. Error	Lower 95% CI	Upper 95% CI	R-hat	Bulk ESS	Tail ESS
1	kinship (scaled)	0.036	0.005	0.026	0.046	1.000	16126	12499
2	affiliation log rate* (scaled)	0.023	0.006	0.011	0.035	1.001	11994	12756
2	kinship (scaled)	0.003	0.005	-0.008	0.013	1.000	18253	12880
2	co-housed (T/F)	0.143	0.013	0.117	0.168	1.000	12402	12363
3	co-housed (T/F)	0.195	0.015	0.164	0.225	1.000	14074	11694
4	affiliation log rate* (scaled)	-0.006	0.020	-0.044	0.033	1.000	9096	9975
5	food sharing log rate* (scaled)	0.074	0.024	0.027	0.122	1.000	11509	11622

855 *Log rate is (natural log (seconds of observation + 1)) / natural log (sampled hours).