Enterobacter sp. SA187-induced coordinated regulation of high-affinity nitrate transporters

and ethylene signaling enhances nitrogen content and plant growth under low nitrate

Amina Ilyas[1,2], Caroline Mauve[1,2], Jose Caius[1,2], Christine Paysant-Leroux[1,2], Michael Hodges[1,2],

Axel de Zelicourt[1,2*]

Affiliations :

1. Université Paris-Saclay, CNRS, INRAE, Université Evry, Institute of Plant Sciences Paris-

Saclay (IPS2), 91190, Gif-sur-Yvette, France.

2. Université Paris Cité, CNRS, INRAE, Institute of Plant Sciences Paris-Saclay (IPS2), 91190,

Gif-sur-Yvette, France

* Correspondence: axel.de-julien-de-zelicourt@universite-paris-saclay.fr

11 **Abstract**

Sustainable crop production demands solutions to reduce the over use of synthetic nitrogen

fertilizers, and plant-growth-promoting bacteria (PGPB) offer a promising alternative by

enhancing nutrient acquisition under limiting conditions. This study shows that the endophytic

bacterium _Enterobacter_ sp. SA187 significantly improved the growth of _Arabidopsis thaliana_

seedlings under low nitrate conditions. SA187 inoculation led to marked increases in biomass,

primary root length, and lateral root density, with growth-promoting effects becoming more

important with the severity of nitrate limitation. Analyses suggested enhanced nitrate uptake and

improved nitrogen partitioning as indicated by reduced shoot carbon:nitrogen ratios.

Transcriptomic profiling identified SA187-driven reprogramming of host gene expression,

including the induction of high-affinity nitrate transporters (HATs) and hormone-responsive

pathways. Functional analyses confirmed that the growth-promoting effect of SA187 required

both ethylene signaling and specific HATs. The _ein2_, _nrt2.5_, and _nrt2.6_ mutants failed to show

SA187-induced growth promotion, and expression analysis revealed that *NRT2.5* was regulated

downstream of ethylene signaling, while *NRT2.6* was partially independent. These findings

position SA187 as a conditionally beneficial endophyte that can improve plant growth through

coordinated, context-dependent regulation of nitrate uptake and allocation. Such results highlight

the potential of SA187 as a bioinoculant for improving plant performance under nitrate-limited

conditions and thus reduce fertilizer dependency.

I. Introduction

Nitrogen (N) is essential for plant growth, development, and productivity. As a core component of nucleic acids, proteins, amino acids, chlorophyll, and secondary metabolites, N coordinates numerous physiological and metabolic processes (Krapp, 2015; O'Brien et al., 2016). Consequently, N-deficiency results in chlorosis, reduced photosynthetic capacity, stunted growth, and yield losses (Ding et al., 2005; Hawkesford et al., 2012; Ahmad Wani et al., 2022). Although atmospheric nitrogen (N_2) is abundant, most plants cannot directly use this inert form, instead they rely on bioavailable N-compounds such as nitrate (NO_3^-), ammonium (NH_4^+), and amino acids. Among these, NO_3^- generally predominates as the available N-form in well-aerated soils (Glass, 2009; Krapp et al., 2014). However, the availability of bioavailable N-compounds is often limited in agricultural systems thus making this a significant factor limiting crop yields worldwide (LeBauer and Treseder, 2008; Sinclair and Rufty, 2012).

To alleviate N-limitations and ensure global food security, agriculture has relied on synthetic N-based fertilizers. While this has significantly improved crop yields, their use remains inefficient, costly and they are a source of greenhouse gases and pollution. N-use-efficiency (NUE) remains remarkably low as less than 50% of applied N is taken up by plants, while the remainder is lost to volatilization, leaching, and denitrification. These losses not only reduce productivity and economic returns for farmers but also cause widespread environmental degradation (Liu et al., 2010; Ali et al., 2025). Adverse impacts include groundwater nitrate contamination, eutrophication, greenhouse gas emissions (such as N_2O), ozone depletion, and long-term soil degradation, with serious consequences for ecosystems and human health (FAO and UNEP, 2021; Sutton et al., 2021; Nath et al., 2023; Govindasamy et al., 2023; de Vries et al., 2024). In parallel, the production of N-based fertilizers via the Haber–Bosch process is energy-intensive

and carbon (C)-emitting, adding further environmental and economic burdens (Cherkasov et al.,

2015; Rouwenhorst and Lefferts, 2020). Therefore, there is a need to improve plant N-uptake and

N-utilization so as to support plant growth under low-N-input conditions in the context of an

environmentally-friendly, sustainable agriculture.

Plant-growth-promoting bacteria (PGPB) have emerged as promising biological tools to reduce

reliance on synthetic inputs. Historically, research has extensively focused on symbiotic N_2-

fixing bacteria (e.g., *Rhizobia*, *Frankia* spp.). However, emerging studies demonstrate that even

non-N_2-fixing endophytic bacteria can significantly improve plant growth directly, by enhancing

nutrient availability, mobilizing soil minerals such as phosphorus, potassium, iron and N, and

indirectly, by modulating hormonal signaling pathways, root architecture, defense and systemic

stress responses (Bakhshandeh et al., 2020; De Zelicourt et al., 2013; Di Benedetto et al., 2017;

Etesami & Adl, 2020; Glick, 2012, 2020; Pathania et al., 2020; Ramakrishna et al., 2019).

Despite increasing interest, the mechanistic details underlying these beneficial plant–microbe

interactions, particularly under N-limiting conditions, remain insufficiently characterized.

Among promising plant-associated endophytes, *Enterobacter* sp. SA187 (SA187) has emerged as

a particularly effective beneficial endophytic bacterium. Initially isolated from the desert-adapted

legume *Indigofera argentea* (Andrés-Barrao et al., 2017), SA187 effectively enhances resilience

across a wide spectrum of abiotic stresses, including drought, salinity, heat, and high light. The

beneficial effects of SA187 are mediated primarily through modulation of critical physiological

and molecular processes such as hormonal homeostasis, particularly involving ethylene signaling

pathways, maintenance of nutrient and redox balances, and activation of stress-responsive

networks (de Zélicourt et al., 2018; Rahman et al., 2025; Rolli et al., 2022; Shekhawat et al.,

2024; Shekhawat et al., 2021). Although previous studies have demonstrated plant growth

improvement by *Enterobacter* species under nutrient-limited conditions, through mechanisms

such as N_2-fixation, improved nutrient acquisition and altered root system architecture (Shankar

et al., 2011; Lin et al., 2012; Madhaiyan et al., 2013; Zhu et al., 2013; Tsegaye et al., 2022), the

role of SA187 under low N conditions has not been investigated.

Nitrate uptake and allocation within higher plants are primarily regulated by specialized nitrate

transporters belonging to two main families: the NRT1/PTR family (NPF) comprising mostly

low-affinity transporters (LATs), and the NRT2 family of high-affinity nitrate transporters

(HATs). NRT2 family members are important under low nitrate availability conditions since they

transport nitrate at very low external concentrations (Krapp et al., 2014; Zhang et al., 2024).

HATs regulate not only soil-derived nitrate uptake but also nitrate distribution within plant

tissues and organs, significantly influencing NUE and plant adaptation under nitrate scarcity

(Wang et al., 2012; Lezhneva et al., 2014; Fan et al., 2017). Moreover, hormonal signals, notably

ethylene, have been implicated in modulating the expression and functional activity of nitrate

transporters (Tian et al., 2009; Zhang et al., 2014; Zheng et al., 2013). Ethylene signaling,

mediated through EIN2 and EIN3, plays a central role in integrating nutrient and environmental

cues, but its role in regulating nitrate acquisition via plant–microbe interactions remains poorly

understood (Iqbal et al., 2017; Khan et al., 2017; Leblanc et al., 2008). Although SA187 is

known to modulate ethylene signaling to enhance stress tolerance (de Zélicourt et al., 2018;

Shekhawat et al., 2024), to date its specific role on nitrate transporter regulation and plant N-

metabolism has not been explored. Since ethylene is known to interact with several nitrate

transporters (including NRT2.1) (Tian et al., 2009; Zheng et al., 2013), it is possible that SA187-

mediated modulation of ethylene signaling could significantly influence nitrate uptake and

nitrate acquisition under low nitrate conditions.

This study investigated the beneficial impact of SA187 on the growth of *Arabidopsis thaliana*

seedlings with respect to nitrate concentration, with a focus on growth responses, nitrate

transporter expression, nitrate content, C:N ratios and ethylene signaling. Using phenotypic,

elemental, transcriptomic, and genetic analyses, including HATs and ethylene-insensitive

mutants, this work revealed that SA187 appeared to improve nitrate uptake and aerial allocation

through a combination of ethylene-dependent and -independent regulatory pathways. These

findings provide mechanistic insights into how a beneficial endophyte can modulate plant

nutrient physiology, offering a path forward for microbial strategies to potentially enhance NUE

under N-limited conditions.

II.   **Materials and Methods**

A.   *Bacterial and plant material*

Arabidopsis thaliana (L.) Heynh., ecotype Columbia-0 (Col-0, wild-type), and selected mutants

in a Col-0 background: *ein2-1* (Guzmán and Ecker, 1990), *nrt2.5* (GK 213H10), *nrt2.6* (SM

3.35179), and a *nrt2.5 x nrt2.6* double mutant (Kechid et al., 2013) were used in this study. The

Enterobacter sp. SA187 strain, originally isolated from root nodules of *Indigofera argentea* in

Jizan, Saudi Arabia (Andrés-Barrao et al., 2017), was used in its GFP-tagged form (de Zélicourt

et al., 2018).

B.   *Plant growth and bacterial colonization*

Arabidopsis thaliana seeds were surface-sterilized and inoculated with SA187 as described

previously (Saad et al., 2018). Briefly, sterilized seeds were sown on modified half-strength

Murashige and Skoog (½MS) medium (Apollo Scientific Catalogue No: PMM531B), containing

basal salts only (0.5 g/L MES, pH 5.8, 9 g/L agar), under different nitrate regimes ranging from

extremely-low to high-nitrate *i.e.* 0.2, 1, 2, 10, 20 mM KNO_3. KCl was added to adjust the

potassium concentration in the growth media so that their composition was identical, except for

the variation in nitrate levels. For SA187 treatments (+SA187), 2×10^5 bacteria/mL were added

directly to the growth media before pouring into square petri plates. For mock controls (Mock),

an equal volume of sterile LB medium was added instead. After stratification for 24 h at 4°C in

the dark, plates were placed vertically in a growth chamber (16-h photoperiod, 18–20°C

night/day, 120 μE/m^2/s light intensity, 50 ± 10% relative humidity) to allow seed germination.

At 6 days post-germination (dpg), seedlings were transferred to fresh modified ½MS agar plates

containing the same nitrate concentrations as in the germination plates and harvested at 16 days

for fresh weight (FW) measurements, nitrate quantification and, molecular and elemental

analyses. Plates were scanned at 16dpg to obtain the images for primary root length (PRL) and

lateral root density (LRD) measurements.

C.   *Growth phenotyping*

FW of individual plants was measured at harvest. PRL and LRD were measured from plate

scans using Fiji/ImageJ software (Schindelin et al., 2012).

D.   *Elemental analyses*

Dried root and shoot tissues (~1 mg per sample) were transferred to tin capsules, weighed and

combusted in a Pyrocube elemental analyzer (Elementar, Lyon, France). C and N contents were

determined via thermal conductivity detection (TCD) after gas separation on selective columns.

Quantification was performed against certified standards: ammonium sulfate (21.2% N), sucrose

(42.11% C), glutamic acid (9.51% N, 40.82% C), and glutamine (19.17% N, 41.09% C). Data

are presented as % dry weight (DW).

E.   *Nitrate content assay*

Flash-frozen tissues (≥10 mg FW) were ground under liquid N. Each sample was extracted with

500 µL preheated Milli-Q water (100°C), vortexed, and boiled for 15 min with intermittent

mixing. After cooling and centrifugation (15,000×g, 15 min), 450 µL of the supernatant was

collected. Nitrate content was measured colorimetrically using the Nitrite/Nitrate Assay Kit

(Sigma-Aldrich, #23479), following the manufacturers protocol. Technical triplicates were

performed per sample.

F.   Transcriptomics

i.   RNA-sequencing and data analyses

Shoot and root samples were collected from 16-day-old mock and SA187-treated plants grown at

20 mM and 1 mM nitrate, with three biological replicates per condition. Each replicate

comprised pooled shoot or root tissues from 4–6 plants. Total RNA was extracted using the

NucleoSpin RNA kit following the manufacturer's recommendations (Macherey-Nagel). RNA-

seq libraries were prepared from 500 ng total RNA using the QuantSeq 3′ mRNA-Seq Library

Prep Kit (Lexogen) and sequenced (75 bp single-end) on an Illumina NextSeq500.

Unique molecular identifiers (UMIs) were removed and appended to the read identifier with the

extract command of UMI-tools (v1.0.1) (Smith et al., 2017). Reads where any UMI base quality

score fell below 10 were removed. To remove adapter sequences, poly(A), poly(G) sequences,

and low quality nucleotides, reads were trimmed with BBduk from the BBmap suite (v38.84)

(Bushnell, 2014) with the options k=13 ktrim=r useshortkmers=t mink=5 qtrim=r trimq=10

minlength=30. Trimmed reads were then mapped onto the reference genome of *Arabidopsis*

thaliana (TAIR10+ARAPORT11+MtCol0) and counted using STAR (v2.7.3a) (Dobin et al.,

2013), with the following parameters --alignIntronMin 5 --alignIntronMax 60000 --

alignMatesGapMax   6000   –alignEndsType   Local--outFilterMultimapNmax   20   --

outFilterMultimapScoreRange 0 –outSAMprimaryFlag AllBestScore –mismatchNoverLmax 0,6.

Reads with identical mapping coordinates and UMI sequences were collapsed to remove PCR

duplicates using the dedup command of UMI-tools with the default directional method

parameter. Deduplicated reads were counted using HTSeq version v0.12.4 (Anders et al., 2015)

(htseq-count mode intersection-nonempty) based on the *A. thaliana* TAIR10-ARAPORT11

annotation. Between 2.9 (86.6% of raw reads) and 2.5 (95% of raw reads) millions of reads were

associated to annotated genes.

Statistical analyses were conducted with R v3.6.2 (R Core Team, 2020) using DiCoExpress V1

(Lambert et al., 2020; Baudry et al., 2022) based on the Bioconductor package edgeR (v 3.28.0)

(Robinson et al., 2010; McCarthy et al., 2012). Two separate statistical analyses were conducted

on data from roots and leaves, starting from the raw count tables.

For each analysis, genes with low counts were filtered using the "filterByExpr" function where

the group argument specifies the biological conditions, the min.count value was set to 3 and the

min.total.count, large.n and min.prop arguments were set to their default values. Libraries were

normalized using the TMM method.

For both analyses, the differential analysis was based on a negative binomial generalized linear

model in which the logarithm of the average gene expression is an additive function of a nitrate

effect (1mM or 20mM), an inoculation effect (SA187 or mock), their interaction and a replicate

effect (3 modalities). Four comparisons were performed (i) the difference between 1 mM and 20

mM nitrate in the mock condition, (ii) the difference between 1 mM and 20 mM nitrate in the

SA187 inoculated condition, (iii) the difference between mock and SA187 inoculated at 1 mM

nitrate, and (iv) the difference between mock and SA187 inoculated at 20 mM nitrate. For each

contrast a likelihood ratio test was applied and raw p-values were adjusted with the Benjamini–

Hochberg procedure to control the false discovery rate. The distribution of the resulting p-values

followed the quality criterion described by Rigaill et al. (2016). A gene was declared

differentially expressed if its adjusted p-value was lower than 0.05.

Hierarchical clustering of differentially expressed genes (DEGs) was performed using the MeV

software (v4.9) with Pearson correlation and average linkage to identify SA187- and nitrate-

responsive gene expression clusters. GO term enrichment analysis of DEG clusters was

conducted using the R package clusterProfiler (Yu et al., 2012). Terms with adjusted p-values

(FDR) < 0.05 were considered significant.

198   *ii.   qRT-PCR*

cDNA synthesis was performed using the ImProm-II™ Reverse Transcription System (Promega)

with oligo-dT primers. Quantitative real-time PCR was performed using SYBR Green I Master

Mix (Roche) on a Bio-Rad CFX384 Real-Time PCR Detection System. Each reaction was run

with technical triplicates. Gene expression was normalized to *ACTIN2* and *YSL8*, and relative

expression levels were calculated using the $\Delta\Delta$Ct method (Livak and Schmittgen, 2001).

G.   Statistical analyses and visualization

qRT-PCR data were analyzed using CFX Manager software version 2.1 (Bio-Rad). All other

statistical analyses were performed in R (v4.2.1). Growth parameters (FW, PRL, and LRD) were

compared using two-tailed unpaired Student's *t*-tests. Elemental composition data (C, N, and

C:N ratio) and nitrate contents were analyzed using the non-parametric Mann–Whitney U test.

Data visualization, including boxplots and bar graphs with error bars, was carried out using

ggplot2, ggpubr, and rstatix packages in R. A threshold of $p < 0.05$ was used to determine

statistical significance, with significance levels indicated directly on plots.

III.   Results

A.   Effect of SA187 on plant growth under varying nitrate conditions

To evaluate the impact of SA187 inoculation on seedling growth with respect to nitrate

concentration, *Arabidopsis thaliana* seedlings were grown on solid media containing various

concentrations of nitrate (20, 10, 2, 1, and 0.2 mM NO_3^-), with or without SA187 inoculation.

Growth responses were quantified by measuring total FW (TFW), PRL, and LRD (Figure 1a–c).

At high nitrate availability (20 mM), SA187 inoculation had no significant effect on TFW and

PRL compared to mock controls (Figure 1a, b), however a significant increase in LRD (+27.2%)

was observed (Figure 1c). As external nitrate concentration declined, SA187-mediated growth

enhancements became progressively more pronounced. At 10 mM, SA187 treatment

significantly increased TFW by 83.1%, PRL by 18.1%, and LRD by 32.9% (Figure 1a–c) and

similar responses were seen at 2 mM nitrate, with SA187 increasing TFW by 90%, PRL by

13.4%, and LRD by 30.8%. At 1 mM nitrate, the beneficial effects of SA187 further intensified

for TFW (+108.5%) and PRL (+29.3%), but the LRD increase (+32%) remained comparable to

that observed at 2 and 10 mM (Figure 1a–c). Under a severe nitrate limitation (0.2 mM), the

effect of SA187 inoculation was exceptionally pronounced, leading to increases of 295.5%

(TFW), 121.6% (PRL), and 274.9% (LRD) compared to mock controls (Figure 1a–c). Taken

together, these results indicate that SA187 substantially improves both shoot and root

development, specifically when nitrate availability is low, with the magnitude of the beneficial

effects inversely correlated with external nitrate levels.

B.   Effect of SA187 on C, N, and nitrate contents

To investigate how SA187 affected nitrate acquisition and internal nitrate partitioning, total C,

total N, the C:N ratio, and nitrate content were measured in both shoots and roots of *Arabidopsis*

thaliana grown under high (20 mM) and low (1 mM) nitrate conditions, with or without SA187

inoculation (Figure 2a–d). SA187 inoculation did not significantly alter shoot C content under

either nitrate condition, but slightly reduced root C content at low nitrate (Figure 2a). N content

exhibited tissue-specific responses: SA187 significantly increased shoot N content but decreased

root N content under low nitrate conditions (Figure 2b). However, no significant changes in N

content were observed under high nitrate conditions (20 mM) (Figure 2b). These variations in C

and N content led to a significantly lower C:N ratio upon SA187 inoculation at 1 mM nitrate,

both in shoots and in roots (Figure 2c). Most notably, SA187 induced a significant increase in

nitrate content under low nitrate (1 mM) in both shoots and roots when compared to controls

(Figure 2d). At high nitrate conditions (20 mM), on the other hand, SA187 had no significant

effect on nitrate content. Collectively, these findings emphasize the role of SA187 inoculation in

improving nitrate uptake, and shifting N-allocation and nitrate translocation towards aerial

tissues under conditions of low nitrate. Such SA187-dependent changes could contribute to the

observed increase in seedling growth.

C.   *SA187 mediated transcriptomic reprogramming of Arabidopsis shoots and roots*

To gain insights into the molecular basis of SA187-mediated growth enhancement in the

presence of low nitrate, transcriptome analyses (RNA-seq) were performed on Arabidopsis

shoots and roots under high (20 mM) and low (1 mM) nitrate conditions with and without SA187

inoculation. Differential gene expression analysis identified 2,280 and 1,005 DEGs in shoots and

roots, respectively (Supplementary Figure S1, Supplementary Table S2 and Table S3).

Hierarchical clustering and Gene Ontology (GO) enrichment analyses were conducted to

categorize the DEGs based on their expression patterns and functional annotations.

In shoots, hierarchical clustering revealed distinct expression patterns. Cluster 4 (517 genes), the

largest shoot cluster, showed significant gene expression induction by SA187 in both nitrate

conditions, although predominantly under high nitrate, and included genes associated with

defense, immune response, oxidative stress, ethylene, salicylic acid (SA) signaling and metabolic

processes, jasmonic acid (JA), cell wall thickening and response to N-compounds. Cluster 10

(239 genes) exhibited a nitrate-dependent regulation, being upregulated under low nitrate alone

but downregulated by SA187 treatment under low nitrate conditions. This cluster was enriched in

genes linked to JA response, oxidative stress, amino acid metabolism, flavonoid biosynthesis,

and glutathione metabolism. Genes in cluster 3 (81 genes) and cluster 9 (284 genes)

demonstrated SA187-specific induction regardless of nitrate concentration. These clusters

included genes involved in systemic resistance, response to N-compounds, lateral root

development, auxin and ethylene signaling, and camalexin biosynthesis (Fig. 3a).

Root transcriptome profiling also revealed distinct clusters of gene expression. Cluster 3 (417

genes), the largest root cluster, exhibited down-regulation under low nitrate but pronounced

induction by SA187 across both nitrate conditions. Genes in this cluster were primarily related to

defense, ethylene, JA, SA, auxin, cytokinin signaling, and oxidative stress responses. Cluster 5

(234 genes) showed an inverse expression pattern to cluster 3, being upregulated under low

nitrate but strongly downregulated by SA187 at both nitrate concentrations. It included genes

involved in amino acid transport and metabolism, N-signaling, and oxidative stress responses.

Notably, cluster 11 (151 genes) was selectively induced by SA187 under low nitrate and enriched

for genes involved in root morphogenesis, nitrate import, and responses to reactive N-species

(Figure 3b), suggesting a condition-specific program optimized for nutrient acquisition under

stress. Targeted examination of HATs highlighted a strong SA187-induced expression of *NRT2.5*

(At1g12940) in shoots, and *NRT2.4* (At5g60770) and *NRT2.6* (At3g45060) in roots, under low

nitrate conditions (Fig. 3c).

Together, these findings suggested that SA187 orchestrated a transcriptional network integrating

nutrient transport, hormone signaling, stress buffering, and developmental remodeling to

promote plant adaptation under a low nitrate condition.

D.   Role of high-affinity nitrate transporters NRT2.5, NRT2.6 and ethylene

signaling in SA187-mediated growth promotion

To evaluate the functional requirement of ethylene signaling and specific HATs in SA187-

induced plant growth benefits under low nitrate conditions, phenotypic analyses were performed

using ethylene-insensitive (*ein2-1*), HAT single (*nrt2.5* , *nrt2.6*) and double (*nrt2.5 × nrt2.6*)

mutants under low nitrate (1 mM). Wild-type plants (Col-0) exhibited substantial growth

enhancement upon SA187 inoculation, as expected. However, none of the tested mutants

displayed significant growth improvements upon SA187 inoculation, indicating that both nitrate

transporters and a functional ethylene signaling were indispensable for the growth-promoting

effects of SA187 under low nitrate (Figure 4a).

To further dissect the regulatory link between ethylene signaling and nitrate transporter

expression, transcript levels of six HAT genes (*NRT2.1–NRT2.6*) were quantified by qRT-PCR in

shoots and roots of Col-0 and *ein2-1* plants at 1mM nitrate, with and without SA187 inoculation

(Figure 4b). NRT2.7 was excluded due to a consistently undetectable expression. In mock-

inoculated *ein2-1* plants, baseline expression of *NRT2.3*, *NRT2.4*, and *NRT2.6* was higher in

shoots compared to Col-0, while in roots, *NRT2.3* and *NRT2.6* were slightly downregulated, and

NRT2.2 was slightly upregulated. To examine how SA187 differentially modulated HATs

expression in Col-0 and *ein2-1*, qRT-PCR analyses were compared following SA187 inoculation

of both genotypes. In shoots, *NRT2.2*, *NRT2.3*, *NRT2.4*, and *NRT2.5* were upregulated by SA187

in Col-0. However, in *ein2-1*, SA187-mediated upregulation was only observed for *NRT2.3* in

both organs and for *NRT2.4* in roots, suggesting a partial disruption of the regulatory response.

Notably, *NRT2.6* was downregulated by SA187 in Col-0 shoots but showed no significant

regulation in *ein2-1*. In roots, SA187 induced the expression of all tested HATs in Col-0. In

contrast, SA187 failed to upregulate *NRT2.1*, *NRT2.2*, and *NRT2.5* in *ein2-1* roots, despite a

normal induction of the other HATs. Amongst the examined HATs, *NRT2.5* exhibited the highest

fold-change in response to SA187 in both organs when compared to mock controls, highlighting

it as a potential major component in SA187-mediated growth enhancement under low nitrate

conditions. Taken together, these results revealed that ethylene signaling was not essential for the

basal expression of most HATs but it was critical for their SA187-induced activation. A major

part of the SA187-induced modulation of HAT expression was observed predominantly in roots,

although some notable effects were also detected in shoots. The *ein2-1* mutation specifically

disrupted SA187-mediated induction of *NRT2.2* and *NRT2.5* in both shoots and roots, *NRT2.4*

and *NRT2.6* in shoots, and *NRT2.1* in roots. These findings support a model in which SA187

leverages ethylene signaling to fine-tune the expression of a subset of nitrate transporters that

improve seedling growth and nitrate content under low nitrate conditions.

E.   Importance of EIN2, NRT2.5, and NRT2.6 in SA187-induced nitrogen

accumulation

To determine whether the SA187-induced upregulation of HATs and growth promotion is

associated with changes in N-allocation, total N content (% DW) was measured in shoots and

roots of Col-0, *ein2-1*, and the *nrt2.5×nrt2.6* double mutant under low nitrate conditions (1 mM

NO_3^-) with and without SA187 inoculation (Figure 4c). In wild-type (Col-0) plants, SA187

inoculation significantly increased shoot N-content compared to mock controls. However, this

beneficial effect was abolished in both *ein2-1* and *nrt2.5×nrt2.6* shoots, with no significant

differences detected between inoculated and mock-treated plants (Fig. 4c). In roots, SA187 did

not significantly alter N-content in wild-type and mutant lines, suggesting that SA187 impacted

N-allocation to the shoot under these experimental conditions (Fig. 4c). These results further

confirmed the involvement of ethylene signaling, *NRT2.5*, and *NRT2.6* in SA187-induced

promotion of shoot N content.

IV.   Discussion

A.   SA187 enhances plant growth under low nitrate conditions

This study demonstrates that SA187 significantly promoted Arabidopsis growth under low

nitrate conditions through a coordinated enhancement of seedling growth and root development,

thus highlighting a potential role in limiting N-based fertilizer use. The observed growth-

promoting effect was tightly linked to external nitrate availability. While only a negligible SA187

effect was seen under high nitrate (20 mM), SA187 significantly increased total FW, PRL, and

LRD under low nitrate regimes (between 0.2–2 mM nitrate) (Figure 1). These findings position

SA187 as a context-dependent growth promoting inoculant that becomes functionally more

advantageous under low nitrate conditions, a hallmark of beneficial plant–microbe interactions to

support host fitness under suboptimal environmental conditions. Indeed, previous studies have

shown that SA187 does not promote plant growth under non-stress conditions but significantly

enhances biomass and root system development under abiotic stresses such as salinity, heat,

drought, and high light stress (Andrés-Barrao et al., 2017; Andres-Barrao et al., 2021; de

Zélicourt et al., 2018; Rolli et al., 2022; Shekhawat et al., 2024).

Interestingly, SA187 consistently increased LRD across all tested nitrate concentrations

including 20 mM when shoot biomass and PRL were unaffected. This constitutive increase in

lateral root formation, regardless of nitrate status, suggests that SA187 may activate a basal root

developmental response. This may improve the plant's capacity to explore heterogeneous soil

environments and exploit transient nutrient patches, an adaptive advantage even when nitrate

supply is sufficient (Garnett et al., 2009). Enhancement of root branching has been observed

previously in other PGPB interactions, where changes in root architecture were found to be

driven by microbial modulation of plant hormone pathways, including auxins, cytokinins, and

ethylene (Spaepen et al., 2008; Vacheron et al., 2013; Chen et al., 2024). The SA187 genome

contains *tnaA*, encoding a tryptophanase that converts tryptophan into indole, a known auxin

precursor, as well as an indole efflux pump (*acrEF*) (Andrés-Barrao et al., 2017; Synek et al.,

2021). Indole and its derivatives contribute to the biosynthesis of indole-3-acetic acid (IAA) and

indole-3-butyric acid (IBA), key auxins involved in lateral root initiation and elongation

(Trujillo-Hernandez et al., 2020; Sun et al., 2023). It is therefore possible that SA187 influences

host auxin dynamics either by directly producing auxin-like metabolites or by stimulation of host

endogenous auxin.

The observed root remodeling may be underpinned by an additional layer of mutualism. It has

been proposed that SA187 preferentially enters plant tissues via natural openings at lateral root

emergence sites (Synek et al., 2021). From this perspective, increased LRD not only benefits the

plant by expanding its nutrient absorption capacity but it will also facilitate bacterial entry and

endophytic establishment. Such a dual-purpose modulation, benefiting both microbe and host,

exemplifies the mutualistic sophistication of beneficial endophytes. Collectively, these results

suggest that SA187 promotes root system remodeling and overall growth to enhance nutrient

acquisition under low N, while also potentially optimizing its own endophytic lifestyle within the

plant host.

B. *SA187 impacts C and N content and improves nitrate accumulation under low nitrate conditions*

SA187 inoculation enhanced Arabidopsis nitrate content and elemental analyses revealed an altered internal N-distribution under low nitrate conditions. This was seen especially under low-nitrate where SA187 increased shoot total N-content while either reducing (Fig. 2b) or maintaining (Fig. 4c) it in the root system. This was accompanied by elevated nitrate levels in both organs in SA187 inoculated seedlings compared to wild-type Col-0. This suggested that SA187 not only facilitated root nitrate uptake but also enhanced its translocation to aerial tissues. Therefore, it appeared that SA187 mediated an altered N-allocation that favored N investment in aerial parts compared to roots. This was only observed under low nitrate, when sustaining photosynthetic efficiency and shoot biomass depends heavily on optimal N-partitioning. The observed shift in N-content from roots to shoots could reflect a strategy to redirect resources to maximize growth under suboptimal N-conditions. The role of PGPB in enhancing both nitrate uptake and root-to-shoot transport has been demonstrated previously in multiple plant-microbe systems. For example, *Achromobacter* sp. inoculation significantly improved biomass production and doubled nitrate uptake efficiency in *Brassica napus* (Bertrand et al., 2000). Likewise, volatile organic compounds from *Bacillus velezensis* SQR9 promoted nitrate accumulation and growth in both Arabidopsis and rice (Chen et al., 2024). Another *Enterobacter* species, *E. cloacae* subsp. *dissolvens*, was shown to improve N-content and yield of teff varieties under field conditions, indicating the broader potential of this genus in improving plant N-nutrition (Tsegaye et al., 2022).

Beyond uptake, the effective transport and assimilation of nitrate into amino acids and proteins is

a key determinant of NUE and overall plant growth and productivity. By enhancing both nitrate

uptake and its allocation to shoots, PGPB may help plants optimize N-use under stress

conditions. Upregulated expression of nitrate transporters involved in xylem loading, such as

AtNRT1.5, has been reported in response to beneficial microbes, facilitating enhanced nitrate

flow to shoot tissues (Calvo et al., 2019). Such enhanced translocation may not only support

vegetative growth but also influence seed development and nutrient remobilization during

reproductive stages. Indeed, microbial inoculation has been shown to improve N-remobilization

efficiency in crop plants, benefiting grain protein content and yield (Adesemoye et al., 2008).

Although the current study focused on vegetative development, it is plausible that SA187-

mediated enhancements in nitrate uptake and N-allocation may also support reproductive success

and seed filling, contributing to overall NUE, a key trait for achieving sustainable crop

productivity under N-limited conditions (Congreves et al., 2021).

The C/N ratio in plant tissues is an indicator of metabolic balance and nutritional status.

Efficient N-assimilation requires the incorporation of C-skeletons into amino acids and structural

components, thereby limiting the buildup of unused C-rich metabolites when N is scarce (Elser

et al., 2010; Baslam et al., 2020; Li et al., 2022). In line with this, C:N ratios were elevated under

low nitrate availability. However, SA187 inoculation significantly reduced C:N ratios under

these conditions, suggesting improved N acquisition and assimilation. Notably, SA187-

inoculated plants under low nitrate exhibited shoot C:N ratios comparable to those observed in

plants grown under a high nitrate supply. This suggests that SA187 enhanced nitrate uptake

sufficiently to restore metabolic balance even in low-nitrate environments. From an agronomic

perspective, plants with low C:N ratios often have high protein contents, improved nutritional

quality, and produce residues that decompose more readily, enhancing soil fertility and nutrient

recycling (Hertzler et al., 2020; De Silva et al., 2023).

Collectively, these results point to a dual contribution of SA187 in facilitating nitrate uptake and

optimizing its internal use and allocation. Such a capacity highlights the potential of SA187 as a

bioinoculant for improving plant N-status, especially in the context of reducing the use of

synthetic fertilizers while maintaining the productivity of low-input agricultural systems.

C. Transcriptomic analysis reveals SA187-induced reprogramming of nitrogen and

hormone signaling pathways under low nitrogen

Transcriptomic profiling revealed that SA187 induced a coordinated and condition-specific

reprogramming of *Arabidopsis thaliana* gene expression (Figure 3). This reprogramming

encompassed multiple layers of regulatory networks including nutrient acquisition, hormone-

mediated development, immune modulation, and cellular stress responses. The observed

expression patterns reflected a potential dynamic microbial modulation of host physiology to

improve growth and resilience in low nitrate environments.

The notable induction of defense- and immune-related genes by SA187, especially under

sufficient nitrate conditions, suggested an underlying mechanism of induced systemic resistance

(ISR). It is a common plant defense strategy triggered by beneficial microbes, allowing plants to

rapidly and robustly respond to eventual biotic stress conditions (Van der Ent et al., 2009;

Pieterse et al., 2014). This priming effect by SA187 aligns closely with previous reports

demonstrating ISR induction by beneficial rhizobacteria (Zamioudis et al., 2013; de Zélicourt et

al., 2018). The largest SA187-induced gene cluster in shoots (Cluster 4) was enriched for genes

associated with defense signaling, oxidative stress responses, and hormone pathways including

ethylene, SA and JA. These genes were strongly induced by SA187 under high nitrate conditions

and to a lesser extent under low nitrate, suggesting a nitrate-independent component of defense

priming. Such transcriptional activation is consistent with microbe-ISR, a process where non-

pathogenic rhizobacteria enhance plant immunity (Zehnder et al., 2001; Choudhary et al., 2007;

Van der Ent et al., 2009; Pieterse et al., 2014). The concurrent induction of SA and JA signaling

components points to a nuanced immune regulation, potentially enabling both basal defense and

biotic stress tolerance. Studies have shown that an increased N-supply can lead to plant

susceptibility to pathogens (Mur et al., 2016; Jeon, 2019; Maina et al., 2024). Interestingly,

although SA187 did not enhance biomass under high nitrate, its activation of immune pathways

suggested it could confer fitness advantages by strengthening plant immunity. Consistent with

this idea, shoot Clusters 3 and 9 displayed SA187-induced expression irrespective of N-status.

These clusters were enriched in genes responding to microbial symbionts, hormone signaling

(SA, JA, auxin, ethylene), N-compounds, and root developmental regulators such as lateral root

initiation and camalexin biosynthesis. The nitrate-independent induction of these clusters

suggested a basal SA187-triggered transcriptional program integrating developmental cues with

immune modulation, likely enabling dual growth-promotion and stress-buffering effects. The co-

enrichment of hypoxia-responsive genes could be linked to bacterial modulation of the

rhizosphere redox environment, potentially altering local oxygen tension or reactive oxygen

species homeostasis, both known to influence root development and immune status (Bailey-

Serres and Voesenek, 2008; Considine and Foyer, 2014).

On the other hand, shoot Cluster 10 displayed a nitrate-sensitive signature where genes were

upregulated under low nitrate but repressed upon SA187 inoculation. This cluster included genes

related to detoxification, glutathione metabolism enzymes, and flavonoid biosynthetic, classical

metabolic markers of cellular stress responses to N-deprivation and oxidative imbalance (Lea et

al., 2007; Krapp et al., 2011; Curci et al., 2017; Liang and He, 2018; Li et al., 2023; Yang et al.,

2024). SA187-mediated suppression of these stress markers under low nitrate again suggested a

buffering effect, whereby SA187 reduced the perceived severity or molecular consequences of

low nitrate.

Root transcriptome profiling revealed a comparable complexity. The largest cluster (Cluster 3)

was strongly induced by SA187 under both nitrate regimes and contained genes involved in

immune signaling, phytohormone pathways (ethylene, SA, JA, auxin, cytokinin), and oxidative

stress responses. SA187 is known to colonize the root endosphere, therefore this widespread

induction probably reflected direct microbe-host signaling at the root interface. Cluster 5 was

characterized by a classical stress signature under low nitrate that was downregulated by SA187.

This included genes related to amino acid metabolism, redox balance, and hypoxia, consistent

with earlier findings that SA187 alleviates cellular stress via metabolic reprogramming (de

Zélicourt et al., 2018; Andres-Barrao et al., 2021; Shekhawat et al., 2024). A strikingly distinct

pattern was observed in Cluster 11, where genes were upregulated by SA187 only under low

nitrate. Functional enrichment revealed root morphogenesis, root hair elongation, nitrate import,

and cellular responses to reactive N species. The exclusive upregulation of this cluster under

SA187–low nitrate conditions suggested a tightly controlled transcriptional module integrating

nutrient signals with bacterial cues to selectively activate root developmental pathways when

most needed.

Among the most functionally relevant findings was the upregulation of high-affinity nitrate

transporter genes (*NRT2.5* in shoots, *NRT2.4* and *NRT2.6* in roots). These transporters are critical

under low external nitrate concentrations (Lezhneva et al., 2014; Krapp et al., 2014). Their

induction under low nitrate conditions suggested that SA187 benefits observed in terms of

growth and nitrate acquisition and partitioning concerned NRT2.4, NRT2.5 and NRT2.6.

Collectively, the transcriptomic data supported a model in which SA187 dynamically rewired

plant gene expression to coordinate developmental and metabolic adaptations, enhance nitrate

uptake and allocation, and buffer physiological stress responses. These multifaceted changes may

enable SA187-inoculated plants to sustain growth and development under low nitrate conditions

and position this endophyte as a potent microbial tool for improving NUE and resilience in

crops.

D. SA187-mediated growth promotion requires coordinated action of ethylene

signaling and high-affinity nitrate transporters

Functional analyses suggested that both ethylene signaling and specific high-affinity nitrate

transporters, particularly *NRT2.5* and *NRT2.6*, were required for SA187-mediated growth

promotion under low nitrate conditions. Phenotypic analyses confirmed that the *ein2-1*, *nrt2.5*,

nrt2.6, *nrt2.5×nrt2.6* mutants failed to show significant growth improvements following SA187

inoculation, unlike wild-type plants. Indeed, previous reports have demonstrated the crucial

regulatory role of ethylene in modulating nitrate transporter expression and root system

architecture under varying N-conditions. Ethylene signaling, mediated through *EIN2/EIN3*

pathways, has been previously shown to regulate critical nitrate transporter genes such as

NRT1.1 and *NRT2.1*, fine-tuning nitrate uptake and root developmental responses to external N-

availability (Tian et al., 2009; Zheng et al., 2013). For instance, ethylene-mediated repression of

lateral root growth under high nitrate conditions required intact *EIN2-1* signaling, and ethylene-

insensitive mutants (*ein2-1*, *etr1-3*) exhibited dysregulated root responses to nitrate signals (Tian

et al., 2009). Additionally, Zheng et al. (2013) identified a feedback regulatory loop whereby low

nitrate availability induced ethylene biosynthesis, subsequently repressing *NRT2.1* expression to

avoid excessive nitrate uptake under prolonged deficiency. Under abiotic stresses, ethylene also

modulated stress-induced nitrate allocation to roots via coordinated regulation of *NRT1.5* (xylem

loading) and *NRT1.8* (xylem unloading), further exemplifying its central role in plant nutrient

homeostasis and stress adaptation (Zhang et al., 2014).

The requirement for ethylene signaling in this context was further supported by SA187-induced

expression of *NRT2.1–NRT2.6* in wild-type roots, and selective induction of *NRT2.2*, *NRT2.3*,

NRT2.4, and *NRT2.5* in shoots. In contrast, *NRT2.6* was downregulated in shoots following

SA187 treatment. Among all transporters, *NRT2.5* displayed the strongest induction, suggesting a

central role in SA187-mediated enhancement of nitrate content and N-allocation. This

transcriptional activation was largely abolished in *ein2-1*, which failed to upregulate key

transporters including *NRT2.1* in roots, and *NRT2.2* and *NRT2.5* in both shoots and roots. These

results clearly position these genes downstream of ethylene signaling in the SA187 response.

Interestingly, SA187-induced *NRT2.6* downregulation in shoots was not observed in *ein2-1*,

while root expression of *NRT2.6* remained upregulated in both genotypes, suggesting a partially

ethylene-dependent regulatory pattern that was dependent in shoots but independent in roots.

The significance of *NRT2.5* in this context is consistent with its critical role under prolonged N-

starvation. Indeed, *NRT2.5* was found to be essential not only for efficient nitrate uptake but also

for phloem loading and N-remobilization, supporting plant growth under sustained N-limitation

(Lezhneva et al., 2014). Similarly, *NRT2.4*, which was also significantly upregulated by SA187

in both organs, has been shown to contribute to nitrate uptake and remobilization, particularly in

lateral roots under severe N-deficiency (Kiba et al., 2012). While a precise role for *NRT2.6* in

nitrate transport remains poorly defined, earlier studies suggested its involvement in pathogen-

induced oxidative stress responses rather than direct nitrate acquisition (Dechorgnat et al., 2012).

However, the requirement of *NRT2.6* for SA187-mediated growth enhancement aligns with

observations from beneficial interactions involving other bacteria such as *Phyllobacterium*

brassicacearum STM196, where mutants of *NRT2.6* (and *NRT2.5*) lose beneficial bacteria

growth effects (Kechid et al., 2013).

The existence of both ethylene-dependent and ethylene-independent regulations of nitrate

transporter expression by SA187 inoculation was observed when comparing wild-type and *ein2-*

1 seedlings. While *NRT2.2*, *NRT2.5*, *NRT2.6* (shoots) and *NRT2.1* (roots) clearly required

ethylene signaling for SA187-induced expression, the regulation of *NRT2.3* in both organs and

NRT2.4 and *NRT2.6* in roots appeared to be independent of ethylene. This dual regulatory system

may provide robustness to nitrate acquisition and translocation strategies under fluctuating

environmental conditions where ethylene signaling may be perturbed, for example, under

combined stress scenarios. Such regulatory flexibility shows the complexity of microbial

modulation of plant nutrient responses and highlights the sophisticated molecular interplay

microbial signals, hormone-mediated transcriptional networks, and nitrate transporter systems.

E.   Ethylene signaling and nitrate transporters coordinate SA187-induced nitrogen

allocation to shoots

The loss of SA187-mediated growth promotion in *ein2-1* and *nrt2.5×nrt2.6* mutants, combined

with the altered expression of HATs in ethylene-insensitive *ein2-1*, led to the hypothesis that both

ethylene signaling and functional NRT2.5 and NRT2.6 were critical for SA187-driven N-

allocation to shoots. In wild-type plants, SA187 significantly increased shoot N-content,

consistent with enhanced nitrate uptake and preferential allocation to aerial tissues. The complete

loss of this effect in *ein2-1* and *nrt2.5×nrt2.6* confirmed that both ethylene signaling and (at least

one of) the HATs were required for this shift in N-allocation induced by SA187. The loss of this

response in *ein2-1* reinforced the idea of ethylene as a central regulator in SA187-induced nitrate

transport and N-allocation. Transcriptomic and qRT-PCR data consistently showed that *NRT2.5*

expression was strongly induced by SA187 in wild-type plants but not in *ein2-1*, demonstrating

that this transporter functioned downstream of ethylene signaling. These findings align with

earlier studies indicating that ethylene can regulate nitrate transporter expression under N-

limitation (Tian et al., 2009; Zheng et al., 2013), and support the view that ethylene functions as

a key signaling hub linking microbial perception with nitrate uptake pathways. Similarly, the

failure of SA187 to enhance shoot N-content in the *nrt2.5×nrt2.6* double mutant confirmed the

essential contribution of these transporters to nitrate acquisition and distribution. SA187-

dependent regulation of *NRT2.5* and *NRT2.6* under low nitrate, as observed in wild-type plants,

indicates their roles in mediating improved N-allocation to shoots. Given that shoot N-

accumulation correlated with growth improvement, this transporter-dependent redistribution is

likely a prerequisite for the observed biomass gains.

These findings collectively support a model in which SA187 promotes nitrate uptake and N-

allocation to the shoot through coordinated regulation of HATs expression via both ethylene-

dependent and ethylene-independent pathways. *NRT2.5* functions downstream of ethylene

signaling in both shoots and roots, while *NRT2.6* shows a dual regulatory pattern; ethylene-

dependent in the shoot, but ethylene-independent in the root (Figure 5). This coordinated spatial

and hormonal regulatory control allows SA187-mediated fine-tuning of nitrate uptake and N-

allocation through multiple entry points in the signaling network, enabling improved plant N-

status and internal redistribution under low nitrate conditions. These findings advance our

understanding of plant–microbe interactions and offer a framework for leveraging beneficial

microbes to enhance NUE and promote sustainable low N-based fertilizer agricultural practices.

Acknowledgments and funding statement

Amina Ilyas was supported by a PhD scholarship from the French Ministry of Higher Education

and Research (Ministère de l'Enseignement supérieur et de la Recherche). This work was

supported by funding from the Indo-French Centre for the Promotion of Advanced Research

(IFCPAR/CEFIPRA) under project 68T06-1. Transcriptomics analyses were performed in

collaboration with the POPS platform (IPS2, University Paris-Saclay) which is supported by

Saclay Plant Sciences (ANR-17-EUR-0007). Seeds of the *nrt2.5*, *nrt2.6*, and *nrt2.5×nrt2.6*

mutant lines were kindly provided by Dr. Guilhem Debrosses (Université de Montpellier). The

authors would like to thank all colleagues and collaborators who contributed to discussions and

technical support throughout the course of this study.

Author Approvals

591 All authors have seen and approved the final version of the manuscript. The work described has

592 not been published previously and is not accepted for publication elsewhere.

Competing Interests

594 The authors declare no competing interests.

References

Adesemoye, A. O., Torbert, H. A., Kloepper, J. W., Adesemoye, A., Torbert, H., and Kloepper, J. (2008). Enhanced Plant Nutrient Use Efficiency with PGPR and AMF in an Integrated Nutrient Management System **562**.

Ahmad Wani, B., Rashid, S., Rashid, K., Javid, H., Ahmad Magray, J., Qadir, R. ul, and Islam, T. (2022). Nitrogen Deficiency in Plants. In *Advances in Plant Nitrogen Metabolism*, pp. 28–37. CRC Press.

Ali, A., Jabeen, N., Farruhbek, R., Chachar, Z., Laghari, A. A., Chachar, S., Ahmed, N., Ahmed, S., and Yang, Z. (2025). Enhancing nitrogen use efficiency in agriculture by integrating agronomic practices and genetic advances. *Front Plant Sci* **16**:1543714.

Anders, S., Pyl, P. T., and Huber, W. (2015). HTSeq—a Python framework to work with high-throughput sequencing data. *Bioinformatics* **31**:166–169.

Andrés-Barrao, C., Lafi, F. F., Alam, I., de Zélicourt, A., Eida, A. A., Bokhari, A., Alzubaidy, H., Bajic, V. B., Hirt, H., and Saad, M. M. (2017). Complete genome sequence analysis of Enterobacter sp. SA187, a plant multi-stress tolerance promoting endophytic bacterium. *Front Microbiol* **8**:285704.

Andres-Barrao, C., Alzubaidy, H., Jalal, R., Mariappan, K. G., De Zelicourt, A., Bokhari, A., Artyukh, O., Alwutayd, K., Rawat, A., Shekhawat, K., et al. (2021). Coordinated bacterial and plant sulfur metabolism in Enterobacter sp. SA187–induced plant salt stress tolerance. *pnas.org* **118**.

Bailey-Serres, J., and Voesenek, L. A. C. J. (2008). Flooding stress: Acclimations and genetic diversity. *Annu Rev Plant Biol* **59**:313–339.

Bakhshandeh, E., Pirdashti, H., Shahsavarpour Lendeh, K., Gilani, Z., Yaghoubi

Khanghahi, M., and Crecchio, C. (2020). Effects of plant growth promoting

microorganisms inoculums on mineral nutrition, growth and productivity of rice (Oryza

sativa L.). *J Plant Nutr* **43**:1643–1660.

Baslam, M., Mitsui, T., Sueyoshi, K., and Ohyama, T. (2020). Recent Advances in Carbon and

Nitrogen Metabolism in C3 Plants. *International Journal of Molecular Sciences 2021, Vol.*

22, Page 318 **22**:318.

Baudry, K., Paysant-Le Roux, C., Colella, S., Castandet, B., and Martin, M.-L. (2022).

Analyzing Multifactorial RNA-Seq Experiments with DiCoExpress. *Journal of Visualized*

Experiments Advance Access published July 29, 2022, doi:10.3791/62566.

Bertrand, H., Plassard, C., Pinochet, X., Touraine, B., Normand, P., and Cleyet-Marel, J. C.

(2000). Stimulation of the ionic transport system in Brassica napus by a plant growth-

promoting rhizobacterium (Achromobacter sp.). *Can. J. Microbiol* **46**:229–236.

Bushnell, B. (2014). BBMap: A fast, accurate, splice-aware aligner. Retrieved from

https://sourceforge.net/projects/bbmap/.

Calvo, P., Zebelo, S., McNear, D., Kloepper, J., and Fadamiro, H. (2019). Plant growth-

promoting rhizobacteria induce changes in Arabidopsis thaliana gene expression of nitrate

and ammonium uptake genes. *J Plant Interact* **14**:224–231.

Chen, Y., Li, Y., Fu, Y., Jia, L., Li, L., Xu, Z., Zhang, N., Liu, Y., Fan, X., Xuan, W., et al.

(2024). The beneficial rhizobacterium Bacillus velezensis SQR9 regulates plant nitrogen

uptake via an endogenous signaling pathway. *J Exp Bot* **75**:3388–3400.

Cherkasov, N., Ibhadon, A. O., and Fitzpatrick, P. (2015). A review of the existing and alternative methods for greener nitrogen fixation. *Chemical Engineering and Processing: Process Intensification* **90**:24–33.

Choudhary, D. K., Prakash, A., and Johri, B. N. (2007). Induced systemic resistance (ISR) in plants: Mechanism of action. *Indian J Microbiol* **47**:289–297.

Congreves, K. A., Otchere, O., Ferland, D., Farzadfar, S., Williams, S., and Arcand, M. M. (2021). Nitrogen Use Efficiency Definitions of Today and Tomorrow. *Front Plant Sci* **12**:637108.

Considine, M. J., and Foyer, C. H. (2014). Redox Regulation of Plant Development. *Antioxid Redox Signal* **21**:1305.

Curci, P. L., Aiese Cigliano, R., Zuluaga, D. L., Janni, M., Sanseverino, W., and Sonnante, G. (2017). Transcriptomic response of durum wheat to nitrogen starvation. *Sci Rep* **7**:1–14.

De Silva, C., Rathor, P., Poudel, H. P., and Thilakarathna, M. S. (2023). Effects of Drought Stress on Red Clover-Grass Mixed Stands Compared to Grass Monoculture Stands in Nitrogen-Deficient Systems. *Nitrogen 2023, Vol. 4, Pages 382-396* **4**:382–396.

de Vries, W., Posch, M., Simpson, D., de Leeuw, F. A. A. M., van Grinsven, H. J. M., Schulte-Uebbing, L. F., Sutton, M. A., and Ros, G. H. (2024). Trends and geographic variation in adverse impacts of nitrogen use in Europe on human health, climate, and ecosystems: A review. *Earth Sci Rev* **253**:104789.

De Zelicourt, A., Al-Yousif, M., and Hirt, H. (2013). Rhizosphere microbes as essential partners for plant stress tolerance. *Mol Plant* **6**:242–245.

de Zélicourt, A., Synek, L., Saad, M. M., Alzubaidy, H., Jalal, R., Xie, Y., Andrés-Barrao, C., Rolli, E., Guerard, F., Mariappan, K. G., et al. (2018). Ethylene induced plant stress

tolerance by Enterobacter sp. SA187 is mediated by 2□keto□4□methylthiobutyric acid production. *journals.plos.org* **14**.

Dechorgnat, J., Patrit, O., Krapp, A., Fagard, M., and Daniel-Vedele, F. (2012). Characterization of the Nrt2.6 Gene in Arabidopsis thaliana: A Link with Plant Response to Biotic and Abiotic Stress. *PLoS One* **7**:e42491.

Di Benedetto, N. A., Corbo, M. R., Campaniello, D., Cataldi, M. P., Bevilacqua, A., Sinigaglia, M., and Flagella, Z. (2017). The role of Plant Growth Promoting Bacteria in improving nitrogen use efficiency for sustainable crop production: a focus on wheat. *AIMS Microbiol* **3**:413.

Ding, L., Wang, K. J., Jiang, G. M., Biswas, D. K., Xu, H., Li, L. F., and Li, Y. H. (2005). Effects of Nitrogen Deficiency on Photosynthetic Traits of Maize Hybrids Released in Different Years. *Ann Bot* **96**:925–930.

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., and Gingeras, T. R. (2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**:15–21.

Elser, J. J., Fagan, W. F., Kerkhoff, A. J., Swenson, N. G., and Enquist, B. J. (2010). Biological stoichiometry of plant production: Metabolism, scaling and ecological response to global change. *New Phytologist* **186**:593–608.

Etesami, H., and Adl, S. M. (2020). Plant Growth-Promoting Rhizobacteria (PGPR) and Their Action Mechanisms in Availability of Nutrients to Plants Advance Access published 2020, doi:10.1007/978-981-15-2576-6_9.

Fan, X., Naz, M., Fan, X., Xuan, W., Miller, A. J., and Xu, G. (2017). Plant nitrate transporters: from gene function to application. *J Exp Bot* **68**:2463–2475.

FAO, and UNEP (2021). *Environmental, health and socio-economic impacts of soil pollution.*

(ed. FAO and UNEP) Rome: FAO and UNEP.

Garnett, T., Conn, V., and Kaiser, B. N. (2009). Root based approaches to improving nitrogen

use efficiency in plants. *Plant Cell Environ* **32**:1272–1283.

Glass, A. D. M. (2009). Nitrate uptake by plant roots. *Botany* **87**:659–667.

Glick, B. R. (2020). Introduction to Plant Growth-Promoting Bacteria. *Beneficial Plant-*

Bacterial Interactions Advance Access published 2020, doi:10.1007/978-3-030-44368-9_1.

Govindasamy, P., Muthusamy, S. K., Bagavathiannan, M., Mowrer, J., Jagannadham, P. T.

K., Maity, A., Halli, H. M., G. K, S., Vadivel, R., T. K, D., et al. (2023). Nitrogen use

efficiency—a key to enhance crop productivity under a changing climate. *Front Plant Sci*

14:1121073.

Guzmán, P., and Ecker, J. R. (1990). Exploiting the triple response of Arabidopsis to identify

ethylene-related mutants. *Plant Cell* **2**:513–523.

Hawkesford, M., Horst, W., Kichey, T., Lambers, H., Schjoerring, J., Møller, I. S., and

White, P. (2012). Functions of Macronutrients. In *Marschner's Mineral Nutrition of Higher*

Plants, pp. 135–189. Elsevier.

Hertzler, S. R., Lieblein-Boff, J. C., Weiler, M., and Allgeier, C. (2020). Plant Proteins:

Assessing Their Nutritional Quality and Effects on Health and Physical Function. *Nutrients*

12:3704.

Iqbal, N., Khan, N. A., Ferrante, A., Trivellini, A., Francini, A., and Khan, M. I. R. (2017).

Ethylene role in plant growth, development and senescence: interaction with other

phytohormones. *Front Plant Sci* **8**:235913.

Jeon, J. (2019). Phytobiome as a Potential Factor in Nitrogen-Induced Susceptibility to the Rice

Blast Disease. *Plant Dis* **25**:103–107.

Kechid, M., Desbrosses, G., Rokhsi, W., Varoquaux, F., Djekoun, A., and Touraine, B.

(2013). The NRT2.5 and NRT2.6 genes are involved in growth promotion of Arabidopsis by

the plant growth-promoting rhizobacterium (PGPR) strain Phyllobacterium brassicacearum

STM196. *New Phytologist* **198**:514–524.

Khan, N. A., Khan, M. I. R., Ferrante, A., and Poor, P. (2017). Editorial: Ethylene: A Key

Regulatory Molecule in Plants ROLE OF ETHYLENE IN DEVELOPMENTAL PROCESS

Flower Development and Fruit Ripening. *Frontiers in Plant Science | www.frontiersin.org*

8:1782.

Kiba, T., Feria-Bourrellier, A. B., Lafouge, F., Lezhneva, L., Boutet-Mercey, S., Orsel, M.,

Bréhaut, V., Miller, A., Daniel-Vedele, F., Sakakibara, H., et al. (2012). The Arabidopsis

Nitrate Transporter NRT2.4 Plays a Double Role in Roots and Shoots of Nitrogen-Starved

Plants. *Plant Cell* **24**:245–258.

Krapp, A. (2015). Plant nitrogen assimilation and its regulation: A complex puzzle with missing

pieces. *Curr Opin Plant Biol* **25**:115–122.

Krapp, A., Berthomé, R., Orsel, M., Mercey-Boutet, S., Yu, A., Castaings, L., Elftieh, S.,

Major, H., Renou, J. P., and Daniel-Vedele, F. (2011). Arabidopsis Roots and Shoots

Show Distinct Temporal Adaptation Patterns toward Nitrogen Starvation. *Plant Physiol*

157:1255.

Krapp, A., David, L. C., Chardin, C., Girin, T., Marmagne, A., Leprince, A. S., Chaillou, S.,

Ferrario-Méry, S., Meyer, C., and Daniel-Vedele, F. (2014). Nitrate transport and

signalling in Arabidopsis. *J Exp Bot* **65**:789–798.

Lambert, I., Paysant-Le Roux, C., Colella, S., and Martin-Magniette, M.-L. (2020).

DiCoExpress: a tool to process multifactorial RNAseq experiments from quality controls to

co-expression analysis through differential analysis based on contrasts inside GLM models.

Plant Methods **16**:68.

Lea, U. S., Slimestad, R., Smedvig, P., and Lillo, C. (2007). Nitrogen deficiency enhances

expression of specific MYB and bHLH transcription factors and accumulation of end

products in the flavonoid pathway. *Planta* **225**:1245–1253.

LeBauer, D. S., and Treseder, K. K. (2008). Nitrogen limitation of net primary productivity in

terrestrial ecosystems is globally distributed. *Wiley Online LibraryDS LeBauer, KK*

TresederEcology, 2008•Wiley Online Library **89**:371–379.

Leblanc, A., Renault, H., Lecourt, J., Etienne, P., Deleu, C., and Le Deunff, E. (2008).

Elongation Changes of Exploratory and Root Hair Systems Induced by Aminocyclopropane

Carboxylic Acid and Aminoethoxyvinylglycine Affect Nitrate Uptake and BnNrt2.1 and

BnNrt1.1 Transporter Gene Expression in Oilseed Rape. *Plant Physiol* **146**:1928–1940.

Lezhneva, L., Kiba, T., Feria-Bourrellier, A. B., Lafouge, F., Boutet-Mercey, S., Zoufan, P.,

Sakakibara, H., Daniel-Vedele, F., and Krapp, A. (2014). The Arabidopsis nitrate

transporter NRT2.5 plays a role in nitrate acquisition and remobilization in nitrogen-starved

plants. *Plant Journal* **80**:230–241.

Li, Q., Ren, Y., Fu, H., Li, Z., Kong, F., and Yuan, J. (2022). Cultivar differences in carbon

and nitrogen accumulation, balance, and grain yield in maize. *Front Plant Sci* **13**:992041.

Li, J., Liu, X., Xu, L., Li, W., Yao, Q., Yin, X., Wang, Q., Tan, W., Xing, W., and Liu, D.

(2023). Low nitrogen stress-induced transcriptome changes revealed the molecular response

752 and tolerance characteristics in maintaining the C/N balance of sugar beet (Beta vulgaris

753 L.). *Front Plant Sci* **14**:1164151.

754 **Liang, J., and He, J.** (2018). Protective role of anthocyanins in plants under low nitrogen stress.

755 *Biochem Biophys Res Commun* **498**:946–953.

756 **Lin, L., Li, Z., Hu, C., Zhang, X., Chang, S., Yang, L., Li, Y., and An, Q.** (2012). Plant

757 Growth-Promoting Nitrogen-Fixing Enterobacteria Are in Association with Sugarcane

758 Plants Growing in Guangxi, China. *Microbes Environ* **27**:391–398.

759 **Liu, J., You, L., Amini, M., Obersteiner, M., Herrero, M., Zehnder, A. J. B., and Yang, H.**

760 (2010). A high-resolution assessment on global nitrogen flows in cropland. *Proc Natl Acad*

761 *Sci U S A* **107**:8035.

762 **Livak, K. J., and Schmittgen, T. D.** (2001). Analysis of Relative Gene Expression Data Using

763 Real-Time Quantitative PCR and the $2-\Delta\Delta CT$ Method. *Methods* **25**:402–408.

764 **Madhaiyan, M., Peng, N., Te, N. S., Hsin, C., Lin, C., Lin, F., Reddy, C., Yan, H., and Ji, L.**

765 (2013). Improvement of plant growth and seed yield in Jatropha curcas by a novel nitrogen-

766 fixing root associated Enterobacter species. *Biotechnol Biofuels* **6**:1.

767 **Maina, A. W., Becker, M., and Oerke, E. C.** (2024). Assessing Interactions between Nitrogen

768 Supply and Leaf Blast in Rice by Hyperspectral Imaging. *Remote Sens (Basel)* **16**:939.

769 **McCarthy, D. J., Chen, Y., and Smyth, G. K.** (2012). Differential expression analysis of

770 multifactor RNA-Seq experiments with respect to biological variation. *Nucleic Acids Res*

771 **40**:4288–4297.

772 **Mur, L. A. J., Simpson, C., Kumari, A., Gupta, A. K., and Gupta, K. J.** (2016). Moving

773 nitrogen to the centre of plant defence against pathogens. *Ann Bot* **119**:703.

Nath, A., Bhuyan, P., Gogoi, N., and Deka, P. (2023). Pesticides and Chemical Fertilizers: Role

in Soil Degradation, Groundwater Contamination, and Human Health. *Xenobiotics in Urban*

Ecosystems: Sources, Distribution and Health Impacts Advance Access published January

1, 2023, doi:10.1007/978-3-031-35775-6_7.

O'Brien, J. A. A., Vega, A., Bouguyon, E., Krouk, G., Gojon, A., Coruzzi, G., and Gutiérrez,

R. A. A. (2016). Nitrate Transport, Sensing, and Responses in Plants. *Mol Plant* **9**:837–856.

Pathania, P., Rajta, A., Singh, P. C., and Bhatia, R. (2020). Role of plant growth-promoting

bacteria in sustainable agriculture. *Biocatal Agric Biotechnol* **30**:101842.

Pieterse, C. M. J., Zamioudis, C., Berendsen, R. L., Weller, D. M., Van Wees, S. C. M., and

Bakker, P. A. H. M. (2014). Induced systemic resistance by beneficial microbes. *Annu Rev*

Phytopathol **52**:347–375.

Rahman, S., Rahmani, A. A., Abdul Aziz, M., Ahmad, M., Sivasankaran, S. K., Hirt, H.,

and Masmoudi, K. (2025). The Enterobacter sp. SA187 stimulates stress-responsive genes

and promotes salt and heat stress tolerance in tomato plants. *Sci Hortic* **342**:114038.

Rigaill, G., Balzergue, S., Brunaud, V., Blondet, E., Rau, A., Rogier, O., Caius, J., Maugis-

Rabusseau, C., Soubigou-Taconnat, L., Aubourg, S., et al. (2016). Synthetic data sets for

the identification of key ingredients for RNA-seq differential analysis. *Brief Bioinform*

Advance Access published October 14, 2016, doi:10.1093/bib/bbw092.

Robinson, M. D., McCarthy, D. J., and Smyth, G. K. (2010). <tt>edgeR</tt>□: a

Bioconductor package for differential expression analysis of digital gene expression data.

Bioinformatics **26**:139–140.

Rolli, E., de Zélicourt, A., Alzubaidy, H., Karampelias, M., Parween, S., Rayapuram, N.,

Han, B., Froehlich, K., Abulfaraj, A. A., Alhoraibi, H., et al. (2022). The Lys□motif

receptor LYK4 mediates Enterobacter sp. SA187 triggered salt tolerance in Arabidopsis

thaliana. *Wiley Online Library* **24**:223–239.

Rouwenhorst, K. H. R., and Lefferts, L. (2020). Feasibility study of plasma-catalytic ammonia

synthesis for energy storage applications. *Catalysts* **10**:1–19.

Saad, M., de Zelicourt, A., Rolli, E., Synek, L., and Hirt, H. (2018). Quantification of Root

Colonizing Bacteria. *Bio Protoc* **8**.

Shankar, M., Ponraj, P., Ilakkiam, D., and Gunasekaran, P. (2011). Root colonization of a

rice growth promoting strain of Enterobacter cloacae. *J Basic Microbiol* **51**:523–530.

Shekhawat, K., Sheikh, A., Mariappan, K., Jalal, R., bioRxiv, H. H.-, and 2020, undefined

(2021a). Enterobacter sp. SA187 mediates plant thermotolerance by chromatin modification

of heat stress genes. *biorxiv.org* Advance Access published 2021,

doi:10.1101/2020.01.16.908756.ABSTRACT.

Shekhawat, K., Saad, M. M., Sheikh, A., Mariappan, K., Al□Mahmoudi, H., Abdulhakim,

F., Eida, A. A., Jalal, R., Masmoudi, K., and Hirt, H. (2021b). Root endophyte induced

plant thermotolerance by constitutive chromatin modification at heat stress memory gene

loci. *EMBO Rep* **22**.

Shekhawat, K., Veluchamy, A., Fatima, A., García-Ramírez, G. X., Reichheld, J. P.,

Artyukh, O., Fröhlich, K., Polussa, A., Parween, S., Nagarajan, A. P., et al. (2024).

Microbe-induced coordination of plant iron–sulfur metabolism enhances high-light-stress

tolerance of Arabidopsis. *Plant Commun* **5**:101012.

Sinclair, T. R., and Rufty, T. W. (2012). Nitrogen and water resources commonly limit crop

yield increases, not necessarily plant genetics. *Glob Food Sec* **1**:94–98.

Smith, T., Heger, A., and Sudbery, I. (2017). UMI-tools: modeling sequencing errors in Unique

Molecular Identifiers to improve quantification accuracy. *Genome Res* **27**:491–499.

Sun, P., Huang, Y., Yang, X., Liao, A., and Wu, J. (2023). The role of indole derivative in the

growth of plants: A review. *Front Plant Sci* **13**:1120613.

Sutton, M. A., Howard, C. M., Kanter, D. R., Lassaletta, L., Móring, A., Raghuram, N., and

Read, N. (2021). The nitrogen decade: mobilizing global action on nitrogen to 2030 and

beyond. *One Earth* **4**:10–14.

Synek, L., Rawat, A., L'Haridon, F., Weisskopf, L., Saad, M. M., and Hirt, H. (2021).

Multiple strategies of plant colonization by beneficial endophytic Enterobacter sp. SA187.

Environ Microbiol **23**:6223–6240.

Tian, Q. Y., Sun, P., and Zhang, W. H. (2009). Ethylene is involved in nitrate-dependent root

growth and branching in Arabidopsis thaliana. *New Phytologist* **184**:918–931.

Trujillo-Hernandez, J. A., Bariat, L., Enders, T. A., Strader, L. C., Reichheld, J. P., and

Belin, C. (2020). A glutathione-dependent control of the indole butyric acid pathway

supports Arabidopsis root system adaptation to phosphate deprivation. *J Exp Bot* **71**:4843–

4857.

Tsegaye, Z., Alemu, T., Desta, F. A., and Assefa, F. (2022). Plant growth-promoting

rhizobacterial inoculation to improve growth, yield, and grain nutrient uptake of teff

varieties. *Front Microbiol* **13**:896770.

Van der Ent, S., Van Wees, S. C. M., and Pieterse, C. M. J. (2009). Jasmonate signaling in

plant interactions with resistance-inducing beneficial microbes. *Phytochemistry* **70**:1581–

1588.

Wang, Y. Y., Hsu, P. K., and Tsay, Y. F. (2012). Uptake, allocation and signaling of nitrate.

Trends Plant Sci **17**:458–467.

Yang, L. Y., -Yang, and -Zhou, X. Y. (2024). TRANSCRIPTOMIC ANALYSIS OF

ADAPTIVE RESPONSES TO NITROGEN DEFICIENCY IN QUINOA Advance Access

published 2024, doi:10.15666/aeer/2206_52375253.

Yu, G., Wang, L. G., Han, Y., and He, Q. Y. (2012). ClusterProfiler: An R package for

comparing biological themes among gene clusters. *OMICS* **16**:284–287.

Zehnder, G. W., Murphy, J. F., Sikora, E. J., and Kloepper, J. W. (2001). Application of

rhizobacteria for induced resistance. *Eur J Plant Pathol* **107**:39–50.

Zhang, G. Bin, Yi, H. Y., and Gong, J. M. (2014). The Arabidopsis Ethylene/Jasmonic Acid-

NRT Signaling Module Coordinates Nitrate Reallocation and the Trade-Off between

Growth and Environmental Adaptation. *Plant Cell* **26**:3984.

Zhang, X., Zhang, Q., Gao, N., Liu, M., Zhang, C., Luo, J., Sun, Y., and Feng, Y. (2024).

Nitrate transporters and mechanisms of nitrate signal transduction in Arabidopsis and rice.

Physiol Plant **176**.

Zheng, D., Han, X., An, Y., Guo, H., Xia, X., and Yin, W. (2013). The nitrate transporter

NRT2.1 functions in the ethylene response to nitrate deficiency in Arabidopsis. *Plant Cell*

Environ **36**:1328–1337.

Zhu, B., Zhou, Q., Lin, L., Hu, C., Shen, P., Yang, L., An, Q., Xie, G., and Li, Y. (2013).

Enterobacter sacchari sp. nov., a nitrogen-fixing bacterium associated with sugar cane

(Saccharum officinarum L.). *Int J Syst Evol Microbiol* **63**:2577–2582.

Figures

Figure 1. The beneficial effect of SA187 on *Arabidopsis thaliana* plant growth parameters

under high and low nitrate. (a) Total fresh weight (mg) per seedling; (b) Primary root length

(cm); (c) Lateral root density (n/cm = number/cm). Boxplots depict the median (horizontal line),

interquartile range (box), and whiskers to 1.5× IQR; individual data points are overlaid as black

dots, and white circles denote group means (n ≥ 40). Percentage above each bracket gives the

change in mean□with +SA187 relative to Mock. Statistical significance between treatments at

each nitrate level was determined by two-sided, unpaired Student's t-test: *P < 0.05; **P < 0.01;

***P < 0.001; ****P < 0.0001; NS, not significant.

Figure 2. The effect of SA187 on *Arabidopsis thaliana* shoot and root C and N composition

under high and low nitrate. Panels show boxplots of (a) total C content (% DW), (b) total N

content (% DW), (c) C:N ratio, and (d) nitrate content (µmol□g⁻¹ FW) in shoots and roots for

mock and SA187 treatments at each nitrate concentration. Boxplots display the median (center

line), interquartile range (box), and whiskers (1.5× IQR); individual samples are plotted as black

dots, white circles mark group means. Mock and +SA187 are distinguished by lighter versus

darker colors. Wilcoxon rank sum test (Mann–Whitney U test) was used to compare treatments

within each nitrate condition (significance: * = p < 0.05, ** = p < 0.01, *** = p < 0.001, NS =

non-significant).

Figure 3. Global transcriptomic responses to SA187 in *Arabidopsis thaliana* shoots and

roots under high and low nitrate. Hierarchical clustering heat maps of normalized gene

expression (genes mean-centered across all samples) in **(a)** shoots and **(b)** roots of wild-type Col-

0 seedlings treated with Mock or SA187 under high (20 mM) and low (1 mM) nitrate. Clustering

was performed in MeV v4 using Pearson correlation and average linkage. Key clusters showing

the most relevant induction or repression patterns are highlighted and annotated with enriched

Gene Ontology (GO) biological processes obtained by G:Profiler. The color scale denotes

relative expression (yellow = high, blue = low). The original mean counts were subjected to data

adjustment by normalizing genes across all samples. Hierarchical clustering was performed using

average linkage under Pearson correlation (MeV version 4). **(c)** Bar plots representing RNA-seq

differential expression of selected HAT candidate genes (AT1G12940, *NRT2.5*; AT3G45060,

NRT2.6; AT5G60770, *NRT2.4*) significantly regulated by SA187 treatment at 1 mM nitrate.

Significance was determined by RNA-seq data analysis, with asterisks (***) denoting

statistically significant differences (p-value < 0.001).

Figure 4. Effect of SA187 inoculation on *Arabidopsis thaliana* wild-type (Col-0) and mutant

seedlings grown under low nitrate. (a) Total FW of seedlings grown with Mock or +SA187

treatments. Boxplots show median (center line), interquartile range (box) and whiskers to $1.5\times$

IQR; individual plants are overlaid as black dots and white circles denote group means ($n \geq 40$).

Percentage above each bracket gives the relative change in mean fresh weight with +SA187

versus Mock. Statistical significance was determined by two-sided unpaired t-tests within each

genotype (**$P < 0.01$, ***$P < 0.001$). (b) qRT-PCR analysis of high-affinity nitrate transporter

genes in shoots (upper panels) and roots (lower panels) of Arabidopsis Col-0 and *ein2-1*

seedlings. Expression was calculated by the $\Delta\Delta Ct$ method and is shown as $\log_2$ fold-change

relative to the Col-0 Mock samples for each tissue. Black asterisks indicate significance between

Mock and +SA187 treatments (*$P < 0.05$; **$P < 0.01$; ***$P < 0.001$); grey asterisks denote

differences between Col-0 and *ein2-1* under the mock treatment. Error bars represent $\pm$ SEM of

three biological replicates ($n \geq 3$). (c) Total N content (% DW) in shoots and roots under mock or

SA187 treatments in Col-0, *ein2-1*, and *nrt2.5×nrt2.6* seedlings grown under 1 mM nitrate.

Boxplots display the median (center line), interquartile range (box), and whiskers (1.5× IQR);

individual samples are plotted as black dots, white circles mark group means (n ≥ 3). Mock and

+SA187 are distinguished by lighter versus darker colored tones. Wilcoxon rank sum test

(Mann–Whitney U test) was used to compare treatments within each nitrate condition

(significance: * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$).

Figure 5. Simplified model of the regulatory influence of SA187 on the expression of

Arabidopsis thaliana **high-affinity nitrate transporter genes in shoots and roots.** SA187

inoculation promotes plant growth under low nitrate by modulating the expression of HATs

through both ethylene-dependent and independent pathways. *NRT2.5* is strongly upregulated by

SA187 in an EIN2-dependent manner in both shoots and roots, while *NRT2.6* shows tissue-

specific regulation: downregulated in shoots and upregulated in roots, with a partial ethylene

independence. Solid arrows indicate confirmed regulatory relationships; dashed arrows denote

hypothesized or indirect links. The model proposes that SA187 enhances nitrate uptake and

aerial N-allocation via NRT2.5 and NRT2.6 thus contributing to improved growth under limited

nitrate conditions.

Supplementary Figure 1. Differential gene expression in *Arabidopsis thaliana* under high or

low nitrate and with or without SA187 inoculation. Transcriptomic comparisons of

Arabidopsis seedlings inoculated with SA187 versus mock-treated controls under low (1 mM)

and high (20 mM) nitrate concentrations. (a) DEGs in shoot tissues. (b) DEGs in root tissues.

Each comparison indicates the number of significantly upregulated (red) and downregulated

(purple) genes identified by pairwise contrasts. Arrows represent the direction of each

comparison.

Figure 1. The beneficial effect of SA187 on *Arabidopsis thaliana* plant growth parameters under high and low nitrate. (a) Total fresh weight (mg) per seedling; (b) Primary root length (cm); (c) Lateral root density (n/cm = number/cm). Boxplots depict the median (horizontal line), interquartile range (box), and whiskers to 1.5× IQR; individual data points are overlaid as black dots, and white circles denote group means (n ≥ 40). Percentage above each bracket gives the change in mean with +SA187 relative to Mock. Statistical significance between treatments at each nitrate level was determined by two-sided, unpaired Student's t-test: *P < 0.05; **P < 0.01; ***P < 0.001; ****P < 0.0001; NS, not significant.

Figure 2. The effect of SA187 on *Arabidopsis thaliana* shoot and root C and N composition under high and low nitrate. Panels show boxplots of (a) total C content (% DW), (b) total N content (% DW), (c) C : N ratio, and (d) nitrate content ($\mu mol\ g^{-1}$ FW) in shoots and roots for mock and SA187 treatments at each nitrate concentration. Boxplots display the median (center line), interquartile range (box), and whiskers (1.5× IQR); individual samples are plotted as black dots, white circles mark group means. Mock and +SA187 are distinguished by lighter versus darker colors. Wilcoxon rank sum test (Mann–Whitney U test) was used to compare treatments within each nitrate condition (significance: * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$, NS = non-significant).

Figure 3. Global transcriptomic responses to SA187 in *Arabidopsis thaliana* shoots and roots under high and low nitrate. Hierarchical clustering heat maps of normalized gene expression (genes mean-centered across all samples) in **(a)** shoots and **(b)** roots of wild-type Col-0 seedlings treated with Mock or SA187 under high (20 mM) and low (1 mM) nitrate. Clustering was performed in MeV v4 using Pearson correlation and average linkage. Key clusters showing the most relevant induction or repression patterns are highlighted and annotated with enriched Gene Ontology (GO) biological processes obtained by G:Profiler. The color scale denotes relative expression (yellow = high, blue = low). The original mean counts were subjected to data adjustment by normalizing genes across all samples. Hierarchical clustering was performed using average linkage under Pearson correlation (MeV version 4). **(c)** Bar plots representing RNA-seq differential expression of selected HAT candidate genes (AT1G12940, *NRT2.5*; AT3G45060, *NRT2.6*; AT5G60770, *NRT2.4*) significantly regulated by SA187 treatment at 1 mM nitrate. Significance was determined by RNA-seq data analysis, with asterisks (***) denoting statistically significant differences (p-value < 0.001).

Figure 4. Effect of SA187 inoculation on *Arabidopsis thaliana* wild-type (Col-0) and mutant seedlings grown under low nitrate. (a) Total FW of seedlings grown with Mock or +SA187 treatments. Boxplots show median (center line), interquartile range (box) and whiskers to 1.5× IQR; individual plants are overlaid as black dots and white circles denote group means (n ≥ 40). Percentage above each bracket gives the relative change in mean fresh weight with +SA187 versus Mock. Statistical significance was determined by two-sided unpaired t-tests within each genotype (**P < 0.01, ***P < 0.001). **(b)** qRT-PCR analysis of high-affinity nitrate transporter genes in shoots (upper panels) and roots (lower panels) of Arabidopsis Col-0 and *ein2-1* seedlings. Expression was calculated by the ΔΔCt method and is shown as $\log_2$ fold-change relative to the Col-0 Mock samples for each tissue. Black asterisks indicate significance between Mock and +SA187 treatments (*P < 0.05; **P < 0.01; ***P < 0.001); grey asterisks denote differences between Col-0 and *ein2-1* under the mock treatment. Error bars represent ± SEM of three biological replicates (n ≥ 3). **(c)** Total N content (% DW) in shoots and roots under mock or SA187 treatments in Col-0, *ein2-1*, and *nrt2.5×nrt2.6* seedlings grown under 1 mM nitrate. Boxplots display the median (center line), interquartile range (box), and whiskers (1.5× IQR); individual samples are plotted as black dots, white circles mark group means (n ≥ 3). Mock and +SA187 are distinguished by lighter versus darker colored tones. Wilcoxon rank sum test (Mann–Whitney U test) was used to compare treatments within each nitrate condition (significance: * = p < 0.05, ** = p < 0.01, *** = p < 0.001).

Figure 5. Simplified model of the regulatory influence of SA187 on the expression of Arabidopsis thaliana high-affinity nitrate transporter genes in shoots and roots. SA187 inoculation promotes plant growth under low nitrate by modulating the expression of HATs through both ethylene-dependent and independent pathways. *NRT2.5* is strongly upregulated by SA187 in an EIN2-dependent manner in both shoots and roots, while *NRT2.6* shows tissue-specific regulation: downregulated in shoots and upregulated in roots, with a partial ethylene independence. Solid arrows indicate confirmed regulatory relationships; dashed arrows denote hypothesized or indirect links. The model proposes that SA187 enhances nitrate uptake and aerial N-allocation via NRT2.5 and NRT2.6 thus contributing to improved growth under limited nitrate conditions.