Bifidobacterium longum subsp. *nexti* subsp. nov., a novel subspecies isolated from infant stool

Dena Ennis[1], Sergio Andreu-Sánchez[2], Yarden Levin[1], Omer Goldberger[1], Cyrus A. Mallon[2], Yuvashankar Kavanal Jayaprakash[2,3], Yan Shao[4], Shijie Zhao[6,7], Trishla Sinha[2], Jingyuan Fu[2], Rinse K. Weersma[5], Eric Alm[6,7,8], Orna Amster-Choder[1], Trevor D. Lawley[4], Hermie J.M. Harmsen[3], Douwe van Sinderen[9], Alexandra Zhernakova[2], Moran Yassour[1,10*]

Affiliations:
1. Microbiology & Molecular Genetics Department, Faculty of Medicine, The Hebrew University of Jerusalem, Jerusalem, Israel
2. Department of Genetics, University of Groningen and University Medical Center Groningen, The Netherlands
3. Department of Microbiology, University of Groningen and University Medical Center Groningen, The Netherlands
4. Host–Microbiota Interactions Laboratory, Wellcome Sanger Institute, Hinxton, UK
5. Department of Gastroenterology and Hepatology, University of Groningen and University Medical Center Groningen, The Netherlands
6. Department of Biological Engineering, Massachusetts Institute of Technology, Cambridge, MA, USA
7. Center for Microbiome Informatics and Therapeutics, Massachusetts Institute of Technology, Cambridge, MA, USA
8. The Broad Institute of MIT and Harvard, Cambridge, MA, USA
9. APC Microbiome Institute and School of Microbiology, University College Cork, Cork T12 YT20, Ireland
10. The Rachel and Selim Benin School of Computer Science and Engineering, The Hebrew University of Jerusalem, Jerusalem, Israel
    *corresponding author: moranta@mail.huji.ac.il

Keywords: Bifidobacterium longum, human gut, infant, microbiome.
Author Notes: The GenBank accession numbers for the 16S rRNA encoding gene and the genome of LL6991 are PV770940 and JBPCCJ000000000, respectively.

Abstract

Bifidobacterium species are well-established members of the human gut microbiome, particularly prominent during infancy, contributing to host health. Within this genus, *Bifidobacterium longum* (*BL.*) is a widespread species found in both infant and adult guts, known for its complexity and functional diversity among its known subspecies: *BL. longum*, *BL. infantis* and *BL. suis*. Here, using genomic and phylogenetic tools we propose a novel subspecies within the *BL.* species, *Bifidobacterium longum* subsp. *nexti* subspecies novel. We analyzed 435 *BL.* genomes using a polyphasic taxonomic approach comprising average nucleotide identity (ANI), digital DNA–DNA hybridization (dDDH), and pangenome analysis. We identified nine *BL.* strains, isolated from human infants and adults stool samples, as members of a distinct lineage within the *BL.* species. The type strain, LL6991, was isolated from the stool of a two-week-old Dutch infant in the Lifelines NEXT birth cohort. Phenotypically, *BL. nexti* exhibits a distinct morphological pattern, predominantly forming rod-shaped cells, often in chains with visible septa, contrasting with the Y-shaped morphology commonly observed for other *BL.* subspecies. Furthermore, *BL. nexti* demonstrates unique metabolic capabilities, including efficient utilization of fructose, and starch, carbohydrates not metabolized well by other tested *BL.* subspecies. This ability may be attributed to specific genes, such as a gene predicted to encode an extracellular amylopullulanase. This characterization expands the known diversity within the *BL.* species and provides insights into *BL. nexti*'s unique adaptations and potential ecological roles within the human gut, especially in infants. Based on the consistent results from genotypic, phylogenetic, and phenotypic analyses, a novel subspecies with the name *Bifidobacterium longum* subsp. *nexti*, with type strain LL6991 (=NCCB 101085 =DSM 120337), is proposed.

Introduction

Bifidobacterium species are well-established members of the human gut microbiome, particularly prominent during infancy, where they contribute to various aspects of host health, including metabolic processes, immune system development, and providing resistance against pathogens[1–6]. Within this genus, *Bifidobacterium longum* (*BL.*) is recognized as a widespread species commonly found in both the infant and adult gut[7–10]. Growing research continues to reveal the complexity and functional diversity within this species in the human gut, where five main subspecies have been reported in several studies: *B. longum* subsp. *longum*, *B. longum* subsp. *infantis*, *B. longum* subsp. *suis*, *B. longum* subsp. *iuvenis* and *B. longum* subsp. *suillum*[11–14]. The original *BL. suis* taxon has recently been divided into three subclusters; *BL. suis*, *B. longum* subsp. *iuvenis*[14], a

subspecies originally identified in Bangladesh infants as transitional *BL.[15]*, and *B. longum* subsp. *suillum*, a *BL. suis* subspecies which is urease negative[16]. These subspecies are increasingly understood to possess distinct functional capabilities. For instance, *BL. infantis* is notably adapted for the utilization of human milk oligosaccharides (HMOs)[17], and therefore more prevalent in the infant gut, while *BL. longum* is associated with the metabolism of plant-derived oligosaccharides[18].

The *Bifidobacterium* genus is famously named for the characteristic bifid or Y-shape exhibited by some strains when observed microscopically[19]. Historical accounts trace the identification of these Y-shaped bacteria back to 1899, when French pediatrician Henri Tissier isolated a bacterium with a Y-shaped morphology from the intestinal microbiota of breast-fed infants[20]. He named it "bifidus" recognizing its distinctive shape, which is now a defining feature of the *Bifidobacterium* genus. This distinctive shape has been visualized using various microscopy techniques, such as phase contrast microscopy and scanning electron microscopy (SEM)[21, 22]. Although this morphology has long been considered a hallmark of *Bifidobacterium*, studies have shown that their cell shape is pleomorphic, with high variability influenced by environmental and growth conditions[22, 23].

Defining bacterial species and subspecies accurately is fundamental to understanding microbial diversity and function. Historically, classification relied heavily on phenotypic characteristics and core gene sequencing, such as the 16S rRNA gene[24, 25]. However, these methods often lack sufficient resolution to delineate closely related taxa such as subspecies. For example, 16S rRNA gene can have a slow evolutionary rate so that even full-length sequences may not provide adequate resolution to distinguish between closely related species or subspecies[26]. Modern prokaryotic taxonomy increasingly employs a polyphasic approach that integrates traditional methods with advanced genome-based analyses[25, 27]. Advancements in whole-genome sequencing (WGS) have facilitated the adoption of genomic metrics such as average nucleotide identity (ANI) and digital DNA–DNA hybridization (dDDH), offering robust, quantitative measures of overall genomic relatedness[28]. These genome-based approaches have become standard tools in microbial taxonomy, providing higher resolution for species and subspecies delineation compared to traditional methods. Commonly accepted thresholds for species delineation using these methods are 95% for ANI and 70% for dDDH, which have shown a strong correlation with one another. Subspecies are typically distinguished by distinct clustering patterns in these analyses, with dDDH values generally falling below 79–80%. By leveraging WGS data, researchers can gain a more accurate view of microbial diversity and delineate finer taxonomic classifications, including subspecies.

The use of modern taxonomic approaches has significantly refined the classification of
complex microbial groups. For example, a recent study combining metagenomic and
genomic data revealed that *Segatella copri* is not a monotypic species but consists of four
distinct clades, each associated with functional diversity[29]. Further, by integrating targeted
isolation with large-scale metagenomic analysis, researchers identified 13 distinct *S. copri*-
related species, expanding the known diversity beyond the initial four clades[30]. In a
different study of *Escherichia coli*, long-read sequencing enabled the application of ANI and
dDDH as robust alternative typing methods, potentially offering higher taxonomic
resolution[31]. In another study conducted in Bangladesh, analysis of *BL.* using
metagenome-derived marker genes and ANI led to the discovery of a novel *BL.* subspecies
known as transitional *BL.[15]*. A recent large-scale phylogenomic analysis of the *BL.* species
identified four clades, including a novel subspecies found exclusively in infants from
industrialised countries[32].

Given the known diversity and functional specialization within the *BL.* species, and the
increasing power of genomic methods to resolve fine-scale taxonomic structures, a
comprehensive analysis of a large collection of *BL.* genomes and associated phenotypic
data is warranted to fully delineate the species and its subspecies. Such an analysis may
reveal new lineages and provide insights into their unique genetic content and potential
ecological roles. Here, we applied comprehensive genomic and phenotypic analyses to
uncover a previously unrecognized lineage within the *BL.* species, which we describe here
as *Bifidobacterium longum* subsp. *nexti* subsp. novel.

122 Methods

123 Bacteria isolation and genomic retrieving

In this study we used 435 *BL.* genomes from public data and from strains we isolated. Public
genomes were selected based on criteria of over 95% completeness and less than 5%
contamination, found by CheckM2 (1.0.2)[33], to minimize analytical noise. A total of 114
genomes were retrieved from NCBI by searching for entries matching *Bifidobacterium*
longum. Similarly, 166 genomes were obtained from the JGI IMG/M (Integrated Microbial
Genomes & Microbiomes) database. Subspecies annotations were extracted from strain
names when available. In addition, four *BL.* genomes from Chilean isolates were included
from Díaz et al.[34].

Genomes from strains we isolated were taken from multiple cohorts and geographical
locations. 49 genomes were sourced from the Baby Biome Study (BBS) in the United

Kingdom[35], with details on isolation, sequencing and assembly as previously described[35,
36]. An additional 99 genomes were obtained from the MicrobeMom study in Ireland, with full
methods described therein[37].

Two isolates were isolated as part of the Dutch Microbiome Project (DMP)[38], as follows:
we collected stool samples from 550 population-based participants from the Lifelines Dutch
Microbiome project. We used two distinct sampling strategies to culture bacterial isolates. In
the first round of culturing, we chose selective culturing conditions for specific taxonomic
groups, in this caseDe Man–Rogosa–Sharpe (MRS, Millipore, 69964) medium under
anaerobic conditions for genus Bifidobacterium (phylum Actinobacterium). We assigned
species for each isolate via multiplexed 16S sequencing for each isolate and performed
whole-genome sequencing (WGS) on a subset (on average two isolates per species per
donor). In the second round of culturing, we selected dozens of isolates per donor based on
diverse morphological characteristics, cultured all in a rich medium, Brain Heart Infusion
(Bacto 237300), and performed WGS. DNA extraction was performed using the PureLink
Pro Genomic DNA Purification (K182104A) Kit.

The type strain, LL6991, was isolated in April 2025 in Groningen, Netherlands (53.2210° N,
6.5770° E) from a fecal sample from a two-weeks old infant as part of the Lifelines NEXT
cohort, a birth cohort designed to study the effects of intrinsic and environmental
determinants on health in mothers and children[39]. A detailed description of the design,
inclusion and exclusion criteria has been described previously[40]. For isolation, a stool
sample that was known to contain the type strain was spread-plated onto Yeast Casitone
Fatty acid agar[41] supplemented with 20 mM Lactose and 0.05% (w/v) cystine (YCFA-
Lactose) were then incubated in an anaerobic chamber (90% nitrogen, 5% hydrogen, 5 %
carbon dioxide) for 72 hours. Each quadrant of the spread plate was subcultured onto fresh
YCFA-Lactose agar plates after which individual colonies were picked up randomly and
subcultured on fresh YCFA-Lactose agar for colony purification. These purified colonies
were identified up to species level using Bruker MALDI TOF (Bruker Daltonics, Germany)
where one of the isolates obtained was identified as *Bifidobacterium longum.* Further, the
DNA of *B. longum* isolate was extracted using QiAmp DNA Mini kit (Qiagen, Hilden,
Germany) and PCR reactions were run using primers that bind to gene loci unique to *BL.*
nexti to verify its subspecies identity.

DNA extraction, sequencing and assembly

For the type strain, LL6991, and for an additional *BL. nexti* strain, MB0413, DNA was
extracted using MagAttract HMW DNA Kit (#67563, Qiagen). Short read sequencing libraries

were prepared using Nextera XT DNA Library Preparation kit (FC-131-1096, Illumina)
according to the manufacturer's recommended protocol with half of the volume and the DNA.
Samples were sequenced using Illumina single-end 150□bp sequencing on a NextSeq 500
device. Long reads were retrieved using native barcoding sequencing kit (SQK-NBD114-24,
ONT) on a MinION device. Basecalling and demultiplexing of long reads was done using
Dorado (Oxford Nanopore Technologies). Hybrid assembly was performed using SPAdes
(v4.0.0)[42, 43].

For the two strains derived from the UK Baby Biome Study infants, 37822_2_275 and
37822_2_213, hybrid genome assemblies were generated. DNA was extracted using the
MasterPure Complete DNA and RNA Purification Kit (Lucigen), then sequenced on the
Illumina NovaSeq 6000 (384-plex, 2 x 151bp, Illumina) and on the Oxford Nanopore GridION
(24-plex) or PromethION (96-plex) using R9.4.1 and R10.4.1 flow cells with basecalling in
'super-accurate (SUP) model using Dorado. Hybrid genome assemblies were generated
using the Dragonflye with Flye for long-read assembly ('--nanohq' with R9.4.1 data and '--
nano_hq --read-error 0.03' with R10.4.1 data), Medaka
(https://github.com/nanoporetech/medaka) for long-read polishing and Polypolish for short-
read polishing.

Genomic and phylogeny analysis

Assessment of genome quality and retrieving genome information was performed using
CheckM2[33] and Bakta[44–47]. Average nucleotide identity between genomes was
estimated using FastANI (1.33)[48]. The selection of the number four for clustering was done
using the Silhouette method, where the optimal number of clusters is the one with the
highest average silhouette width. Gene annotations were created using Prokka (1.14.6)[49],
which uses BioPerl[50], GNU Parallel[51], BLAST+[52] and Prodigal[46], and pangenomes
were analyzed with Roary (3.13.1)[53]. 16s rRNA gene sequence was retrieved from
genomes using Barrnap (0.9)[54]. Multiple sequence alignment and distance calculation of
16S rRNA gene was performed using Clustal Omega (1.2.4)[55]. 16s rRNA and core gene
phylogenetic trees were created using RAxML with raxmlHPC-SSE3[56]. Genome synteny
was assessed using minimap2 (2-2.24)[57] with -ax flag asm20, and visualised using the
gggenomes (1.0.1) R package[58]. Gene clusters were searched using Integrative
Genomics Viewer (IGV)[59] and visualised using gggenomes[58]. Genes annotated as
hypothetical proteins by Prokka[49] were further analyzed using BLASTx[52] against non-
redundant protein sequences (nr) database and HHpred[60] to infer potential functions.
Biosynthetic gene clusters (BGCs) were searched using antiSMASH (8.0.1)[61].

dDDH values were calculated using the genome-to-genome distance web-based calculator
(GGDC 3.0[62, 63]). Genomes were uploaded to the web server, genomes were aligned
using BLAST, distances and dDDH values were calculated using the d2 formula (identities /
HSP length) of GGDC, which uses a generalized linear model (GLM) inferred from an
empirical reference dataset comprising real DDH values and genome sequence[63].

BL. nexti-specific gene sequences were mapped against the Uniref90 database[64, 65]
using uniref_annotator.py script[66], which utilizes DIAMOND[67] for alignment. Uniref90
annotations were then assigned a gene Gene Ontology (GO) annotations using UniProt.

Further analysis was done using an in-house R (4.4.0) script utilizing dplyr[68] (1.1.4),
tidyr[69] (1.3.1), stringr[70] (1.5.1) and tidyverse[71] (2.0.0). Plots were created using
ggplot2[72] (3.5.1) and ggforce[73] (0.4.2), colors were used from RColorBrewer[74] (1.1.3).
Heatmaps were created using pheatmap[75] (1.0.12) using the cutree option to divide into
clusters. Distance was calculated using the dist function with euclidean distance.
Dendrograms were constructed using a neighbour joining algorithm with the nj function from
the ape package[76] (5.8.1) and visualized using ggtree[77] (3.12.0). T-tests were
pregormed using using R.

Culture Conditions and Microscopy

Culturing and microscopy was performed on a representative of *BL. longum*, *BL. infantis* and
three *BL. nexti* isolates; MB0401, MB0413 and LL6991. *BL. infantis* and *BL. longum* were
obtained from the BMB cohort[7].

Strains were grown routinely at 37ºC in an anaerobic chamber (COY) that was maintained at
> 2.5% H_2 by flushing with a gas mix containing 20% H_2, 5% CO_2, and 75% N_2. Strains
were grown in supplemented Brain Heart Infusion (sBHI) media, supplemented with 50 ml/L
fetal bovine serum (FBS; Sigma, F2442), 10 ml/L trace vitamins (ATCC® MD-VS™), 10 ml/L
trace minerals (ATCC® MD-TMS™), 10 ml/L vitamin K1 and hemin (BBL, 212354), 1 g/L D-
(+)-Cellobiose (Alfa Aesar, 528507), 1 g/L D-(+)-Maltose (Caisson, 6363537), 1 g/L D-(+)-
Fructose (Sigma Aldrich, 1286504), and 0.5 g/L L-Cysteine (Acros Organics, 52904). For
imaging, strains were cultured overnight and then diluted 30-fold in fresh medium and grown
to exponential phase. Next, they were stained with DAPI (4-,6-diamidino-2-phenylindole,
Sigma-Aldrich Cat #D8417) and FM 4-64 Dye (N-(3-Triethylammoniumpropyl)-4-(6-(4-
(Diethylamino) Phenyl) Hexatrienyl) Pyridinium Dibromide, Invitrogen Cat# T13320). Cells
were placed on an agarose pad with uncoated coverslips and imaged using a Nikon Eclipse
Ti-E inverted microscope equipped with Perfect Focus System (PFS) and ORCA Flash 4

camera (Hamamatsu photonics). Images were processed, two-dimensional (2D)
deconvolution and bacteria cell length measurement was performed using NIS Elements-AR
software.

Additionally, to test growth on various carbohydrates, a 2% w/v sugar solution was mixed
with 2x Man-Rogosa-Sharpe (mMRS) medium at a 1:1 ratio (1.5 mL of each) to yield a final
3 mL solution of 1x mMRS medium containing 1% sugar. 2x mMRS medium was made with
20 g/L peptone from casein, tryptic digest; 10 g/L yeast extract; 4 g/L potassium phosphate;
10 g/L sodium acetate; 4 g/L ammonium citrate; 0.4 g/L magnesium sulfate; 0.05 g/L
manganese sulfate; and 2 g/L (2 mL/L) Tween 80. 0.5 mg/ml L-cysteine hydrochloride was
added before use. The strains were inoculated into this medium at a 1:100 dilution from
cultures grown in liquid supplemented BHI overnight. These cultures were prepared by
picking colonies off of CHG plates incubated for 72 hours and transferring them into liquid
CHG to incubate overnight. The initial plates were seeded using glycerol stocks of each
strain preserved in a -80 degree freezer. Two biological replicates were used for each strain
from two isolated colonies on their respective plates.

250 Results

251 Phylogeny of the *Bifidobacterium longum* clade

To explore the taxonomic structure of *Bifidobacterium longum* (*BL.*) species, we compared
the genomic sequence of 435 genomes, 284 of which were obtained from public databases
and 151 newly generated from isolated strains obtained from stool samples. These new
isolates originated from several studies involving British infants[35, 36], Irish mothers[78]and
infants[37] and both adults and infants in the Netherlands[38–40] (**Supplementary table 1**).
Using these genomes, we estimated the pairwise average nucleotide identity (ANI), and
observed four distinct clusters of genomes (**Fig. 1A, Methods**). Three of these clusters had
genomes that were previously annotated to one of *BL.* clades (37 belonged to *BL. infantis*,
122 to *BL. longum* and 13 from the *BL. suis* clade – which included genomes annotated as
BL. suis, *BL. suillum* and *BL. iuvenis*). Interestingly, the remaining cluster had nine
unannotated genomes, including two public genomes and seven new genomes from
internally isolated strains. We hereby propose the name *Bifidobacterium longum* subsp. *nexti*
novel for strains belonging to this novel clade, since the strain was isolated from an infant in
the Lifelines NEXT cohort. Some publicly available genomes appeared to be misannotated
according to their ANI-based clustering, specifically three *BL. longum* genomes clustered
within the *BL. suis* group, and one *BL. infantis* genome was found within the *BL. longum*

cluster. Previous reports have noted such mis-annotations, including for genome assembly
GCA_000092325, which is likely misclassified as *BL. longum* and instead represents *BL.*
suis[34, 79]. We next compared the ANI across the various clades, and found that *BL. nexti*
had an inter-subspecies ANI of 99.3% (**Fig. 1B**). The lowest ANI value observed among all
genomes across the clades was 94.7%, supporting the classification of all subspecies within
the *BL.* species (**Fig. 1B**).

To further support the classification of *BL. nexti* as a distinct subspecies, we applied gene-
level characterization across all genomes described above. Overall, we examined 18,591
genes in the pangenome of the *BL.* species. We identified 789 core genes present in >99%
of genomes, and 17,802 accessory genes. Using alignment of the core genes we
constructed a phylogenetic tree on a subset of 24 representative genomes, and found four
distinct clades with genomes belonging to *BL. nexti* forming their own group (**Fig. 1C**). In
addition, using the accessory gene presence/absence profiles of all annotated genomes we
constructed a dendrogram revealing the same result (**Supplementary Fig. 1A**).

To further validate the existence of the novel clade, we employed digital DNA-DNA
hybridization (dDDH), a widely used *in silico* method for assessing genomic similarity
between sequences (**Methods**). The dDDH values between *BL. nexti* and the other
subspecies ranged from 62% to 75.9%, with *BL. longum* being its closest subspecies. Inter-
subspecies dDDH values ranged from 94% to 99.3% (**Methods, Supplementary Fig. 1B**).
These values meet the 79-80% dDDH suggested threshold for delineating subspecies[80].
We next generated a dendrogram based on these values, showing that *BL. nexti* clusters
within the *BL.* clade, however being distinct enough to be a novel discrete subspecies (**Fig.**
1D). Interestingly, *BL. infantis* was shown to exhibit low dDDH values when compared to
other subspecies (62-69.4%), supporting the notion that it is more distinct than other *BL.*
subspecies, as has previously been suggested[81, 82]. The dDDH results showed strong
correlation with the estimated ANI values, collectively indicating that *BL. nexti* is most closely
related to *BL. longum* and most distantly related to *BL. infantis* (**Fig. 1D**).

Finally, we conducted a phylogenetic analysis based on the complete 16S rRNA gene
sequence, using the *Bifidobacterium breve* type strain genome as an outgroup. This
confirmed that all 24 selected genomes share their most recent common ancestor, while *B.*
breve forms, as expected, a sister branch (**Supplementary Fig. 1C**). Although some
genomes from the same subspecies clustered closely, the 16S rRNA gene lacked sufficient
phylogenetic resolution to clearly distinguish the known *BL.* subspecies from the novel clade
(**Supplementary Fig. 1C**).

Genomic features of *Bifidobacterium longum* subsp. *nexti* nov.

To further investigate *BL. nexti*, we analyzed the genomic features of both newly isolated strains and publicly available genomes. The *BL. nexti* strains were isolated from human stool; five from infants in the Netherlands, Ireland, and the United Kingdom, and two from adults in the Netherlands. Publicly available *BL. nexti* genomes were originally isolated from human milk[83] and vaginal microbiomes[84], sourced from Italy and Canada, respectively. The average genome length of *BL. nexti* was 2,400,483 base pairs (bp), comparable to that of *BL. longum* and *BL. suis*, but shorter than *BL. infantis* (t-test, p= $4*10^{-6}$; **Supplementary Fig. 2A**). The guanine+cytosine (G+C) content across all *BL. nexti* isolates ranged from 59-60 mol%, consistent with other *BL.* subspecies (t-test, p>0.3, **Table 1**).

To examine genome structure conservation and variation across *BL.* subspecies, we performed whole-genome synteny analysis using the type strain from each subspecies. We revealed that *BL. nexti* exhibits a high level of genomic similarity to the other *BL.* subspecies (**Fig. 2**). However, we observed a few notable genomic inversions between subspecies, reflecting apparent genome plasticity of the *BL.* genome[81]. Interestingly, *BL. infantis* was shown to contain genomic regions that are absent from genomes of the other subspecies, consistent with its overall larger genome size. In addition, *BL. nexti* and *BL. suis* were determined to lack a genomic region that is present in *BL. longum* and *BL. infantis.*

To investigate *BL. nexti* specific genes, we searched for genes that are unique to the *BL. nexti* genomes. We identified 112 genes that are uniquely present in all *BL. nexti* genomes and were found in less than 1% of other subspecies (**Fig. 3A**). These genes are distributed throughout the entire *BL. nexti* genome (**Fig. 2**). They corresponded to 222 UniRef90 gene clusters and were primarily linked to Gene Ontology (GO) categories involved in carbohydrate metabolic processes (GO:0005975), carbohydrate transport (GO:0008643), and amino acid transport (GO:0006865). Among these unique genes, we identified a gene predicted to encode an extracellular amylopullulanase, previously characterized in *B. breve[85]*, suggesting that *BL. nexti* strains possess the ability to degrade starch, a phenotypic trait not observed in other *BL.* subspecies[86–88]. This is consistent with a recent publication that showed enrichment for starch utilisation in an unknown *BL.* subspecies, based on KEGG (K01200) and COG (COG1523) analysis[32]. The predicted starch utilization trait suggests an adaptation of *BL. nexti* to metabolize starch-like polysaccharides.

We next searched for *BL. nexti*-specific gene-clusters, namely genes that are located in close proximity on the genome. We found two distinct clusters, each consisting of four or more adjacent genes (**Fig. 3B**). Gene annotations indicate that these clusters are/were

involved in carbohydrate metabolism. The first cluster appears to be a genetically decayed
remnant of a sugar utilization operon. While most genes are likely non-functional, the
remaining MFS transporter may still be active and potentially specific for melibiose, which is
a component of raffinose and stachyose, though this remains a prediction based on
sequence similarity. The second cluster includes genes predicted to encode a carbohydrate-
specific ABC-type transport system. The GH35 enzyme encoded within the cluster lacks a
signal peptide, suggesting it functions intracellularly, likely acting on a substrate imported by
the afore-mentioned-ABC transporter. Notably, some genes within these clusters encode
hypothetical proteins, suggesting the presence of novel functionalities that may contribute to
the unique physiological traits of *BL. nexti*.

Description of *Bifidobacterium longum* subsp. *nexti* subsp. nov.

Bifidobacterium longum subsp. *nexti* (nex'ti. N.L. gen. n. nexti, since the type strain was
isolated from an infant in the Lifelines NEXT Dutch cohort).

Cells are Gram-stain-positive. Colonies are circular, with entire margins, smooth, convex,
beige and approximately 1 mm in diameter after 3 days cultivation at 37 °C under
anaerobic conditions. Growth in supplemented BHI and modified MRS with lactose is good
under anaerobic conditions and weak under aerobic conditions. Cells show growth on
lactose, fructose, galactose, melibiose, lactulose, mlatose, sucrose, glucose, raffinose and
turanose.

The type strain, LL6991 (=NCCB 101085 =DSM 120337) was isolated from the stool of a
two-week old Dutch infant born vaginally and breastfed. The DNA G+C content of this strain
is 60% mol.

The GenBank accession numbers for the 16S rRNA encoding gene and the genome of
LL6991 are PV770940 and JBPCCJ000000000, respectively.

Subsequently, to explore the microscopic phenotype of *BL. nexti*, we performed fluorescent
staining targeting the cell wall and DNA to visualize cell shape during the exponential phase.
Using microscopic imaging we revealed that *BL. nexti* cells predominantly exhibit a rod-
shaped morphology, frequently forming chains with a visible septum between cells (**Fig. 4A**).
This suggests active cell division with incomplete separation of daughter cells. In many
cases, we observed accumulation of the cell wall at the cell poles, possibly indicating sites of
upcoming division or budding and further cell splitting. In contrast, a significant proportion of
BL. infantis and *BL. longum* cells displayed Y-shaped morphologies (**Fig. 4A**), a feature
previously documented as characteristic of *Bifidobacterium* species, and particularly

visualized in *BL.[22, 23]*. Notably, Y-shaped cells were considerably less prevalent in *BL.*

nexti strains (<1% of cells), indicating a distinct morphological pattern under similar growth

conditions. These observations suggest pleomorphism within the *BL.* species, with

morphological variations manifesting at the subspecies level.

The average cell length of *BL. nexti* strains we examined was 1.64 ± 0.41 µm,

significantly shorter than that of *BL. infantis* (p = 8.3e-12) and slightly shorter than *BL.*

longum (p = 0.017, **Figure 4B**). This difference may be related to the larger genome size of

BL. infantis[89, 90], as well as to the distinct cellular morphologies observed, however this

may vary between strains.

To further explore the functional capabilities of *BL. nexti*, we assessed its growth on various

carbohydrates and compared the results with the growth capabilities of *BL. infantis* and *BL.*

longum. All tested strains of *BL. nexti*, *BL. longum*, and *BL. infantis* exhibited robust growth

on lactose (OD600 > 1, **Fig. 4C**). In contrast, growth on fructose and maltodextrin was

significantly stronger in *BL. nexti* strains, whereas *BL. longum* and *BL. infantis* displayed only

low to moderate growth (OD600 0.2-1, **Fig. 4C**). Notably, while *BL. longum* and *BL. infantis*

showed no growth on starch, *BL. nexti* strains demonstrated vigorous growth (**Fig. 4C**). This

unique metabolic capability may be attributed to specific genes present in *BL. nexti*, such as

the identified amylopullulanase-encoding gene, a homologue of which is known to enable

starch utilization in *B. breve[85]*.

Discussion

Here, using comprehensive genomic and phenotypic analyses we showed that *BL. nexti*

subsp nov. constitutes a distinct subspecies within the *BL.* species. This classification is

robustly supported by multiple complementary genomic approaches, including average

nucleotide identity (ANI), digital DNA–DNA hybridization (dDDH), and pangenome analysis.

These findings align with modern prokaryotic taxonomy, which employs a polyphasic

approach integrating traditional methods with advanced genome-based analyses to resolve

finer taxonomic classifications like subspecies[25, 27].

One notable finding is the distinct morphological pattern observed in *BL. nexti*. While the

Bifidobacterium genus is famously named for its characteristic bifid or Y-shape[19, 20],

microscopic examination revealed that *BL. nexti* cells predominantly exhibited a rod-shaped

morphology, frequently forming chains with visible septa, when cultured in rich

supplemented BHI media. In contrast, *BL. infantis* and *BL. longum* cells under similar

conditions displayed a significant proportion of Y-shaped morphologies (>30% of cells). This

suggests pleomorphism within the *BL.* species, with morphological variations evident at the
subspecies level. Pleomorphism has previously been observed in *Bifidobacterium* species
and is known to vary not only between subspecies but also between individual strains,
influenced by environmental and growth conditions[21]. Recent research on *Bifidobacterium*
breve indicates that while the Y-shape can be observed, cells are primarily rod-shaped in
nutritionally rich media and *in vivo*, with the branched morphology becoming more frequent
in sub-optimal environments[91].

We found that *BL. nexti* exhibits unique metabolic capabilities, particularly its unique
capability to efficiently utilize fructose, and starch-like carbohydrates (maltodextrin and
starch). These carbohydrates could not be metabolized efficiently by *BL. infantis* or *BL.*
longum in this study and in most strains in previous studies[86, 92]. This distinct metabolic
profile is likely attributed to specific genes present in *BL. nexti*, such as the identified
amylopullulanase-encoding gene, a homologue of which is known to enable starch utilization
in *B. breve[85]*. The ability to utilize fructose is particularly interesting in the context of the
infant gut, where *BL. nexti* was originally isolated. In early life, infants lack the GLUT-5
transporter, which is specific for fructose absorption, implying that any consumed fructose is
likely metabolized by the gut microbiota, thereby altering its composition[93–95].
Furthermore, fructoselysine, a food process contaminant prevalent in infant formula due to
heating processes, seems to lead to an increased fructoselysine degradation activity in the
gut microbiota of infants, indicating adaptation towards dietary fructoselysine exposure[96].
We did not identify specific genes directly responsible for fructose utilization in the *BL. nexti*
genomes. However, we observed a broader enrichment of genes associated with
carbohydrate metabolism. Further research is required to uncover the underlying
mechanisms driving fructose utilization in this subspecies.

To deepen our understanding of *Bifidobacterium* diversity, particularly at the subspecies
level, it is crucial to carry out systematic studies across varied geographical regions and host
populations. In this study, we isolated *BL. nexti* strains from infants in the Netherlands,
Ireland, and the United Kingdom, as well as from adults in the Netherlands, highlighting the
importance of sampling across different ages and regions. Such targeted isolation, combined
with comprehensive genomic and phylogenetic analysis, lays the foundation for identifying
novel taxa.

In conclusion, the characterization o*f BL. nexti* subsp. nov. not only expands our
understanding of the diversity within the *BL.* species but also highlights its unique
morphological and metabolic adaptations, particularly its ability to utilize various

carbohydrates like starch and fructose. These findings provide insights into its potential
ecological roles within the human gut microbiome, especially in infants.

439 Ethical approval

For the DMP cohort, research approvals were obtained from MIT (COUHESprotocol
#1612797956) and from the Medical Ethical Committee of the University Medical Center
Groningen,the Netherlands; IRB number: 2017.1520. For Lifelines Next protocol has been
approved by the local Institutional Review Board (METc 2015/600). For MicrobeMom ethical
approval was received from National Maternity Hospital research ethics committee in
February 2016 (EC 35.2015). BBS study was approved by the NHS London – City and East
Research Ethics Committee (REC reference 12/LO/1492). All participants provided informed
consent upon enrollment in all studies, either by themselves or by a parent or legal guardian.

448 Funding and conflict of interests

YS and TL acknowledge core funding from the Wellcome Trust (206194 and
220540/Z/20/A). COI: T.D.L. is the co-founder and CSO of Microbiotica. AZ received travel
funding from Nestle to attend the Nestlé Nutrition Institute Workshop 101. AZ is supported by
the Netherlands Organization for Scientific Research NWO-VICI grant VI.C.232.074, the
NWO Gravitation grant ExposomeNL 024.004.017, the NWO KIC grant
KICH1.LWV04.21.01.,the ZonMW ME/CFS grant 10091012110015, EU Horizon Europe
Program grant INITIALISE (101094099), and EU Horizon Europe Program grant DarkMatter
(ID-DarkMatter-NCD, 101136582). MY is the Rosalind, Paul and Robin Berlin Faculty
Development Chair in Perinatal Research, and was also funded by the Azrieli Family
Foundation.

459 Acknowledgements

460 We would like to acknowledge Anouk Toxopeus for her technical assistance with bacterial
461 isolation. We would also like to thank all the mothers and infants who participated in the
462 studies included here for their generous time and engagement.

463 References

464 1. **Turroni F, Milani C, Duranti S, Lugli GA, Bernasconi S, *et al.*** The infant gut

465 microbiome as a microbial organ influencing host well-being. *Ital J Pediatr*
466 2020;46:16.

2. **Arboleya S, Watkins C, Stanton C, Ross RP**. Gut bifidobacteria populations in
    human health and aging. *Front Microbiol* 2016;7:1204.

3. **Derrien M, Turroni F, Ventura M, van Sinderen D**. Insights into endogenous
    Bifidobacterium species in the human gut microbiota during adulthood. *Trends Microbiol* 2022;30:940–947.

4. **Yassour M, Jason E, Hogstrom LJ, Arthur TD, Tripathi S, *et al.*** Strain-Level
    Analysis of Mother-to-Child Bacterial Transmission during the First Few Months
    of Life. *Cell Host Microbe* 2018;24:146–154.e4.

5. **Taft DH, Liu J, Maldonado-Gomez MX, Akre S, Huda MN, *et al.***
    Bifidobacterial dominance of the gut in early life and acquisition of antimicrobial
    resistance. *mSphere*;3. Epub ahead of print 26 September 2018. DOI:
    10.1128/mSphere.00441-18.

6. **Laforest-Lapointe I, Arrieta M-C**. Patterns of early-life gut microbial
    colonization during human immune development: An ecological perspective.
    Front Immunol 2017;8:788.

7. **Ennis D, Shmorak S, Jantscher-Krenn E, Yassour M**. Longitudinal
    quantification of Bifidobacterium longum subsp. infantis reveals late colonization
    in the infant gut independent of maternal milk HMO composition. *Nat Commun*
    2024;15:894.

8. **Xiao Y, Yang C, Yu L, Tian F, Wu Y, *et al.*** Human gut-derived B. longum
    subsp. longum strains protect against aging in a D-galactose-induced aging
    mouse model. *Microbiome* 2021;9:180.

9. **Zhang C, Yu Z, Zhao J, Zhang H, Zhai Q, *et al.*** Colonization and probiotic
    function of Bifidobacterium longum. *J Funct Foods* 2019;53:157–165.

10. **Rivière A, Selak M, Lantin D, Leroy F, De Vuyst L**. Bifidobacteria and
    butyrate-producing colon bacteria: Importance and strategies for their
    stimulation in the human gut. *Front Microbiol* 2016;7:979.

11. **Parte AC, Sardà Carbasse J, Meier-Kolthoff JP, Reimer LC, Göker M**. List of
    Prokaryotic names with Standing in Nomenclature (LPSN) moves to the DSMZ.
    Int J Syst Evol Microbiol 2020;70:5607–5612.

12. **Odamaki T, Bottacini F, Kato K, Mitsuyama E, Yoshida K, *et al.*** Genomic
    diversity and distribution of Bifidobacterium longum subsp. longum across the
    human lifespan. *Sci Rep* 2018;8:85.

13. **Mattarelli P, Bonaparte C, Pot B, Biavati B**. Proposal to reclassify the three
    biotypes of Bifidobacterium longum as three subspecies: Bifidobacterium
    longum subsp. longum subsp. nov., Bifidobacterium longum subsp. infantis
    comb. nov. and Bifidobacterium longum subsp. suis comb. nov. *Int J Syst Evol Microbiol* 2008;58:767–772.

14. **Modesto M, Ngom-Bru C, Scarafile D, Bruttin A, Pruvost S, *et al.***
Bifidobacterium longum subsp. iuvenis subsp. nov., a novel subspecies isolated
from the faeces of weaning infants. *Int J Syst Evol Microbiol*;73. Epub ahead of
print October 2023. DOI: 10.1099/ijsem.0.006013.

15. **Vatanen T, Ang QY, Siegwald L, Sarker SA, Le Roy CI, *et al.*** A distinct clade
of Bifidobacterium longum in the gut of Bangladeshi children thrives during
weaning. *Cell* 2022;185:4280–4297.e12.

16. **Yanokura E, Oki K, Makino H, Modesto M, Pot B, *et al.*** Subspeciation of
Bifidobacterium longum by multilocus approaches and amplified fragment length
polymorphism: Description of B. longum subsp. suillum subsp. nov., isolated
from the faeces of piglets. *Syst Appl Microbiol* 2015;38:305–314.

17. **Sela DA, Garrido D, Lerno L, Wu S, Tan K, *et al.*** Bifidobacterium longum
subsp. infantis ATCC 15697 α-fucosidases are active on fucosylated human
milk oligosaccharides. *Appl Environ Microbiol* 2012;78:795–803.

18. **Kelly SM, Munoz-Munoz J, van Sinderen D**. Plant glycan metabolism by
bifidobacteria. *Front Microbiol* 2021;12:609418.

19. **Young KD**. The selective value of bacterial shape. *Microbiol Mol Biol Rev*
2006;70:660–703.

20. **Tissier G Carre H, Naud C**. *Recherches sur la Flore Intestinale des*
Nourrissons (e'tat Normal et Pathologique). Paris; 1900.

21. **Bauer H, Sigarlakie E, Faure JC**. Scanning and transmission electron
microscopy of three strains of Bifidobacterium. *Can J Microbiol* 1975;21:1305–
1316.

22. **Husain I, Poupard JA, Norris RF**. Influence of nutrition on the morphology of a
strain of Bifidobacterium bifidum. *J Bacteriol* 1972;111:841–844.

23. **Dhanashree, Rajashekharan S, Krishnaswamy B, Kammara R**. Bifid Shape
Is Intrinsic to Bifidobacterium adolescentis. *Front Microbiol* 2017;8:478.

24. **Schleifer KH**. Classification of Bacteria and Archaea: past, present and future.
Syst Appl Microbiol 2009;32:533–542.

25. **Prakash O, Verma M, Sharma P, Kumar M, Kumari K, *et al.*** Polyphasic
approach of bacterial classification - An overview of recent advances. *Indian J*
Microbiol 2007;47:98–108.

26. **Kiepas AB, Hoskisson PA, Pritchard L**. 16S rRNA phylogeny and clustering is
not a reliable proxy for genome-based taxonomy in Streptomyces. *Microb*
Genom;10. Epub ahead of print September 2024. DOI:
10.1099/mgen.0.001287.

27. **Vandamme P, Pot B, Gillis M, de Vos P, Kersters K, *et al.*** Polyphasic
taxonomy, a consensus approach to bacterial systematics. *Microbiol Rev*
1996;60:407–438.

28. **Riesco R, Trujillo ME**. Update on the proposed minimal standards for the use
of genome data for the taxonomy of prokaryotes. *Int J Syst Evol Microbiol*;74.
Epub ahead of print March 2024. DOI: 10.1099/ijsem.0.006300.

29. **Tett A, Huang KD, Asnicar F, Fehlner-Peach H, Pasolli E, *et al.*** The
Prevotella copri complex comprises four distinct clades underrepresented in
Westernized populations. *Cell Host Microbe* 2019;26:666–679.e7.

30. **Blanco-Míguez A, Gálvez EJC, Pasolli E, De Filippis F, Amend L, *et al.***
Extension of the Segatella copri complex to 13 species with distinct large
extrachromosomal elements and associations with host conditions. *Cell Host*
Microbe 2023;31:1804–1819.e9.

31. **Versmessen N, Mispelaere M, Vandekerckhove M, Hermans C, Boelens J,**
et al. Average Nucleotide Identity and digital DNA-DNA Hybridization analysis
following PromethION nanopore-based whole genome sequencing allows for
accurate prokaryotic typing. *Diagnostics (Basel)* 2024;14:1800.

32. **Shao Y, Wang S, Gichuki B, Stares MD, Rozday T, *et al.*** Genomic atlas of
early life bifidobacteria for rational development of next-generation probiotics for
infant health. *Social Science Research Network*. Epub ahead of print 2025. DOI:
10.2139/ssrn.5095902.

33. **Chklovski A, Parks DH, Woodcroft BJ, Tyson GW**. CheckM2: a rapid,
scalable and accurate tool for assessing microbial genome quality using
machine learning. *Nat Methods* 2023;20:1203–1212.

34. **Díaz R, Torres-Miranda A, Orellana G, Garrido D**. Comparative Genomic
Analysis of Novel Bifidobacterium longum subsp. longum Strains Reveals
Functional Divergence in the Human Gut Microbiota. *Microorganisms*;9. Epub
ahead of print 8 September 2021. DOI: 10.3390/microorganisms9091906.

35. **Shao Y, Forster SC, Tsaliki E, Vervier K, Strang A, *et al.*** Stunted microbiota
and opportunistic pathogen colonization in caesarean-section birth. *Nature*
2019;574:117–121.

36. **Shao Y, Garcia-Mauriño C, Clare S, Dawson NJR, Mu A, *et al.*** Primary
succession of Bifidobacteria drives pathogen resistance in neonatal microbiota
assembly. *Nat Microbiol* 2024;9:2570–2582.

37. **Feehily C, O'Neill IJ, Walsh CJ, Moore RL, Killeen SL, *et al.*** Detailed
mapping of Bifidobacterium strain transmission from mother to infant via a dual
culture-based and metagenomic approach. *Nat Commun* 2023;14:3015.

38. **Gacesa R, Kurilshikov A, Vich Vila A, Sinha T, Klaassen MAY, *et al.***
Environmental factors shaping the gut microbiome in a Dutch population. *Nature*
2022;604:732–739.

39. **Sinha T, Brushett S, Fernández-Pato A, Garmaeva S, Andreu-Sánchez S, *et***
al. Pregnancy and early life gut microbiome: Influencing factors and health
implications. *Research Square*. Epub ahead of print 31 October 2024. DOI:
10.21203/rs.3.rs-5334252/v1.

40. **Warmink-Perdijk WDB, Peters LL, Tigchelaar EF, Dekens JAM,**
Jankipersadsing SA, *et al.* Lifelines NEXT: a prospective birth cohort adding
the next generation to the three-generation Lifelines cohort study. *Eur J*
Epidemiol 2020;35:157–168.

41. **Duncan SH, Louis P, Flint HJ.** Lactate-utilizing bacteria, isolated from human
feces, that produce butyrate as a major fermentation product. *Appl Environ*
Microbiol 2004;70:5810–5817.

42. **Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A**. Using
SPAdes DE Novo Assembler. *Curr Protoc Bioinformatics* 2020;70:e102.

43. **Antipov D, Korobeynikov A, McLean JS, Pevzner PA**. hybridSPAdes: an
algorithm for hybrid assembly of short and long reads. *Bioinformatics*
2016;32:1009–1015.

44. **Schwengers O, Jelonek L, Dieckmann MA, Beyvers S, Blom J,** *et al.* Bakta:
rapid and standardized annotation of bacterial genomes via alignment-free
sequence identification. *Microb Genom*;7. Epub ahead of print November 2021.
DOI: 10.1099/mgen.0.000685.

45. **Chan PP, Lin BY, Mak AJ, Lowe TM**. tRNAscan-SE 2.0: improved detection
and functional classification of transfer RNA genes. *Nucleic Acids Res*
2021;49:9077–9096.

46. **Hyatt D, Chen G-L, Locascio PF, Land ML, Larimer FW,** *et al.* Prodigal:
prokaryotic gene recognition and translation initiation site identification. *BMC*
Bioinformatics 2010;11:119.

47. **Larralde M**. Pyrodigal: Python bindings and interface to Prodigal, an efficient
method for gene prediction in prokaryotes. *J Open Source Softw* 2022;7:4296.

48. **Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S**. High
throughput ANI analysis of 90K prokaryotic genomes reveals clear species
boundaries. *Nat Commun* 2018;9:5114.

49. **Seemann T**. Prokka: rapid prokaryotic genome annotation. *Bioinformatics*
2014;30:2068–2069.

50. **Stajich JE, Block D, Boulez K, Brenner SE, Chervitz SA,** *et al.* The Bioperl
toolkit: Perl modules for the life sciences. *Genome Res* 2002;12:1611–1618.

51. **Tange O**. GNU Parallel - The Command-Line Power Tool, ;login. *The USENIX*
Magazine, Feb 2011;42–47.

52. **Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J,** *et al.*
BLAST+: architecture and applications. *BMC Bioinformatics* 2009;10:421.

53. **Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S,** *et al.* Roary: rapid
large-scale prokaryote pan genome analysis. *Bioinformatics* 2015;31:3691–
3693.

54. **Seemann T**. *barrnap: :microscope: Bacterial ribosomal RNA predictor.* Github.
https://github.com/tseemann/barrnap (accessed 10 June 2025).

55. **Sievers F, Higgins DG**. Clustal Omega for making accurate alignments of
many protein sequences. *Protein Sci* 2018;27:135–145.

56. **Stamatakis A**. RAxML version 8: a tool for phylogenetic analysis and post-
analysis of large phylogenies. *Bioinformatics* 2014;30:1312–1313.

57. **Li H**. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics*
2018;34:3094–3100.

58. **Hackl T, Ankenbrand M, van Adrichem B, Wilkins D, Haslinger K**.
gggenomes: effective and versatile visualizations for comparative genomics.
arXiv. Epub ahead of print 2024. DOI: 10.48550/arXiv.2411.13556.

59. **Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, *et al.***
Integrative genomics viewer. *Nat Biotechnol* 2011;29:24–26.

60. **Söding J, Biegert A, Lupas AN**. The HHpred interactive server for protein
homology detection and structure prediction. *Nucleic Acids Res* 2005;33:W244–
8.

61. **Blin K, Shaw S, Vader L, Szenei J, Reitz ZL, *et al.*** antiSMASH 8.0: extended
gene cluster detection capabilities and analyses of chemistry, enzymology, and
regulation. *Nucleic Acids Res*. Epub ahead of print 25 April 2025. DOI:
10.1093/nar/gkaf334.

62. **Meier-Kolthoff JP, Carbasse JS, Peinado-Olarte RL, Göker M**. TYGS and
LPSN: a database tandem for fast and reliable genome-based classification and
nomenclature of prokaryotes. *Nucleic Acids Res* 2022;50:D801–D807.

63. **Meier-Kolthoff JP, Auch AF, Klenk H-P, Göker M**. Genome sequence-based
species delimitation with confidence intervals and improved distance functions.
BMC Bioinformatics 2013;14:60.

64. **Suzek BE, Wang Y, Huang H, McGarvey PB, Wu CH, *et al.*** UniRef clusters: a
comprehensive and scalable alternative for improving sequence similarity
searches. *Bioinformatics* 2015;31:926–932.

65. **UniProt Consortium**. UniProt: The universal protein knowledgebase in 2025.
Nucleic Acids Res 2025;53:D609–D617.

66. **Beghini F, McIver LJ, Blanco-Míguez A, Dubois L, Asnicar F, *et al.***
Integrating taxonomic, functional, and strain-level profiling of diverse microbial
communities with bioBakery 3. *Elife*;10. Epub ahead of print 4 May 2021. DOI:
10.7554/eLife.65088.

67. **Buchfink B, Reuter K, Drost H-G**. Sensitive protein alignments at tree-of-life
scale using DIAMOND. *Nat Methods* 2021;18:366–368.

68. **Wickham H, François R, Henry L, Müller K, Vaughan D**. dplyr: A Grammar of

661   Data Manipulation.

662 69. **Wickham H, Vaughan D, Girlich M**. tidyr: Tidy Messy Data.

663 70. **Wickham H**. stringr: Simple, Consistent Wrappers for Common String
664   Operations. https://CRAN.R-project.org/package=stringr (2023).

665 71. **Wickham H, Averick M, Bryan J, Chang W, McGowan LD, *et al.*** Welcome to
666   the tidyverse. *Journal of Open Source Software* 2019;4:1686.

667 72. **Wickham H**. Data Analysis. In: Wickham H (editor). *ggplot2: Elegant Graphics*
668   *for Data Analysis*. Cham: Springer International Publishing; 2016. pp. 189–201.

669 73. **Pedersen TL**. ggforce: Accelerating 'ggplot2'.

670 74. **Neuwirth E**. RColorBrewer: ColorBrewer Palettes. https://CRAN.R-
671   project.org/package=RColorBrewer (2022).

672 75. **Kolde R**. pheatmap: Pretty Heatmaps. https://CRAN.R-
673   project.org/package=pheatmap (2019).

674 76. **Paradis E, Schliep K**. ape 5.0: an environment for modern phylogenetics and
675   evolutionary analyses in R. *Bioinformatics* 2019;35:526–528.

676 77. **Xu S, Li L, Luo X, Chen M, Tang W, *et al.*** *Ggtree* : A serialized data object for
677   visualization of a phylogenetic tree and annotation data. *Imeta*;1. Epub ahead of
678   print December 2022. DOI: 10.1002/imt2.56.

679 78. **Díaz R, Garrido D**. Screening competition and cross-feeding interactions during
680   utilization of human milk oligosaccharides by gut microbes. *Microbiome Res*
681   *Rep* 2024;3:12.

682 79. **Andreu-Sánchez S, Blanco-Míguez A, Wang D, Golzato D, Manghi P, *et al.***
683   Global genetic diversity of human gut microbiome species is related to
684   geographic location and host health. *Cell*;0. Epub ahead of print 30 April 2025.
685   DOI: 10.1016/j.cell.2025.04.014.

686 80. **Meier-Kolthoff JP, Hahnke RL, Petersen J, Scheuner C, Michael V, *et al.***
687   Complete genome sequence of DSM 30083(T), the type strain (U5/41(T)) of
688   Escherichia coli, and a proposal for delineating subspecies in microbial
689   taxonomy. *Stand Genomic Sci* 2014;9:2.

690 81. **Tarracchini C, Milani C, Lugli GA, Mancabelli L, Fontana F, *et al.***
691   Phylogenomic disentangling of the Bifidobacterium longum subsp. infantis
692   taxon. *Microb Genom*;7. Epub ahead of print July 2021. DOI:
693   10.1099/mgen.0.000609.

694 82. **Sela DA, Chapman J, Adeuya A, Kim JH, Chen F, *et al.*** The genome
695   sequence of Bifidobacterium longum subsp. infantis reveals adaptations for milk
696   utilization within the infant microbiome. *Proc Natl Acad Sci U S A*
697   2008;105:18964–18969.

83. **Duranti S, Lugli GA, Mancabelli L, Armanini F, Turroni F, *et al.*** Maternal
inheritance of bifidobacterial communities and bifidophages in infants through
vertical transmission. *Microbiome* 2017;5:66.

84. **Freitas AC, Hill JE**. Quantification, isolation and characterization of
Bifidobacterium from the vaginal microbiomes of reproductive aged women.
Anaerobe 2017;47:145–156.

85. **O'Connell Motherway M, Fitzgerald GF, Neirynck S, Ryan S, Steidler L, *et***
al. Characterization of ApuB, an extracellular type II amylopullulanase from
Bifidobacterium breve UCC2003. *Appl Environ Microbiol* 2008;74:6271–6279.

86. **Ryan SM, Fitzgerald GF, van Sinderen D**. Screening for and identification of
starch-, amylopectin-, and pullulan-degrading activities in bifidobacterial strains.
Appl Environ Microbiol 2006;72:5289–5296.

87. **Millar ME, Abele M, Harris HC, Koev TT, Telatin A, *et al.*** Novel amylase
genes enable utilisation of resistant starch by bifidobacteria relevant to early-life
microbiome development. *Microbiology*.
https://www.biorxiv.org/content/10.1101/2024.10.09.617373v1.full (2024).

88. **Rhim SL, Park MS, Ji GE**. Expression and secretion of Bifidobacterium
adolescentis amylase by Bifidobacterium longum. *Biotechnol Lett* 2006;28:163–
168.

89. **Sabath N, Ferrada E, Barve A, Wagner A**. Growth temperature and genome
size in bacteria are negatively correlated, suggesting genomic streamlining
during thermal adaptation. *Genome Biol Evol* 2013;5:966–977.

90. **Mueller RL**. Genome biology and the evolution of cell-size diversity. *Cold Spring*
Harb Perspect Biol 2015;7:a019125.

91. **Shiver AL, Sun J, Culver R, Violette A, Wynter C, *et al.*** Genome-scale
resources in the infant gut symbiont Bifidobacterium breve reveal genetic
determinants of colonization and host-microbe interactions. *Cell*
2025;188:2003–2021.e19.

92. **Liu S, Ren F, Zhao L, Jiang L, Hao Y, *et al.*** Starch and starch hydrolysates
are favorable carbon sources for bifidobacteria in the human gut. *BMC Microbiol*
2015;15:54.

93. **Ferraris RP, Choe J-Y, Patel CR**. Intestinal absorption of fructose. *Annu Rev*
Nutr 2018;38:41–67.

94. **Mokhtari P, Holzhausen EA, Chalifour BN, Schmidt KA, Babaei M, *et al.***
Associations between dietary sugar and fiber with infant gut microbiome
colonization at 6 mo of age. *J Nutr* 2024;154:152–162.

95. **Ramne S, Brunkwall L, Ericson U, Gray N, Kuhnle GGC, *et al.*** Gut
microbiota composition in relation to intake of added sugar, sugar-sweetened
beverages and artificially sweetened beverages in the Malmö Offspring Study.
Eur J Nutr 2021;60:2087–2097.

96. **van Dongen KCW, Ioannou A, Wesseling S, Beekmann K, Belzer C**.
Differences in gut microbial fructoselysine degradation activity between breast-
fed and formula-fed infants. *FEMS Microbiol Ecol* 2022;99:fiac145.

741 Figure captions

Figure 1: Analysis of *Bifidobacterium longum* genomes reveals four distinct clades.
(A) Average nucleotide identity (ANI) distances between all 435 B. longum genomes
analyzed in this study. Genome annotation data is shown for genomes with available public
annotations. *BL. nexti* genomes are highlighted in orange. **(B)** Median pairwise average
nucleotide identity (ANI) within and between the four *B. longum* subspecies. **(C)**
Phylogenetic tree of 24 selected genomes across subspecies constructed from the core
genome alignment. **(D)** Dendrogram of selected genomes generated from digital DNA-DNA
hybridization (dDDH) distances among 24 selected *BL.* genomes.

Figure 2: Genomic synteny analysis between type strains of *Bifidobacterium longum*
subspecies. Each line represents a genome, when multiple lines are present each one is a
single contig. Green shading between genomes denote conserved collinear regions
identified via sequence alignment. On the *BL. nexti* genome, *BL. nexti*-specific genes are
highlighted in orange.

Figure 3: Unique genomic features of *BL. nexti*. (A) Presence/absence heatmap of
accessory genes in the *B. longum* pangenome. Genomes are annotated by subspecies
where available (from NCBI or JGI IMG/M). **(B)** Genomic regions containing clusters of ≥4
BL. nexti-specific genes are shown in brown; non-*BL. nexti*-specific genes are shown with
transparent shading. Gene annotations appear above/below each feature.

Figure 4. Morphology and phenotypic characteristics of *Bifidobacterium longum*
strains. (A) Representative microscopic images of *BL. infantis*, *BL. longum*, and three *BL.*
nexti strains during the logarithmic growth phase. Cells are stained to visualize the cell
membrane and DNA. Scale bar, 2.5 μm. **(B)** Distribution of bacterial cell lengths (μm) for
each strain. Boxes indicate the interquartile range (25th–75th percentiles), with the median
marked; whiskers extend to 1.5 times the interquartile range. Colors denote subspecies. **(C)**
Growth rates of *BL. infantis*, *BL. longum*, and three *BL. nexti* strains on various
carbohydrates, measured by OD600. OD600 values are categorized as no growth (0-0.2),
low (0.2–0.6), medium (0.6–1), and high (>1).

Table 1. Metadata for *BL. nexti* strains and genomes. This table summarizes key

metadata for the nine *BL. nexti* genomes analyzed in this study, including isolation source of

the strains, geographic origin, and relevant genomic features.

Supplementary Figure 1: (A) Dendrogram of *B. longum* genomes with subspecies

annotations (172 genomes), constructed using presence/absence patterns of accessory

genes in the *B. longum* pangenome. **(B)** Heatmap showing digital DNA–DNA hybridization

(dDDH) values among selected genomes. **(C)** Phylogenetic tree of selected genomes based

on the 16S rRNA gene nucleotide sequences, with *B. breve* used as the outgroup.

Subspecies clustering is not as clear in this case due to lack of resolution in the 16s rRNA

gene sequence.

Supplementary Figure 2: Histogram showing the genome lengths (in base pairs) of all

analyzed genomes.

Supplementary Table 1: List of all genomes used in the study, including their sources and

subspecies annotations when publicly available (from NCBI or JGI IMG/M). Cohorts included

are: Baby Biome Study (BBS)[35, 36], Diaz et al.[34], Dutch microbiome project (DMP)[38],

Vatanen et al. 2022[15], MicrobeMom[37], Lifelines NEXT (LLNEXT)[39, 40].

Figure 1

Figure 2

Figure 3

Figure 4

Supplementary Figure 1

Supplementary Figure 2