Title: Heterosynaptic interactions between dorsal and ventral hippocampus in individual medium

spiny neurons of the nucleus accumbens ventromedial shell

Abbreviated title: Interactions between Hipp inputs in the NAc

Authors: Ashley E. Copenhaver[1], Sydnee Vance[1], Sarah A. Snider[1], Kaela Befano[1], Tara A.

LeGates[1,2]

[1] Department of Biological Sciences, University of Maryland, Baltimore County (UMBC)

[2] Department of Physiology and Pharmacology, University of Maryland School of Medicine

Corresponding author:

Tara LeGates, Ph.D.

1000 Hilltop Circle, Interdisciplinary Life Sciences Building 315

Baltimore, MD 21250

tlegates@umbc.edu

Number of pages:

Number of figures: 5 + 1 supplemental figure

Number of words, abstract: 242

Number of words, introduction: 649

Number of words, discussion: 1459

Conflict of interest statement: The authors declare no competing financial interests.

Acknowledgments: We would like to thank Dr. Tagide deCarvalho for her support with microscopy

and image analysis, and Dr. Ramesh Chandra for his help with viruses. We would also like to

thank Jennifer Pham for her help with injections and Dr. Brian Mathur for his helpful suggestions

on this project. This work was supported by NSF-IOS2402645, T32GM144876-02,

T34GM136497, and startup funds provided by UMBC.

ABSTRACT

Establishing learned associations between rewarding stimuli and the context under which those rewards are encountered is critical for survival. Hippocampal input to the nucleus accumbens (NAc) is a key connection involved in integrating environmental information and reward processing to facilitate goal-directed behaviors. This connection consists of two independent pathways originating from the dorsal (dHipp) or ventral (vHipp) hippocampus, which have previously been considered functionally and anatomically distinct. Here, we show overlap in dHipp and vHipp terminal fields in the NAc, which led us to reconsider this view and raise new questions regarding the potential interactions between dHipp and vHipp pathways in the NAc. Using optogenetics, electrophysiology, and transsynaptic labeling in adult male and female mice, we investigated anatomical and functional convergence of dHipp and vHipp in the NAc. We identified a subpopulation of dually innervated cells in the NAc medial shell where dHipp and vHipp inputs are located near one another along dendritic branches. We independently manipulated dHipp and vHipp inputs via two-color optogenetic manipulation during whole-cell electrophysiology recordings to confirm functional dual innervation of individual neurons and revealed heterosynaptic interactions between the two pathways. Altogether, these results demonstrate that dHipp and vHipp dually innervate a subset of neurons in the NAc, suggesting integration of these inputs at the level of individual neurons. Exploring the physiological and behavioral implications of this convergence will offer new insights into how individual neurons incorporate information from distinct inputs and how this integration may shape learning.

SIGNIFICANCE STATEMENT

Forming associations between rewards and the circumstances under which they are experienced is vital for survival. Hipp input to the NAc is essential for associating rewards with their environmental context to effectively guide motivated behaviors. This connection consists of two separate pathways originating from dHipp and vHipp that have long been considered distinct.

Here, we reveal a subpopulation of neurons in the NAc shell innervated by both Hipp subregions

as well as heterosynaptic interactions that occur between dHipp and vHipp synapses. These

findings suggest that integration of distinct hippocampal information occurs at the single-neuron

level, providing a critical mechanism underlying learning and motivated behavior while also

opening new avenues for understanding how diverse contextual and reward signals shape

decision-making.

INTRODUCTION

The integration of diverse information across neural systems is crucial for driving complex

behaviors. The brain dynamically integrates information from multiple sources, enabling

organisms to react to stimuli and establish learned associations crucial for survival. Plasticity of

and interactions between synapses enables coordinated regulation across multiple inputs, critical

for orchestrating cell-wide functional responses (Bailey et al., 2000; Govindarajan et al., 2006;

Harvey & Svoboda, 2007; Redondo & Morris, 2011; Schuman & Madison, 1994). However, the

mechanisms underlying how convergent inputs interact to shape neuronal responses remain

unclear.

The nucleus accumbens (NAc) is a key brain region that integrates diverse information from

several brain areas to effectively mediate goal-directed behaviors (Francis & Lobo, 2017; French

& Totterdell, 2002; Gruber et al., 2009; Mannella et al., 2013; Marinescu & Labouesse, 2024;

Salimpoor et al., 2013; Soares-Cunha et al., 2020). Hippocampal (Hipp) input is a major source

of excitatory drive that influences NAc neuron activity (Brog et al., 1993; Calhoon & O'Donnell,

2013; Groenewegen et al., 1987; Kelley & Domesick, 1982; Li et al., 2018; O'Donnell & Grace,

1995; Phillipson & Griffiths, 1985; Trouche et al., 2019), and because of the Hipp's essential role

in contextual and spatial learning (Duarte-Guterman et al., 2015; Lamsa & Lau, 2019; Safari et

al., 2021; Savage, 2004; Squire, 1992), Hipp-NAc synapses are also key sites for integrating

rewards with related contextual and spatial information. Furthermore, the modulation of the

strength of Hipp-NAc synapses is a critical mediator of motivated behaviors and is necessary for

supporting contextual- and spatial-based reward learning (Huntley et al., 2020; Ibrahim et al.,

2024; Ito et al., 2008, 2008; LeGates et al., 2018; Patterson et al., 2025, p. 202; Pennartz et al.,

2011; Sjulson et al., 2018; Trouche et al., 2019; F.-C. Yang & Liang, 2014).

The connection between the Hipp and NAc consists of two independent pathways originating

from the dorsal and ventral Hipp (dHipp and vHipp), which are considered functionally and

anatomically distinct (Amaral & Witter, 1989; Bannerman et al., 2002; Fanselow & Dong, 2010;

Friedman et al., 2002; K. B. Kjelstrup et al., 2008; K. G. Kjelstrup et al., 2002; Roberts et al., 2007;

Strange et al., 2014). Notably, the NAc is one of the few regions that receives input from both the

dHipp and vHipp (Friedman et al., 2002; Kelley & Domesick, 1982; Phillipson & Griffiths, 1985;

Trouche et al., 2019). Although dHipp and vHipp are interconnected within the Hipp (Ishizuka et

al., 1990; Swanson & Cowan, 1977; Tao et al., 2021), interactions through connections with other

brain regions have been proposed as a mechanism linking cognitive and emotional processes

(Fanselow & Dong, 2010). In a multifaceted environment, integration of diverse information is

important for properly mediating complex behaviors, and the behavioral complexity governed by

these pathways warrants synaptic integration to support learning and memory. These ideas are

supported by recent findings demonstrating a key role for convergence of dHipp and vHipp input

to the amygdala in observational contextual fear memory (Terranova et al., 2022), yet the synaptic

basis of this convergence and whether it is also observed in the NAc remains unknown.

Using a combination of microscopy, electrophysiology, and optogenetics, we identified and

characterized a subpopulation of NAc medium spiny neurons (MSNs) that are functionally

innervated by both the dHipp and vHipp. dHipp- and vHipp-MSN synapses on these neurons have

similar basal synaptic properties, and these synapses are close enough to engage in

heterosynaptic interactions likely via presynaptic mechanisms. These findings suggest that

convergent Hipp inputs interact at the synaptic level, which may contribute to the integration of

spatial and contextual information to modulate motivated behaviors. Overall, understanding the

synaptic integration of dHipp and vHipp inputs in the NAc not only enhances our knowledge of

reward-related behaviors, but also offers insight into the fundamental task of neuronal processing

and provides a foundation for developing targeted interventions for disorders where these

mechanisms are dysregulated.

METHODS

Animals

Male and female C57BL/6J, Drd1a-tdTomato, Ai14, and Ai195 mice were bred in-house. Mice

were housed with same-sex cage mates in a temperature- and humidity-controlled environment

under a 12:12 light cycle (lights on at 07:00), with food available *ad libitum*. We did not track

estrous cycle in females. All experiments were performed in accordance with the regulations set

forth by the Institutional Animal Care and Use Committee at the University of Maryland, Baltimore

County.

Stereotaxic surgeries

Mice (6-8 weeks old) were anesthetized with 3% isoflurane and underwent stereotaxic

surgery. A Hamilton syringe was used to inject adeno-associated viruses (AAV) into the dHipp (in

millimeters: from bregma, anterior-posterior (AP), -3.7; medial-lateral (ML), ±2.4; dorsal-ventral

(DV), -2.8), vHipp (from bregma, AP, -3.7; ML, ±3.0; DV, -4.8), and NAc (in millimeters: from

bregma, anterior-posterior (AP), +1.8; medial-lateral (ML), ±0.6; dorsal-ventral (DV), -4.7). Viruses

were injected bilaterally and were infused at a rate of 0.1uL per minute. The injection needle was

left in place for 10 minutes following the infusion. After surgery, mice were placed in their home

cage on a heating pad for recovery. Carprofen (5mg/kg) was subcutaneously injected daily for 3

days after surgery. Mice recovered for 5-7 weeks for viral expression before anatomical or

electrophysiological experiments. Viral constructs pAAV-hSyn-EGFP (Addgene #50465-AAV5)

and pAAV-hSyn-mCherry (Addgene #114472-AAV5) were used to visualize dHipp and vHipp

axon terminal fields. Anatomical labeling of dually innervated neurons involved the use of

pENN.AAV.hSyn.Cre.WPRE.hGH (AAV1, 1013 titer, UMB Viral Vector Core), pAAV-EF1a-Flpo

(AAV1, 1013 titer, UMB Viral Vector Core), and pAAV-hSyn-DIO-EGFP (Addgene #50457-AAV2).

Specifically, pENN.AAV.hSyn.Cre.WPRE.hGH and pAAV-EF1a-Flpo were injected into the dHipp

or vHipp of Ai195 mice for selective labeling of dually innervated neurons.

pENN.AAV.hSyn.Cre.WPRE.hGH and pAAV-EF1a-Flpo were injected into the dHipp or vHipp

and pAAV-hSyn-DIO-EGFP was injected in the NAc of Ai14 mice to label neurons single and

dually innervated neurons. For optogenetic experiments, viral vectors pAAV-Syn-ChrimsonR-tdT

(Addgene ID: 59171-AAV5) and either pAAV-Syn-Chronos-GFP (Addgene ID: 59170-AAV5) or

pAAV-hSyn-hChR2(H134R)-EYFP (Addgene ID: 26973-AAV5) were used. For all experiments,

injection sites were verified by fluorescence microscopy, and brains with off target expression

were removed from the experiment.

Visualization of axon terminal fields and dually innervated neurons

Mice expressing viruses for anatomical tracing experiments were first anesthetized with

isoflurane and then perfused with 1X phosphate buffer solution (PBS), followed by 4%

paraformaldehyde (PFA). Brains were removed and post-fixed in 4% PFA overnight at 4ºC and

were then transferred to 1X PBS and stored at 4ºC for up to 3 weeks. Brains were sectioned

(100µm) using a VT1000S Leica Microsystems vibratome. Slices were then mounted with

Vectashield Antifade Mounting Medium on slides for imaging.

Images were captured with either a Zeiss LSM 900 Confocal microscope or Zeiss Axio

Zoom.V16, both equipped with Zeiss software. To prevent cross talk among channels, images

were obtained using sequential scanning to detect multiple fluorophores. To reduce background

noise and graininess of images, 2x frame-averaging was used and gain was kept at or below 700.

The laser powers used were: EGFP <15% and mCherry or tdTomato <15%. For imaging of

dendritic segments, z-stacks with intervals of 0.2µm were acquired and further analyzed using

Imaris.

To quantify individual and dually innervated neurons, images of dorsomedial NAcSh,

ventromedial NacSh, ventrolateral NacSh, and ventromedial NAc core were captured on Zeiss

LSM 900 Confocal microscope at 40x magnification, equipped with Zeiss software. Images were

captured throughout the dorsomedial NAcSh, ventromedial NacSh, ventrolateral NacSh, and

ventromedial NAc core. Cells were then manually counted and recounted a day apart. Analysis

and counting were done by an experimenter blinded to conditions.

Acute brain slice preparation for electrophysiology

Acute coronal slices containing the nucleus accumbens or hippocampus were prepared for

whole-cell patch-clamp electrophysiology. Hipp-containing slices were kept to verify injection sites

before proceeding with experiments in the NAc. Animals were deeply anesthetized with

isoflurane, decapitated, and brains were quickly dissected and submerged in ice-cold, bubbled

(carbogen: 95% O_2/5% CO_2) *N*-methyl-D-glucamine (NMDG) recovery solution containing the

following (in mM): 93 NMDG, 2.5 KCl, 1.2 NaH_2PO_4, 11 glucose, 25 $NaHCO_3$, 1.2 $MgCl_2$, and 2.4

$CaCl_2$, pH=7.3-7.4, osmolarity=300-310 mOsm. Using a vibratome (VT1000S, Leica

Microsystems), coronal slices (400 µm) were cut in cold, oxygenated NMDG. Slices were

transferred to 32-34°C NMDG for 7-12 minutes to recover and were then transferred to room-

temperature artificial cerebrospinal fluid (aCSF) containing the following (in mM): 120 NaCl, 3 KCl,

1.0 NaH_2PO_4, 20 glucose, 25 $NaHCO_3$, 1.5 $MgCl_2 \cdot 7H_2O$, and 2.5 $CaCl_2$, pH=7.3-7.4. Slices were

allowed to recover for 1-hour at room-temperature before beginning electrophysiological

recordings.

Whole-cell recordings

We performed whole-cell patch-clamp recordings using an Axopatch 200B amplifier (Axon

Instruments, Molecular Devices) and a Digidata 1550B digitizer (Axon Instruments). Slices were

placed in a submersion-type recording chamber and superfused with room-temperature aCSF

(flow rate 0.5-1mL/min). Patch pipettes (4-8MΩ) were made from borosilicate glass (World

Precision Instruments) using a Sutter Instruments P-97 model puller. Cells were visualized using

a 60x water immersion objective (Nikon Eclipse FN-1). All recordings were performed in voltage-

clamp conditions from MSNs in the NAc ventromedial shell.

Hipp axon terminals expressing Chrimson, Chronos, or Channelrhodopsin were

optogenetically stimulated to record evoked excitatory postsynaptic currents (EPSC). Unless

otherwise noted, alternating stimulation was performed every 10s to record EPSCs evoked by

either dHipp or vHipp stimulation. For most experiments shown, patch pipettes were filled with a

solution containing 130 mM K-gluconate, 5 mM KCl, 2 mM $MgCl_6$-H_2O, 10 mM HEPES, 4 mM

Mg-ATP, 0.3 mM Na_2-GTP, 10 mM Na_2-phosphocreatine, and 1 mM EGTA; pH=7.3-7.4;

osmolarity=285-295mOsm. High-frequency stimulation of either dHipp or vHipp afferents was

performed to induce long-term potentiation of synapses and examine possible long-lasting

heterosynaptic plasticity. For experiments determining I-V relationship and AMPA:NMDA receptor

ratios, the patch pipette solution was composed of 135 mM CsCl, 2 mM $MgCl_6$-H_2O, 10 mM

HEPES, 4 mM Mg-ATP, 0.3 mM Na_2-GTP, 10 mM Na_2-phosphocreatine, 1 mM EGTA, 5 mM

QX-314 bromide, and 100 µM spermine; pH=7.3-7.4; osmolarity=285-295mOsm. EPSCs were

collected from holding potentials ranging from -80mV to +40mV to create an I-V curve. To

calculate AMPA:NMDA receptor ratios, the AMPA component was defined by the peak amplitude

at -70mV while the NMDA component was defined by the amplitude at +40mV at 50ms after

stimulation. We used the following exclusion criteria to eliminate unhealthy cells and unreliable

recordings: 1) We only proceeded with experiments on cells with series resistances <10MΩ, 2)

Cells were excluded if their series resistance changed by >20% (comparing the resistance at the

beginning and end of the experiment), 3) Cells in poor health or poor recording status were

excluded (i.e. cell partially or fully sealed up, a decrease in holding current >100pA that is

consistent with the cell dying, an increase in jitter post HFS, and/or an increase in response failure

rate to > 50%).

To minimize the likelihood of cross-contamination between Chrimson and Chronos/ChR2, we

conducted control experiments to identify light stimulation intensities that would not cause

activation of Chrimson by 470nm light. We found that using light intensities ≤0.5mW prevented

cross-contamination (Figure S1), aligning with other published studies (Christoffel et al., 2021;

Klapoetke et al., 2014; Xia et al., 2020).

Visualization of dHipp/vHipp terminals with dually innervated neurons

Using a small subset of slices from electrophysiological experiments, dually innervated MSNs

were patched and filled with CF633 dye (Biotium). Upon completion of electrophysiology

experiments, the 400μm-thick slices were fixed in 4% PFA for 4-8 hours at 4ºC. Slices were then

mounted in Invitrogen ProLong Glass Antifade Mountant (Thermo Fisher Scientific, #P36980) on

slides with spacers (400μm-thick) for imaging. Images were captured with a Zeiss LSM 900

Confocal microscope equipped with Zeiss software. To prevent cross talk among channels,

images were obtained using sequential scanning to detect multiple fluorophores. To reduce

background noise and graininess of images, 2x frame-averaging was used and gain was kept at

or below 700. The laser powers used were: EGFP <15%, mCherry or tdTomato <15%, CF633

<10%. For imaging of dendritic segments, z-stacks with intervals of 0.2μm were acquired and

further analyzed using Imaris.

Quantification, statistical analysis, and reproducibility

We found no statistically significant difference between males and females, so data from male

and females are plotted together. Both the number of cells and number of mice is reported for

each experiment. Plots represent the mean ± SEM. The sample size (n) per condition represents

the number of cells unless otherwise indicated in the figure caption. All statistical analyses were

performed using Graphpad Prism 9/10 software. Statistical significance (p<0.05) was assessed

using parametric or non-parametric versions of paired t tests, or one-way or two-way ANOVA with

multiple comparisons. For box plots, the line in the middle of the box is plotted at the median. The

box extends from the 25th to 75th percentiles. Whiskers represent minimum and maximum.

Figure schematics and experimental timelines were created with BioRender.com.

RESULTS

dHipp and vHipp axon terminal fields overlap in the NAc medial shell

Using viral-mediated labeling to visualize Hipp neurons and their terminal fields, we examined

innervation of the NAc by the posterior dHipp and vHipp. We injected an adenoassociated virus

(AAV) expressing eYFP into the dHipp and an AAV expressing mCherry into the vHipp (Figure

1a-b). After allowing 5-7 weeks for expression, brains were fixed, sectioned, and imaged using

confocal microscopy. We observed dHipp axon terminal fields in the NAc core and lateral shell

while the vHipp axon terminal fields were primarily found in the NAc medial shell, in line with

previous work (Figure 1c). However, we also observed strong overlap of dHipp and vHipp axon

terminal fields in the NAc ventromedial shell (Figure 1c-d), which aligns with the results presented

by Brog et al., 1993. 3-D reconstruction of z-stack images of these overlapping axon fields

revealed adjacent dHipp and vHipp axons (Figure 1d), highlighting the close proximity of dHipp

and vHipp afferents within the ventromedial shell.

Identification of NAc neurons dually innervated by the dHipp and vHipp

Given the apposition of dHipp and vHipp terminals, we sought to determine whether individual

neurons were innervated by both hippocampal regions. To do this, we utilized transsynaptic

expression of Cre and Flp recombinases to selectively target NAc neurons innervated by dHipp

and vHipp pathways (Figure 2a). We injected AAV1-hSyn-Cre into the dHipp and AAV1-Ef1a-

Flpo into the vHipp of Ai195 mice in which GcaMP6s is only expressed in the presence of Cre

and Flp. This approach allowed for targeted expression of GCaMP6s only in neurons that are

innervated by both dHipp and vHipp (Figure 2b). After allowing 5-7 weeks for expression, brains

were fixed, sectioned and imaged with confocal microscopy using GCaMP6s to visualize dually

innervated neurons. Similar to our observations of terminal field overlap, we observed GCaMP6s-

expressing cells in the NAc ventromedial shell (Figure 2c), suggesting dual innervation of NAc

neurons by both dHipp and vHipp. Furthermore, examination of GcaMP6s-expressing cells

revealed morphology similar to MSNs (Figure 2d), suggesting that primarily NAc MSNs are

targeted by both the dHipp and vHipp.

To gain better insight into the proportions of cells innervated by dHipp, vHipp, or both, we

used our transsynaptic strategy to differentially label cells innervated by dHipp and vHipp.

Specifically, we injected AAV1-hSyn-Cre into the dHipp and AAV1-Ef1a-Flpo into the vHipp of

Ai14 mice in which there is Cre-dependent expression of tdTomato. We also injected AAV9-Ef1a-

fDIO-eYFP into the NAc of these mice (Figure 2e). As a result, cells innervated by dHipp express

tdTomato, cells innervated by vHipp express eYFP, and those innervated by both will express

both reporters (Figure 3f-g). Quantification of cells in medial shell revealed approximately 13% of

cells were dually innervated. The remainder of cells were innervated by one Hipp subregion

(singly innervated). Of these, 26% cells were innervated by dHipp, 61% were innervated by vHipp

(Figure 3h).

To determine whether our anatomical results translated into functional convergence, we

explored whether MSNs in the NAc ventromedial shell were functionally innervated by the dHipp

and vHipp (AKA "dually innervated"). We focused on MSNs because they comprise 95% of the

NAc neuron population, align with the morphology of cells identified by our anatomical tracing

studies (Figure 2), and have been shown to receive functional input from either dHipp or vHipp in

separate studies. To independently stimulate dHipp and vHipp inputs, we used a dual wavelength

optogenetic strategy involving viral expression of Chronos and Chrimson, which are maximally

sensitive to different wavelengths of light (480nm and 660nm, respectively) (Klapoetke et al.,

2014). AAV2-Syn-Chronos-eYFP or AAV2-Syn-Chrimson-mCherry were injected into the dHipp

or vHipp. At least 5 weeks after surgery, coronal brain slices containing the NAc were prepared

for electrophysiology. Using whole-cell patch-clamp electrophysiology, we recorded from MSNs

in the NAc ventromedial shell and used alternating 473 and 635nm light to measure dHipp and

vHipp evoked responses in the MSNs. Keeping the light intensity below 1mW prevented cross-

contamination, allowing us to measure and independently manipulate dHipp and vHipp inputs

with 473 and 635nm light (Figure S1 and 3a). Using this strategy, we identified a subpopulation

of MSNs in the ventromedial shell that responded to both dHipp and vHipp stimulation,

demonstrating dual functional innervation of individual NAc MSNs by both dHipp and vHipp

(Figure 3a).

Synaptic properties and strengths influence how a neuron responds to and integrates

information from distinct inputs, leading us to examine and compare the synaptic profiles of dHipp-

MSN and vHipp-MSN synapses in dually innervated neurons. To examine excitatory synaptic

strength, we measured AMPAR:NMDAR ratios at dHipp-MSN and vHipp-MSN synapses and

found no difference, suggesting that the strength of dHipp- and vHipp-MSN synapses is similar

(Figure 3b). We then stimulated each pathway with paired pulses, alternating stimulation such

that each pathway was stimulated every 20s. Calculation of paired-pulse ratios (PPR) revealed a

significantly greater PPR at vHipp-MSN synapses as compared to dHipp-MSN synapses (Figure

3c), suggesting potential differences in release probability, though this may be driven by a

subpopulation of recorded neurons.

Calcium-permeable AMPA receptors (CP-AMPAR) have been identified at ventral subiculum-

NAc synapses (Boxer et al., 2023) and have been implicated in NAc plasticity and related

behaviors (Carr, 2020; Mameli et al., 2009; McCutcheon et al., 2011; Wolf & Tseng, 2012). Given

their role in plasticity and behavior, we examined whether CP-AMPARs were present at dHipp-

or vHipp-MSN synapses. Using their characteristic inward rectification and non-linear current-

voltage relationship (Cull-Candy et al., 2006; Liu & Zukin, 2007), we performed

electrophysiological experiments to identify the predominant population of AMPARs at dHipp and

vHipp-MSN synapses. Our results revealed that neither type of synapse contained a significant

population of CP-AMPARs (Figure 3d). Altogether, these results indicate that dHipp and vHipp

synapses in dually innervated NAc MSNs have similar basal synaptic properties.

We then evaluated the potential for synaptic or dendritic-level interactions between dHipp and

vHipp synapses by comparing rise times and the spatial locations of putative dHipp and vHipp

synapses. Our rise time analysis revealed similar rise times of dHipp- and vHipp-evoked

responses, suggesting that dHipp-MSN and vHipp-MSN synapses are located a similar distance

away from the cell body (Figure 3e). We filled a subset of cells electrophysiologically confirmed

to be dually innervated with dye (CF633) (Figure 4a), and slices were fixed and prepared for

confocal microscopy. We used confocal microscopy and image analysis to measure the spatial

location of potential dHipp and vHipp synapses along the dendrites of a dually innervated MSN

using the eYFP co-expressed with Chronos, the mCherry co-expressed with Chrimson, and the

CF633. From this analysis, we found several examples of dHipp and vHipp puncta that are

spatially close (<2um) along distal dendrites of dually innervated MSNs (Figure 4b). Three-

dimensional reconstruction of z-stack images revealed overlap of dHipp axons, vHipp axons, and

the MSN dendrite, providing further evidence for dHipp and vHipp synapses being spatially close

(Figure 4c). Altogether, these results demonstrate functional innervation of individual MSNs by

both dHipp and vHipp, where synapses may be close enough to each other to interact at a

synaptic or dendritic level.

Heterosynaptic interactions between dHipp and vHipp synapses in NAc MSNs

Synapses that are sufficiently close (<2um apart) can interact and influence each other's activity (Fu et al., 2012; Harvey & Svoboda, 2007; Murakoshi et al., 2011; Xia et al., 2020; Yuste, 2011). Though activity in dHipp and vHipp is not typically temporally coordinated, learning has been found to increase synchrony between dHipp and vHipp (Biane et al., 2023; Morici et al., 2025). Synchronous activity has been shown to induce plasticity through heterosynaptic interactions between adjacent synapses, raising the question of whether the spatial proximity of dHipp and vHipp synapses facilitates heterosynaptic interactions. To address this, we alternated 473nm and 635nm stimulation for 2 minutes to measure EPSCs evoked from independent stimulation of each Hipp pathway. We then synchronized 473nm and 635nm stimulations for 2 minutes to record an evoked EPSC from the simultaneous stimulation of both dHipp and vHipp inputs (Figure 5a,b). If there are no heterosynaptic interactions occurring, then the EPSC amplitudes from the alternating stimulation should be the linear sum of the EPSC amplitude evoked by simultaneous stimulation. Instead, our data show a supralinear summation of separately measured EPSCs (Figure 5b), indicating that heterosynaptic potentiation occurs between dHipp and vHipp synapses. We conducted the same experiment in cells that were not dually innervated and found no difference between EPSC amplitudes evoked by alternating and synchronous stimulation, demonstrating that this effect is specific to dually innervated neurons and not a nonspecific artifact resulting from cross-contamination (Figure 5c).

To better understand the site of synaptic interaction, we compared the failure rates in a separate experiment using minimal stimulation to evoke pathway-specific responses only 50% of the time so that we could appropriately assess an increase or decrease in failure rate during synchronous stimulation. We measured EPSCs evoked from alternating and synchronous stimulation of dHipp-MSN and vHipp-MSN synapses. We then calculated the theoretical probability of getting a response in both the dHipp- and vHipp-MSN pathways, P_{dual}, if they were

stimulated at the same time. Assuming the pathways are independent/non-interacting, we can

calculate this probability as the product of their individual success properties: $P_{dual}^{theoretical} =$

$P_{dHipp-MSN}^{success} \cdot P_{vHipp-MSN}^{success}$. We compared this theoretical probability to the observed success rate of

sync responses that were classified as dual-evoked responses, finding a significant difference in

the observed and theoretical values (Figure 5d) suggesting presynaptic mechanisms may

facilitate heterosynaptic interactions between dHipp- and vHipp-MSN synapses.

DISCUSSION

The experiments in this study identified a subpopulation of neurons within the NAc that are

dually innervated by dHipp and vHipp. These neurons are primarily located in the NAc

ventromedial shell with morphological characteristics that align with those of MSNs. We found

that dHipp-MSN and vHipp-MSN synapses in dually innervated neurons have similar basal

synaptic properties including comparable release probabilities, synaptic strength, and expression

of calcium-impermeable AMPARs. Examination of EPSCs evoked by individual or synchronous

stimulation of dHipp and vHipp revealed supralinear summation of EPSCs, suggesting that the

synchronous activation of dHipp and vHipp inputs induces short-term heterosynaptic potentiation

of dHipp-MSN and vHipp-MSN synapses likely due to presynaptic mechanisms. Altogether, these

results reveal mechanisms of dHipp and vHipp synaptic integration within the NAc medial shell,

which may be key to supporting the integration of cognitive and emotional processes needed to

effectively mediate goal-directed behaviors.

Innervation of the NAc by dHipp and vHipp

Our viral-mediated axon terminal field labeling strategy revealed overlapping dHipp and vHipp

innervation patterns within the NAc ventromedial shell. At first glance, this finding seems to

contradict previous reports that have suggested that the dHipp and vHipp preferentially innervate

distinct parts of the NAc. However, these reported innervation patterns appear to depend upon

the anterior/posterior coordinate used for Hipp viral injections. Most studies have targeted the

posterior vHipp and anterior dHipp, showing vHipp projections in the NAc core and medial shell,

with dHipp projections in the NAc core and lateral shell (Groenewegen et al., 1987; Ibrahim et al.,

2024; Li et al., 2018; Mulder et al., 1998; Scofield et al., 2016). But when the posterior dHipp was

targeted with an anterograde tracer, dHipp axon terminal field expression was observed in the

NAc ventromedial shell (Ibrahim et al., 2024). Furthermore, the injection of a retrograde tracer

into the NAc ventromedial shell showed that neurons located in the posterior dHipp and vHipp

were projecting to this area of the NAc (Groenewegen et al., 1987). Our findings align with the

latter two studies, as we injected into the posterior Hipp (A/P: -3.7mm) and found overlapping

dHipp and vHipp axon terminal fields and dually innervated neurons primarily in the NAc

ventromedial shell.

Using a transsynaptic viral-mediated labeling technique, we were able to visualize a small

subpopulation of dually innervated neurons that had morphological characteristics similar to

MSNs (Figure 2). Because the viruses used for this technique have low efficiency for

transsynaptic transfer (Zingg et al., 2017), and the labeling of a dually innervated neuron requires

successful transport of both Cre and Flipase virus payloads, it is likely that this anatomical labeling

approach underestimates the number of dually innervated neurons in the NAc. This is supported

by our electrophysiology experiments, as within a single slice, we were able to record from a

larger number of dually innervated neurons than were indicated from the anatomical tracing study.

Optimization of this transsynaptic approach would likely provide a better estimate of the number

of dually innervated neurons and could aid in identifying whether interneurons, which are sparsely

located in the NAc, can receive dual input from dHipp and vHipp.

dHipp and vHipp synapses have similar properties in dually innervated NAc neurons

We found that dHipp- and vHipp-MSN synapses contain primarily calcium-impermeable AMPARs and have similar excitatory synaptic strengths, as measured by AMPA:NMDA receptor ratios (Figure 3). These similarities suggest that, in basal conditions, both pathways may contribute equally to downstream circuit activity. Since the dHipp and vHipp convey different types of behavioral information, this may mean that Hipp-related contextual and sensory cues are equally capable of influencing motivated behaviors. While these similarities are present at baseline, several studies have highlighted changes that occur to dHipp-MSN and vHipp-MSN synapses in response to hormones, neuromodulators, cocaine, external stress, and experience (Floresco et al., 2001; LeGates et al., 2018; Peleg-Raibstein & Feldon, 2006; R. Yang & J. Mogenson, 1984; Sjulson et al., 2018; Williams et al., 2020).

Synaptic interactions between distinct Hipp inputs

The NAc is widely regarded as a critical integrator of several major excitatory inputs and several studies have examined how convergent inputs interact at the synaptic level within the NAc. These studies have revealed intricate interactions mediated by a combination of presynaptic and postsynaptic mechanisms which influence behavioral output (O'Donnell & Grace, 1995; Xia et al., 2020; F.-C. Yang & Liang, 2014; Yu et al., 2022). Further, vHipp input to the NAc can influence how NAc MSNs respond to other excitatory inputs (Calhoon & O'Donnell, 2013; O'Donnell & Grace, 1995), however it is unknown whether these vHipp inputs exert similar regulatory control over dHipp inputs to the NAc. Since the dHipp and vHipp are known to convey distinct types of behavioral information which are critical for driving NAc-mediated motivated behaviors (Ibrahim et al., 2024; Ito et al., 2008; LeGates et al., 2018; Muir et al., 2020; Patterson et al., 2025; Sjulson et al., 2018; Sosa et al., 2020; Trouche et al., 2019; Williams et al., 2020; A. K. Yang et al., 2020; F.-C. Yang & Liang, 2014), we were interested in examining whether dHipp and vHipp inputs are regulated at the synaptic level. Our results revealed that dHipp and vHipp

inputs form synaptic connections with individual MSNs within the NAc ventromedial shell, and

these synapses rapidly interact with one another to enhance MSN responses to both regions

(Figure 5). Interestingly, activity in dHipp and vHipp is not typically temporally coordinated,

however increased synchrony between dHipp and vHipp has been associated with learning(Biane

et al., 2023; Morici et al., 2025).The rapid response we observed may provide a basis by which

distinct information simultaneously conveyed by dHipp and vHipp can strengthen the NAc's

response to external stimuli important for establishing and maintaining learned associations.

Implications of integration that occurs at the synaptic level

Because of the Hipp's essential role in contextual and spatial learning across species and the

role of the NAc in reward processing (Fanselow, 2000; Nakazawa et al., 2003; Rudy et al., 2002;

Squire, 1992; Steele & Morris, 1999), the connection between the Hipp and NAc is crucial for

integrating reward with related contextual and spatial information (Britt et al., 2012; Floresco &

Phillips, 1999; Humphries & Prescott, 2010; Ito et al., 2008; Lansink et al., 2008, 2009; LeGates

et al., 2018; Pennartz et al., 2011; Sjulson et al., 2018). Functional imaging studies in humans

have revealed that greater Hipp-NAc connectivity is associated with positive reinforcement

learning (Davidow et al., 2016), increased reward seeking in adolescents (Huntley et al., 2020),

and greater positive affect in adults (Heller et al., 2020), implicating a strong link between the

strength of Hipp-NAc connectivity and reward-related behaviors. Similar findings have been

demonstrated in rodent models which examined Hipp-NAc connectivity using in vivo

electrophysiology during spatial and contextual-based reward tasks (Sjulson et al., 2018; Sosa et

al., 2020; Trouche et al., 2019). The causal relationship between Hipp-NAc connectivity and

reward behaviors has been established with lesion and optogenetic studies, demonstrating that

Hipp-NAc synapses are required for contextual- and spatial-based reward tasks (Floresco &

Phillips, 1999; Ito et al., 2008, p. 200; LeGates et al., 2018; Trouche et al., 2019; Zhou et al.,

2019). This includes our previously published work, which demonstrated that the strength of

ventral Hipp-NAc synapses is a critical mediator of reward-related behaviors (LeGates et al.,

2018).

While the dHipp and vHipp are interconnected within the hippocampus (Ishizuka et al., 1990;

Swanson & Cowan, 1977; Tao et al., 2021), their interactions in other areas of the brain have

been proposed as a mechanism linking cognitive and emotional processes (Fanselow & Dong,

2010) and may be crucial to establishing and retrieving memories. This has been demonstrated

in the amygdala, where dHipp and vHipp activate and reactive, respectively, a specific ensemble

of neurons to support contextual fear memory (Terranova et al., 2022). The NAc serves as another

key brain region that receives both dHipp and vHipp input, where we have revealed rapid synaptic

interactions that occur between these pathways. Since Hipp input uniquely exerts such a strong

influence over NAc neuron activity and information flow, it is likely these interactions encourage

integration of distinct information conveyed by the dHipp and vHipp, critically impacting behavioral

outputs.

Altogether, the results presented in this paper provide evidence for the integration of dHipp

and vHipp information within individual MSNs of the NAc ventromedial shell. The combination of

techniques enabling identification of similar synaptic profiles and rapid interactions occurring

between dHipp- and vHipp-MSN synapses provides a foundation for similar studies investigating

the integration of distinct inputs at the level of individual neurons. Ultimately, these findings

advance our understanding of how spatial and contextual information may contribute to reward-

related behaviors that are mediated by the NAc shell while also raising further questions as to

how each of these distinct pathways are regulated.

References

Amaral, D. G., & Witter, M. P. (1989). The three-dimensional organization of the hippocampal

    formation: A review of anatomical data. *Neuroscience*, *31*(3), 571–591.

    https://doi.org/10.1016/0306-4522(89)90424-7

Bailey, C. H., Giustetto, M., Huang, Y. Y., Hawkins, R. D., & Kandel, E. R. (2000). Is

    heterosynaptic modulation essential for stabilizing Hebbian plasticity and memory?

    Nature Reviews. Neuroscience, *1*(1), 11–20. https://doi.org/10.1038/35036191

Bannerman, D. M., Deacon, R. M. J., Offen, S., Friswell, J., Grubb, M., & Rawlins, J. N. P.

    (2002). Double dissociation of function within the hippocampus: Spatial memory and

    hyponeophagia. *Behavioral Neuroscience*, *116*(5), 884–901.

    https://doi.org/10.1037/0735-7044.116.5.884

Biane, J. S., Ladow, M. A., Stefanini, F., Boddu, S. P., Fan, A., Hassan, S., Dundar, N.,

    Apodaca-Montano, D. L., Zhou, L. Z., Fayner, V., Woods, N. I., & Kheirbek, M. A. (2023).

    Neural dynamics underlying associative learning in the dorsal and ventral hippocampus.

    Nature Neuroscience, *26*(5), 798–809. https://doi.org/10.1038/s41593-023-01296-6

Boxer, E. E., Kim, J., Dunn, B., & Aoto, J. (2023). Ventral Subiculum Inputs to Nucleus

    Accumbens Medial Shell Preferentially Innervate D2R Medium Spiny Neurons and

    Contain Calcium Permeable AMPARs. *The Journal of Neuroscience*, *43*(7), 1166–1177.

    https://doi.org/10.1523/JNEUROSCI.1907-22.2022

Britt, J. P., Benaliouad, F., McDevitt, R. A., Stuber, G. D., Wise, R. A., & Bonci, A. (2012).

    Synaptic and Behavioral Profile of Multiple Glutamatergic Inputs to the Nucleus

    Accumbens. *Neuron*, *76*(4), 790–803. https://doi.org/10.1016/j.neuron.2012.09.040

Brog, J. S., Salyapongse, A., Deutch, A. Y., & Zahm, D. S. (1993). The patterns of afferent

    innervation of the core and shell in the "Accumbens" part of the rat ventral striatum:

    Immunohistochemical detection of retrogradely transported fluoro-gold. *Journal of*

    Comparative Neurology, *338*(2), 255–278. https://doi.org/10.1002/cne.903380209

Calhoon, G. G., & O'Donnell, P. (2013). Closing the Gate in the Limbic Striatum: Prefrontal

      Suppression of Hippocampal and Thalamic Inputs. *Neuron*, *78*(1), 181–190.

      https://doi.org/10.1016/j.neuron.2013.01.032

Carr, K. D. (2020). Homeostatic regulation of reward via synaptic insertion of calcium-permeable

      AMPA receptors in nucleus accumbens. *Physiology & Behavior*, *219*, 112850.

      https://doi.org/10.1016/j.physbeh.2020.112850

Christoffel, D. J., Walsh, J. J., Hoerbelt, P., Heifets, B. D., Llorach, P., Lopez, R. C.,

      Ramakrishnan, C., Deisseroth, K., & Malenka, R. C. (2021). Selective filtering of

      excitatory inputs to nucleus accumbens by dopamine and serotonin. *Proceedings of the*

      National Academy of Sciences, *118*(24), e2106648118.

      https://doi.org/10.1073/pnas.2106648118

Cull-Candy, S., Kelly, L., & Farrant, M. (2006). Regulation of Ca2+-permeable AMPA receptors:

      Synaptic plasticity and beyond. *Current Opinion in Neurobiology*, *16*(3), 288–297.

      https://doi.org/10.1016/j.conb.2006.05.012

Davidow, J. Y., Foerde, K., Galván, A., & Shohamy, D. (2016). An Upside to Reward Sensitivity:

      The Hippocampus Supports Enhanced Reinforcement Learning in Adolescence.

      Neuron, *92*(1), 93–99. https://doi.org/10.1016/j.neuron.2016.08.031

Duarte-Guterman, P., Yagi, S., Chow, C., & Galea, L. A. M. (2015). Hippocampal learning,

      memory, and neurogenesis: Effects of sex and estrogens across the lifespan in adults.

      Hormones and Behavior, *74*, 37–52. https://doi.org/10.1016/j.yhbeh.2015.05.024

Fanselow, M. S. (2000). Contextual fear, gestalt memories, and the hippocampus. *Behavioural*

      Brain Research, *110*(1–2), 73–81. https://doi.org/10.1016/S0166-4328(99)00186-2

Fanselow, M. S., & Dong, H.-W. (2010). Are The Dorsal and Ventral Hippocampus functionally

      distinct structures? *Neuron*, *65*(1), 7. https://doi.org/10.1016/j.neuron.2009.11.031

Floresco, S. B., Blaha, C. D., Yang, C. R., & Phillips, A. G. (2001). Modulation of Hippocampal

      and Amygdalar-Evoked Activity of Nucleus Accumbens Neurons by Dopamine: Cellular

Mechanisms of Input Selection. *The Journal of Neuroscience*, *21*(8), 2851–2860.

https://doi.org/10.1523/JNEUROSCI.21-08-02851.2001

Floresco, S. B., & Phillips, A. G. (1999). Dopamine and hippocampal input to the nucleus

accumbens play an essential role in the search for food in an unpredictable environment.

Psychobiology, *27*(2), 277–286. https://doi.org/10.3758/BF03332121

Francis, T. C., & Lobo, M. K. (2017). Emerging Role for Nucleus Accumbens Medium Spiny

Neuron Subtypes in Depression. *Biological Psychiatry*, *81*(8), 645–653.

https://doi.org/10.1016/j.biopsych.2016.09.007

French, S. J., & Totterdell, S. (2002). Hippocampal and prefrontal cortical inputs

monosynaptically converge with individual projection neurons of the nucleus accumbens.

The Journal of Comparative Neurology, *446*(2), 151–165.

https://doi.org/10.1002/cne.10191

Friedman, D. P., Aggleton, J. P., & Saunders, R. C. (2002). Comparison of hippocampal,

amygdala, and perirhinal projections to the nucleus accumbens: Combined anterograde

and retrograde tracing study in the Macaque brain. *Journal of Comparative Neurology*,

450(4), 345–365. https://doi.org/10.1002/cne.10336

Fu, M., Yu, X., Lu, J., & Zuo, Y. (2012). Repetitive motor learning induces coordinated formation

of clustered dendritic spines in vivo. *Nature*, *483*(7387), 92–95.

https://doi.org/10.1038/nature10844

Govindarajan, A., Kelleher, R. J., & Tonegawa, S. (2006). A clustered plasticity model of long-

term memory engrams. *Nature Reviews. Neuroscience*, *7*(7), 575–583.

https://doi.org/10.1038/nrn1937

Groenewegen, H. J., Der Zee, E. V.-V., Te Kortschot, A., & Witter, M. P. (1987). Organization of

the projections from the subiculum to the ventral striatum in the rat. A study using

anterograde transport of Phaseolus vulgaris leucoagglutinin. *Neuroscience*, *23*(1), 103–

120. https://doi.org/10.1016/0306-4522(87)90275-2

Gruber, A. J., Hussain, R. J., & O'Donnell, P. (2009). The Nucleus Accumbens: A Switchboard

    for Goal-Directed Behaviors. *PLoS ONE*, *4*(4), e5062.

    https://doi.org/10.1371/journal.pone.0005062

Harvey, C. D., & Svoboda, K. (2007). Locally dynamic synaptic learning rules in pyramidal

    neuron dendrites. *Nature*, *450*(7173), 1195–1200. https://doi.org/10.1038/nature06416

Heller, A. S., Shi, T. C., Ezie, C. E. C., Reneau, T. R., Baez, L. M., Gibbons, C. J., & Hartley, C.

    A. (2020). Association between real-world experiential diversity and positive affect

    relates to hippocampal–striatal functional connectivity. *Nature Neuroscience*, *23*(7), 800–

    804. https://doi.org/10.1038/s41593-020-0636-4

Humphries, M. D., & Prescott, T. J. (2010). The ventral basal ganglia, a selection mechanism at

    the crossroads of space, strategy, and reward. *Progress in Neurobiology*, *90*(4), 385–

    417. https://doi.org/10.1016/j.pneurobio.2009.11.003

Huntley, E. D., Marusak, H. A., Berman, S. E., Zundel, C. G., Hatfield, J. R. B., Keating, D. P., &

    Rabinak, C. A. (2020). Adolescent substance use and functional connectivity between

    the ventral striatum and hippocampus. *Behavioural Brain Research*, *390*, 112678.

    https://doi.org/10.1016/j.bbr.2020.112678

Ibrahim, K. M., Massaly, N., Yoon, H.-J., Sandoval, R., Widman, A. J., Heuermann, R. J.,

    Williams, S., Post, W., Pathiranage, S., Lintz, T., Zec, A., Park, A., Yu, W., Kash, T. L.,

    Gereau, R. W., & Morón, J. A. (2024). Dorsal hippocampus to nucleus accumbens

    projections drive reinforcement via activation of accumbal dynorphin neurons. *Nature*

    Communications, *15*(1), 750. https://doi.org/10.1038/s41467-024-44836-9

Ishizuka, N., Weber, J., & Amaral, D. G. (1990). Organization of intrahippocampal projections

    originating from CA3 pyramidal cells in the rat. *Journal of Comparative Neurology*,

    295(4), 580–623. https://doi.org/10.1002/cne.902950407

Ito, R., Robbins, T. W., Pennartz, C. M., & Everitt, B. J. (2008). Functional Interaction between

    the Hippocampus and Nucleus Accumbens Shell Is Necessary for the Acquisition of

588   Appetitive Spatial Context Conditioning. *Journal of Neuroscience*, *28*(27), 6950–6959.

589   https://doi.org/10.1523/JNEUROSCI.1615-08.2008

590 Kelley, A. E., & Domesick, V. B. (1982). The distribution of the projection from the hippocampal

591   formation to the nucleus accumbens in the rat: An anterograde and retrograde-

592   horseradish peroxidase study. *Neuroscience*, *7*(10), 2321–2335.

593   https://doi.org/10.1016/0306-4522(82)90198-1

594 Kjelstrup, K. B., Solstad, T., Brun, V. H., Hafting, T., Leutgeb, S., Witter, M. P., Moser, E. I., &

595   Moser, M.-B. (2008). Finite Scale of Spatial Representation in the Hippocampus.

596   *Science*, *321*(5885), 140–143. https://doi.org/10.1126/science.1157086

597 Kjelstrup, K. G., Tuvnes, F. A., Steffenach, H.-A., Murison, R., Moser, E. I., & Moser, M.-B.

598   (2002). Reduced fear expression after lesions of the ventral hippocampus. *Proceedings*

599   *of the National Academy of Sciences*, *99*(16), 10825–10830.

600   https://doi.org/10.1073/pnas.152112399

601 Klapoetke, N. C., Murata, Y., Kim, S. S., Pulver, S. R., Birdsey-Benson, A., Cho, Y. K.,

602   Morimoto, T. K., Chuong, A. S., Carpenter, E. J., Tian, Z., Wang, J., Xie, Y., Yan, Z.,

603   Zhang, Y., Chow, B. Y., Surek, B., Melkonian, M., Jayaraman, V., Constantine-Paton,

604   M., … Boyden, E. S. (2014). Independent optical excitation of distinct neural populations.

605   *Nature Methods*, *11*(3), 338–346. https://doi.org/10.1038/nmeth.2836

606 Lamsa, K., & Lau, P. (2019). Long-term plasticity of hippocampal interneurons during in vivo

607   memory processes. *Current Opinion in Neurobiology*, *54*, 20–27.

608   https://doi.org/10.1016/j.conb.2018.08.006

609 Lansink, C. S., Goltstein, P. M., Lankelma, J. V., Joosten, R. N. J. M. A., McNaughton, B. L., &

610   Pennartz, C. M. A. (2008). Preferential Reactivation of Motivationally Relevant

611   Information in the Ventral Striatum. *Journal of Neuroscience*, *28*(25), 6372–6382.

612   https://doi.org/10.1523/JNEUROSCI.1054-08.2008

Lansink, C. S., Goltstein, P. M., Lankelma, J. V., McNaughton, B. L., & Pennartz, C. M. A. (2009). Hippocampus Leads Ventral Striatum in Replay of Place-Reward Information. *PLoS Biology*, *7*(8), e1000173. https://doi.org/10.1371/journal.pbio.1000173

LeGates, T. A., Kvarta, M. D., Tooley, J. R., Francis, T. C., Lobo, M. K., Creed, M. C., & Thompson, S. M. (2018). Reward behaviour is regulated by the strength of hippocampus–nucleus accumbens synapses. *Nature, 564*(7735), 258–262. https://doi.org/10.1038/s41586-018-0740-8

Li, Z., Chen, Z., Fan, G., Li, A., Yuan, J., & Xu, T. (2018). Cell-Type-Specific Afferent Innervation of the Nucleus Accumbens Core and Shell. *Frontiers in Neuroanatomy*, *12*, 84. https://doi.org/10.3389/fnana.2018.00084

Liu, S. J., & Zukin, R. S. (2007). Ca2+-permeable AMPA receptors in synaptic plasticity and neuronal death. *Trends in Neurosciences*, *30*(3), 126–134. https://doi.org/10.1016/j.tins.2007.01.006

Mameli, M., Halbout, B., Creton, C., Engblom, D., Parkitna, J. R., Spanagel, R., & Lüscher, C. (2009). Cocaine-evoked synaptic plasticity: Persistence in the VTA triggers adaptations in the NAc. *Nature Neuroscience*, *12*(8), 1036–1041. https://doi.org/10.1038/nn.2367

Mannella, F., Gurney, K., & Baldassarre, G. (2013). The nucleus accumbens as a nexus between values and goals in goal-directed behavior: A review and a new hypothesis. *Frontiers in Behavioral Neuroscience*, *7*. https://doi.org/10.3389/fnbeh.2013.00135

Marinescu, A.-M., & Labouesse, M. A. (2024). The nucleus accumbens shell: A neural hub at the interface of homeostatic and hedonic feeding. *Frontiers in Neuroscience*, *18*, 1437210. https://doi.org/10.3389/fnins.2024.1437210

McCutcheon, J. E., Wang, X., Tseng, K. Y., Wolf, M. E., & Marinelli, M. (2011). Calcium-Permeable AMPA Receptors Are Present in Nucleus Accumbens Synapses after Prolonged Withdrawal from Cocaine Self-Administration But Not Experimenter-

Administered Cocaine. *The Journal of Neuroscience*, *31*(15), 5737–5743.

https://doi.org/10.1523/JNEUROSCI.0350-11.2011

Morici, J. F., Paiva-Lima, I., Silva, A., Pronier, É., & Girardeau, G. (2025). *Dorso-ventral*

hippocampus neural assemblies reactivate during sleep following an aversive

experience (p. 2024.06.13.598918). bioRxiv. https://doi.org/10.1101/2024.06.13.598918

Muir, J., Tse, Y. C., Iyer, E. S., Biris, J., Cvetkovska, V., Lopez, J., & Bagot, R. C. (2020).

Ventral Hippocampal Afferents to Nucleus Accumbens Encode Both Latent Vulnerability

and Stress-Induced Susceptibility. *Biological Psychiatry*, *88*(11), 843–854.

https://doi.org/10.1016/j.biopsych.2020.05.021

Mulder, A. B., Hodenpijl, M. G., & Lopes Da Silva, F. H. (1998). Electrophysiology of the

Hippocampal and Amygdaloid Projections to the Nucleus Accumbens of the Rat:

Convergence, Segregation, and Interaction of Inputs. *The Journal of Neuroscience*,

18(13), 5095–5102. https://doi.org/10.1523/JNEUROSCI.18-13-05095.1998

Murakoshi, H., Wang, H., & Yasuda, R. (2011). Local, persistent activation of Rho GTPases

during plasticity of single dendritic spines. *Nature*, *472*(7341), 100–104.

https://doi.org/10.1038/nature09823

Nakazawa, K., Sun, L. D., Quirk, M. C., Rondi-Reig, L., Wilson, M. A., & Tonegawa, S. (2003).

Hippocampal CA3 NMDA Receptors Are Crucial for Memory Acquisition of One-Time

Experience. *Neuron*, *38*(2), 305–315. https://doi.org/10.1016/S0896-6273(03)00165-X

O'Donnell, P., & Grace, A. (1995). Synaptic interactions among excitatory afferents to nucleus

accumbens neurons: Hippocampal gating of prefrontal cortical input. *The Journal of*

Neuroscience, *15*(5), 3622–3639. https://doi.org/10.1523/JNEUROSCI.15-05-

03622.1995

Patterson, D., Khan, N., Collins, E. A., Stewart, N. R., Sassaninejad, K., Yeates, D., Lee, A. C.

H., & Ito, R. (2025). Ventral hippocampus to nucleus accumbens shell circuit regulates

approach decisions during motivational conflict. *PLOS Biology*, *23*(1), e3002722.

https://doi.org/10.1371/journal.pbio.3002722

Paxinos, G., & Franklin, K. B. J. (2019). *Paxinos and Franklin's The mouse brain in stereotaxic*

coordinates (Fifth edition). Academic Press, an imprint of Elsevier.

Peleg-Raibstein, D., & Feldon, J. (2006). Effects of dorsal and ventral hippocampal NMDA

stimulation on nucleus accumbens core and shell dopamine release.

Neuropharmacology, *51*(5), 947–957. https://doi.org/10.1016/j.neuropharm.2006.06.002

Pennartz, C. M. A., Ito, R., Verschure, P. F. M. J., Battaglia, F. P., & Robbins, T. W. (2011). The

hippocampal–striatal axis in learning, prediction and goal-directed behavior. *Trends in*

Neurosciences, *34*(10), 548–559. https://doi.org/10.1016/j.tins.2011.08.001

Phillipson, O. T., & Griffiths, A. C. (1985). The topographic order of inputs to nucleus

accumbens in the rat. *Neuroscience*, *16*(2), 275–296. https://doi.org/10.1016/0306-

4522(85)90002-8

R. Yang, C., & J. Mogenson, G. (1984). Electrophysiological responses of neurones in the

nucleus accumbens to hippocampal stimulation and the attenuation of the excitatory

responses by the mesolimbic dopaminergic system. *Brain Research*, *324*(1), 69–84.

https://doi.org/10.1016/0006-8993(84)90623-1

Redondo, R. L., & Morris, R. G. M. (2011). Making memories last: The synaptic tagging and

capture hypothesis. *Nature Reviews. Neuroscience*, *12*(1), 17–30.

https://doi.org/10.1038/nrn2963

Roberts, A. C., Tomic, D. L., Parkinson, C. H., Roeling, T. A., Cutter, D. J., Robbins, T. W., &

Everitt, B. J. (2007). Forebrain connectivity of the prefrontal cortex in the marmoset

monkey (*Callithrix jacchus*): An anterograde and retrograde tract-tracing study. *Journal*

of Comparative Neurology, *502*(1), 86–112. https://doi.org/10.1002/cne.21300

Rudy, J. W., Barrientos, R. M., & O'Reilly, R. C. (2002). Hippocampal formation supports

conditioning to memory of a context. *Behavioral Neuroscience*, *116*(4), 530–538.

https://doi.org/10.1037/0735-7044.116.4.530

Safari, S., Ahmadi, N., Mohammadkhani, R., Ghahremani, R., Khajvand-Abedeni, M., Shahidi,

S., Komaki, A., Salehi, I., & Karimi, S. A. (2021). Sex differences in spatial learning and

memory and hippocampal long-term potentiation at perforant pathway-dentate gyrus

(PP-DG) synapses in Wistar rats. *Behavioral and Brain Functions*, *17*(1), 9.

https://doi.org/10.1186/s12993-021-00184-y

Salimpoor, V. N., Van Den Bosch, I., Kovacevic, N., McIntosh, A. R., Dagher, A., & Zatorre, R.

J. (2013). Interactions Between the Nucleus Accumbens and Auditory Cortices Predict

Music Reward Value. *Science*, *340*(6129), 216–219.

https://doi.org/10.1126/science.1231059

Savage, L. (2004). The effects of hippocampal lesions on learning, memory, and reward

expectancies. *Neurobiology of Learning and Memory*, *82*(2), 109–119.

https://doi.org/10.1016/j.nlm.2004.05.002

Schuman, E. M., & Madison, D. V. (1994). Locally distributed synaptic potentiation in the

hippocampus. *Science (New York, N.Y.)*, *263*(5146), 532–536.

https://doi.org/10.1126/science.8290963

Scofield, M. D., Heinsbroek, J. A., Gipson, C. D., Kupchik, Y. M., Spencer, S., Smith, A. C. W.,

Roberts-Wolfe, D., & Kalivas, P. W. (2016). The Nucleus Accumbens: Mechanisms of

Addiction across Drug Classes Reflect the Importance of Glutamate Homeostasis.

Pharmacological Reviews, *68*(3), 816–871. https://doi.org/10.1124/pr.116.012484

Sjulson, L., Peyrache, A., Cumpelik, A., Cassataro, D., & Buzsáki, G. (2018). Cocaine Place

Conditioning Strengthens Location-Specific Hippocampal Coupling to the Nucleus

Accumbens. *Neuron*, *98*(5), 926-934.e5. https://doi.org/10.1016/j.neuron.2018.04.015

Soares-Cunha, C., de Vasconcelos, N. A. P., Coimbra, B., Domingues, A. V., Silva, J. M.,

Loureiro-Campos, E., Gaspar, R., Sotiropoulos, I., Sousa, N., & Rodrigues, A. J. (2020).

Nucleus accumbens medium spiny neurons subtypes signal both reward and aversion.

Molecular Psychiatry, *25*(12), 3241–3255. https://doi.org/10.1038/s41380-019-0484-3

Sosa, M., Joo, H. R., & Frank, L. M. (2020). Dorsal and Ventral Hippocampal Sharp-Wave

Ripples Activate Distinct Nucleus Accumbens Networks. *Neuron*, *105*(4), 725-741.e8.

https://doi.org/10.1016/j.neuron.2019.11.022

Squire, L. R. (1992). Memory and the hippocampus: A synthesis from findings with rats,

monkeys, and humans. *Psychological Review*, *99*(2), 195–231.

https://doi.org/10.1037/0033-295X.99.2.195

Steele, R. J., & Morris, R. G. M. (1999). Delay-dependent impairment of a matching-to-place

task with chronic and intrahippocampal infusion of the NMDA-antagonist D-AP5.

Hippocampus, *9*(2), 118–136. https://doi.org/10.1002/(SICI)1098-

1063(1999)9:2<118::AID-HIPO4>3.0.CO;2-8

Strange, B. A., Witter, M. P., Lein, E. S., & Moser, E. I. (2014). Functional organization of the

hippocampal longitudinal axis. *Nature Reviews Neuroscience*, *15*(10), 655–669.

https://doi.org/10.1038/nrn3785

Swanson, L. W., & Cowan, W. M. (1977). An autoradiographic study of the organization of the

efferet connections of the hippocampal formation in the rat. *Journal of Comparative*

Neurology, *172*(1), 49–84. https://doi.org/10.1002/cne.901720104

Tao, S., Wang, Y., Peng, J., Zhao, Y., He, X., Yu, X., Liu, Q., Jin, S., & Xu, F. (2021). Whole-

Brain Mapping the Direct Inputs of Dorsal and Ventral CA1 Projection Neurons. *Frontiers*

in Neural Circuits, *15*, 643230. https://doi.org/10.3389/fncir.2021.643230

Terranova, J. I., Yokose, J., Osanai, H., Marks, W. D., Yamamoto, J., Ogawa, S. K., & Kitamura,

T. (2022). Hippocampal-amygdala memory circuits govern experience-dependent

observational fear. *Neuron*, *110*(8), 1416-1431.e13.

https://doi.org/10.1016/j.neuron.2022.01.019

Trouche, S., Koren, V., Doig, N. M., Ellender, T. J., El-Gaby, M., Lopes-dos-Santos, V., Reeve,

H. M., Perestenko, P. V., Garas, F. N., Magill, P. J., Sharott, A., & Dupret, D. (2019). A

Hippocampus-Accumbens Tripartite Neuronal Motif Guides Appetitive Memory in Space.

Cell, *176*(6), 1393-1406.e16. https://doi.org/10.1016/j.cell.2018.12.037

Williams, E. S., Manning, C. E., Eagle, A. L., Swift-Gallant, A., Duque-Wilckens, N.,

Chinnusamy, S., Moeser, A., Jordan, C., Leininger, G., & Robison, A. J. (2020).

Androgen-Dependent Excitability of Mouse Ventral Hippocampal Afferents to Nucleus

Accumbens Underlies Sex-Specific Susceptibility to Stress. *Biological Psychiatry*, *87*(6),

492–501. https://doi.org/10.1016/j.biopsych.2019.08.006

Wolf, M. E., & Tseng, K. Y. (2012). Calcium-permeable AMPA receptors in the VTA and nucleus

accumbens after cocaine exposure: When, how, and why? *Frontiers in Molecular*

Neuroscience, *5*. https://doi.org/10.3389/fnmol.2012.00072

Xia, S., Yu, J., Huang, X., Sesack, S. R., Huang, Y. H., Schlüter, O. M., Cao, J.-L., & Dong, Y.

(2020). Cortical and Thalamic Interaction with Amygdala-to-Accumbens Synapses. *The*

Journal of Neuroscience, *40*(37), 7119–7132. https://doi.org/10.1523/JNEUROSCI.1121-

20.2020

Yang, A. K., Mendoza, J. A., Lafferty, C. K., Lacroix, F., & Britt, J. P. (2020). Hippocampal Input

to the Nucleus Accumbens Shell Enhances Food Palatability. *Biological Psychiatry*,

87(7), 597–608. https://doi.org/10.1016/j.biopsych.2019.09.007

Yang, F.-C., & Liang, K. C. (2014). Interactions of the dorsal hippocampus, medial prefrontal

cortex and nucleus accumbens in formation of fear memory: Difference in inhibitory

avoidance learning and contextual fear conditioning. *Neurobiology of Learning and*

Memory, *112*, 186–194. https://doi.org/10.1016/j.nlm.2013.07.017

Yu, J., Sesack, S. R., Huang, Y., Schlüter, O. M., Grace, A. A., & Dong, Y. (2022). Contingent

Amygdala Inputs Trigger Heterosynaptic LTP at Hippocampus-To-Accumbens

Synapses. *The Journal of Neuroscience*, *42*(34), 6581–6592.

https://doi.org/10.1523/JNEUROSCI.0838-22.2022

Yuste, R. (2011). Dendritic Spines and Distributed Circuits. *Neuron*, *71*(5), 772–781.

https://doi.org/10.1016/j.neuron.2011.07.024

Zhou, Y., Zhu, H., Liu, Z., Chen, X., Su, X., Ma, C., Tian, Z., Huang, B., Yan, E., Liu, X., & Ma,

L. (2019). A ventral CA1 to nucleus accumbens core engram circuit mediates

conditioned place preference for cocaine. *Nature Neuroscience*, *22*(12), 1986–1999.

https://doi.org/10.1038/s41593-019-0524-y

Zingg, B., Chou, X., Zhang, Z., Mesik, L., Liang, F., Tao, H. W., & Zhang, L. I. (2017). AAV-

Mediated Anterograde Transsynaptic Tagging: Mapping Corticocollicular Input-Defined

Neural Pathways for Defense Behaviors. *Neuron*, *93*(1), 33–47.

https://doi.org/10.1016/j.neuron.2016.11.045

FIGURES

Figure 1. Overlapping dHipp and vHipp axon terminal fields in the NAc. a) Viral injection

strategy and experimental timeline. b) Injection sites in the hippocampus. Scale bar=500μm. Atlas

image from (Paxinos & Franklin, 2019) c) Overlap of dHipp and vHipp axon terminal fields in the

NAc. Scale bar=500μm. Atlas image from (Paxinos & Franklin, 2019) d) Close-up image of

overlapping dHipp and vHipp axon terminal fields within the NAc ventromedial shell (top). Scale

bar=20μm. With 3-dimensional reconstruction of overlapping axon terminals (bottom). Scale bar=

0.5μm

Figure 2. dHipp⁺/vHipp⁺-innervated neurons in the NAc ventromedial shell. a) Viral targeting

strategy to selectively label dually innervated neurons b) Cartoon representation of singly vs

dually innervated neurons. c) Image of NAc shell depicting GCaMP-expressing cells in the NAc

ventromedial shell. Scale bar=100μm. Atlas image from (Paxinos & Franklin, 2019) d) Close-up

image of GCaMP-expressing cells in the NAc ventromedial shell, where neuron morphology is

consistent with medium spiny neurons. Scale bar=100μm. e) Viral targeting strategy and

experimental timeline to label cells innervated by dHipp and vHipp. f) Cartoon representation of

singly vs dually innervated neurons. g) Image of cells innervated by dHipp (green) and vHipp

(magenta). Dually innervated neurons express both reporters and appear white. Arrowheads point

out examples of dually innervated neurons. Scale bar = 100μm h) Quantification of cells

innervated single and dually innervated neurons in the NAc medial shell across 4 slices.

Figure 3. Individual NAc neurons are functionally innervated by dHipp and vHipp. a) Viral

targeting and optogenetic approach. Electrophysiological recording strategy and representative

traces from a dually innervated NAc MSN. b) Similar AMPA:NMDA receptor ratios between dHipp-

and vHipp-MSN synapses (n=8 cells/4 mice, p=0.1953, Wilcoxon test). Scale bars on

representative traces = 40pA/10ms c) Rectification curve reveals linear relationship, indicating a

lack of calcium-permeable AMPA receptors (n=8 cells/4 mice). d) PPRs recorded from dHipp-

MSN and vHipp-MSN synapses on dually innervated neurons (n=16 cells/10 mice, *p=0.0394,

Paired t-test). e) Rise time was calculated for dHipp- and vHipp-evoked EPSCs, showcasing

similar rise times (n=31 cells/12 mice, p=0.2877, Wilcoxon test).

Figure 4. Spatial proximity of dHipp and vHipp puncta along distal dendrites of MSN. a)

Cartoon depicting optogenetic, electrophysiology, and cell-filling strategy. b) Combined and

separated images depicting an MSN filled with CF633 confirmed to have functional dHipp and

vHipp synapses, dHipp puncta (green), vHipp puncta (red). c) Three-dimensional reconstruction

of an MSN dendritic segment (cyan) with overlapping/merged dHipp (red) and vHipp (green)

afferents.

Figure 5. Heterosynaptic potentiation of dHipp-MSN and vHipp-MSN synapses. a) Viral

expression, optogenetic, and electrophysiology strategy. b) Recording from dually innervated

MSNs, sequential and synchronous stimulation assay reveals non-linear summation of responses

(n=16 cells/10 mice, **p=0.0027, Wilcoxon test). c) Sequential and synchronous stimulation assay

performed in singly innervated MSNs reveals similar responses to a single wavelength of light or

both wavelengths of light (n=16 cells/10 mice, p=0.4037, Wilcoxon test). d) Theoretical probability

of dual success, P_{exp}, compared with observed dual response success, P_{obs}, during sync period

(n=9 cells/5 mice, *p=0.0273, Wilcoxon test).

Figure S1. Prevention of cross-contamination when using 470nm light and the optogenetic

protein, Chrimson.

a) Mice were injected with only a Chrimson-containing viral construct and slices were taken

at least 5 weeks after surgery to perform electrophysiology. b) A response curve was generated

by sampling Chrimson-evoked responses using 470nm light, showing minimal responses at a

laser intensity of 0.75mW, and no Chrimson-evoked response at laser intensities ≤0.5mW. EPSC

amplitudes were normalized to the maximum EPSC amplitude.

Figure 1

a.

AAV1-hSyn-Cre
AAV1-Ef1a-Flpo

dHipp
vHipp
NAc
Ai195

b.

dHipp
NAc
vHipp
NAc
singly-innervated

dHipp   vHipp
NAc
dually-innervated

c.

d.

e.

AAV9-Ef1a-
fDIO-eYFP

AAV1-hSyn-Cre
AAV1-Ef1a-Flpo

dHipp
vHipp
NAc
Ai14

f.

dHipp
NAc
vHipp
NAc
singly-innervated

dHipp   vHipp
NAc
dually-innervated

g.

NAc neurons innervated
by dHipp

NAc neurons innervated
by vHipp

Merge

h.

28%
8%
65%

■ dH innervated (28%, 68 cells)
■ vH innervated (65%, 159 cells)
■ Dually innervated (8%, 19 cells)

Figure 2

Figure 3

Figure 4

Figure 5

a. Surgery — >= 5 weeks — Electrophysiology

AAV-CAMKII-Chrimson-mCherry

dHipp
vHipp
NAc

b. **Chrimson responses to 470nm light**

EPSC Amplitude (norm) vs Laser Intensity (mW)

Supplemental Figure 1