Title: Heterosynaptic interactions between dorsal and ventral hippocampus in individual medium

spiny neurons of the nucleus accumbens ventromedial shell

Abbreviated title: Interactions between Hipp inputs in the NAc

Authors: Ashley E. Copenhaver[1], Sydnee Vance[1], Sarah A. Snider[1], Kaela Befano[1], J. Branwen

She[1], Tara A. LeGates[1,2]

[1] Department of Biological Sciences, University of Maryland, Baltimore County (UMBC)

[2] Department of Pharmacology and Physiology, University of Maryland School of Medicine

Corresponding author:

Tara LeGates, Ph.D.

1000 Hilltop Circle, Interdisciplinary Life Sciences Building 315

Baltimore, MD 21250

tlegates@umbc.edu

Number of pages:

Number of figures: 5 + 2 supplemental figures

Number of words, abstract: 249

Number of words, introduction: 647

Number of words, discussion: 1474

Conflict of interest statement: The authors declare no competing financial interests.

Acknowledgments: We would like to thank Dr. Tagide deCarvalho for her support with microscopy

and image analysis, and Dr. Ramesh Chandra for his help with viruses. We would also like to

thank Jennifer Pham for her help with injections and Dr. Brian Mathur for his helpful suggestions

27 on this project. This work was supported by NSF-IOS2402645, T32GM144876-02,

28 T34GM136497, the Merck Academic Fellowship Program, and startup funds provided by UMBC.

ABSTRACT

Establishing learned associations between rewarding stimuli and the context under which those rewards are encountered is critical for survival. Hippocampal input to the nucleus accumbens (NAc) provides important environmental context to reward processing to support goal-directed behaviors. This connection consists of two independent pathways originating from the dorsal (dHipp) or ventral (vHipp) hippocampus, which have previously been considered functionally and anatomically distinct. Here, we show overlap in dHipp and vHipp terminal fields in the NAc, leading us to reconsider this view and raise new questions regarding the potential interactions between dHipp and vHipp pathways in the NAc. Using optogenetics, electrophysiology, and transsynaptic labeling in mice, we investigated anatomical and functional convergence of dHipp and vHipp inputs in the NAc. Transsynaptic labeling revealed a subpopulation of dually innervated cells in the NAc medial shell, confirmed by independent optogenetic manipulation of dHipp and vHipp inputs during whole-cell electrophysiological recordings. Further analysis revealed closely apposed dHipp and vHipp inputs along dendritic branches, and simultaneous stimulation of both inputs elicited heterosynaptic potentiation. Comparison of observed and theoretical success rates suggests interactions between dHipp-NAc and vHipp-NAc synapses may occur presynaptically. Altogether, these results demonstrate that inputs originating from dHipp and vHipp converge onto a subset of NAc neurons with synapses positioned to enable rapid heterosynaptic interactions, indicating integration of these inputs at the single-neuron level. Exploring the physiological and behavioral implications of this convergence will offer new insights into how individual neurons incorporate information from distinct inputs and how this integration may shape learning.

SIGNIFICANCE STATEMENT

Forming associations between rewards and the circumstances under which they are experienced is vital for survival. Hipp input to the NAc is essential for associating rewards with

their environmental context to effectively guide motivated behaviors. This connection consists of

two separate pathways originating from dHipp and vHipp that have long been considered distinct.

Here, we reveal a subpopulation of neurons in the NAc shell innervated by both Hipp subregions

as well as heterosynaptic interactions that occur between dHipp and vHipp synapses. These

findings suggest that integration of distinct hippocampal information occurs at the single-neuron

level, providing a critical mechanism underlying learning and motivated behavior while also

opening new avenues for understanding how diverse contextual and reward signals shape

decision-making.

INTRODUCTION

The integration of diverse information across neural systems is crucial for driving complex

behaviors. The brain dynamically integrates information from multiple sources, enabling

organisms to react to stimuli and establish learned associations crucial for survival. Plasticity of

and interactions between synapses enables coordinated regulation across multiple inputs, critical

for orchestrating cell-wide functional responses (Bailey et al., 2000a; Govindarajan et al., 2006a;

Harvey & Svoboda, 2007a; Redondo & Morris, 2011a; Schuman & Madison, 1994a). However,

the mechanisms underlying how convergent inputs interact to shape neuronal responses remain

unclear.

The nucleus accumbens (NAc) is a key brain region that integrates diverse information from

several brain areas to effectively mediate goal-directed behaviors (Francis & Lobo, 2017; French

& Totterdell, 2002; Gruber et al., 2009; Mannella et al., 2013; Marinescu & Labouesse, 2024;

Salimpoor et al., 2013; Soares-Cunha et al., 2020). Hippocampal (Hipp) input is a major source

of excitatory drive that influences NAc neuron activity (Brog et al., 1993; Calhoon & O'Donnell,

2013; Groenewegen et al., 1987; Kelley & Domesick, 1982; Li et al., 2018; O'Donnell & Grace,

1995; Phillipson & Griffiths, 1985; Trouche et al., 2019), and because of the Hipp's essential role

in contextual and spatial learning (Duarte-Guterman et al., 2015; Lamsa & Lau, 2019; Safari et

al., 2021; Savage, 2004; Squire, 1992), Hipp-NAc synapses are also key sites for integrating

rewards with related contextual and spatial information. Furthermore, the modulation of the

strength of Hipp-NAc synapses is a critical mediator of motivated behaviors and is necessary for

supporting contextual- and spatial-based reward learning (Huntley et al., 2020; Ibrahim et al.,

2024; Ito et al., 2008, 2008; LeGates et al., 2018; Patterson et al., 2025, p. 202; Pennartz et al.,

2011; Sjulson et al., 2018; Trouche et al., 2019; F.-C. Yang & Liang, 2014).

The connection between the Hipp and NAc consists of two independent pathways originating

from the dorsal and ventral Hipp (dHipp and vHipp), which are considered functionally and

anatomically distinct (Amaral & Witter, 1989; Bannerman et al., 2002; Fanselow & Dong, 2010;

Friedman et al., 2002; K. B. Kjelstrup et al., 2008; K. G. Kjelstrup et al., 2002; Roberts et al., 2007;

Strange et al., 2014). Notably, the NAc is one of the few regions that receives input from both the

dHipp and vHipp (Friedman et al., 2002; Kelley & Domesick, 1982; Phillipson & Griffiths, 1985;

Trouche et al., 2019). Although dHipp and vHipp are interconnected within the Hipp (Ishizuka et

al., 1990; Swanson & Cowan, 1977; Tao et al., 2021), interactions through connections with other

brain regions have been proposed as a mechanism linking cognitive and emotional processes

(Fanselow & Dong, 2010). In a multifaceted environment, integration of diverse information is

important for properly mediating complex behaviors, and the behavioral complexity governed by

these pathways warrants synaptic integration to support learning and memory. This is supported

by recent findings demonstrating a key role for convergence of dHipp and vHipp input to the

amygdala in observational contextual fear memory (Terranova et al., 2022), yet the synaptic basis

of this convergence and whether it is also observed in the NAc remains unknown.

Using a combination of microscopy, electrophysiology, and optogenetics, we identified and

characterized a subpopulation of medium spiny neurons (MSNs) in the medial NAc shell (NAcSh)

that are functionally innervated by both the dHipp and vHipp. dHipp and vHipp synapses on dually

innervated MSNs have similar basal synaptic properties and are positioned at distances close

enough to enable heterosynaptic interactions. These findings suggest that convergent Hipp inputs

interact at the synaptic level, which may contribute to the integration of spatial and contextual

information to modulate motivated behaviors. Overall, understanding the synaptic integration of

dHipp and vHipp inputs in the NAc not only enhances our knowledge of reward-related behaviors,

but also offers insight into the fundamental task of neuronal processing and provides a foundation

for developing targeted interventions for disorders where these mechanisms are dysregulated.

METHODS

Animals

Male and female C57BL/6J, Drd1a-tdTomato, Ai14, and Ai195 mice were bred in-house. Mice

were housed with same-sex cage mates in a temperature- and humidity-controlled environment

under a 12:12 light cycle (lights on at 07:00), with food available *ad libitum*. We did not track

estrous cycle in females. All experiments were performed in accordance with the regulations set

forth by the Institutional Animal Care and Use Committee at the University of Maryland, Baltimore

County.

Stereotaxic surgeries

Mice (6-8 weeks old) were anesthetized with 3% isoflurane and underwent stereotaxic

surgery. A Hamilton syringe was used to inject adeno-associated viruses (AAV) into the dHipp (in

millimeters: from bregma, anterior-posterior (AP), -3.7; medial-lateral (ML), ±2.4; dorsal-ventral

(DV), -2.8), vHipp (from bregma, AP, -3.7; ML, ±3.0; DV, -4.8), and NAc (in millimeters: from

bregma, anterior-posterior (AP), +1.8; medial-lateral (ML), ±0.6; dorsal-ventral (DV), -4.7). Viruses

were injected unilaterally or bilaterally and were infused at a rate of 0.1uL per minute. The injection

needle was left in place for 10 minutes following the infusion. After surgery, mice were placed in

their home cage on a heating pad for recovery. Carprofen (5mg/kg) was subcutaneously injected

daily for 3 days after surgery. Mice recovered for 5-7 weeks for viral expression before anatomical

or electrophysiological experiments. Viral constructs pAAV-hSyn-EGFP (Addgene #50465-AAV5)

and pAAV-hSyn-mCherry (Addgene #114472-AAV5) were used to visualize dHipp and vHipp

axon terminal fields. Anatomical labeling of dually innervated neurons involved the use of

pENN.AAV.hSyn.Cre.WPRE.hGH (AAV1, 1013 titer, UMB Viral Vector Core), pAAV-EF1a-Flpo

(AAV1, 1013 titer, UMB Viral Vector Core), and pAAV-hSyn-DIO-EGFP (Addgene #50457-AAV2).

Specifically, pENN.AAV.hSyn.Cre.WPRE.hGH and pAAV-EF1a-Flpo were injected into the dHipp

or vHipp of Ai195 mice for selective labeling of dually innervated neurons.

pENN.AAV.hSyn.Cre.WPRE.hGH and pAAV-EF1a-Flpo were injected into the dHipp or vHipp

and pAAV-hSyn-DIO-EGFP was injected in the NAc of Ai14 mice to label neurons single and

dually innervated neurons. For optogenetic experiments, viral vectors pAAV-Syn-ChrimsonR-tdT

(Addgene ID: 59171-AAV5) and either pAAV-Syn-Chronos-GFP (Addgene ID: 59170-AAV5) or

pAAV-hSyn-hChR2(H134R)-EYFP (Addgene ID: 26973-AAV5) were used. For all experiments,

injection sites were verified by fluorescence microscopy, and brains with off target expression

were removed from the experiment.

147 Visualization of axon terminal fields and dually innervated neurons

Mice expressing viruses for anatomical tracing experiments were first anesthetized with

isoflurane and then perfused with 1X phosphate buffer solution (PBS), followed by 4%

paraformaldehyde (PFA). Brains were removed and post-fixed in 4% PFA overnight at 4ºC and

were then transferred to 1X PBS and stored at 4ºC for up to 3 weeks. Brains were sectioned

(100µm) using a VT1000S Leica Microsystems vibratome. Slices were then mounted with

Vectashield Antifade Mounting Medium on slides for imaging.

Images were captured with either a Zeiss LSM 900 Confocal microscope or Zeiss Axio

Zoom.V16, both equipped with Zeiss software. To prevent cross talk among channels, images

were obtained using sequential scanning to detect multiple fluorophores. To reduce background

noise and graininess of images, 2x frame-averaging was used and gain was kept at or below 700.

The laser powers used were: EGFP <15% and mCherry or tdTomato <15%. For imaging of

dendritic segments, z-stacks with intervals of 0.2μm were acquired and further analyzed using

Imaris.

To quantify individual and dually innervated neurons, images of dorsomedial NAcSh,

ventromedial NAcSh, ventrolateral NAcSh, and ventromedial NAc core were captured on Zeiss

LSM 900 Confocal microscope at 40x magnification, equipped with Zeiss software. Images were

captured throughout the dorsomedial NAcSh, ventromedial NacSh, ventrolateral NacSh, and

ventromedial NAc core. Cells were then manually counted and recounted a day apart. MSNs were

identified based on soma size and the presence of highly branched, spiny dendrites, as compared

to interneurons, which have larger somas and fewer spines. Analysis and counting were done by

an experimenter blinded to conditions.

Acute brain slice preparation for electrophysiology

Acute coronal slices containing the nucleus accumbens or hippocampus were prepared for

whole-cell patch-clamp electrophysiology. Hipp-containing slices were kept to verify injection sites

before proceeding with electrophysiological experiments in the NAc. Animals were deeply

anesthetized with isoflurane, decapitated, and brains were quickly dissected and submerged in

ice-cold, bubbled (carbogen: 95% O_2/5% CO_2) N-methyl-D-glucamine (NMDG) recovery solution

containing the following (in mM): 93 NMDG, 2.5 KCl, 1.2 NaH_2PO_4, 11 glucose, 25 $NaHCO_3$, 1.2

$MgCl_2$, and 2.4 $CaCl_2$, pH=7.3-7.4, osmolarity=300-310 mOsm. Using a vibratome (VT1000S,

Leica Microsystems), coronal slices (400 μm) were cut in cold, oxygenated NMDG. Slices were

transferred to 32-34°C NMDG for 7-12 minutes to recover and were then transferred to room-

temperature artificial cerebrospinal fluid (aCSF) containing the following (in mM): 120 NaCl, 3 KCl,

1.0 NaH_2PO_4, 20 glucose, 25 $NaHCO_3$, 1.5 $MgCl_2 \cdot 7H_2O$, and 2.5 $CaCl_2$, pH=7.3-7.4. Slices were

allowed to recover for 1-hour at room-temperature before beginning electrophysiological

recordings.

Whole-cell recordings

We performed whole-cell patch-clamp recordings using an Axopatch 200B amplifier (Axon Instruments, Molecular Devices) and a Digidata 1550B digitizer (Axon Instruments). Slices were placed in a submersion-type recording chamber and superfused with room-temperature aCSF (flow rate 0.5-1mL/min). Patch pipettes (4-8MΩ) were made from borosilicate glass (World Precision Instruments) using a Sutter Instruments P-97 model puller. Cells were visualized using a 60x water immersion objective (Nikon Eclipse FN-1). All recordings were performed in voltage-clamp conditions from MSNs in the NAc ventromedial shell.

470nm and 635nm light was used for optogenetic stimulation of Hipp axon terminals in the NAc to record evoked excitatory postsynaptic currents (EPSC). We used light intensities ≤0.5mW to prevent non-specific activation of Chrimson by 470nm light as shown in Figure S1a-b, which aligns with other published studies (Christoffel et al., 2021; Klapoetke et al., 2014; Xia et al., 2020). Similar light intensity ranges were used for 635nm stimulation, and we did not observe non-specific stimulation with this wavelength at any light intensity (Figure S1c-d).

For experiments determining I-V relationship and AMPA:NMDA receptor ratios, the patch pipette solution was composed of 135 mM CsCl, 2 mM MgCl6-H2O, 10 mM HEPES, 4 mM Mg-ATP, 0.3 mM Na2-GTP, 10 mM Na2-phosphocreatine, 1 mM EGTA, 5 mM QX-314 bromide, and 100 µM spermine; pH=7.3-7.4; osmolarity=285-295mOsm. EPSCs were collected from holding potentials ranging from -80mV to +40mV to create an I-V curve. To calculate AMPA:NMDA receptor ratios, the AMPA component was defined by the peak amplitude at -70mV while the NMDA component was defined by the amplitude at +40mV at 50ms after stimulation. Paired pulse stimulation (100ms ISI) of dHipp or vHipp terminals was alternated every 10s to record EPSCs at dHipp-MSN and vHipp-MSN synapses. PPR was determined by calculating 5-minute averages of EPSC amplitudes from our paired pulses (EPSC1 and EPSC2) and dividing EPSC2 by EPSC1 (ie: EPSC2/EPSC1 = PPR). For all other experiments, patch pipettes were filled with a solution

containing 130 mM K-gluconate, 5 mM KCl, 2 mM MgCl6-H2O, 10 mM HEPES, 4 mM Mg-ATP,

0.3 mM Na2-GTP, 10 mM Na2-phosphocreatine, and 1 mM EGTA; pH=7.3-7.4; osmolarity=285-

295mOsm. We used the following exclusion criteria to eliminate unhealthy cells and unreliable

recordings: 1) We only proceeded with experiments on cells with series resistances <10MΩ, 2)

Cells were excluded if their series resistance changed by >20% (comparing the resistance at the

beginning and end of the experiment), 3) Cells in poor health or poor recording status were

excluded (i.e. cell partially or fully sealed up, a decrease in holding current >100pA that is

consistent with the cell dying, an increase in jitter post HFS, and/or an increase in response failure

rate to > 50%).

Visualization of dHipp/vHipp terminals with dually innervated neurons

Using a small subset of slices from electrophysiological experiments, dually innervated MSNs

were patched and filled with CF633 or CF647 dye (Biotium). Upon completion of electrophysiology

experiments, the 400μm-thick slices were fixed in 4% PFA for 4-8 hours at 4ºC. Slices were then

mounted in Invitrogen ProLong Glass Antifade Mountant (Thermo Fisher Scientific, #P36980) on

slides with spacers (400μm-thick) for imaging. Images were captured with a Zeiss LSM 910

confocal microscope equipped with Zeiss software. To prevent cross talk among channels,

images were obtained using sequential scanning to detect multiple fluorophores. To reduce

background noise and graininess of images, 2x frame-averaging was used and gain was kept at

or below 700. The laser powers used were: EGFP <15%, mCherry or tdTomato <15%, CF633

<10%. For imaging of dendritic segments and analysis of synapse distributions, z-stacks (0.2μm

slices) of distal dendrites were acquired. Surface rendering was conducted using Imaris to create

3-dimensional reconstructions of dHipp and vHipp inputs and MSN dendrites. These images were

then analyzed in Imaris to identify putative dHipp-MSN and vHipp-MSN synapses and calculate

Eucledian distances (edge-to-edge) between nearest neighboring dHipp-MSN and vHipp-MSN

synapses along 100μm dendritic segments.

Quantification, statistical analysis, and reproducibility

We found no statistically significant difference between males and females, so data from male and females are plotted together. Both the number of cells and number of mice are reported for each experiment. Plots represent the mean ± SEM. The sample size (n) per condition represents the number of cells unless otherwise indicated in the figure caption. Potentiation scores were calculated as the amplitude of the EPSC evoked from synchronous stimulation divided by the sum of the amplitudes from the alternating stimulations. All statistical analyses were performed using Graphpad Prism 9/10 software. Statistical significance ($p < 0.05$) was assessed using parametric or non-parametric versions of paired t-tests, or one-way or two-way ANOVA with multiple comparisons. Distributions of nearest neighbor distances were compared to a uniform distribution using a Chi-square goodness of fit test. For box plots, the line in the middle of the box is plotted at the median. The box extends from the 25th to 75th percentiles. Whiskers represent minimum and maximum, and individual data points are shown. Figure schematics and experimental timelines were created with BioRender.com.

RESULTS

dHipp and vHipp axon terminal fields overlap in the NAc medial shell

Using viral-mediated labeling to visualize Hipp neurons and their terminal fields, we examined innervation of the NAc by the posterior dHipp and vHipp. We injected an adenoassociated virus (AAV) expressing eYFP into the dHipp and an AAV expressing mCherry into the vHipp (Figure 1a-b). After allowing 5-7 weeks for expression, brains were fixed, sectioned, and imaged using confocal microscopy. We observed dHipp axon terminal fields in the NAc core and lateral shell while the vHipp axon terminal fields were primarily found in the medial shell, in line with previous work (Figure 1c). However, we also observed strong overlap of dHipp and vHipp axon terminal fields in the medial NAcSh (Figure 1c-d), which aligns with the results presented by Brog et al.,

1993. 3-D reconstruction of z-stack images of these overlapping axon fields revealed adjacent

dHipp and vHipp axons (Figure 1d), highlighting the close proximity of dHipp and vHipp afferents

within the medial NAcSh.

Identification of NAc neurons dually innervated by the dHipp and vHipp

Given the apposition of dHipp and vHipp terminals, we sought to determine whether individual

neurons were innervated by both hippocampal regions. To do this, we utilized transsynaptic

expression of Cre and Flp recombinases to selectively target NAc neurons innervated by dHipp

and vHipp pathways (Figure 2a). We injected AAV1-hSyn-Cre into the dHipp and AAV1-Ef1a-

Flpo into the vHipp of Ai195 mice in which GcaMP6s is only expressed in the presence of Cre

and Flp. This approach allowed for targeted expression of GCaMP6s only in neurons that are

innervated by both dHipp and vHipp (Figure 2b). After allowing 5-7 weeks for expression, brains

were fixed, sectioned and imaged with confocal microscopy using GCaMP6s to visualize dually

innervated neurons. Similar to our observations of terminal field overlap, we observed GCaMP6s-

expressing cells in the medial NAcSh (Figure 2c), suggesting dual innervation of NAc neurons by

both dHipp and vHipp. Furthermore, examination of GCaMP6s-expressing cells revealed

morphology similar to MSNs (based on soma size and spiny, highly branched dendrites) (Figure

2d), suggesting that individual NAc MSNs receive input from both dHipp and vHipp.

To gain better insight into the proportions of cells innervated by dHipp, vHipp, or both, we

used our transsynaptic strategy to differentially label cells innervated by dHipp and vHipp.

Specifically, we injected AAV1-hSyn-Cre into the dHipp and AAV1-Ef1a-Flpo into the vHipp of

Ai14 mice in which there is Cre-dependent expression of tdTomato. We also injected AAV9-Ef1a-

fDIO-eYFP into the NAc of these mice (Figure 2e). As a result, cells innervated by dHipp express

tdTomato, cells innervated by vHipp express eYFP, and those innervated by both will express

both reporters (Figure 2f-g). Quantification of cells in medial NAcSh revealed approximately 8%

of cells were innervated by dHipp and vHipp (dually innervated). The remainder of cells were

innervated by one Hipp subregion (singly innervated). Of these, 28% cells were innervated by

dHipp, 65% were innervated by vHipp (Figure 2h). This suggests that a small proportion of

neurons in the medial NAcSh are dually innervated. However, it is worth noting that previous work

has reported low efficiency of transsynaptic transport, leading us to believe that our visualization

strategy underrepresents the number of dually innervated neurons (Zingg et al., 2017).

To independently verify whether individual NAc neurons were dually innervated and determine

whether our anatomical results translated into functional convergence, we used optogenetics and

whole-cell electrophysiology to independently stimulate dHipp and vHipp inputs while recording

from individual NAc neurons. We focused on MSNs because they comprise 95% of the NAc

neuron population, align with the morphology of cells identified by our anatomical tracing studies

(Figure 2), and have been shown to receive functional input from either dHipp or vHipp in separate

studies. To independently stimulate dHipp and vHipp inputs, we used a dual wavelength

optogenetic strategy involving viral expression of Chronos and Chrimson, which are maximally

sensitive to different wavelengths of light (480nm and 660nm, respectively) (Klapoetke et al.,

2014). AAV2-Syn-Chronos-eYFP or AAV2-Syn-Chrimson-mCherry were injected into the dHipp

or vHipp. At least 5 weeks after surgery, coronal brain slices containing the NAc were prepared

for electrophysiology. Using whole-cell patch-clamp electrophysiology, we recorded from MSNs

in the medial NAcSh and used alternating 473 and 635nm light to measure dHipp and vHipp

evoked responses in the MSNs. Keeping the light intensity below 1mW prevented cross-

contamination, allowing us to measure and independently manipulate dHipp and vHipp inputs

with 473 and 635nm light (Figure S1 and 3a). Using this strategy, we identified a subpopulation

of MSNs in the medial NAcSh that responded to both dHipp and vHipp stimulation, demonstrating

dual functional innervation of individual NAc MSNs by both dHipp and vHipp (Figure 3a). Dually

innervated neurons accounted for ~20-50% of neurons surveyed, a greater proportion than that

observed with our transsynaptic labeling (Figure 2). The variation in dually innervated neurons

appeared to depend on recording site, with higher proportions detected in regions that overlapped

with those identified by our labeling approach.

Synaptic properties and strengths influence how a neuron responds to and integrates

information from distinct inputs, leading us to examine and compare the synaptic profiles of dHipp-

MSN and vHipp-MSN synapses in dually innervated neurons. To examine excitatory synaptic

strength, we measured AMPAR:NMDAR ratios at dHipp-MSN and vHipp-MSN synapses and

found no difference, suggesting that the strength of dHipp- and vHipp-MSN synapses is similar

(Figure 3b). We then used paired pulse stimulation, alternating paired pulses such that each

pathway was stimulated every 20s, to examine paired-pulse ratios (PPR). PPR was significantly

greater at vHipp-MSN synapses as compared to dHipp-MSN synapses (Figure 3c), suggesting

potential differences in release probability, though this may be driven by a subpopulation of

recorded neurons.

Calcium-permeable AMPA receptors (CP-AMPAR) have been identified at ventral subiculum-

NAc synapses (Boxer et al., 2023) and have been implicated in NAc plasticity and related

behaviors (Carr, 2020; Mameli et al., 2009; McCutcheon et al., 2011; Wolf & Tseng, 2012). Given

their role in plasticity and behavior, we examined whether CP-AMPARs were present at dHipp-

or vHipp-MSN synapses. Using their characteristic inward rectification and non-linear current-

voltage relationship (Cull-Candy et al., 2006; Liu & Zukin, 2007), we performed

electrophysiological experiments to identify the predominant population of AMPARs at dHipp and

vHipp-MSN synapses. Our results revealed that neither Hipp-MSN synapse contained a

significant population of CP-AMPARs (Figure 3d). Collectively, these results indicate that dHipp

and vHipp synapses in dually innervated NAc MSNs have comparable basal postsynaptic

properties with presynaptic differences in release probability.

We then evaluated the potential for synaptic or dendritic-level interactions between dHipp and

vHipp synapses by comparing rise times and the spatial locations of putative dHipp and vHipp

synapses. Our rise time analysis revealed similar rise times of dHipp- and vHipp-evoked

responses, suggesting that dHipp-MSN and vHipp-MSN synapses are located a similar distance

away from the cell body (Figure 3e). We filled a subset of cells electrophysiologically confirmed

to be dually innervated with a fluorescent dye (Figure 4a), and slices were fixed and prepared for

confocal microscopy. We used confocal microscopy and image analysis to measure the spatial

location of potential dHipp and vHipp synapses along the dendrites of a dually innervated MSN

using the eYFP co-expressed with Chronos, the mCherry co-expressed with Chrimson, and the

dye. From this analysis, we found several examples of dHipp and vHipp puncta that are spatially

close (<2µm) along distal dendrites of dually innervated MSNs (Figure 4b-c). We analyzed 3-

dimensional reconstructions of z-stack images to identify putative dHipp and vHipp inputs on

MSNs dendrites and conducted nearest neighbor analysis to measure the spatial relationships

between dHipp-MSN and vHipp-MSN synapses. We found that many (>50%) dHipp-MSN and

vHipp-MSN synapses are located within a 0-2µm distance (Figure 4d-e), suggesting that dHipp–

MSN and vHipp–MSN synapses cluster preferentially near each other. Altogether, these results

demonstrate functional innervation of individual MSNs by both dHipp and vHipp, where synapses

are frequently observed in close spatial proximity along MSN dendrites.

Heterosynaptic interactions between dHipp and vHipp synapses in NAc MSNs

Synapses that are sufficiently close (<2µm apart) can interact and influence each other's

activity (Fu et al., 2012; Harvey & Svoboda, 2007a; Murakoshi et al., 2011; Xia et al., 2020; Yuste,

2011). Though activity in dHipp and vHipp is not typically temporally coordinated, learning has

been found to increase synchrony between dHipp and vHipp (Biane et al., 2023; Morici et al.,

2025). Synchronous activity has been shown to induce plasticity through heterosynaptic

interactions between adjacent synapses, raising the question of whether the spatial proximity of

dHipp and vHipp synapses facilitates heterosynaptic interactions. To address this, we alternated

473nm and 635nm stimulation for 2 minutes to measure EPSCs evoked from independent

stimulation of each Hipp pathway. We then synchronized 473nm and 635nm stimulations for 2

minutes to record an evoked EPSC from the simultaneous stimulation of both dHipp and vHipp

inputs (Figure 5a,b). If no heterosynaptic interactions occur, then the EPSC amplitudes from the

alternating stimulation should be the linear sum of the EPSC amplitude evoked by simultaneous

stimulation. Instead, our data show a supralinear summation of separately measured EPSCs

(Figure 5b), indicating that heterosynaptic potentiation occurs between dHipp and vHipp

synapses, an effect observed across multiple light intensities in most cells (Figure S2). We

conducted the same experiment in cells that were not dually innervated and found no difference

between EPSC amplitudes evoked by alternating and synchronous stimulation, demonstrating

that this effect is specific to dually innervated neurons and not a nonspecific artifact resulting from

cross-contamination (Figure 5c).

To better understand the site of synaptic interaction, we compared the failure rates in a

separate experiment using minimal stimulation to evoke pathway-specific responses ~50% of the

time so that we could appropriately assess an increase or decrease in failure rate during

synchronous stimulation (Figure 5d). We measured EPSCs evoked from alternating and

synchronous stimulation of dHipp-MSN and vHipp-MSN synapses. We then calculated the

theoretical probability of eliciting a response in both the dHipp- and vHipp-MSN pathways, P_{dual},

if they were stimulated at the same time. Assuming the pathways are independent/non-

interacting, we can calculate this probability as the product of their individual success properties:

$P_{dual}^{theoretical} = P_{dHipp-MSN}^{success} \cdot P_{vHipp-MSN}^{success}$. We compared this theoretical probability to the observed

success rate of sync responses that were classified as dual-evoked responses, finding a

significant difference in the observed and theoretical values (Figure 5e) suggesting presynaptic

mechanisms may facilitate heterosynaptic interactions between dHipp- and vHipp-MSN

synapses.

DISCUSSION

The experiments in this study identified a subpopulation of neurons within the NAc that are dually innervated by dHipp and vHipp. These neurons are primarily located in the NAc ventromedial shell with morphological characteristics that align with those of MSNs. We found that dHipp-MSN and vHipp-MSN synapses in dually innervated neurons have similar basal synaptic properties including comparable release probabilities, synaptic strength, and expression of calcium-impermeable AMPARs. Examination of EPSCs evoked by individual or synchronous stimulation of dHipp and vHipp revealed supralinear summation of EPSCs, suggesting that the synchronous activation of dHipp and vHipp inputs induces short-term heterosynaptic potentiation of dHipp-MSN and vHipp-MSN synapses likely due to presynaptic mechanisms. Altogether, these results reveal mechanisms of dHipp and vHipp synaptic integration within the NAc medial shell, which may be key to supporting the integration of cognitive and emotional processes needed to effectively mediate goal-directed behaviors.

Innervation of the NAc by dHipp and vHipp

Our viral-mediated axon terminal field labeling strategy revealed overlapping dHipp and vHipp innervation patterns within the NAc ventromedial shell. At first glance, this finding seems to contradict previous reports that have suggested that the dHipp and vHipp preferentially innervate distinct parts of the NAc. However, these reported innervation patterns appear to depend upon the anterior/posterior coordinate used for Hipp viral injections. Most studies have targeted the posterior vHipp and anterior dHipp, showing vHipp projections in the NAc core and medial shell, with dHipp projections in the NAc core and lateral shell (Groenewegen et al., 1987; Ibrahim et al., 2024; Li et al., 2018; Mulder et al., 1998; Scofield et al., 2016). But when the posterior dHipp was targeted with an anterograde tracer, dHipp axon terminal field expression was observed in the NAc ventromedial shell (Ibrahim et al., 2024). Furthermore, the injection of a retrograde tracer into the NAc ventromedial shell showed that neurons located in the posterior dHipp and vHipp

were projecting to this area of the NAc (Groenewegen et al., 1987). Our findings align with the

latter two studies, as we injected into the posterior Hipp (A/P: -3.7mm) and found overlapping

dHipp and vHipp axon terminal fields and dually innervated neurons primarily in the NAc

ventromedial shell.

Using a transsynaptic viral-mediated labeling technique, we were able to visualize a small

subpopulation of dually innervated neurons that had morphological characteristics similar to

MSNs (Figure 2). We confirmed this electrophysiologically, demonstrating that dHipp and vHipp

inputs form synaptic connections with individual MSNs within the medial NAcSh (Figure 3).

Because this labeling method has low transsynaptic efficiency (Zingg et al., 2017) and requires

successful transfer of both Cre and Flp payloads, it likely underrepresents the number of dually

innervated neurons. This is supported by our electrophysiology recordings, as within a single

slice, we were able to record from a larger number of dually innervated neurons than were

indicated from the anatomical tracing study. Further optimization of this transsynaptic approach

could improve detection and determine whether sparsely distributed NAc interneurons also

receive dual hippocampal input.

dHipp and vHipp synapses have similar properties in dually innervated NAc neurons

We found that dHipp- and vHipp-MSN synapses contain primarily calcium-impermeable

AMPARs and have similar excitatory synaptic strengths, as measured by AMPA:NMDA receptor

ratios (Figure 3). These similarities suggest that, in basal conditions, both pathways may

contribute equally to downstream circuit activity. Since the dHipp and vHipp convey different types

of behavioral information, this may mean that Hipp-related contextual and sensory cues are

equally capable of influencing motivated behaviors. While these similarities are present at

baseline, several studies have highlighted changes that occur to dHipp-MSN and vHipp-MSN

synapses in response to hormones, neuromodulators, cocaine, external stress, and experience

(Floresco et al., 2001; LeGates et al., 2018; Peleg-Raibstein & Feldon, 2006; R. Yang & J.

Mogenson, 1984; Sjulson et al., 2018; Williams et al., 2020).

Synaptic interactions between distinct Hipp inputs

The NAc is a critical integrator of several major excitatory inputs, many of which exhibit

coordinated or synchronous activation (Mahon et al., 2006; Stern et al., 1998). While dHipp and

vHipp activity is not typically temporally coordinated, increased synchrony has been associated

with learning (Biane et al., 2023; Morici et al., 2025). We found that dHipp-MSN and vHipp-MSN

synaptic interactions during simultaneous stimulation elicit heterosynaptic potentiation (Figure 5).

Convergent afferent activity and heterosynaptic interactions can alter plasticity thresholds or

propagate plastic changes across synapses, fundamentally influencing neuronal information

processing and storage (Chistiakova et al., 2015; Reissner et al., 2010). Several mechanisms

underlying heterosynaptic plasticity have been identified, including neuromodulator-dependent

forms, synaptic tagging and capture, and spatially clustered plasticity, each engaging distinct

molecular pathways that ultimately promote plasticity across synapses (Bailey et al., 2000b;

Govindarajan et al., 2006b; Harvey & Svoboda, 2007b; Redondo & Morris, 2011b; Schuman &

Madison, 1994b). Within the NAc, convergent inputs interact at the synaptic level via a

combination of presynaptic and postsynaptic mechanisms to influence behavioral output

(O'Donnell & Grace, 1995; Xia et al., 2020; F.-C. Yang & Liang, 2014; Yu et al., 2022). Our

analysis of observed and theoretical success rates suggests that dHipp/vHipp heterosynaptic

potentiation may occur presynaptically (Figure 5). Axo-axonic synapses in the striatum are rare,

but recent work has highlighted heterosynaptic interactions between the basolateral amygdala

(BLA), medial prefrontal cortex (mPFC) and paraventricular nucleus of the thalamus (PVT) during

co-activation that occur due to presynaptic innervation of mPFC and PVT by BLA (Kornhuber &

Kornhuber, 1983; Xia et al., 2020). Additionally, co-activation of vHipp-MSN and BLA-MSN

synapses induces long-term potentiation that requires the activity of dopaminergic terminals and

D1 receptors (Yu et al., 2022). While this may be due to activity of postsynaptic receptors given

that this form of plasticity is more prominent on D1-expressing MSNs, D1 receptors are also

expressed presynaptically on glutamatergic terminals and astrocytes where their signaling can

alter release probability (Corkrum et al., 2020; Nicola & Malenka, 1997; Zhang et al., 2014).

Implications of integration that occurs at the synaptic level

The connection between the Hipp and NAc is crucial for integrating reward with related

contextual and spatial information (Britt et al., 2012; Floresco & Phillips, 1999; Humphries &

Prescott, 2010; Ito et al., 2008; Lansink et al., 2008, 2009; LeGates et al., 2018; Pennartz et al.,

2011; Sjulson et al., 2018). Functional imaging studies in humans have revealed greater Hipp-

NAc connectivity associated with positive reinforcement learning (Davidow et al., 2016), increased

reward seeking in adolescents (Huntley et al., 2020), and greater positive affect in adults (Heller

et al., 2020), implicating a strong link between the strength of Hipp-NAc connectivity and reward-

related behaviors. Similar findings have been demonstrated in rodent models which examined

Hipp-NAc connectivity using *in vivo* electrophysiology during spatial and contextual-based reward

tasks (Sjulson et al., 2018; Sosa et al., 2020; Trouche et al., 2019). The causal relationship

between Hipp-NAc connectivity and reward behaviors has been established with lesion and

optogenetic studies, demonstrating that Hipp-NAc synapses are required for contextual- and

spatial-based reward tasks (Floresco & Phillips, 1999; Ito et al., 2008, p. 200; LeGates et al.,

2018; Trouche et al., 2019; Zhou et al., 2019). This includes our previously published work, which

demonstrated that the strength of vHipp-NAc synapses is a critical mediator of reward-related

behaviors (LeGates et al., 2018).

dHipp and vHipp convergence within other areas of the brain has been proposed to be a key

circuit mechanism connecting cognitive and emotional processes (Fanselow & Dong, 2010) and

may be crucial to establishing and retrieving memories. The dHipp and vHipp are known to convey

distinct information critical for driving NAc-mediated motivated behaviors (Ibrahim et al., 2024; Ito

et al., 2008; LeGates et al., 2018; Muir et al., 2020; Patterson et al., 2025; Sjulson et al., 2018;

Sosa et al., 2020; Trouche et al., 2019; Williams et al., 2020; A. K. Yang et al., 2020; F.-C. Yang

& Liang, 2014). Therefore, the convergence and plasticity we observed may provide a basis by

which distinct information conveyed by dHipp and vHipp can shapes the NAc's response to more

nuanced contextual and associative information. Similar convergence has been observed in the

amygdala, where dHipp and vHipp activate and reactivate, respectively, specific neuron

ensembles to mediate contextual fear memory (Terranova et al., 2022). Since Hipp input exerts

a strong influence over NAc neuron activity and information flow, it is likely these interactions

encourage integration of distinct information conveyed by the dHipp and vHipp, critically impacting

behavioral outputs.

Altogether, the results presented here provide evidence for integration of dHipp and vHipp

information within individual MSNs of the NAc ventromedial shell. The combination of techniques

enabling identification of similar synaptic profiles and rapid interactions occurring between dHipp-

and vHipp-MSN synapses provides a foundation for similar studies investigating the integration

of distinct inputs at the level of individual neurons. Ultimately, these findings advance our

understanding of how spatial and contextual information may contribute to reward-related

behaviors while also raising further questions regarding how each of these distinct pathways are

regulated.

References

Amaral, D. G., & Witter, M. P. (1989). The three-dimensional organization of the hippocampal

formation: A review of anatomical data. *Neuroscience*, *31*(3), 571–591.

https://doi.org/10.1016/0306-4522(89)90424-7

Bailey, C. H., Giustetto, M., Huang, Y. Y., Hawkins, R. D., & Kandel, E. R. (2000a). Is

heterosynaptic modulation essential for stabilizing Hebbian plasticity and memory?

Nature Reviews. Neuroscience, *1*(1), 11–20. https://doi.org/10.1038/35036191

Bailey, C. H., Giustetto, M., Huang, Y.-Y., Hawkins, R. D., & Kandel, E. R. (2000b). Is

Heterosynaptic modulation essential for stabilizing hebbian plasiticity and memory.

Nature Reviews Neuroscience, *1*(1), 11–20. https://doi.org/10.1038/35036191

Bannerman, D. M., Deacon, R. M. J., Offen, S., Friswell, J., Grubb, M., & Rawlins, J. N. P.

(2002). Double dissociation of function within the hippocampus: Spatial memory and

hyponeophagia. *Behavioral Neuroscience*, *116*(5), 884–901.

https://doi.org/10.1037/0735-7044.116.5.884

Biane, J. S., Ladow, M. A., Stefanini, F., Boddu, S. P., Fan, A., Hassan, S., Dundar, N.,

Apodaca-Montano, D. L., Zhou, L. Z., Fayner, V., Woods, N. I., & Kheirbek, M. A. (2023).

Neural dynamics underlying associative learning in the dorsal and ventral hippocampus.

Nature Neuroscience, *26*(5), 798–809. https://doi.org/10.1038/s41593-023-01296-6

Boxer, E. E., Kim, J., Dunn, B., & Aoto, J. (2023). Ventral Subiculum Inputs to Nucleus

Accumbens Medial Shell Preferentially Innervate D2R Medium Spiny Neurons and

Contain Calcium Permeable AMPARs. *The Journal of Neuroscience*, *43*(7), 1166–1177.

https://doi.org/10.1523/JNEUROSCI.1907-22.2022

Britt, J. P., Benaliouad, F., McDevitt, R. A., Stuber, G. D., Wise, R. A., & Bonci, A. (2012).

Synaptic and Behavioral Profile of Multiple Glutamatergic Inputs to the Nucleus

Accumbens. *Neuron*, *76*(4), 790–803. https://doi.org/10.1016/j.neuron.2012.09.040

Brog, J. S., Salyapongse, A., Deutch, A. Y., & Zahm, D. S. (1993). The patterns of afferent

innervation of the core and shell in the "Accumbens" part of the rat ventral striatum:

Immunohistochemical detection of retrogradely transported fluoro-gold. *Journal of*

Comparative Neurology, *338*(2), 255–278. https://doi.org/10.1002/cne.903380209

Calhoon, G. G., & O'Donnell, P. (2013). Closing the Gate in the Limbic Striatum: Prefrontal

Suppression of Hippocampal and Thalamic Inputs. *Neuron*, *78*(1), 181–190.

https://doi.org/10.1016/j.neuron.2013.01.032

Carr, K. D. (2020). Homeostatic regulation of reward via synaptic insertion of calcium-permeable

AMPA receptors in nucleus accumbens. *Physiology & Behavior*, *219*, 112850.

https://doi.org/10.1016/j.physbeh.2020.112850

Chistiakova, M., Bannon, N., Chen, J.-Y., Bazhenov, M., & Volgushev, M. (2015). Homeostatic

role of heterosynaptic plasticity: Models and experiments. *Frontiers in Computational*

Neuroscience, *9*. https://www.frontiersin.org/articles/10.3389/fncom.2015.00089

Christoffel, D. J., Walsh, J. J., Hoerbelt, P., Heifets, B. D., Llorach, P., Lopez, R. C.,

Ramakrishnan, C., Deisseroth, K., & Malenka, R. C. (2021). Selective filtering of

excitatory inputs to nucleus accumbens by dopamine and serotonin. *Proceedings of the*

National Academy of Sciences, *118*(24), e2106648118.

https://doi.org/10.1073/pnas.2106648118

Corkrum, M., Covelo, A., Lines, J., Bellocchio, L., Pisansky, M., Loke, K., Quintana, R.,

Rothwell, P. E., Lujan, R., Marsicano, G., Martin, E. D., Thomas, M. J., Kofuji, P., &

Araque, A. (2020). Dopamine-Evoked Synaptic Regulation in the Nucleus Accumbens

Requires Astrocyte Activity. *Neuron*, *105*(6), 1036-1047.e5.

https://doi.org/10.1016/j.neuron.2019.12.026

Cull-Candy, S., Kelly, L., & Farrant, M. (2006). Regulation of Ca2+-permeable AMPA receptors:

Synaptic plasticity and beyond. *Current Opinion in Neurobiology*, *16*(3), 288–297.

https://doi.org/10.1016/j.conb.2006.05.012

Davidow, J. Y., Foerde, K., Galván, A., & Shohamy, D. (2016). An Upside to Reward Sensitivity:

The Hippocampus Supports Enhanced Reinforcement Learning in Adolescence.

Neuron, *92*(1), 93–99. https://doi.org/10.1016/j.neuron.2016.08.031

Duarte-Guterman, P., Yagi, S., Chow, C., & Galea, L. A. M. (2015). Hippocampal learning,

memory, and neurogenesis: Effects of sex and estrogens across the lifespan in adults.

Hormones and Behavior, *74*, 37–52. https://doi.org/10.1016/j.yhbeh.2015.05.024

Fanselow, M. S., & Dong, H.-W. (2010). Are The Dorsal and Ventral Hippocampus functionally

distinct structures? *Neuron*, *65*(1), 7. https://doi.org/10.1016/j.neuron.2009.11.031

Floresco, S. B., Blaha, C. D., Yang, C. R., & Phillips, A. G. (2001). Modulation of Hippocampal

and Amygdalar-Evoked Activity of Nucleus Accumbens Neurons by Dopamine: Cellular

Mechanisms of Input Selection. *The Journal of Neuroscience*, *21*(8), 2851–2860.

https://doi.org/10.1523/JNEUROSCI.21-08-02851.2001

Floresco, S. B., & Phillips, A. G. (1999). Dopamine and hippocampal input to the nucleus

accumbens play an essential role in the search for food in an unpredictable environment.

Psychobiology, *27*(2), 277–286. https://doi.org/10.3758/BF03332121

Francis, T. C., & Lobo, M. K. (2017). Emerging Role for Nucleus Accumbens Medium Spiny

Neuron Subtypes in Depression. *Biological Psychiatry*, *81*(8), 645–653.

https://doi.org/10.1016/j.biopsych.2016.09.007

French, S. J., & Totterdell, S. (2002). Hippocampal and prefrontal cortical inputs

monosynaptically converge with individual projection neurons of the nucleus accumbens.

The Journal of Comparative Neurology, *446*(2), 151–165.

https://doi.org/10.1002/cne.10191

Friedman, D. P., Aggleton, J. P., & Saunders, R. C. (2002). Comparison of hippocampal,

amygdala, and perirhinal projections to the nucleus accumbens: Combined anterograde

and retrograde tracing study in the Macaque brain. *Journal of Comparative Neurology*,

450(4), 345–365. https://doi.org/10.1002/cne.10336

585 Fu, M., Yu, X., Lu, J., & Zuo, Y. (2012). Repetitive motor learning induces coordinated formation

586   of clustered dendritic spines in vivo. *Nature*, *483*(7387), 92–95.

587   https://doi.org/10.1038/nature10844

588 Govindarajan, A., Kelleher, R. J., & Tonegawa, S. (2006a). A clustered plasticity model of long-

589   term memory engrams. *Nature Reviews. Neuroscience*, *7*(7), 575–583.

590   https://doi.org/10.1038/nrn1937

591 Govindarajan, A., Kelleher, R. J., & Tonegawa, S. (2006b). A clustered plasticity model of long-

592   term memory engrams. *Nature Reviews Neuroscience*, *7*(7), 575–583.

593   https://doi.org/10.1038/nrn1937

594 Groenewegen, H. J., Der Zee, E. V.-V., Te Kortschot, A., & Witter, M. P. (1987). Organization of

595   the projections from the subiculum to the ventral striatum in the rat. A study using

596   anterograde transport of Phaseolus vulgaris leucoagglutinin. *Neuroscience*, *23*(1), 103–

597   120. https://doi.org/10.1016/0306-4522(87)90275-2

598 Gruber, A. J., Hussain, R. J., & O'Donnell, P. (2009). The Nucleus Accumbens: A Switchboard

599   for Goal-Directed Behaviors. *PLoS ONE*, *4*(4), e5062.

600   https://doi.org/10.1371/journal.pone.0005062

601 Harvey, C. D., & Svoboda, K. (2007a). Locally dynamic synaptic learning rules in pyramidal

602   neuron dendrites. *Nature*, *450*(7173), 1195–1200. https://doi.org/10.1038/nature06416

603 Harvey, C. D., & Svoboda, K. (2007b). Locally dynamic synaptic learning rules in pyramidal

604   neuron dendrites. *Nature*, *450*(7173), 1195–1200. https://doi.org/10.1038/nature06416

605 Heller, A. S., Shi, T. C., Ezie, C. E. C., Reneau, T. R., Baez, L. M., Gibbons, C. J., & Hartley, C.

606   A. (2020). Association between real-world experiential diversity and positive affect

607   relates to hippocampal–striatal functional connectivity. *Nature Neuroscience*, *23*(7), 800–

608   804. https://doi.org/10.1038/s41593-020-0636-4

Humphries, M. D., & Prescott, T. J. (2010). The ventral basal ganglia, a selection mechanism at

the crossroads of space, strategy, and reward. *Progress in Neurobiology*, *90*(4), 385–

417. https://doi.org/10.1016/j.pneurobio.2009.11.003

Huntley, E. D., Marusak, H. A., Berman, S. E., Zundel, C. G., Hatfield, J. R. B., Keating, D. P., &

Rabinak, C. A. (2020). Adolescent substance use and functional connectivity between

the ventral striatum and hippocampus. *Behavioural Brain Research*, *390*, 112678.

https://doi.org/10.1016/j.bbr.2020.112678

Ibrahim, K. M., Massaly, N., Yoon, H.-J., Sandoval, R., Widman, A. J., Heuermann, R. J.,

Williams, S., Post, W., Pathiranage, S., Lintz, T., Zec, A., Park, A., Yu, W., Kash, T. L.,

Gereau, R. W., & Morón, J. A. (2024). Dorsal hippocampus to nucleus accumbens

projections drive reinforcement via activation of accumbal dynorphin neurons. *Nature*

Communications, *15*(1), 750. https://doi.org/10.1038/s41467-024-44836-9

Ishizuka, N., Weber, J., & Amaral, D. G. (1990). Organization of intrahippocampal projections

originating from CA3 pyramidal cells in the rat. *Journal of Comparative Neurology*,

295(4), 580–623. https://doi.org/10.1002/cne.902950407

Ito, R., Robbins, T. W., Pennartz, C. M., & Everitt, B. J. (2008). Functional Interaction between

the Hippocampus and Nucleus Accumbens Shell Is Necessary for the Acquisition of

Appetitive Spatial Context Conditioning. *Journal of Neuroscience*, *28*(27), 6950–6959.

https://doi.org/10.1523/JNEUROSCI.1615-08.2008

Kelley, A. E., & Domesick, V. B. (1982). The distribution of the projection from the hippocampal

formation to the nucleus accumbens in the rat: An anterograde and retrograde-

horseradish peroxidase study. *Neuroscience*, *7*(10), 2321–2335.

https://doi.org/10.1016/0306-4522(82)90198-1

Kjelstrup, K. B., Solstad, T., Brun, V. H., Hafting, T., Leutgeb, S., Witter, M. P., Moser, E. I., &

Moser, M.-B. (2008). Finite Scale of Spatial Representation in the Hippocampus.

Science, *321*(5885), 140–143. https://doi.org/10.1126/science.1157086

Kjelstrup, K. G., Tuvnes, F. A., Steffenach, H.-A., Murison, R., Moser, E. I., & Moser, M.-B.

(2002). Reduced fear expression after lesions of the ventral hippocampus. *Proceedings*

of the National Academy of Sciences, *99*(16), 10825–10830.

https://doi.org/10.1073/pnas.152112399

Klapoetke, N. C., Murata, Y., Kim, S. S., Pulver, S. R., Birdsey-Benson, A., Cho, Y. K.,

Morimoto, T. K., Chuong, A. S., Carpenter, E. J., Tian, Z., Wang, J., Xie, Y., Yan, Z.,

Zhang, Y., Chow, B. Y., Surek, B., Melkonian, M., Jayaraman, V., Constantine-Paton,

M., … Boyden, E. S. (2014). Independent optical excitation of distinct neural populations.

Nature Methods, *11*(3), 338–346. https://doi.org/10.1038/nmeth.2836

Kornhuber, J., & Kornhuber, M. E. (1983). Axo-axonic synapses in the rat striatum. *European*

Neurology, *22*(6), 433–436. https://doi.org/10.1159/000115598

Lamsa, K., & Lau, P. (2019). Long-term plasticity of hippocampal interneurons during in vivo

memory processes. *Current Opinion in Neurobiology*, *54*, 20–27.

https://doi.org/10.1016/j.conb.2018.08.006

Lansink, C. S., Goltstein, P. M., Lankelma, J. V., Joosten, R. N. J. M. A., McNaughton, B. L., &

Pennartz, C. M. A. (2008). Preferential Reactivation of Motivationally Relevant

Information in the Ventral Striatum. *Journal of Neuroscience*, *28*(25), 6372–6382.

https://doi.org/10.1523/JNEUROSCI.1054-08.2008

Lansink, C. S., Goltstein, P. M., Lankelma, J. V., McNaughton, B. L., & Pennartz, C. M. A.

(2009). Hippocampus Leads Ventral Striatum in Replay of Place-Reward Information.

PLoS Biology, *7*(8), e1000173. https://doi.org/10.1371/journal.pbio.1000173

LeGates, T. A., Kvarta, M. D., Tooley, J. R., Francis, T. C., Lobo, M. K., Creed, M. C., &

Thompson, S. M. (2018). Reward behaviour is regulated by the strength of

hippocampus–nucleus accumbens synapses. *Nature*, *564*(7735), 258–262.

https://doi.org/10.1038/s41586-018-0740-8

Li, Z., Chen, Z., Fan, G., Li, A., Yuan, J., & Xu, T. (2018). Cell-Type-Specific Afferent

Innervation of the Nucleus Accumbens Core and Shell. *Frontiers in Neuroanatomy*, *12*,

84. https://doi.org/10.3389/fnana.2018.00084

Liu, S. J., & Zukin, R. S. (2007). Ca2+-permeable AMPA receptors in synaptic plasticity and

neuronal death. *Trends in Neurosciences*, *30*(3), 126–134.

https://doi.org/10.1016/j.tins.2007.01.006

Mahon, S., Vautrelle, N., Pezard, L., Slaght, S. J., Deniau, J.-M., Chouvet, G., & Charpier, S.

(2006). Distinct Patterns of Striatal Medium Spiny Neuron Activity during the Natural

Sleep–Wake Cycle. *Journal of Neuroscience*, *26*(48), 12587–12595.

https://doi.org/10.1523/JNEUROSCI.3987-06.2006

Mameli, M., Halbout, B., Creton, C., Engblom, D., Parkitna, J. R., Spanagel, R., & Lüscher, C.

(2009). Cocaine-evoked synaptic plasticity: Persistence in the VTA triggers adaptations

in the NAc. *Nature Neuroscience*, *12*(8), 1036–1041. https://doi.org/10.1038/nn.2367

Mannella, F., Gurney, K., & Baldassarre, G. (2013). The nucleus accumbens as a nexus

between values and goals in goal-directed behavior: A review and a new hypothesis.

Frontiers in Behavioral Neuroscience, *7*. https://doi.org/10.3389/fnbeh.2013.00135

Marinescu, A.-M., & Labouesse, M. A. (2024). The nucleus accumbens shell: A neural hub at

the interface of homeostatic and hedonic feeding. *Frontiers in Neuroscience*, *18*,

1437210. https://doi.org/10.3389/fnins.2024.1437210

McCutcheon, J. E., Wang, X., Tseng, K. Y., Wolf, M. E., & Marinelli, M. (2011). Calcium-

Permeable AMPA Receptors Are Present in Nucleus Accumbens Synapses after

Prolonged Withdrawal from Cocaine Self-Administration But Not Experimenter-

Administered Cocaine. *The Journal of Neuroscience*, *31*(15), 5737–5743.

https://doi.org/10.1523/JNEUROSCI.0350-11.2011

Morici, J. F., Paiva-Lima, I., Silva, A., Pronier, É., & Girardeau, G. (2025). *Dorso-ventral*

hippocampus neural assemblies reactivate during sleep following an aversive

experience (p. 2024.06.13.598918). bioRxiv. https://doi.org/10.1101/2024.06.13.598918

Muir, J., Tse, Y. C., Iyer, E. S., Biris, J., Cvetkovska, V., Lopez, J., & Bagot, R. C. (2020).

Ventral Hippocampal Afferents to Nucleus Accumbens Encode Both Latent Vulnerability

and Stress-Induced Susceptibility. *Biological Psychiatry*, *88*(11), 843–854.

https://doi.org/10.1016/j.biopsych.2020.05.021

Mulder, A. B., Hodenpijl, M. G., & Lopes Da Silva, F. H. (1998). Electrophysiology of the

Hippocampal and Amygdaloid Projections to the Nucleus Accumbens of the Rat:

Convergence, Segregation, and Interaction of Inputs. *The Journal of Neuroscience*,

18(13), 5095–5102. https://doi.org/10.1523/JNEUROSCI.18-13-05095.1998

Murakoshi, H., Wang, H., & Yasuda, R. (2011). Local, persistent activation of Rho GTPases

during plasticity of single dendritic spines. *Nature*, *472*(7341), 100–104.

https://doi.org/10.1038/nature09823

Nicola, S. M., & Malenka, R. C. (1997). Dopamine Depresses Excitatory and Inhibitory Synaptic

Transmission by Distinct Mechanisms in the Nucleus Accumbens. *Journal of*

Neuroscience, *17*(15), 5697–5710. https://doi.org/10.1523/JNEUROSCI.17-15-

05697.1997

O'Donnell, P., & Grace, A. (1995). Synaptic interactions among excitatory afferents to nucleus

accumbens neurons: Hippocampal gating of prefrontal cortical input. *The Journal of*

Neuroscience, *15*(5), 3622–3639. https://doi.org/10.1523/JNEUROSCI.15-05-

03622.1995

Patterson, D., Khan, N., Collins, E. A., Stewart, N. R., Sassaninejad, K., Yeates, D., Lee, A. C.

H., & Ito, R. (2025). Ventral hippocampus to nucleus accumbens shell circuit regulates

approach decisions during motivational conflict. *PLOS Biology*, *23*(1), e3002722.

https://doi.org/10.1371/journal.pbio.3002722

Paxinos, G., & Franklin, K. B. J. (2019). *Paxinos and Franklin's The mouse brain in stereotaxic*

coordinates (Fifth edition). Academic Press, an imprint of Elsevier.

Peleg-Raibstein, D., & Feldon, J. (2006). Effects of dorsal and ventral hippocampal NMDA

stimulation on nucleus accumbens core and shell dopamine release.

Neuropharmacology, *51*(5), 947–957. https://doi.org/10.1016/j.neuropharm.2006.06.002

Pennartz, C. M. A., Ito, R., Verschure, P. F. M. J., Battaglia, F. P., & Robbins, T. W. (2011). The

hippocampal–striatal axis in learning, prediction and goal-directed behavior. *Trends in*

Neurosciences, *34*(10), 548–559. https://doi.org/10.1016/j.tins.2011.08.001

Phillipson, O. T., & Griffiths, A. C. (1985). The topographic order of inputs to nucleus

accumbens in the rat. *Neuroscience*, *16*(2), 275–296. https://doi.org/10.1016/0306-

4522(85)90002-8

R. Yang, C., & J. Mogenson, G. (1984). Electrophysiological responses of neurones in the

nucleus accumbens to hippocampal stimulation and the attenuation of the excitatory

responses by the mesolimbic dopaminergic system. *Brain Research*, *324*(1), 69–84.

https://doi.org/10.1016/0006-8993(84)90623-1

Redondo, R. L., & Morris, R. G. M. (2011a). Making memories last: The synaptic tagging and

capture hypothesis. *Nature Reviews. Neuroscience*, *12*(1), 17–30.

https://doi.org/10.1038/nrn2963

Redondo, R. L., & Morris, R. G. M. (2011b). Making memories last: The synaptic tagging and

capture hypothesis. *Nature Reviews Neuroscience*, *12*(1), 17–30.

https://doi.org/10.1038/nrn2963

Reissner, K. J., Pu, L., Schaffhausen, J. H., Boyle, H. D., Smith, I. F., Parker, I., & Carew, T. J.

(2010). A Novel Postsynaptic Mechanism for Heterosynaptic Sharing of Short-Term

Plasticity. *Journal of Neuroscience*, *30*(26), 8797–8806.

https://doi.org/10.1523/JNEUROSCI.4767-09.2010

Roberts, A. C., Tomic, D. L., Parkinson, C. H., Roeling, T. A., Cutter, D. J., Robbins, T. W., &

Everitt, B. J. (2007). Forebrain connectivity of the prefrontal cortex in the marmoset

monkey (*Callithrix jacchus*): An anterograde and retrograde tract-tracing study. *Journal*

of Comparative Neurology, *502*(1), 86–112. https://doi.org/10.1002/cne.21300

Safari, S., Ahmadi, N., Mohammadkhani, R., Ghahremani, R., Khajvand-Abedeni, M., Shahidi,

S., Komaki, A., Salehi, I., & Karimi, S. A. (2021). Sex differences in spatial learning and

memory and hippocampal long-term potentiation at perforant pathway-dentate gyrus

(PP-DG) synapses in Wistar rats. *Behavioral and Brain Functions*, *17*(1), 9.

https://doi.org/10.1186/s12993-021-00184-y

Salimpoor, V. N., Van Den Bosch, I., Kovacevic, N., McIntosh, A. R., Dagher, A., & Zatorre, R.

J. (2013). Interactions Between the Nucleus Accumbens and Auditory Cortices Predict

Music Reward Value. *Science*, *340*(6129), 216–219.

https://doi.org/10.1126/science.1231059

Savage, L. (2004). The effects of hippocampal lesions on learning, memory, and reward

expectancies. *Neurobiology of Learning and Memory*, *82*(2), 109–119.

https://doi.org/10.1016/j.nlm.2004.05.002

Schuman, E. M., & Madison, D. V. (1994a). Locally distributed synaptic potentiation in the

hippocampus. *Science (New York, N.Y.)*, *263*(5146), 532–536.

https://doi.org/10.1126/science.8290963

Schuman, E. M., & Madison, D. V. (1994b). Locally distributed synaptic potentiation in the

hippocampus. *Science (New York, N.Y.)*, *263*(5146), 532–536.

https://doi.org/10.1126/science.8290963

Scofield, M. D., Heinsbroek, J. A., Gipson, C. D., Kupchik, Y. M., Spencer, S., Smith, A. C. W.,

Roberts-Wolfe, D., & Kalivas, P. W. (2016). The Nucleus Accumbens: Mechanisms of

Addiction across Drug Classes Reflect the Importance of Glutamate Homeostasis.

Pharmacological Reviews, *68*(3), 816–871. https://doi.org/10.1124/pr.116.012484

Sjulson, L., Peyrache, A., Cumpelik, A., Cassataro, D., & Buzsáki, G. (2018). Cocaine Place

762   Conditioning Strengthens Location-Specific Hippocampal Coupling to the Nucleus

763   Accumbens. *Neuron*, *98*(5), 926-934.e5. https://doi.org/10.1016/j.neuron.2018.04.015

764   Soares-Cunha, C., de Vasconcelos, N. A. P., Coimbra, B., Domingues, A. V., Silva, J. M.,

765   Loureiro-Campos, E., Gaspar, R., Sotiropoulos, I., Sousa, N., & Rodrigues, A. J. (2020).

766   Nucleus accumbens medium spiny neurons subtypes signal both reward and aversion.

767   *Molecular Psychiatry*, *25*(12), 3241–3255. https://doi.org/10.1038/s41380-019-0484-3

768   Sosa, M., Joo, H. R., & Frank, L. M. (2020). Dorsal and Ventral Hippocampal Sharp-Wave

769   Ripples Activate Distinct Nucleus Accumbens Networks. *Neuron*, *105*(4), 725-741.e8.

770   https://doi.org/10.1016/j.neuron.2019.11.022

771   Squire, L. R. (1992). Memory and the hippocampus: A synthesis from findings with rats,

772   monkeys, and humans. *Psychological Review*, *99*(2), 195–231.

773   https://doi.org/10.1037/0033-295X.99.2.195

774   Stern, E. A., Jaeger, D., & Wilson, C. J. (1998). Membrane potential synchrony of

775   simultaneously recorded striatal spiny neuronsin vivo. *Nature*, *394*(6692), 475–478.

776   https://doi.org/10.1038/28848

777   Strange, B. A., Witter, M. P., Lein, E. S., & Moser, E. I. (2014). Functional organization of the

778   hippocampal longitudinal axis. *Nature Reviews Neuroscience*, *15*(10), 655–669.

779   https://doi.org/10.1038/nrn3785

780   Swanson, L. W., & Cowan, W. M. (1977). An autoradiographic study of the organization of the

781   efferet connections of the hippocampal formation in the rat. *Journal of Comparative*

782   *Neurology*, *172*(1), 49–84. https://doi.org/10.1002/cne.901720104

783   Tao, S., Wang, Y., Peng, J., Zhao, Y., He, X., Yu, X., Liu, Q., Jin, S., & Xu, F. (2021). Whole-

784   Brain Mapping the Direct Inputs of Dorsal and Ventral CA1 Projection Neurons. *Frontiers*

785   *in Neural Circuits*, *15*, 643230. https://doi.org/10.3389/fncir.2021.643230

Terranova, J. I., Yokose, J., Osanai, H., Marks, W. D., Yamamoto, J., Ogawa, S. K., & Kitamura,

T. (2022). Hippocampal-amygdala memory circuits govern experience-dependent

observational fear. *Neuron*, *110*(8), 1416-1431.e13.

https://doi.org/10.1016/j.neuron.2022.01.019

Trouche, S., Koren, V., Doig, N. M., Ellender, T. J., El-Gaby, M., Lopes-dos-Santos, V., Reeve,

H. M., Perestenko, P. V., Garas, F. N., Magill, P. J., Sharott, A., & Dupret, D. (2019). A

Hippocampus-Accumbens Tripartite Neuronal Motif Guides Appetitive Memory in Space.

Cell, *176*(6), 1393-1406.e16. https://doi.org/10.1016/j.cell.2018.12.037

Williams, E. S., Manning, C. E., Eagle, A. L., Swift-Gallant, A., Duque-Wilckens, N.,

Chinnusamy, S., Moeser, A., Jordan, C., Leininger, G., & Robison, A. J. (2020).

Androgen-Dependent Excitability of Mouse Ventral Hippocampal Afferents to Nucleus

Accumbens Underlies Sex-Specific Susceptibility to Stress. *Biological Psychiatry*, *87*(6),

492–501. https://doi.org/10.1016/j.biopsych.2019.08.006

Wolf, M. E., & Tseng, K. Y. (2012). Calcium-permeable AMPA receptors in the VTA and nucleus

accumbens after cocaine exposure: When, how, and why? *Frontiers in Molecular*

Neuroscience, *5*. https://doi.org/10.3389/fnmol.2012.00072

Xia, S., Yu, J., Huang, X., Sesack, S. R., Huang, Y. H., Schlüter, O. M., Cao, J.-L., & Dong, Y.

(2020). Cortical and Thalamic Interaction with Amygdala-to-Accumbens Synapses. *The*

Journal of Neuroscience, *40*(37), 7119–7132. https://doi.org/10.1523/JNEUROSCI.1121-

20.2020

Yang, A. K., Mendoza, J. A., Lafferty, C. K., Lacroix, F., & Britt, J. P. (2020). Hippocampal Input

to the Nucleus Accumbens Shell Enhances Food Palatability. *Biological Psychiatry*,

87(7), 597–608. https://doi.org/10.1016/j.biopsych.2019.09.007

Yang, F.-C., & Liang, K. C. (2014). Interactions of the dorsal hippocampus, medial prefrontal

cortex and nucleus accumbens in formation of fear memory: Difference in inhibitory

avoidance learning and contextual fear conditioning. *Neurobiology of Learning and*

Memory, *112*, 186–194. https://doi.org/10.1016/j.nlm.2013.07.017

Yu, J., Sesack, S. R., Huang, Y., Schlüter, O. M., Grace, A. A., & Dong, Y. (2022). Contingent

Amygdala Inputs Trigger Heterosynaptic LTP at Hippocampus-To-Accumbens

Synapses. *The Journal of Neuroscience*, *42*(34), 6581–6592.

https://doi.org/10.1523/JNEUROSCI.0838-22.2022

Yuste, R. (2011). Dendritic Spines and Distributed Circuits. *Neuron*, *71*(5), 772–781.

https://doi.org/10.1016/j.neuron.2011.07.024

Zhang, L., Bose, P., & Warren, R. A. (2014). Dopamine Preferentially Inhibits NMDA Receptor-

Mediated EPSCs by Acting on Presynaptic D1 Receptors in Nucleus Accumbens during

Postnatal Development. *PLoS ONE*, *9*(5), e86970.

https://doi.org/10.1371/journal.pone.0086970

Zhou, Y., Zhu, H., Liu, Z., Chen, X., Su, X., Ma, C., Tian, Z., Huang, B., Yan, E., Liu, X., & Ma,

L. (2019). A ventral CA1 to nucleus accumbens core engram circuit mediates

conditioned place preference for cocaine. *Nature Neuroscience*, *22*(12), 1986–1999.

https://doi.org/10.1038/s41593-019-0524-y

Zingg, B., Chou, X., Zhang, Z., Mesik, L., Liang, F., Tao, H. W., & Zhang, L. I. (2017). AAV-

Mediated Anterograde Transsynaptic Tagging: Mapping Corticocollicular Input-Defined

Neural Pathways for Defense Behaviors. *Neuron*, *93*(1), 33–47.

https://doi.org/10.1016/j.neuron.2016.11.045

 FIGURES

 Figure 1. Overlapping dHipp and vHipp axon terminal fields in the NAc. a) Viral injection

strategy and experimental timeline. b) Injection sites in the hippocampus. Scale bar = 500µm.

Atlas image from (Paxinos & Franklin, 2019) c) Overlap of dHipp and vHipp axon terminal fields

in the NAc. Scale bar = 500µm. Atlas image from (Paxinos & Franklin, 2019) d) Close-up image

of overlapping dHipp and vHipp axon terminal fields within the NAc ventromedial shell (top). Scale

bar = 20µm. With 3-dimensional reconstruction of overlapping axon terminals (bottom). Scale bar

= 0.5µm

Figure 2. dHipp$^+$/vHipp$^+$-innervated neurons in the medial NAc shell. a) Viral targeting

strategy to selectively label dually innervated neurons b) Cartoon representation of singly vs

dually innervated neurons. c) Image of medial NAcSh depicting GCaMP-expressing cells. Scale

bar = 100µm. Atlas image from (Paxinos & Franklin, 2019) d) Close-up image of GCaMP-

expressing cells in the medial NAcSh, where neuron morphology is consistent with medium spiny

neurons. Scale bar = 100µm. e) Viral targeting strategy and experimental timeline to label cells

innervated by dHipp and vHipp. f) Cartoon representation of singly vs dually innervated neurons.

g) Image of cells innervated by dHipp (green) and vHipp (magenta). Dually innervated neurons

express both reporters and appear white. Arrowheads point out examples of dually innervated

neurons. Scale bar = 100µm h) Quantification of cells innervated single and dually innervated

neurons in the medial NAcSh across 4 slices.

Figure 3. Individual NAc neurons are functionally innervated by dHipp and vHipp. a) Viral

targeting and optogenetic approach. Electrophysiological recording strategy and representative

traces from a dually innervated NAc MSN. b) Similar AMPA:NMDA receptor ratios between dHipp-

and vHipp-MSN synapses (n=8 cells/4 mice, p=0.1953, Wilcoxon test). Scale bars on

representative traces = 40pA/10ms. c) PPRs recorded from dHipp-MSN and vHipp-MSN

synapses on dually innervated neurons (n=16 cells/10 mice, *p=0.0394, Paired t-test). Scale bars

on representative traces = 20pA/20ms. d) Rectification curve reveals linear relationship, indicating

a lack of calcium-permeable AMPA receptors (n=8 cells/4 mice). e) Rise time was calculated for

dHipp- and vHipp-evoked EPSCs, showcasing similar rise times (n=31 cells/12 mice, *p*=0.2877,

Wilcoxon test).

Figure 4. Spatial proximity of dHipp and vHipp puncta along distal dendrites of MSN. a)

Cartoon depicting optogenetic, electrophysiology, and cell-filling strategy. b) Combined and

separated images depicting an MSN filled with CF633 confirmed to have functional dHipp and

vHipp synapses, dHipp puncta (green), vHipp puncta (red). Scale bar = 5µm. c) Three-

dimensional reconstruction of an MSN dendritic segment (cyan) with overlapping/merged dHipp

(red) and vHipp (green) afferents. Scale bar = 1µm. d) Summary data showing the proportions of

nearest neighbor distances in a stacked bar graph. e) Nearest neighbor distances shown in 2µm

bins (n=9 cells/6 mice). The observed distribution differed significantly from a uniform distribution

(p<0.0001; Chi-square goodness-of-fit test). Dotted line indicates expected values from uniform

distribution.

Figure 5. Heterosynaptic potentiation of dHipp-MSN and vHipp-MSN synapses. a) Viral

expression, optogenetic, and electrophysiology strategy. b) Recording from dually innervated

MSNs, sequential and synchronous stimulation assay reveals non-linear summation of responses

(n=16 cells/10 mice, **p*=0.0027, Wilcoxon test). Scale bars on representative traces =

20pA/10ms. c) Sequential and synchronous stimulation assay performed in singly innervated

MSNs reveals similar responses to a single wavelength of light or both wavelengths of light (n=16

cells/10 mice, *p*=0.4037, Wilcoxon test). Scale bars on representative traces = 20pA/10ms. d)

Raster plot showing failures and successes from minimal stimulation assay. Time period 300-

420s is synchronized light pulses. Representative traces from Cell 9 are shown above the plot.

Scale bars on representative traces = 30pA/20ms. e) Theoretical probability of dual success, P_{exp},

compared with observed dual response success, P_{obs}, during sync period (n=9 cells/5 mice,

*p=0.0273, Wilcoxon test).

Figure S1. Prevention of cross-contamination when using 470nm light and the optogenetic

protein, Chrimson. a) Mice were injected with a Chrimson-containing viral construct and slices

were taken at least 5 weeks after surgery to perform electrophysiology. b) A response curve was

generated by sampling Chrimson-evoked responses using 470nm light, showing minimal

responses at a laser intensity of 0.75mW, and no Chrimson-evoked response at laser intensities

≤0.5mW. EPSC amplitudes were normalized to the maximum EPSC amplitude. c) Mice were

injected with a Chronos-containing viral construct and slices were taken at least 5 weeks after

surgery to perform electrophysiology. d) No responses were observed at any of the sampled light

intensities. Responses were also recorded in response to 470nm (1mW) to verify successful

targeting and stimulation (shown in green on the left).

Figure S2. Heterosynaptic potentiation across multiple light intensities. Recording from

dually innervated MSNs, sequential and synchronous stimulation assay reveals non-linear

summation of responses at multiple light intensities (0.5-0.7mW). (n=11 cells/3 mice, p=0.0029,

Friedman test with Dunn's multiple comparisons test: 0.7 vs 0.4: **p=0.0031, 0.6 vs 0.4:

*p=0.0300, 0.5 vs 0.4: *p=0.0494). Each line represents a cell. Black highlights a cell with no

change across light intensities. Red highlights a cell where potentiation score decreased with

increasing light intensity. Scale bars on representative traces = 20pA/20ms.

a.

AAV1-hSyn-Cre
AAV1-Ef1a-Flpo

dHipp
vHipp
NAc
Ai195

b.

dHipp
NAc

vHipp
NAc
singly-innervated

dHipp   vHipp
NAc
dually-innervated

c.

d.

e.

AAV9-Ef1a-
fDIO-eYFP

AAV1-hSyn-Cre
AAV1-Ef1a-Flpo

dHipp
vHipp
NAc
Ai14

f.

dHipp
NAc

vHipp
NAc
singly-innervated

dHipp   vHipp
NAc
dually-innervated

g.

NAc neurons innervated
by dHipp

NAc neurons innervated
by vHipp

Merge

h.

28%
8%
65%

■ dH innervated (28%, 68 cells)
■ vH innervated (65%, 159 cells)
■ Dually innervated (8%, 19 cells)

Figure 2

Figure 3

Figure 4

$$P_{dual}^{theoretical} = P_{dHipp-MSN}^{success} \cdot P_{vHipp-MSN}^{success}$$

Figure 5

a.

AAV-CAMKII-Chrimson-mCherry

dHipp

vHipp

NAc

Surgery

Electrophysiology

>= 5 weeks

b.

**Chrimson responses to
470nm light**

EPSC Amplitude (norm)

Laser Intensity (mW)

c.

AAV-CAMKII-Chronos-eYFP

dHipp

vHipp

NAc

Surgery

Electrophysiology

>= 5 weeks

d.

**Chronos responses to
635nm light**

EPSC Amplitude (norm)

470nm   635nm

Laser Intensity (mW)

Supplemental Figure 1