Male-Biased Cyp17a2 Orchestrates Antiviral Sexual Dimorphism in Fish via STING Stabilization and Viral Protein Degradation

Long-Feng Lu[a,c,1], Bao-jie Cui[a,e,1], Sheng-Chi Shi[a,c,1], Yang-Yang Wang[a,c], Can Zhang[a,c], Xiao Xu[a,e], Meng-Ze Tian[a,c], Zhen-Qi Li[a,c], Na Xu[a,c], Zhuo-Cong Li[a,c], Dan-Dan Chen[a,c], Li Zhou[a,c], Gang Zhai[a,c,2], Zhan Yin[f,a,c,2], Shun Li[a,b,c,d,e,2]

[a]Key Laboratory of Breeding Biotechnology and Sustainable Aquaculture, Institute of Hydrobiology, Chinese Academy of Sciences, Wuhan, Hubei 430072, China

[b]Laboratory for Marine Biology and Biotechnology, Qingdao Marine Science and Technology Center

[c]University of Chinese Academy of Sciences, Beijing, China

[d]Key Laboratory of Aquaculture Disease Control, Ministry of Agriculture, Wuhan, China

[e]College of Fisheries and Life Science, Dalian Ocean University, Dalian, China

[f]Key Laboratory of Mariculture, Ministry of Education, Ocean University of China, Qingdao 266003, China

[1]These authors have contributed equally to this work

[2]To whom correspondence may be addressed. Email: bob@ihb.ac.cn, zyin@ihb.ac.cn and zhaigang@ihb.ac.cn

Abstract

Differences in immunity between males and females in living organisms are generally thought to be due to sex hormones and sex chromosomes, and it is often assumed that males have a weaker immune response. Here we report that in fish, males exhibit stronger antiviral immune responses, the male-biased gene *cyp17a2* as a critical mediator of this enhanced response. First, we observed that male zebrafish exhibit enhanced antiviral resistance compared to females, and notably, zebrafish lack sex chromosomes. Through transcriptomic screening, we found that *cyp17a2* was specifically highly expressed in male fish. *Cyp17a2* knockout males were equivalent to wild-type males in terms of sex organs and androgen secretion, but the ability to upregulate IFN as well as antiviral resistance was greatly reduced. Then, Cyp17a2 is identified as a positive IFN regulator which located at endoplasmic reticulum, and specifically interacts with and enhances STING mediated antiviral responses. Mechanistically, Cyp17a2 stabiles STING expression by recruiting the E3 ubiquitin ligase bloodthirsty-related gene family member 32 (btr32), which facilitates K33-linked polyubiquitination. The capacity of IFN induction of Cyp17a2 was abolished when STING is knockdown. Meanwhile, Cyp17a2 also attenuates viral infection directly to strengthen the antiviral capacity as an antiviral protein, Cyp17a2 degrades the spring viremia of carp virus (SVCV) P protein by utilizing USP8 to reduce its K33-linked polyubiquitination. These findings reveal a sex-based regulatory mechanism in teleost antiviral immunity, broadening our understanding of sexual dimorphism in immune responses beyond the conventional roles of sex chromosomes and hormones.

Keywords

Cyp17a2; Sexual Dimorphism; STING; SVCV; Ubiquitination

Introduction

Research has demonstrated that innate immune responses to viruses differ significantly between the sexes. Males exhibit heightened susceptibility to viral pathogens, correlating with their generally weaker innate immune activation compared to females[1]. This immunological dimorphism manifests clinically as elevated infection intensity (quantified by intra-host viral load) and prevalence (population-level infection rates) in males[2]. Notably, sex-specific clinical outcomes are exemplified by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infections, where epidemiological data suggest males face twice the risk of intensive care unit (ICU) admission and a 30% higher mortality rate compared to females[3]. Similar trends extend to other viral pathogens including Dengue virus, hantaviruses, and hepatitis B/C viruses, all exhibiting male-biased infection prevalence[4, 5, 6].

Several factors have been identified as contributors to these sex-specific disparities in antiviral immunity, including hormonal variations and sex chromosomes. For instance, androgens such as testosterone exhibit broad immunosuppressive effects, including reduced NK cell cytotoxicity, impaired neutrophil chemotaxis, and attenuated macrophage proinflammatory cytokine production *in vitro*[7]. In addition, the Y chromosome harbors only two annotated miRNAs, while the X chromosome contains numerous immune-relevant miRNAs, and emerging evidence implicates X-escapee miRNAs as key modulators of sex-biased immune responses[8, 9].

It is well known that the interferon (IFN) response is an important host immune response to viral infection, in both mammals and fish[10]. The signaling pathways that activate IFN expression, although slightly different in fish and mammals, are generally conserved in both[11]. The endoplasmic reticulum (ER) protein stimulator of IFN genes (STING) is an important antiviral protein capable of resisting both DNA and RNA viruses by activating TANK binding kinase 1 (TBK1) during the induction of IFN production, which in turn phosphorylates IFN regulatory Factor 3 (IRF3) in the nucleus and activates the signaling of IFN transcription[12, 13]. For instance, STING

deficiency has been shown to impair innate immune responses against multiple RNA

viruses, such as dengue virus, West Nile virus, and Japanese encephalitis virus[14, 15, 16].

Notably, oncogenes of the DNA tumor viruses, including human papillomavirus E7

and adenovirus E1A, function as potent and specific inhibitors of the cyclic

GMP-AMP synthase (cGAS)-STING signaling pathway[17].

Cyp17a is classified within the cytochrome P450 (CYP450) family. In mammals,

there is only one form, cyp17a1, which possesses both hydroxylase and lyase

functions. Mutations in the human *cyp17a1* gene are associated with congenital

adrenal hyperplasia (CAH)[18]. Comparative genomic analyses reveal that teleost fishes

possess two distinct *cyp17a* isoforms, the evolutionarily conserved *cyp17a1* and a

phylogenetically divergent *cyp17a2* paralog, with the latter representing a

teleost-specific innovation absent in mammalian genomes[19]. Current investigations of

Cyp17a2 in teleost species have predominantly centered on its mechanistic roles

governing sexual development and differentiation, for example, with knockout of

zebrafish *cyp17a2* leads to significant impairment of sperm motility[20]. Although

Cyp17a2 has been characterized in sexual development, its non-canonical

physiological roles have yet to be fully elucidated.

In most vertebrates, the XY sex chromosomes exhibit significant physiological

differences between males and females, as exemplified in humans. However, the sex

determination mechanisms in fish are more complex. For instance, zebrafish lack sex

chromosomes, which confers a natural advantage. This absence allows for the

exclusion of sex chromosome-related differences in the study of immunological

disparities between males and females. Consequently, we selected zebrafish as the

model organism for this investigation. This enabled us to more precisely identify the

genes that are differentially expressed in male and female autosomes, which is crucial

for understanding the immunological differences between the sexes. This study

elucidates the role of the teleost-specific gene *cyp17a2* in antiviral immunity.

Furthermore, the discovery that the male-biased expression of *cyp17a2* confers

enhanced resistance to infection in males provides valuable insight into the complex

mechanisms underlying sexual immune dimorphism.

Results

1. Male fish exhibit enhanced resistance to viral infection compared to female fish

To determine whether sex differences exist in the antiviral response capacity of fish, age-matched female and male zebrafish were intraperitoneally (i.p.) injected with SVCV. Male fish exhibited a significantly higher survival rate than female fish within seven days post-infection (Fig. 1A). Concurrently, females demonstrated a markedly higher incidence of cutaneous hemorrhages relative to males (Fig. 1B). Consistent with the reduced mortality, male fish showed less tissue damage in the liver and spleen following viral infection (Fig. 1C). IF analysis revealed that the SVCV N protein, represented by green fluorescence, was barely detectable in male tissues but was prominently present in female tissues (Fig. 1D). At the mRNA level, viral transcripts were quantified in these tissues, and the abundance of *svcv-n* gene was significantly lower in male tissues (Fig. 1E). At the protein level, SVCV N and G proteins were less frequently detected in male tissues compared to female tissues in 7 samples (Fig. 1F). Additionally, viral titers were significantly lower in the hearts of male fish compared to female fish (Fig. 1G). These findings demonstrate a stronger antiviral response capacity in male fish compared to female fish.

2. Male-biased gene cyp17a2 orchestrates antiviral immunity through RLR pathway

To explore the potential causes of the observed differences in virus resistance between male and female fish, transcriptome sequencing was performed on head-kidney tissues of healthy adult male and female zebrafish. Differential expression analysis identified 1511 upregulated genes and 1117 downregulated genes (Fig. 2A and Table S2). From these, we focused on a subset of known or putative sex-related genes. Among eight sex-related genes, *cyp17a2* exhibited the most significant male-biased upregulation, which was subsequently confirmed by qPCR

(Fig. 2B and S1A). Furthermore, we systematically quantified Cyp17a2 mRNA and

protein expression in multiple tissues of male and female zebrafish, demonstrating a

consistent male-biased expression pattern of Cyp17a2 (Fig. 2C and 2D). Thus,

Cyp17a2 was selected for further study. We first sought to understand whether it was

associated with host antiviral immunity, and the significant upregulation of the

Cyp17a2 at the mRNA and protein level in response to viral stimulation suggested its

possible involvement in the antiviral process (Fig. S1B and S1C). Following

generation of *cyp17a2* knockout zebrafish, previous study revealed that male

cyp17a2$^{-/-}$ mutants displayed normal development of male-typical secondary sexual

characteristics comparable to WT males. However, female *cyp17a2$^{-/-}$* mutants

exhibited significantly underdeveloped genital papillae compared to their WT

counterparts. To eliminate potential confounding effects of androgen-mediated

pathways on immune parameters, subsequent experiments were conducted

exclusively using male specimens from both genotypes. The male *cyp17a2$^{-/-}$* mutants

exhibited a higher mortality rate than WT males after viral infection (Fig. 2E).

Significant tissue damage and strong green fluorescence representing the SVCV N

protein was observed in the *cyp17a2$^{-/-}$* tissues, virus titers assay also confirmed that

viral proliferation was higher in *cyp17a2$^{-/-}$* group (Fig. S1D-S1F). Consistent with

these observations, the replication of viral genes transcription and protein levels were

increased in *cyp17a2$^{-/-}$* homozygote tissues, these data demonstrated that Cyp17a2 is

crucial for host antiviral defense (Fig. 2F and S1G). To further understand their

functional mechanisms, a transcriptomic analysis contains total of 3,114 differentially

expressed genes (DEGs) were identified, there are 1,217 genes upregulated and 1,897

were downregulated in the liver of *cyp17a2* knockout male zebrafish compared to WT

males after viral infection (Fig. S1H). Kyoto encyclopedia of genes and genomes

(KEGG) pathway analysis revealed RIG-I-like receptors (RLRs), NOD-like receptors

(NLRs), and Toll-like receptors (TLRs) signaling pathways were significantly affected

(Fig. S1I). Gene set enrichment analysis (GSEA) showed that RLRs were

downregulated in *cyp17a2$^{-/-}$* group under SVCV infection, combining with numerous

IFN-stimulated genes (ISGs) decreases (Fig. S1J and S2). To validate the

transcriptome data, *ifnφ1* levels were assessed and identified significantly

downregulated in the *cyp17a2$^{-/-}$* group (Fig. 2G). Collectively, these data suggest that

viral proliferation was enhanced and IFN response were reduced in *cyp17a2$^{-/-}$* fish.

3. Cyp17a2 enhances IFN expression and suppresses viral proliferation

Given the critical role of the IFN response in host defense against viral infections,

we investigated the impact of Cyp17a2 on IFN expression and viral infection *in vitro*.

Overexpression of Cyp17a2 significantly upregulated the IFN promoter and ISRE

activities in response to poly I:C or SVCV stimulation (Fig. 3A and S3A-S3B). An

effective sh-*cyp17a2*#2 was generated and confirmed at both the mRNA and protein

levels, and knockdown of *cyp17a2* inhibited both the IFN promoter and ISRE activity

(Fig. 3B-3E and S3C-S3D). RNA sequencing analysis of the cell line revealed that

SVCV infection activated numerous ISGs in Cyp17a2-overexpressing cells, whereas

these ISGs were significantly suppressed upon *cyp17a2* knockdown (Fig. 3F and

Table S3). The induction or attenuation of IFN and ISGs upon Cyp17a2

overexpression or knockdown were confirmed by qPCR analysis, respectively (Fig.

3G and S3E). For antiviral capacity assays, Cyp17a2 from zebrafish and gibel carp

both inhibited the proliferation of host-specific viruses, conversely, *cyp17a2*

knockdown significantly enhanced viral proliferation (Fig. 3H and S3F). Regarding

viral mRNA and proteins, overexpression or knockdown Cyp17a2 also displayed the

similar opposite findings (Fig. 3I-3J and S3G-S3H). IF assays indicated a lower

intensity of green signals representing the viral protein in the Cyp17a2 overexpression

group compared to the control, while a higher green signal was observed in the

cyp17a2 knockdown group relative to the normal group (Fig. 3K and S3I). These

results suggest that Cyp17a2 upregulates IFN expression and enhances antiviral

capacity in the host.

4. Cyp17a2 interacts with STING and stabilizes its expression

Given the critical role of the RLR signaling pathway in fish IFN activation, the

relationship between Cyp17a2 and RLR signaling adaptors was investigated.

Co-transfection with Myc-Cyp17a2 and Flag-tagged RLR components (MAVS, TBK1,
STING, IRF3, IRF7) followed by anti-Flag immunoprecipitation identified specific
Cyp17a2-STING interaction via anti-Myc detection (Fig. 4A). Reciprocal Co-IP and
semi-endogenous IP assays confirmed this interaction (Fig. 4B and 4C). Truncated
STING mutants were generated to map binding domains (Fig. 4D). The
transmembrane domain (TM)-deficient STING-ΔTM-HA failed to bind Cyp17a2,
indicating TM domain dependence (Fig. 4E). Cyp17a2-GFP overlapped with ER
markers in confocal microscopy and colocalized with STING but failed to colocalize
with the ΔTM mutant (Fig. 4F-4G and S4A). To identify the effect of Cyp17a2 on
STING regulation, the mRNA level of sting was first analyzed, minimal change was
observed in *sting* upon Cyp17a2 overexpression (Fig. 4H). Protein-level
studies demonstrated Cyp17a2 remarkably enhanced STING abundance and
confirmed by confocal microscopy (Fig. 4I and S4B). Endogenous STING was
similarly stabilized dependent on Cyp17a2 under untreated or stimulated conditions
(Fig. 4J and S4C). Cycloheximide (CHX) chase assay showed prolonged exogenous
and endogenous STING half-life with Cyp17a2 co-expression, and it was also
confirmed by knockdown experiments (Fig. 4K and S4D-S4E). These data
demonstrate that Cyp17a2 binds STING via its TM and enhances STING
posttranslational stability.

5. Cyp17a2 enhances STING-mediated IFN expression and antiviral responses

Considering that Cyp17a2 targets STING, a positive regulator of IFN production,
we investigated whether Cyp17a2 influences the IFN expression induced by STING.
The IFNφ1pro activity induced by STING overexpression was significantly
enhanced by Cyp17a2, with similar amplification observed for ISRE promoter
activation (Fig. 5A and S5A). Conversely, *cyp17a2* knockdown was shown to impair
STING-mediated IFNφ1pro and ISRE activation (Fig. 5B and S5B). Corresponding
increases in *ifn* and *vig1* transcripts were detected in Cyp17a2 overexpressing cells,
while Cyp17a2 knockdown suppressed these responses (Fig. 5C-5D and S5C-S5D).
The subsequent investigation focused on the potential modulatory effect of Cyp17a2
on the STING-induced cellular antiviral response. STING overexpression reduced
cytopathogenic effect (CPE), with viral titers decreased 336-fold compared to the

control group. This protection was augmented by Cyp17a2 co-expression, achieving

an additional 20-fold reduction (Fig. 5E and S5E). IB and IF analysis confirmed

that Cyp17a2 potentiated STING-mediated suppression of viral protein expression

(Fig. 5F and S5F). SVCV gene transcripts were suppressed by STING, with enhanced

inhibition upon Cyp17a2 co-expression (Fig. 5G). STING dependency was

validated using shRNA knockdown (Fig. S5G). The antiviral capacity enhanced by

Cyp17a2 was markedly attenuated in cells with *sting* knockdown, as evidenced by

restored viral titers, gene and protein expression (Fig. 5H-5J and S5H-S5I). These

results demonstrate that Cyp17a2 potentiates STING activity to enhance IFN

production and antiviral responses.

6. Cyp17a2 stabilizes STING through btr32-mediated K33-linked

polyubiquitination

To elucidate the molecular mechanism by which Cyp17a2 augments STING

stabilization, proteasomal and lysosomal degradation were pharmacologically

inhibited using MG132 and Chloroquine (CQ), respectively. Accelerated STING

degradation caused by *cyp17a2* knockdown was rescued by MG132 but not CQ,

suggesting knockdown of *cyp17a2* promoted STING degradation via the proteasome

pathway (Fig. 6A). Then, enhanced polyubiquitination of STING was detected upon

Cyp17a2 co-expression (Fig. 6B). Ubiquitin linkage analysis revealed selective

enhancement of K33-linked chains, with *cyp17a2* knockdown shown to reduce both

WT and K33-specific ubiquitination (Fig. 6C and S6A). Subsequently, candidate E3

ligases were screened, revealing btr32 and SOCS3a as STING interactors (Fig. 6D).

btr32 was specifically identified as a Cyp17a2 binding partner through reciprocal

Co-IP (Fig. 6E-6G). Then the subcellular localization of btr32 was monitored,

btr32-GFP signal overlapped with the red signals of the ER marker, indicating that

btr32 colocalizes with the ER (Fig. 6H). Since the results identify that both STING

and Cyp17a2 are localized to the ER, btr32 was examined for cellular localization

with both STING and Cyp17a2, and the anticipated colocalization pattern was

confirmed (Fig. 6I and 6J). Cyp17a2 was found to strengthen btr32-STING

interaction (Fig. 6K and S6B). Functional assays demonstrated that btr32 augmented

of Cyp17a2-mediated IFNφ1pro activity and *vig1* mRNA induction (Fig. 6L and 6M).

btr32 potentiated the Cyp17a2-dependent stabilization of STING in both endogenous

and exogenous systems (Fig. 6N). Two shRNAs target btr32 were designed and

generated, and after validating the knockdown efficiencies for both endogenous and

exogenous targets, sh-*btr32*#2 was selected for the subsequent assay (Fig. S6C-S6D).

The knockdown of btr32 impaired Cyp17a2-enhanced STING activity, as shown by

reduced IFNφ1pro response and *vig1* transcripts (Fig. S6E and S6F). Additionally,

btr32 knockdown abolished Cyp17a2-mediated STING upregulation in endogenous

and overexpression system (Fig. S6G). These findings establish that Cyp17a2 recruits

btr32 to catalyze K33-linked polyubiquitination, thereby stabilizing STING protein.

7. STING K203 is essential for Cyp17a2-mediated ubiquitination

The mechanism between btr32 and Cyp17a2 in STING stabilization was further

characterized. btr32 overexpression was shown to intensify Cyp17a2-driven STING

ubiquitination, while *btr32* knockdown attenuated this effect (Fig. 7A and S7A).

Specific enhancement of K33-linked ubiquitination by btr32 was demonstrated, with

reciprocal reduction upon btr32 abrogation (Fig. 7B and S7B). Mass spectrometry

analysis identified K203 and K221 as candidate ubiquitination sites (Fig. 7C). Among

generated lysine-to-arginine mutants (K203R, K221R), STING stabilization by

Cyp17a2 was abolished specifically in K203R mutant (Fig. 7D). Corresponding loss

of IFN induction and ubiquitination capacity were observed in K203R mutant (Fig.

7E-7G). K33-linked ubiquitination was specifically impaired in K203R but not

K221R variants (Fig. S7C). btr32-mediated enhancement of WT and K33-specific

ubiquitination was eliminated in K203R mutant (Fig. 7H and S7D). btr32 domain

analysis was performed using RING/BBOX/SPRY deletion mutants (Fig. 7I). The

SPRY domain was required for btr32-STING-Cyp17a2 interactions (Fig. 7J-7L). Both

RING and SPRY domains were essential for btr32-driven ubiquitination (Fig. 7M and

S7E). These data establish K203 as the critical ubiquitination site in STING and

define RING/SPRY domains of btr32 as necessary for Cyp17a2-mediated

stabilization.

8. Cyp17a2 Degrades SVCV P Protein Through Modulation of K33-Linked
Polyubiquitination

To determine whether Cyp17a2 directly engage in antiviral activity beyond
regulating host immune responses, Co-IP assays were performed. Transient
transfection experiments revealed specific interaction between Cyp17a2 and SVCV P
protein, which was confirmed reciprocally (Fig. 8A and 8B). This interaction was
further observed in SVCV-infected cells using P protein-specific antibody (Fig. 8C).
In our antiviral trials, overexpression of Cyp17a2 led to a reduction in P protein levels
during SVCV infection. Consequently, the investigation focused on whether Cyp17a2
could directly target the SVCV P protein for degradation. As anticipated, the
co-expression of Cyp17a2 resulted in a decrease in the level of SVCV P protein
Overexpression of Cyp17a2 reduced P protein levels during SVCV infection (Fig. 8D).
Fluorescence microscopy confirmed diminished red fluorescent signals corresponding
to SVCV P protein in Cyp17a2-expressing cells (Fig. 8E). Confocal microscopy
analysis of SVCV P protein subcellular distribution revealed that the fluorescent
signals from P-mCherry fusion constructs (red) showed partial co-localization with
Cyp17a2 (green) in cytoplasmic compartments (Fig. 8F). To investigate degradation
mechanisms, inhibitors targeting distinct pathways were tested. Cyp17a2-mediated P
protein degradation was blocked by the proteasome inhibitor MG132, but not by
autophagy inhibitors (3-MA, Baf-A1, CQ) (Fig. 8G). Ubiquitination assays
demonstrated that Cyp17a2 significantly reduced P protein polyubiquitination (Fig.
8H). Mutation analysis using ubiquitin variants revealed that Cyp17a2 selectively
decreased K33-linked polyubiquitination, as K33R mutants exhibited no significant
changes, while K33-specific ubiquitination was markedly reduced (Fig. 8I and 8J). In
contrast, knockdown of *cyp17a2* exhibited minimal impact on both WT-Ub and
K33-linked polyubiquitination levels of the P protein (Fig. 8K). These findings
demonstrate that Cyp17a2 promotes proteasomal degradation of the SVCV P protein
by selectively attenuating K33-linked polyubiquitination, defining its role as an
antiviral protein.

9. The K12 site of P protein is essential for Cyp17a2-mediated degradation

To investigate the specific molecular mechanism underlying
ubiquitination-mediated degradation of the P protein, mass spectrometry analysis
revealed ubiquitin-specific protease 7 (USP7) and USP8 as candidate deubiquitinating
enzymes (Fig. 9A). Both USP7 and USP8 were found to associate with the P protein,
however, only USP8 interacted with Cyp17a2. Therefore, USP8 became the focus of
subsequent study (Fig. 9B and 9C). The validity of these interactions was further
confirmed through reciprocal experiments (Fig. S8A). Confocal microscopy analysis
of USP8 subcellular localization demonstrated overlapping cytoplasmic
co-localization between USP8-GFP (green fluorescence) and mCherry-tagged
Cyp17a2 or the P protein (red fluorescence) (Fig. 9D and S8B). Cyp17a2
overexpression enhanced USP8 and P protein binding, whereas *cyp17a2* knockdown
disrupted this interaction (Fig. 9E and S8C). USP8 overexpression amplified
Cyp17a2-mediated P protein reduction (Fig. 9F). Knockdown of USP8 abolished
Cyp17a2-dependent P protein degradation (Fig. 9G and S8D). USP8 facilitated
Cyp17a2-induced deubiquitination of P protein, with knockdown reversing this effect
(Fig. 9H and S8E). This result was also observed in the K33-linked ubiquitination
assay (Fig. 9I and S8F). Mass spectrometry identified K12 as a critical lysine
modification site (Fig. 9J). The K12R mutant resisted Cyp17a2-mediated degradation
and showed impaired deubiquitination in WT ubiquitin and K33-linked assays (Fig.
9K-9L and S8G). USP8 failed to enhance Cyp17a2-mediated deubiquitination in
K12R mutants (Fig. 9M and S8H). Truncation analysis demonstrated that USP8
residues 1-301 aa or 734-1067 aa were essential for interactions with P protein and
Cyp17a2 (Fig. 9N and 9O). USP8 mutants only residues 1-301 aa or 302-733 aa lost
the ability to support Cyp17a2-mediated P protein deubiquitination (Fig. 9P and S8I).
These data establish that Cyp17a2 requires USP8 and the P protein K12 site to
mediate deubiquitination-dependent degradation, with USP8 residues 734-1067 aa
being indispensable for this process.

Discussion

The influence of biological sex on antiviral immunity shows both evolutionary

parallels and divergences across species. While mammals typically exhibit stronger

female immunity through sex chromosome and hormonal mechanisms, our study

reveals an intriguing reversal in zebrafish, where males show enhanced antiviral

responses via an autosomal gene-mediated pathway. This supplements conventional

views and highlights the evolutionary plasticity of teleost immune systems.

Teleost display remarkable diversity in sex determination mechanisms, including

XX/XY, ZZ/ZW, and environmental systems[21, 22, 23]. This diversity provides a unique

framework for studying sexual dimorphism independent of sex chromosomes. Using

zebrafish, which lack conserved sex chromosomes, allows us to exclude the

confounding effects of divergent sex-chromosome gene dosage and focus on the role

of autosomal genes and their differential regulation in shaping sex-biased immunity.

Our study leverages this unique context to demonstrate that enhanced antiviral

immunity in males is mediated by the male-biased expression of the autosomal

gene *cyp17a2*.

Mechanistically, Cyp17a2 operates through dual antiviral pathways, it stabilizes

STING via btr32-mediated K33-linked ubiquitination, enhancing IFN production, and

recruits USP8 to remove K33-linked chains from viral P protein, promoting its

degradation (Fig. 10). This establishes Cyp17a2 as a central regulator of sex-specific

antiviral immunity independent of sex chromosome pathways. Particularly

noteworthy is the demonstration that sexual dimorphism in immunity can arise

through differential regulation of autosomal genes rather than direct sex chromosome

effects - a mechanism distinct from mammalian X-linked immune gene dosage effects

or estrogen-mediated immunomodulation.

A key question emerging from our findings is whether the Cyp17a2-mediated

antiviral mechanism is conserved across the teleost lineage or represents an adaptation

specific to cyprinids such as zebrafish. The teleost-specific nature of *cyp17a2*

suggests functions shaped by aquatic environmental challenges. While further studies

in diverse teleost families are needed, the presence of *cyp17a2* orthologs across fish

genomes indicates potential conservation. It is proposed that the recruitment of a core

component of the steroidogenic enzyme machinery into the innate immune network

may represent a significant evolutionary adaptation in teleost. Aquatic environments

have been shown to have a persistently high pathogen load. The co-option of a gene

such as *cyp17a2*, which is already subject to sex-biased regulatory control, offers an

efficient and rapidly deployable strategy for establishing sexually dimorphic immunity,

obviating the necessity for de novo evolution of entirely new pathways.

Sexual dimorphism in viral susceptibility represents a conserved yet

mechanistically complex phenomenon across vertebrates[24]. While males generally

exhibit increased vulnerability to numerous viral pathogens such as dengue virus and

influenza A virus, notable exceptions exist where females demonstrate higher

susceptibility, including infections by herpes simplex virus and measles virus[2]. These

divergent patterns underscore a dual dependency of sex-biased infection outcomes on

both host immunological determinants and viral evolutionary strategies[1]. For instance,

the male predominance observed in severe dengue virus cases may be explained by

both gender-specific occupational exposure risks and sexual dimorphism in

Ab-dependent enhancement mechanisms, potentially linked to androgen-mediated

modulation of antiviral immune responses[25]. Whereas female susceptibility to herpes

simplex virus could be related to ovarian sex hormones, including estradiol and

progesterone[26]. Critically, sex differences in antiviral immunity are not static but

evolve across the lifespan and are tightly linked to age and reproductive status[27]. For

example, sex disparities in hantavirus infection outcomes emerge post-puberty,

coinciding with hormonal maturation[28]. Similarly, during the 2009 H1N1 pandemic in

Japan, morbidity rates exhibited a triphasic sex bias: males<20 and >80 years old

showed higher susceptibility than females, whereas females of reproductive age

(20-49 years) faced elevated risks[29]. This pattern aligns with the hypothesis that

estrogen peaks during reproductive years enhance antiviral defenses in females, while

age-related declines in gonadal hormones may diminish this protective effect in

postmenopausal populations[30]. Our study extends these principles to teleost species,

revealing novel sex-specific resistance patterns in fish-virus interactions. Male

zebrafish demonstrated heightened resistance to SVCV, while male crucian carp

exhibited reduced susceptibility to CyHV-2[31]. These findings suggest that male-biased

resistance may represent an evolutionarily conserved strategy in certain fish-virus

system. This may reflect evolutionary adaptations to unique aquatic pressures. We

propose several non-mutually exclusive hypotheses, which are rooted in the unique

life histories and ecological pressures faced by aquatic vertebrates[32]. Firstly, the

aquatic environment acts as a potent reservoir for pathogens, constantly exposing fish

to diverse microbial challenges. This high pathogen pressure may select for

specialized and potent immune strategies. If males and females encounter different

levels or types of parasites or infections due to their behavior or morphology, this

could lead to the evolution of sex-specific immune adaptations. Secondly, this

phenomenon can be viewed through the lens of life-history trade-offs. Investment in

immunity is costly in terms of energy and must be balanced against other demands,

such as reproduction. In many teleost species, male reproductive success depends

heavily on courtship displays, nest guarding and aggressive competition with other

males, all of which are high-energy activities that can compromise immune function.

Finally, the prevalence of external fertilization in teleost is a crucial factor. Unlike

mammals, where the embryo develops internally and is protected by the maternal

immune system, fish embryos are exposed to the pathogen-rich aquatic environment

from the moment of fertilization. Therefore, a male's contribution to offspring survival

includes not only his genetic material, but also the intrinsic pathogen resistance of his

sperm and the antimicrobial properties of the seminal fluid or nest site.

STING was originally identified as a critical adaptor molecule mediating

antiviral signaling cascades[33, 34]. Subsequential studies have established its

indispensable role in orchestrating innate immune responses against both DNA and

RNA viral pathogens. Current mechanistic understanding reveals that upon detection

of cytoplasmic pathogen-derived DNA, STING operates through two distinct

molecular pathways: serving as a downstream effector of essential

single-stranded/double-stranded DNA sensors or functioning as a direct receptor for

microbial cyclic dinucleotides (CDNs). Following activation, this TM protein recruits

and activates TBK1, which subsequently phosphorylates IRF3 to facilitate its nuclear

translocation and initiate IFN production. Notably, the protein demonstrates

functional pleiotropy in RNA viral infections through selective interaction with RIG-I

rather than melanoma differentiation-associated protein 5 (MDA5), thereby activating

parallel IRF3-dependent antiviral signaling pathways. Numerous studies indicate

remarkable conservation of STING-mediated IFN induction mechanisms across

teleost species[35, 36, 37]. Mechanism studies further reveal that fish STING maintains

dual antiviral functionality against both nucleic acid virus types, while simultaneously

being subjected to targeted manipulation by various viral pathogens as an immune

evasion strategy.

Beyond its steroidogenic functions, Cyp17a2 represents a novel scaffold protein

in ubiquitination pathways. Despite the absence of clearly identifiable domains that

are directly implicated in ubiquitination (e.g., RING, HECT, or U-box domains), it is

proposed that the subcellular localization and protein-interaction capability of

Cyp17a2 are pivotal to its novel function. As an ER-resident protein, Cyp17a2 is

strategically co-localized with central innate immune sensors such as STING. This

shared localization facilitates specific protein-protein interactions, enabling Cyp17a2

to serve as an essential platform that recruits the E3 ubiquitin ligase btr32 to stabilize

STING via K33-linked ubiquitination. In contrast, Cyp17a2 engages the

deubiquitinate USP8 to remove K33-linked chains from the viral P protein, thereby

targeting it for degradation. This bifunctional and scaffolding role, which orchestrates

both the addition and removal of specific ubiquitin chains to fine-tune antiviral

responses, represents a fascinating evolutionary adaptation in teleost.

The CYP450 enzyme superfamily exhibits multifaceted roles across species,

encompassing physiological processes and immune regulation. In human immune

cells, pharmacological inhibition of CYP1A has been demonstrated to upregulate the

expression of stem cell factor receptor and interleukin 22, directly linking

CYP1-dependent metabolism of environmental small molecules to immune

modulation[38]. Evolutionary conservation of this regulatory mechanism is evident in

teleost, where grass carp (*Ctenopharyngodon idella*) CYP1A demonstrates dynamic

expression during grass carp reovirus (GCRV) infection, implicating its role in

antiviral immunity[39]. Our study further identifies Cyp17a2 as a sexually dimorphic

regulator of IFN signaling in fish. This finding expands the recognized roles of the

CYP450 superfamily in immune regulation, revealing a novel mechanism underlying

sex-specific immunity in lower vertebrates.

In conclusion, zebrafish model with its lack of conserved sex chromosomes and

environmental sex determination plasticity, provides unique insights into the

evolutionary continuum of sexual dimorphism. Our results suggest that the

relationship between sexual differentiation and immune competence may be more

fluid in teleost than in mammals, potentially reflecting adaptive responses to aquatic

pathogens and variable mating systems. Future investigations should explore whether

this autosomal-centered mechanism represents a teleost specific adaptation or an

ancestral feature subsequently modified in terrestrial vertebrates, while also

examining potential interactions between environmental sex determinants and immune gene regulation.

Acknowledgment

We thank Fang Zhou (Institute of Hydrobiology, Chinese Academy of Sciences) for assistance with confocal microscopy analysis and Dr. Feng Xiong (China Zebrafish Resource Center, Institute of Hydrobiology, Chinese Academy of Sciences) for assistance with qPCR analysis.

This work was supported by the Strategic Priority Research Program of the Chinese Academy of Sciences (XDB0730300), Biological Breeding-National Science and Technology Major Project (2023ZD04065), National Excellent Youth Science Fund (32322086) and the Youth Innovation Promotion Association provided funding to Shun Li. National Key Research and Development Program of China (2023YFD2400201) provided funding to Dan-Dan Chen. National Natural Science Foundation of China (32173023) provided funding to Long-Feng Lu.

Declaration of interests

The authors declare no competing interests.

STAR Methods

Ethics statement

The experiments involved in this study were conducted in compliance with ethical regulations. The fish experiments were carried out under the guidance of the European Union Guidelines for the Handling of Laboratory Animals (2010/63/EU) and approved by the Ethics Committee for Animal Experiments of the Institute of Hydrobiology, Chinese Academy of Sciences (CAS) (No. 2024-069).

Fish, cells, and viruses

Mature zebrafish individuals 2.5 months after hatching (0.4 ± 0.1 g) were

selected in this study. Wild-type (WT) zebrafish (AB strain; *Danio rerio*) were procured from the China Zebrafish Resource Center (CZRC), while *cyp17a2* mutant lines were generously provided by Dr. Zhan Yin's laboratory at the Institute of Hydrobiology, Chinese Academy of Sciences. All specimens were bred using standardized procedures. In accordance with the ethical requirements and national animal welfare guidelines, all experimental fish were required to undergo a two-week acclimatization period in the laboratory and have their health assessed prior to the commencement of the study. Only fish that appeared healthy and were mobile were selected for scientific investigation. ZF4 cells (American Type Culture Collection, ATCC) were cultured in Ham's F-12 medium (Invitrogen) supplemented with 10% fetal bovine serum (FBS) at 28°C and 5% CO_2. EPC cells were obtained from the Chinese Culture Collection Centre for Type Cultures (CCTCC), Gibel carp brain (GiCB) cells were provided by Ling-Bing Zeng (Yangtze River Fisheries Research Institute, Chinese Academy of Fishery Sciences), these cells were maintained at 28°C in 5% CO_2 in medium 199 (Invitrogen) supplemented with 10% FBS. Spring viremia of carp virus (SVCV) was propagated in EPC cells until a CPE was observed, and then cell culture fluid containing SVCV was harvested and centrifuged at 4×10^3 *g* for 20 min to remove the cell debris, and the supernatant was stored at -80°C until used. Cyprinid herpesvirus 2 (CyHV-2, obtained from Yancheng city, Jiangsu province, China) was provided by Prof. Liqun Lu (Shanghai Ocean University). CyHV-2 was propagated in GICB cells and harvested in a similar way to SVCV.

Plasmid construction and reagents

The sequences of zebrafish and gibel carp (*Carassius gibelio*) Cyp17a2 (GenBank accession number: NM_001105670.1 and XM_026199804.1) were obtained from the National Centre for Biotechnology Information (NCBI) website. Cyp17a2 was amplified by polymerase chain reaction (PCR) using cDNA from adult zebrafish or gibel carp tissues as a template and cloned into the expression vector pCMV-HA or pCMV-Myc (Clontech) vectors. Zebrafish MAVS (NM_001080584.2), TBK1 (NM_001044748.2), STING (NM_001278837.1) and the truncated mutants of STING, IRF3 (NM_001143904.1), IRF7 (BC058298.1), btr32 (XM_021470074.1)

and the truncated mutants of btr32, TRAIP (NM_205607.1), SOCS3a (NM_199950.1),

USP7 (XM_021473871.1), USP8 (XM_009293353.4), and N, P, and G protein

(DQ097384.2) of SVCV were cloned into pCMV-Myc and pCMV-Tag2C vectors.

The shRNA of *Pimephales promelas* Cyp17a2 (XM_039679078.1), STING

(HE856620.1), and USP8 (XM_039680290.1) were designed by BLOCK-iT RNAi

Designer and cloned into the pLKO.1-TRC Cloning vector. For subcellular

localization experiments, Cyp17a2 was constructed onto pEGFP-N3 (Clontech), while

STING and P protein were constructed onto pCS2-mCherry (Clontech). The plasmids

containing zebrafish IFNφ1pro-Luc and ISRE-Luc in the pGL3-Basic luciferase

reporter vector (Promega) were constructed as described previously. The *Renilla*

luciferase internal control vector (pRL-TK) was purchased from Promega. The

ubiquitin mutant expression plasmids Lys-33/63 (all lysine residues were mutated

except Lys-33 or Lys-63) and Lys-33R/63R (only Lys-33 or Lys-63 mutated) were

ligated into the pCMV-HA vectors named Ub-K33O/K63O-HA and Ub-K33R/K63R.

All constructs were confirmed by DNA sequencing. Polyinosinic-polycytidylic acid

(poly I:C) was purchased from Sigma-Aldrich used at a final concentration of 1 µg/µl.

MG132 (Cat. No. M7449), 3-Methyladenine (3-MA) (Cat. No. M9281), CQ (Cat. No.

C6628) were obtained from Sigma-Aldrich. Bafilomycin A1 (Baf-A1) (Cat. No.

S1413) and CHX (NSC-185) were obtained from Selleck.

Transcriptomic analysis

Total RNA was extracted using the TRIzol method and evaluated for RNA purity

and quantification using a NanoDrop 2000 spectrophotometer (Thermo Scientific,

Waltham, U.S.A.) and RNA integrity was assessed using an Agilent 2100 Bioanalyzer

(Agilent Technologies, Santa Clara, U.S.A.). Transcriptome sequencing and data

analysis were conducted by OE Biotech (Shanghai, China). The raw sequencing data

was submitted to the NCBI (GEO accession number: GSE286486).

Transient transfection and virus infection

EPC cells were transfected in 6-well and 24-well plates using transfection

reagents from FishTrans (MeiSenTe Biotechnology) according to the manufacturer's

protocol. Antiviral assays were conducted in 24-well plates by transfecting EPC cells

with the plasmids shown in the figure. At 24 h post-transfection, cells were infected

with SVCV at a multiplicity of infection (MOI = 0.001). After 24 h, supernatant

aliquots were harvested for detection of virus titers, the cell monolayers were fixed by

4% paraformaldehyde (PFA) and stained with 1% crystal violet for visualizing CPE.

For virus titration, 200 µl of culture medium were collected at 48 h post-infection and

used for detection of virus titers according to the method of Reed and Muench. The

supernatants were subjected to 3-fold or 10-fold serial dilutions and then added (100

µl) onto a monolayer of EPC cells cultured in a 96-well plate. After 48 or 72 h, the

medium was removed and the cells were washed with phosphate-buffered saline

(PBS), fixed by 4% PFA, and stained with 1% crystal violet. The virus titer was

expressed as 50% tissue culture infective dose ($TCID_{50}$/ml). For viral infection, fish

were anesthetized with methanesulfonate (MS-222) and intraperitoneally (i.p.)

injected with 5 µl of M199 containing SVCV (5×10^8 $TCID_{50}$/ml). The i.p. injection

of PBS was used as mock infection. Then the fish were migrated into the aquarium

containing new aquatic water.

Luciferase activity assay

EPC cells were cultured overnight in 24-well plates and subsequently

co-transfected with the expression plasmid and luciferase reporter plasmid. The cells

were infected with SVCV or transfected with poly I:C for 24 h prior to harvest. At 24

h post-stimulation, cells were washed with PBS and lysed for measuring luciferase

activity by the Dual-Luciferase Reporter Assay System (Promega) according to the

manufacturer's instructions. Firefly luciferase activity was normalized based on the

Renilla luciferase activity.

RNA extraction, reverse transcription, and quantitative PCR (qPCR)

The RNA was extracted using TRIzol reagent (Invitrogen), and first-strand

cDNA was synthesized with a PrimeScript RT kit with gDNA Eraser (Takara). qPCR

was performed on the CFX96 Real-Time System (Bio-Rad) using SYBR green PCR

Master Mix (Yeasen). The PCR conditions were as follows: 95°C for 5 min and then

40 cycles of 95°C for 20 s, 60°C for 20 s, and 72°C for 20 s. The primers utilized for

the qPCRs are presented in Table S1, and the *β-actin* gene was utilized as the internal

control. The relative fold changes were calculated by comparison to the corresponding

controls using the $2^{-\Delta\Delta Ct}$ method (where CT was the threshold cycle).

Co-immunoprecipitation (Co-IP) assay

EPC cells were cultured in 10-cm^2 dishes overnight and subsequently transfected

with 10 μg plasmid as illustrated. At 24 h post-transfection, the medium was removed

and cells were washed with PBS. Then the cells were lysed in 1 ml

radioimmunoprecipitation (RIPA) lysis buffer [1% NP-40, 50 mM Tris-HCl, pH 7.5,

150 mM NaCl, 1 mM EDTA, 1 mM NaF, 1 mM sodium orthovanadate (Na$_3$VO$_4$), 1

mM phenyl-methylsulfonyl fluoride (PMSF), 0.25% sodium deoxycholate] containing

protease inhibitor cocktail (Sigma-Aldrich) at 4°C for 1 h on a rocker platform. The

cellular debris was removed by centrifugation at 12,000 × g for 15 min at 4°C. The

supernatant was transferred to a fresh tube and incubated with 20 μl

anti-Flag/HA/Myc affinity gel (Sigma-Aldrich) overnight at 4°C with constant

rotating incubation. These samples were further analyzed by immunoblotting (IB).

Immunoprecipitated proteins were collected by centrifugation at 5000 × g for 1 min at

4°C, washed three times with lysis buffer and resuspended in 50 μl 2 × SDS sample

buffer. The immunoprecipitates and whole cell lysates (WCLs) were analyzed by IB

with the indicated antibodies (Abs).

***In vivo* ubiquitination assay**

The transfected EPC cells were washed twice with 10 mL ice-cold PBS and

subsequently digested with 1 mL 0.25% trypsin-EDTA (1×) (Invitrogen) for 2-3 min

until the cells were dislodged. 100 μL FBS was added to neutralize the trypsin and the

cells were resuspended into 1.5 mL centrifuge tube, centrifuged at 2000 × g for 5 min.

The supernatant was discarded and the cell precipitations were resuspended using 1

mL PBS and centrifuged at 2000 × g for 5 min. The collected cell precipitations were

lysed using 100 μL PBS containing 1% SDS and denatured by heating for 10 min.

The supernatants were diluted with lysis buffer until the concentration of SDS was

reduced to 0.1%. The diluted supernatants were incubated with 20 μL anti-Myc

affinity gel overnight at 4°C with constant agitation. Subsequently, these samples

were subjected to further analysis by IB. The immunoprecipitated proteins were

collected by centrifugation at 5000 × *g* for 1 min at 4°C, washed for three times with

lysis buffer and resuspended in 100 μL 1× SDS sample buffer.

Immunoblot analysis

Immunoprecipitates or WCLs were analyzed as described previously. Abs were

diluted as follows: anti-*β-actin* (ABclonal, AC026) at 1:10000, anti-Flag

(Sigma-Aldrich, F1804) at 1:3000, anti-HA (Covance, MMS-101R) at 1:3000,

anti-Myc (Santa Cruz Biotechnology, sc-40) at 1:3000, anti- btr32 (ABclonal, A13887)

at 1:1000, anti-USP8 (proteintech, 27791-1-AP) at 1:1000, and HRP-conjugated

anti-mouse/rabbit IgG (Thermo Scientific, 31430/31460) at 1:5000,

anti-N/P/G/STING/ (prepared and purified in our lab) at 1:2000.

Immunofluorescence (IF)

EPC cells were plated onto glass coverslips in 6-well plates and infected with

SVCV (MOI = 1) 24 h. The cells were then washed with PBS and fixed in 4% PFA at

room temperature for 1 h and permeabilized with 0.2% Triton X-100 in ice-cold PBS

for 15 min. The samples were incubated for 1 h at room temperature in PBS

containing 2% bovine serum albumin (BSA, Sigma-Aldrich). After additional PBS

washing, the samples were incubated with anti-N Ab in PBS containing 2% BSA for

2-4 h at room temperature. After three times washed by PBS, the samples were

incubated with secondary Ab (Goat anti-Rabbit IgG (H+L) Highly Cross-Adsorbed

Secondary Antibody, Alexa Fluor™ Plus 488) (Thermo Scientific, A32731, 1:5000) in

PBS containing 2% BSA for 1 h at room temperature. After additional PBS washing,

the cells were finally stained with 1 μg/ml 4′, 6-diamidino-2-phenylindole (DAPI,

Beyotime Institute of Biotechnology, Shanghai, China) for 10 min in the dark at room

temperature. Finally, the coverslips were washed and observed with a confocal

microscope under a 10× immersion objective (SP8; Leica).

Fluorescent microscopy

666 EPC cells were plated onto coverslips in 6-well plates and transfected with the

667 plasmids indicated for 24 h. Thereafter, the cells were washed twice with PBS and

668 fixed with 4% PFA for 1 h. After being washed three times with PBS, the cells were

669 stained with 1 μg/ml DAPI for 15 min in the dark at room temperature. Subsequently,

the coverslips were washed and observed with a confocal microscope under a 63× oil immersion objective (SP8; Leica).

Histopathology

Liver and spleen tissues from three individuals of control or virus-infected fish at 2 days post infection (dpi) were dissected, and fixed in Bouin's Fixative overnight. Subsequently, the samples were dehydrated in ascending grades of alcohol and embedded into paraffin. Sections at 5 μm thickness were taken and stained with hematoxylin and eosin (H&E). Histological changes were examined by optical microscopy at ×40 magnification and were analyzed by the Aperio ImageScope software.

Statistics analysis

For fish survival analysis, Kaplan-Meier survival curves were generated and subsequently subjected to a log-rank test. For the bar graph, one representative experiment of at least three independent experiments is shown, and each was done in triplicate. For the dot plot graph, each dot point represents one independent biological replicate. Unpaired Student's t test was used for statistical analysis. Data are presented as mean ± standard error of the mean (SEM). A p value < 0.05 was considered statistically significant.

References

1. Klein, S.L. & Huber, S. Sex Differences in Susceptibility to Viral Infection. In: Klein, S.L. & Roberts, C. (eds). *Sex Hormones and Immunity to Infection*. Springer Berlin Heidelberg: Berlin, Heidelberg, 2010, pp 93-122.

2. Klein, S.L. Sex influences immune responses to viruses, and efficacy of prophylaxis and treatments for viral diseases. *BioEssays : news and reviews in molecular, cellular and developmental biology* **34**, 1050-1059 (2012).

3. Health, G., Icrw, Aphrc & Gautam, A. *The-COVID-19-Sex-Disaggregated-Data-Tracker-Update*, 2020.

4. Guha-Sapir, D. & Schimmer, B. Dengue fever: new paradigms for a changing epidemiology. *Emerg Themes Epidemiol* **2**, 1 (2005).

5. Tsay, P.K. *et al.* Impact of gender, viral transmission and aging in the prevalence of hepatitis B
surface antigen. *Chang Gung medical journal* **32**, 155-164 (2009).

6. Burguete-García, A.I. *et al.* Hepatitis C seroprevalence and correlation between viral load and
viral genotype among primary care clients in Mexico. *Salud publica de Mexico* **53 Suppl 1**, S7-12
(2011).

7. Liva, S.M. & Voskuhl, R.R. Testosterone acts directly on CD4[+]T lymphocytes to increase IL-10
production. *Journal of immunology* **167**, 2060-2067 (2001).

8. Pinheiro, I., Dejager, L. & Libert, C. X-chromosome-located microRNAs in immunity: might they
explain male/female differences? The X chromosome-genomic context may affect X-located miRNAs
and downstream signaling, thereby contributing to the enhanced immune response of females.
BioEssays : news and reviews in molecular, cellular and developmental biology **33**, 791-802 (2011).

9. Sharma, S. & Eghbali, M. Influence of sex differences on microRNA gene regulation in disease.
Biology of sex differences **5**, 3 (2014).

10. Akira, S., Uematsu, S. & Takeuchi, O. Pathogen recognition and innate immunity. *Cell* **124**,
783-801 (2006).

11. Chen, S.N., Zou, P.F. & Nie, P. Retinoic acid-inducible gene I (RIG-I)-like receptors (RLRs) in
fish: current knowledge and future perspectives. *Immunology* **151**, 16-25 (2017).

12. Ishikawa, H. & Barber, G.N. STING is an endoplasmic reticulum adaptor that facilitates innate
immune signalling. *Nature* **455**, 674-678 (2008).

13. Ni, G.X., Ma, Z. & Damania, B. cGAS and STING: At the intersection of DNA and RNA
virus-sensing networks. *PLoS pathogens* **14** (2018).

14. Aguirre, S. *et al.* DENV Inhibits Type I IFN Production in Infected Cells by Cleaving Human
STING. *PLoS pathogens* **8** (2012).

15. Nazmi, A., Mukhopadhyay, R., Dutta, K. & Basu, A. STING Mediates Neuronal Innate Immune
Response Following Japanese Encephalitis Virus Infection. *Scientific reports* **2** (2012).

16. You, F.P. *et al.* ELF4 is critical for induction of type I interferon and the host antiviral response.
Nature immunology **14**, 1237-+ (2013).

17. Lau, L., Gray, E.E., Brunette, R.L. & Stetson, D.B. DNA tumor virus oncogenes antagonize the

cGAS-STING DNA-sensing pathway. *Science* **350**, 568-571 (2015).

18. Kim, Y.M. *et al.* A review of the literature on common CYP17A1 mutations in adults with

17-hydroxylase/17,20-lyase deficiency, a case series of such mutations among Koreans and functional

characteristics of a novel mutation. *Metabolism* **63**, 42-49 (2014).

19. Zhou, L.Y. *et al.* A novel type of P450c17 lacking the lyase activity is responsible for C21-steroid

biosynthesis in the fish ovary and head kidney. *Endocrinology* **148**, 4282-4291 (2007).

20. Shi, S.C. *et al.* Characterization of the Interrenal Gland and Sexual Traits Development in

cyp17a2-Deficient Zebrafish. *Front Endocrinol* **13** (2022).

21. Matsuda, M. *et al.* DMY is a Y-specific DM-domain gene required for male development in the

medaka fish. *Nature* **417**, 559-563 (2002).

22. Yoshida, K. *et al.* B Chromosomes Have a Functional Effect on Female Sex Determination in

Lake Victoria Cichlid Fishes. *Plos Genet* **7** (2011).

23. Hayashi, Y. *et al.* High Temperature Causes Masculinization of Genetically Female Medaka by

Elevation of Cortisol. *Mol Reprod Dev* **77**, 679-686 (2010).

24. Zajitschek, S.R. *et al.* Sexual dimorphism in trait variability and its eco-evolutionary and

statistical implications. *Elife* **9** (2020).

25. Gan, E.S., Ting, D.H.R. & Chan, K.R. The mechanistic role of antibodies to dengue virus in

protection and disease pathogenesis. *Expert Rev Anti-Infe* **15**, 111-119 (2017).

26. Wald, A. Herpes Simplex Virus Type 2 Transmission: Risk Factors and Virus Shedding. *Herpes :*

the journal of the IHMF **11 Suppl 3**, 130A-137A (2004).

27. Klein, S.L. & Flanagan, K.L. Sex differences in immune responses. *Nature Reviews Immunology*

16, 626-638 (2016).

28. Vial, P.A. *et al.* Hantavirus in humans: a review of clinical aspects and management. *Lancet Infect*

Dis **23**, E371-E382 (2023).

29. Eshima, N. *et al.* Sex- and Age-Related Differences in Morbidity Rates of 2009 Pandemic
Influenza A H1N1 Virus of Swine Origin in Japan. *PloS one* **6** (2011).

781
782   30. Cui, J., Shen, Y. & Li, R. Estrogen synthesis and signaling pathways during aging: from periphery
783   to brain. *Trends Mol Med* **19**, 197-209 (2013).

784
785   31. Lu, L.F. *et al.* Fish female-biased gene leads to female antiviral response attenuation between
786   sexes by autophagic degradation of MITA. *PLoS pathogens* **18** (2022).

787
788   32. Gui, J.-F., Zhou, L. & Li, X.-y. Rethinking fish biology and biotechnologies in the challenge era
789   for burgeoning genome resources and strengthening food security. *Water Biology and Security* **1**
790   (2021).

791
792   33. Woznica, A. *et al.* STING mediates immune responses in the closest living relatives of animals.
793   *Elife* **10** (2021).

794
795   34. Ran, Y., Shu, H.B. & Wang, Y.Y. MITA/STING: a central and multifaceted mediator in innate
796   immune response. *Cytokine Growth F R* **25**, 631-639 (2014).

797
798   35. Biacchesi, S., Merour, E., Lamoureux, A., Bernard, J. & Bremont, M. Both STING and MAVS
799   fish orthologs contribute to the induction of interferon mediated by RIG-I. *PloS one* **7**, e47737 (2012).

800
801   36. Lu, L.-F. *et al.* Fish herpesvirus KLP manipulates Beclin1 to selectively degrade MITA through a
802   precise autophagic manner for immune evasion. *Water Biology and Security* **2**, 100092 (2023).

803
804   37. Li, S., Lu, L.-F. & Zhang, C. The LxIS motif is the language between virus and host RLRs: From
805   fish to mammalian viruses. *Water Biology and Security* **3**, 100246 (2024).

806
807   38. Effner, R. *et al.* Cytochrome P450s in human immune cells regulate IL-22 and c-Kit via an AHR
808   feedback loop. *Scientific reports* **7** (2017).

809
39. Chu, P.F. *et al.* Identification, expression and functional characterisation of in grass carp
(Ctenopharyngodon idella). *Fish Shellfish Immun* **95**, 35-43 (2019).

Figure legends

Figure 1. Male fish show stronger viral resistance than females.

(A) Survival (Kaplan-Meier Curve) of male and female zebrafish (n = 15 per group)

at various days after i.p. injected with SVCV (5×10^8 TCID$_{50}$/ml, 5 μL/individual). (B)

Sex-specific morphological alterations in zebrafish given i.p. injection of SVCV for

48 h (n = 3 per group). (C) Microscopy of H&E-stained liver and spleen sections from

male and female zebrafish treated with SVCV for 48 h. (D) IF analysis of SVCV-N

protein in liver and spleen of male and female zebrafish treated with SVCV for 48 h.

(E) qPCR analysis of *svcv-n* mRNA in the liver and spleen of male and female

zebrafish (n = 26 per group) given i.p. injection of SVCV for 48 h. (F) IB analysis of

SVCV proteins in the liver and spleen sections of male and female zebrafish (n = 7

per group) treated with SVCV for 48 h. (G) The viral titer of heart in male and female

zebrafish (n = 40 per group) treated with SVCV for 48 h.

Figure 2. Male-biased cyp17a2 regulates RLR-mediated antiviral immunity.

(A) The differently expressed gene number of mRNA variations presented as a

volcano plot in head-kidney of male and female zebrafish. (B) Heatmap view of

mRNA variations of eight sex-regulated genes in the head-kidney of male and female

zebrafish. Values are represented as $\log_2^{(FPKM+1)}$. (C) Heatmap view of *cyp17a2*

mRNA in the brain, heart, head-kidney, gill, liver, spleen, body-kidney, gonad, gut,

muscle, and skin of male and female zebrafish (n = 5 per group). Expression values

$[\log_2^{(fold\ change)}]$ were transformed to Z-scores across rows for each gene to highlight

patterns of relative expression. (D) IB analysis of Cyp17a2 in the heart, head-kidney,

gill, liver, spleen, body-kidney, gonad, and skin of male and female zebrafish (n = 5

per group). (E) Survival (Kaplan-Meier Curve) of *cyp17a2$^{+/+}$* and *cyp17a2$^{-/-}$* male

zebrafish (n = 16 per group) at various days after i.p. injected with SVCV (5×10^8

TCID50/ml, 5 μl/individual). (F) qPCR analysis of *svcv-n* mRNA in the liver and

spleen of *cyp17a2$^{+/+}$* and *cyp17a2$^{-/-}$* male zebrafish (n = 20 per group) given i.p.

injection of SVCV for 48 h. (G) qPCR analysis of *ifnφ1* mRNA in the liver and spleen

of *cyp17a2$^{+/+}$* and *cyp17a2$^{-/-}$* male zebrafish (n = 20 per group) given i.p. injection of

SVCV for 48 h.

Figure 3. Cyp17a2 upregulates IFN expression and inhibits viral replication.

(A and E) Luciferase activity of IFNφ1pro in EPC cells transfected with indicated

plasmids for 24 h, and then untreated or transfected with poly I:C (0.5 μg) or infected

with SVCV (MOI = 1) for 24 h before luciferase assays. (B) qPCR analysis of

cyp17a2 in EPC cells transfected with indicated plasmids for 24 h, and then untreated

or transfected with poly I:C (2 μg) or infected with SVCV (MOI = 1) for 24 h. (C and

D) IB analysis of proteins in EPC cells transfected with indicated plasmids for 24 h.

(F) Heatmap view of mRNA variations of SVCV-activated ISG sets in the

Cyp17a2-overexpressing cells or *cyp17a2* knockdown cells and infected with SVCV

for 24 h. Expression values are $\log_2^{(FPKM+1)}$ transformed and presented as row

Z-scores. (G) qPCR analysis of *ifn* in EPC cells transfected with indicated plasmids

for 24 h, and then untreated or infected with SVCV (MOI = 1) for 24 h. (H) Plaque

assay of virus titers in EPC cells or GICB cells transfected with indicated plasmids for

24 h, followed by SVCV or CyHV-2 challenge for 24 h or 48 h. (I and J) qPCR and

IB analysis of SVCV genes in EPC cells transfected with indicated plasmids for 24 h,

followed by SVCV challenge for 24 h. (K) IF analysis of SVCV proteins in EPC cells

transfected with indicated plasmids for 24 h, followed by SVCV challenge for 24 h.

Figure 4. Cyp17a2 physically interacts with STING and stabilizes its expression.

(A-C and E) IB analysis of WCLs and proteins immunoprecipitated with anti- Flag or

HA Ab-conjugated agarose beads from EPC cells transfected with indicated plasmids

for 24 h. (D) Schematic representation of full-length STING and its mutants. (F and G)

Confocal microscopy of Cyp17a2 and ER or STING and its mutants in EPC cells

transfected with indicated plasmids for 24 h. The coefficient of colocalization was

determined by qualitative analysis of the fluorescence intensity of the selected area in

Merge. (H) qPCR analysis of *sting* in EPC cells transfected with indicated plasmids

for 24 h, followed by untreated or infected with SVCV (MOI = 1) for 24 h. (I) IB

analysis of proteins in EPC cells transfected with indicated plasmids for 24 h. (J) IB

analysis of proteins in EPC cells transfected with indicated plasmids for 24 h,

followed by untreated or infected with SVCV (MOI = 1) for 24 h. (K) IB analysis of

proteins in EPC cells transfected with indicated plasmids for 18 h, then treated with

CHX for 4 h and 8 h.

Figure 5. Cyp17a2 potentiates STING-induced IFN production and antiviral

responses.

(A and B) Luciferase activity of IFNφ1pro in EPC cells transfected with indicated

plasmids for 24 h before luciferase assays. (C and D) qPCR analysis of *ifn* in EPC

cells transfected with indicated plasmids for 24 h. (E and H) Detection of virus titers

in EPC cells transfected with indicated plasmids for 24 h and then infected with

SVCV (MOI = 1) for 24 h. (F and J) IB analysis of proteins in EPC cells transfected

with indicated plasmids for 24 h and then untreated or infected with SVCV (MOI = 1)

for 24 h. (G and I) qPCR analysis of SVCV genes in EPC cells transfected with

indicated plasmids for 24 h, followed by SVCV challenge for 24 h.

Figure 6. Cyp17a2 stabilizes STING by orchestrating btr32-catalyzed K33-linked

polyubiquitination.

(A) IB analysis of proteins in EPC cells transfected with indicated plasmids for 18 h,

then treated with CHX and DMSO, MG132, or CQ for 4 h and 8 h. (B and C) STING

ubiquitination assays in EPC cells transfected with indicated plasmids for 24 h. (D-G,

K) IB analysis of WCLs and proteins immunoprecipitated with anti-Flag

Ab-conjugated agarose beads from EPC cells transfected with indicated plasmids for

24 h. (H and I) Confocal microscopy of btr32 and ER or STING or Cyp17a2 in EPC

cells transfected with indicated plasmids for 24 h. The coefficient of colocalization

was determined by qualitative analysis of the fluorescence intensity of the selected

area in Merge. (J) Colocalization analyses of figure 6I were performed by calculating

Pearson correlation coefficient. (L and M) Luciferase activity of IFNφ1pro and qPCR

analysis of *vig1* in EPC cells transfected with indicated plasmids for 24 h. (N) IB

analysis of proteins in EPC cells transfected with indicated plasmids for 24 h.

Figure 7 Cyp17a2-mediated ubiquitination of STING is dependent on the K203.

(A and B, G and H, and M) STING ubiquitination assays in EPC cells transfected

with indicated plasmids for 24 h. (C) Mass spectrometry analysis of a peptide derived

from ubiquitinated STING-Myc. (D) IB analysis of proteins in EPC cells transfected

with indicated plasmids for 24 h. (E and F) Luciferase activity of IFNφ1pro and qPCR

analysis of *ifn* in EPC cells transfected with indicated plasmids for 24 h. (I) Schematic

representation of full-length btr32 and its mutants. (J-L) IB of WCLs and proteins

immunoprecipitated with anti-Flag/HA Ab-conjugated agarose beads from EPC cells

transfected with indicated plasmids for 24 h.

Figure 8 Cyp17a2 induces proteasomal degradation of the SVCV P protein by

catalyzing K33-linked polyubiquitination.

(A-C) IB analysis of WCLs and proteins immunoprecipitated with anti-Flag, anti-HA,

or anti-Myc Ab-conjugated agarose beads from EPC cells transfected with indicated

plasmids for 24 h. (D) IB analysis of proteins in EPC cells transfected with indicated

plasmids for 24 h. (E) Fluorescent analysis of proteins in EPC cells transfected with

indicated plasmids for 24 h. (F) Confocal microscopy of P and Cyp17a2 in EPC cells

transfected with indicated plasmids for 24 h. The coefficient of colocalization was

determined by qualitative analysis of the fluorescence intensity of the selected area in

Merge. (G) IB analysis of proteins in EPC cells transfected with indicated plasmids

for 18 h, followed by treatments of MG132 (10 μM), 3-MA (2 mM), Baf-A1 (100

nM), and CQ (50 μM) for 6 h, respectively. (H-K) P protein ubiquitination assays in

EPC cells transfected with indicated plasmids for 18 h, followed by MG132

treatments for 6 h.

Figure 9 Cyp17a2 targets the K12 site of the P protein for K33-linked

polyubiquitination.

(A) List of proteins interacting with P protein detected by mass spectrometry. (B and

C, E, O) IB analysis of WCLs and proteins immunoprecipitated with anti-HA/Myc

Ab-conjugated agarose beads from EPC cells transfected with indicated plasmids for

24 h. (D) Confocal microscopy of USP8 and Cyp17a2 or P protein in EPC cells

transfected with indicated plasmids for 24 h. The coefficient of colocalization was

determined by qualitative analysis of the fluorescence intensity of the selected area in

Merge. (F and G, K) IB analysis of proteins in EPC cells transfected with indicated

plasmids for 24 h. (H and I, L and M, P) P protein ubiquitination assays in EPC cells

transfected with indicated plasmids for 18 h, followed by MG132 treatments for 6 h.

(J) Mass spectrometry analysis of a peptide derived from ubiquitinated P-Myc. (N)

Schematic representation of full-length USP8 and its mutants.

Figure 10 A mechanistic model illustrating the dual regulatory roles of Cyp17a2

in upregulating STING expression and degrading SVCV P protein.

Upon virus infection, male-biased autosomal gene *cyp17a2* enhances antiviral

responses in males. Mechanistically, Cyp17a2, an ER-localized protein, stabilizes

STING expression through recruitment of the E3 ubiquitin ligase btr32, promoting

K33-linked polyubiquitination. Furthermore, Cyp17a2 facilitates proteasomal

degradation of the SVCV P protein by engaging USP8 to reduce its K33-linked

polyubiquitination, thereby amplifying antiviral IFN responses.

Figure S1 The male-biased expression of Cyp17a2 governs RLR antiviral

signaling.

(A) qPCR analysis of *cyp17a2* mRNA in the head-kidney of male and female

zebrafish (n = 50 per group). (B) qPCR analysis of *cyp17a2* mRNA in the ZF4 or EPC

cells infected with SVCV for the indicated times. (C) IB analysis of Cyp17a2 in ZF4

and EPC cells infected with SVCV for the indicated times. (D) Microscopy of

H&E-stained liver and spleen sections from *cyp17a2*$^{+/+}$ and *cyp17a2*$^{-/-}$ male zebrafish

treated with SVCV for 48 h. (E) IF analysis of SVCV-N protein in the liver and spleen

of *cyp17a2*$^{+/+}$ and *cyp17a2*$^{-/-}$ male zebrafish treated with SVCV for 48 h. (F) The viral

titer of heart in *cyp17a2*$^{+/+}$ and *cyp17a2*$^{-/-}$ male zebrafish (n = 15 per group) treated

with SVCV for 48 h. (G) IB analysis of SVCV proteins in the liver and spleen

sections of *cyp17a2*$^{+/+}$ and *cyp17a2*$^{-/-}$ male zebrafish (n = 3 per group) treated with

SVCV for 48 h. (H) The differently expressed gene number of mRNA variations in

liver of *cyp17a2*$^{+/+}$ and *cyp17a2*$^{-/-}$ male zebrafish infected with SVCV. (I) KEGG

pathway enrichment map showing the top 10 signaling pathways of total. (J) GSEA of

differentially expressed genes in the liver of *cyp17a2*$^{+/+}$ and *cyp17a2*$^{-/-}$ male zebrafish

infected with SVCV and enrichment of RLR signaling pathway. The *p*-values were

corrected for the False Discovery Rate (FDR) method, and modules with *p*-values less

than 0.05 were selected as significant.

Figure S2 Heatmap view of mRNA variations of RLR-mediated ISGs sets in the

liver of *cyp17a2*$^{+/+}$ and *cyp17a2*$^{-/-}$ male zebrafish infected with SVCV. Expression

values are $\log_2^{(FPKM+1)}$ transformed and presented as row Z-scores.

Figure S3 Cyp17a2 enhances IFN production to inhibit viral replication. (A-D)

Luciferase activity of IFNφ3pro and ISRE in EPC cells transfected with indicated

plasmids for 24 h, and then untreated or transfected with poly I:C (0.5 μg) or infected

with SVCV (MOI = 1) for 24 h before luciferase assays. (E) qPCR analysis of *vig1* in

EPC cells transfected with indicated plasmids for 24 h, and then untreated or infected

with SVCV (MOI = 1) for 24 h. (F) Plaque assay of virus titers in EPC cells or GICB

cells transfected with indicated plasmids for 24 h, followed by SVCV or CyHV-2

challenge for 24 h or 48 h. (G and H) qPCR and IB analysis of SVCV genes in EPC

cells transfected with indicated plasmids for 24 h, followed by SVCV challenge for 24

h. (I) IF analysis of SVCV proteins in EPC cells transfected with indicated plasmids

for 24 h, followed by SVCV challenge for 24 h.

Figure S4 STING stability is enhanced by Cyp17a2. (A) Colocalization analyses of

figure 4G were performed by calculating Pearson correlation coefficient. (B)

Fluorescent analysis of proteins in EPC cells transfected with indicated plasmids for

24 h. The fluorescence intensity (arbitrary unit, a.u.) was recorded by the LAS X

software, and the data were expressed as mean ± SD, n = 5. (C) IB analysis of

proteins in EPC cells transfected with indicated plasmids for 24 h, followed by

untreated or transfected with poly I:C (2 μg) for 24 h. (D and E) IB analysis of

proteins in EPC cells transfected with indicated plasmids for 18 h, then treated with

CHX for 4 h and 8 h.

Figure S5 Cyp17a2 enhances STING-mediated IFN activation and antiviral

defense. (A and B) Luciferase activity of ISRE in EPC cells transfected with indicated

plasmids for 24 h before luciferase assays. (C and D) qPCR analysis of *vig1* in EPC

cells transfected with indicated plasmids for 24 h. (E and H) Plaque assay in EPC cells

transfected with indicated plasmids for 24 h and then infected with SVCV (MOI = 1)

for 24 h. (F and I) IF analysis of N protein and P protein in EPC cells transfected with

indicated plasmids for 24 h, followed by SVCV challenge for 24 h. The fluorescence

intensity (arbitrary unit, a.u.) was recorded by the LAS X software, and the data were

expressed as mean ± SD, n = 5. (G) IB analysis of proteins in EPC cells transfected

with indicated plasmids for 24 h.

1008

Figure S6 Cyp17a2 promotes STING stabilization by facilitating btr32-catalyzed

K33-linked polyubiquitination. (A) STING ubiquitination assays in EPC cells

transfected with indicated plasmids for 24 h. (B) IB analysis of WCLs and proteins

immunoprecipitated with anti-Flag Ab-conjugated agarose beads from EPC cells

transfected with indicated plasmids for 24 h. (C and D, G) IB analysis of proteins in

EPC cells transfected with indicated plasmids for 24 h. (E and F) Luciferase activity

of IFNφ1pro and qPCR analysis of *vig1* in EPC cells transfected with indicated

plasmids for 24 h.

1017

1018   **Figure S7 The K203 residue is essential for the ubiquitination of STING**

1019   **mediated by Cyp17a2.** (A-E) STING ubiquitination assays in EPC cells transfected

1020   with indicated plasmids for 24 h.

1021

Figure S8 Cyp17a2 mediates K33-linked polyubiquitination of the P protein at

K12. (A and C) IB analysis of WCLs and proteins immunoprecipitated with

anti-Flag/Myc Ab-conjugated agarose beads from EPC cells transfected with

indicated plasmids for 24 h. (B) Colocalization analyses of figure 9D were performed

by calculating Pearson correlation coefficient. (D) IB analysis of proteins in EPC cells

transfected with indicated plasmids for 24 h. (E-I) P protein ubiquitination assays in

EPC cells transfected with indicated plasmids for 18 h, followed by MG132

treatments for 6 h.

Table S1 List of primer sequences used in this study.

Table S2 List of all identified genes in FHK-vs-MHK in transcriptomic analysis.

Table S3 Heatmap view of mRNA variations of SVCV-activated ISG sets in the

Cyp17a2-overexpressing cells or *cyp17a2* knockdown cells.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

Figure 10