miRNA/mRNA analysis of increased TGF-β pathways drive epithelial-mesenchymal

transition and regulatory T cell differentiation

Running title: miRNA/mRNA analysis links TGFβ to EMT and Tregs in Chlamydia

Authors: Toni Darville[1#], Xuejun Sun[2#], Yu Zhang[2], Catherine M. O'Connell[1], Neha Mokashi[1,2],

Weiming Tang[3], Aakash Bhardwaj[1], Bryce Duncan[1], Charles W. Andrews, Jr.[4], Harold

Weisenfeld[5], and Xiaojing Zheng[1,2*]

  1. Department of Pediatrics, University of North Carolina at Chapel Hill, Chapel Hill, NC
  2. Department of Biostatistics, University of North Carolina at Chapel Hill, Chapel Hill, NC
  3. Department of Medicine, University of North Carolina at Chapel Hill, Chapel Hill, NC
  4. Pathology consultant, Newburyport, MA
  5. The University of Pittsburgh School of Medicine and the Magee-Womens Research Institute, Pittsburgh, PA

#: shared first authorship

*Correspondence: Xiaojing Zheng, PhD, University of North Carolina at Chapel Hill, 4018A Mary Ellen Jones Building, CB# 7509, 110 Manning Drive, Chapel Hill, NC 27599-7509, USA

19
20 *This revision updates the preprint to a non-reusable license that restricts redistribution and reuse; no changes were made to the scientific content*
21

Abstract – 249 words (limit 250 words)

Chlamydia trachomatis genital tract infection is linked to severe reproductive complications in women, including ectopic pregnancy, infertility, and adverse pregnancy outcomes. Mouse models of infection suggest that chlamydia-induced dysregulation of microRNAs (miRNAs) can drive harmful cytokine responses, pathogenic epithelial-mesenchymal transition (EMT), and fibrosis. To investigate these mechanisms in humans, we profiled miRNA and mRNA expression in endometrial biopsies from women with endometrial infection (Endo+) and compared them to profiles from women with cervix-only infection (Endo-) or no infection. Ingenuity Pathway Analysis (IPA) revealed that Endo+ tissues had upregulated genes associated with innate and adaptive immune response pathways, as well as EMT regulation, while downregulated genes were linked to cell cycle control. An integrative miRNA-mRNA analysis, which combined a review of published miRNA regulation in human infections and immune responses with IPA's miRNA target filter, identified differentially expressed miRNAs that modulate these pathways in the endometrium of Endo+ women. Functional annotation of these miRNAs showed a predominance of downregulated miRNAs that typically suppress EMT and regulatory T cell (Treg) differentiation, along with miRNAs that usually enhance Th17 responses. Comparisons with previously identified mRNA pathways in blood samples from women with endometrial *Chlamydia* infection indicated that alterations in TGF-β signaling and EMT were specific to the endometrium. Overall, the miRNA-mRNA interactions inferred from Endo+ tissue suggest increased activity in TGF-β pathways that promote enhanced EMT and Treg differentiation, while reducing Th17 activation. These changes highlight a dual potential for promoting tissue scarring while dampening inflammatory responses that could otherwise limit infection.

Introduction

Sexually transmitted *Chlamydia trachomatis* (Ct) genital infections represent a global public health challenge due to their high prevalence and severe reproductive health consequences. These include pelvic inflammatory disease (PID), chronic pelvic pain, infertility, and ectopic pregnancy (1, 2). Infertility and ectopic pregnancy after Ct infection is primarily caused by fibrotic scarring of the oviducts. Additionally, studies have linked Ct with adverse pregnancy outcomes such as stillbirth, infant death, spontaneous abortion, preterm labor, small-for-gestational-age infants, and postpartum endometritis (3). This suggests that Ct may cause lasting endometrial damage.

In the female reproductive tract, immune cells in the vagina and cervix provide the first line of defense against infection, whereas immune cells in the endometrium play a dual role: eliminating pathogens that breach the cervical barrier and maintaining immune tolerance of the embryo during pregnancy. Regulatory T (Treg) cells in the uterus are critical for immune tolerance to foreign fetal antigens(4, 5). Their secretion of anti-inflammatory cytokines such as IL-10 and TGF-β, and expression of inhibitory molecules like CTLA-4 and PD-1, suppress maternal immune responses that could otherwise target the fetus. In contrast, Th17 cells drive proinflammatory responses that can be harmful. The balance between these cell types is crucial for a healthy pregnancy. During Ct infection, the induction of Tregs may hinder the development of effective T cell immunity (6), whereas the activation of chlamydial-specific Th17 cells appears to enhance resistance to reinfection (7). Therefore, while a higher Treg/Th17 ratio supports reproductive health, it may compromise host defense against Ct.

Animal models have also shown that pro-inflammatory signaling and immune cell recruitment are initiated upon Ct infection of host epithelial cells (8, 9). Engagement of pathogen recognition receptors (PRRs) elicits a rapid influx of neutrophils that fail to kill the bacteria while releasing

tissue-damaging molecules such as reactive oxygen species and matrix metalloproteases (10-15).

Eventually, chlamydia-specific IFN-γ-producing CD4 T cells, activated by dendritic cells, resolve

infection. However, danger-associated molecular patterns (DAMPs) released by chlamydiae or

dying epithelial cells can perpetuate inflammation through induction of production of TNFα and

IL-1α by adjacent epithelium or influxing innate cells, creating a feed-forward loop of tissue

damage (11, 16, 17). Furthermore, epithelial-mesenchymal transition (EMT)—a dysregulated

tissue repair process controlled by microRNAs (miRNAs)—has been implicated in oviduct fibrosis

in chlamydia-infected mice (18). During EMT, epithelial cells lose cell-cell contacts and acquire

mesenchymal characteristics, including elevated production of extracellular matrix proteins,

before transitioning into myofibroblasts.

Mouse models of chlamydial infection demonstrated that the induction of EMT involves TNFα

signaling, caspase activation, and cleavage inactivation of Dicer (18, 19) an RNase III enzyme that

processes RNA into microRNAs. MicroRNAs (miRNAs) are small non-coding RNAs that play a

crucial role in post-transcriptional gene regulation, primarily through translational repression and

mRNA degradation. However, miRNAs can also enhance gene expression and translation under

certain conditions (20, 21). miRNAs are predicted to control the activity of approximately 30% to

50% of all protein-coding genes. A single miRNA can target multiple mRNAs, while multiple

miRNAs can collaborate to finely tune the expression of a single mRNA target (22). Chlamydia

infection in mice was associated with reduced expression of miRNAs that normally suppress EMT,

fibrosis and tumorigenesis, alongside increased expression of proteins associated with EMT and

fibrosis (18, 19)

Human salpingeal tissues are challenging to obtain but minimally invasive endometrial suction

catheter biopsies can be collected in an office setting (23). Histological evidence of endometritis

correlates with salpingitis observed via laparoscopy (24) and is associated with an increased risk

of infertility in women with symptomatic or asymptomatic Ct infection (25). Histologic features

resembling endometritis have also been described in surgically removed Fallopian tubes (26),

supporting the use of endometrial biopsies for investigating pathogenic mechanisms underlying

Ct-induced disease.

In this study, we conducted an integrated analysis of endometrial mRNA and miRNA profiles in

cisgender women with high exposure to Ct. Our findings identified an infection-driven mRNA

signature that highlights active innate and adaptive immune signaling pathways, along with

epithelial-mesenchymal transition (EMT), in women with endometrial Ct infection compared to

those with cervical Ct infection only and uninfected women. A parallel miRNA analysis revealed

significant downregulation of several miRNAs that typically suppress mRNAs involved in

activating TGF-β-related pathways, which drive EMT and regulatory T cell (Treg) differentiation.

Additionally, a partially overlapping subset of miRNAs that typically enhance proinflammatory

Th17 differentiation was also downregulated. The suppression of these miRNAs during

endometrial Ct infection appears to release their regulatory effects, facilitating EMT and increasing

the potential for tissue scarring. However, the miRNA-mediated shift in the Treg/Th17 balance,

favoring Tregs, may simultaneously act to dampen inflammation.

Materials and Methods

Ethics statement. This study adhered to the Declaration of Helsinki guidelines, and all participants provided written informed consent prior to participation. The study protocols were approved by the Institutional Review Boards for Human Subjects Research at the University of North Carolina and the University of Pittsburgh.

Study population. Endometrial biopsy samples were obtained from cisgender female participants enrolled in two independent cohorts with high exposure to Ct infection. Anaerobes and Clearance of Endometritis (ACE) Cohort: Participants were women clinically diagnosed with pelvic inflammatory disease (PID) (27, 28). Diagnostic criteria for enrollment included cervical motion tenderness, uterine tenderness, or adnexal tenderness observed during pelvic examination in sexually active young women experiencing pelvic or lower abdominal pain. T Cell Response Against Chlamydia (TRAC) Cohort: Participants were asymptomatic women identified as being at high risk for Ct infection (29). At enrollment, demographic, behavioral, and medical history data were collected. General physical and pelvic exams were conducted, and blood samples were obtained for immune studies. Participants underwent endometrial biopsy sampling using suction catheters. Additional assessments included Gram-stained vaginal smears for bacterial vaginosis using Nugent scores (30), as well as testing for Ct, *Neisseria gonorrhoeae* (Ng), and *Mycoplasma genitalium* (Mg) in cervical swabs and endometrial biopsies using nucleic acid amplification tests. Histopathological analyses of biopsies were performed (25, 31).

Participants were categorized into three primary groups based on infection status: (1) Endo+: Ct detected in both the endometrium and cervix. Endo+ women were further categorized into subgroups of women with clinical PID (Endo+, PID+), and asymptomatic women (Endo+, PID-). (2) Endo-: Ct detected in the cervix only, without endometrial infection. (3) Uninfected: No Ct

infection. For miRNA analyses, women with PID-like symptoms with negative tests for Ct, Ng,

and Mg were included as an additional comparison group.

Study design and workflow. The study workflow is depicted in **Figure 1**. Endometrial biopsy

samples were analyzed for mRNA and miRNA expression. Principal Component Analysis (PCA)

was used to explore transcriptomic relationships among cohort participants. Differentially

expressed (DE) mRNAs and miRNAs were identified between Endo+ and Endo-/Uninfected

groups. Functional annotations of DE mRNAs and miRNA-mRNA pairs were conducted using

Ingenuity Pathway Analysis (IPA). Endometrial mRNA data were also compared with previously

published blood mRNA data from the same cohorts (32).

mRNA and miRNA data collection and processing. mRNA Extraction and Profiling: Total RNA

was extracted from endometrial biopsies stored at -80°C in tubes containing RNA/DNA Shield

(Zymo Research, Irvine, CA). After thawing, samples were weighed, minced, and processed for

simultaneous DNA and RNA extraction using the Quick DNA/RNA™ isolation kit according to

the manufacturers protocol (Zymo) with on column DNAse I treatment prior to total RNA elution.

Libraries were prepared using the Ovation SoLo RNA-Seq kit (NuGen Technologies) and

sequenced on an Illumina HiSeq2500 platform in the High Throughput Sequencing Facility (HTSF)

at the University of North Carolina. Gene expression quantification used BBMap (v37.25) and

samtools (v1.4.1). Blood mRNA pathway data had been previously generated by microarray

hybridization analysis using samples obtained from ACE and TRAC participants (31, 32). miRNA

Extraction and Profiling: miRNAs were isolated using HTG EdgeSeq reagents according to the

user manuscript, and libraries were synthesized at the HTSF at the University of North Carolina.

Barcoded samples were pooled, and sequencing was performed on an Illumina NextSeq 500

platform using a 75-cycle High Output v2 kit. Count data were generated by the EdgeSeq parser software (HTG Molecular Diagnostics, Inc.).

Data accession. All endometrial mRNA and miRNA profiles have been deposited in the Gene Expression Omnibus database (https://www.ncbi.nlm.nih.gov/geo/) GEO accession number: GSE290615 for endometrial mRNA, and GSE289941 for endometrial miRNAs. The blood mRNA profiles were retrieved from the previous study (32) with GEO accession number GSE110106.

Statistical analysis. Demographic Data: One-way ANOVA was used for numerical variables, while Fisher's exact test or Chi-square test was applied for categorical variables. RNA-Seq Data: PCA with ComBat-seq (33) correction was applied to address batch effects. mRNAs with expression levels <20 in >50% of samples were excluded, leaving 16,183 mRNAs for analysis. miRNAs with expression levels below mean +3 standard deviations of negative controls were filtered, retaining 253 miRNAs. Normalization was performed using DESeq2 (v1.36.0) (34). PCA of variance by R was used to identify the inherent pattern of samples. Differential Expression Analysis: DE mRNAs and miRNAs were identified using DESeq2 with false discovery rate (FDR) <0.05. All analyses, unless specified, were completed in R (version 4.1.0). Within-sample group clustering heatmaps were generated to visualize DE miRNA expression across groups.

Functional annotation and pathway enrichment analysis of mRNAs. Ingenuity Pathway Analysis (IPA) (QIAGEN Inc., https://digitalinsights.qiagen.com/IPA) identified enriched pathways for DE mRNAs, using Fisher's exact test. Regulatory T cell-related mRNAs not included in IPA were sourced from Gene Ontology (GO) (35) and manually curated from literature.

Integrated miRNA-mRNA analysis. Putative target mRNAs of DE miRNAs were identified through IPA's "miRNA target filter" which incorporates both published experimental and high-

confidence computational data. mRNA targets were further filtered to include only those that were

also identified as DE mRNAs (FDR<0.05) in this study. We performed a functional annotation of

the negatively correlated miRNA-mRNA pairs to identify significantly enriched pathways.

Systematic review of miRNA functions. A systematic review identified studies examining DE

miRNA functions in human immune-related processes. PubMed searches from 1990 to 2024 used

the following search query: all miRNAs AND (infect OR immune OR autoimmune) AND (human

OR patient OR healthy donor) AND ("T cell" OR "B cell" OR lymphocyte OR CD4 OR CD8 OR

Treg OR Th1 OR Th2 OR Th17 OR Tfh OR neutrophil OR monocyte OR macrophage OR "NK

cell" OR "epithelial cell" OR "dendritic cell") AND (expression OR function OR Differentiation

OR activated OR activation OR inhibit OR suppress OR stimulation) NOT ("beta-cell" OR

methylation OR stem OR endothelial OR pregnancy). Tumor-related and non-immune studies

were excluded, leaving 361 full-text studies curated for miRNA effects on immune and epithelial

cell responses including directionality of their effects.

Comparison of DE mRNA enriched pathways detected in the endometrium versus blood. We

previously identified blood DE mRNAs from women with Ct-induced pelvic inflammatory disease

(PID) (Endo+) to those from asymptomatic women with cervical infection only and uninfected

women (Endo-/Uninfected) (32). Using Bonferroni-adjusted P values <0.05 as the significance

threshold, we compared the significantly enriched pathways for DE mRNAs from this endometrial

study and the aforementioned blood study.

Results

Characteristics of study participants. The characteristics of study participants included in the mRNA (N = 15) and miRNA (N = 23) analyses are summarized in **Supplementary Tables 1 and 2**. Participant group sizes were determined by the quality of mRNA and miRNA isolated from tissue specimens. Most participants were young, African American, unmarried, had some college education, and were insured by Medicaid. A majority reported previous Ct infection, had anti-Ct antibodies, and did not have bacterial vaginosis (BV) by Nugent's criteria (30) or coinfection with Ng and/or Mg. Demographic variables such as age, race, education, insurance status, and contraceptive use did not differ significantly among groups.

Distinct gene expression profiles in Endo+ women contrast with minimal impact of cervical Ct infection. Unsupervised principal component analysis (PCA) was performed to assess whether global gene expression profiles were associated with infection status or disease extent. PCA of mRNA data (**Figure 2A**) showed that most Endo+ women formed a distinct cluster, separate from the Endo- and Uninfected groups, with one outlier in the Endo+ group. Similarly, PCA of miRNA data (**Figure 2B**) indicated that Endo+ women clustered apart from the Endo- and Uninfected groups. However, PCA did not detect differences between endometrial mRNA or miRNA expression profiles of women with cervical Ct infection or those who were uninfected. The lack of separation between Endo- and Uninfected groups suggests that cervical Ct infection does not significantly impact endometrial gene expression.

Endometrial inflammation was detected in Endo+ and non-STI PID cases. Biopsied tissues from Endo+ women, both symptomatic and asymptomatic, and from women with non-STI-induced PID showed plasma cell and lymphocyte infiltrates in the endometrial stroma. Neutrophils were also observed within the basal lamina of the epithelium and in gland lumens in some Endo+ biopsies.

In contrast, 6 evaluable biopsis from women with cervical Ct infection only and 9 evauable

biopsies from uninfected women either lacked inflammatory cells or contained rare mononuclear

cells. The remaining biopsies had insufficient tissue for histological evaluation (**Figure 3A-D**).

These pathological data align with the data from PCA analysis indicating clear separation of

endometrial transcriptional profiles of Endo+ women from those with Ct infection limited to the

cervix.

Key transcriptional differences in Ct endometritis highlight induction of innate and adaptive

immune responses, EMT, and cell cycle dysregulation that is confirmed by functional analysis

of miRNA-mRNA interactions. Since PCA showed minimal variance between the Endo- and

Uninfected groups, their transcriptional data were combined and compared to the Endo+ group. A

total of 2,000 mRNAs were significantly DE between Endo+ and Endo-/Uninfected groups (FDR

< 0.05), with 1046 upregulated, and 954 downregulated genes in the Endo+ group. The top 100

upregulated and downregulated genes with FDR <0.05 are listed in **Supplementary Tables 3 and**

4, respectively and displayed in a volcano plot (**Figure 4**).

Several of the most strongly upregulated genes play crucial roles in local antibody production and

germinal center formation. These include transcription factors and coactivators such

as *BCL6* and *POU2AF1*, *CXCL13*, a key germinal center chemokine, and antibody Fc receptors

including *FCAR*, the IgA receptor, and *FCRL2*, which supports B cell antibody production

(**Supplementary Table 3**). This gene expression pattern was consistent with the marked plasma

cell infiltrates observed in women with PID, underscoring the hallmark histology of chronic

endometritis (**Figure 3A-B**) (36-40). Other highly upregulated genes are involved in the acute

phase response and cytokine cascade along with secondary mediators, including chemokines,

colony-stimulating factors, and prostaglandins, which amplify leukocyte recruitment and local

innate immunity (*IL1B*, *IL1RN*, *IRAK3*, *GSDMC*, *C2CD4A*, *CSF3R*, *CXCL1*, *SOD2*, *PTGS2*)

(**Supplementary Table 3**). *PTGS2* which encodes a cyclooxygenase enzyme that catalyzes

prostaglandin synthesis and is linked to pain, was up-regulated in Endo+ women, some of whom

exhibited PID symptoms.

Upregulated pathways identified by IPA included non-specific inflammatory responses (acute

phase response, PPARa/RXRa activation, *IL-6*, *PI3K/AKT*, *JAK/STAT*), innate (*TREM1*, NK cell,

nitrous oxide, phagosome formation, Toll-like receptor (TLR) and pattern recognition receptor

(PRR) signaling, granulocyte adhesion, and HMGB-1 signaling), and adaptive immune response

pathways (*IL-10*, Th1/Th2, CD40 signaling, IL-9 signaling) (**Figure 5A, Supplementary Table**

5).

IPA also identified EMT pathway enrichment, driven by upregulation of *TGFB2*, *TGFBR2*,

SMAD3, and *ZEB2* (**Figure 5A, Supplementary Table 5**). *TGFB2* interacts with its receptor to

activate SMAD proteins, including SMAD3. This activation promotes the expression of EMT-

related genes such as *ZEB2*, a transcription factor which represses epithelial markers like E-

cadherin and enhances mesenchymal markers, playing a pivotal role in EMT progression.

Downregulated genes included membrane adhesion components (*GJB6*, *CDH10*, *PCDH10*) and

tissue remodeling enzymes (*MMP26*, *FBN3*, *PLOD1*), as well as genes involved in transcription

regulation (*RXRG* and *SPDEF*), protein synthesis (*DHFR*, *GSTZ1*) and modification (*APRT*,

RYCR1), and cellular metabolism (*KMO*, *NDUFB7*, *UQCRC1*) (Supplementary Table 4). Multiple

genes related to histone metabolism were also decreased, potentially altering chromatin structure

and impacting gene expression regulation (**Figure 4 and Supplementary Tables 4 and 6**).

Downregulated pathways were enriched in genes involved with cell cycle regulation,

mitochondrial dysfunction, and nucleotide biosynthesis (**Figure 5B, Supplementary Table 6**).

Consistent with the DE mRNA results (**Figure 5A**), functional enrichment analysis using IPA

revealed that downregulated miRNA-mRNA interactions in Endo+ women enriched pathways

such as "Regulation of EMT by Growth Factors" (**Figure 5C**, **Supplementary Table 7**),

highlighting EMT promotion. Additionally, these interactions enriched pathways containing

upregulated innate and adaptive immune signaling genes (**Figure 5C**). In contrast, upregulated

miRNAs associated with downregulated mRNAs were enriched in pathways involving cell cycle

regulation and DNA damage signaling (**Figure 5D**), aligning with mRNA findings (**Figure 5B**

and Supplementary Table 8). These results suggest that Ct infection decreases the transcription

of miRNAs that normally suppress EMT and immune signaling genes, potentially accelerating

scarring and inflammation. Simultaneously, the infection increases the transcription of miRNAs

that downregulate mRNAs involved in mucosal epithelial homeostasis, possibly further impairing

uterine healing processes.

Differential expression of miRNAs highlights EMT promotion and Treg/Th17 modulation in

endometrial Ct infection. A total of 89 miRNAs were significantly DE (FDR<0.05) between

Endo+ and Endo- women. Among these, 47 miRNAs were upregulated, and 42 were

downregulated (**Supplementary Tables 9 and 10).** miRNAs with fold changes >2 and <1/2 are

depicted in **Figure 6A**. The majority of (53 of 89) DE miRNAs were associated with EMT with

32 miRNAs that normally dampen EMT being downregulated and 4 that increase EMT being

upregulated (**Figure 6B**), leading to an overall effect of enhancing EMT (**Table 1**).

IPA's target filter and literature review identified target mRNAs linked to TGF-β signaling, crucial

for EMT induction. These included *JAK1/STAT3* (103), Smad-dependent and (104) independent

TGF-β pathways, and the autocrine signaling network for EMT involving *TGF-β* and *ZEB2* (104).

Transcription factors (e.g., *ZEB2*, *FOXO1*) and pathways like PI3K-AKT were upregulated,

emphasizing EMT promotion. Among the 25 most significantly downregulated miRNAs in Endo+

samples, seven (miR-141-3p, miR-200a-3p, miR-106b-5p, miR-224-5p, miR-20a-5p, miR-21-5p,

and miR-223-3p) were identified through literature review as directly suppressing *TGF-β*

production. Consequently, their downregulation in Endo+ women likely enhances *TGF-β* activity,

which may suppress chronic inflammation but also promote EMT, potentially contributing to

adverse pathological outcomes.

Among the 89 differentially expressed (DE) miRNAs, 32 were associated with T cell regulation,

including subclasses such as Tregs, Th17, and Th1. *TGF-β* plays a key role in the differentiation

of both Tregs and Th17 cells. However, the presence of additional cytokines, particularly

interleukin-6 (*IL-6*), shifts differentiation toward Th17 cells. Specifically, low *TGF-β*

concentrations combined with *IL-6* promote Th17 differentiation, whereas high *TGF-β* levels

suppress *IL-6* signaling and favor Treg differentiation (105).

Twenty miRNAs were linked to Treg differentiation, with 12 downregulated miRNAs typically

functioning to suppress Tregs (**Figure 6C, Table 2**). Notably, miR-146a-5p, which promotes

Tregs, was upregulated, while miR-148a-3p, another Treg-enhancing miRNA, was reduced. These

expression patterns suggest an overall increase in Treg differentiation in Endo+ women.

For Th17 differentiation, 25 miRNAs were identified (**Figure 6D, Table 3**). Of these, 14 miRNAs

that typically enhance Th17 cells were significantly reduced, 10 of which also function to suppress

Tregs. Additionally, two miRNAs that normally inhibit Th17 differentiation were upregulated.

Conversely, six miRNAs that suppress Th17 differentiation were downregulated, and miR-149-

3p, which promotes Th17 cells, was increased. Overall, 16 of the 25 DE miRNAs are predicted to reduce Th17 differentiation, while 7 are associated with increased Th17 differentiation.

Several target mRNAs were linked to promoting Treg differentiation over Th17 cells. These include transcription factors such as *FOXO1*, *RUNX1*, *IRF1*, and *BATF*, as well as key *TGF-β* pathway genes (*TGFBR2*, *SMAD3*, *STAT5*, *STAT3*, and *IL-10R*). Collectively, these findings indicate a shift toward increased Treg activity and reduced Th17 differentiation. This shift may help limit tissue-damaging neutrophil responses but could come at the cost of impaired Th17/Th1-mediated protection against reinfection (7).

A within-sample group clustering heatmap of DE miRNAs (**Figure 7**) revealed distinct expression patterns across groups: Endo+ women with and without symptoms, Endo-, Uninfected, and women with non-STI-induced pelvic pain. One cluster of miRNAs was downregulated in Endo+ women (symptomatic and asymptomatic) compared to Endo-, Uninfected, and women with non-STI-induced pelvic pain. Literature review suggests these miRNAs typically suppress EMT (**Table 1**), implying their downregulation in Endo+ women may promote EMT, regardless of symptom presence. Notably, this downregulation was not observed in non-STI pelvic pain cases, suggesting a link to active Ct infection (**Figure 7**).

However, a minor cluster of miRNAs (mir-1273, mir-1285-5p, mir-1254, mir-4734, and mir-566) that normally dampen mRNAs promoting EMT (**Table 1, Figure 7**), was downregulated in Endo+, 4 of 5 Endo-, and 5 of 7 non-STI pelvic pain cases with recent prior Ct infection, indicating potential enhancement of EMT processes irrespective of current Ct ascension or symptom status. Each of these miRNAs have been studied in various cancer types, and reportedly have both positive (miR-1273 (137), miR-4734) and negative roles (miR-1285-5p (138), miR-1254 (139), miR-566 (140)) in promoting cell proliferation, invasion, and metastasis. Furthermore, four miRNAs (miR-

320b (102, 141), let-7c-5p and let-7b-5p (79)) reported to induce EMT and promote oncogenesis were upregulated in Endo+ women and those with non-STI-induced pelvic pain (**Table 1, Figure 7**).

Distinct immune and fibrosis-related pathways were detected in endometrial vs. blood mRNA profiles in women with chlamydial endometritis. In a previous study (32), we compared blood-derived mRNA profiles from women with Ct-induced pelvic inflammatory disease (PID) (Endo+) to those from asymptomatic women with cervical infection only and uninfected women (Endo-/Uninfected). A comparison of the top enriched pathways identified in blood with those found in the endometrium of Endo+ women revealed notable differences (**Table 4**). Shared pathways related to innate immune responses, including granulocyte adhesion, *TREM1* signaling, Toll-like receptor (TLR) signaling, integrin signaling, *IL-1*, *IL-10*, and *IL-6* response pathways, as well as *JAK/STAT* signaling. Endometrium--specific responses included natural killer (NK) cell signaling pathways, *IL-23*, *STAT3*, and Th17 activation pathways which can lead to chronic inflammation if excessively activated (142), multiple adaptive T cell response pathways (Th1/Th17, Th2, and *IL-7*), and fibrosis-related pathways, *TGF-β* signaling and EMT. These sharply contrasted with the blood profiles of Endo+ women, where TCR signaling pathways—including mTOR, which integrates immune signals and metabolic cues for T cell maintenance and activation, and *ICOS-ICOSL* signaling which delivers the co-stimulatory signals that promote T cell activation and differentiation—were significantly downregulated, and activation of EMT pathways was undetected. Conversely, type I interferon signaling pathways were prominent in blood-borne profiles, but displayed minimal enrichment in the endometrium.

Discussion

The findings from this study improve understanding of how Ct infection modulates the molecular landscape of the human endometrium, emphasizing the interplay between immune responses, EMT, and regulatory miRNA activity. These results provide insight into how Ct infection may contribute to adverse pregnancy outcomes, including stillbirth and preterm labor, as well as the Ct-induced complication of chronic pelvic pain.

The integrated transcriptomic and miRNA analyses revealed distinct molecular signatures in the endometrial tissues of women with endometrial Ct infection (Endo+), highlighting active innate and adaptive immune signaling pathways alongside EMT promotion. The transcriptional patterns of upregulated genes associated with the acute phase response as well as genes that promote antibody-producing cells and cell-mediated immunity correlate with histological findings of endometritis, marked by a predominance of plasma cell infiltrates mixed with subepithelial neutrophils and lymphocytes. Other upregulated genes in Endo+ women, such as *TGFB2*, *TGFBR2*, and *SMAD3*, collectively drive EMT through Smad-dependent and independent mechanisms, while transcription factors like *ZEB2* suppress epithelial markers to facilitate mesenchymal transitions. The significant downregulation of EMT-suppressing miRNAs, including miR-141-3p and miR-200a-3p, further supports this shift toward EMT, aligning with prior observations in murine models (18, 19, 143). These molecular alterations are consistent with enhanced tissue remodeling, fibrosis, and scarring—hallmarks of the Ct-induced complications of tubal factor infertility and ectopic pregnancy. The additional detection of upregulated miRNAs that depress gene pathways engaged in nucleotide synthesis and cell cycle control indicate a further mechanism whereby miRNA dysregulation could inhibit healing of infected endometrial tissue. While this study focused on endometrial biopsy tissues, it is highly likely that similar mechanisms

are at work in the delicate oviduct. EMT and fibrosis in the oviduct could impair critical

reproductive processes, including gamete transport, fertilization, early embryo development,

immune defense, and structural integrity—all of which are essential for successful reproduction.

The enrichment of *TGF-β* signaling and its downstream transcription factors, such as *ZEB2*, in

Endo+ samples underscores the role of this pathway in driving EMT. Downregulated miRNAs that

typically suppress *TGF-β* signaling, including miR-141-3p and miR-200a-3p, likely amplify this

effect. These molecular changes are consistent with previous studies in mouse models, which

demonstrated that dysregulated EMT contributes to fibrosis and oviductal scarring. Igietseme et

al. (18) showed that *Chlamydia muridarum*-induced TNFα signaling and caspase activation were

linked to the downregulation of miRNAs that typically inhibit EMT and fibrosis. Similarly,

alterations in *TGF-β*, *TNF*, *ZEB*, and the miR-200 family observed in Endo+ women were also

identified in infertile mice, with multiple differentially expressed miRNAs involved in EMT

overlapping between the two studies (143, 144) demonstrated significant downregulation of

miRNAs that suppress fibrosis, including members of the miR-200 family in mice with oviduct

pathology following *C. muridarum* infection. This overlap in miRNA dysregulation between

human and mouse studies suggests a conserved *Chlamydia*-induced EMT mechanism across

species. The downregulation of these miRNAs, combined with the activation of *TGF-β* signaling,

likely creates a microenvironment conducive to tissue remodeling and scarring. This process may

represent a critical step in the progression from infection to tubal factor infertility.

A notable finding was the miRNA-driven shift in the Treg/Th17 balance. Among the 89 DE

miRNAs, many were linked to immune cell regulation, particularly promoting Treg differentiation

and function. The downregulation of miRNAs that typically suppress Tregs (e.g., miR-200a-

3p, miR-374b-5p) suggests an enhanced Treg response, potentially favoring immune tolerance at

the expense of effective Th1/Th17-mediated immunity. This interpretation is bolstered by the

upregulation of miR-146a-5p, known to support Treg function, and the increased expression of

Treg-associated mRNAs, such as TGFBR2 and Smad3.

Conversely, the downregulation of Th17-promoting miRNAs, such as miR-21-5p, miR-19b-

3p, and miR-223-3p, indicates a reduction in Th17 activity. Since Th17 cells enhance neutrophil

recruitment, decreased Th17 activity could mitigate tissue-damaging neutrophil responses, key

drivers of chlamydial disease pathogenesis (15). However, given Ct's intracellular developmental

cycle, and its production of Chlamydial Protease Activation Factor (CPAF), which impairs

neutrophil function (145), neutrophils have limited efficacy in eliminating Ct, and reduced

neutrophil activation may have little impact on infection resolution. However, Th17 cells exhibit

plasticity, transitioning between Th17 and Th1 phenotypes, and have been strongly associated with

resistance to chlamydial reinfection (7). Thus, the observed shift in the Treg/Th17 balance has

contrasting implications. Enhanced Treg/Th17 activity may reduce chronic inflammation and

preserve endometrial function; while simultaenously, impeding development of protective

immunity, potentially contributing to the high rates of reinfection observed in some cohort

participants (7). This delicate trade-off underscores the challenges of therapeutically targeting

these pathways, as interventions could disrupt the balance between immune suppression and

effective defense.

Comparative analyses between endometrial and previously published blood mRNA profiles

highlight tissue-specific responses to Ct infection. While blood profiles revealed activation of

interferon-mediated signaling and downregulation of adaptive T cell pathwys, endometrial

responses were dominated by activation of T cells (e.g., Treg, Th1, Th17), EMT, and *TGF-β*-

driven immune regulation. These differences underscore the localized nature of Ct-induced

endometrial inflammation and fibrosis, which may not be readily detectable through systemic biomarkers. This localized EMT induction aligns with the histological evidence of fibrosis observed in Ct-affected Fallopian tubes, suggesting a direct link between Ct gene regulation in the upper genital tract and long-term reproductive sequelae. These findings underscore the importance of localized studies for understanding the distinct molecular landscape of Ct-infected reproductive tissues.

We acknowledge several limitations of this study. The sample size for the endometrial transcriptome was constrained by the difficulty of obtaining sufficient high-quality endometrial tissue for mRNA and miRNA profiling. Additionally, the cross-sectional design of the study limits our ability to assess the temporal progression of infection and associated pathology. Future studies with larger sample sizes and longitudinal designs are needed to validate these findings and provide a more comprehensive understanding into the progression of Ct-induced pathology and the persistence of molecular changes post-infection. Extending these analyses to other sexually transmitted infections could determine whether similar miRNA-mediated mechanisms drive reproductive tract inflammation and scarring across different pathogens.

Despite these limitations, our study provides a new and comprehensive view of molecular changes in the endometrium during Ct infection, highlighting critical pathways that drive tissue remodeling, fibrosis, and immune dysregulation. The findings also raise concerns about asymptomatic Endo+ women, who exhibited similar molecular profiles to symptomatic individuals, suggesting that subclinical endometrial inflammation may contribute to reproductive morbidity, which supports prior studies indicating that asymptomatic women with Ct-induced endometritis have an increased risk for infertility (25).

In summary, this study demonstrates that endometrial Ct infection induces a miRNA-mediated

regulatory network that promotes EMT and alters immune cell balance, favoring Treg

differentiation over Th17 responses. These molecular changes likely contribute to tissue scarring

and impaired immune protection, offering new insights into the pathogenesis of Ct-induced

reproductive morbidity. Targeting these pathways may provide a foundation for innovative

therapeutic strategies to mitigate the long-term consequences of Ct infection.

Disclosures

The authors have no financial conflicts of interest.

Acknowledgment

We thank the women who agreed to participate in this study; Ingrid Macio, Melinda Petrina, Carol

Priest, Abi Jett, and Lorna Rabe, for their efforts in the clinic and the microbiology laboratory; and

the staff at the Allegheny County Health Department STD Clinic. We gratefully acknowledge the

technical support from the UNC High Throughput Sequencing Facility.

Footnotes

This work was supported by the National Institute of Allergy and Infectious Diseases
U19AI144181 and R01AI170959.

References

1. Chow, J. M., M. L. Yonekura, G. A. Richwald, S. Greenland, R. L. Sweet, and J. Schachter. 1990. The association between Chlamydia trachomatis and ectopic pregnancy. A matched-pair, case-control study. *JAMA* 263: 3164-3167.
2. Passos, L. G., P. Terraciano, N. Wolf, F. D. S. Oliveira, I. Almeida, and E. P. Passos. 2022. The Correlation between Chlamydia Trachomatis and Female Infertility: A Systematic Review. *Rev Bras Ginecol Obstet* 44: 614-620.
3. Tang, W., J. Mao, K. T. Li, J. S. Walker, R. Chou, R. Fu, W. Chen, T. Darville, J. Klausner, and J. D. Tucker. 2020. Pregnancy and fertility-related adverse outcomes associated with Chlamydia trachomatis infection: a global systematic review and meta-analysis. *Sex Transm Infect* 96: 322-329.
4. Huang, N., H. Chi, and J. Qiao. 2020. Role of Regulatory T Cells in Regulating Fetal-Maternal Immune Tolerance in Healthy Pregnancies and Reproductive Diseases. *Front Immunol* 11: 1023.
5. La Rocca, C., F. Carbone, S. Longobardi, and G. Matarese. 2014. The immunology of pregnancy: regulatory T cells control maternal immune tolerance toward the fetus. *Immunol Lett* 162: 41-48.
6. Stary, G., A. Olive, A. F. Radovic-Moreno, D. Gondek, D. Alvarez, P. A. Basto, M. Perro, V. D. Vrbanac, A. M. Tager, J. Shi, J. A. Yethon, O. C. Farokhzad, R. Langer, M. N. Starnbach, and U. H. von Andrian. 2015. VACCINES. A mucosal vaccine against Chlamydia trachomatis generates two waves of protective memory T cells. *Science* 348: aaa8205.
7. Yount, K. S., C. J. Chen, A. Kollipara, C. Liu, N. V. Mokashi, X. Zheng, C. B. Bagwell, T. B. Poston, H. C. Wiesenfeld, S. L. Hillier, C. M. O'Connell, N. Stanley, and T. Darville. 2024. Unique T cell signatures associated with reduced Chlamydia trachomatis reinfection in a highly exposed cohort. *bioRxiv*.
8. Darville, T., and T. J. Hiltke. 2010. Pathogenesis of genital tract disease due to Chlamydia trachomatis. *J Infect Dis* 201 Suppl 2: S114-125.
9. Rank, R. G., and J. A. Whittum-Hudson. 2010. Protective immunity to chlamydial genital infection: evidence from animal studies. *J Infect Dis* 201 Suppl 2: S168-177.
10. Hart, K. M., A. J. Murphy, K. T. Barrett, C. R. Wira, P. M. Guyre, and P. A. Pioli. 2009. Functional expression of pattern recognition receptors in tissues of the human female reproductive tract. *J Reprod Immunol* 80: 33-40.
11. Darville, T., J. M. O'Neill, C. W. Andrews, Jr., U. M. Nagarajan, L. Stahl, and D. M. Ojcius. 2003. Toll-like receptor-2, but not Toll-like receptor-4, is essential for development of oviduct pathology in chlamydial genital tract infection. *J Immunol* 171: 6187-6197.
12. Frazer, L. C., C. M. O'Connell, C. W. Andrews, Jr., M. A. Zurenski, and T. Darville. 2011. Enhanced neutrophil longevity and recruitment contribute to the severity of oviduct pathology during Chlamydia muridarum infection. *Infect Immun* 79: 4029-4041.
13. Lee, H. Y., J. H. Schripsema, I. M. Sigar, C. M. Murray, S. R. Lacy, and K. H. Ramsey. 2010. A link between neutrophils and chronic disease manifestations of Chlamydia muridarum urogenital infection of mice. *FEMS Immunol Med Microbiol* 59: 108-116.

14. Rank, R. G., J. Whittimore, A. K. Bowlin, S. Dessus-Babus, and P. B. Wyrick. 2008.
Chlamydiae and polymorphonuclear leukocytes: unlikely allies in the spread of
chlamydial infection. *FEMS Immunol Med Microbiol* 54: 104-113.

15. Ramsey, K. H., I. M. Sigar, J. H. Schripsema, N. Shaba, and K. P. Cohoon. 2005.
Expression of matrix metalloproteinases subsequent to urogenital Chlamydia muridarum
infection of mice. *Infect Immun* 73: 6962-6973.

16. Murthy, A. K., W. Li, B. K. Chaganty, S. Kamalakaran, M. N. Guentzel, J. Seshu, T. G.
Forsthuber, G. Zhong, and B. P. Arulanandam. 2011. Tumor necrosis factor alpha
production from CD8+ T cells mediates oviduct pathological sequelae following primary
genital Chlamydia muridarum infection. *Infect Immun* 79: 2928-2935.

17. Gyorke, C. E., A. Kollipara, J. t. Allen, Y. Zhang, J. A. Ezzell, T. Darville, S. A.
Montgomery, and U. M. Nagarajan. 2020. IL-1alpha Is Essential for Oviduct Pathology
during Genital Chlamydial Infection in Mice. *J Immunol* 205: 3037-3049.

18. Igietseme, J. U., Y. Omosun, O. Stuchlik, M. S. Reed, J. Partin, Q. He, K. Joseph, D.
Ellerson, B. Bollweg, Z. George, F. O. Eko, C. Bandea, H. Liu, G. Yang, W. J. Shieh, J.
Pohl, K. Karem, and C. M. Black. 2015. Role of Epithelial-Mesenchyme Transition in
Chlamydia Pathogenesis. *PLoS One* 10: e0145198.

19. Igietseme, J. U., Y. Omosun, T. Nagy, O. Stuchlik, M. S. Reed, Q. He, J. Partin, K.
Joseph, D. Ellerson, Z. George, J. Goldstein, F. O. Eko, C. Bandea, J. Pohl, and C. M.
Black. 2018. Molecular Pathogenesis of Chlamydia Disease Complications: Epithelial-
Mesenchymal Transition and Fibrosis. *Infect Immun* 86.

20. Filipowicz, W., S. N. Bhattacharyya, and N. Sonenberg. 2008. Mechanisms of post-
transcriptional regulation by microRNAs: are the answers in sight? *Nat Rev Genet* 9: 102-
114.

21. O'Brien, J., H. Hayder, Y. Zayed, and C. Peng. 2018. Overview of MicroRNA
Biogenesis, Mechanisms of Actions, and Circulation. *Front Endocrinol (Lausanne)* 9:
402.

22. Lewis, B. P., I. H. Shih, M. W. Jones-Rhoades, D. P. Bartel, and C. B. Burge. 2003.
Prediction of mammalian microRNA targets. *Cell* 115: 787-798.

23. Dacco, M. D., M. Moustafa, D. Papoutsis, D. Georgantzis, G. Halmos, and A. Magos.
2012. An audit of using the H Pipelle for endometrial sampling at outpatient
hysteroscopy and literature review comparison with the Pipelle de Cornier. *Eur J Obstet*
Gynecol Reprod Biol 165: 299-301.

24. Kiviat, N. B., P. Wolner-Hanssen, D. A. Eschenbach, J. N. Wasserheit, J. A. Paavonen,
T. A. Bell, C. W. Critchlow, W. E. Stamm, D. E. Moore, and K. K. Holmes. 1990.
Endometrial histopathology in patients with culture-proved upper genital tract infection
and laparoscopically diagnosed acute salpingitis. *Am J Surg Pathol* 14: 167-175.

25. Wiesenfeld, H. C., S. L. Hillier, L. A. Meyn, A. J. Amortegui, and R. L. Sweet. 2012.
Subclinical pelvic inflammatory disease and infertility. *Obstet Gynecol* 120: 37-43.

26. Hunt, J. L., and A. A. Lynn. 2002. Histologic features of surgically removed fallopian
tubes. *Arch Pathol Lab Med* 126: 951-955.

27. Wiesenfeld, H. C., L. A. Meyn, T. Darville, I. S. Macio, and S. L. Hillier. 2021. A
Randomized Controlled Trial of Ceftriaxone and Doxycycline, With or Without
Metronidazole, for the Treatment of Acute Pelvic Inflammatory Disease. *Clin Infect Dis*
72: 1181-1189.

28. Workowski, K. A., G. A. Bolan, C. Centers for Disease, and Prevention. 2015. Sexually
transmitted diseases treatment guidelines, 2015. *MMWR Recomm Rep* 64: 1-137.
29. Russell, A. N., X. Zheng, C. M. O'Connell, B. D. Taylor, H. C. Wiesenfeld, S. L. Hillier,
W. Zhong, and T. Darville. 2016. Analysis of Factors Driving Incident and Ascending
Infection and the Role of Serum Antibody in Chlamydia trachomatis Genital Tract
Infection. *J Infect Dis* 213: 523-531.
30. Nugent, R. P., M. A. Krohn, and S. L. Hillier. 1991. Reliability of diagnosing bacterial
vaginosis is improved by a standardized method of gram stain interpretation. *J Clin*
Microbiol 29: 297-301.
31. Wiesenfeld, H. C., R. L. Sweet, R. B. Ness, M. A. Krohn, A. J. Amortegui, and S. L.
Hillier. 2005. Comparison of acute and subclinical pelvic inflammatory disease. *Sex*
Transm Dis 32: 400-405.
32. Zheng, X., C. M. O'Connell, W. Zhong, U. M. Nagarajan, M. Tripathy, D. Lee, A. N.
Russell, H. Wiesenfeld, S. Hillier, and T. Darville. 2018. Discovery of Blood
Transcriptional Endotypes in Women with Pelvic Inflammatory Disease. *J Immunol* 200:
2941-2956.
33. Zhang, Y., G. Parmigiani, and W. E. Johnson. 2020. ComBat-seq: batch effect
adjustment for RNA-seq count data. *NAR Genom Bioinform* 2: lqaa078.
34. Love, M. I., W. Huber, and S. Anders. 2014. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol* 15: 550.
35. Gene Ontology, C., S. A. Aleksander, J. Balhoff, S. Carbon, J. M. Cherry, H. J. Drabkin,
D. Ebert, M. Feuermann, P. Gaudet, N. L. Harris, D. P. Hill, R. Lee, H. Mi, S. Moxon, C.
J. Mungall, A. Muruganugan, T. Mushayahama, P. W. Sternberg, P. D. Thomas, ..., and
M. Westerfield. 2023. The Gene Ontology knowledgebase in 2023. *Genetics* 224.
36. Rangel-Moreno, J., J. Moyron-Quiroz, K. Kusser, L. Hartson, H. Nakano, and T. D.
Randall. 2005. Role of CXC chemokine ligand 13, CC chemokine ligand (CCL) 19, and
CCL21 in the organization and function of nasal-associated lymphoid tissue. *J Immunol*
175: 4904-4913.
37. Carlsen, H. S., E. S. Baekkevold, F. E. Johansen, G. Haraldsen, and P. Brandtzaeg. 2002.
B cell attracting chemokine 1 (CXCL13) and its receptor CXCR5 are expressed in
normal and aberrant gut associated lymphoid tissue. *Gut* 51: 364-371.
38. Tolnay, M. 2022. Lymphocytes sense antibodies through human FCRL proteins:
Emerging roles in mucosal immunity. *J Leukoc Biol* 111: 477-487.
39. King, H. W., K. L. Wells, Z. Shipony, A. S. Kathiria, L. E. Wagar, C. Lareau, N. Orban,
R. Capasso, M. M. Davis, L. M. Steinmetz, L. K. James, and W. J. Greenleaf. 2021.
Integrated single-cell transcriptomics and epigenomics reveals strong germinal center-
associated etiology of autoimmune risk loci. *Sci Immunol* 6: eabh3768.
40. Wang, H., S. Jain, P. Li, J. X. Lin, J. Oh, C. Qi, Y. Gao, J. Sun, T. Sakai, Z. Naghashfar,
S. Abbasi, A. L. Kovalchuk, S. Bolland, S. L. Nutt, W. J. Leonard, and H. C. Morse, 3rd.
2019. Transcription factors IRF8 and PU.1 are required for follicular B cell development
and BCL6-driven germinal center responses. *Proc Natl Acad Sci U S A* 116: 9511-9520.
41. Di Ruocco, F., V. Basso, M. Rivoire, P. Mehlen, J. Ambati, S. De Falco, and V. Tarallo.
2018. Alu RNA accumulation induces epithelial-to-mesenchymal transition by
modulating miR-566 and is associated with cancer progression. *Oncogene* 37: 627-637.
42. Feng, J., S. Hu, K. Liu, G. Sun, and Y. Zhang. 2022. The Role of MicroRNA in the
Regulation of Tumor Epithelial-Mesenchymal Transition. *Cells* 11.

43.   Wang, S., M. Zhang, T. Zhang, J. Deng, X. Xia, and X. Fang. 2020. microRNA-141
inhibits TGF-beta1-induced epithelial-to-mesenchymal transition through inhibition of
the TGF-beta1/SMAD2 signalling pathway in endometriosis. *Arch Gynecol Obstet* 301:
707-714.
44.   Sun, Y., S. Shen, X. Liu, H. Tang, Z. Wang, Z. Yu, X. Li, and M. Wu. 2014. MiR-429
inhibits cells growth and invasion and regulates EMT-related marker genes by targeting
Onecut2 in colorectal carcinoma. *Mol Cell Biochem* 390: 19-30.
45.   Zhang, L., Y. Chen, H. Wang, X. Zheng, C. Li, and Z. Han. 2018. miR-376a inhibits
breast cancer cell progression by targeting neuropilin-1 NR. *Onco Targets Ther* 11: 5293-
5302.
46.   Lin, S., L. Tan, D. Luo, X. Peng, Y. Zhu, and H. Li. 2019. Linc01278 inhibits the
development of papillary thyroid carcinoma by regulating miR-376c-3p/DNM3 axis.
Cancer Manag Res 11: 8557-8569.
47.   Zhao, X., X. Zhang, X. Zhang, T. Jiang, J. Zhai, H. Wang, M. Huang, R. Lang, and Q.
He. 2021. MiR-374b-5p inhibits KDM5B-induced epithelial-mesenchymal transition in
pancreatic cancer. *Am J Cancer Res* 11: 3907-3920.
48.   Song, K., Y. Jiang, Y. Zhao, Y. Xie, J. Zhou, W. Yu, and Q. Wang. 2020. Members of
the miR-30 family inhibit the epithelial-to-mesenchymal transition of non-small-cell lung
cancer cells by suppressing XB130 expression levels. *Oncol Lett* 20: 68.
49.   Lu, J., Z. Zhao, and Y. Ma. 2020. miR-186 Represses Proliferation, Migration, Invasion,
and EMT of Hepatocellular Carcinoma via Directly Targeting CDK6. *Oncol Res* 28: 509-
518.
50.   Tian, X., Q. Fei, M. Du, H. Zhu, J. Ye, L. Qian, Z. Lu, W. Zhang, Y. Wang, F. Peng, J.
Chen, B. Liu, Q. Li, X. He, and L. Yin. 2019. miR-130a-3p regulated TGF-beta1-induced
epithelial-mesenchymal transition depends on SMAD4 in EC-1 cells. *Cancer Med* 8:
1197-1208.
51.   Du, W., H. Tang, Z. Lei, J. Zhu, Y. Zeng, Z. Liu, and J. A. Huang. 2019. miR-335-5p
inhibits TGF-beta1-induced epithelial-mesenchymal transition in non-small cell lung
cancer via ROCK1. *Respir Res* 20: 225.
52.   Fang, L. L., X. H. Wang, B. F. Sun, X. D. Zhang, X. H. Zhu, Z. J. Yu, and H. Luo. 2017.
Expression, regulation and mechanism of action of the miR-17-92 cluster in tumor cells
(Review). *Int J Mol Med* 40: 1624-1630.
53.   Qi, J. C., Z. Yang, Y. P. Zhang, B. S. Lu, Y. W. Yin, K. L. Liu, W. Y. Xue, C. B. Qu, and
W. Li. 2019. miR-20b-5p, TGFBR2, and E2F1 Form a Regulatory Loop to Participate in
Epithelial to Mesenchymal Transition in Prostate Cancer. *Front Oncol* 9: 1535.
54.   Khalilian, S., H. Abedinlou, B. M. Hussen, S. Z. H. Imani, and S. Ghafouri-Fard. 2022.
The emerging role of miR-20b in human cancer and other disorders: Pathophysiology and
therapeutic implications. *Front Oncol* 12: 985457.
55.   Morizane, R., S. Fujii, T. Monkawa, K. Hiratsuka, S. Yamaguchi, K. Homma, and H.
Itoh. 2014. miR-34c attenuates epithelial-mesenchymal transition and kidney fibrosis
with ureteral obstruction. *Sci Rep* 4: 4578.
56.   Li, C., J. Xia, W. Yao, G. Yang, Y. Tian, Y. Qi, and C. Hao. 2022. Mechanism of
LncRNA XIST/ miR-101-3p/ZEB1 axis in EMT associated with silicosis. *Toxicol Lett*
360: 11-19.

57.   Dong, P., M. Kaneuchi, H. Watari, J. Hamada, S. Sudo, J. Ju, and N. Sakuragi. 2011.
655   MicroRNA-194 inhibits epithelial to mesenchymal transition of endometrial cancer cells
656   by targeting oncogene BMI-1. *Mol Cancer* 10: 99.

657   58.   Chen, J., S. Gao, C. Wang, Z. Wang, H. Zhang, K. Huang, B. Zhou, H. Li, Z. Yu, J. Wu,
658   and C. Chen. 2016. Pathologically decreased expression of miR-193a contributes to
659   metastasis by targeting WT1-E-cadherin axis in non-small cell lung cancers. *J Exp Clin*
660   *Cancer Res* 35: 173.

661   59.   Li, J., Y. Song, Y. Wang, J. Luo, and W. Yu. 2013. MicroRNA-148a suppresses
662   epithelial-to-mesenchymal transition by targeting ROCK1 in non-small cell lung cancer
663   cells. *Mol Cell Biochem* 380: 277-282.

664   60.   Hu, B., Z. Chen, X. Wang, F. Chen, Z. Song, and C. Cao. 2021. MicroRNA-148a-3p
665   Directly Targets SERPINE1 to Suppress EMT-Mediated Colon Adenocarcinoma
666   Progression. *Cancer Manag Res* 13: 6349-6362.

667   61.   Miscianinov, V., A. Martello, L. Rose, E. Parish, B. Cathcart, T. Mitic, G. A. Gray, M.
668   Meloni, A. Al Haj Zen, and A. Caporali. 2018. MicroRNA-148b Targets the TGF-beta
669   Pathway to Regulate Angiogenesis and Endothelial-to-Mesenchymal Transition during
670   Skin Wound Healing. *Mol Ther* 26: 1996-2007.

671   62.   Jia, Z., Y. Zhang, Q. Xu, W. Guo, and A. Guo. 2018. miR-126 suppresses epithelial-to-
672   mesenchymal transition and metastasis by targeting PI3K/AKT/Snail signaling of lung
673   cancer cells. *Oncol Lett* 15: 7369-7375.

674   63.   Jiang, R., C. Zhang, G. Liu, R. Gu, and H. Wu. 2017. MicroRNA-126 Inhibits
675   Proliferation, Migration, Invasion, and EMT in Osteosarcoma by Targeting ZEB1. *J Cell*
676   *Biochem* 118: 3765-3774.

677   64.   Yao, Y., Q. Xu, L. Yan, Y. Jiao, Q. Su, X. Li, C. Liu, and F. Zhao. 2020. MiRNA-128
678   and MiRNA-142 Regulate Tumorigenesis and EMT in Oral Squamous Cell Carcinoma
679   Through HOXA10. *Cancer Manag Res* 12: 9987-9997.

680   65.   Pao, S. I., L. T. Lin, Y. H. Chen, C. L. Chen, and J. T. Chen. 2021. Repression of Smad4
681   by MicroRNA-1285 moderates TGF-beta-induced epithelial-mesenchymal transition in
682   proliferative vitreoretinopathy. *PLoS One* 16: e0254873.

683   66.   Jiang, M., L. Shi, C. Yang, Y. Ge, L. Lin, H. Fan, Y. He, D. Zhang, Y. Miao, and L.
684   Yang. 2019. miR-1254 inhibits cell proliferation, migration, and invasion by down-
685   regulating Smurf1 in gastric cancer. *Cell Death Dis* 10: 32.

686   67.   Yan, S., M. Cheng, Q. Duan, Z. Wang, W. Gao, B. Ren, and D. Xu. 2019. MiR-6803-5p
687   Promotes Cancer Cell Proliferation and Invasion via PTPRO/NF-kappaB Axis in
688   Colorectal Cancer. *Mediators Inflamm* 2019: 8128501.

689   68.   Zhu, X., S. Ju, F. Yuan, G. Chen, Y. Shu, C. Li, Y. Xu, J. Luo, and L. Xia. 2017.
690   microRNA-664 enhances proliferation, migration and invasion of lung cancer cells. *Exp*
691   *Ther Med* 13: 3555-3562.

692   69.   You, X., Y. Wang, J. Meng, S. Han, L. Liu, Y. Sun, J. Zhang, S. Sun, X. Li, W. Sun, Y.
693   Dong, and Y. Zhang. 2021. Exosomal miR-663b exposed to TGF-beta1 promotes
694   cervical cancer metastasis and epithelial-mesenchymal transition by targeting MGAT3.
695   *Oncol Rep* 45.

696   70.   Hamada, S., K. Satoh, S. Miura, M. Hirota, A. Kanno, A. Masamune, K. Kikuta, K.
697   Kume, J. Unno, S. Egawa, F. Motoi, M. Unno, and T. Shimosegawa. 2013. miR-197
698   induces epithelial-mesenchymal transition in pancreatic cancer cells by targeting p120
699   catenin. *J Cell Physiol* 228: 1255-1263.

71. Miao, L. J., S. Yan, Q. F. Zhuang, Q. Y. Mao, D. Xue, X. Z. He, and J. P. Chen. 2019. miR-106b promotes proliferation and invasion by targeting Capicua through MAPK signaling in renal carcinoma cancer. *Onco Targets Ther* 12: 3595-3607.

72. Luan, W., Y. Ding, H. Xi, H. Ruan, F. Lu, S. Ma, and J. Wang. 2021. Exosomal miR-106b-5p derived from melanoma cell promotes primary melanocytes epithelial-mesenchymal transition through targeting EphA4. *J Exp Clin Cancer Res* 40: 107.

73. Yang, C., R. Dou, C. Wei, K. Liu, D. Shi, C. Zhang, Q. Liu, S. Wang, and B. Xiong. 2021. Tumor-derived exosomal microRNA-106b-5p activates EMT-cancer cell and M2-subtype TAM interaction to facilitate CRC metastasis. *Mol Ther* 29: 2088-2107.

74. Wang, S., X. Zhang, C. Yang, and S. Xu. 2019. MicroRNA-198-5p inhibits the migration and invasion of non-small lung cancer cells by targeting fucosyltransferase 8. *Clin Exp Pharmacol Physiol* 46: 955-967.

75. Wang, H., D. Kanmangne, R. Li, Z. Qian, X. Xia, X. Wang, and T. Wang. 2020. miR-30a-3p suppresses the proliferation and migration of lung adenocarcinoma cells by downregulating CNPY2. *Oncol Rep* 43: 646-654.

76. Wang, X., J. Wang, G. Huang, Y. Li, and S. Guo. 2021. miR-320a-3P alleviates the epithelial-mesenchymal transition of A549 cells by activation of STAT3/SMAD3 signaling in a pulmonary fibrosis model. *Mol Med Rep* 23.

77. Liu, X., J. Song, Y. Kang, Y. Wang, and A. Chen. 2022. CircPDSS1 promotes the proliferation, invasion, migration, and EMT of breast cancer cell via regulating miR-320c/CKAP5 axis. *Cancer Cell Int* 22: 238.

78. Shi, S., X. Hu, J. Xu, H. Liu, and L. Zou. 2018. MiR-320d suppresses the progression of breast cancer via lncRNA HNF1A-AS1 regulation and SOX4 inhibition. *RSC Adv* 8: 19196-19207.

79. Hou, B., H. Ishinaga, K. Midorikawa, S. Nakamura, Y. Hiraku, S. Oikawa, N. Ma, K. Takeuchi, and M. Murata. 2018. Let-7c inhibits migration and epithelial-mesenchymal transition in head and neck squamous cell carcinoma by targeting IGF1R and HMGA2. *Oncotarget* 9: 8927-8940.

80. Liang, L., Y. W. Zheng, and Y. L. Wang. 2020. miR-4429 Regulates the Proliferation, Migration, Invasion, and Epithelial-Mesenchymal Transition of Cervical Cancer by Targeting FOXM1. *Cancer Manag Res* 12: 5301-5312.

81. Pan, H., Y. Hong, B. Yu, L. Li, and X. Zhang. 2019. miR-4429 Inhibits Tumor Progression and Epithelial-Mesenchymal Transition Via Targeting CDK6 in Clear Cell Renal Cell Carcinoma. *Cancer Biother Radiopharm* 34: 334-341.

82. Dong, P., M. Karaayvaz, N. Jia, M. Kaneuchi, J. Hamada, H. Watari, S. Sudo, J. Ju, and N. Sakuragi. 2013. Mutant p53 gain-of-function induces epithelial-mesenchymal transition through modulation of the miR-130b-ZEB1 axis. *Oncogene* 32: 3286-3295.

83. Li, B. L., C. Lu, W. Lu, T. T. Yang, J. Qu, X. Hong, and X. P. Wan. 2013. miR-130b is an EMT-related microRNA that targets DICER1 for aggression in endometrial cancer. *Med Oncol* 30: 484.

84. Chang, R. M., J. F. Xu, F. Fang, H. Yang, and L. Y. Yang. 2016. MicroRNA-130b promotes proliferation and EMT-induced metastasis via PTEN/p-AKT/HIF-1alpha signaling. *Tumour Biol* 37: 10609-10619.

85. Arisan, E. D., O. Rencuzogullari, C. Cieza-Borrella, F. Miralles Arenas, M. Dwek, S. Lange, and P. Uysal-Onganer. 2021. MiR-21 Is Required for the Epithelial-Mesenchymal Transition in MDA-MB-231 Breast Cancer Cells. *Int J Mol Sci* 22.

86.   Ghosh, A., N. Ranjan, L. Jiang, A. H. Ansari, N. Degyatoreva, S. Ahluwalia, D. P. Arya,
and S. Maiti. 2022. Fine-tuning miR-21 expression and inhibition of EMT in breast
cancer cells using aromatic-neomycin derivatives. *Mol Ther Nucleic Acids* 27: 685-698.
87.   Zhao, N., H. Sun, B. Sun, D. Zhu, X. Zhao, Y. Wang, Q. Gu, X. Dong, F. Liu, Y. Zhang,
and X. Li. 2016. miR-27a-3p suppresses tumor metastasis and VM by down-regulating
VE-cadherin expression and inhibiting EMT: an essential role for Twist-1 in HCC. *Sci*
Rep 6: 23091.
88.   Pan, G., Y. Liu, L. Shang, F. Zhou, and S. Yang. 2021. EMT-associated microRNAs and
their roles in cancer stemness and drug resistance. *Cancer Commun (Lond)* 41: 199-217.
89.   Zhang, J., X. Hua, N. Qi, G. Han, J. Yu, Y. Yu, X. Wei, H. Li, X. Chen, C. Leng, Q. Liu,
Y. Lu, and Y. Li. 2020. MiR-27b suppresses epithelial-mesenchymal transition and
chemoresistance in lung cancer by targeting Snail1. *Life Sci* 254: 117238.
90.   Suzuki, H. I., A. Katsura, H. Mihira, M. Horie, A. Saito, and K. Miyazono. 2017.
Regulation of TGF-beta-mediated endothelial-mesenchymal transition by microRNA-27.
J Biochem 161: 417-420.
91.   Lu, Y., W. Huang, H. Chen, H. Wei, A. Luo, G. Xia, X. Deng, and G. Zhang. 2019.
MicroRNA-224, negatively regulated by c-jun, inhibits growth and epithelial-to-
mesenchymal transition phenotype via targeting ADAM17 in oral squamous cell
carcinoma. *J Cell Mol Med* 23: 4913-4920.
92.   Zang, C. S., H. T. Huang, J. Qiu, J. Sun, R. F. Ge, and L. W. Jiang. 2020. MiR-224-5p
targets EGR2 to promote the development of papillary thyroid carcinoma. *Eur Rev Med*
Pharmacol Sci 24: 4890-4900.
93.   Hsiao, S. Y., S. M. Weng, J. R. Hsiao, Y. Y. Wu, J. E. Wu, C. H. Tung, W. L. Shen, S. F.
Sun, W. T. Huang, C. Y. Lin, S. H. Chen, T. M. Hong, Y. L. Chen, and J. Y. Chang.
2023. MiR-455-5p suppresses PDZK1IP1 to promote the motility of oral squamous cell
carcinoma and accelerate clinical cancer invasion by regulating partial epithelial-to-
mesenchymal transition. *J Exp Clin Cancer Res* 42: 40.
94.   Meng, C., K. Liu, X. Cai, and Y. Chen. 2021. Mechanism of miR-455-3 in suppressing
epithelial-mesenchymal transition and angiogenesis of non-small cell lung cancer cells.
Cell Stress Chaperones 27: 107-117.
95.   Chen, Y., D. Sun, D. Shang, Z. Jiang, P. Miao, and J. Gao. 2022. miR-223-3p alleviates
TGF-beta-induced epithelial-mesenchymal transition and extracellular matrix deposition
by targeting SP3 in endometrial epithelial cells. *Open Med (Wars)* 17: 518-526.
96.   Ke, Y., W. Zhao, J. Xiong, and R. Cao. 2013. miR-149 Inhibits Non-Small-Cell Lung
Cancer Cells EMT by Targeting FOXM1. *Biochem Res Int* 2013: 506731.
97.   Wang, J., and L. Liu. 2021. MiR-149-3p promotes the cisplatin resistance and EMT in
ovarian cancer through downregulating TIMP2 and CDKN1A. *J Ovarian Res* 14: 165.
98.   Sun, Y., Y. Yi, S. Gan, R. Ye, C. Huang, M. Li, J. Huang, and Y. Guo. 2021. miR-574-5p
mediates epithelial-mesenchymal transition in small cell lung cancer by targeting
vimentin via a competitive endogenous RNA network. *Oncol Lett* 21: 459.
99.   Zhang, K. J., Y. Hu, N. Luo, X. Li, F. Y. Chen, J. Q. Yuan, and L. Guo. 2020.
miR-574-5p attenuates proliferation, migration and EMT in triple-negative breast cancer
cells by targeting BCL11A and SOX2 to inhibit the SKIL/TAZ/CTGF axis. *Int J Oncol*
56: 1240-1251.
100.   Wang, C., W. Zhang, L. Zhang, X. Chen, F. Liu, J. Zhang, S. Guan, Y. Sun, P. Chen, D.
Wang, E. Un Nesa, Y. Cheng, and G. M. Yousef. 2016. miR-146a-5p mediates epithelial-

mesenchymal transition of oesophageal squamous cell carcinoma via targeting Notch2. *Br J Cancer* 115: 1548-1554.

101. Wang, F., L. J. Ye, F. J. Wang, H. F. Liu, and X. L. Wang. 2020. miR-146a promotes proliferation, invasion, and epithelial-to-mesenchymal transition in oral squamous carcinoma cells. *Environ Toxicol* 35: 1050-1057.

102. Liang, Y., S. Li, and L. Tang. 2021. MicroRNA 320, an Anti-Oncogene Target miRNA for Cancer Therapy. *Biomedicines* 9.

103. Liu, R. Y., Y. Zeng, Z. Lei, L. Wang, H. Yang, Z. Liu, J. Zhao, and H. T. Zhang. 2014. JAK/STAT3 signaling is required for TGF-beta-induced epithelial-mesenchymal transition in lung cancer cells. *Int J Oncol* 44: 1643-1651.

104. Gregory, P. A., C. P. Bracken, E. Smith, A. G. Bert, J. A. Wright, S. Roslan, M. Morris, L. Wyatt, G. Farshid, Y. Y. Lim, G. J. Lindeman, M. F. Shannon, P. A. Drew, Y. Khew-Goodall, and G. J. Goodall. 2011. An autocrine TGF-beta/ZEB/miR-200 signaling network regulates establishment and maintenance of epithelial-mesenchymal transition. *Mol Biol Cell* 22: 1686-1698.

105. Bettelli, E., Y. Carrier, W. Gao, T. Korn, T. B. Strom, M. Oukka, H. L. Weiner, and V. K. Kuchroo. 2006. Reciprocal developmental pathways for the generation of pathogenic effector TH17 and regulatory T cells. *Nature* 441: 235-238.

106. Guan, T., C. X. Dominguez, R. A. Amezquita, B. J. Laidlaw, J. Cheng, J. Henao-Mejia, A. Williams, R. A. Flavell, J. Lu, and S. M. Kaech. 2018. ZEB1, ZEB2, and the miR-200 family form a counterregulatory network to regulate CD8(+) T cell fates. *J Exp Med* 215: 1153-1168.

107. Wang, X. Y., X. Y. Chen, J. Li, H. Y. Zhang, J. Liu, and L. D. Sun. 2017. MiR-200a expression in CD4+ T cells correlates with the expression of Th17/Treg cells and relevant cytokines in psoriasis vulgaris: A case control study. *Biomed Pharmacother* 93: 1158-1164.

108. Naghavian, R., K. Ghaedi, A. Kiani-Esfahani, M. Ganjalikhani-Hakemi, M. Etemadifar, and M. H. Nasr-Esfahani. 2015. miR-141 and miR-200a, Revelation of New Possible Players in Modulation of Th17/Treg Differentiation and Pathogenesis of Multiple Sclerosis. *PLoS One* 10: e0124555.

109. Bahmani, L., M. Baghi, M. Peymani, A. Javeri, and K. Ghaedi. 2021. MiR-141-3p and miR-200a-3p are involved in Th17 cell differentiation by negatively regulating RARB expression. *Hum Cell* 34: 1375-1387.

110. Yu, X., X. Fan, X. Zhang, P. Wei, H. Zhou, D. Liu, and B. Chen. 2023. miR-429 inhibits the formation of an immunosuppressive microenvironment to counteract hepatocellular carcinoma immune escape by targeting PD-L1. *Funct Integr Genomics* 23: 312.

111. Jebbawi, F., H. Fayyad-Kazan, M. Merimi, P. Lewalle, J. C. Verougstraete, O. Leo, P. Romero, A. Burny, B. Badran, P. Martiat, and R. Rouas. 2014. A microRNA profile of human CD8(+) regulatory T cells and characterization of the effects of microRNAs on Treg cell-associated genes. *J Transl Med* 12: 218.

112. Cruz, L. O., S. S. Hashemifar, C. J. Wu, S. Cho, D. T. Nguyen, L. L. Lin, A. A. Khan, and L. F. Lu. 2017. Excessive expression of miR-27 impairs Treg-mediated immunological tolerance. *J Clin Invest* 127: 530-542.

113. Zhou, Y., Y. Li, J. Lu, X. Hong, and L. Xu. 2019. MicroRNA-30a controls the instability of inducible CD4+ Tregs through SOCS1. *Mol Med Rep* 20: 4303-4314.

114. Li, J. Q., J. M. Tian, X. R. Fan, Z. Y. Wang, J. Ling, X. F. Wu, F. Y. Yang, and Y. L.
Xia. 2020. miR-106b-5p induces immune imbalance of Treg/Th17 in immune
thrombocytopenic purpura through NR4A3/Foxp3 pathway. *Cell Cycle* 19: 1265-1274.
115. Jiao, P., X. P. Wang, Z. M. Luoreng, J. Yang, L. Jia, Y. Ma, and D. W. Wei. 2021. miR-
223: An Effective Regulator of Immune Cell Differentiation and Inflammation. *Int J Biol*
Sci 17: 2308-2322.
116. Chang, C. C., Q. Y. Zhang, Z. Liu, R. A. Clynes, N. Suciu-Foca, and G. Vlad. 2012.
Downregulation of inflammatory microRNAs by Ig-like transcript 3 is essential for the
differentiation of human CD8(+) T suppressor cells. *J Immunol* 188: 3042-3052.
117. Lu, L. F., M. P. Boldin, A. Chaudhry, L. L. Lin, K. D. Taganov, T. Hanada, A.
Yoshimura, D. Baltimore, and A. Y. Rudensky. 2010. Function of miR-146a in
controlling Treg cell-mediated regulation of Th1 responses. *Cell* 142: 914-929.
118. Zhou, Q., S. Haupt, J. T. Kreuzer, A. Hammitzsch, F. Proft, C. Neumann, J. Leipe, M.
Witt, H. Schulze-Koops, and A. Skapenko. 2015. Decreased expression of miR-146a and
miR-155 contributes to an abnormal Treg phenotype in patients with rheumatoid arthritis.
Ann Rheum Dis 74: 1265-1274.
119. Li, C., P. J. Ebert, and Q. J. Li. 2013. T cell receptor (TCR) and transforming growth
factor beta (TGF-beta) signaling converge on DNA (cytosine-5)-methyltransferase to
control forkhead box protein 3 (foxp3) locus methylation and inducible regulatory T cell
differentiation. *J Biol Chem* 288: 19127-19139.
120. Yang, H. Y., J. Barbi, C. Y. Wu, Y. Zheng, P. D. Vignali, X. Wu, J. H. Tao, B. V. Park,
S. Bandara, L. Novack, X. Ni, X. Yang, K. Y. Chang, R. C. Wu, J. Zhang, C. W. Yang,
D. M. Pardoll, H. Li, and F. Pan. 2016. MicroRNA-17 Modulates Regulatory T Cell
Function by Targeting Co-regulators of the Foxp3 Transcription Factor. *Immunity* 45: 83-
93.
121. de Kouchkovsky, D., J. H. Esensten, W. L. Rosenthal, M. M. Morar, J. A. Bluestone, and
L. T. Jeker. 2013. microRNA-17-92 regulates IL-10 production by regulatory T cells and
control of experimental autoimmune encephalomyelitis. *J Immunol* 191: 1594-1605.
122. Skinner, J. P., A. A. Keown, and M. M. Chong. 2014. The miR-17 approximately 92a
cluster of microRNAs is required for the fitness of Foxp3+ regulatory T cells. *PLoS One*
9: e88997.
123. Dong, L., X. Wang, J. Tan, H. Li, W. Qian, J. Chen, Q. Chen, J. Wang, W. Xu, C. Tao,
and S. Wang. 2014. Decreased expression of microRNA-21 correlates with the imbalance
of Th17 and Treg cells in patients with rheumatoid arthritis. *J Cell Mol Med* 18: 2213-
2224.
124. Li, D., X. Du, M. Zhu, S. Yang, and W. Zhao. 2020. MiR-374b-5p Regulates T Cell
Differentiation and Is Associated with rEg.P29 Immunity. *Biomed Res Int* 2020:
8024763.
125. Sun, R., P. P. Zhang, X. Q. Weng, X. D. Gao, C. X. Huang, L. Wang, X. X. Hu, P. P. Xu,
L. Cheng, L. Jiang, D. Fu, B. Qu, Y. Zhao, Y. Feng, H. J. Dou, Z. Zheng, and W. L.
Zhao. 2022. Therapeutic targeting miR130b counteracts diffuse large B-cell lymphoma
progression via OX40/OX40L-mediated interaction with Th17 cells. *Signal Transduct*
Target Ther 7: 80.
126. Liu, S. Q., S. Jiang, C. Li, B. Zhang, and Q. J. Li. 2014. miR-17-92 cluster targets
phosphatase and tensin homology and Ikaros Family Zinc Finger 4 to promote TH17-
mediated inflammation. *J Biol Chem* 289: 12446-12456.

127. Honardoost, M. A., R. Naghavian, F. Ahmadinejad, A. Hosseini, and K. Ghaedi. 2015.
Integrative computational mRNA-miRNA interaction analyses of the autoimmune-
deregulated miRNAs and well-known Th17 differentiation regulators: An attempt to
discover new potential miRNAs involved in Th17 differentiation. *Gene* 572: 153-162.
128. Wei, Y., S. Chen, D. Sun, X. Li, R. Wei, X. Li, and H. Nian. 2019. miR-223-3p promotes
autoreactive T(h)17 cell responses in experimental autoimmune uveitis (EAU) by
inhibiting transcription factor FOXO3 expression. *FASEB J* 33: 13951-13965.
129. Angelou, C. C., A. C. Wells, J. Vijayaraghavan, C. E. Dougan, R. Lawlor, E. Iverson, V.
Lazarevic, M. Y. Kimura, S. R. Peyton, L. M. Minter, B. A. Osborne, E. L.
Pobezinskaya, and L. A. Pobezinsky. 2019. Differentiation of Pathogenic Th17 Cells Is
Negatively Regulated by Let-7 MicroRNAs in a Mouse Model of Multiple Sclerosis.
Front Immunol 10: 3125.
130. Qu, X., J. Zhou, T. Wang, J. Han, L. Ma, H. Yu, D. Geng, H. Fan, Q. Zhang, F. Hua, and
R. Yao. 2016. MiR-30a inhibits Th17 differentiation and demyelination of EAE mice by
targeting the IL-21R. *Brain Behav Immun* 57: 193-199.
131. Zhu, E., X. Wang, B. Zheng, Q. Wang, J. Hao, S. Chen, Q. Zhao, L. Zhao, Z. Wu, and Z.
Yin. 2014. miR-20b suppresses Th17 differentiation and the pathogenesis of
experimental autoimmune encephalomyelitis by targeting RORgammat and STAT3. *J*
Immunol 192: 5599-5609.
132. Chen, J., X. Shi, Y. Deng, J. Dang, Y. Liu, J. Zhao, R. Liang, D. Zeng, W. Wu, Y. Xiong,
J. Yuan, Y. Chen, J. Wang, W. Lin, X. Chen, W. Huang, N. Olsen, Y. Pan, Q. Fu, and S.
G. Zheng. 2024. miRNA-148a-containing GMSC-derived EVs modulate Treg/Th17
balance via IKKB/NF-kappaB pathway and treat a rheumatoid arthritis model. *JCI*
Insight 9.
133. Pacholewska, A., M. F. Kraft, V. Gerber, and V. Jagannathan. 2017. Differential
Expression of Serum MicroRNAs Supports CD4(+) T Cell Differentiation into Th2/Th17
Cells in Severe Equine Asthma. *Genes (Basel)* 8.
134. Cao, Y., Z. Wang, Y. Yan, L. Ji, J. He, B. Xuan, C. Shen, Y. Ma, S. Jiang, D. Ma, T.
Tong, X. Zhang, Z. Gao, X. Zhu, J. Y. Fang, H. Chen, and J. Hong. 2021.
Enterotoxigenic Bacteroidesfragilis Promotes Intestinal Inflammation and Malignancy by
Inhibiting Exosome-Packaged miR-149-3p. *Gastroenterology* 161: 1552-1566 e1512.
135. Ahmadian-Elmi, M., A. Bidmeshki Pour, R. Naghavian, K. Ghaedi, S. Tanhaei, T. Izadi,
and M. H. Nasr-Esfahani. 2016. miR-27a and miR-214 exert opposite regulatory roles in
Th17 differentiation via mediating different signaling pathways in peripheral blood CD4+
T lymphocytes of patients with relapsing-remitting multiple sclerosis. *Immunogenetics*
68: 43-54.
136. Cho, S., C. J. Wu, T. Yasuda, L. O. Cruz, A. A. Khan, L. L. Lin, D. T. Nguyen, M.
Miller, H. M. Lee, M. L. Kuo, D. H. Broide, K. Rajewsky, A. Y. Rudensky, and L. F. Lu.
2016. miR-23 approximately 27 approximately 24 clusters control effector T cell
differentiation and function. *J Exp Med* 213: 235-249.
137. Akimniyazova, A., A. Pyrkova, V. Uversky, and A. Ivashchenko. 2021. Predicting
Associations of miRNAs and Candidate Gastric Cancer Genes for Nanomedicine.
Nanomaterials (Basel) 11.
138. Hironaka-Mitsuhashi, A., K. Otsuka, L. Gailhouste, A. Sanchez Calle, M. Kumazaki, Y.
Yamamoto, Y. Fujiwara, and T. Ochiya. 2020. MiR-1285-5p/TMEM194A axis affects
cell proliferation in breast cancer. *Cancer Sci* 111: 395-405.

139.   Zhang, J., X. Li, J. Yang, and Y. Zhang. 2022. MiR-1254 suppresses the proliferation and
930   invasion of cervical cancer cells by modulating CD36. *J Transl Med* 19: 531.

931   140.   Zhao, M. C., M. M. Zhang, T. Li, Z. H. Tao, Y. Q. Du, L. P. Wang, J. Zhang, B. Y.
932   Wang, and X. C. Hu. 2020. MiR-566 protects the malignant progression of breast cancer
933   by negatively regulating WNT6. *Eur Rev Med Pharmacol Sci* 24: 6185-6194.

934   141.   Zhou, J., M. Zhang, Y. Huang, L. Feng, H. Chen, Y. Hu, H. Chen, K. Zhang, L. Zheng,
935   and S. Zheng. 2015. MicroRNA-320b promotes colorectal cancer proliferation and
936   invasion by competing with its homologous microRNA-320a. *Cancer Lett* 356: 669-675.

937   142.   Pastor-Fernandez, G., I. R. Mariblanca, and M. N. Navarro. 2020. Decoding IL-23
938   Signaling Cascade for New Therapeutic Opportunities. *Cells* 9.

939   143.   Yeruva, L., D. L. Pouncey, M. R. Eledge, S. Bhattacharya, C. Luo, E. W. Weatherford,
940   D. M. Ojcius, and R. G. Rank. 2017. MicroRNAs Modulate Pathogenesis Resulting from
941   Chlamydial Infection in Mice. *Infect Immun* 85.

942   144.   Yeruva, L., A. K. Bowlin, N. Spencer, A. T. Maurelli, and R. G. Rank. 2015. Chlamydial
943   variants differ in ability to ascend the genital tract in the guinea pig model of chlamydial
944   genital infection. *Infect Immun* 83: 3176-3183.

945   145.   Rajeeve, K., S. Das, B. K. Prusty, and T. Rudel. 2018. Chlamydia trachomatis paralyses
946   neutrophils to evade the host innate immune response. *Nat Microbiol* 3: 824-835.

975   **Figures**

976
Figure 1. Workflow depicting sample processing, mRNA and miRNA profiling and

subsequent analytical methods. Endometrial biopsy samples were processed to profile mRNA

from 7 Endo (+) and 8 Endo (-) women, including 4 cervix (+) only and 4 uninfected, and

miRNA from 9 Endo (+) and 14 Endo (-) women, including 5 cervix(+) only and 9 uninfected.

Relationships among cohort participants based on their transcriptomes were revealed by

Principal Component Analysis (PCA). Differentially expressed (DE) mRNAs and miRNAs

between Endo+ and Endo- biopsies were determined and functional annotation of DE mRNA

and miRNA-mRNA pairs were provided by IPA. Additionally, endometrial biopsy generated

mRNA data were compared to previously published blood mRNA data generated from

participants in the T Cell Response Against Chlamydia (TRAC) cohort (32).

Figure 2. Principal component analysis (PCA) of endometrial transcriptome (A) mRNA

and (B) miRNA. Each dot represents one subject, with infection status indicated by color. Red

denotes endometrial infection (Endo+), light blue denotes absence of endometrial infection but

positive cervical infection (Endo-), and dark blue (Uninfected). The x-axis represents the first

principal component, PC1, which accounts for the largest variance of mRNA (A) or miRNA

(B) expression, and the y axis, PC2, explains the second largest variance.

Figure 3. **Histopathology of *Chlamydia trachomatis* endometritis.** (A, 20X; B, 40X)

Histological hematoxylin and eosin-stained endometrial tissue sections from a biopsy taken from

a woman positive for *C. trachomatis* in their endometrium. White arrows indicate plasma cells;

white arrowheads indicate neutrophils. (C, 20X; D, 40X) Endometrial biopsy section from a

woman who was negative for endometrial infection but tested positive for *C. trachomatis*

infection at their cervix.

1002

Figure 4. The significance and fold change of DE mRNAs (FDR<0.05) in Endo+ versus

Endo-/Uninfected biopsies depicted by volcano plot. Each dot represents one mRNA gene,

with red indicating upregulation and blue indicating downregulation in the Endo+ group

compared to the Endo-/Uninfected group. Horizontal dashed line indicates FDR=0.05 (-

$\log_{10}$P=2.2). Left vertical dashed line indicates log2 fold change=-2 (fold change= 0.25) and

right vertical dashed line indicates log2 fold change=2 (fold change=4).

1009

Figure 5. Ingenuity canonical pathways enriched by DE mRNAs and miRNAs in Endo+ compared to Endo-/Uninfected biopsies. (A) Top 20 signaling pathways enriched by significantly upregulated mRNAs in Endo+ biopsies. (B) All 18 pathways enriched by significantly downregulated mRNAs in Endo+ biopsies. (C) Top 20 signaling pathways enriched by downregulated miRNA-upregulated mRNA pairs in Endo+ biopsies. (D) Top 20 signaling pathways enriched by significantly upregulated miRNA-downregulated mRNA pairs in Endo+ biopsies. The percentage is the number of the DE genes present in each pathway, divided by the total number of genes in that pathway according to the IPA database, as listed on the right-hand side of each figure panel.

Figure 6. The significance and fold change of DE miRNAs (FDR<0.05) in Endo+ versus Endo-/Uninfected biopsies depicted by (A) volcano plot. Each dot represents one miRNA gene with colors indicating directions of dysregulated expression in the Endo+ group compared to the Endo-/Uninfected group. Horizontal dashed line indicates FDR=0.05 (-log₁₀P=2.2). Left vertical dashed line indicates log2 fold change=-1 (fold changes= 0.5) and right vertical dashed line indicates log2 fold change=1 (fold change=2). All significant DE miRNAs (FDR<0.05) with

1026 fold changes>2 or <0.5 are highlighted. **DE miRNAs linked to (B) EMT, (C) Tregs, and (D)**

1027 **Th17 differentiation are depicted by pie charts.** Red, blue and white colors indicate overall

1028 enhancing, suppressing, and enhancing/suppressing effects of miRNAs, respectively, in Endo+

1029 women compared to Endo-/Uninfected women.

1030

 Figure 7. A within-sample group hierarchical clustering heatmap of DE miRNAs revealed

distinct expression patterns across groups. Heatmap columns indicate individual patients with

red: Endo+ with pelvic pain; yellow: Endo+ without symptoms; green: Endo- without symptoms;

blue: uninfected without symptoms; pink: pelvic pain without sexually transmitted infection

(STI). Heatmap rows indicate genes. Blue: low expression miRNAs, white: intermediately

expressed miRNAs, and red: highly expressed miRNAs.

Table 1. DE miRNAs in Endo+ compared to Endo- associated with EMT.

DE miRNAs	Fold Change	P value	FDR	Expression changes in Endo+ compared to Endo-	Function of miRNAs	Overall effects	Citations
miR-566	0.32	0.002	0.009	Decrease (-)	↓EMT		(41)
miR-200a-3p	0.32	<0.001	<0.001	Decrease (-)	↓EMT		(42)
miR-200c-3p	0.59	0.013	0.039	Decrease (-)	↓EMT	**Promote EMT**	(42)
miR-141-3p	0.22	<0.001	<0.001	Decrease (-)	↓EMT		(43)
miR-429	0.29	<0.001	<0.001	Decrease (-)	↓EMT		

miR-376a-3p	0.51	0.016	0.048	Decrease (-)	↓EMT		(44)
miR-376c-3p	0.45	0.002	0.009	Decrease (-)	↓EMT		(45)
miR-374b-5p	0.39	0.002	0.008	Decrease (-)	↓EMT		(46)
miR-30b-5p	0.4	0.002	0.009	Decrease (-)	↓EMT		(47)
miR-30c-5p	0.68	0.015	0.045	Decrease (-)	↓EMT		(47)
miR-30e-5p	0.47	0.002	0.008	Decrease (-)	↓EMT		(47)
miR-186-5p	0.62	0.002	0.009	Decrease (-)	↓EMT		(48)
miR-130a-3p	0.63	0.001	0.006	Decrease (-)	↓EMT		(49)
miR-335-5p	0.44	0.001	0.007	Decrease (-)	↓EMT		(50)
miR-17-5p	0.6	<0.001	0.002	Decrease (-)	↓EMT		(51)
miR-18a-5p	0.47	0.001	0.004	Decrease (-)	↓EMT		(51)
miR-19a-3p	0.37	0.002	0.009	Decrease (-)	↓EMT		(51)
miR-19b-3p	0.4	<0.001	0.003	Decrease (-)	↓EMT		(51)
miR-20a-5p	0.5	0.005	0.019	Decrease (-)	↓EMT		(51)
miR-20b-5p	0.51	<0.001	<0.001	Decrease (-)	↓EMT		(52, 53)
miR-34c-5p	0.36	0.011	0.036	Decrease (-)	↓EMT		(54)
miR-101-3p	0.39	0.01	0.035	Decrease (-)	↓EMT		(55)
miR-194-5p	0.46	<0.001	0.002	Decrease (-)	↓EMT		(56)
miR-193a-3p	0.57	0.004	0.017	Decrease (-)	↓EMT		(57)
miR-148a-3p	0.53	0.001	0.004	Decrease (-)	↓EMT		(58, 59)
mir-148b-3p	0.73	0.002	0.011	Decrease (-)	↓EMT		(60)
miR-126-5p	0.32	<0.001	0.004	Decrease (-)	↓EMT		(61, 62)
miR-142-5p	0.26	0.009	0.032	Decrease (-)	↓EMT		(63)
miR-1285-5p	0.15	<0.001	0.003	Decrease (-)	↓EMT		(64)
mir-1254	0.24	<0.001	0.002	Decrease (-)	↓EMT		(65)
miR-6803-5p	2.26	<0.001	0.003	Increase (+)	↑EMT		(66)
miR-664a-3p	1.82	0.001	0.004	Increase (+)	↑EMT		(67)

miR-663b	1.98	0.001	0.005	Increase (+)	↑EMT		(68)
miR-197-3p	1.67	0.012	0.039	Increase (+)	↑EMT		(69)
miR-106b-5p	0.49	<0.001	0.001	Decrease (-)	↑EMT	Suppress EMT	(70-72)
miR-198	3.54	<0.001	0.001	Increase (+)	↓EMT		(73)
mir-30a-3p	1.67	0.008	0.027	Increase (+)	↓EMT		(74)
miR-320a	1.9	<0.001	<0.001	Increase (+)	↓EMT		(75)
miR-320c	1.88	<0.001	<0.001	Increase (+)	↓EMT		(76)
miR-320d	1.85	<0.001	0.001	Increase (+)	↓EMT		(77)
let-7c-5p	2.08	<0.001	0.002	Increase (+)	↓EMT		(78)
miR-4429	1.9	<0.001	<0.001	Increase (+)	↓EMT		(79, 80)
miR-130b-3p	0.63	0.007	0.026	Decrease (-)	↓↑EMT		(81-83)
miR-21-5p	0.6	0.024	0.05	Decrease (-)	↓↑EMT		(84, 85)
miR-27a-3p	0.55	<0.001	0.001	Decrease (-)	↓↑EMT		(86, 87)
miR-27b-3p	0.66	0.004	0.017	Decrease (-)	↓↑EMT		(88, 89)
miR-224-5p	0.37	0.001	0.004	Decrease (-)	↓↑EMT		(90, 91)
miR-455-5p	0.56	0.011	0.035	Decrease (-)	↓↑EMT	Suppress/Promote EMT	(92, 93)
miR-223-3p	0.19	0.008	0.027	Decrease (-)	↓↑EMT		(94)
miR-149-3p	2.74	0.006	0.024	Increase (+)	↓↑EMT		(95, 96)
miR-574-5p	2.28	0.005	0.021	Increase (+)	↓↑EMT		(97, 98)
miR-146a-5p	2.09	0.017	0.048	Increase (+)	↓↑EMT		(99, 100)
miR-320b	1.91	<0.001	<0.001	Increase (+)	↓↑EMT		(75, 101)

1047

1048   Table 3. DE miRNAs in Endo+ compared to Endo-, associated with Th17 cell differentiation.

DE miRNAs	Fold change	P value	FDR	Expression changes in Endo+ compared to Endo-	Function of miRNAs	Overall effects	Citations
miR-200a-3p	0.32	8.02E-07	2.25E-05	Reduced (-)	↑ Th17		(105, 106)

miR-141-3p	0.22	5.87E-08	3.71E-06	Reduced (-)	↑ Th17	Suppress Th17 differentiation	(108)
miR-200c-3p	0.59	1.26E-02	3.93E-02	Reduced (-)	↑ Th17		(105)
miR-429	0.29	8.84E-06	2.03E-04	Reduced (-)	↑ Th17		
miR-374b-5p	0.39	1.64E-03	7.98E-03	Reduced (-)	↑ Th17		(123)
miR-106b-5p	0.49	1.17E-04	1.34E-03	Reduced (-)	↑ Th17		(113)
miR-130b-3p	0.63	7.17E-03	2.59E-02	Reduced (-)	↑ Th17		(124)
miR-17	0.60	1.65E-04	1.59E-03	Reduced (-)	↑ Th17		(125)
miR-18a	0.47	5.44E-04	3.53E-03	Reduced (-)	↑ Th17		(125)
miR-19a	0.37	1.88E-03	8.76E-03	Reduced (-)	↑ Th17		(125)
miR-19b	0.4	4.12E-04	2.89E-03	Reduced (-)	↑ Th17		(125)
miR-20a	0.50	4.86E-03	1.92E-02	Reduced (-)	↑ Th17		(125)
mir-30c-5p	0.68	1.48E-02	4.45E-02	Reduced (-)	↑ Th17		(126)
mir-223-3p	0.19	7.53E-03	2.68E-02	Reduced (-)	↑ Th17		(114, 127)
Let-7b	2.3	5.66E-05	7.54E-04	Increased (+)	↓ Th17		(128)
Let-7c	2.1	1.52E-04	1.54E-03	Increased (+)	↓ Th17		(128)
miR-30a-3p	1.7	8.51E-03	2.95E-02	Reduced (-)	↓ Th17	Promote Th17 differentiation	(129)
miR-20a	0.50	4.86E-03	1.92E-02	Reduced (-)	↓ Th17		(125)
miR-20b	0.51	7.39E-08	3.74E-06	Reduced (-)	↓ Th17		(130)
miR-21-5p	0.44	6.90E-03	2.53E-02	Reduced (-)	↓ Th17		(122, 126)
miR-148a	0.53	5.15E-04	3.52E-03	Reduced (-)	↓ Th17		(131) (125)
miR-148b	0.73	2.50E-03	1.07E-02	Reduced (-)	↓ Th17		(132)
miR-149-3p	2.7	6.32E-03	2.35E-02	Increased (+)	↑ Th17		(133)
miR-27 a	0.55	9.72E-05	1.17E-03	Reduced (-)	↓↑ Th17	Suppress/promote Th17 differentiation	(134, 135)
miR-27 b	0.66	4.10E-03	1.67E-02	Reduced (-)	↓↑ Th17		(134, 135)

1049

Table 4. Comparison of pathways enriched by significantly differentially expressed (DE) mRNAs in Endo+ versus Endo- women, as determined from either endometrial biopsy or blood samples#.

IPA Pathway	Endometrium	Blood

	Regulation direction in Endo+	Adjusted P*	Regulation direction in Blood	Adjusted P**
Up-regulated in Endometrium and Blood				
IL-10 Signaling	↑	3.83E-15	↑	1.29E-06
IL-6 Signaling	↑	3.83E-13	↑	1.32E-05
Granulocyte Adhesion and Diapedesis	↑	3.04E-11	↑	6.92E-03
TREM1 Signaling	↑	6.07E-11	↑	7.78E-09
Production of Nitric Oxide and Reactive Oxygen Species in Macrophages	↑	6.07E-11	↑	6.15E-04
PI3K/AKT Signaling	↑	6.07E-11	↑	1.48E-03
Phagosome Formation	↑	1.92E-09	↑	1.66E-04
Toll-like Receptor Signaling	↑	2.42E-09	↑	3.47E-05
IL-8 Signaling	↑	9.84E-08	↑	5.13E-07
IL-3 Signaling	↑	1.71E-06	↑	3.47E-02
IL-17 Signaling	↑	1.30E-05	↑	4.16E-02
Leukocyte Extravasation Signaling	↑	4.20E-05	↑	1.86E-04
IL-1 Signaling	↑	5.29E-03	↑	1.38E-02
Dendritic Cell Maturation	↑	1.63E-02	↑	2.57E-05
Integrin Signaling	↑	1.81E-02	↑	9.34E-04
T Helper Cell Differentiation	↑	2.05E-02	↑	1.02E-03
Interferon Signaling	↑	3.88E-02	↑	1.95E-12
Up-regulated Only in Endometrium				
Th1 and Th2 Activation Pathway	↑	3.04E-14		##NS
Natural Killer Cell Signaling	↑	6.07E-12		NS
Regulation Of The Epithelial Mesenchymal Transition	↑	1.21E-10		NS
IL-23 Signaling Pathway	↑	7.47E-08		NS
IL-7 Signaling Pathway	↑	4.10E-07		NS
STAT3 Pathway	↑	4.50E-07		NS
TGF-β Signaling	↑	1.67E-06		NS
IL-13 Signaling Pathway	↑	3.83E-03		NS
T Cell Exhaustion Signaling Pathway	↑	2.47E-02		NS
Th17 Activation Pathway	↑	3.74E-02		NS
Up-regulated in Endometrium and Down regulated in Blood				
T Cell Receptor Signaling	↑	5.29E-03	↓	8.52E-03
ICOS-ICOSL Signaling in T Helper Cells	↑	2.36E-02	↓	7.78E-04
mTOR Signaling	↑	NS	↓	4.90E-12
Regulation of eIF4 and p70S6K Signaling	↑	NS	↓	3.89E-11

Data from reference (32).
* Bonferroni adjusted P values in endometrium: P*482 for endometrium (482 total pathways).
**Bonferroni adjusted P values in blood: P*389 for blood (389 total pathways).
##NS: Not Significant.

Light gray indicates pathways that were enriched by significantly upregulated genes only in endometrial
biopsy specimens.
Dark gray indicates pathways that were enriched by significantly down-regulated genes only in blood
specimens.
1058