Clonal memory of cell division in humans diverges between healthy haematopoiesis and acute myeloid leukaemia

Donada A[1]**, Hermange G[2,3,4], Tocci T[1], Midoun A[1], Prevedello G[5,6], Hadj Abed L[1,7], Dupré D[7], Sun W[1], Milo I[1], Tenreira Bento S[1], Pospori C[8,9], Innes A[10], Willekens C[11], Vargaftig J[12], Michonneau D[13,14], Lo Celso C[8,9,10], Servant N[7,15], Duffy KR[16], Isambert H[1], Cournède PH[2], Laplane L[17,18] and Perié L[1,6,19,20,21]*

1. Institut Curie, PSL Research University, Sorbonne University, CNRS UMR168, Physics of Cells and Cancer lab, Paris, France

2. Laboratory of Mathematics and Informatics (MICS), Université Paris-Saclay, CentraleSupélec, Gif-sur-Yvette, France

3. IHU-National PRecISion Medicine Center in Oncology, Villejuif, France

4. Université Paris-Saclay, Gustave Roussy, CentraleSupélec, INSERM, Cancer Data Science unit, Gif-sur-Yvette, France

5. Sony Computer Science Laboratories, Paris, France

6. Enrico Fermi's Research Center (CREF), via Panisperna 89A, 00184, Rome, Italy

7. Single Cell Initiative, Institut Curie, Paris, France

8. The Francis Crick Institute, London, UK

9. Department of Life Sciences, Imperial College, London UK

10. Centre for Haematology, Imperial College, London UK

11. Department of Haematology, Gustave Roussy Cancer Campus, Villejuif, France

12. Department of Haematology, Institut Curie, Saint-Cloud, France

13. Haematology and Transplantation Unit, Saint Louis Hospital, AP-HP, Paris, France

14. INSERM U1342, Saint Louis Research Institute, IHU Leukemia Institute Paris Saint Louis, SIRIC InSitu, Université Paris Cité, Paris, France

15. Institut Curie, INSERM U900, Mines Paris Tech, PSL Research University, Paris, France

16. Northeastern University, Boston, MA, USA

17. Hematopoietic Tissue Aging, INSERM UMR 1287, Gustave Roussy Cancer Campus, Villejuif, France

18. Institut d'Histoire et Philosophie des Sciences et des Techniques, CNRS UMR 8590, University Paris I Pantheon-Sorbonne, Paris, France

19. Biogipuzkoa Health Research Institute, San Sebastián, Spain.

20. IKERBASQUE: The Basque Foundation for Science, Bilbao, Spain

21. Lead Contact

* Correspondence: leila.perie@bio-gipuzkoa.eus

** Correspondence: alessandro.donada@curie.fr

Summary

Clonal memory, a cellular property inherited across at least two divisions, has emerged as a
key driver of cell heterogeneity. To uncover its roles in human haematopoiesis, we developed
high-resolution *ex vivo* tools that track both division and fate commitment of individual
primary human haematopoietic stem and progenitor cells (HSPCs). We show that human
HSPCs display a clonal memory of division, as cells descending from the same ancestor cell
divide synchronously over multiple generations. In parallel, HSPCs inherit a clonal memory of
fate commitment, independently of lineage identity. Both forms of clonal memory persist
over at least two divisions, across different HSPC commitment stages and cell culture
conditions. In contrast, malignant haematopoiesis exhibits lower synchronicity, revealing a
disruption of clonal memory in leukemic cells. Epigenetic remodelling using a bromodomain
inhibitor partially restores the clonal memory in division in leukemic HSPCs, highlighting the
plasticity of this trait and its potential for therapeutic modulation. Our findings position clonal
memory as a key regulator of human haematopoietic stem cell behaviour. Demonstrating that
clonal memory can be modulated opens new avenues for tuning cell heterogeneity in healthy
and pathological tissues.

Introduction

Stem cells, and how they proliferate, play a key role in tissue development, homeostasis, and
regeneration. However, not all stem cells divide or die at the same rate[1]. These proliferation
differences impact the cellular composition of tissues and thereby tissue formation and
renewal over time. In cancer, changes in cell division directly affect the fitness of cancer cells,
leading to their clonal expansion, increased malignant phenotypes and poorer prognosis[2].
Currently, we still know little about what drives differences in stem cell proliferation in
healthy or diseased tissues.

Recently, clonal memory has emerged as a driver of stem cell heterogeneity[3–7]. Clonal
memory is defined as a cellular property inherited over at least two successive rounds of cell
division. Despite its importance in stem cell heterogeneity, the role of clonal memory in
proliferation heterogeneity is poorly understood. Additionally, fundamental questions about
clonal memory remain, including how long it persists over time (e.g. hours[8], days[9], months[10])
and its long-term functional impact. Although clonal memory is an intrinsic property, whether
and how extrinsic factors (e.g. microenvironment, soluble factors) could modulate it remains
unknown. Modulating clonal memory is crucial as it could promote the stability required for
homeostasis by transmitting cellular traits across multiple cell generations. However, this
stability could come at the expense of adaptability to environment changes. Thus,
understanding how clonal memory contributes to heterogeneity in stem cell division is critical
to determining its role in balancing stability and adaptability in both homeostasis and disease.

Haematopoiesis is a good model to address the role of clonal memory. It is a hierarchically
organized differentiating system that generates over 13 different cell types across tens of
divisions[11]. HSPCs are a heterogeneous population with diverse division and differentiation

abilities[12]. In mice, clonal memory of differentiation partly contributes to this heterogeneity.
Indeed, HSPCs show heterogeneous differentiation biases that are maintained and
transmitted throughout cell division and differentiation[13], as shown by single-cell lineage
tracing[3] and serial transplantations[10]. However, most clonal memory studies have focused on
differentiation, while its influence on HSPC proliferation[14], in particular in human cells,
remains unexplored. Moreover, as differentiation tends to be coupled with division[15,16] and
both properties are central to stem cell biology, the clonal memory of both processes needs
to be studied together. Critical for this aim is the ability to simultaneously track division and
differentiation within individual cell lineages, using lineage tracing tools such as the MultiGen
assay[14,17–19].

In pathological human haematopoiesis, evidence for clonal memory of division and fate in
HSPCs remains largely indirect. While stem cell bone marrow transplantation induces cell
proliferation to repopulate the recipient bone marrow, the observation that stem cells
accumulate mutations at similar rates in both recipient and donor support the hypothesis that
stem cell proliferation is an intrinsic properties[20–22]. In clonal haematopoiesis, known genetic
drivers account only for approximately 20% of all age-related clonal expansions, indicating
that while genetic mutations contribute to proliferation heterogeneity, other intrinsic and
extrinsic factors are at play[23]. In humanized mouse models, cellular memory was
demonstrated to impact the heterogeneity in clonal fitness for otherwise genetically identical
acute myeloid leukaemia (AML) stem cells, shaping their clonal evolution[3], and leading to
differences in treatment resistance[24,25]. However, whether clonal memory can be modulated
in leukaemia and leveraged for therapeutic benefit remains unexplored.

Here, we address whether clonal memory plays a role in human haematopoiesis, by studying
clonal memory in primary human HSPCs during healthy and malignant haematopoiesis,
focusing on division. We developed several tools to follow division, differentiation, and
lineage of individual human HSPC *ex vivo*. Using live-cell microscopy, we establish that human
HSPCs display a clonal memory of cell division over at least two division and three cell
generations, across different HSPCs commitment stages and several culture conditions.
Further, we demonstrate that there are two distinct inherited clonal memory properties, one
for division and one for fate commitment. Counterintuitively, we find that HSPCs in malignant
AML haematopoiesis show altered clonal memory in division, with a decreased level of
synchronicity within cell families. By manipulating clonal memory of division through
inhibition of epigenetic remodelling in AML cells, we observe increased clonal memory but no
changes in the total number of divisions, suggesting that clonal memory is regulated
independently of the number of divisions. Altogether, we demonstrate the existence of
distinct clonal memories that govern division and lineage commitment in healthy human
primary HSPCs. Furthermore, we show that while the clonal memory of division is altered in
AML, it remains amenable to modulation. This ability to modulate clonal memory opens new
avenues for tuning cell heterogeneity.

Results

Clonal memory of division in human HSPC families

To measure the heterogeneity in division in human HSPCs and their descendant cells, we analysed three types of HSPCs, via single-cell live-cell imaging. We isolated hundreds of immature progenitors, defined as CD90$^+$ (Lin$^-$CD34$^+$CD38$^-$CD90$^+$CD45RA$^-$), CD90$^-$ (Lin$^-$CD34$^+$CD38$^-$CD90$^-$CD45RA$^-$), and haematopoietic progenitor cells (HPCs; Lin$^-$CD34$^+$CD38$^+$) from umbilical cord blood and adult bone marrow from unrelated donors **(Extended Data Fig. 1a).** We then cultured single HSPCs for 96 hours in two serum-free media: one to promote cell division and differentiation towards the myeloid and erythroid lineages (Diff)[26], and one to maintain the immature CD34$^+$ phenotype (gene therapy medium, GT)[27]. Analysing 1973 cell families, each composed of the descendants of a single HSPC, we manually extracted the time necessary to complete the first three divisions from more than 5000 cells **(Fig. 1a)**. First, we observed differences of proliferation between cell types, with HPCs dividing faster and forming bigger colonies at 96 hours than CD90$^+$ or CD90$^-$, for multiple donors in both culture media **(Fig. 1b-c, Extended Data Fig. 2m)**. Past the first division **(Extended Data Fig. 2a-b)**, the time to divide shortened not only for CD90$^+$ and CD90$^-$ **(Extended Data Fig. 2d-e, 2g-h),** as previously described[28], but also for HPCs. Therefore, cellular quiescence alone was not responsible for this shortened division time, as HPCs are not quiescent[29,30] **(Extended Data Fig. 2j-k)**. Additionally, cells of the same type displayed heterogenous division times, as previously reported[31]. The heterogeneity was of similar amplitude between cell types (i.e., CD90$^+$s, CD90$^-$s, and HPCs) and culture media, although adult bone marrow HSPCs **(Extended Data Fig. 2c, 2f, 2i, 2l)** displayed more heterogeneity than foetal cord blood HSPCs.

To assess the existence of clonal memory of division, we next explored the heterogeneity in division times within cell families. To this end, we measured the difference in division timing between cells in the same generation: sister cells (cells sharing the same mother cell) and cousin cells (cells sharing the same grandmother cell). Cells within a family divided synchronously **(Fig. 1d),** as sister cell division times were highly correlated for all tested progenitors (CD90$^+$s, CD90$^-$s, and HPCs) and culture media, accordingly to the calculated Spearman ranked correlation coefficient **(Extended Data Fig. 3a)**. Furthermore, cousin cells also divided synchronously **(Fig. 1f)**, with the four division for a given set of cousin cells displaying a lower standard deviation compared to permuted families composed of unrelated cells. This synchronicity in division was present in all HSPC types, irrespective of their origin **(Fig. 1e, Extended Data Fig. 3c)** or culture media **(Extended Data Fig. 3b)**, suggesting a clonal memory of division across commitment stages. However, cell division time across generations correlated weakly **(Fig. 1g),** as in other cell systems. This was similar for each HSPC type, irrespective of culture media **(Extended Data Fig. 3d-e, 3g-h, 3j)** or cell origin **(Extended Data Fig. 3f, 3i, 3k)**. In summary, we observed a significant and conserved intragenerational synchrony in cell division timing, over different HSPC commitment stages, and across different culture conditions.

Figure 1. Division in synchronicity within cell family. *a) Overview of the live cell imaging experimental workflow: individual progenitors were index sorting via FACS in 96-well tissue plates and kept in culture for at least 96 hours. Division dynamics were manually annotated.* ***b)*** *Colony size, measured as number of light-refringent cells in the well at T = 96 hours, for foetal cord blood HSPCs expanded in Diff medium, in 7 independent experiments (n = 166 for CD90$^+$, n = 369 for CD90$^-$ and n = 157 for HPC).* ***c)*** *Colony size, measured as number of light-refringent cells in the well at T = 96 hours, for bone marrow HSPCs, in 8 independent experiments (n = 356 for CD90$^+$, n = 328 for CD90$^-$ and n = 222 for HPCs). Dunn post-hoc test with Hochberg correction, ****p < 1x10^{-4}.* ***d-e)*** *Correlation in division time (expressed in hours) between two sister cells in generation 1, for* ***d)*** *foetal HSPCs isolated from cord blood and expanded in Diff medium (7 independent experiments, n = 149 for CD90$^+$, n = 335 for CD90$^-$ and n = 144 for HPC), and* ***e)*** *adult bone marrow cells (8 independent experiments, n = 255 for CD90$^+$, n = 269 for CD90$^-$ and n = 205 for HPC). Spearman's rank correlation coefficient R.* ***f)*** *Mean SD of third division time (T3) of cells derived from the same ancestor cells (black line), and permuted families (red histograms), for HSPCs from foetal cord blood and expanded in Diff medium. Only families with four recorded T3 divisions were used. Seven independent experiments; n = 65 for CD90$^+$, n = 170 for CD90$^-$ and n = 99 for HPC), p values < 1x10^{-4}.* ***g)*** *Correlation in division time (expressed in hours) between the mean division time T2 and the mean division time T3, for HSPCs isolated from foetal cord blood and expanded in Diff medium. Spearman's rank correlation coefficient R.*

Clonal memory of division and lineage commitment in human HSPCs

To determine if human HSPCs display a clonal memory of lineage commitment, we performed the MultiGen assay[14,17,18] to track simultaneously the cell family, the number of cell divisions and the phenotype of each cell produced by individual HSPCs **(Fig. 2a)**. We analysed 1634 individual CD90$^+$- and CD90$^-$-derived families, from human cord blood, after 72 hours and 96 hours and using both Diff and GT media. Following the MultiGen assay, we seeded 4 individual progenitors per each well, all labelled with a different division tracer colour, to allow the correct identification of the cells sharing the same ancestry. We assigned the progeny phenotype using an antibody panel that identifies six different HSPC subtypes **(Extended Data Fig. 1b)**. At the population level, we observed that CD90$^-$ families divided and differentiated more often than CD90$^+$ families **(Fig. 2b-c, Extended Data Fig. 4a-b)**, as expected from their position in the haematopoietic hierarchy. Consistent with previous observations[14], cells that divided more were also more likely to be more differentiated than the ones that divided less **(Extended Data Fig. 4c-j)**. Moreover, we observed only a minority of cells retaining their initial phenotype (CD90$^+$ or CD90$^-$) after the first two divisions, confirming that most cells undergo differentiation instead of self-renewal. The vast majority of immature HSPCs (i.e., CD90$^+$s and CD90$^-$s) were associated with fewer divisions, while committed progenitors and precursors emerged with more divisions **(Fig. 2d-e)**. We observed substantial heterogeneity in lineage commitments and divisions between families of the same progenitor cell type **(Fig. 2c, Extended Data Fig. 4k-r)**, in line with the current state of the art[32]. Therefore, our experimental set-up recapitulates the heterogeneity of the lineage commitment and division found in human HSPCs.

Next, we explored the heterogeneity in lineage commitment within cell families. Cells from the same family consistently co-localized on the UMAP representation of the HSPC subsets detected via flow cytometry **(Fig. 2f)**. Most of the families displayed only one subset **(Fig. 2g)**, despite the detection of multiple cell types in our culture system and the multipotent nature[33] of CD90$^+$ and CD90$^-$. Permutation testing revealed significant intrafamilial similarities in lineage commitment, more than expected if each cell was committing independently of the sister and cousin cells. This observation was independent of the ancestor cell type, the timing of the analysis, and the culture condition **(Extended Data Fig. 5a-c)**. We concluded that a clonal memory of lineage commitment in human HSPCs is present over different HSPC commitment stages and across culture conditions.

We also observed clonal memory of division with the MultiGen assay, consistent with our results using live-imaging microscopy **(Fig. 1)**. For each family, we compiled a metric called "range", which is the difference in the number of divisions performed by cells sharing the same ancestor. Most dividing families displayed high concordance in division: for 786 CD90$^+$ and 848 CD90$^-$ families, all cells within the same family had the same division number (range = 0) after 72 and 96 hours. Only 25% of families were spread over more than one generation (range > 1), over all our datasets **(Fig. 2h, Extended Data Fig. 5d-f)**. As cell death and sampling effects could potentially make families look more concordant in division, we developed an improved version of the previously published mathematical model[14] to account for sampling and cell death. We did not detect cell death during the first 96 h, using live-imaging data, and we estimated a recovery rate of 0.6 **(Extended Data Fig. 1c)**. We fitted the model to the family distribution over range, to estimate the correlation in division progression decisions among

cells within a family. High correlation coefficients (70–85%) **(Fig. 2h)** resulted in the best fit to the measured familial ranges, demonstrating that cell families progress through division together. We used permutation testing to assess if families sharing the same microwell would influence each other, for their properties of cell division and differentiation. We did not see such an effect, as families sharing the same well retained both different division times and cell type composition **(Extended Data Table 1).** These results suggest the existence of a cell-intrinsic clonal memory of division and lineage commitment over several divisions.

Figure 2. Clonal memory of fate and division in human HSPCs a) *MultiGen experimental workflow: HSPCs stained with four combinations of CFSE and CTV were sorted in 96-wells tissue plates: four cells per well, each one from a different CTV/CFSE combination. After 72 and 96 hours in culture, plates were analysed by flow cytometry.* ***b)*** *Distribution of the maximum division reached by individual families, derived from CD90$^+$ and CD90$^-$ and expanded in two media (Diff and GT) after 72 hours (7 independent experiments, n = 235 for CD90$^+$ GT, n = 248 for CD90$^-$ GT, n = 135 for CD90$^+$ Diff, n = 167 for CD90$^-$ Diff).* ***c)*** *Proportions of cell types per families derived from CD90$^+$ and CD90$^-$, cultured in the GT and Diff media for 72 hours.* ***d-f),*** *UMAP projection of 6148 individual cells from 13 independent experiments, colour coded by* ***d)*** *division (based on CTV/CFSE fluorescence),* ***e)*** *surface markers-based phenotype and* ***f)*** *family (displaying the ten largest families).* ***g)*** *Distribution of the number of cell types per family, compared with 250,000 permutations of the experimental data (CD90$^+$ and CD90$^-$ in Diff medium, after 72 hours).* ***h)*** *Percentage of families per division range, defined as the difference between divisions*

Two distinct clonal memories for division and lineage commitment within cell families

While we observed clonal memory in division and lineage commitment within HSPC families,
we also observed that lineage commitment progresses with division. Therefore, these factors
could influence each other: for example, the concordance in division being a consequence of
lineage commitment progressing with division or vice-versa. In this line of thought, while
concordant and discordant families showed the same diversity of cell subsets **(Fig. 2i,**
Extended Data Fig. 5g-i), individual discordant families (range > 0) displayed more progenitor
subsets than concordant families **(Fig. 2j, Extended Data Fig. 5j-l)**. We next tested whether
the presence of one clonal memory of division or lineage commitment or both would best
explain the similarity in lineage commitment and concordance in division. To this end, we
applied a stochastic modelling framework with approximate Bayesian computation (ABC)[34–
36] to the live-cell imaging and the MultiGen data combined. Briefly, we built a generic model
of the HSPCs division and differentiation over 96 h **(Fig. 3a)** starting from one cell of a known
phenotype in a well and considering four possible phenotypes: CD90$^+$, CD90$^-$, HPC, or CD34$^-$.
In the model cells divided at random times, according to log-normal probability
distributions[37,38] for each cell type. Differentiation only happened after division[39] and de-
differentiation was not allowed. We tested distinct modelling scenarios, whereby cells from
the same family could be similar (or not) in lineage commitment and/or concordant in division
(Fig. 3b). In all scenarios, symmetric and asymmetric division were allowed. In the scenarios
where we modelled the similarity in lineage commitment, we introduced - as a parameter -
the strength of this similarity in lineage commitment between daughter cells. To compare
how the different modelling scenarios fitted the entire dataset, we used a distance that
aggregated the derivation of many summary statistics. We ran simulations with different
parametrizations of the models and then compared the posterior probability of those models.
We found that the models lacking similarity in commitment within the cell family were
inconsistent with the data. The best model, with the higher posterior probability, displayed
both similarity in commitment and concordance in division for cells from the same family. The
model in which cells from the same family were only similar in commitment but not
concordant in division had a lower posterior probability **(Fig. 3c).** As a control, we tested if a
minimal scenario with no correlation between division and differentiation and no clonal
memory could explain the data, we assumed that the probability distributions of the time to
divide was identical across cell types. Unsurprisingly, this scenario had a lower posteriori
probability, making it unlikely to explain our data. Overall, these results demonstrated that
two distinct clonal memories for division and lineage commitment underlie the similarity in
lineage commitment and concordance in division.

***Figure 3. Two distinct clonal memories of division and lineage commitment** a)* *Overview of the*
modelling strategy: proliferation and differentiation for each well were simulated by agent-based modelling for
different scenarios. For each simulation, simulated data were extracted to be compared with both the live-cell
and MultiGen datasets. From the live-cell imaging datasets, the experimental parameters used were the values
of time of 1st (T_1), 2nd ($T_{2,1}$ and $T_{2,2}$) and 3rd divisions ($T_{3,1}$ to $T_{3,4}$) and the colony size at T_{max}. From the MultiGen
datasets, the relative percentages of cell types and the associated cell divisions were used. Green, cyan, orange
*and dark blue correspond to CD90⁺, CD90⁻, HPC, and CD34⁻ types, respectively. **b)** The four distinct modelling*
scenarios tested, based on whether there is similarity in fate (H=1) or not (H=0), and whether there is concordance
*in division (C=1) or not (C=0). **c)** Value of the posterior probability of each scenario $\mathbb{P}[(H,C)|\mathcal{D}]$, given the*
experimental observations $\mathcal{D}$, after running the ABC (Approximate Bayesian Computation) procedure. We
included also a fifth modelling scenario assuming no cell memory of cell division and cell differentiation and time-
to-divide distributions that are independent of the dividing cell type.

Clonal transcriptional programs of clonal memory in human HSPCs

To explore the transcriptional programs involved in the clonal memory of human HSPCs, we
developed an experimental protocol integrating lineage tracing and molecular profiling with
the division history of each cell. Briefly, we FACS-sorted individual CD90⁻s from two cord blood
samples and cultured them *ex vivo* in the Diff media while recording their division history by
live-cell imaging over 90 hours. We then re-isolated individual cells from randomly selected
wells, using low throughput, multi-fluorescence image sorting[40], and profiled each cell with
high-coverage single-cell RNA sequencing (scRNA-seq) via SMART-seq3[41] **(Fig. 4a)**. We
obtained two multidimensional datasets containing information on division, family, and
transcriptome for 142 individual cells from 30 families after QC and filtering **(Fig. 4b, Extended**
Data Fig. 6d, Extended Data Table 2). After supervised annotation using known marker genes,
we identified several cell subsets at different stages of myelo-erythroid differentiation:
immature progenitors (HSPCs), erythroid-megakaryocytic progenitors (EryP/MkP) and
myelomonocytic committed progenitors (MyP and GraP) **(Fig. 4c-d, Extended Data Fig. 6e-g)**.
These subsets globally matched the subsets we identified via CITE-seq (cellular indexing of
transcriptomes and epitopes by sequencing) profiling of 8,580 cord blood HSPCs after 4 days
in Diff media **(Extended Data Fig. 6a-c)**.

We next explored the similarity of cell families in the transcriptomics space. We observed that cell families co-localized within the same UMAP clusters, suggesting that cell from the same family share a similar transcriptome **(Fig. 4b-c, Extended Data Fig. 6d-e)**. To quantify the transcriptomic similarity between cells from the same family, we compared the median cosine distance[42] distributions between the observed cell families and artificial families made of permutated cells. Cell families had lower median cosine distance than permutated families **(Fig. 4e)**, demonstrating that cells within a family are more transcriptionally similar than cells between families, in line with the MultiGen results. Moreover, this result held after permutating cells within the same cell type subsets **(Fig. 4e),** suggesting that transcripts associated with cell subsets alone were insufficient to explain this similarity. Thus, biological processes other than those associated with lineage commitment contributed to the similarities between the cells within a family.

To characterize the transcriptional programs associated with the familial similarity[9,43], and therefore with clonal memory, we extracted the genes that varied the least within a family and the most between families **(Fig. 4f, Extended Data Fig. 6i)**. We could then identify transcriptional fingerprints specific for each family, contributing to the unique intrafamilial similarities beyond fate commitment. These signatures contained genes associated to many biological processes, including membrane trafficking (e.g., *RAB31, CAVIN2, VAMP8*), cell proliferation (e.g., *CASP3, IGFBP2*), cytoskeleton structure (e.g., *LESP1, ANK1, MYH10)*, cell signalling (e.g., *IL17RA, IL1RL1, LAT2*), and metabolism and biosynthesis (e.g., *DLK1, CALR, XBP1*) **(Fig. 4f; Extended Data Table 3)**. To identify common genes between families that contributed to clonal memory, we ran the Multivariate Information-based Inductive Causation (MIIC) causal network reconstruction algorithm[44] on both transcriptomic datasets using the transcriptomes, the family and the division information from live imaging of each cell **(Fig. 4g; Extended Data Fig. 7a, Extended Data Table 4)**. The MIIC algorithm is a multivariate information-based network reconstruction method[45,46] that reconstructs causal and non-causal graphical models from heterogeneous data. The MIIC analysis identified the variables "family" (sharing the same ancestry), "predicted cell type" and "distance" (transcriptomic similarity) as key hubs in the networks **(Extended Data Fig. 7b)**. Gene expression of *IGLL1* and surface markers like *CD36* and *CD84*, amongst others, were strongly associated with the "family" hub **(Fig. 4g)**. Notably, the transmembrane protein STAB1, which directly connected with the "family" hub, emerged as a central hub within the gene network, orchestrating interactions with a broad array of other genes **(Extended Data Fig. 7c)**. In contrast, variables associated with division properties, such as division time and colony size, were only directly connected to less than 5 genes, and were strongly related to the "family" variable **(Extended Data Fig. 7d)** as all these families were synchronous in division **(Fig. 1d-f)**. The limited number of genes associated with division-related properties suggests that the transcriptome is not a suitable proxy for these properties, or that the relevant genes are expressed at low levels and thus remain undetected. Finally, the genes associated to the variable "family" were not equally expressed by all families **(Fig. 4g-h, Extended Data Fig. 7e-f)**, suggesting the existence of different transcriptomic program between families. Some genes were expressed in all families but at different levels **(Fig. 4h)**, while others were specific to some families **(Fig. 4i)**. In summary, human HSPC families displayed heritable

transcriptomes linked to several cellular processes, including cell commitment, that contributed to clonal memory.

Figure 4. Clonal transcriptional programs of clonal memory in human HSPCs *a) Scheme of the experimental workflow: individual CD90⁻ were FACS-sorted in 96-well plates and live-imaged for 90 hours (Diff medium). Then, individual cells from 30 randomly selected families were re-sorted by image-based sorting into 384-wells plates and processed by SMART-Seq3. **b-c-d)** UMAP representations of a transcriptomic dataset (70 cells), colour-coded by **b)** family and **c)** annotated cell type, one experiment representative of two and **d)** pseudotime. **e)** Distribution of the gene expression median cosine distance for real cell families (red) and two permutation-generated distributions, either retaining (purple) or discarding (green) the cell type of the permuted cells. Wilcoxon rank-sum test, p value < 0.0035 for family/permutation discarding cell type and p value < 0.02 for family/permutation retaining cell type. **f)** Heatmap displaying the z-score of the 371 most variant genes (rows) between families. 24 randomly selected gene names are displayed on the right side. Cells (columns) are colour-coded according to their original family. **g)** Snapshot of the MIIC network generated using transcriptomic and divisional data, highlighting the features directly connected to Family. Variables in orange are causal regarding Family, those in green are consequential. Edges represent an association between two variables, while edges with an arrow represent a causal relationship. Red edges highlight positive association, blue edges represent negative associations, and grey edges represent categorical association with no sign. A green arrow indicates genuine causality, while a red, blue or grey arrow stands for putative causality (i.e. the presence of an unobserved common cause between the two variables cannot be ruled out). Full network with detailed legend is shown in Extended Data Figure 8a. **h-i)** Dataset A expression of a selection of "family"-associated genes accordingly to the*

The clonal memory of division is disrupted in leukemic cells

AML is a proliferative disease with a differentiation blockade, resulting in a disproportionate production of blast cells. To test if this disruption affects the cellular clonal memory, we analysed primary CD34[+] blasts from AML patients **(Extended Data Table 5),** that had been isolated from bone marrow or peripheral blood and cultured for at least 72 hours in Diff media supplemented with IL-1β[47], using live-cell imaging and MultiGen. By live-cell imaging, the intragenerational correlations in the division time between daughter and cousin cells were diminished **(Fig. 5a, Extended Data Fig. 8a).** As expected, blasts displayed increased number of divisions **(Extended Data Fig. 8c)** and division time **(Extended Data Fig. 8d-f)** when compared with cells from healthy controls. The MultiGen assay showed that CD34[+] AML cells possessed a lower clonal memory of division than healthy HSPCs, as the concordance in division ρ was lower than for control cells from healthy blood donors **(Fig. 5b).** Stratifying the analysis per ancestor cell types (i.e., CD90[+]s, CD90[-]s, HPCs), we measured clonal memory of division reduction for all cell types **(Extended Data Fig. 8b, 8i-j),** even if we did not observe significant changes in the differentiation profile **(Extended Data Fig. 8g)** or in the number of divisions over the 72 hours **(Extended Data Fig. 8h)**. We excluded the impact of genetic heterogeneity, as we measured synchronicity for sisters/cousin cells, all sharing the same ancestor genomic background. These results revealed that primary CD34[+] blasts from AML patients had a lower synchronicity of division within families.

We observed some variability in the synchronicity in division between patient samples **(Fig. 5c),** with best-fitted correlation coefficients of division within families varying between 53% and 72%. Nevertheless, this variability was uncorrelated to the patient mutation profile, clinical history **(Extended Data Table 5)** or the ability of the blasts to expand *ex vivo* **(Extended Data Fig. 8k-n)**. These findings demonstrated a reduction in the synchronicity of primary AML blast divisions, suggesting a disruption of the clonal memory that orchestrates cell division. Decreasing clonal memory could be a strategy for AML blasts to burst their heterogeneity in division and promote adaptability over stability.

Clonal memory of division within AML HSPC families can be modulated

Finally, we assessed if the clonal memory in division was a plastic trait that can be modulated. As we observed that the clonal memory was independent of genetic mutations, we hypothesized an epigenetic mechanism as central to clonal memory[3,7,9,43]. To this end, we tested the effects of JQ-1, a pan-bromodomain and extra-terminal domain (BET) inhibitor[48] on the clonal memory of AML cells.

First, we used a leukaemia stem cell (LSCs) model of differentiating AML[49,50], the OCI_8227, to test the effect of JQ-1 on clonal memory **(Extended Data Fig. 9a)**. We selected the drug dose of 5 nM, as it induced little death and did not cause significant changes in cell cycle

progression and cell division number **(Fig. 5d, Extended Data Fig. 9b)**. Without JQ-1, the OCI_8227 cells displayed a decreased clonal memory in division, similarly to primary CD34[+] blasts, with fewer families in range 0 and low correlation coefficients of division within families. After exposure to JQ-1 (5 nM), the cell families that divided simultaneously increased by approximately 10%, and the best fitted correlation coefficient by 1.5-fold **(Fig. 5e).** We concluded that the clonal memory of cell division could be modulated, in this case increased with JQ-1. As the JQ-1 treatment affected the differentiation profile **(Extended Data Fig. 9c)** but not the total number of divisions **(Extended Data Fig. 9d)** or the clonogenicity **(Extended Data Fig. 9e),** we concluded that JQ-1 inhibition modulated only the division clonal memory but not the division capacity of cells, suggesting that these two processes are regulated by different mechanisms[51].

We obtained similar results in primary CD34[+] blasts from AML patients. JQ-1 treatment increased the synchronicity between sister cells during their second divisions **(Fig. 5e)** without significantly modifying their division time **(Extended Data Fig. 9f-j).** Using the MultiGen assay, JQ-1 treatment also increased the percentage of families in range 0 and the correlation coefficients of division within families **(Fig. 5g)** as compared to vehicle, without changing either the family progression through divisions **(Fig. 5h)** or their surface phenotype **(Extended Data Fig. 9j).** These results were confirmed using samples from two other patients **(Extended Data Fig. 9k-r)**. Thus, as the clonal memory in division increased *ex vivo* in primary AML CD34[+] blasts when exposed to an agent acting on the cellular epigenetic landscape, we concluded that clonal memory can be modulated.

Figure 5. Reduced concordance in division of leukemic families can be restored *a) Correlation*
in division time (in hours) between sister cells after 1 division, for CD34$^+$-derived families from AML patient #30-
*001-20. Spearman's rank correlation coefficient R (n = 154). **b)** Percentage of CD34$^+$-derived families per division*
range, for the same patient and a healthy control (HC, #200-061), after 72 hours. The red bar represents the
data, the dashed blue bar represents the correlation coefficient ρ best fit, the filled blue bars represent
*simulations for different parametrization of ρ, n = 120. **c)** Summary of correlation coefficients as in **a)** and **b)** for*
9 AML samples (red) and three healthy controls (green), analysed with live-cell imaging, MultiGen or both. NA
*for not available. **d)** Proportions of OCI_8227 cells in G1-, S- and G2/M-phase, after 72 hours exposure to JQ-1 (5*
*and 50 nM), 2-way ANOVA test error bars indicate SD, **** for p < 0.001, n = 3. **e)** Percentage of families per*
division range for OCI_8227 treated for 72h with PBS (control) or JQ-1 5 nM (treatment). The red bars correspond
to the data (n = 250 and 327 respectively, four independent experiments), the dashed blue bar represents the ρ
*best fit. We used a recovery rate of 0.6 and a survival rate of 0.95. **f)** Correlation in division time (in hours)*
between sister cells after 1 division, for LMPP-derived families for patient #127-002-23 exposed to PBS and JQ-1
*for 96 hours (n = 32 for both). Spearman's rank correlation coefficient R. **g)** Percentage of families per division*
range for LMPP--derived families from patient #127-002-23, treated for 72h with sublethal doses of JQ-1 or PBS
(n = 69 and n = 68 respectively). The red bars correspond to the data, the dashed blue bar represents the ρ best
*fit. **h)** Distribution of the maximum division reached by individual families treated for 72h with sublethal doses of*
*JQ-1 and PBS (n = 69 and 68 respectively). For the panels **b, e, g** and **h** the error bars indicate 95% confidence*
intervals calculated via basic bootstrap. The p values are calculated using permutation testing, ns for p > 0.05.

Discussion

Our study revealed that human HSPCs display two distinct forms of clonal memory— one for
division and one for fate commitment. These clonal memories were consistently observed
across multiple progenitor types, for more than two successive divisions, and diverse culture
conditions, suggesting that clonal memory is a fundamental and conserved feature of early
haematopoiesis. To the best of our knowledge, this is the first demonstration of both forms
of clonal memory in human HSPCs. Clonal memory of differentiation has been previously
described in murine haematopoiesis[52,53], and we previously identified clonal memory of
division in mouse[14]. They also align with indirect *in vivo* evidence of proliferation inheritance
in twin patients developing the same neoplasms[54,55] and in donor and recipient of bone
marrow transplants [20–22]. Although clonal memory of division has been reported in other cell
systems[56–58], such memory has not yet been described in differentiating systems.

As differentiation and division are linked, both factors could influence each other and produce
synchronicity or similarity in fate within cells without clonal memory. Using our mathematical
model, we demonstrated that clonal memory is required to explain the observed synchrony
and similarity in fate. In fact, the presence of two distinct forms of memory for division and
differentiation explained best the data, raising the possibility that they are governed by
separate regulatory mechanisms. This echoes findings in B cell differentiation, where division
and differentiation are independently clonally inherited[59]. Whether these two memory types
are independent or coordinated remains an open question.

Interestingly, the division time of cells from the same family across generations did not
correlate. One might think that it is not the division time itself that is transferred across
generations, but the ability to divide at the same time. This mechanism was observed accross
cell type, suggesting that it could be independent from commitment, but we do not exclude

the fact that cells sharing the same differentiating phenotype could also share this mechanism. Advanced experimental approaches allowing to assess simultaneously the differentiation status and the division times[60] will help elucidate this point.

We also identified that transcriptional fingerprints, gene modules shared by each member of that family, extend beyond differentiation states[43], implicating clonal identity in shaping gene expression. Proliferative parameters such as clone size and division timing are poorly captured at the transcriptomic level, indicating that these features may be regulated through epigenetic or post-transcriptional mechanisms. Our results confirm that clonal analysis by transcriptomics[3,4,56–58,61] is an effective method when the cells analysed display a strong clonal memory (cells from the same family display high transcriptional similarity). This should be confirmed beforehand, as we could observe a non-negligible fraction of discordant, more heterogeneous families, which could bias the clonal analysis by transcriptomics.

The existence of clonal memory in healthy haematopoiesis represents a form of non-genetic inheritance under non-pathological conditions. For this reason, we use the term *family* rather than *clone* to describe cell genealogies independently of the underlying cellular genetics. Our work focused on early haematopoiesis, the stage at which initial commitment decisions are made. Further studies are needed to determine how long clonal memory persists through subsequent stages of haematopoietic differentiation.

Functionally, clonal memory of division may serve to maintain proliferative heterogeneity among haematopoietic stem cells, a feature critical for homeostasis and stress-induced haematopoiesis. Such heterogeneity could enable distinct cell families to contribute to steady-state haematopoiesis versus responses to environmental fluctuations (e.g. inflammation, blood loss). Unexpectedly, we observed a marked reduction in clonal memory of division in AML, which could be partially restored by treatment with the epigenetic modifier JQ-1. This demonstrates that clonal memory is not only regulated but can also be tuned pharmacologically. While AML is composed of genetically distinct clones, our data reveal that proliferation rates vary even within genetically defined families. This intra-clonal variability may reflect a strategy by which leukemic cells reduce memory to enhance adaptability. Whether this reduction is a dynamic, reversible process or a fixed trait of selected clones remains to be determined.

The ability to modulate clonal memory pharmacologically opens new therapeutic avenues. By shifting the balance from plasticity toward stability, it may be possible to reduce intra-tumour heterogeneity and sensitize malignant cells to treatment. More broadly, our findings position clonal memory as a key regulatory layer in stem cell biology, with implications for both normal development and disease.

Limitations of study

To enhance the reproducibility of our findings, we employed several culture conditions, including at least two distinct media and multiple timepoints, to ensure the reproducibility of

our observations. However, future studies should aim to better capture the influence of the microenvironment, which we could only partially addressed here. We prioritized the analysis of many individual families per sample, particularly for the pathological ones, and the inclusion of a representative, yet incomplete selection of AML samples. We acknowledge the small number of samples displayed, which are not necessarily recapitulating the genetic and phenotypic heterogeneity associated to AML. The magnitude of the decrease in cell division concordance, or the JQ-1-associated increases, differed between samples, underlying different sensitivities to JQ-1 and unknown biological determinants associated to cell division synchronicity. Furthermore, our study faced the classical technical challenges with primary AML samples, notably: (1) the limited capacity for *ex vivo* expansion of primary leukaemia samples, especially in the absence of supportive stromal cells, and (2) the inherent constraints of single-cell culture systems. Continued development of advanced culture platforms will enable to expand our analysis of primary AML cell proliferation and differentiation.

Material and Methods

Ethical approval, banking, and processing of human samples

De-identified umbilical cord blood units were collected accordingly to the guidelines defined by the Saint-Louis Hospital Umbilical Cord Blood biobank (authorization AC-2016-2759) and in accordance with the Declaration of Helsinki.

Human Bone Marrow samples were collected accordingly to the guidelines defined by the Saint-Louis Hospital Institutional Review board (IRB 00003888), authorization 21-799.

Human AML primary samples (bone marrow or peripheral blood) were obtained from patients with informed consent for sample collection and use in research were obtained accordingly to the IRB-approved procedures from Institut Curie (Saint-Cloud, France), Gustave Roussy (Department of Clinical Haematology and Drug Development Department DITEP, Villejuif, France) and Hammersmith Hospital (London, UK). A summary of the patient samples used for this study can be found in **Extended Data Table 5**.

Mononuclear Cells (MNC) were isolated via Ficoll density gradient. Human bone marrow cells were systematically processed fresh, while umbilical cord blood units and human AML primary samples underwent CD34-enrichment using immunomagnetic microbeads (MACS Miltenyi). Total MNCs or enriched CD34$^+$ fractions were immediately processed or frozen in Fetal Bovine Serum (FBS, Eurobio Scientific) supplemented with 10% dimethyl sulfoxide (DMSO, Sigma) for further analysis.

Fluorescence Activated Cell-Sorting (FACS) and single-cell isolation

Total MNCs or enriched CD34$^+$ fractions were prepared for immunostaining as previously described[17]. The list of antibodies used with the respective fluorochromes can be found in Extended Data Table 7. Single-cell FACS was performed using BD Aria III or BD Fusion instruments (Becton Dickson), in 96-wells U-bottom tissue culture vessels (Falcon). Cells were isolated using Single Cell sorting precision when individually isolated or Purity precision when isolated in bulk. The sorted cells were indexed for all the fluorescence parameters, using the Index sorting function, to record the fluorescence levels for each marker. Data analysis was performed using the software FlowJo v10.1 (BD Biosciences).

MultiGen cell dye labelling

The cell dye labelling step was performed as previously described[17]. Briefly, cells were stained with four combinations of 5-(and 6)-carboxyfluorescein diacetate succinimidyl ester (CFSE, ThermoFisher Scientific) and CellTrace Violet (CTV, ThermoFisher Scientific): CFSE only, CFSEhigh + CTVlow, CFSElow + CTVhigh and CTV only.

For umbilical cord blood HSPCs, the labelling was performed in two sequential steps, starting with the CFSE staining and concluding with the CTV staining. The final concentration used for the labelling are respectively CFSE 5 µM (CFSE only and CFSEhigh CTVlow), CFSE 2.5 µM

(CFSElow CTVhigh), CTV 5 μM (CTV only and CFSElow CTVhigh) and CTV 2.5 μM (CFSEhigh
CTVlow).

For OCI_8227 cells, the labelling was performed in a single step, preparing 2X solutions for
each of the cell dye combination. The final concentration used for the labelling are
respectively CFSE 2.5 μM (CFSE only and CFSEhigh CTVlow), CFSE 1.25 μM (CFSElow CTVhigh),
CTV 2.5 μM (CTV only and CFSElow CTVhigh) and CTV 1.25 μM (CFSEhigh CTVlow).

For primary human AML cells and healthy adult peripheral blood, the labelling was performed
in a single step, preparing 2X solutions for each of the cell dye combination. The final
concentration used for the labelling are respectively CFSE 5 μM (CFSE only and CFSEhigh
CTVlow), CFSE 2.5 μM (CFSElow CTVhigh), CTV 5 μM (CTV only and CFSElow CTVhigh) and CTV
2.5 μM (CFSEhigh CTVlow).

Short-term liquid culture of primary cells

Progenitors isolated from umbilical cord blood or healthy bone marrow, either as single cells
or in bulk, have been cultured in two types of cell culture medium, named Gene Therapy
medium (GT)[27] and Differentiation medium (Diff)[26]. Both media are based on StemPro-34 SFM
(SP34, ThermoFisher Scientific) supplemented with Penicillin – Streptomycin 1X (Pen/Strep,
ThermoFisher Scientific), L-Glutamine 2 mM (ThermoFisher Scientific) and human Low-
Density Lipoprotein 500 ng/mL (LDL, Stem Cells). For GT medium we added the following
cytokines: SCF 300 ng/mL (Peprotech), FLT3-ligand 300 ng/mL (FLT3l, Peprotech), TPO 100
ng/mL (Peprotech), IL-3 60 ng/mL (Peprotech). For Diff medium: SCF 100 ng/mL, TPO 100
ng/mL, IL-6 50 ng/mL (Peprotech), GM-CSF (Miltenyi) 25 ng/mL, FLT3l 20 ng/mL, EPO 1 U/mL
(Janssen) and IL-3 10 ng/mL. Progenitors isolated from AML samples, both as single cells
orbulks, were cultured in a slightly modified version of the Diff medium, containing IL-1β 50
ng/mL (Peprotech). A complete list of references for cytokines and small molecules is
available in **Extended Data Table 6**.

MultiGen flow cytometry assessment of cell division, cell differentiation and cell family

Flow cytometry experiments, both using umbilical cord blood progenitors, healthy and
leukemic peripheral blood and 8227 cells, were performed as previously described[17]. The list
of antibodies used with the respective fluorochromes can be found in **Extended Data Table
7**. Data analysis was performed using the software FlowJo v10.1 (BD Biosciences) and custom
Python scripts. The recovery rate was experimentally measured following the same protocol,
except that each well was seeded with a range of CTV-stained 8227 cells (4 to 16 cells).
Immediately before the flow cytometry analysis, each well was manually assessed and the
number of light-refringent cells assessed via an inverted microscope. The recovery rate was
therefore calculated as a ratio between the value from the flow cytometry analysis (CTV+ cells
in the Single Cells gate) and the microscope value (Extended Data Figure 1c-d).

Time-lapse imaging

Haematopoietic progenitors from umbilical cord blood or adult bone marrow were, as previously mentioned, individually sorted via FACS in 96-wells U bottom plates (Falcon) and then continuously imaged for 96 hours, using an Incucyte Zoom system (Sartorius). Cell culture was performed using the same culture media previously described. Time-lapse experiments were conducted at 37 °C, 5% CO_2, with time intervals of bright field image acquisition of 1-2 hours. Cell viability and cell division were assessed manually: high refractility, membrane integrity and cell size were the criteria used to assess viability, and cell death time was measured as the first image when visible blebbing could be observed, together with cell fragmentation, loss of circularity and lack of cell movement. The first cell division time was measured as the time between the beginning of the experiment and the time point at which two cells could be clearly distinguished. For the second and third divisions, the time was measured as the interval between the end of the former division and the time point at which two novel cells could be clearly distinguished and were adjacent in the same region. We stopped to measure the time for cell division after the third division, due to cell motility making impossible to reliably track each individual cell manually. For the analysis of the synchrony in division for the second divisions, we randomised the order of annotation of the two division times, to avoid any bias in the two distributions.

Cell line culture and *ex vivo* treatments

OCI_8227 cells were maintained in culture for no more than eight weeks at the time. Cells were passaged every 5 days in T-25 or T-75 cell culture-grade flasks, at a cell concentration of 1.5-3 x 10^5 cells/mL. Cells were cultured in StemSpan II (Stem Cells) supplemented with Pen/Strep 1x, SCF 50 ng/mL, TPO 50 ng/mL, FLT3l 50 ng/mL, G-CSF 10 ng/mL, IL-6 10 ng/mL and IL-3 10 ng/mL. 8227 cells were treated *ex vivo* using sublethal doses of JQ-1 (Sigma): the working solutions were prepared accordingly to the manufacturer's instructions.

Confidence Intervals for MultiGen analysis

The confidence intervals at 95% level were calculated via basic bootstrap[62] with 250,000 bootstrap datasets. Following this procedure, each bootstrap dataset was constructed by sampling with replacement as many cellular families as were in the original data. The distribution of the statistics, each calculated from one bootstrap dataset, then provided a reference from which the confidence interval was derived. Formally, given the statistics ϑ^ calculated from the original data, and $\vartheta*_{(0.025)}$ and $\vartheta*_{(0.975)}$ percentiles, respectively derived from the bootstrapped distribution, the confidence interval of ϑ^ at 95% level was calculated as follows:

CI(95%) = $(2\vartheta - \vartheta*_{(0.975)}, 2\vartheta - \vartheta*_{(0.025)})$

Cell division range analysis

To quantify how correlated were the cell decisions to divide, for cells from the same generation and descending from the same ancestor, we employed a previously used stochastic model[14,18,19], upgraded to account for cell death. Biologically, each cell reaching a certain generation have a chance of surviving that is independent from other cells. All surviving cells can then divide or stop dividing, with a given probability that is correlated between cells from the same family and generation. When sampling, cells that survive but divide less tend to be also less detected, simply because they are less abundant. Thus, it is important to take both survival and sampling into account when measuring the range of division within a family. In this model, cells have a certain probability to survive before dividing, as defined by the survival rate r_{surv}. Cells have also a certain probability to be measured, defined by the recovery rate r. The correlation coefficient that links the cell division within the same family was fitted to the data and was used to evaluate the degree of correlation in division within families, otherwise called concordance in division. The model was directly parameterized by the data, apart from one variable that encapsulated the correlation in decision-making that fitted to the data. With n being the maximum generation recorded, let $p_i \in [0, 1]$ for $i = 0, …, n$ denoted the proportion of cells that divided from generation i to the next, which was determined from the data as follows. Set z_i the total number of cells recovered in generation I, the p_i was estimated by

$$p_i = \frac{\sum_{j=i+1}^{n} z_j\, 2^{-j}}{\sum_{j=i}^{n} z_j\, 2^{-j}}$$

For $i = 0, …, n$-1 and by $p^{\wedge}_n = 0$.

In this model, given k_i the number of cells from a particular family that reach generation i, the number of cells that continued dividing to $i + 1$ followed a beta-binomial distribution with a number of trials that was a random variable K_i (independent of all the others) and parameters $a_i = p_i (1 − \rho)/\rho$, and $b_i = (1 − p_i)(1 − \rho)/\rho$, namely $\beta(K_i, a_i, b_i)$, where $\rho \in [0, 1]$ was a free parameter.

To account for cell death, $K_i = B(k_i, r_{surv})$ followed a binomial distribution with k_i that reached generation i and survived to divide. r_{surv} was estimated from time-lapse microscopy experiments for each cell type and specified in the results section. Finally, one family was generated recursively by setting $k_o = 1$ and defining $k_{i+1} = 2\beta(K_i, a_i, b_i)$. The recovery rate r was determined experimentally as previously described, and set at 0.6. In the model, we accounted therefore for the sampling effect by sub-sampling each cell from every artificially generated family with a probability of 0.6 and independently of all other cells. The beta-binomial model interpolated between surviving cells that decided to divide independently of each other ($\rho = 0$) and cells that were perfectly aligned in their decision to divide ($\rho = 1$). A value between 0 and 1 reflected the concordance level within each family in division, but, by design irrespective of the values of p_i, that determined the population distribution among generations. Defining the family range as the difference between maximum and minimum generation for all recovered surviving cells per family, the best fit $p^{\wedge}$ of p was determined to be the value from 0 to 1 that maximized the likelihood of recapitulating the family range distribution in the data. The 95% confidence interval for $p^{\wedge}$ was calculated following the basic

bootstrap procedure previously described, that was by sampling with replacement the
cellular families and setting the maximum-likelihood estimation as $p^\wedge$ the statistic to be
calculated on the bootstrapped datasets. To assess the significance of $p^\wedge$, the statistical
hypothesis $H_0 : \rho = 0$ was tested against the alternative $H_1 : 0 < \rho \leq 1$ using the likelihood-ratio
statistic

$\lambda_{LR} = -2\,(L(0) - L(\hat{\rho}))$

where $L(\rho)$ was the log-likelihood of the cellular family ranges under the beta-binomial model
with parameter p. The p-value for the test was derived from the asymptotic distribution of
λ_{LR} that was a χ^2 with 1 degree of freedom.

Mathematical modelling and statistical inference of the HSPC proliferation and
differentiation dynamics

The detailed description of the mathematical model, the parameter estimation and model
selection procedure, and supplementary results are available in the Supplementary
Information (SI).

Model assumptions: we considered a stochastic model, at the scale of a family, continuous in
time (denoted by t), over a short period ($T_{max} = 96h$), and where each cell was characterized
by a phenotype p, with four different possibilities: $p \in \mathcal{P} = (CD90^+, CD90^-, HPC, CD34^-)$,
where $\mathcal{P}$ was naturally provided with an order relation (e.g., a CD90$^-$ being more
differentiated than a CD90$^+$). Therefore, we confounded $\mathcal{P}$ with the set [1,2,3,4]. $p = 3$ would
then correspond to an HPC. At $t = 0$, the process begun with a single cell of a known phenotype
p_1 **(Fig. 3a)**. The initial cell division occurred at a time T_1, randomly drawn from the empirical
distribution of observed first division times (of cells of type p_1) (for further details, see **SI –**
B.3.1). We treated separately the first division from the subsequent ones since we observed
that the initial division typically exhibits a longer duration **(Extended Data Figure 2j-l)**. After
division, two daughter cells were considered. No cell death was considered in our model, and
each of the daughter cells was assigned a cell phenotype at least as differentiated as the
mother cell (we assume no de-differentiation). The probability for a daughter cell to be of
type $p = i$, given its mother's type $p_m = a$, was denoted by $p_{a \to i} = \mathbb{P}[\, p = i \mid p_m = a]$. Then,
the cells pursued the proliferation process, with division times randomly sampled from Log-
Normal distributions ($\mathcal{LN}$) where the distribution parameters (μ_p, σ_p) varied according to the
dividing cell's phenotype: $T \sim \mathcal{LN}(\mu_p, \sigma_p)$. We noted that a Gamma distribution would have
been appropriate as well, as we describe in Supplementary Material B.3.2.

Possible scenarios: The generic model of HSPC proliferation and differentiation could be
declined into several scenarios to explore different biological hypotheses **(Fig. 3b)**. These
scenarios modify the interactions between sister cells, to reflect distinct biological hypotheses
that we evaluated for their agreement with our experimental data. Scenarios were defined
based on two criteria: homogeneity in fate (H) and concordance in division timing (C).
Homogeneity (H) assesses whether sister cells tended to share the same fate (H=1) or not
(H=0). Concordance (C) evaluated whether sister cells tended to divide simultaneously, with

C = 1 for synchronised divisions and C = 0 for independent timing. Each scenario was identified

by a pair of values for (H, C), creating a matrix of possibilities. The default scenario,

characterized by H = 0 and C = 0, posited that sister cells were fully independent from each

other (conditioning on their mother). This scenario served as a baseline, against which the

implications of fate homogeneity and division concordance were assessed. When we

hypothesized a homogeneity in fate (H = 1), we considered in our model a higher probability

to have a symmetrical division compared to the scenario where H = 0. To do that, we

introduced the parameter $\rho_H \in [0,1]$ such that the probability of having two daughter cells

of the same type b, given their mother's type a, was equal to: $\rho_H\, p_{a\to b} + (1 - \rho_H)p_{a\to b}^2$. In

the limiting case where $\rho_H = 1$, all divisions were symmetrical. The case $\rho_H = 0$

corresponded to the hypothesis H = 0 of absence of homogeneity in fate. When we

hypothesised a concordance in division (C = 1), we considered that the division times of two

sister cells were not independent anymore but correlated. Let us denote by T_i and T_j the

division times of two daughter cells of type p_i and p_j, respectively. Then, in our model, the

division times of the two sister cells followed a bivariate log-normal distribution:

751
$$\left(T_i, T_j\right)^t \sim \mathcal{LN}\left(\left(\mu_{p_i}, \mu_{p_j}\right)^t, \Sigma\right),$$

where the covariance matrix was expressed by: $\Sigma = \begin{pmatrix} \sigma_{p_i}^2 & \rho_C\,\sigma_{p_i}\sigma_{p_j} \\ \rho_C\,\sigma_{p_j}\sigma_{p_i} & \sigma_{p_j}^2 \end{pmatrix}$.

With parameter $\rho_C \in [0,1]$ quantifying to which extent sister cells were concordant.

The limiting case $\rho_C = 0$ would correspond to the hypothesis C = 0, which was absence of

concordance in division.

In addition to our four main scenarios, we also studied more complex ones, by considering

that ρ_H would depend on the mother cell type (SI – B.2.3), or that there could be concordance

in division between cousin cells, or that the concordance between sister cells would depend

on their types (SI – B.3.4). We also tested a minimal – more parsimonious – scenario, still

without concordance in division timing and without homogeneity in fate, but also without

differentiating the distribution of the division time, according to the type of cell which divides.

That is, in this minimal scenario, we assume that $\mu_4 = \mu_3 = \mu_2 = \mu_1$ and $\sigma_4 = \sigma_3 =$

$\sigma_2 = \sigma_1$.

We denote by:

$\theta = (p_{1\to 1}, p_{1\to 2}, p_{1\to 3}, p_{2\to 2}, p_{2\to 3}, p_{3\to 3}, \mu_1, \sigma_1, \mu_2, \sigma_2, \mu_3, \sigma_3, \mu_4, \sigma_4, \rho_H, \rho_C)^t$

our parameter vector.

Comparing the model simulation with the experimental observations: To evaluate which

scenario is the most likely, we compared the output of our model simulations to the

experimental observations, defining a distance between the data and simulations. We

denoted by $\mathcal{D}$ our dataset (for further details about its definition, the inclusion and pre-

treatment step, and the uncertainties in the observations, see **Supplemental Information –**

A). $\mathcal{D}$ is of size 1,740, meaning that we had the observations for 1740 different families, either

from the MultiGen or the single-cell live imaging assay, for different initial conditions, and for

different observation times. Because of the large size of our dataset, and the heterogeneity in the observations, defining a proper distance was challenging. Our methodology involved the derivation of an extensive set of summary statistics, denoted as S_i, where *i* indexes the specific summary statistic within the set. Each summary statistic can be computed from experimental observations, in that case we noted: $\hat{S}_i = S_i(\mathcal{D})$.

Given a parameter vector θ, we simulated the model to generate a virtual dataset of equivalent size to the number of experimental observations, from which we calculated each theoretical (i.e., given by our model) summary statistic $S_i(\theta)$. Summary statistics synthesised the information from the observations (or from the simulations) making comparisons possible. This was accompanied by a potential loss of information, which we first minimised by defining 340 summary statistics. Ideally, we aimed to find parameter values such that the simulations of the model lead to computed summary statistics as close as possible to the experimental ones, that is, find θ^* such that, $\forall i \in \{1, \dots, 340\}$, $S_i(\theta^*) \approx \hat{S}_i$.

However, given the number of summary statistics considered, and the potential redundancy and high correlation among them, we refined our approach using Principal Component Analysis (PCA), thus reducing the dimensionality of our observation space. We ended-up with 20 principal components, accounting for 80% of the variance, denoted as $\left(S_{p,j}\right)_{j=1}^{20}$ where each $S_{p,j}$ was a linear combination of the original 340 summary statistics.

The distance d(θ) was then defined as a mean squared error (MSE) between the simulated and observed values of these principal summary statistics:

$$d(\theta) = \sum_{j=1}^{20} \left(S_{p,j}(\theta) - \hat{S}_{p,j}\right)^2$$

This approach provided a practical balance between simulation accuracy and computational cost, making it well-suited for methods like Approximate Bayesian Computation (ABC).

The ABC method: to compare the experimental observations to the output of our model, through the construction of a relevant distance d(θ), we ran the ABC method to determine the posterior probability of each scenario, given our experimental data. First, we defined our prior distributions. We considered *a priori* our four scenarios to be equiprobable. Then, we defined the prior distributions of our model parameters. Except for ρ_C and ρ_H, all the parameters did not depend, *a priori*, on the scenario. We considered them uniformly distributed (see **Supplemental Information – B.1 and B.4** for further information on the possible range of values of the different parameters). For our parameter associated with the hypothesis of homogeneity in fate, we had: $\mathbb{P}[\rho_H = 0 \,|H = 0] = 1$ and $\rho_H| H = 1 \sim \mathcal{U}([0.3, 1])$ ($\mathcal{U}$ denotes a uniform distribution) such that we enforced a clear distinction between our scenarios of absence (H=0) or presence (H=1) of homogeneity in fate. We did similarly for ρ_C. In a Bayesian framework, the objective was to update our prior, through the integration of the data, to get the posterior distribution. Given the complexity of our model, the likelihood was intractable, hence the need to use an approximate Bayesian Computation method where the posterior distribution might be approximated. Instead of using the

expression of a likelihood, we relied on the definition of our distance d(θ) between the model
and the observations.

The ABC-rejection sampling method: For $i \in \{1, \cdots, N\}$, we sampled *a* modelling scenario (H_i,
C_i) and *a* parameter vector θ_i from the prior distributions. We ran as many simulations as
observations (1,740), computed the principal summary statistics $S_{p,i}(\theta_i)$ and then the distance
to the observation $d_i = d(\theta_i)$. We kept only the samples $i' \in I$ such that $d_{i'} < \varepsilon$, where ε
corresponded to the 0.1 percentile computed over the $(d_i)_{1 \leq i \leq N}$. That is, we kept the 0.1%
best samples. These samples were used to approximate our posterior distribution **(Extended**
Data Fig. 10a). Finally, the posterior distribution of each of our four scenarios – when using
the ABC-rejection algorithm - was approximated by:

822
$$\mathbb{P}[H = i, C = j \mid \mathcal{D}] \approx \frac{1}{0.001N} \sum_{i' \in I} \{H_{i'} = i\} \cap \{C_{i'} = j\}$$

That is, we counted how many times we selected each scenario among the 0.1% best samples.
With N=1,733,106, the results of this procedure led us to select the scenario (H=1, C=1) as the
most likely **(Supplemental Information - D.1)**. Using the ABC-rejection sampling algorithm to
explore how more complex modelling scenarios performed, and testing nearly 11 million of
different parametrisations, we found no evidence that more complex scenarios would be
more appropriate to describe our experimental observations **(Supplemental Information –**
E.1-3). In addition to the four main scenarios, we considered a minimal scenario with a single
time-to-division distribution (i.e., without distinguishing between cell types). Running
~500,000 simulations under this scenario (this number was chosen to consider a priori all five
scenarios equiprobable) has almost no impact on the previous results. Despite involving six
fewer parameters, this minimal scenario reaches a very low posterior probability (0.0152)
after ABC-rejection sampling and can therefore be discarded **(Supplementary Information -**
E.4).

The ABC-SMC method: The ABC rejection algorithm requires to sample parameter vectors a
high number of times to converge. Then, to refine our results, and ensure their robustness
and the convergence of the algorithm, we implemented the ABC-SMC sampler from Toni and
colleagues[63] (details of the implementation in **Supplemental Information - D.2**). With this
method, we considered N = 2000 parameter vectors called particles, to approximate the
posterior distribution. The ABC-SMC algorithm is sequential, so we iteratively sampled the N
particles from intermediate distributions, starting from the prior distribution to the posterior
distribution. At each iteration i, we sampled a parameter vector from the previous
intermediate distribution (sampling the previously selected particles, according to some
weights, and using pre-defined perturbation kernels) and ran the simulation to compute the
distance to the observations. As for the ABC-rejection algorithm, particles were accepted if
they are below a given threshold, discarded otherwise. In the ABC-SMC method, there was a
decreasing sequence of tolerances. The ABC-SMC method ran for many iterations, until
convergence, that was, until the distributions of the distances did not evolve for new
iterations **(Extended Data Fig. 10b)**. Over the iterations, those distributions evolved until
reaching a stable value, which corresponded to the approximated posterior distributions of

the modelling scenarios **(Extended Data Fig. 10c).** To ensure the robustness of the ABC-SMC algorithm that we used, also regarding the choice of the perturbation kernels used, we ran different configurations, leading to the same results **(Extended Data Fig. 10e)**. ABC-SMC approach refined the results from the ABC-rejection sampling approach, from iteration 9, allowing to better explore the parameter space, progressively focusing on the region where the parameter vectors were more likely to have generated the experimental data.

We focused on the discrimination between modelling scenarios, rather than the estimation of the parameters of the most likely model. Yet, we also got the intermediate posterior distributions of the model parameters and saw how each iteration allowed to be more informative on the different parameter values, and how the parameters for which the model converged faster than other ones **(Extended Data Fig. 10f** and **Supplemental Information - D.4**). Some parameters stayed undetermined (or weakly determined) at this stage, meaning that their posteriors did not move away from the prior. When comparing the summary statistics computed from the models – sampling from the posterior – to the experimental values, we observed for most of them already a good agreement **(Extended Data Fig. 10d** and **Supplemental Information - D.5**)

CITE-seq profiling of short-term liquid culture

Antibody labelling: magnetic sorted CD34$^+$ cells from umbilical cord blood were plated at a cell density of 1×10^5 cells/mL, using the same culture condition described for the condition Diff. Cells were harvested after 96 hours, counted with a haemocytometer, and resuspended in cold PBS 1X (ThermoFisher Scientific) containing 2% Bovine Serum Albumin (BSA, Euromedex) and 0.01% Tween 20 (Sigma), to a final concentration of 10^6 cells/mL. 100 µL of this cell suspension was incubated with a mix of previously prepared CITE-seq antibodies. Cells were incubated for 30 minutes at 4 °C, washed three times with the same solution used for the incubation and resuspended in 16.5 µL, at a final concentration of 1000 cells/mL.

Library sequencing: approximately 1.65×10^4 cells were loaded, to obtain a final yield of 1×10^4 recovered cells. Cells were processed accordingly to the Chromium Single Cell 3' library v3.1 (10x Genomics). Libraries were sequenced using a Novaseq s1 (Illumina), allocating at least 20,000 reads per cell for the mRNA library, and 5,000 reads for the Antibody Derived Tags (ADT) library **(Extended Data Table 8)**.

Computation methods: Raw sequencing reads were processed using Cellranger v3.1.0. To obtain a reads/cell/gene count table, reads were mapped to the human GRCh38 reference genome. Cells with less than 1000 genes per cell and with a high percentage (> 30%) of mitochondrial genes were removed from downstream analyses. Following our filtering procedures, the average UMI count per cell was 4406. The median number of genes detected per cell was 1662, 2.2% mapped to mitochondrial genes. Data normalization and integration were performed using the default Seurat v4 approach FindIntegrationAnchors() followed by IntegrateData(), and differentially expressed genes were determined using a logistic regression in Seurat on the non-integrated data using the FindConservedMarkers() function. Pathway based analyses were performed using the enrichR package. Unsupervised clustering

was performed on the significant variable genes using the 10 first PCA followed by the nonlinear dimensionality reduction technique UMAP. Annotation of the data was obtained by mapping published signatures using the AddModuleScore() method of Seurat.

Single-cell re-distribution

Single cell distribution of individual cell families was performed using the cellenONE X1 platform (Cellenion), in two independent experiments. Briefly, we followed the procedure describe in the section "Time-lapse imaging", for three tissue culture plates per experiment. Then, we randomly selected individual wells displaying at least four cells from the three 96-wells plates (10 families per plate). We harvested the content of the entire well, centrifugated for 8' at 300g and resuspended in 6 µL of PBS 1X (Sigma). Then, the whole volume was loaded into the machine, but due to the dead volume, effectively only 4 µL of this cell suspension could be redistributed. Individual cells were each sorted directly into single wells of 96-wells plates (90 cells per plate) at 4 °C, each well containing 3 µL of lysis buffer. The lysis buffer contains 0.03 µL Triton X100 (ThermoFisher Scientific), 0.0375 µL Recombinant RNAse Inhibitor (Takara Bio), 0.2 µL dNTP 10mM (ThermoFisher Scientific), 0.5 µL PEG 40% (Sigma), 0.04 µL SMARTseq3 oligo-dT 100 µM (/5BiosG/ACGAGCATCAGCAGCATACGATTTTTTTTTTTTTTTTTTTTTTTTTTTTTTVN; IDT) and ultrapure RNAse water (Invitrogen).

SMART-Seq3 library preparation and sequencing

Reverse transcription (RT) and the subsequent steps were performed accordingly to the SMART-seq3 technique[41]. Briefly, lysates in each well underwent RT (42 °C for 90 mins, 10 cycles at 50 °C 2 mins + 42 °C 2 mins, and finally 85 °C for 5 mins) on ice, in presence of template switch oligonucleotide at 100 µM (/5BiosG/AGAGACAGATTGCGCAATGNNNNNNNNrGrGrGIDT), 0.04 µL Maxima H minus reverse transcriptase (ThermoFisher Scientific), 0.1 µl TRIS-HCl 1M PH 8.3 (Invitrogen), 0.12µL NaCl 1M (Invitrogen), 0.04 µL GTP 100mM (ThermoFisher Scientific), 0,32 µL DTT 100mM (ThermoFisher Scientific), 0.2 µL MgCl2 50mM (ThermoFisher Scientific) and 0.05 µL Recombinant RNAse Inhibitor (Takara Bio) and water. The following PCR step was performed immediately after, using 5 µL Kapa Hifi HotStart Ready Mix (Roche), and forward (100 µM TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGATTGCGCAATG) and reverse (25 µM ACGAGCATCAGCAGCATACGA) primers (IDT technologies) for 25 cycles. After DNA purification using AMPure XP microbeads (ratio 0.8:1) (Beckman Coulter, A63881), the cDNA was tagmented in a final volume of 2 µL, using the Nextera XT DNA Library preparation kit (Illumina). The following library amplification was performed adding 1.5 µL of UDI primers and 2,5 µL of PCR mix including the Phusion High Fidelity DNA polymerase (ThermoFisher Scientific). After tagmentation, the different samples were pooled, purified using AMPure XP microbeads (ratio 0.9:1) and sequenced at 100-bp paired end on a high-output flow cell, using a NovaSeq SP, 800M reads (Illumina).

Single-cell 3' mRNA sequencing pre-processing

FASTQ files for each single cell were generated using bcl2fastq (version 2.20.0), with default parameters. The Read 1 contain the TSO (constant tag sequence and a UMI) followed by cDNA or only cDNA. The Read 2 contain cDNA only. Index1 and index2 reads correspond to cell-specific barcode. to generate UMI count tables from FASTQ files, the scRNA-SmartSeq3 pipeline v1.2.1 (https://doi.org/10.5281/zenodo.15056215) was used. Briefly, reads were aligned to the human genome (hg38) using STAR (version 2.7.8a) and assigned to genes using FeatureCounts (deeptools version 3.5.0). UMIs were extracted, deduplicated and counted for each gene using umiTools (version 1.1.1). Low-quality cells were filtered out based on a minimum of 500 genes detected and a percentage of mitochondrial RNA lower than 20. Families composed of only one cell were also removed.

Dimensionality reduction, clustering and pseudotime analysis

Cell fate was determined based on known marker gene expression. For each dataset the Seurat workflow is implemented by applying SCT normalization, PCA dimensionality reduction, clustering cells and running UMAP reduction [1]. The cells cluster in three groups, namely CD34+CD38-/HPC, MyP (Myeloid Progenitors) and EryP/MkP (Erythroid/Megakaryocyte Progenitors). The ELANE-expressing cells are labelled as GraP (Granulocyte Progenitors). The pseudotime analysis was performed using the Slingshot package, specifying the cluster CD34$^+$CD38$^-$/HPC as the starting cluster. For cells assigned multiple pseudotime values across different lineages, the maximum value was retained for each cell.

Cosine distance

The variable "distance" was a measure of how cells of the same family tended to be transcriptomically similar, regardless of their cell fate. The cosine distance between two cells was defined as follows[42]:

$$cosine\ distance\ (A, B) = 1 - \frac{A \cdot B}{\|A\| \|B\|}$$

where A represented the gene expression vector of the first cell and B represented the gene expression vector of the second cell. For each family, the cosine distance was computed for each pair of cells and the median was assigned. The cosine distance distributions were computed on the log-normalized UMIs for real families and both types of random families.

Identification of Variably Expressed Genes Across Families

Single-cell RNA sequencing (scRNA-seq) data from Smart-seq3 was analysed to investigate
gene expression variability across families. The pre-processed and normalized data were used
for the analysis.

Cells were grouped into families based on well information. Families with fewer than two cells
were excluded. An analysis of variance (ANOVA) was performed for each gene to assess
differential expression across families:

974
$$x_{ij} = \mu + \alpha_j + \epsilon_{ij}$$

975   where x_{ij} represents the expression of gene ii in family j, μ is the overall mean expression, α_j
976   represents family-specific effects, and ϵ_{ij} denotes residual error.

977   The total variance was decomposed into between-family variance (σ_B^2) and within-family
978   variance (σ_W^2), and an F-test was applied to determine whether the between-group variance
979   was significantly greater than the within-group variance:

980
$$F = \frac{\sigma_B^2}{\sigma_W^2}$$

where a high F-value indicates greater variance between families than within families. To
control for multiple hypothesis testing, p-values were adjusted using the Benjamini-Hochberg
procedure to limit the false discovery rate (FDR). Genes with adjusted p-values (FDR < 0.05)
were considered significantly homogeneous within families. To visualize gene expression
patterns across families and clusters, a heatmap was generated using pheatmap R package.
The expression matrix was prepared by including the significantly family conserved genes. The
samples were ordered by family information, while genes were clustered separately for cells
and genes using hierarchical clustering with Euclidean distance and the complete linkage
method. The genes were clustered into modules by cutting the hierarchical cluster tree.

MIIC analysis

MIIC was used to investigate which genes were significantly associated with specific families,
their recorded division times, colony size, predicted cell types, and transcriptomic dispersion
"distance", using transcriptomics data of 142 cells obtained via SMART-Seq3[41]. In the
reconstructed graphical networks, the variables were represented as nodes and the edges
encoded for dependency. An edge with an arrow stands for causality, an edge without an
arrow stands for association, a bi-directed dotted edge represented an unobserved common
cause between two connected variables[45]. The green arrows represented genuine causality
while the non-green arrows represented putative causality. In the case of putative causality,
it was not possible to exclude the presence of an unobserved common cause. The link-colours
between two nodes were assigned based on the Spearman's correlation coefficient: red stand
for positive correlation and blue for negative correlation. Gray was used for categorical
variables without natural order, such as the family IDs. The networks were reconstructed from
two scRNA-seq datasets, corresponding to two different experiments: 70 cells belonging to
14 families and 28,750 genes for the first, 72 cells belonging to 16 families and 30,682

sequenced genes for the second. The metadata for each dataset was the following: family of origin, times of division, colony size at 90 hours. The network was reconstructed from the 142 cells with their respective raw UMI counts of selected genes and additional features of interest. The gene selection was performed based on their high mutual information with the cellular features of interest: "family", "T1", "avg dT2", "avg dT3", "predicted.celltype", "Colony size", "exp", "distance". The genes selected to reconstruct the networks shared the highest mutual information with these properties of interest (i.e. we retained all genes above some tuneable mutual information threshold). Mitochondrial, Ribosomal Protein Large subunit (RPL), Ribosomal Protein Small subunit (RPS), Large Subunit Mitochondrial Ribosomal Proteins (MRPL) and Small Subunit Mitochondrial Ribosomal Proteins (MRPS) genes were filtered out from the analysis. A confidence parameter of 1% was used. This parameter sets the probability upper bound that this edge could be spurious due to sampling noise. Hence, the confidence parameter allowed to filter the least robust edges[44].

Two networks were obtained either including or excluding the "Family" variable:

• A network with the "family" variable and a total of 1009 variables (including 147 regulatory genes or variables and 862 consequence genes):

https://miic.curie.fr/showcase.php?id=Donada_wtfamily

• A network without the "family" variable and a total of 716 variables (including 113 regulatory genes or variables and 603 consequence genes):

https://miic.curie.fr/showcase.php?id=Donada_wofamily

Statistical analysis

Statistical analyses are detailed in figure legends and performed using R (version 4.3) or Python (version 3.10). The number of independent replicates, experiments and donors is stated for each figure in the legend.

Data availability

Raw and pre-processed single-cell RNA-sequencing dataset are deposited here:

CITE-Seq: https://zenodo.org/records/15720816

SMART-Seq3: https://zenodo.org/records/15721710 -
https://zenodo.org/records/15722093 - https://zenodo.org/records/15722300 -
https://zenodo.org/records/15722418 - https://zenodo.org/records/15722505

Additional data for reanalysis and reproducibility effort is available upon request to the lead author.

Code availability

All the scripts used in this paper are freely accessible on GitHub:

Live-cell imaging analysis: https://github.com/TeamPerie/Live-cell_imaging.git

MultiGen analysis: https://github.com/TeamPerie/MultiGen.git

Mathematical model of division and differentiation: https://gitlab-research.centralesupelec.fr/2012hermangeg/HSPC_proliferation

CITE-seq analysis: https://github.com/TeamPerie/CITEseq_analysis_pipeline.git

Smartseq3 pre-processing:

https://github.com/bioinfo-pf-curie/scRNA-SmartSeq3/releases/tag/v1.2.1

Smartseq3 quality control: https://github.com/TeamPerie/smartseq3_familly_analysis

Single-cell 3' mRNA sequencing pre-processing and analysis:

https://github.com/tizianatocci/Donada2025

https://github.com/TeamPerie/Donada-et-al2025-family-ANOVA.git

Acknowledgments

We would like to thank all the members of the Perié lab for helpful discussion, particularly Anais Lamoureux, Jason Cosgrove, Lucie Hustin. We would like to thank the members of the Institut Curie Flow Facility for their help with setting up the flow cytometry experiments. We thank the members of the Institut Curie Imaging Platform for their assistance with the live-cell imaging experiments. We thank the Saint Louis hospital cord blood biobank for providing the biological resources necessary for this project. This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 847718 and from Fondation pour la Rercherche Médicale (FRM) under the "Fin de thèse ARC FRM" programme (dossier No. ARC202506021018). The study was supported by grants from the Labex CelTisPhyBio (ANR-10-LBX-0038) (to L.P. and A.D.), the Canceropole INCA Emergence (2021-1- EMERG-54b-ICR-1, to L.P.), and the ITMO MIIC grant (21CM044, to L.P. and P.H.C.). As well as funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program ERC StG 758170-Microbar (to L.P.), A.D. was supported by a fellowship from Fondation de France (00119150 / WB-2021-36197).

Authors Contributions

A.D. and A.M. performed, analysed and interpreted the live-cell imaging data. A.D., I.M. and C.P. performed, analysed and interpreted the MultiGen assay. G.P. developed the statistical framework for the MultiGen analysis, supervised directly by K.D. and with feedback from A.D.

and L.P. G.H. developed the mathematical framework, under the supervision of P.H.C. and with feedback from A.D. and L.P. A.D., S.T.B. and D.D. performed the transcriptomics experiments. L.H.A., W.S. and T.T. analysed and interpreted the transcriptomics data, supervised directly by H.I. and L.P. with feedback from N.S. and A.D. T.T. performed and interpreted the network analysis, supervised directly by H.I. A.I., C.W., J.V., D.M., C.P. and C.L.C. provided the biological and clinical samples. A.D. prepared the paper final figures, with contributions from T.T., A.M. and W.S. A.D. and L.P. wrote the manuscript, with substantial contribution from L.L. and all the co-authors.

Declaration of interests

D.M. received research funding from Novartis, Sanofi and CSL Behring, and consulting fee from Incyte, Novartis, Sanofi, CSL Behring, Jazz pharmaceuticals and Mallinckrodt. None of this COI are directly linked to this research.

The other authors have nothing to declare.

References

1.   Till, J. E., McCulloch, E. a. & Siminovitch, L. A Stochastic Model of Stem Cell Proliferation, Based on the Growth of Spleen Colony-Forming Cells*. *Proc. Natl. Acad. Sci. U. S. A.* **51**, 29–36 (1964).

2.   Laplane, L. & Maley, C. C. The evolutionary theory of cancer: challenges and potential solutions. *Nat. Rev. Cancer* **24**, 718–733 (2024).

3.   Fennell, K. A. *et al.* Non-genetic determinants of malignant clonal fitness at single-cell resolution. *Nature* **601**, 125–131 (2022).

4.   Weinreb, C., Rodriguez-Fraticelli, A., Camargo, F. D. & Klein, A. M. Lineage tracing on transcriptional landscapes links state to fate during differentiation. *Science (80-.).* **367**, eaaw3381 (2020).

5.   Marsolier, J. *et al.* H3K27me3 conditions chemotolerance in triple-negative breast cancer. *Nat. Genet.* **54**, 459–468 (2022).

6.   Reyes Hueros, R. A., Gier, R. A. & Shaffer, S. M. Non-genetic differences underlie variability in proliferation among esophageal epithelial clones. *PLOS Comput. Biol.* **20**, e1012360 (2024).

7.   Shaffer, S. M. *et al.* Rare cell variability and drug-induced reprogramming as a mode of cancer drug resistance. *Nature* **546**, 431–435 (2017).

8.   Sigal, A. *et al.* Variability and memory of protein levels in human cells. *Nature* **444**, 643–646 (2006).

9.   Shaffer, S. M. *et al.* Memory Sequencing Reveals Heritable Single-Cell Gene Expression Programs Associated with Distinct Cellular Behaviors. *Cell* **182**, 947-

959.e17 (2020).

10. Sieburg, H. B. *et al.* The hematopoietic stem compartment consists of a limited number of discrete stem cell subsets. *Blood* **107**, 2311–2316 (2006).

11. Cosgrove, J., Hustin, L. S. P., de Boer, R. J. & Perié, L. Hematopoiesis in numbers. *Trends in Immunology* **42**, 1100–1112 (2021).

12. Laurenti, E. & Göttgens, B. From haematopoietic stem cells to complex differentiation landscapes. *Nature* **553**, 418–426 (2018).

13. Dykstra, B. *et al.* Long-Term Propagation of Distinct Hematopoietic Differentiation Programs In Vivo. *Cell Stem Cell* **1**, 218–229 (2007).

14. Tak, T. *et al.* Hspcs display within-family homogeneity in differentiation and proliferation despite population heterogeneity. *Elife* **10**, (2021).

15. Kueh, H. Y. *et al.* Positive feedback between PU.1 and the cell cycle controls myeloid differentiation. *Science* **341**, 670–3 (2013).

16. Treichel, S. & Filippi, M. D. Linking cell cycle to hematopoietic stem cell fate decisions. *Frontiers in Cell and Developmental Biology* **11**, (2023).

17. Donada, A. *et al.* Simultaneous Assessment of Kinship, Division Number, and Phenotype via Flow Cytometry for Hematopoietic Stem and Progenitor Cells. *J. Vis. Exp.* 1–26 (2023). doi:10.3791/64918

18. Horton, M. B. *et al.* Multiplexed Division Tracking Dyes for Proliferation-Based Clonal Lineage Tracing. *J. Immunol.* **201**, 1097–1103 (2018).

19. Marchingo, J. M. *et al.* T-cell stimuli independently sum to regulate an inherited clonal division fate. *Nat. Commun.* **7**, 13540 (2016).

20. Spencer Chapman, M. *et al.* Clonal dynamics after allogeneic haematopoietic cell transplantation. *Nature* (2024). doi:10.1038/s41586-024-08128-y

21. Oshima, M. U. *et al.* Characterization of clonal dynamics using duplex sequencing in donor-recipient pairs decades after hematopoietic cell transplantation. *Sci. Transl. Med.* **16**, eado5108 (2024).

22. Aldoss, I., Clark, M., Marcucci, G. & Forman, S. J. Donor derived leukemia in allogeneic transplantation. *Leuk. Lymphoma* **62**, 2823–2830 (2021).

23. Calabria, A. *et al.* Long-term lineage commitment in haematopoietic stem cell gene therapy. *Nature* **636**, 162–171 (2024).

24. Oliveira, E. A. *et al.* Epigenetic Heritability of Cell Plasticity Drives Cancer Drug Resistance through a One-to-Many Genotype-to-Phenotype Paradigm. *Cancer Res.* **85**, 2921–2938 (2025).

25. Bell, C. C. *et al.* Targeting enhancer switching overcomes non-genetic drug resistance in acute myeloid leukaemia. *Nat. Commun.* **10**, 1–15 (2019).

26. Laurenti, E. *et al.* CDK6 Levels Regulate Quiescence Exit in Human Hematopoietic Stem Cells. *Cell Stem Cell* **16**, 302–313 (2015).

27. Aiuti, A. *et al.* Lentiviral Hematopoietic Stem Cell Gene Therapy in Patients with
Wiskott-Aldrich Syndrome. *Science (80-.).* **341**, 1233151 (2013).

28. Knapp, D. J. H. F. *et al.* Dissociation of Survival, Proliferation, and State Control in
Human Hematopoietic Stem Cells. *Stem Cell Reports* **8**, 152–162 (2017).

29. Passegué, E., Wagers, A. J., Giuriato, S., Anderson, W. C. & Weissman, I. L. Global
analysis of proliferation and cell cycle gene expression in the regulation of
hematopoietic stem and progenitor cell fates. *J. Exp. Med.* **202**, 1599–1611 (2005).

30. Johnson, C., Belluschi, S. & Laurenti, E. Beyond "to divide or not to divide": Kinetics
matters in hematopoietic stem cells. *Experimental Hematology* **92**, 1-10.e2 (2020).

31. Hammond, C. A., Wu, S. W., Wang, F., MacAldaz, M. E. & Eaves, C. J. Aging alters the
cell cycle control and mitogenic signaling responses of human hematopoietic stem
cells. *Blood* **141**, 1990–2002 (2023).

32. Gutch, S. *et al.* Protocol for enrichment and functional analysis of human
hematopoietic cells from umbilical cord blood. *STAR Protoc.* **5**, 103024 (2024).

33. Majeti, R., Park, C. Y. & Weissman, I. L. Identification of a hierarchy of multipotent
hematopoietic progenitors in human cord blood. *Cell Stem Cell* **1**, 635–45 (2007).

34. Grelaud, A., Roberty, C. P., Marinz, J. M., Rodolphe, F. & Talys, J. F. ABC likelihood-
free methods for model choice in Gibbs random fields. *Bayesian Anal.* **4**, 317–335
(2009).

35. Toni, T. & Stumpf, M. P. H. Simulation-based model selection for dynamical systems
in systems and population biology. *Bioinformatics* **26**, 104–110 (2009).

36. Sunnåker, M. *et al.* Approximate Bayesian Computation. *PLOS Comput. Biol.* **9**,
e1002803 (2013).

37. Dowling, M. R., Milutinović, D. & Hodgkin, P. D. Modelling cell lifespan and
proliferation: is likelihood to die or to divide independent of age? *J. R. Soc. Interface*
2, 517–526 (2005).

38. Hawkins, E. D., Turner, M. L., Dowling, M. R., Van Gend, C. & Hodgkin, P. D. A model
of immune regulation as a consequence of randomized lymphocyte division and
death times. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 5032–5037 (2007).

39. Upadhaya, S. *et al.* Kinetics of adult hematopoietic stem cell differentiation in vivo. *J.
Exp. Med.* **215**, 2815–2832 (2018).

40. Chen, T. N., Gupta, A., Zalavadia, M. D. & Streets, A. µCB-seq: microfluidic cell
barcoding and sequencing for high-resolution imaging and sequencing of single cells.
Lab Chip **20**, 3899–3913 (2020).

41. Hagemann-Jensen, M. *et al.* Single-cell RNA counting at allele and isoform resolution
using Smart-seq3. *Nat. Biotechnol.* **38**, 708–714 (2020).

42. Smets, T. *et al.* Evaluation of Distance Metrics and Spatial Autocorrelation in Uniform
Manifold Approximation and Projection Applied to Mass Spectrometry Imaging Data.
Anal. Chem. **91**, 5706–5714 (2019).

43. Mold, J. E. *et al.* Clonally heritable gene expression imparts a layer of diversity within
cell types. *Cell Syst.* **15**, 149-165.e10 (2024).

44. Verny, L., Sella, N., Affeldt, S., Singh, P. P. & Isambert, H. Learning causal networks
with latent variables from multivariate information in genomic data. *PLoS Comput.*
Biol. **13**, 1–25 (2017).

45. Ribeiro-Dantas, M. da C. *et al.* Learning interpretable causal networks from very large
datasets, application to 400,000 medical records of breast cancer patients. *iScience*
27, (2024).

46. Cabeli, V. *et al.* Learning clinical networks from medical records based on information
estimates in mixed-type data. *PLOS Comput. Biol.* **16**, e1007866 (2020).

47. Carey, A. *et al.* Identification of Interleukin-1 by Functional Screening as a Key
Mediator of Cellular Expansion and Disease Progression in Acute Myeloid Leukemia.
Cell Rep. **18**, 3204–3218 (2017).

48. Filippakopoulos, P. *et al.* Selective inhibition of BET bromodomains. *Nature* **468**,
1067–1073 (2010).

49. Lechman, E. R. *et al.* MiR-126 Regulates Distinct Self-Renewal Outcomes in Normal
and Malignant Hematopoietic Stem Cells. *Cancer Cell* **29**, 214–228 (2016).

50. Schoof, E. M. *et al.* Quantitative single-cell proteomics as a tool to characterize
cellular hierarchies. *Nat. Commun.* **12**, 1–15 (2021).

51. Horton, M. B. *et al.* Lineage tracing reveals B cell antibody class switching is
stochastic, cell-autonomous, and tuneable. *Immunity* **55**, 1843-1855.e6 (2022).

52. Müller-Sieburg, C. E., Cho, R. H., Thoman, M., Adkins, B. & Sieburg, H. B. Deterministic
regulation of hematopoietic stem cell self-renewal and differentiation. *Blood* **100**,
1302–1309 (2002).

53. Sieburg, H. B., Rezner, B. D. & Muller-Sieburg, C. E. Predicting clonal self-renewal and
extinction of hematopoietic stem cells. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 4370–4375
(2011).

54. Valdés-Mas, R. *et al.* Transplacental transfer of essential thrombocythemia in
monozygotic twins. *Blood* **128**, 1894–1896 (2016).

55. Sousos, N. *et al.* In utero origin of myelofibrosis presenting in adult monozygotic
twins. *Nat. Med.* **28**, 1207–1211 (2022).

56. Singh, I., Fernandez-Perez, D., Sanchez, P. S. & Rodriguez-Fraticelli, A. E. Pre-existing
stem cell heterogeneity dictates clonal responses to the acquisition of leukemic driver
mutations. *Cell Stem Cell* **32**, 564-580.e6 (2025).

57. Schönberger, K. *et al.* Multilayer omics analysis reveals a non-classical retinoic acid
signaling axis that regulates hematopoietic stem cell identity. *Cell Stem Cell* **29**, 131-
148.e10 (2022).

58. Bonora, M. *et al.* A mitochondrial NADPH-cholesterol axis regulates extracellular
vesicle biogenesis to support hematopoietic stem cell fate. *Cell Stem Cell* **31**, 359-

377.e10 (2024).

59.   Duffy, K. R. *et al.* Activation-induced B cell fates are selected by intracellular
1232   stochastic competition. *Science (80-.).* **335**, 338–341 (2012).

1233   60.   Loeffler, D. *et al.* Asymmetric organelle inheritance predicts human blood stem cell
1234   fate. *Blood* **139**, 2011–2023 (2022).

1235   61.   Eisele, A. S., Tarbier, M., Dormann, A. A., Pelechano, V. & Suter, D. M. Gene-
1236   expression memory-based prediction of cell lineages from scRNA-seq datasets. *Nat.*
1237   *Commun.* **15**, 1–16 (2024).

1238   62.   Davison, A. C. & Hinkley, D. V. Bootstrap Methods and their Application. *Bootstrap*
1239   *Methods their Appl.* (1997). doi:10.1017/CBO9780511802843

1240   63.   Toni, T., Welch, D., Strelkowa, N., Ipsen, A. & Stumpf, M. P. H. Approximate Bayesian
1241   computation scheme for parameter inference and model selection in dynamical
1242   systems. *J. R. Soc. Interface* **6**, 187–202 (2009).

1243