1   **Oscillatory markers of interoceptive attention: beta suppression as a neural**

2   **signature of heartbeat processing**

3   Kristina Pultsina[1], Tiina Parviainen[1,2], Suvi Karjalainen[1,2]

1 - Centre for Interdisciplinary Brain Research, University of Jyväskylä, PO Box 35,

Jyväskylä FI-40014, Finland

2 - Department of Psychology, University of Jyväskylä, PO Box 35, Jyväskylä FI-40014,

Finland

Author Note

We have no known conflict of interest to disclose.

Correspondence concerning this article should be addressed to Kristina I. Pultsina,

Centre for Interdisciplinary Brain Research, University of Jyväskylä, PO Box 35,

Jyväskylä FI-40014, Finland

Email: kpultsin@jyu.fi

Abstract

Interoceptive attention—the ability to selectively focus on internal bodily signals—has been

linked to distinct neural responses, yet the contribution of oscillatory dynamics to this process

remains underexplored. This study investigates the neural mechanisms underlying

interoceptive attention by examining beta-band power suppression during heartbeat and

auditory discrimination tasks. Fifty-one healthy participants engaged in interoceptive

(heartbeat detection) and exteroceptive (auditory discrimination) tasks while their brain activity

was measured using magnetoencephalography (MEG). The results revealed significant beta

suppression time-locked to the R-peak in the somatosensory cortex, anterior cingulate cortex,

mid-cingulate cortex, and dorsolateral prefrontal cortex from 310 to 530 ms post-R-peak. Beta

suppression was more pronounced during interoceptive attention, correlating positively with

interoceptive accuracy. The findings support the notion that beta suppression in fronto-cingulo-

somatosensory network may serve as a neural marker of interoceptive processing, contributing

to predictive coding models of interoception. This study highlights the potential for using beta

suppression as an objective measure of interoceptive accuracy and suggests that neural

oscillations play a critical role in the brain's regulation of heartbeat-related information.

Furthermore, the study proposes that interoceptive attention involves a top-down mechanism

that dynamically adjusts the brain's response to cardiac afferent signals, enhancing the

precision of interoceptive processing. These findings have implications for understanding how

the brain integrates interoceptive signals and may provide insights into clinical applications

targeting interoceptive dysfunctions.

Keywords: Interoceptive attention, interoception, heartbeat processing, neural oscillations,

beta power

Introduction

Interoception refers to the perception of sensory information originating from the viscera and internal milieu. The ability to attend to and consciously process internal bodily signals is considered crucial for fundamental aspects of selfhood and emotion regulation, as well as for optimal cognitive functioning from basic goal-directed behavior to higher-order executive tasks. A broader definition of interoception involves not only low-level processing of individual visceral signals, such as changes in the biochemical composition of blood, but also the integration of information about physiological state, which in turn can influence cognitive processes such as attention and memory, as well as emotions (1–3).

Cardiac interoceptive sensitivity, specifically the ability to discriminate one's own heartbeats, serves as a paradigmatic example of this phenomenon. Elucidating the physiological pathways underlying cardiac interoceptive sensitivity is crucial for comprehending the complex interplay between cardiac and neural systems, and its role in individual differences in experience. A potential source of neural responses to heartbeats is the activation of mechanoreceptors in the heart wall, aortic arch, and carotid sinus, along with cutaneous somatosensory mapping and neurovascular coupling in the cortex (4,5). This information travels via the nucleus of the solitary tract and then mainly via thalamus, functionally connected to forebrain regions, including amygdala, insular cortex, anterior and mid cingulate, and orbitomedial prefrontal cortex (2).

Neuroimaging studies have implicated the insular cortex and anterior cingulate cortex (ACC) as pivotal structures in the representation of interoceptive states, including the modulation of cardiac activity (6). The ACC's involvement in both central autonomic control and peripheral sympathetic responses across various contexts suggests its role in integrating visceral states with behavioral output (1,7).While the insula is predominantly associated with generating emotional arousal and the cortical input of interoceptive signals into the brain, the

anterior cingulate cortex, known as the 'limbic motor cortex' is crucially involved in generating

autonomic responses to emotionally salient stimuli (6,8,9). Notably, different studies have

shown increased activity in these brain structures when participants were asked to attend to

internal bodily responses, for example, to their heartbeats in Heartbeat Detection task (10–12).

Recent research suggests the insula and ACC are essential for interoceptive awareness,

but a study by Khalsa challenges this view (13). Using isoproterenol to induce heartbeat

sensations, they found that a patient with bilateral insula and ACC damage could still perceive

his heartbeat, like healthy controls. However, when his skin sensations were blocked by a

topical anesthetic, his interoceptive awareness disappeared. These results suggest that both

visceral and somatosensory pathways contribute to interoception, rather than the insula alone.

This is consistent with neuroimaging findings showing that activation of the somatosensory

cortex correlates with interoceptive awareness in tasks requiring focused attention on the

heartbeats (14,15).

The cumulative evidence supports the existence of multiple, semi-independent

pathways subserving cardiac interoceptive awareness and suggests a complex interplay

between somatosensory processing and interoceptive phenomena. This is consistent with the

idea of neurovisceral integration that interoception is part of a broad multisensory integration

used for self-regulation and adaptability of the organism (16,17). The precise neural

mechanisms involved in integrating interoceptive signals remain largely unexplored.

According to the influential predictive coding theory, top-down predictions about the body's

state are thought to be continuously compared with incoming bottom-up sensory information

(18).

The most common marker of heart–brain communication is heart beat evoked potential

(HEP), which reflects the brain's processing of cardiac signals. Even though evoked potential

predominantly reflects bottom-up processing of sensory signals, in this case, the brain

processes heartbeat signals through baroreceptor activity (4,19). There is also evidence that

attention focus and affective predictions about external stimuli can influence HEP amplitude

(12,20,21).

There is growing evidence that bodily sensations have been linked to modulations of

induced activity. For example, breathing cycles modulate alpha rhythms in the brain, both at

rest and during both spontaneous and deep breathing (22–24). Similarly, arousal has been

shown to modulate the cortical processing of heartbeats, with higher arousal states amplifying

HEPs. Notably, alpha oscillations influence individual variability in interoceptive sensitivity,

suggesting a potential gating mechanism for cardiac awareness. This highlights the crucial role

of oscillatory activity in shaping the brain's dynamic integration of bodily signals (25).

Moreover, recent evidence highlights the role of cortical oscillatory dynamics in cardiac

interoception. Zaccaro et al. demonstrated that heartbeat-related alpha and theta band

oscillations, particularly during exhalation, are associated with increased interoceptive

accuracy during heartbeat counting tasks, with these oscillations – reflected in power, inter-

trial coherence, and functional connectivity – being most prominent over right fronto-centro-

parietal regions and independent of peripheral cardiac physiology (24). Importantly, alpha-

band modulations showed a significant correlation with heartbeat-evoked potentials,

suggesting that oscillatory activity may support the top-down attentional allocation to cardiac

signals by enhancing the precision-weighting of prediction errors.

The study by Crivelli & Balconi found that interoceptive attention during the Heartbeat

Counting Task (HCT) was associated with increased alpha oscillatory activity in the right

parahippocampal gyrus, suggesting that alpha rhythms play a role in top-down attentional

control, helping to suppress irrelevant external information while focusing on internal bodily

signals (26). Additionally, beta oscillations in the cingulate cortex were positively correlated

with interoceptive accuracy, indicating that beta activity reflects active processing of

interoceptive signals rather than just passive bodily awareness. Notably, these neural markers

were not linked to heart rate variability, reinforcing the idea that interoceptive attention is

guided by cognitive control rather than direct cardiac input. These findings support predictive

coding models of interoception, highlighting oscillatory induced activity as a potential

objective marker for interoceptive accuracy and regulation.

Thus, the analysis of induced oscillatory activity may provide a better understanding of

the mechanisms of predictive processing, adaptation, and integration of interoceptive

information over time. Specifically, we focused on oscillatory dynamics in the beta frequency

bands, hypothesizing that directing attention to heartbeats would modulate these rhythms – in

particular, that it would produce R-locked power changes as a neural signature of interoceptive

attention in the heartbeat and auditory discrimination task. The primary focus is on beta-band

power, given its critical role in the top-down control of predictions, with a decrease in beta

power associated with updating predictions and processing prediction errors (27).

Results

To assess the presence of R-peak-related changes in beta band, we performed a one-sample

cluster t-test on all vertices for the collapsed exteroceptive and interoceptive tasks.

A significant TFCE cluster was identified between 310 ms and 530 ms post-R-peak

with a pronounced suppression at beta frequency band in anterior (ACC) and middle cingulate

cortex (MCC), somatosensory and premotor cortex (S & PMC), as well as the dorsolateral

prefrontal cortex (dlPFC) (Fig.2).

Further refinement of the preselected ROI boundaries (see Methods) identified

statistically significant vertices in all regions except for the insula and frontal operculum cortex,

which were excluded from further analysis (Fig. 2A). The time-frequency analysis (Fig.2 B,C)

showed that besides modulation at beta band, also alpha band (8 - 14 Hz) suppression was

related to heartbeats. We performed the same analysis to check for significant alpha suppression but found none.

Fig.2. R-peak related β suppression in all conditions collapsed. **A** - Source distribution of β power in the TFCE defined cluster time interval 0.310 to 0.530 s (only significant vertices shown) for all tasks combined; ROIs with newly defined borders thresholded by significant vertices; Time course of β power changes across all tasks and ROIs. **B** - Time-frequency distribution across all pooled ROIs. The dashed line indicates the area that differs significantly from the baseline (one-sample t-test, pFDR<0.05). **C** - Phase-locked distribution across all pooled ROIs. **D** – Beta power changes following the R peak, averaging over 100-ms intervals. Only vertices that significantly differ from the baseline are shown (one-sample t-test, pFDR < 0.05).

Beta power during interoception and exteroception tasks

After identifying the main ROIs with significant suppression, we performed a comparison of

oscillatory activity at beta band between the exteroceptive and interoceptive tasks using the

average power in the time window of 0.310 to 0.530 s after the R-peak.

We found significant differences between the interoception and exteroception tasks for

all the selected ROIs (MCC: $t(50) = 3.967$, $p<0.001$, $d = 0.556$; S & PMC: $t(50) =2.933$,

$p=0.005$, $d = 0.411$; dlPFC: $t(50) =3.872$ $p<0.001$, $d = 0.542$; ACC: $t(50) =3.857$, $p<0.001$, $d =$

0.540). During interoception, participants demonstrated more prominent suppression in the

somatosensory cortex and premotor cortex, dlPFC, ACC, and midcingulate cortex (Fig.3A).

Also, to validate that the observed difference in suppression was not whole-brain related, we

performed the same comparison between the tasks on the whole-brain averaged data and didn't

reveal any significant differences ($t (50) =-0.517$, $p=0.608$, $d = 0.072$).

To rule out the possibility that the observed differences between tasks were due to

difference in duration in heartbeat intervals, we compared the heart interbeat intervals (IBI)

during interoception and exteroception task (Fig.1B). The t-test did not show significant

difference between exteroception and interoception task ($t (50) = - 0.458$, $p=0.649$, $d =0.064$)

in IBI. The average IBI for the exteroception task was 0.926 s ± 0.018 and 0.930 s ± 0.018 for

the interoception task.

Fig.3. Beta power change in the ROIs averaged in 0.310 to 0.530 s after R-peak onset. **A** -

Boxplots comparing mean beta power in exteroceptive and interoceptive attention tasks (**p

≤ 0.01, ***p ≤ 0.001). **B** - Correlation between beta power during interoceptive attention task

and the interoception accuracy.

Correlation between interoception accuracy and beta suppression during the interoception

task

To test whether beta suppression in ROIs related to interoceptive accuracy, we performed a

correlation analysis. A significant correlation between accuracy and beta power suppression

was found for the S & PMC (r_s (49) = -0.43, p=0.002) and ACC (r_s (49) = -0.35, p = 0.013)

ROIs. Higher accuracy was related to greater suppression in these brain areas. The same trend

was observed in the anterior and mid-cingulate cortex; however, the correlation values did not

reach statistical significance.

Beta suppression during resting state condition

We further tested if the observed suppression is specifically related to the tasks requiring

attention to exteroceptive or interoceptive signals, or whether the task instead modulates R-

peak suppression that is already present at rest?

To investigate this, we used resting state data. We averaged beta power within the

regions of interest (ROIs) and the time interval that exhibited significant suppression during

both the interoceptive and exteroceptive tasks. We then compared these values to the baseline

using a one-sample t-test.

The results revealed significant beta suppression in somatosensory and premotor ROI

$(t (50) = -3.058, p=0.004, d=0.428)$, MCC ROI $(t (50) = -2.982, p=0.004, d=0.418)$ and dlPFC

ROI $(t (50) = -3.068, p=0.003, d=0.430)$. However, in the ACC, beta suppression did not reach

statistical significance $(t (50) = -1.947, p=0.057, d=0.273)$ (S1 Fig).

The comparison of resting-state beta suppression and during interoception and

exteroception tasks was analyzed using a rmANOVA. The results showed a significant effect

of condition on beta power in the S & PMC $(F (2,50)=14.76, p<0.001, \eta p2= 0.228)$, MCC $(F$

$(2,50) =21.39, p<0.001, \eta p2= 0.3)$ and dlPFC $(F (2,50)=25.18, p<0.001, \eta p2 =0.335)$. Post-hoc

comparisons revealed the lowest beta activation in the resting state (S & PMC: Rest vs

Exteroception p=0.011, Rest vs Interoception p<0.001; MCC: Rest vs Exteroception p=0.021,

Rest vs Interoception p<0.001; dlPFC Rest vs Exteroception p<0.001, Rest vs Interoception

p<0.001) (S2 Fig.).

Temporal Specificity of R-peak related beta suppression

To determine whether the observed post-R-peak beta-band suppression is time-locked to

cardiac events – reflecting neural responses to afferent signals following heart contraction—

rather than being related to motor activity or spontaneous neural fluctuations, we generated

surrogate epochs for each participant (41,42). This approach allowed us to test whether the beta

suppression was specifically aligned with actual cardiac events rather than with efferent

processes preceding the heartbeat or random fluctuations. However, when using the surrogate

R-peak epochs, this post-R-peak beta suppression vanished in all ROIs, demonstrating that the

effect requires precise alignment with true heartbeats (S3 Fig). In other words, randomly

shifting the timing of heartbeats negated the beta suppression effect, yielding no significant

post-trigger power drop in the beta band.

We further examined whether the timing or magnitude of post-R-peak beta suppression

depended on the length of the cardiac cycle, which would be expected if the brain were

anticipating the next heartbeat. Heartbeat events were divided into two groups based on the

interbeat interval (short vs. long, via a median split of IBI durations). We then compared the

post-R-peak beta activity between these groups. The results showed no significant difference

in the onset, duration, or amplitude of beta suppression between short-IBI and long-IBI

heartbeats – the suppression followed the preceding R-peak on a consistent timeline in both

cases (S4 Fig). This temporal invariance directly contradicts an anticipatory explanation. If the

beta suppression were an anticipatory response to the upcoming beat, we would expect it to

adjust based on the expected timing of that beat (for example, occurring earlier or ending sooner

when the next heartbeat is imminent). Instead, the beta suppression appears to be locked to the

previous heartbeat's afferent signals and does not shift in accordance with the varying interval

to the next beat. Such consistency indicates that the suppression is driven by the physiological

consequences of the preceding cardiac cycle (e.g. baroreceptor input from that heartbeat) rather

than by a countdown or prediction of the next heartbeat's timing.

Discussion

Our study, using a heartbeat and an auditory discrimination task, provides evidence that

paying attention to internal bodily signals – in this case, one's heartbeat – engages distinct

neural dynamics characterized by suppression of beta-band oscillatory activity in the premotor

and somatosensory cortex, anterior and medial cingulate cortex, and dorsolateral prefrontal

cortex. In other words, when participants focused on feeling their heartbeat, the beta rhythms

(16 – 29 Hz) significantly decreased relative to when they attended to external sounds.

Intriguingly, the depth of beta suppression positively correlated with interoceptive accuracy.

This finding supports existing evidence that the focus of attention modulates

interoceptive signal processing (11,12). Compared to previous studies that primarily examined

phase-locked heartbeat-evoked responses or focused on alpha- and theta-band modulations

during interoceptive attention (12,24,43), our findings extend this body of work by

demonstrating that dynamically shifting attention focus - specifically when integrating internal

cardiac signals with external auditory information – engages distinct oscillatory mechanisms,

particularly beta-band suppression. This interpretation is consistent with previous evidence

linking beta suppression with integrating sensory input with existing priors and the

maintenance of ongoing cognitive processes (44–46).

While several studies have reported alpha-band changes during interoceptive attention,

particularly in heartbeat counting tasks that require sustained internal focus (24,26,47), our

findings instead revealed prominent beta-band suppression. The observed differences in results

can be attributed to several factors. Primarily, they stem from variations in the experimental

task: participants engaged in a heartbeat counting task, which relies on internally directed, top-

down attention and the inhibition of brain areas not directly involved in the task (24,26). This

form of attention has been associated with increased alpha-band activity, reflecting the

suppression of external input. The alpha synchronization observed in visual and parietal regions

during interoceptive attention, relative to attention to external stimuli, likely reflects a neural

gating mechanism that filters out task-irrelevant information (47,48).

By contrast, our heartbeat and an auditory discrimination task required splitting

attention between internal signals and external tones, reflect a shift away from a stable, "status

quo" state toward dynamic processing (27). The R-peak locked beta suppression we observed

reflects this active engagement, where each heartbeat-tone pair demands evaluation, eliciting

transient beta desynchronization. Within the predictive coding framework, heartbeat-related

beta suppression may reflect a comparison between priors about interoceptive signals and

incoming sensations (sound). For instance, hearing one's own heartbeat evokes a smaller

auditory N1 response than hearing external sounds, reflecting predictive attenuation of

expected input (49). In this context, beta suppression may serve as a neural mechanism for

minimizing the impact of predictable interoceptive stimuli, consistent with classical sensory

gating and predictive coding accounts. Notably, the fact that suppression was independent of

cardiac cycle length further supports its top-down, cognitive nature, rather than a purely

physiological reflex.

Finally, an important piece of evidence linking neural mechanisms to function is that

the strength of heartbeat-related beta suppression in ACC, somatosensory, and premotor areas

correlated with interoceptive accuracy. This supports the interpretation that beta suppression

indexes interoceptive signal processing. Prior studies have similarly linked interoceptive

accuracy to oscillatory brain activity (24,50). For example, prior studies have shown that

individuals with higher heartbeat detection accuracy tend to exhibit larger HEP amplitudes,

and that beta oscillation changes in the cingulate gyrus during heartbeat counting tasks are

significantly associated with interoceptive performance (26,50). Together, these findings

support the notion that beta suppression may serve as a candidate neural marker of

interoceptive sensitivity, reflecting the intentional engagement of attention toward subtle

internal sensations rather than basic, low-level sources of information, such as heartbeat-related

sensations; however, this interpretation requires further validation.

The spatial distribution of beta suppression – especially in the anterior cingulate cortex,

somatosensory cortex, and prefrontal areas – partially aligns with earlier fMRI data on brain

structures engaged during interoceptive attention (11,15). One notable divergence from prior

neuroimaging studies is the absence of significant heartbeat-related effects in the insular cortex.

This may reflect the insula's role in sustained or integrative interoceptive processes that are not

captured by time-locked analyses. fMRI, for instance, can detect overall increases in insular

activity during interoceptive tasks, but lacks the temporal resolution to resolve rapid, R-peak-

locked oscillatory changes. It is conceivable that insular activation reflects a meta-

representation of bodily state or subjective feeling of the heartbeat, rather than transient

responses to individual cardiac afferent bursts (6,51). Second, technical factors could have

limited our ability to detect insular signals. Additionally, the insula's deep and folded anatomy

may limit MEG/EEG sensitivity, further contributing to the absence of detectable effects (52).

Therefore, our findings do not preclude insular involvement but highlight the need for

complementary imaging approaches to fully capture its contribution to interoception.

Importantly, the cortical sources we identified for heartbeat-related beta suppression

are consistent with the proposed "second pathway" of cardiac interoception that involves

somatosensory processing (4,13). Our results resonate with the case of Khalsa's patient and

other evidence suggesting that cardioceptive signals can be routed through somatosensory

cortices (13). Observed activation in the somatosensory and motor cortex, MCC, and ACC may

be related not only to the direct monitoring of heartbeats through baroreceptors – connected to

the nucleus tractus solitarius (NTS) – but also to peripheral changes associated with heartbeats

(2). Thus, our findings reinforce the view that interoceptive awareness of heartbeats is the

product of an integrated viscerosensory and somatosensory process.

Recent neurophysiological studies provide compelling evidence for a bidirectional

interplay between central neural oscillations and peripheral autonomic activity, particularly

involving the dlPFC, ACC and somatosensory cortex (53,54). These findings suggest a

functional loop wherein cortical regions modulate sympathetic outflow, and peripheral

autonomic signals, in turn, influence central processing.

Specifically, during heartbeats, spontaneous bursts of muscle sympathetic nerve

activity (MSNA) – which regulate arteriolar diameter in skeletal muscles and contribute to

beat-to-beat blood pressure control – exhibit strong temporal coupling within the dlPFC and

MCC activation, as measured using concurrent microneurography and fMRI (54). This

interaction suggests a potential top-down modulation of sympathetic output to skeletal muscle

vasculature. Furthermore, transcranial alternating current stimulation of the dlPFC in awake

humans has been shown to partially entrain MSNA, heart rate, and blood pressure,

underscoring the dlPFC's role in autonomic regulation and cardiovascular control (53).

Complementing these findings, research by Riaz et al. has demonstrated a correlation

between MSNA inhibition and beta-band rebound activity in the ACC and somatosensory

cortex (55). This suggests that sensorimotor processing contributes to the modulation of

sympathetic responses, possibly through the integration of afferent somatosensory input with

autonomic regulation.

The observed synchronization in response to MSNA suppression supports the notion

that beta suppression may be related to the brain's engagement in processing sensory input

about cardiac sensations, which are generated by a cascade of peripheral reactions. This

engagement in afferent information processing may be particularly pronounced during

interoceptive attention – a cognitive state known to modulate cortical oscillations, particularly

in the beta frequency band, thereby enhancing the integration of internal bodily signals with

autonomic regulation (56).

The convergence of these results strengthens the idea that oscillatory responses tied to

the cardiac cycle carry behavioral significance. In our study, the fact that the beta suppression

tracks interoceptive accuracy implies that this neural measure indexes the efficacy of

interoceptive signal processing. Our findings suggests beta suppression reflects individual

variability in interoceptive processing, potentially serving as a candidate neural marker for

further investigation. Those with a more pronounced suppression appear to have more precise

or sensitive internal monitoring, resulting in higher interoceptive performance. This insight has

potential implications for using neural oscillatory markers to assess or even train interoceptive

abilities in the future.

Materials and methods

Participants

51 volunteers participated in the experiment (21 men and 30 women), aged 33 ± 6.7 years

($M \pm SD$) with $19 - 62$ range. Exclusion criteria included a history of cardiovascular or

respiratory disease, head trauma, intellectual disability, neurological or psychiatric condition,

use of medication affecting the nervous system, and claustrophobia. Additionally, participants

included in this study had normal or corrected-to-normal hearing and vision.

Ethical approval was received from the local Ethics Committee of the University of

Jyväskylä and all the study procedures were conducted in accordance with the Declaration of

Helsinki. Prior to any study procedures, participants were informed about the study, and a

written informed consent was obtained from all participants.

Procedure

Participants performed modified heartbeat and an auditory discrimination task (HDT) (11,28).

The task included randomized trials that required participants to focus attention on their own

heartbeat (interoceptive attention task) or a sound stimulus (exteroceptive attention task). Both

task conditions were based on monitoring a sequence of sounds that were played either

simultaneously or with a delay with respect to individual's own heartbeat. Additionally, in all

the task trials the pitch of the sound was constant (800 Hz, 100 ms).

During each trial, a visual cue was displayed on the screen to indicate the task condition.

A heart icon signaled the interoceptive attention task, instructing participants to focus on

whether in given trial the sounds were played simultaneously with their heartbeat. After the

sound sequence ended (20 heart beats), they responded by pressing the yellow button if the

sound was subjectively felt to be in sync with their heartbeat and the blue button if it was out

of sync (asynchronous).

A musical note icon indicated the exteroceptive attention task, requiring participants to

focus on the pitch of the sound. After the sequence ended, they pressed the yellow button if

they detected a pitch deviation and the blue button if the pitch remained unchanged (Fig. 1A).

Importantly, auditory sounds were present during both types of attention blocks to

prevent any differences related to sound response between tasks. The auditory stimulus was

triggered by the onset of the participant's heartbeat either simultaneously (synchronized

condition: 200 ms delay after R-peak), or with a delay (asynchronized condition: 500 ms delay

after R-peak). The 200 ms delay for the simultaneous condition was selected based on

subjective assessments, as it corresponded to the highest synchronicity judgments, while the

500 ms delay corresponded to the lowest (29,30).This delay also corresponds to the time

interval between the R-wave peak and the base of the leading edge of the finger pulse wave

(31). In both tasks, participants completed 6 blocks each containing 20 heartbeats.

Fig.1. Schematic overview of the block design in the heartbeat detection and an auditory

discrimination task. **A** The task alternated between two types of attention blocks: the

interoceptive condition, where participants focused on their heartbeat, and the exteroceptive

condition, where they attended to the sound presented through headphones. **B** Distribution of

Interbeat Intervals (IBI) Across Rest, Exteroception, and Interoception Conditions.

In this study, the MEG data analysis was based on the interoceptive and exteroceptive

tasks with simultaneous sound presentation to avoid overlap between induced activity and

phase-locked activity related to the auditory response. Asynchronous and synchronous

conditions in the interoceptive task were used to assess interoceptive accuracy.

ECG data recording

ECG was measured using Ag/AgCl electrodes placed at the fifth left intercostal space at the

midclavicular line and the right sternum (with the ground electrode on the participant's right

front side below the ribs). The position of the electrode on the sternum was adjusted to ensure

a clear R-peak of the QRS complex was detectable in the ECG signal, which was acquired

using a specially designed device controlled by ARDUINO® software. Custom software script

detected individual R-peaks and controlled the presentation of heart-beat locked auditory

stimuli. The device also sent a trigger marking each heartbeat to the trigger channel in MEG

data. The resulting triggers associated with the R-peak were then applied to define epochs for

further signal analysis.

MEG data recording

MEG data was collected using a whole-head 306-sensor (102 magnetometer channels and 204

planar gradiometer channels) Elekta Neuromag TRIUX system (MEGIN Oy, Helsinki,

Finland) in a magnetically shielded sound-attenuated room. Data were sampled at 1000 Hz and

filtered at 0.1-330 Hz. We measured participants' head shapes using five head position

indicator (HPI) coils that were attached to the scalp (three HPI coils placed on the forehead and

one behind each ear).

MEG data preprocessing and epoch selection

The MEG signal was preprocessed using MaxFilter software (v.2.2) to reduce external noise

through the temporal signal-space separation (tSSS) method (32). Additionally, head

movements were compensated by realigning the head position at each time point to an

"optimal" common reference position (head origin).

For further offline analysis, we utilized the MNE-Python (v 1.7) software (33). The raw

data contaminated with myogenic artifacts were automatically detected using MNE-python

function 'annotate_muscle_zscore' with default parameters (ch_type = 'mag', threshold = 5.0,

min_length_good = 0.2, filter_freq = [110, 140]) and excluded from analysis. After its raw data

were band-passed in 1 - 40 Hz range and resampled offline to 300 readings per second.

The biological artifacts were removed from the continuous data using the fastICA

method implemented in MNE-Python (34). Because temporal signal space separation induces

rank deficiency, we defined the maximum number of ICA components as the rank of raw data.

We used a semi-automated approach implemented in the MNE-python for detection and

removal of electrocardiographic (ECG) and vertical electrooculographic (EOG) artefacts. To

ensure that the automatically detected ICA components were caused by the artefacts, we

checked their timecourses and spatial distributions. The number of cardiac components

removed from the data was 2.3+0.2. We used only gradiometers for analysis because they are

less sensitive to fluctuations in stroke volume and cardiac field artefacts (35).

We extracted R peak locked epochs from −0.500 to 1.500 s relative to the R peaks of

the heart beats. Epochs with a cardiac cycle shorter than 0.7 s were excluded from analysis.

Time-frequency analysis of the MEG data at the source space

Source reconstruction was performed in a template brain (FreeSurfer fsaverage-5.1.0 template) (36). Coregistration between the T1-weighted MRI template scalp and the digitized scalp was performed for each participant using a three-axis scaling mode.

To compute the β-band (16–28 Hz) power we used 'mne.time_frequency.tfr_morlet' function (implemented in the MNE-Python open-source software), with Morlet wavelets using adaptive Morlet constant for each frequency bin (frequency/2) and 2 Hz steps.

A source reconstruction was performed using 'mne.minimum_norm.apply_inverse_tfr_epochs' function, and standardized low-resolution brain electromagnetic tomography (sLORETA) localization method with a source space consisting of 4,098 vertices per hemisphere (37).

The empty room recording was used to construct the noise covariance matrix. The empty room MEG recordings were spatially filtered using the tSSS method and signals were passband 1–40 Hz.

The data were baseline-corrected by dividing β-power at each time point by the averaged power computed over the pre-stimulus interval (from −0.300 to −0.100 s relative to R-peak onset). The resulting values were converted to dB by log10-transformation and multiplication by 10 (38).

Statistical analysis. Regions of interest and time interval selection

The selection of regions of interest (ROIs) we employed a hybrid approach initially identifying anatomical ROIs based on existing literature and subsequently conducting statistical analyses to verify the presence of significant changes in these regions.

It is known that regions such as the insula, anterior and mid-cingulate cortex, medial prefrontal areas, and somatosensory and motor cortex are involved in processing interoceptive

signals (2,6,9,13). Therefore, we selected these regions as potential areas involved in cardiac

interoception.

We then tested whether power in these ROIs significantly differed from baseline. This

was done using a fully flattened average approach applied to data from both the exteroception

and interoception tasks under the simultaneous condition (39). To rule out the possibility of

missing areas with significant R-peak-related power changes, we combined the preselected

ROI approach with a data-driven approach. Therefore, we performed spatiotemporal analysis

on all vertices to identify the time and source distribution of the peak activity. We applied one-

sample permutation test with spatiotemporal threshold-free cluster enhancement (TFCE)

across time (from 0.100 to 1.0 s) and space (8196 vertices) using

mne.stats.permutation_cluster_1samp_test (n_permutations=1000) and starting threshold h0 =

0, step size dh = 0. 4. To minimize computational costs, we applied a stringent cluster-forming

threshold based on the t-distribution with a p-value of 0.05 for data with spatial adjacency.

Next, to precisely refine the boundaries of the preselected ROIs, we overlaid the

obtained activity distribution in the significant TFCE cluster with functional anatomical

regions from the multi-modal atlas from the Human Connectome Project

HCPMMP1_combined and retained only statistically significant vertices (40).

To compare the activity between the interoceptive and exteroceptive tasks, a paired t-

test was used with effect size estimation using Cohen's d. For comparing activity at rest and

during the performance of exteroceptive and interoceptive tasks, repeated measures ANOVA

(rmANOVA) was used with partial eta squared ($\eta p2$) for effect size estimation. Pairwise

comparisons were conducted using Holm–Bonferroni post-hoc tests.

Interoception accuracy

During the experiment, as part of the interoceptive task, participants were required to determine

whether the auditory tones were played simultaneously with their heartbeat or not after each

block of 20 consecutive tones. Interoceptive accuracy was assessed as the ratio of correct

responses to the total number of interoceptive blocks. To investigate the relationship between

interoceptive accuracy and MEG signal power, Spearman's correlation was used.

Acknowledgments

We thank Doris Hernández Barros for her assistance with data collection.

References

1. Craig AD. How do you feel - now? The anterior insula and human awareness. Vol. 10, Nature Reviews Neuroscience. 2009.

2. Critchley HD, Harrison NA. Visceral Influences on Brain and Behavior. Vol. 77, Neuron. 2013.

3. Tsakiris M, Critchley H. Interoception beyond homeostasis: Affect, cognition and mental health. Vol. 371, Philosophical Transactions of the Royal Society B: Biological Sciences. 2016.

4. Park HD, Blanke O. Heartbeat-evoked cortical responses: Underlying mechanisms, functional roles, and methodological considerations. Neuroimage. 2019;197.

5. Tallon-Baudry C, Campana F, Park HD, Babo-Rebelo M. The neural monitoring of visceral inputs, rather than attention, accounts for first-person perspective in conscious vision. Vol. 102, Cortex. 2018.

6. Craig AD. How do you feel - now? The anterior insula and human awareness. Vol. 10, Nature Reviews Neuroscience. 2009.

7. Critchley HD, Nagai Y, Gray MA, Mathias CJ. Dissecting axes of autonomic control in humans: Insights from neuroimaging. Auton Neurosci. 2011;161(1–2).

8. Medford N, Critchley HD. Conjoint activity of anterior insular and anterior cingulate cortex: awareness and response. Vol. 214, Brain structure & function. 2010.

508   9.   Uddin LQ. Salience processing and insular cortical function and dysfunction. Vol. 16,

509   Nature Reviews Neuroscience. Nature Publishing Group; 2015. p. 55–61.

510   10.   Pollatos O, Kirsch W, Schandry R. Brain structures involved in interoceptive

511   awareness and cardioafferent signal processing: A dipole source localization study.

512   Hum Brain Mapp. 2005;26(1).

513   11.   Critchley HD, Wiens S, Rotshtein P, Öhman A, Dolan RJ. Neural systems supporting

514   interoceptive awareness. Nat Neurosci. 2004;7(2).

515   12.   Petzschner FH, Weber LA, Wellstein K V., Paolini G, Do CT, Stephan KE. Focus of

516   attention modulates the heartbeat evoked potential. Neuroimage. 2019;186.

517   13.   Khalsa SS, Rudrauf D, Feinstein JS, Tranel D. The pathways of interoceptive

518   awareness. Nat Neurosci. 2009;12(12).

519   14.   Critchley HD, Wiens S, Rotshtein P, Öhman A, Dolan RJ. Neural systems supporting

520   interoceptive awareness. Nat Neurosci. 2004;7(2).

521   15.   Pollatos O, Traut-Mattausch E, Schroeder H, Schandry R. Interoceptive awareness

522   mediates the relationship between anxiety and the intensity of unpleasant feelings. J

523   Anxiety Disord. 2007;21(7).

524   16.   Thayer JF, Lane RD. A model of neurovisceral integration in emotion regulation and

525   dysregulation. J Affect Disord. 2000;61(3).

526   17.   Pezzulo G, Rigoli F, Friston K. Active Inference, homeostatic regulation and adaptive

527   behavioural control. Vol. 134, Progress in Neurobiology. 2015.

528   18.   Owens AP, Allen M, Ondobaka S, Friston KJ. Interoceptive inference: From

529   computational neuroscience to clinic. Vol. 90, Neuroscience and Biobehavioral

530   Reviews. 2018.

19.   Suarez-Roca H, Mamoun N, Sigurdson MI, Maixner W. Baroreceptor modulation of

the cardiovascular system, pain, consciousness, and cognition. Compr Physiol.

2021;11(2).

20.   Gentsch A, Sel A, Marshall AC, Schütz-Bosbach S. Affective interoceptive inference:

Evidence from heart-beat evoked brain potentials. Hum Brain Mapp. 2019;40(1).

21.   Al E, Iliopoulos F, Forschack N, Nierhaus T, Grund M, Motyka P, et al. Heart-brain

interactions shape somatosensory perception and evoked potentials. Proc Natl Acad

Sci U S A. 2020;117(19).

22.   Kluger DS, Balestrieri E, Busch NA, Gross J. Respiration aligns perception with

neural excitability. Elife. 2021;10.

23.   Karjalainen S, Kujala J, Parviainen T. Neural activity is modulated by spontaneous and

volitionally controlled breathing. Biol Psychol [Internet]. 2025 Apr 1 [cited 2025 May

13];197:109026. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0301051125000444

24.   Zaccaro A, della Penna F, Bubbico F, Bayram B, Parrotta E, Perrucci MG, et al.

Cardio-respiratory interactions in interoceptive perception: The role of heartbeat-

modulated cortical oscillations [Internet]. 2025. p. 1–55. Available from:

http://biorxiv.org/lookup/doi/10.1101/2025.04.14.648690

25.   Luft CDB, Bhattacharya J. Aroused with heart: Modulation of heartbeat evoked

potential by arousal induction and its oscillatory correlates. Sci Rep. 2015;5.

26.   Crivelli, D., Balconi M. Electrophysiological signature of interoceptive attention: a

spectral and source localization analysis. Cogn Affect Behav Neurosci. 2025;402–12.

27.   Engel AK, Fries P. Beta-band oscillations-signalling the status quo? Vol. 20, Current

Opinion in Neurobiology. 2010.

28. Lyyra P, Parviainen T. Behavioral inhibition underlies the link between interoceptive

sensitivity and anxiety-related temperamental traits. Front Psychol. 2018;9(JUN).

29. Garfinkel SN, Seth AK, Barrett AB, Suzuki K, Critchley HD. Knowing your own

heart: Distinguishing interoceptive accuracy from interoceptive awareness. Biol

Psychol. 2015;104.

30. Wiens S, Palmer SN. Quadratic trend analysis and heartbeat detection. Biol Psychol.

2001;58(2).

31. Payne RA, Symeonides CN, Webb DJ, Maxwell SRJ. Pulse transit time measured

from the ECG: An unreliable marker of beat-to-beat blood pressure. J Appl Physiol.

2006;100(1).

32. Taulu S, Simola J, Kajola M. Applications of the signal space separation method.

IEEE Transactions on Signal Processing. 2005;53(9).

33. Gramfort A, Luessi M, Larson E, Engemann DA, Strohmeier D, Brodbeck C, et al.

MEG and EEG data analysis with MNE-Python. Front Neurosci. 2013;(7 DEC).

34. Hyvärinen A. Fast and robust fixed-point algorithms for independent component

analysis. IEEE Trans Neural Netw. 1999;10(3).

35. Buot A, Azzalini D, Chaumon M, Tallon-Baudry C. Does stroke volume influence

heartbeat evoked responses? Vol. 165, Biological Psychology. 2021.

36. Fischl B. FreeSurfer. Vol. 62, NeuroImage. 2012.

37. Pascual-Marqui RD. Standardized low-resolution brain electromagnetic tomography

(sLORETA): Technical details. In: Methods and Findings in Experimental and Clinical

Pharmacology. 2002. p. 5–12.

38. Keil A, Bernat EM, Cohen MX, Ding M, Fabiani M, Gratton G, et al.

Recommendations and publication guidelines for studies using frequency domain and

time-frequency domain analyses of neural time series. Psychophysiology. 2022;59(5).

39.   Bowman H, Brooks JL, Hajilou O, Zoumpoulaki A, Litvak V. Breaking the circularity

in circular analyses: Simulations and formal treatment of the flattened average

approach. PLoS Comput Biol. 2020;16(11).

40.   Glasser MF, Coalson TS, Robinson EC, Hacker CD, Harwell J, Yacoub E, et al. A

multi-modal parcellation of human cerebral cortex. Nature. 2016;536(7615).

41.   Park HD, Correia S, Ducorps A, Tallon-Baudry C. Spontaneous fluctuations in neural

responses to heartbeats predict visual detection. Nat Neurosci. 2014;17(4).

42.   Steinfath P , Herzog N , Fourcade A , Sander C , Nikulin V VA. Validating genuine

changes in Heartbeat Evoked Potentials using Pseudotrials and Surrogate

Procedures.itle. bioRxiv. 2024;

43.   Lee W, Kim E, Park J, Eo J, Jeong B, Park HJ. Heartbeat-related spectral perturbation

of electroencephalogram reflects dynamic interoceptive attention states in the trial-by-

trial classification analysis. Neuroimage [Internet]. 2024 Oct 1 [cited 2025 May

13];299:120797. Available from:

https://www.sciencedirect.com/science/article/pii/S1053811924002945#fig6

44.   Kilavik BE, Zaepffel M, Brovelli A, MacKay WA, Riehle A. The ups and downs of

beta oscillations in sensorimotor cortex. Exp Neurol [Internet]. 2013 Jul 1 [cited 2025

May 14];245:15–26. Available from:

https://www.sciencedirect.com/science/article/abs/pii/S0014488612003767?via%3Dih

ub

45.   Sherman MA, Lee S, Law R, Haegens S, Thorn CA, Hämäläinen MS, et al. Neural

mechanisms of transient neocortical beta rhythms: Converging evidence from humans,

computational modeling, monkeys, and mice. Proc Natl Acad Sci U S A.

2016;113(33).

46. Arnal LH, Giraud AL. Cortical oscillations and sensory predictions. Vol. 16, Trends in
Cognitive Sciences. 2012.

47. Villena-González M, Moënne-Loccoz C, Lagos RA, Alliende LM, Billeke P, Aboitiz
F, et al. Attending to the heart is associated with posterior alpha band increase and a
reduction in sensitivity to concurrent visual stimuli. Psychophysiology. 2017;54(10).

48. Pfurtscheller G, Stancák A, Neuper C. Event-related synchronization (ERS) in the
alpha band - An electrophysiological correlate of cortical idling: A review.
International Journal of Psychophysiology. 1996;24(1–2).

49. Van Elk M, Salomon R, Kannape O, Blanke O. Suppression of the N1 auditory evoked
potential for sounds generated by the upper and lower limbs. Biol Psychol [Internet].
2014 Oct 1 [cited 2025 May 15];102(1):108–17. Available from:
https://www.sciencedirect.com/science/article/abs/pii/S0301051114001173?via%3Dih
ub

50. Pollatos O, Schandry R. Accuracy of heartbeat perception is reflected in the amplitude
of the heartbeat-evoked brain potential. Psychophysiology. 2004;41(3).

51. Corbetta M, Miezin FM, Dobmeyer S, Shulman GL, Petersen SE. Selective and
divided attention during visual discriminations of shape, color, and speed: Functional
anatomy by positron emission tomography. Journal of Neuroscience. 1991;11(8).

52. Hauk O, Stenroos M, Treder MS. Towards an objective evaluation of EEG/MEG
source estimation methods – The linear approach. Neuroimage. 2022;255.

53. Sesa-Ashton G, Wong R, McCarthy B, Datta S, Henderson LA, Dawood T, et al.
Stimulation of the dorsolateral prefrontal cortex modulates muscle sympathetic nerve
activity and blood pressure in humans. Cereb Cortex Commun. 2022;3(2).

54.   James C, Macefield VG, Henderson LA. Real-time imaging of cortical and subcortical

628   control of muscle sympathetic nerve activity in awake human subjects. Neuroimage.

629   2013;70.

630   55.   Riaz B, Eskelin JJ, Lundblad LC, Wallin BG, Karlsson T, Starck G, et al. Brain

631   structural and functional correlates to defense-related inhibition of muscle sympathetic

632   nerve activity in man. Sci Rep. 2022;12(1).

633   56.   Angioletti L, Balconi M. EEG brain oscillations are modulated by interoception in

634   response to a synchronized motor vs. cognitive task. Front Neuroanat. 2022;16.

656