Host iron deficiency protects against *Plasmodium* infection and drives parasite molecular

reprofiling.

Authors

Danielle Clucas, PhD[1,2,3]*, Cavan Bennett, PhD[1,2]*, Rebecca Harding, PhD[1]*, Anne Pettikiriarachchi,

PhD[1], Andrew Baldi, PhD[1], Louise M. Randall, PhD[1], Ryan Steel, PhD[1,2], Ronan Mellin, PhD[1,2], Melissa

Hobbs, BSc[1], Sabrina Caiazzo, PhD[1,2], Martin N. Mwangi, PhD[4], Katherine L. Fielding, MBBS[1], Peter F.

Hickey, PhD[2,5], Tracey M. Baldwin, BAppBiolBiotech[5], Daniela Amann-Zalcenstein, PhD[2,5], Samantha J.

Emery-Corbin, PhD[5,6], Glory Mzembe, MD[7], Ernest Moya, PhD[7], Sabine Braat, MSc[1,10,12], Aaron Jex,

PhD[1], Ayse Y Demir, PhD[8], Hans Verhoef, PhD[8], Kamija S. Phiri, PhD[7,9], Beverley-Ann Biggs, PhD[10],

Wai-Hong Tham, PhD[1,2,11], Justin A. Boddey, PhD[1,2], Sant-Rayn Pasricha, PhD[1,2,3,12]#, Ricardo Ataíde,

PhD[1]#‡

[1]Infection and Global Health, funding

, Parkville, Victoria, Australia.

[2]Department of Medical Biology, The University of Melbourne, Parkville, Victoria, Australia.

[3]Diagnostic Haematology, The Royal Melbourne Hospital, Melbourne, Victoria, Australia.

[4]The Micronutrient Forum, Healthy Mothers Healthy Babies Consortium, Washington, DC, USA.

[5]Advanced Biology and Technology Division, The Walter and Eliza Hall Institute of Medical Research,

Parkville, Victoria, Australia.

[6]Monash Proteomics and Metabolomics Platform, Department of Biochemistry and Molecular Biology,

Monash Biomedicine Discovery Institute, Monash University, Clayton, VIC, Australia.

[7]Training and Research Unit of Excellence, Blantyre, Malawi.

[8]Division of Human Nutrition and Health, Wageningen University, Wageningen, The Netherlands

[9]School of Public and Global Health, Kamuzu University of Health Sciences, Blantyre, Malawi.

[10]Department of Infectious Diseases, Peter Doherty Institute, University of Melbourne, Melbourne,

Victoria, Australia

[11]Research School of Biology, The Australian National University, Canberra, ACT, Australia

[12]School of Population and Global Health, University of Melbourne, Melbourne, Victoria, Australia.

*,# Authors contributed equally

‡Lead contact: Ricardo Ataíde, PhD, The Walter and Eliza Hall Institute, 1G Royal Parade, Parkville, ,

VIC 3052, Australia, ataide.r@wehi.edu.au

Author contributions

S-RP and KSP conceptualized, designed and secured funding for the REVAMP trial. GM, EM and MNM

led data collection in the field. DC, CB, S-RP and RA planned this study. DC, CB, AP, AB, LMR, RS,

RM, SC, KLF, PH, TMB, DA-Z, SJE-C, AY-D, HV, S-RP and RA conducted experimental data

collection. DC, CB, RH, PH, SB and RA led the statistical analysis of the trial. DC, CB, RH, PH, SJE-C,

RA had full access to and analyzed the data. DC, CB, RH, JAB, S-RP and RA interpreted the data. DC,

CB, RH, S-RP and RA prepared the initial draft of the manuscript. All authors critically revised the

manuscript for intellectual content, edited and approved the final version, and agreed to be accountable

for all aspects of the work.

Abstract

Background: Iron deficiency, anemia and *Plasmodium* infection represent significant global health

challenges with overlapping geographical distributions, particularly affecting pregnant women in Africa.

Previous evidence suggests complex interactions between iron status and malaria susceptibility. However,

the mechanisms and clinical implications of this relationship remain poorly understood.

Methods: We employed a multilayered integrated approach to clarify the association between iron

deficiency and malaria infection risk. First, we analyzed clinical data from Malawian pregnant women

(n=711) participating in the REVAMP clinical trial—an RCT of intravenous iron versus oral iron—to

assess associations between iron status and *P. falciparum* parasitemia detected by ultra-sensitive qPCR.

All eligible women received intermittent preventive anti-malaria treatment. Then, we utilized a genetic

mouse model (*Tmprss6*-knockout) to isolate the effect of iron deficiency on *P. berghei* infection and

progression to clinical disease. Finally, we explored direct effects of iron chelation on cultured *P.*

falciparum parasites through transcriptomic and proteomic analyses.

Results: In REVAMP, iron deficiency was associated with a 50% reduced probability of *P. falciparum*

qPCR positivity at baseline (95% CI [30%-64%], p<0.0001). Iron intervention given at baseline did not

significantly modify the probability of subsequent parasitemia across the pregnancy. In the murine model,

iron-deficient *Tmprss6*-knockout mice exhibited significantly improved survival compared to controls

(median survival 15.5 vs. 7.0 days) and protection from cerebral malaria (survival 83% vs 17%). Iron

chelation in *P. falciparum* cultures induced substantial transcriptomic (1,397 differentially expressed

genes) and proteomic changes (121 differentially expressed proteins; 46 matched differentially expressed

gene/protein pairs), primarily affecting processes involved in host cell invasion, protein export, and

nutrient acquisition.

Conclusion: Our findings consistently demonstrate that iron deficiency protects against *Plasmodium*

infection across clinical, pre-clinical, and *in vitro* models. Importantly, in the presence of adequate

malaria prevention intravenous iron supplementation did not significantly increase subsequent parasitemia

prevalence. These results provide mechanistic insights into iron and malaria interactions and support

current WHO recommendations for iron supplementation in pregnant women in malaria-endemic regions

when coupled with adequate malaria prevention strategies.

Funding: Bill and Melinda Gates Foundation, National Health and Medical Research Council of

Australia, and the Victorian State Government Operational Infrastructure Support.

Introduction

Iron deficiency anemia and *Plasmodium* infection remain critical global health problems with significant overlap in their geographical distribution. In 2021, an estimated 1.92 billion people worldwide experienced anemia, while approximately 247 million cases of malaria and 619,000 malaria-related deaths were reported in the same year.[1,2] Across much of sub-Saharan Africa, both iron deficiency and *Plasmodium* infection coexist as significant public health challenges, with anemia commonly resulting from either condition.[3–5]

These conditions appear to interact in complex ways that complicate public health interventions. There are concerns that iron supplementation might increase susceptibility to *Plasmodium* infection. These concerns stem from evidence such as a large randomized controlled trial (RCT) of ~24,000 Tanzanian children in a malaria-endemic area, which demonstrated an increased risk of hospitalization, death, and malaria-specific hospitalization in children receiving oral iron compared to those receiving placebo.[6] In addition, observational evidence suggests that iron deficiency may protect against malaria in children. For example, in a cohort of 785 Tanzanian children followed for up to three years, iron-deficient children had lower odds of subsequent parasitemia and reduced malaria-associated mortality.[7]

The biological mechanisms underlying these interactions are multifaceted. *Plasmodium* infection may induce iron deficiency through several pathways, including hemolysis,[8] inflammation-induced hepcidin upregulation[9], and impaired iron absorption, while, simultaneously, iron deficiency may modify host susceptibility to the parasite.[10–12] *In vitro* studies have demonstrated reduced *P. falciparum* growth and invasion in red blood cells from anemic participants — a protective effect that was reversed when iron supplementation was administered.[8,13–15]

Pregnant women represent a particularly vulnerable population for both iron deficiency and *Plasmodium* infection.[16,17] Approximately one-third of pregnancies in moderate and high malaria transmission

countries in the African region are exposed to malaria. A similar proportion of pregnant women globally

experience anemia,[18] with half of these anemia cases expected to be iron-responsive.[19] Both anemia and

malaria during pregnancy are associated with adverse maternal and neonatal outcomes, including low

birth weight (LBW), which increases the risk of neonatal and infant mortality.[17,20–23] Importantly, iron

supplementation has been linked to increases in birth weight.[24] These complex risk-benefit considerations

underlie the World Health Organization's (WHO) current recommendation for universal iron

supplementation for pregnant women in areas where anemia prevalence exceeds 40%, many of which are

also malaria-endemic.[25]

Despite these recommendations, significant knowledge gaps remain regarding the safety and efficacy of

iron interventions in malaria-endemic regions, particularly for pregnant women. Previous studies have

been limited by challenges in accurately diagnosing iron deficiency during inflammation, confounding by

malaria prevention measures, and difficulty isolating iron status' specific effects on *Plasmodium* infection

from other factors. [26–28]

In this study, we employed a multi-system approach to clarify the association between iron deficiency and

risk of malaria infection. First, we examined clinical data from the REVAMP clinical trial[29] in Malawian

anemic pregnant women to assess the relationship between iron status and *P. falciparum* parasitemia

detected by ultra-sensitive PCR. Second, we utilized genetic mouse model to isolate the effect of iron

deficiency on the progression of *Plasmodium* infection. Finally, we explored the direct effects of iron

chelation on *P. falciparum* parasites *in vitro* through transcriptomic and proteomic analyses. This

comprehensive approach allowed us to test our hypothesis that iron deficiency confers protection against

Plasmodium infection through direct effects on parasite metabolism, which would have implications for

the safety of iron supplementation in malaria-endemic regions.

Methods

Study design and participants

Samples were derived from the REVAMP randomized controlled trial (ANZCTR registration number

ACTRN12618001268235).[29] REVAMP was an open-label, randomized controlled trial comparing a

single dose of intravenous ferric carboxymaltose at enrolment to standard-of-care oral iron for the

duration of pregnancy for anemia recovery in 862 anemic (capillary hemoglobin <10.0 g/dL) second-

trimester pregnant women in Blantyre and Zomba districts in Malawi.[29] All eligible pregnant women

received insecticide treated nets, and intermittent preventative treatment with sulfadoxine/pyrimethamine

(IPTp-SP, standard malaria prophylaxis during pregnancy) as per the Malawian national guidelines—

HIV-positive women received cotrimoxazole.[21] The trial was approved by ethics committees at the

College of Medicine, University of Malawi, Zomba, Malawi; and The Walter and Eliza Hall Institute of

Medical Research, Melbourne, Victoria, Australia.

Procedures

Before enrolment in REVAMP (i.e., pre-screening), women were screened for parasitemia by rapid

diagnostic test (RDT) (SD Bioline Malaria AG P.F/PAN, Standard Diagnostics, Inc.). RDT-positive

women were excluded from enrolment and were treated with artemether-lumefantrine (AL); however, if

they met all other inclusion criteria, they could be rescreened after one week later and enter the trial if

they were malaria microscopy-negative (Figure 1a). Venous blood was collected at enrolment (baseline,

prior to randomization and treatment administration), at 28 days post-treatment, 36 weeks gestation,

delivery, 28 days postpartum, and unscheduled visits when participants presented with illness during the

trial.[30] EDTA whole blood was banked for *P. falciparum* DNA analysis.

Clinical laboratory measurements

Hemoglobin (Hb) concentration was measured on venous blood using a Sysmex XP-300 automated

analyzer (Sysmex, Japan). Serum ferritin and C-reactive protein (CRP) were measured at the Meander

Medical Centre laboratory (accreditation number M040, EN ISO 15189:2012, Amersfoort, The

Netherlands) using the Architect System (Abbott Ireland, Longford, Ireland). The lower limit of detection

for ferritin was 3.1 µg/L, and for CRP was 0.2 mg/L.

Parasitemia detection

DNA was extracted from 400μL of EDTA whole blood using the QIAamp 96 Blood kit (Qiagen). DNA

was also extracted from 200μL of reconstituted WHO International Standard for *Plasmodium falciparum*

DNA Nucleic Acid Amplification Techniques (NIBSC code: 04/176, National Institute for Biological

Standards and Control, UK).[31] Pf-VarATS ultra-sensitive PCR [32] was performed using published probe

and primer sequences and Qiagen QuantiNova Probe PCR master mix, with master mix preparation and

cycling conditions as per manufacturer recommendations. The lower limit of detection was 0.012

parasites/μL of blood. qPCR was performed on the LightCycler 480 II thermocycler (Roche) with

samples run in triplicate.

Pre-clinical models of iron deficiency

Animals and Ethics

The iron deficiency model comprised *Tmprss6*-knockout (KO) mice on a C57Bl/6 background[33].

Tmprss6-heterozygous (Het) littermates and wild-type (WT) animals were used as controls. All animal

experiments were conducted in accordance with the recommendations in the National Statement on

Ethical Conduct in Animal Research of the National Health and Medical Research Council and under

approved requirements set out by the Walter and Eliza Hall Institute of Medical Research (WEHI) Animal

Ethics Committee, Melbourne, Australia (approvals 2017.031, 2019.013, 2021.064, and 2020.034).

Animals were housed in specific pathogen-free conditions with access to standard chow (180mg/kg iron)

and water *ad libitum*.

Mosquito colony maintenance

Anopheles stephensi mosquitoes originally imported from Johns Hopkins School of Public Health, USA,

were reared and maintained in the insectary at the WEHI, Melbourne, Australia according to standard

methods. The mosquitoes were stored in BugDorm insect cages as mixed genders with an environment of

26-27 °C, relative humidity 70%–80%, light/dark photoperiod 12 h:12 h including 30 min ramping to

imitate dawn and dusk. Mosquitoes were fed on reverse osmosis filtered water via cotton wicks and sugar

cubes (sucrose-CSR).

P. berghei sporozoite production

To produce *P. berghei* sporozoites expressing mCherry and luciferase reporters (PbmcherryLuci)[34],

BALB/c 'donor' mice were infected via the intraperitoneal (i.p.) route with blood-stage parasites and 4

days later infected erythrocytes from the donor mice were transferred to naïve "acceptor" mice via i.p. or

i.v. injection. Mice with ≥1% parasitemia and exhibiting exflagellation of microgametes by microscopy at

40x magnification were anesthetized with ketamine/xylazine via i.p. inoculation and individually placed

on top of a single container of 50 female *An. stephensi* (3–5 days old) mosquitoes, allowing them to feed

on mice for 15-30 min, after which any unfed mosquitoes were collected and discarded. Infected

mosquitoes were maintained at 21 °C with 80% humidity and 12 h:12 h light:dark photoperiod with 30

min ramping. Midgut oocysts were determined at 14 days post□blood feeding and at 18 days post□blood

feeding salivary glands were dissected from cold-anesthetized and ethanol killed mosquitoes into

Schneider's Insect Media (Sigma-Aldrich) pH 7.0[35] and quantified in a hemocytometer counting chamber

(Assistent, Neubauer improved) under phase at 400x magnification.

P. berghei infection and IVIS imaging

Mice were infected by intravenous injection of 6500–10000 PbmCherryLuci sporozoites (Day 0). Mice

were euthanized upon reaching any one of the following endpoints: signs of cerebral malaria (loss of self-

righting reflex and hind-limb paresis); >20% parasitemia; or at least two of the following: weight loss

>10%, hunched posture, piloerection, blanching of the tail and footpads, or palpable splenomegaly; or at

Day 20 post-infection (the experiment was terminated at Day 16 post-infection as there were no surviving

WT or HET mice). Liver-stage infection loads were assessed after injecting them with D-Luciferin in PBS

(IVISbrite, PerkinElmer), and imaging using an IVIS Lumina S5 (PerkinElmer).[36] WT and *Tmprss6*-Het

mice were shaved around the abdomen prior to IVIS imaging to account for *Tmprss6*-KO mice having

alopecia. Tail blood smears were performed daily from Day 3 post-infection for blood-stage parasitemia

assessment. Following euthanasia, blood was collected by cardiac puncture and livers were harvested for

RNA extraction and liver iron quantitation.

Gene expression analysis

500ng of RNA was reverse transcribed to cDNA using the SensiFAST cDNA synthesis kit (Bioline)

following the manufacturer's protocol. Reverse transcription-qPCR (RT-qPCR) using the SensiFAST

SYBR No-ROX kit (Bioline) on a LightCycler 480 II thermocycler (Roche) was used to quantify gene

expression levels. Primers and probes used for gene expression analysis and RT-qPCR are listed below:

Species	Gene	Forward sequence	Reverse sequence
Mus Musculus (C57BL6)	*Hamp*	CAGGGCAGACATTGCGATAC	TGCAACAGATACCACACTGGG
Mus Musculus (C57BL6)	*Hrpt*	ACAGGCCAGACTTTGTTGGA	ACTTGCGCTCATCTTAGGCT
Mus Musculus (C57BL6)	*Tmprss6*	TCACCTCGGTCTATGGGTGC	CCCTCCAGTGTCAGGTTCAC
Mus Musculus (C57BL6)	*Gapdh*	GGAGAGTGTTTCCTCGTCCC	ATGAAGGGGTCGTTGATGGC
Plasmodium falciparum (3D7)	*Pf-varATS*	CCCATACACAACCAAYTGGA	TTCGCACATATCTCTATGTCTATCT
Plasmodium falciparum (3D7)	*Pf-varATS-probe*	FAM-TRTTCCATAAATGGT-NFQ-MGB	
Plasmodium falciparum (3D7)	*Pf-18SrRNA DNA*	TATTGCTTTTGAGAGGTTTTGTTG	ACCTCTGACATCTGAATACGAATGC
Plasmodium falciparum (3D7)	*Pf-18SrRNA-probe*	FAM-ACGGGTAGTCATGATTGAGTT-MGB-NFQ	

Pre-clinical specimen analysis

Liver taken into PBS and then snap frozen at -80°C was used to determine hepatic iron content as

previously described.[37] EDTA anticoagulated murine whole blood was analyzed on an Advia2120i for full

blood count determination. Murine plasma iron was determined as previously described.[38]

In vitro effects of iron restriction on P. falciparum

Parasite culture and growth assays

Wild-type *P. falciparum 3D7* asexual blood stage parasites were cultured in Group O RhD+ red blood

cells from healthy donors. Growth assays with desferrioxamine (DFO) (Sigma-Aldrich) were set up using

sorbitol synchronized ring-stage cultures with a starting parasitemia of ~1%, assessed by flow cytometry,

at 1% hematocrit (HCT) and 50µL volume. At 48 hours parasites were stained with Hoechst and

parasitemia analyzed by flow cytometry on a Verse flow cytometer (BD Biosciences, San Jose, USA);

Flow cytometry data were analyzed using FlowJo software version 10.0.8 (Tree Star).

Samples for mass spectrometry and RNA sequencing were obtained from parasite cultures synchronized

using a combination of serial sorbitol treatments and heparin exposure prior to being set up at early ring-

stage (4-6 hours post-invasion) with a ~1% starting parasitemia, 1% HCT and total volume of 10mL with

concentrations of 0 and 12.5µM of DFO. After 30 hours, cultures were treated with saponin to obtain

parasite pellets which were frozen at -80°C for proteomic analysis or resuspended in 50µL of RNAlater

and stored at -20°C for RNA extraction and sequencing. Four biological replicates were performed at

least one week apart, and all samples from the replicates were subsequently processed together.

Mass spectrometry

Parasite pellets were lysed in 8 M urea, 100 mM trisaminomethane TRIS (pH 8.8), with protein

concentrations determined and normalized by BCA assay (Pierce). Proteins were reduced (10 mM DTT),

alkylated (15 mM IAM), and digested overnight with trypsin (1:100). Peptides were acidified, desalted

using C18 tips (in house), vacuum-dried, and reconstituted in 2% ACN, 0.1% TFA for LC-MS analysis.

Mass spectrometry was performed on an Orbitrap Fusion Lumos with FAIMS Pro interface (CVs: –40 V

and –60 V). MS1 scans were acquired at 120k resolution with AGC target $4 \times 10\square$ (normalized AGC

100%) and max injection time (maxIT) of 50 ms. MS2 scans were acquired at 60k resolution with AGC

target $1 \times 10\square$ (normalized AGC 100%) and 50 ms maxIT, All spectra were acquired in positive mode

with full scan MS spectra scanning from m/z 300-1600 using a ±1.6 m/z isolation window and CID at

35% collision energy. Peptide separation was performed by nanoflow reversed-phase HPLC (Ultimate

3000 RSLC, Dionex) using an Acclaim PepMap C18 nano-trap column (75 μm × 2 cm) and analytical

column (75 μm × 50 cm). The gradient was: 3–23% buffer B over 29 min, 23–40% B in 10 min, 40–80%

B in 5 min, held at 80% B for 5 min, then re-equilibrated at 3% B for 10 min. Buffer A: 0.1% FA; buffer

B: 80% ACN, 0.1% FA. The mass spectrometry proteomics data have been deposited to the

ProteomeXchange Consortium via the PRIDE[39] partner repository with the dataset identifier PXD064348.

RNA sequencing

RNA was extracted using the Isolate II RNA mini kit (Bioline) as per the manufacturer's protocol. RNA

was quantified using the Qubit 2.0 HS RNA assay (Thermo Fisher) and the quality assessed using the

2200 TapeStation System High Sensitivity RNA Screen Tape (Agilent). After normalisation to RNA

concentrations of 10ng/µL, sequencing was performed using an in-house mini-bulk sequencing protocol.

In short, single-cell transcriptome libraries were generated by adapting the CelsSeq2 protocol[40] as

follows: samples were pooled after first strand cDNA synthesis, treated with Exonuclease 1 for 30

minutes, followed by a 1.2X bead clean-up. Second strand synthesis was performed using NEBNext

Second Strand Synthesis module (NEB) in a final reaction volume of 20 µL and NucleoMag NGS Clean-

up and Size select magnetic beads (Macherey-Nagel) were used for all DNA purification and size

selection steps. Sequencing was performed on the Illumina NextSeq2000 instrument (San Diego, USA),

using a 100 cycle kit.

Bioinformatic analysis

Briefly, database searching for mass spectrometry data was performed using MaxQuant version

1.6.10.43.[41] Each raw file was split into two mzXML files using FAIMS-MzXML-Generator for the two

CVs used. We assigned the two CVs per raw file as fraction 1 (-40 CV) and fraction 2 (-60CV) across all

the files. The *Plasmodium falciparum* 3D7 isolate genome was downloaded from PlasmoDB (release

51)[42] and used for database searching. For RNA sequencing, expression was quantified by counting the

number of unique molecular identifiers (UMIs) and reads mapped to each gene (*P. falciparum* 3D7

genome downloaded from PlasmoDB release 51, and spike-in transcripts (ERCC)) using scPipe

(v1.14.0).[43] Only exonic reads were included in the differential expression analysis.

In vitro analyses

Data analysis for mass spectrometry was performed as previously outlined.[44] Peptide intensities were

summed, and non-unique peptides, reverse peptide decoys and contaminants removed. Unique peptides

detected in a minimum of 18 samples were log-transformed and quartile-normalised using limma package

version 3.46.0 in R,[45] and missing values imputed using MSImpute using the v1 method.[46] Differentially

expressed peptides between conditions were identified using linear models with empirical Bayes

moderated t-statistics. Differentially expressed proteins (DEPs) were estimated using peptide-set

enrichment analysis in a gene-set enrichment analysis framework.[47] For RNA sequencing analysis,

differential expression was performed using the negative binomial GLM framework workflow in edgeR

(v3.34.0), with enrichment analyses were performed in PlasmoDB. These identified Gene Ontology

(GO)[48] terms and metabolic pathways (from pathway databases - Kyoto Encyclopedia of Genes and

Genomes (KEGG)[49] and MetaCyC[50].

Statistical analysis

Clinical data

Iron deficiency was defined as ferritin <15µg/L, or <30µg/L if CRP was >5mg/L. Associations between

iron status and *P. falciparum* infection and parasitemia at baseline and 28 days post-treatment, as well as

299 the effect of supplementation with IV iron at baseline on the subsequent risk of parasitemia, were analized

300 with Poisson regression models with a log link and robust error variance. Results are presented as point

301 estimates and two-sided 95% confidence intervals (CIs). Additional analyses included adjusted models for

302 pre-specified covariates with the use of directed acyclic graphs as presented in Supplementary figure 1

303 and 2 and sensitivity analysis excluding women who were malaria RDT positive at screening and were

304 rescreened and deemed malaria microscopy-slide negative at enrolment. For analysis of parasite densities

305 across timepoints by treatment group, we calculated median values with interquartile ranges (IQRs). Due

306 to the skewed nature of the data, parasite densities were log-transformed and groups were compared using

307 appropriate statistical testing. Analyses were performed using Stata SE, version 18.0 (StataCorp, College

308 Station, TX).

310 *Pre-clinical models*

311 Sample sizes for animal experiments were calculated based on expected differences in survival between

312 groups with 80% power at an alpha of 0.05, accounting for expected effect sizes based on preliminary

313 data. Statistical testing of continuous variables was performed using Ordinary One-way ANOVA with

314 Dunnet correction for multiple comparisons or Two-way Repeated Measures ANOVA with Dunnett's

315 correction for multiple comparisons. Survival analyses were performed using Gehan-Breslow-Wilcoxon

316 tests. Statistical analyses were performed in Prism version 10.4.1 (GraphPad software).

Results

Sample characteristics

We evaluated a total of 3,141 samples, collected from 711 women in Zomba, Southern Malawi, across 5

timepoints during the trial (Figure 1a, Supplementary Table 1). All enrolled women were anemic by

capillary hemoglobin (Hb), with Hb<10 g/dL. Cohort characteristics were similar between time points,

despite not all participants having samples at all timepoints (Supplementary Table 1). At enrolment,

55.9% (358/641) of women with available samples were primigravid and there was a relatively high

percentage of women with HIV infection (14.2%; 90/634), which was expected in this setting. Iron-

deficiency and inflammation were common, with 39.1% (245/626) of women being iron-deficient and

52.7% (330/626) being inflamed.

Plasmodium falciparum infection by malaria RDT at enrolment

One third (33.1%, 212/641) of the women included in this study were malaria positive at pre-screening by

RDT performed on a finger-prick sample. These women were treated with artemether-lumefantrine (AL),

rescreened (no sooner than one week after AL treatment) and malaria-free by malaria microscopy slide

(Figure 1a, Supplementary Table 1). Once enrolled, all women had a venous sample taken, which was

subsequently used to repeat the malaria RDT—1.4% (9/627) of samples had a venous sample positive

RDT result.

Plasmodium falciparum infection by qPCR

We assessed the prevalence of *P. falciparum* on venous whole blood at all time points by ultra-sensitive

qPCR (Figure 1b). A total of 284/641 (44.3%) of samples were *P. falciparum* positive by qPCR at

enrolment. Excluding from this analysis all the women who were malaria RDT positive at screening and

allowed to enter the trial after being rescreened (after AL treatment and with possible dead parasites or

parasite DNA still in circulation) that prevalence dropped to 35.9% (154/429) (Figure 1b). As all women

who were not rescreened entered the trial after a negative malaria RDT, this shows that a high-proportion

of *P. falciparum* infections go undetected by RDT at this time point in pregnancy in this area of Malawi.

After enrolment, and receipt of the anti-malarial sulfadoxine-pyrimethamine (IPTp-SP), the overall

prevalence of positive qPCR for *P. falciparum* dropped to 18.2% (120/659) at 28 days post-treatment and

11.9% (46/386) at delivery (Figure 1b). The prevalence was similar between those who entered the trial

directly or after rescreen. The overall prevalence of *P. falciparum* positive qPCR remained at 10.8%

(32/295) after delivery and into the 28 days postpartum time point. It is worth noting that 81/284 (28.5%)

of women with a positive qPCR result at baseline were also qPCR positive 28 days post-treatment. We

were not able to assess if these were persisting infections or new infections. At baseline, parasite densities

in those who were qPCR positive, were expectedly low (median (IQR) 1.01 (0.22-9.48) parasites/mL) and

remained so throughout all time points (Figure 1c). Of those enrolled with a negative RDT (not

rescreened), 11.6% (20/171) were positive by qPCR, possibly representing RDT false-negatives

(Supplementary Figure 3 – blue dots). Those who entered the study with a microscopy-negative slide after

rescreening but with a positive qPCR, may have had true sub-microscopic infections, or likely had

circulating dead parasites or parasite DNA (Supplementary Figure 3 – red dots).

Impact of iron status on prevalence and density of parasitemia

We sought to understand the associations between iron status and *P.* falciparum infection and parasitemia.

In the REVAMP women, at baseline, iron-deficiency was associated with a 63% reduction in the

probability of being *P. falciparum* qPCR positive (95% CI [51%-73%]) (Table 1 and Supplementary table

2). This association remained strong when adjusted for gestational age at baseline, gravidity status, HIV,

rescreened + AL treatment, maternal age, education, income source, and religion (50% reduced risk,

[30%-64%]) (Table 1). This association was maintained when excluding the women who were rescreened

(55% adjusted reduced prevalence, 95% CI [34%-69%]) (Table 1 and Supplementary table 3).

We looked at the association of *P. falciparum* qPCR positivity at 28 days post treatment, considering iron-

deficiency status at baseline (Table 1). The strength and direction of these associations were maintained

after adjustment but did not reach statistical significance (prevalence reductions of 34% [-0.8%, 60%],

and 41% [-0.1%, 66%], respectively) (Table 1). Sensitivity analysis was performed to include the WHO

definition of iron deficiency (serum ferritin <15 μg/L or ferritin <70 μg/L if CRP>5 mg/L) revealed

similar patterns (Supplementary Table 2 and 4). We found similar strength and direction of associations

when compared to the main analysis. Additionally, we conducted sensitivity analysis of the association

between qPCR positivity at 28 days post treatment and iron status at baseline in only those women

entering the trial with a negative qPCR result at baseline, thus assessing the risk of becoming infected

(Supplementary table 5). We found no evidence of an effect of iron status at baseline on the risk of

becoming qPCR positive (risk reductions ID versus IR: 5% [-119%, 59%], p=0.91).

Overall quantification of parasitemia in those who were qPCR positive, revealed similar parasite densities

irrespective of iron status at both enrolment (mean difference [95%CI] -0.16 [-0.91, 0.60], p = 0.69) and

28 days post treatment (-0.17 [-2.75, 0.40], p=0.14) (Figure 1d). Similar analysis of parasitemia on those

individuals not rescreened or in those coming into the trial after being rescreened revealed baseline iron-

status was not associated with parasite densities at baseline or at the 28 days post-treatment visit (Figure 1

e-f). Sensitivity analysis of the parasite densities at 28 days post-treatment by iron status at baseline in

those women with a qPCR negative result at baseline revealed a higher median parasite density in iron

replete versus iron deficient women (19.60 (0.33-558.50) parasites/mL versus 3.27 (0.70-439.66)

parasites/mL) (Supplementary table 6).

Impact of iron intervention on prevalence and density of parasitemia

We next investigated the effect of supplementation with IV iron at baseline compared to oral iron on the

subsequent prevalence of parasitemia (Table 2). Treatment with IV Iron was not associated with increased

prevalence of parasitemia positivity by qPCR at any of the subsequent visits when compared to oral iron.

Adjusting the analysis for iron-status at previous visit and time between visits (among others) did not alter

the risk of becoming qPCR positive. The same pattern was observed when excluding from the analysis

those individuals who were enrolled after rescreening (Table 2).

An analysis of the parasite densities across all visits by treatment group revealed higher parasite densities

in the IV Iron group at 28 days post-treatment (N, median (IQR) N=86, 5.66 (0.12 – 601.13)) compared to

the oral iron group (N=85, 0.36 (0.08 – 33.23) (Figure 1g). However, this pattern was inverted at delivery

and 28 days postpartum, with individuals in the IV Iron group presenting reduced parasite densities

(Figure 1g).

Genetic dissection of iron homeostasis in pre-clinical models of malaria

Interpreting the effects of iron status on *Plasmodium* risk in a field cohort is complex. Animal models of

Plasmodium infection allow us to better dissect the timing and effect of iron status. To isolate the effects

of iron status on *Plasmodium* infection and disease burden, we used *P. berghei* infection to induce liver

and blood-stage infection, as well as cerebral malaria, in a mouse model of iron deficiency (*Tmprss6*-KO

mice) (Figure 2a).

Inactivation of *Tmprss6* causes an inability to suppress hepcidin expression, leading to reduced iron

absorption and utilization, plasma iron depletion and iron deficiency phenotype.[51] Uninfected *Tmprss6*-

KO mice had ablated hepatic *Tmprss6* mRNA expression (Figure 2b), leading to increased hepcidin

expression (Figure 2c). *Tmprss6*-KO mice showed reduced plasma iron (Figure 2d), however, hepatic iron

remained unchanged (Figure 2e). As expected, *Tmprss6*-KO mice displayed iron restricted erythropoiesis

with decreased hemoglobin (HGB) concentrations and mean corpuscular volumes (MCVs) and increased

red cell distribution width (RDW) compared to WT mice (Figure 2f-h). *Tmprss6*-Het mice have hepatic

Tmprss6 expression mid-way between WT and *Tmprss6*-KO animals with a complementary increase in

hepcidin (*Hamp*) mRNA mid-way between WT and *Tmprss6*-KO (Figure 2b-c). However, *Tmprss6*-Het

mice display iron and erythroid parameters almost identical to WT animals.

IVIS imaging of animals infected with *P. berghei* sporozoites expressing mCherry and luciferase revealed

Tmprss6-KO mice had comparable intensity of liver infection to that of both WT and *Tmprss6*-Het mice

from 36 to 55 hours post-infection (Figure 2i). In this model of *P. berghei* infection, animals display

cerebral malaria symptoms between days 7 and 10 post-infection. *Tmprss6*-KO mice experienced

significantly improved survival compared to *Tmprss6*-Het and WT animals. The median survival for *P.*

berghei infected *Tmprss6*-KO mice was 15.5 days compared with 7.5 days and 7 days for *Tmprss6*-Het

and WT mice, respectively (p<0.002) (Figure 2j). Survival from cerebral malaria was also significantly

improved in *Tmprss6*-KO mice (83% survival in *Tmprss6*-KO animals vs 17% WT; Figure 2k) and time

to succumb to cerebral malaria was delayed (median time to cerebral malaria: 10 days *Tmprss6*-KO vs 7

days WT; p<0.005 Figure 2l). However, parasitemia from day 3 to day 7 (the timepoint at which most WT

and *Tmprss6*-Het mice succumbed to cerebral malaria), was similar between *Tmprss6*-KO and WT

animals (Figure 2m). Parasitemia was higher in the only WT mouse that survived cerebral malaria, when

compared with the 5 surviving *Tmprss6*-KO mice (Supplementary figure 4).

Direct effects of iron deprivation on Plasmodium falciparum in vitro

Having observed a protective effect of host iron deficiency on *Plasmodium* infection in both our clinical

and pre-clinical models, we finally sought to establish whether iron chelation in parasite cultures

functionally influences *Plasmodium* molecular profiles.

The antimalarial actions of iron chelators on blood-stage *Plasmodium* parasites have been previously

identified but the full molecular effects of this chelation have not been thoroughly defined.[52] We used the

iron chelator desferrioxamine (DFO) in 3D7 *P. falciparum* blood-stage cultures followed by

transcriptomic and proteomic analysis to investigate the effect of iron restriction (Figure 3a). To confirm

an antimalarial action of DFO, we first exposed *in vitro* cultures of 3D7 *P. falciparum* to DFO for 48

hours; a dose dependent inhibition of parasite growth was evident (Figure 3b). Similar to previous reports,

we found the IC50 for DFO to be 16.16µM (95% CI 14.99 – 17.33).[53,54]

To characterise the effect of iron chelation on the transcriptome and proteome of the parasite, we

performed RNA sequencing and quantitative proteomics on synchronous WT 3D7 *P. falciparum* cultures

exposed to 12.5µM of DFO for 30 hours from early ring-stage (4-6 hours post invasion, Figure 3a), a

concentration of DFO expected to affect parasite growth, without causing excessive parasite death that

could render results difficult to interpret.

A total of 1,397 genes were significantly differentially expressed (FDR<0.05) in cultures exposed to DFO

for 30 hours compared with control parasites in media alone, of which 889 were upregulated and 508

downregulated (Figure 3c). At the protein level, 121 proteins were differentially expressed in DFO treated

parasites compared with controls comprising 81 proteins significantly downregulated and 40 upregulated

(Figure 3d). Of these, 46 were differentially expressed at both the RNA and protein level. Most

commonly this differential expression was in the same direction – 32 downregulated in both, 9

upregulated in both (Figure 3e). Among the 46 gene/protein pairs differentially expressed in both

analyses, 15 are associated with merozoite invasion,[55] 5 are characterized as exported proteins,[56] 6 are

associated with the inner membrane complex,[57,58] and 4 are associated with the food vacuole.[59]

Enrichment analysis performed for genes/proteins differentially expressed at both an RNA and protein

level revealed enriched biological process GO terms associated with interaction with the host and entry

into host (Figure 3f).

Discussion

Our multilayered approach to understanding the relationship between iron-deficiency and *P. falciparum* infection has yielded consistent findings across clinical, pre-clinical and *in vitro* data. In Malawian pregnant women in the REVAMP trial—an RCT of IV iron versus standard of care oral iron in anemic women—, iron deficiency at baseline in the second trimester, and prior to iron intervention, was associated with a 50% reduced probability of being qPCR positive for *P. falciparum*. Additionally, iron-deficiency at baseline was associated with a lower probability of *P. falciparum* parasitemia at 28 days post enrolment. These associations were maintained when looking exclusively at women who were malaria RDT-negative at pre-screening, suggesting an effect that is independent from recent malaria treatment. Treatment with IV iron was not associated with an increased prevalence of parasitemia at any timepoint during the trial when compared with the standard of care, oral iron treatment. Complementing these clinical observations, our pre-clinical models using iron-deficient *Tmprss6*-knockout mice exhibited improved survival and reduced cerebral malaria following *P. berghei* infection. Finally, *in vitro* cultures using 3D7 parasites confirmed these antimalarial effects at the molecular level with iron chelation resulting in substantial inhibition of parasite growth and substantial changes in the transcriptome and proteome of *P. falciparum*. Together, our data all converges to support the protective role of iron deficiency against *Plasmodium* infection and disease progression.

Previous studies in pregnant women have been contradictory about the protective effects of iron deficiency.[60,61] Some of that contradiction may be derived from the definition of iron deficiency used in any particular study, as well as from the underlying characteristics of the women; however, a protective association of iron deficiency against *Plasmodium* infection is consistently observed when using ferritin as a biomarker of iron stores.[61] The strong protective association we observed for maternal peripheral infections in the second trimester (50% reduced risk, 95% CI [30%-64%], p<0.0001) is of similar magnitude to the one found in a case-control study conducted in a similar area of Malawi in which placental malaria infections (measured at delivery) were found to be less frequent in women who were

iron-deficient (OR 0.4 [0.2-0.8], p = 0.01).[62] Also, a secondary analysis of an antimalarial RCT conducted

in Papua New Guinea—where the mean hemoglobin at enrolment during the second trimester was 9.7

g/dL— found women who were iron deficient at enrolment were 50% less likely to be *Plasmodium*

positive by microscopy (adjusted odds ratio [aOR] 0.50 [0.38, 0.66], P□<□0.001).[28] While most studies

make concurrent observations of iron deficiency and *Plasmodium* infection—or look at association

between time points that are extremely far apart (second trimester versus delivery)—and are thus unable

to determine the direction of the association, we determined the influence of iron deficiency at baseline

with risk of *Plasmodium* infection 28 days after. Although all women in our cohort received iron

treatment (either IV or oral) the iron utilization over four weeks would still be different between iron-

deficient and replete women. In women overall, as well as excluding those women who received AL

treatment before entering the study, iron deficiency at baseline was still more likely to protect from

Plasmodium infection 28 days after (Table 1). Interestingly, among women who were qPCR negative at

baseline, iron status did not influence the risk of qPCR positivity at 28 days post treatment, likely

reflecting the stronger effect of being on trial (receipt of insecticide treated nets, and antimalarial drugs)

compared to the effect of iron status (Supplementary table 3). Importantly, while iron deficiency was

protective overall, our analysis of IV iron versus oral iron supplementation showed no significant

difference in subsequent parasitemia risk by qPCR across treatment groups. This mirrors the findings of

the main REVAMP trial[29] (which evaluated clinical malaria cases) and those of other iron-

supplementation RCTs in malaria endemic areas[24] and suggests that when appropriate malaria prevention

measures are in place (all women received IPTp-SP or cotrimoxazole), iron supplementation—

independently of the route of administration—can be given safely, supporting current WHO

recommendations for universal iron supplementation in high-anemia-prevalence settings.[63] However, the

transiently higher parasite densities observed in the IV iron group at 28 days post-treatment highlight the

need for vigilant malaria surveillance in the period immediately following iron repletion.

Our genetic mouse model of iron deficiency — genetic knockout of *Tmprss6, a* gene regulating systemic

iron homeostasis[33,51]—provides important mechanistic insights that are difficult to disentangle in human

populations. Additionally, our pre-clinical genetic animal model of infection with sporozoites, replicates

human infection—unlike previous studies that have bypassed liver-stage infection using infected red

cells.[65-69] After infection with *P. berghei* sporozoites, *Tmprss6*-knockout mice displayed comparable liver

infection intensity (Figure 2i), followed by improved survival and protection from cerebral malaria. In

Tmprss6-KO livers—where iron levels are not significantly changed (Figure 2e)—parasite development

seemed unhindered (Figure 2i). Blood-stage infection in these mice, in conditions where iron availability

is restricted and erythropoiesis shows an iron-deficient phenotype (Figure 2d,f-h), also did not differ

(Figure 2m). This mirrors the similar parasite densities observed in women who were iron deficient

compared to those who were iron replete in our clinical cohort (Figure 1d). The low numbers of WT mice

surviving past the cerebral malaria window in this model of infection (N=1) do not allow us to make

conclusions regarding a possible parasitemia control in *Tmprss6*-KO mice past day 7 (Supplementary

Figure 4). We observed a significant increase in survival (Figure 2j-k) and a delayed time to cerebral

malaria in *Tmprss6*-KO mice (median 10 days versus 7 days in WT mice, p<0.005) suggesting that iron

restriction leading to iron-restricted erythropoiesis impacts the pathogenic processes leading to severe

disease manifestations (a finding that can have important clinical implications, as previously reported).[64]

The mechanisms of action of iron-restriction on clinical symptoms could involve altered cytoadherence

properties, reduced inflammatory responses, or changes in endothelial activation—all critical factors in

cerebral malaria pathogenesis.[65,66] This lack of effect on parasitemia in our mouse experiments mirrors

what we observed in our clinical cohort, where iron-deficiency was associated with less risk of infection,

but not with parasite density.

Iron chelation with DFO is not only an *in vitro* tool but also a clinically relevant one.[64] Our transcriptomic

and proteomic analyses of *P. falciparum* cultured under iron-chelated conditions reveal the molecular

adaptations that may underlie the parasite's response to iron restriction. Our transcriptome and proteome

wide analysis of DFO-exposed *P. falciparum* parasite cultures identified substantial changes in mRNA

and protein expression by iron restriction. We identified 1,397 differentially expressed genes and 121

differentially expressed proteins in DFO-treated parasites, with 46 entities showing mostly concordant

changes at both RNA and protein levels (41/46 pairs). Changes were identified in proteins and genes

involved in processes including merozoite invasion, protein export and RNA binding, and associated with

the food vacuole – processes and organelles important for parasite survival, virulence and nutrient

acquisition. The enrichment of GO terms related to host interaction and cell entry further supports the

hypothesis that iron deficiency primarily protects by impairing the parasite's ability to invade and

establish infection within host cells. This molecular *in vitro* evidence bridges our clinical observation of

reduced infection prevalence in iron-deficient women with the improved survival seen in iron-deficient

mice. These data highlight iron as an important nutrient for *P. falciparum* and indicate proteins of interest

for future study.

While our manuscript focussed primarily on the effects of iron on *Plasmodium* biology, there is an

element of interaction between iron and immunity that represents another interface that can affect the

relationship between iron-deficiency and infection. Outside pregnancy, iron's essential role in immune

function is well-established [67,68]. Recent evidence shows that iron status influences antibody responses to

vaccines [69] and may affect antibody functionality. [70] In pregnancy, both the inflammatory and antibody

response to *Plasmodium* are key elements that determine the course of the infection. [21,71] In our pregnant

clinical cohort, pre-existing immunity (or lack of) from previous malaria exposures may interact with iron

deficiency status. The reduced parasitemia risk we observed in iron-deficient women might reflect an

optimal balance where modest reductions in adaptive immunity are offset by direct inhibitory effects on

parasite growth and virulence. Additionally, our transcriptomic and proteomic analyses under iron-

restricted conditions revealed significant changes in parasite-exported proteins, as well as proteins

essential for erythrocyte invasion and intraerythrocytic development (Supplementary table 3). Many of

these proteins are known to interact with the host immune system and modify infected erythrocyte

surfaces, thereby affecting immunogenicity and cytoadherence properties. [72,73] While these potential

immune-related mechanisms provide additional context for our findings, they remain largely speculative

and highlight important areas for future research.

Our findings have several important implications for public health policy and clinical practice. Our data

provide further evidence for the protective effect of iron deficiency on *Plasmodium* infection. The

absence of increased infection risk in our IV iron treatment group (compared to standard of care oral iron)

suggests that rapidly correcting iron deficiency can be done safely in anemic women when appropriate

malaria control measures are in place. However, this reinforces the necessity of careful consideration

when implementing iron supplementation programs in malaria endemic areas. While correcting anemia

remains a critical goal, particularly during pregnancy, iron repletion may transiently increase

susceptibility to malaria highlighting the need for enhanced vigilance and possibly intensified malaria

prevention during the period immediately following iron administration. Finally, our pre-clinical and

molecular findings suggest that under iron-restriction conditions there are substantial transcriptomic and

proteomic changes that are essential for parasite survival and pathogenicity. These can potentially be

targeted as new therapeutics avenues.

Our study has several limitations that should be considered. In our clinical cohort, all women enrolled

were anemic and received one form of iron supplementation. While our models were adjusted for

measured well-known confounders of the associations, estimates may be biased by the presence of

unmeasured confounders. Additionally, the numbers of women with longitudinal data available across the

all trial did not allow for the necessary statistical power to model the interaction between iron-deficiency

and *Plasmodium* infection throughout. We also did not have enough infected women to evaluate the

influence of new infections versus persistent infections in this cohort. In our pre-clinical model, while the

Tmprss6-knockout approach offers several advantages for isolating the impact of iron deficiency, it does

not perfectly recapitulate the complex and often multifactorial iron deficiency observed in pregnant

human populations. Future studies could explore a range of iron deficiency models, including pregnancy

models, with varying severity and mechanisms to better understand dose-response relationships. Our in

vitro experiments used DFO as an iron chelator, which may have effects beyond just iron restriction and

may not replicate the iron-restricted erythropoiesis that is characteristic of iron deficiency. Alternative

approaches using direct manipulation of parasite iron-related genes would strengthen the evidence

regarding the role of iron in parasite biology.

In conclusion, our study demonstrates that iron deficiency protects against *Plasmodium* infection across

clinical, pre-clinical, and molecular contexts. In anemic pregnant Malawian women, iron deficiency

reduced *P. falciparum* parasitemia risk, while iron-deficient mice showed improved survival against

cerebral malaria. Molecular analyses of DFO-exposed parasite cultures revealed iron restriction

significantly alters parasite transcriptomes and proteomes, particularly affecting invasion pathways.

Future research should explore iron-dependent parasite metabolic pathways as potential therapeutic

targets and optimise iron supplementation protocols to balance maternal health benefits with malaria risk

in vulnerable populations.

Role of the funding source

Acknowledgments

The REVAMP trial was funded by the Bill and Melinda Gates Foundation (INV-010612). S.-R.P. is

funded by a National Health and Medical Research Council of Australia (NHMRC) Fellowship

(GNT1158696 and GNT2009047). C.B. is funded by the NHMRC Fellowship (GNT2033705). JAB was

funded by the NHMRC (1139153, 1176955) and is an NHMRC Leadership Fellow. W-H.T. is supported

by an NHMRC Investigator Grant GNT2016908, Victorian State Government Operational Infrastructure

Support and an NHMRC Independent Research Institutes Infrastructure Support Scheme (IRIISS). We

thank Malawian health workers, participants and their families for their involvement in the REVAMP

study. We thank the WEHI animal house staff and the WEHI Insectary facility.

Data availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the

PRIDE[39] partner repository with the dataset identifier PXD064348. Underlying de-identified individual

participant data encompassing the clinical data results and a data dictionary are accessible at figshare

(data available from the corresponding author upon request). All in vitro transcriptomic RNA-Seq data

relevant to the paper are available on Gene Expression Omnibus (*data available from the corresponding*

author upon request).

Conflict of Interest

637   S-RP declares paid advisory board roles for Vifor Pharma for iron and immunity and a role for Vamifeport

638   in sickle cell disease. S-RP reports consultancy for ITL Biomedical on point-of-care devices in iron and

639   research support from WHO. S-RP reports unpaid roles as Director (WHO Collaborating Centre for

640   Anemia Detection and Control). RA declares holding shares in CSL Limited, Australia. All other authors

641   declare no competing interests.

Table 1 - Comparison by ID/IR at baseline and qPCR positivity

	Iron deficient at baseline		Iron replete at baseline		Unadjusted*		Adjusted*	
PCR positive	N	n/N (%)	N	n/N (%)	Effect size (95% CI)	p-value	Effect size (95% CI)	p-value
Baseline*	245	53/245 (21.6)	381	225/381 (59.1)	0.37 (0.27, 0.49)	<0.0001	0.50 (0.36, 0.70)	<0.0001
28 days post treatment+	263	31/263 (11.8)	380	84/380 (22.1)	0.53 (0.35, 0.80)	0.0027	0.66 (0.40, 1.08)	0.096
Excluding those malaria RDT-positive at screening, treated with AL and enrolled after microscopy-negative slide (rescreened)								
Baseline*	225	44/225 (19.6)	191	105/191 (55.0)	0.36 (0.25, 0.51)	<0.0001	0.45 (0.31, 0.66)	<0.0001
28 days post treatment+	242	27/242 (11.2)	189	47/189 (24.9)	0.45 (0.28, 0.72)	0.0009	0.59 (0.34, 1.01)	0.053

Iron deficient indicates serum ferritin <15 μg/L or ferritin <30 μg/L if C-reactive protein >5 mg/L.

*Unadjusted: N/A;

Adjusted: gestational age at baseline, gravidity status, HIV, rescreened + AL treatment, maternal age, education, income source, and religion.

+Unadjusted: treatment group:

Adjusted: treatment group, gestational age at baseline, gravidity status, HIV, rescreened + AL treatment, maternal age, education, income source, and religion.

*Poisson regression model with robust error variance was fit.

Table 2 - Comparison of qPCR positivity by treatment group (IV iron versus Oral iron)

	IV Iron		Oral Iron		Unadjusted *		Adjusted*	
PCR positive	N	n/N (%)	N	n/N (%)	Effect size (95% CI)	p-value	Effect size (95% CI)	p-value
28 days post treatment	334	65/334 (19.5)	325	55/325 (16.9)	1.15 (0.83, 1.59)	0.40	1.14 (0.81, 1.61)	0.46
36 weeks' gestation	182	26/182 (14.3)	195	23/195 (11.8)	1.21 (0.72, 2.05)	0.47	1.45 (0.77, 2.74)	0.25
Delivery	191	17/191 (8.9)	195	29/195 (14.9)	0.60 (0.34, 1.05)	0.072	0.56 (0.30, 1.02)	0.059
28 days postpartum	138	10/138 (7.2)	157	22/157 (14.0)	0.52 (0.25, 1.05)	0.070	0.52 (0.24, 1.14)+	0.10
Excluding those malaria RDT-positive at screening, treated with AL and enrolled after microscopy-negative slide (rescreened)								
28 days post treatment	226	40/226 (17.7)	219	38/219 (17.4)	1.02 (0.68, 1.53)	0.92	0.95 (0.62, 1.44)	0.81
36 weeks' gestation	117	13/117 (11.1)	131	14/131 (10.7)	1.04 (0.51, 2.12)	0.91	1.86 (0.65, 5.31)	0.25
Delivery	120	10/120 (8.3)	131	17/131 (13.0)	0.64 (0.31, 1.35)	0.24	0.67 (0.31, 1.45)	0.31
28 days postpartum	85	5/85 (5.9)	106	16/106 (15.1)	0.39 (0.15, 1.02)	0.056	0.38 (0.14, 1.08)+	0.070

Adjusted: ID at previous visit, anemia at previous visit, HIV status, rescreened + AL treatment, gravidity, maternal age, concurrent gestational age, time between visits.

+: As "Adjusted", excluding concurrent gestational age.

*Poisson regression model with robust error variance was fit (by timepoint). The model estimates the direct effect, not total effect given ID at previous visit, and anemia at previous visit are mediators in the model.

Figure Legends

Figure 1- Trial cohort and *P. falciparum prevalence and parasite density.* (a) Flowchart depicting participant enrolment and sample collection timeline in the REVAMP trial. Women who tested positive by malaria rapid diagnostic test (RDT) at pre-screening received artemether-lumefantrine treatment and were re-screened after at least one week before enrolment. Created with Biorender.com (b) Prevalence of *P. falciparum* infection by ultrasensitive qPCR across study timepoints. All women (blue) or excluding those who were malaria RDT-positive at screening, treated with AL and enrolled after microscopy-negative slide (re-screened) (white). (c) Parasite densities (parasites/µL) in qPCR-positive samples across study timepoints. Box plots show median, interquartile range, and range. The theoretical limits of detection (LOD) for RDT (black) and microscopy (red) are shown on the graph. (d) Parasite densities stratified by iron status (iron-replete (IR) – full circles; iron-deficient (ID) – empty circles) at baseline (magenta) and 28 days post-treatment (green) for all women. (e) Parasite densities stratified by iron status (iron-replete (IR) – full circles; iron-deficient (ID) – empty circles) at baseline (magenta) and 28 days post-treatment (green) excluding those who were malaria RDT-positive at screening, treated with AL and enrolled after microscopy-negative slide (re-screened). (f) Parasite densities stratified by iron status (iron-replete (IR) – full circles; iron-deficient (ID) – empty circles) at baseline (magenta) and 28 days post-treatment (green) in those who were re-screened. (g) Parasite densities stratified by treatment group (IV iron (orange) vs. oral iron (blue)) across timepoints. Lines represent median values with 95% confidence intervals. dpt – days post treatment; wk – weeks; dpp – days postpartum. P-values represent two-sample t-tests (d) and (e) or Mann-Whitney (f).

Figure 2- Iron deficiency in Tmprss6-knockout mice provides protection against P. berghei infection. (a) Experimental design for P. berghei infection in C57BL/6 wild-type (WT), Tmprss6-heterozygous (Het), and Tmprss6-knockout (KO) mice. Mice reach cerebral malaria within 7-10 days post-infection. Created with Biorender.com (b) Hepatic Tmprss6 mRNA expression in uninfected mice.

(c) Hepatic hepcidin (Hamp) mRNA expression. (d) Plasma iron concentration. (e) Non-heme liver iron content. (f-h) Hematological parameters: (f) hemoglobin concentration (HGB), (g) mean corpuscular volume (MCV), and (h) red cell distribution width (RDW). (i) IVIS imaging of liver-stage infection showing parasite burden in all mice at indicated hours post-infection. Representative IVIS images of select mice are shown below. (j) Kaplan-Meier survival curves following P. berghei infection. (k) Percentage of mice succumbing to cerebral malaria (black), high parasitemia (grey) or not reaching humane endpoint (white). (l) Kaplan-Meier survival curve of mice succumbing to cerebral malaria. (m) Blood-stage parasitemia. Data represented as mean ± SD, n=6-9 mice per group (b-c and e), n=4-5 (d), n=6-7 (f-h and j-m), and n=6-8 (I).

Figure 3. Iron chelation slows parasite development and induces extensive transcriptomic and proteomic changes in *P. falciparum*. (a) Experimental design for in vitro analysis of iron chelation on P. berghei. Created with Biorender.com (b) Dose-dependent growth inhibition of 3D7 *P. falciparum* parasites exposed to the iron chelator desferrioxamine (DFO) for 48 hours. (c) Volcano plot showing differentially expressed genes in parasites treated with 12.5μM DFO for 30 hours (889 upregulated [red], 508 downregulated [blue] genes; log□ fold change ≥1, false discovery rate <0.05). (d) Volcano plot of differentially expressed proteins (40 upregulated, 81 downregulated; log□ fold change ≥1, false discovery rate <0.05). (e) Overlap between differentially expressed genes and proteins, showing concordant and discordant regulation. (f) Enriched Gene Ontology (GO) biological process terms for the 46 differentially expressed gene-protein pairs, highlighting processes involved in host cell interaction and invasion. (b) Data represented as mean ± SD, n=6.

References

1.   Gardner, W. M. *et al.* Prevalence, years lived with disability, and trends in anemia burden by severity and cause, 1990–2021: findings from the Global Burden of Disease Study 2021. *Lancet Haematol* **10**, e713–e734 (2023).

2.   *World Malaria Report 2022*. (2022).

3.   Pasricha, S. R., Tye-Din, J., Muckenthaler, M. U. & Swinkels, D. W. Iron deficiency. *The Lancet* **397**, 233–248 (2021).

4.   Camaschella, C. Iron-Deficiency Anemia. *New England Journal of Medicine* **372**, 1832–1843 (2015).

5.   Haldar, K. & Mohandas, N. Malaria, erythrocytic infection, and anemia. *Hematology* 87–93 (2009).

6.   Sazawal, S. *et al.* Effects of routine prophylactic supplementation with iron and folic acid on admission to hospital and mortality in preschool children in a high malaria transmission setting: community-based, randomised, placebo-controlled trial. *The Lancet* **367**, 133–143 (2006).

7.   Gwamaka, M. *et al.* Iron deficiency protects against severe Plasmodium falciparum malaria and death in young children. *Clin Infect Dis* **54**, 1137–1144 (2012).

8.   Clark, M. A. *et al.* Host iron status and iron supplementation mediate susceptibility to erythrocytic stage Plasmodium falciparum. *Nat Commun* **5**, 4446 (2014).

9.   Mohandas, N. & Hillyer, C. D. The iron fist: Malaria and hepcidin. *Blood* **123**, 3217–3218 (2014).

10. Atkinson, S. H. *et al.* Combinatorial effects of malaria season, iron deficiency, and inflammation determine plasma hepcidin concentration in African children. *Blood* **123**, 3221–3229 (2014).

11. Bundi, C. K. *et al.* Iron Deficiency Is Associated With Reduced Levels of Plasmodium falciparum-specific Antibodies in African Children. *Clinical Infectious Diseases* **73**, 43–49 (2020).

12. Muriuki, J. M. *et al.* Malaria is a cause of iron deficiency in African children. *Nat Med* **27**, 653–658 (2021).

13. Goheen, M. M. *et al.* Anemia Offers Stronger Protection Than Sickle Cell Trait Against the Erythrocytic Stage of Falciparum Malaria and This Protection Is Reversed by Iron Supplementation. *EBioMedicine* **14**, 123–130 (2016).

14. Clark, M. A., Goheen, M. M. & Cerami, C. Influence of host iron status on Plasmodium falciparum infection. *Front Pharmacol* **5 MAY**, 1–12 (2014).

15. Goheen, M. M. *et al.* Host iron status and erythropoietic response to iron supplementation determines susceptibility to the RBC stage of falciparum malaria during pregnancy. *Sci Rep* **7**, 17674 (2017).

16. Rogerson, S. J. *et al.* Burden, pathology, and costs of malaria in pregnancy: new developments for an old problem. *Lancet Infect Dis* (2018).

17. Ataide, R., Fielding, K., Pasricha, S.-R. & Bennett, C. Iron deficiency, pregnancy, and neonatal development. *Int J Gynaecol Obstet* **162 Suppl 2**, 14–22 (2023).

18. Desai, M. *et al.* The global burden of malaria in pregnancy: what's known and where are the gaps? *Lancet Infectious Diseases* **7**, 93–104 (2007).

19. WHO 2011. The global prevalence of anemia in 2011. *Who* 1–48 (2011).

20. Umbers, A. J., Aitken, E. H. & Rogerson, S. J. Malaria in pregnancy: small babies, big problem. *Trends Parasitol* **27**, 168–75 (2011).

21. Rogerson, S. J., Hviid, L., Duffy, P. E., Leke, R. F. & Taylor, D. W. Malaria in pregnancy: pathogenesis and immunity. *Lancet Infect Dis* **7**, 105–117 (2007).

22. Levy, A., Fraser, D., Katz, M., Mazor, M. & Sheiner, E. Maternal anemia during pregnancy is an independent risk factor for low birthweight and preterm delivery. *Eur J Obstet Gynecol Reprod Biol* **122**, 182–186 (2005).

23. Moore, K. A., Simpson, J. A., Scoullar, M. J. L., McGready, R. & Fowkes, F. J. I. Quantification of the association between malaria in pregnancy and stillbirth: a systematic review and meta-analysis. *Lancet Glob Health* **5**, e1101–e1112 (2017).

24.   Mwangi, M. N. *et al.* Effect of daily antenatal iron supplementation on plasmodium infection in kenyan women: A randomized clinical trial. *JAMA - Journal of the American Medical Association* **314**, 1009–1020 (2015).

25.   WHO recommendations on antenatal care for a positive pregnancy experience. Preprint at https://apps.who.int/iris/bitstream/handle/10665/250796/9789241549912-eng.pdf?sequence=1 (2016).

26.   Fowkes, F. J. I. *et al.* Iron deficiency during pregnancy is associated with a reduced risk of adverse birth outcomes in a malaria-endemic area in a longitudinal cohort study. *BMC Med* **16**, 1–10 (2018).

27.   Fowkes, F. J. I., Davidson, E., Agius, P. A. & Beeson, J. G. Understanding the interactions between iron supplementation, infectious disease and adverse birth outcomes is essential to guide public health recommendations. *BMC Med* **17**, 1–4 (2019).

28.   Unger, H. W., Bleicher, A., Kaius, M. O., Aitken, E. H. & Rogerson, S. J. Associations of maternal iron deficiency with malaria infection in a cohort of pregnant Papua New Guinean women. *Malar J* 1–11 (2022) doi:10.1186/s12936-022-04177-8.

29.   Pasricha, S.-R. *et al.* Ferric carboxymaltose versus standard-of-care oral iron to treat second-trimester anemia in Malawian pregnant women: a randomised controlled trial. *Lancet* **401**, 1595–1609 (2023).

30.   Mwangi, M. N. *et al.* Protocol for a multicentre, parallel-group, open-label randomised controlled trial comparing ferric carboxymaltose with the standard of care in anemic Malawian pregnant women: the REVAMP trial. *BMJ Open* **11**, e053288 (2021).

31.   World Health Organization. WHO International Standard 1st WHO International Standard for Plasmodium falciparum DNA Nucleic Acid Amplification Techniques NIBSC code: 04/176 Instructions for use. *Medicines & Healthcare products Regulatory Agency* (2012).

32.   Hofmann, N. *et al.* Ultra-Sensitive Detection of Plasmodium falciparum by Amplification of Multi-Copy Subtelomeric Targets. *PLoS Med* **12**, e1001788 (2015).

33. Folgueras, A. R. *et al.* Membrane-bound serine protease matriptase-2 (Tmprss6) is an essential regulator of iron homeostasis. *Blood* **112**, 2539–2545 (2008).

34. Prado, M. *et al.* Long-term live imaging reveals cytosolic immune responses of host hepatocytes against Plasmodium infection and parasite escape mechanisms. *Autophagy* **11**, 1561–1579 (2015).

35. Roth, A. *et al.* Unraveling the Plasmodium vivax sporozoite transcriptional journey from mosquito vector to human host. *Sci Rep* **8**, (2018).

36. Ebert, G. *et al.* Targeting the Extrinsic Pathway of Hepatocyte Apoptosis Promotes Clearance of Plasmodium Liver Infection. *Cell Rep* **30**, 4343-4354.e4 (2020).

37. Bennett, C. *et al.* Serum iron and transferrin saturation variation are circadian regulated and linked to the harmonic circadian oscillations of erythropoiesis and hepatic Tfrc expression in mice. *Am J Hematol* (2024) doi:10.1002/ajh.27447.

38. Bennett, C. *et al.* Iron homeostasis governs erythroid phenotype in Polycythemia Vera. *Blood* (2023) doi:10.1182/blood.2022016779.

39. Perez-Riverol, Y. *et al.* The PRIDE database at 20 years: 2025 update. *Nucleic Acids Res* **53**, D543–D553 (2025).

40. Hashimshony, T. *et al.* CEL-Seq2: Sensitive highly-multiplexed single-cell RNA-Seq. *Genome Biol* **17**, (2016).

41. Tyanova, S., Temu, T. & Cox, J. The MaxQuant computational platform for mass spectrometry-based shotgun proteomics. *Nat Protoc* **11**, 2301–2319 (2016).

42. Amos, B. *et al.* VEuPathDB: the eukaryotic pathogen, vector and host bioinformatics resource center. *Nucleic Acids Res* **50**, D898–D911 (2021).

43. Tian, L. *et al.* scPipe: A flexible R/Bioconductor preprocessing pipeline for single-cell RNA-sequencing data. *PLoS Comput Biol* **14**, e1006361 (2018).

44. Emery-Corbin, S. J. *et al.* Eukaryote-Conserved Methylarginine Is Absent in Diplomonads and Functionally Compensated in Giardia. *Mol Biol Evol* **37**, 3525–3549 (2020).

45.   Ritchie, M. E. *et al.* Limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res* **43**, e47 (2015).

46.   Hediyeh-zadeh, S., Webb, A. I. & Davis, M. J. MSImpute: Imputation of label-free mass spectrometry peptides by low-rank approximation. *bioRxiv* 2020.08.12.248963 (2020) doi:10.1101/2020.08.12.248963.

47.   Alhamdoosh, M. *et al.* Easy and efficient ensemble gene set testing with EGSEA. *F1000Res* **6**, 2010 (2017).

48.   Edgar, R., Domrachev, M. & Lash, A. E. *Gene Expression Omnibus: NCBI Gene Expression and Hybridization Array Data Repository*. vol. 30 http://www.ninds.nih.gov/ (2002).

49.   Kanehisa, M., Goto, S., Kawashima, S. & Nakaya, A. The KEGG databases at GenomeNet. *Nucleic Acids Res* **30**, 42–46 (2002).

50.   Caspi, R. *et al.* The MetaCyc database of metabolic pathways and enzymes - a 2019 update. *Nucleic Acids Res* **48**, D445-d453 (2020).

51.   Finberg, K. E., Whittlesey, R. L., Fleming, M. D. & Andrews, N. C. Down-regulation of Bmp/Smad signaling by Tmprss6 is required for maintenance of systemic iron homeostasis. *Blood* **115**, 3817–3826 (2010).

52.   Moormann, A. M., Hossler, P. A. & Meshnick, S. R. *Deferoxamine Effects on Plasmodium Falciparum Gene Expression. Molecular and Biochemical Parasitology* vol. 98 (1999).

53.   van Zyl, R. L., Havlik, I. & Monteagudo, F. S. The combined effect of iron chelators and classical antimalarials on the in-vitro growth of Plasmodium falciparum. *J Antimicrob Chemother* **30**, 273–278 (1992).

54.   van Zyl, R. L., Havlik, I., Hempelmann, E., MacPhail, A. P. & McNamara, L. Malaria pigment and extracellular iron. Possible target for iron chelating agents. *Biochem Pharmacol* **45**, 1431–1436 (1993).

55. Ginsburg, H., Gilberger, T. & Pierce, A. Malaria Parasite Metabolic Pathways: Functional annotation of merozoite invasion-related proteins. vol. 2023 https://mpmp.huji.ac.il/maps/Merozoiteproteins.html.

56. Proellocks, N. I., Coppel, R. L., Mohandas, N. & Cooke, B. M. Chapter One - Malaria Parasite Proteins and Their Role in Alteration of the Structure and Function of Red Blood Cells. in *Advances in Parasitology* (eds. Rollinson, D. & Stothard, J. R.) vol. 91 1–86 (Academic Press, 2016).

57. Wichers, J. S. *et al.* Identification of novel inner membrane complex and apical annuli proteins of the malaria parasite Plasmodium falciparum. *Cell Microbiol* **23**, e13341 (2021).

58. Ferreira, J. L. *et al.* The Dynamic Roles of the Inner Membrane Complex in the Multiple Stages of the Malaria Parasite. *Front Cell Infect Microbiol* **10**, 611801 (2020).

59. Lamarque, M. *et al.* Food vacuole proteome of the malarial parasite Plasmodium falciparum. *Proteomics Clin Appl* **2**, 1361–1374 (2008).

60. Moya-Alvarez, V. *et al.* Does iron increase the risk of malaria in pregnancy. *Open Forum Infect Dis* **2**, (2015).

61. Sangaré, L., van Eijk, A. M., Ter Kuile, F. O., Walson, J. & Stergachis, A. The association between malaria and iron status or supplementation in pregnancy: a systematic review and meta-analysis. *PLoS One* **9**, e87743 (2014).

62. Senga, E. L., Harper, G., Koshy, G., Kazembe, P. N. & Brabin, B. J. Reduced risk for placental malaria in iron deficient women. *Malar J* **10**, 47 (2011).

63. WHO. *Iron and Folate Supplementation: Intergrated Management of Pregnancy and Childbirth (IMPAC). Standards for Maternal and Neonatal Care developed by the Department of Making Pregnancy Safer* www.who.int/making_pregnancy_safer/publications/en (2006).

64. Gordeuk, V. *et al.* Effect of Iron Chelation Therapy on Recovery from Deep Coma in Children with Cerebral Malaria. *New England Journal of Medicine* **327**, 1473–1477 (1992).

65. Carvalho, L. J. M. Murine cerebral malaria: how far from human cerebral malaria? *Trends Parasitol* **26**, 271–2 (2010).

66. Rénia, L., Grüner, A. C. & Snounou, G. Cerebral malaria: in praise of epistemes. *Trends Parasitol* **26**, 275–7 (2010).

67. Ni, S., Yuan, Y., Kuang, Y. & Li, X. Iron Metabolism and Immune Regulation. *Frontiers in Immunology* vol. 13 Preprint at https://doi.org/10.3389/fimmu.2022.816282 (2022).

68. Nairz, M. & Weiss, G. Iron in infection and immunity. *Molecular Aspects of Medicine* vol. 75 Preprint at https://doi.org/10.1016/j.mam.2020.100864 (2020).

69. Stoffel, N. U. *et al.* Iron Deficiency Anemia at Time of Vaccination Predicts Decreased Vaccine Response and Iron Supplementation at Time of Vaccination Increases Humoral Vaccine Response: A Birth Cohort Study and a Randomized Trial Follow-Up Study in Kenyan Infants. *Front Immunol* **11**, (2020).

70. Hassan, T. H. *et al.* Impact of iron deficiency anemia on the function of the immune system in children. *Medicine (United States)* **95**, (2016).

71. Aitken, E. H., Mahanty, S. & Rogerson, S. J. Antibody effector functions in malaria and other parasitic diseases: a few needles and many haystacks. *Immunology and Cell Biology* vol. 98 264–275 Preprint at https://doi.org/10.1111/imcb.12320 (2020).

72. Zelter, T. *et al.* Neutrophils impose strong immune pressure against PfEMP1 variants implicated in cerebral malaria. *EMBO Rep* **23**, (2022).

73. Sampaio, N. G. *et al.* Extracellular vesicles from early stage Plasmodium falciparum-infected red blood cells contain PfEMP1 and induce transcriptional changes in human monocytes. *Cell Microbiol* **20**, (2018).

Figure 1

Figure 2

Figure 3